Action does not enhance but attenuates predicted touch

Xavier Job[1] and Konstantina Kilteni[1,2]*

[1]Department of Neuroscience, Karolinska Institute, Stockholm, Sweden. [2]Department of Sensorimotor Neuroscience, Donders Institute, Radboud University, Netherlands

Abstract

Dominant motor control theories propose that the brain predicts and attenuates the somatosensory consequences of actions, referred to as somatosensory attenuation. Support comes from psychophysical and neuroimaging studies showing that touch applied on a passive hand elicits attenuated perceptual and neural responses if it is actively generated by one's other hand, compared to identical touch from an external origin. However, recent experimental findings have challenged this view by providing psychophysical evidence that the perceived intensity of touch on the passive hand is enhanced if the active hand does not receive touch simultaneously with the passive hand (somatosensory enhancement) and by further attributing attenuation to the double tactile stimulation of the hands upon contact. Here, we directly contrasted the hypotheses of the attenuation and enhancement models regarding how action influences somatosensory perception by manipulating whether the active hand contacts the passive hand. We further assessed somatosensory perception in the absence of any predictive cues in a condition that turned out to be essential for interpreting the experimental findings. In three preregistered experiments, we demonstrate that action does not enhance the predicted touch (Experiment 1), that the previously reported "enhancement" effects are driven by the reference condition used (Experiment 2), and that self-generated touch is robustly attenuated regardless of whether the two hands make contact (Experiment 3). Our results provide conclusive evidence that action does not enhance but attenuates predicted touch and prompt a reappraisal of recent experimental findings upon which theoretical frameworks proposing a perceptual enhancement by action prediction are based.

Highlights

- Dominant motor control theories propose that action attenuates or cancels predicted touch.
- Recent theories propose that action enhances predicted touch.
- We show that action does not enhance but attenuates predicted touch.

Keywords

somatosensory attenuation; prediction; cancellation; sharpening; enhancement

Introduction

Dominant motor control theories propose that the brain uses an internal forward model in combination with a copy of the motor command (efference copy) to predict the sensory consequences of our movements [1–3]. For example, the brain predicts the upcoming touch as one reaches towards an object. These predictions allow for the correction of motor errors without relying on sensory feedback that suffers from intrinsic delays [4], thereby improving the estimation of the current state of our body by combining the predicted touch with the actual sensory input [1,5,6]. These predictions *attenuate* the perception of the self-generated input (sensory reafference) compared to that of externally generated input [7–9] and infer whether the cause of the sensory input is the self or the environment [10–12]. A classic example of this attenuation is that we are unable to tickle ourselves with our own touch, yet we are easily tickled by the touch of others [13]. The attenuation of sensory reafference – also referred to as sensory cancelation – is thought to be necessary to compensate for the limited capacity of the sensory systems by optimally prioritising the perception of more informative externally generated stimuli [2,14]. Thus, the attenuation model proposes that we dampen perceptual representations of expected self-generated stimuli to reduce redundancy and to highlight behaviourally relevant unexpected externally generated stimuli.

In contrast, an alternative theoretical framework proposes that predictions, including those arising from our motor commands, should not attenuate but instead enhance sensory signals, thereby allowing for sharper (*i.e.,* more accurate) representations of predicted compared to unpredicted sensory events [15–17]. This *enhancement* account – also referred to as the sharpening account – posits that predictions based on our motor commands are equivalent to expectations formed by statistical regularities in sensory input or from prior knowledge (*e.g.*, at the North Pole, one expects to see a polar bear rather than an elephant) and that these predictions should bias our perception towards our expectations. The proposal mainly stems from experimental research outside the domain of action and argues that weak, noisy, or ambiguous sensory input that is in line with prior expectations should be enhanced to achieve, on average, more accurate representations. For example, we are more biased to report the presence of visual events that are statistically likely to occur rather than unlikely events [18,19], more sensitive to low-level visual features that are in line with prior expectations [20,21], and show greater biases when perceiving visual events that are congruent with our expectations [22,23]. Such effects are thought to result from mechanisms that increase the "gain" of expected information by altering the weights of different sensory signals [24]. Thus, the enhancement model proposes that we amplify the perceptual representations of expected compared to unexpected sensory input.

In the somatosensory domain, evidence supporting the attenuation model has repeatedly shown that touch delivered to one (passive) hand by the other (active) hand (*i.e.*, self-generated touch) is perceived as weaker or less ticklish [25–39] and evokes attenuated somatosensory cortical activation [40–44] compared to touch of identical intensity applied on the passive hand that is externally generated. In contrast, evidence supporting the enhancement model has shown that the action of the active hand results in an increase in

the perceived intensity of touch on the passive hand, provided that the active hand never receives touch simultaneously with the passive hand (*i.e.*, hands do not make contact) [45]. This enhancement finding has been recently used to support the sharpening model and to argue that attenuation effects are due to unspecific gating processes caused by the simultaneous double tactile stimulation of the two hands [46].

The attenuation (or cancellation) and enhancement (or sharpening) models present strikingly different hypotheses regarding how action influences the perception of sensory input and are supported by contradictory experimental evidence, leading to debates between researchers [11,15,27,46,47]. Clarifying how predictions about the sensory consequences of our movements affect our perception is fundamental to understanding the interaction between perception and action [1–3] but also for clinical and neurobiological theories of psychosis spectrum disorders, such as schizophrenia [44,48–54] and schizotypy [25], as well as functional movement disorders [55], Parkinson's disease [56] and ageing [29].

The present study aimed to contribute to this debate by directly contrasting the enhancement and attenuation models within the same paradigm. To this end, the same force discrimination task employed in earlier studies reporting attenuation [25–27,32,34,35,39,57] and enhancement [45] was used to determine (a) whether movement of the right hand enhances or attenuates the perceived magnitude of touch applied on the left hand when the two hands do not make contact (Experiments 1 and 2) and (b) whether attenuation effects are due to double tactile stimulation caused by the contact of the two hands or action prediction (Experiment 3). Capitalizing on the fact that *any* conclusion about whether action prediction "attenuates" or "enhances" the perception of the somatosensory input needs to be made with a comparison to one's somatosensory perception in the absence of action, we also included a condition in which participants passively received externally generated touch, with which we compared the participants' perception in all experimental conditions. Consequently, if participants perceive a touch as less or more intense during action than in the absence of action, we can infer that the received touch was attenuated or enhanced, respectively. This is a critical methodological novelty compared to previous studies [32,45] because if such baseline conditions are missing, the same patterns of results can be incorrectly attributed to "attenuation" or "enhancement": for example, if one condition produces less attenuation than another, it may be interpreted as enhancement, and vice versa. Indeed, it was this baseline condition that turned out to be critical for distinguishing between attenuation and enhancement effects. All studies and analysis plans were preregistered on the Open Science Framework prior to data collection (**Methods**).

Results

Experiment 1. Action does not enhance predicted touch.

Thirty naïve participants moved their right index finger towards their left index finger to generate the touch on their left index finger with (*contact* condition) or without (*no-contact* condition) simultaneous stimulation on their active finger. A *baseline* condition in which the participants did not move their right index finger and received touch on the

left index finger passively (externally generated touch) was included in order to distinguish between effects of attenuation or enhancement (**Figure 1A**).

During the force-discrimination task, participants judged the intensity of a *'test'* force and a *'comparison'* force (100 ms duration each) separated by a random interval ranging from 800 ms to 1200 ms. The forces were delivered to the pulp of the left index finger by a cylindrical probe attached to a lever driven by a DC electric motor. In each trial, participants reported which force they felt was stronger (the *test* or the *comparison*). The intensity of the *test* force was fixed at $2\,N$, while the intensity of the *comparison* force was systematically varied among seven force levels (1, 1.5, 1.75, 2, 2.25, 2.5 or $3\,N$). In the *contact* condition, participants moved their right index finger towards their left index finger after an auditory cue and actively tapped on a force sensor placed on top of, but not in direct contact with, the probe. The participant's active tap on the force sensor triggered the *test* force on their left index finger, thereby producing simultaneous stimulation of both fingers in the *contact* condition and the sensation of pressing with the right index finger against the left index finger through a rigid object. Participants were told that they would always make contact between their fingers (through the probe) in this condition. In the *no-contact* condition, following the same auditory cue, participants moved their right index finger towards their left index finger. At the beginning of the condition, the force sensor was removed and replaced with a distance sensor that detected the relative distance of their active finger as it approached their left index finger to trigger the *test* force, similar to [32,45]. The distance threshold was set such that the position of the right index finger when triggering the *test* force was equivalent to that in the *contact* condition. Participants were told that they would never make contact between their fingers. Before the experiment, participants were trained to make similar movements with their right index finger in the *contact* and *no-contact* conditions. Finally, in the *baseline* condition (externally generated touch), participants were told to relax their right hand, and each trial began with the same auditory cue followed by the *test* force delivered to the participants' left index finger 800 ms after the cue. In all trials, the *comparison* force was delivered after the *test* force, and participants verbally reported their judgement (**Methods**).

Participants' responses were fitted with logistic psychophysics curves, and we extracted the point of subjective equality (PSE), which represents the intensity of the comparison force at which the test force feels equally as strong. Consequently, a PSE in a movement condition (*contact* or *no-contact*) that is lower than the PSE of the *baseline* condition indicates attenuation, while a PSE that is higher than the PSE of the *baseline* condition indicates enhancement (**Methods**). According to the attenuation model (**Figure 1B**), attenuation of the somatosensory input on the left hand in the *contact* condition compared to the *baseline* condition should be observed (*i.e.*, lower PSEs) due to the *test* force being a predicted consequence of the action of the right hand. However, regarding the *no-contact* condition, earlier studies have shown that the mere movement of one hand is not sufficient to produce predictions of somatosensory input simultaneously applied on the other hand [26,36–38,42,56,58]. Instead, a bimanual sensorimotor context is needed such as during bimanual object manipulation and bimanual contact of the hands (Blakemore et al., 1998). Thus, while touch in the *no-contact* condition is *self-generated* in the sense

that it is a consequence of one's movement, the efference copy and the sensory feedback are inconsistent with the sensorimotor context of touching oneself; in this case, the context of applying force on one's finger with another finger. In other words, the attenuation model suggests that not any voluntary movement will attenuate upcoming sensory input. Based on this, neither attenuation nor enhancement should be observed in the *no-contact* condition (*i.e.*, no change in PSEs from *baseline*). In contrast, according to the enhancement model, if action enhances the received sensation [45], then higher PSEs in the *no-contact* condition compared to the *baseline* condition should be observed (**Figure 1C**). Finally, since the enhancement model proposes that attenuation effects are attributed to unspecific (nonpredictive) gating effects caused by the simultaneous tactile stimulation of the two hands, touch in the *contact* condition should also be perceived as weaker, albeit not due to action prediction. In summary, Experiment 1 explicitly assessed whether action enhances the received touch when the index fingers of the two hands do not make contact.

The results showed a robust attenuation of the perceived touch when the two fingers made contact (*contact* condition) (**Figure 1D, E, F**): the PSEs were significantly lower in the *contact* condition than in the *baseline* condition ($W = 422.00$, $p < 0.001$, $rrb = 0.82$, $CI^{95} = [0.08\ 0.25]$). Similarly, the PSEs were significantly lower in the *contact* condition than in the *no-contact* condition ($W = 441.00$, $p < 0.001$, $rrb = 0.90$, $CI^{95} = [0.13\ 0.26]$ (**Figure 1D, E, G**). Critically, however, in the comparison that contrasts the hypotheses of the two models, the *no-contact* condition did not produce any significant change in the perceived magnitude of the touch compared to the *baseline* condition ($W = 187.00$, $p = 0.360$, $rrb = -0.20$, $CI^{95} = [-0.08\ 0.030]$) (**Figure 1D, E, H**). A Bayesian factor analysis provided moderate support for the absence of any difference ($BF_{01}=3.58$). (See also **Supplementary Material** for individual fits and additional analyses).

In summary, we used both frequentist and Bayesian statistics and found no evidence that action of the right index finger produces an enhancement of the touch received on the left index finger when the fingers did not make contact. Thus, Experiment 1 does not support the hypothesis of the enhancement model that action enhances predicted touch but supports the hypothesis of the attenuation model.

Figure 1. Experimental methods, hypotheses, and results of Experiment 1. A. In the *contact* condition (magenta), participants tapped with their right index finger (R) on a force sensor placed above the probe that delivered a *test* force of 2 N to their left index finger (L), followed by a

comparison force randomly varying between 1 and 3 N. In the *no-contact* condition (green),
participants approached a distance sensor with their right index finger, which triggered the test
force on their left index finger, thus receiving no touch on their active finger. In the *baseline*
condition (blue), participants relaxed both hands and passively received the forces on their left
index finger. **B.** Hypotheses based on the attenuation model. Touch in the *contact* condition
(magenta) should be perceived as weaker than in the *baseline* (blue), but touch in the *no-contact*
condition (green) should be perceived similarly to that in the *baseline* (blue). **C.** Hypotheses
based on the enhancement model. Touch in the *no-contact* condition (green) should be perceived
as stronger than in the *baseline* (*i.e.*, enhanced) (blue), but touch in the *contact* condition
(magenta) should be perceived weaker than the *baseline* (blue). Note that attenuation effects in
the *contact* condition are predicted both by the attenuation and the enhancement model with the
difference that the attenuation model attributes these effects to action prediction, while the
enhancement model attributes these effects to the simultaneous touch on the active hand. **D-H.**
Data are colour-coded per condition. **D.** Box plots show the median and interquartile ranges for
the PSE values per condition, black circles and error bars show the mean PSE ± s.e.m, and the
raincloud plots show the individual PSE values and their distributions. No enhancement effects
were observed in the *no-contact* condition. **E.** Group psychometric functions for each condition.
The leftward shift of the curves in the *contact* condition indicates attenuated PSE values
compared to the other two conditions. **F-H**. Line plots for individual participant PSE values
illustrating significantly lower PSEs for the *contact* versus *baseline* (**F**) and *no-contact* (**G**)
conditions, but no significant differences in the PSE values between *no-contact* and *baseline* (**H**).
(***$p <.001$)

Experiment 2. Previous 'enhancement' effects are driven by the baseline used.

Having found no evidence for somatosensory enhancement in Experiment 1, Experiment
2 aimed to understand the potential source of the previously reported enhancement effects
[45]. One critical methodological difference between Experiment 1 and previous evidence
for enhancement concerns the *baseline* condition, as Thomas et al. [45] compared the
participants' perception in the *no-contact* condition to a condition where the participants
prepared their right index movement but received a NOGO cue to inhibit the movement,
and it was that comparison that revealed enhancement effects. However, motor inhibition
(i.e. planning the hand movement but not executing it) can lead to suppression of
somatosensory input both on the hand that is planned to move [60-62] and the hand that
would receive touch if the movement was executed [38]; moreover, such conditions result
in a competition for attentional resources to inhibit the motor response with processes
that encode sensory stimuli into memory [63-65] (*e.g.*, the *test* force in this paradigm).
Therefore, if the motor inhibition condition used by Thomas et al. [45] results in a
suppression of the perceived touch, a comparison of the *no-contact* condition with such a
baseline would produce an apparent enhancement effect.

Thirty new naïve participants participated in Experiment 2, which included a block of
contact trials and a block of *no-contact* trials in which participants moved their right
index finger, but each block also contained randomly intermixed NOGO trials where
participants were cued to withhold their movement, identical to the design of Thomas et
al. [45] (**Figure 2A**). Participants were trained to make similar movements with their right
index finger in the *contact* and *no-contact* trials. An externally generated touch condition
was also included as an action-free *baseline* condition (*i.e.*, no action execution or

inhibition). If the previously reported enhancement effects are due to the baseline used, attenuated perception of touch during the motor inhibition condition (*i.e.*, NOGO trials) should be found compared to the *baseline*, which would then lead to apparent enhancement effects upon comparison with the *no-contact* trials (**Figure 2B, C**) (**Methods**).

This hypothesis (**Figure 2B**) was confirmed. First, all the effects of Experiment 1 were replicated in the new sample (**Figure 2D, E, F & H**). The *contact* trials yielded significant attenuation (*i.e.*, lower PSEs) compared to the *baseline* condition ($t(29) = 6.79$, $p < 0.001$, $d = 1.24$, $CI^{95} = [0.18\ 0.33]$) (**Figure 2E & H**). Once again, there was no enhancement in the *no-contact* trials compared to the *baseline* condition ($t(29) = 0.45$, $p = 0.658$, $d = 0.08$, $CI^{95} = [-0.05\ 0.07]$) (**Figure 2D & F),** and the Bayesian analysis again yielded moderate evidence for the absence of any effects ($BF_{01} = 4.69$). Importantly, the PSEs in the *NOGO* (motor inhibition) trials were significantly lower than the *baseline* condition both for the *contact NOGO* ($t(29) = 2.99$, $p = 0.006$, $d = 0.55$, $CI^{95} = [0.02\ 0.09]$) and the *no-contact NOGO* trials ($t(29) = 4.44$, $p < 0.001$, $d = 0.81$, $CI^{95} = [0.05\ 0.13]$) (**Figure 2D, E & G**). This demonstrates that NOGO trials resulted in a suppression of perceived touch on the left hand. Critically, this led to an apparent increase in the PSE from the *no-contact* trials to the NOGO trials in the no-contact block ($t(29) = -2.98$, $p = 0.006$, $d = -0.54$, $CI^{95} = [-0.12\ -0.02]$), mimicking an 'enhancement' effect. Finally, PSEs were significantly lower in the *contact* trials than in the NOGO trials ($t(29) = 5.91$, $p < 0.001$, $d = 1.08$, $CI^{95} = [0.13\ 0.27]$), while the NOGO trials in the *contact* and *no-contact* blocks did not significantly differ ($t(29) = 1.58$, $p = 0.126$, $d = 0.29$, $CI^{95} = [-0.01, 0.08]$, $BF_{01} = 1.697$). (See also **Supplementary Material** for individual fits and additional analyses).

In summary, identical to Experiment 1, we used both frequentist and Bayesian statistics and did not find any evidence that action of the right index finger produces an enhancement of the touch received on the left index finger when the fingers do not make contact. Moreover, we showed that the purported "enhancement" effect [45] is driven by a suppression of the perceived intensity of touch following a cue to inhibit the planned movement. Thus, Experiment 2 does not support the hypothesis that action enhances predicted touch.

Figure 2. Experimental methods, hypotheses, and results of Experiment 2. A. The *contact* (magenta) and *no-contact* (green) trials were identical to those of Experiment 1, with the only difference that in 50% of the trials, the participants had to inhibit their movement (*NOGO* trials - yellow), and the test force was delivered automatically. The *baseline* condition (blue) was identical to that of Experiment 1. **B.** Hypotheses based on the attenuation model. Touch in the *contact* condition (magenta) should be perceived as weaker than in the *baseline* (blue), but touch in the *no-contact* condition (green) should be perceived similarly to that in the *baseline* (blue). Critically, touch may also be perceived as weaker than *baseline* in the *NOGO* trials, resulting in a 'false enhancement' of the *no-contact* trials. **C.** Hypotheses based on the enhancement model. Touch in the *no-contact* condition (green) should be perceived as stronger than in the *baseline* (*i.e.*, enhanced) (blue), but touch in the *contact* condition (magenta) should be perceived weaker than the *baseline* (blue). **D-H.** Data are colour-coded per condition. **D-E.** Box plots show the median and interquartile ranges for the PSE values in the *baseline* and *no-contact* blocks (**D**) and in the *baseline* and *contact* blocks (**E**). Black circles and error bars show the mean PSE ± s.e.m, and the raincloud plots show the individual PSE values and their distributions. **F-H.** Line plots for individual participant PSE values illustrating no significant differences in the PSE values between *no-contact* and *baseline* (**F**), significantly higher PSEs for the *no-contact* versus *NOGO* trials in the same block (**G**) and significantly lower PSEs in the *contact* versus *baseline* trials (**H**). (*******p* < *0.01*, ********p* < 0.001)

Experiment 3. Action attenuates predicted touch, even without simultaneous stimulation of the active hand.

Within the framework of internal forward models, the absence of attenuation in the *no-contact* conditions of Experiments 1 and 2 is expected, given the lack of a sensorimotor context conducive to perceiving the touch as self-generated: when contact is never made, the forces applied on the passive left hand are only arbitrarily, and not causally, associated with the movement of the active right hand. Alternatively, it could be argued that the absence of attenuation in the *no-contact* conditions is caused by the lack of simultaneous tactile stimulation of the active hand rather than by predictive mechanisms.

In Experiment 3, this hypothesis was explicitly tested with thirty new naïve participants. The same *contact* and *no-contact* conditions were included as in Experiment 1, but their relative frequency was manipulated within the same block (*contact* trials: 80%, *no-contact* trials: 20%) (**Figure 3A**). The force sensor the participants tapped in the *contact* trials was now attached to a platform that could be automatically retracted. In the *contact* trials, participants tapped the force sensor to trigger the *test* force identically to Experiments 1 and 2, but in the *no-contact* trials, the platform automatically retracted before trial onset, unbeknownst to the participants, leading them to unexpectedly miss the contact with the sensor but still trigger the *test* force only by the position of their right index finger. Participants' vision was occluded so that they could not know whether the next trial would be a *contact* or a *no-contact* trial. According to the attenuation model, providing a bimanual sensorimotor context in 80% of the trials should lead participants to form predictions about the somatosensory consequences of their movement in most of the trials and thus attenuate the received touch on their passive left hand compared to the *baseline*, even if the touch of their active hand was unexpectedly missed (**Figure 3B**). In contrast, if attenuation is a nonpredictive process caused by the mere simultaneous tactile

stimulation of the active finger, no attenuation effects in the *no-contact* trials should be
observed with respect to the *baseline* (**Figure 3C**).

The results showed a robust attenuation in both *contact* and *no-contact* trials with respect
to the *baseline* condition, regardless of whether contact was made (**Figure 3D, E, F &**
H): the PSEs were significantly lower in the *contact* condition than in the *baseline*
condition ($t(29) = 8.06$, $p < 0.001$, $d = 1.47$, $CI^{95} = [0.16\ 0.27]$) and lower in the *no-*
contact condition than in the *baseline* condition ($t(29) = 4.45$, $p < 0.001$, $d = 0.81$, $CI^{95} =$
$[0.07\ 0.20]$). The magnitude of attenuation was larger in the *contact* condition than in the
no-contact condition, with significantly lower PSEs in the *contact* condition than in the
no-contact condition ($t(29) = 3.94$, $p < 0.001$, $d = 0.72$, $CI^{95} = [0.04\ 0.12]$) (**Figure 3G)**
(see also **Supplementary Material** for individual fits and additional analyses). The
difference in the magnitude of attenuation was modest ($\cong 35\%$). These findings indicate
that the perceived intensity of touch on the passive left hand was significantly attenuated
both when the active hand received touch or not, thereby ruling out the possibility that
attenuation is merely due to simultaneous tactile stimulation [45,46].

The results of Experiment 3 emphasise the predictive nature of somatosensory
attenuation, which is observed only when the sensorimotor context allows the formation
of such predictions. In Experiments 1 and 2 attenuation was only observed in the *contact*
conditions, whereas attenuation was also observed in the *no-contact* condition of
Experiment 3, where participants expected that they would make contact (i.e., the
sensorimotor context was conducive to predicting self-touch). Thus, larger differences
between the PSEs in the *contact* and *no-contact* conditions were observed for
Experiments 1 and 2 compared to Experiment 3. To further strengthen our interpretation,
we performed an ANOVA on the difference in the PSEs between the *contact* and *no-*
contact conditions across all three experiments. The ANOVA revealed a significant main
effect of Experiment ($F(2, 87) = 8.05$, $p<0.001$, $\eta p2 = 0.156$), with Bonferroni corrected
post hoc comparisons indicating significant differences between Experiments 1 and 3
($t(2, 87) = 3.10$, $p = 0.008$, $d = 0.80$, CI $[0.03\ 0.23]$) as well as Experiments 2 and 3 ($t(2,$
$87) = 3.76$, $p < 0.001$, $d = 0.97$, CI $[0.06\ 0.26]$), but no significant differences between
Experiments 1 and 2 ($t(2, 87) = -0.66$, $p = 1.000$, $d = -0.17$, CI $[-0.13\ 0.07]$, $BF_{01} =$
3.308). Thus, *no-contact* trials elicited attenuated perception only when the sensorimotor
context allowed for a prediction of self-touch (Experiment 3) and not when the action
was only arbitrarily associated with the touch (Experiments 1 and 2).

Figure 3. Experimental methods, hypotheses, and results of Experiment 3. A. The *contact* (magenta) and *no-contact* (green) conditions were identical to those of Experiment 1, with the only difference being their relative proportion (*contact* trials 80%, *no-contact* trials 20%). In the *no-contact* trials, the force sensor was automatically retracted, unbeknownst to the participant, revealing the distance sensor placed below. The *baseline* condition (blue) was identical to that of Experiments 1 and 2. **B.** Hypotheses based on the attenuation model. If attenuation is due to action prediction, then the perceived magnitude of touch should be reduced in both the *contact* (magenta) and *no-contact* conditions (green) compared to the baseline (blue). **C.** Hypotheses based on the enhancement model. If attenuation effects are driven by simultaneous touch on the active hand, then the perceived magnitude of touch should be reduced only in the *contact* condition (magenta) compared to the *baseline* (blue) and not in the *no-contact* condition (green) which should be similar to the *baseline* (blue). **D-H.** Data are colour-coded per condition. **D.** Box plots show the median and interquartile ranges for the PSE values per condition, black circles and error bars show the mean PSE ± s.e.m, and the raincloud plots show the individual PSE values and their distributions. **E.** Group psychometric functions for each condition. The leftward shift of the curves in the *contact* and *no-contact* conditions indicates attenuated PSE values compared to the *baseline*. **F.** Line plots for individual participant PSE values illustrating significantly lower PSEs for the *contact* versus *baseline* condition, **G.** significantly lower PSEs in the *contact* versus *no-contact* condition and **H.** significantly lower PSEs in the *no-contact* versus *baseline*. (***$p < 0.001$)

Discussion

In the present study, two opposing hypotheses regarding how action influences somatosensory perception were contrasted: the attenuation model and the enhancement model. Our findings demonstrate that action does not enhance (Experiments 1 and 2) but attenuates the predicted touch (Experiment 3). Before discussing the findings, it is important to emphasise that to draw conclusions about whether perception is attenuated or enhanced in an experimental condition including action, it is necessary to include a baseline condition without action. Only comparing conditions that include action prevents differentiating a genuine effect of enhancement (or attenuation) from an effect of reduced attenuation (or enhancement) in one of the two conditions. This also applies to experimental manipulations that contrast predicted with unpredicted somatosensory stimuli during action (see [45] for such comparisons). To this end, a *baseline* condition of pure somatosensory exafference (*i.e.*, externally generated touch) was included in all the present experiments that allowed us to distinguish the direction of the effects.

The results of Experiment 1 showed a robust attenuation of the touch applied on the passive left hand when the two hands made contact (*contact* condition), but neither attenuation nor enhancement was caused by the mere movement of the right hand (*no-contact* condition). In contrast, the perceived intensity of touch in the *no-contact* condition was similar to that of the *baseline* (*i.e.*, externally generated touch). These findings are in line with the results of Bays et al. [32], who found no change in the participants' somatosensory perception when the two hands did not make contact, but do not replicate those of Thomas et al. [45], who observed enhancement of touch in a *no-contact* condition. Experiment 2 further investigated the source of the previously reported enhancement effects and showed that they are in fact driven by the baseline condition

used: enhancement was observed only relative to a condition in which the touch was
applied rapidly following a cue to inhibit the movement (*i.e.*, a "do not move" cue), as in
previous research [45], but not relative to an externally generated touch condition (our
baseline). In support of this claim, Experiment 2 showed that a cue to inhibit the
movement results in a significant reduction in the perceived intensity of the imperative
stimulus (*i.e.*, touch on the passive hand) compared to baseline perception. This is in line
with previous evidence showing reduced amplitude of somatosensory evoked potentials
for tactile stimuli presented shortly following a cue to inhibit a movement [62], reduced
perceived amplitude for tactile stimuli under the mere expectation to move [60,66], and
reduced encoding of sensory stimuli following motor inhibition [63–65]. Therefore, rather
than participants' perception being enhanced in the *no-contact* condition, it is a reduction
of the perceived intensity following the cue to inhibit the movement that leads to an
apparent enhancement. In contrast, by including a novel externally generated touch
condition as a *baseline* that involved neither motor planning nor response inhibition, it
became clear that there were no enhancement effects.

Experiment 3 demonstrated that action attenuates the predicted touch even if the active
hand does not receive simultaneous tactile stimulation with the passive hand. When
participants simply moved their right hand to trigger the touch on their left hand (*no-
contact* trials in Experiments 1 & 2), no change in their somatosensory perception was
found. This suggests that an arbitrary mapping between the movement of the right hand
and the delivery of touch on the left hand is insufficient to elicit attenuation. In contrast,
when participants expected to touch their own hand (Experiment 3), significant
attenuation was observed, even when the active hand unexpectedly missed the contact
(*no-contact* trials, 20%). This pattern of results can be explained by the attenuation
model, which suggests that self-generation of a stimulus is in itself not sufficient for
predictive attenuation, but that the action and it's predicted sensory feedback are
consistent with touching one's own body [59]. Thus, a sensorimotor context that closely
resembles tapping directly on the left index finger with the right (self-touch) was critical
for predictive attenuation [32]. The findings of Experiment 3 also contradict the suggestion
that attenuation on a passive hand reflects a nonpredictive gating of tactile input during
movement of the active hand [45,46]. Indeed, several earlier studies showed gating effects
only on the moving limb (movement effector) and not on the passive limb [67–72], and we
have also recently shown that experimental paradigms identical to the one used in the
present study produce attenuation effects but not gating effects on the passive hand [27].
Overall, the results suggest that attenuation effects are driven by action prediction and not
the simultaneous tactile stimulation of the two hands. From an ecological point of view,
in every self-touch behaviour, we necessarily receive somatosensory input on the active
hand and the body part that passively receives touch ("touchant-touché" [73]), and it is
within these sensorimotor contexts that the brain forms predictions about the
somatosensory consequences on multiple body parts [59].

Interestingly, the magnitude of attenuation was greater in the *contact* compared to the *no-
contact* condition by approximately 35% in Experiment 3. This could suggest that the
attenuation effect observed in the *contact* condition arises from multiple sources, one
being the motor-based prediction and another source being the simultaneous touch on the

active hand. However this interpretation is inconsistent with the findings that somatosensory attenuation has been previously observed in the absence of simultaneous tactile stimulation, for example, when imagining but not executing the right hand movement [38] or just before the hands make contact [26]. Similarly, no somatosensory attenuation is observed in the absence of action prediction, even if the two hands received simultaneous tactile stimulation; for example, the passive displacement of the right hand towards the left hand that is accompanied by double touch [35] or the mere delivery of simultaneous tactile stimulation on both hands [26] does not produce attenuation. Instead, our finding that the magnitude of attenuation was greater in the *contact* than the *no-contact* condition in Experiment 3 may instead reflect a decrease of attenuation in the *no-contact* trials, due to the unexpected omission of contact influencing the perceived magnitude of touch in a postdictive manner. Specifically, the unexpected omission of contact on the minority of trials (20%) could be seen as a form of stimulus omission akin to so called 'silent oddballs' that are known to generate prediction errors [74–77]. Although participants clearly attenuated the touch predicted by their movement in the *no-contact* trials, their expectation of contact was necessarily violated. This violation could be considered a novel event, given its infrequency. Novel events can have several consequences for cognition, including transient enhancements of perception[78], facilitated encoding of information into working memory [79] as well as changes in the allocation of attentional resources in a postdictive manner (for a review of effects of novelty on cognition see[80].

Some authors have criticised the comparison of perceived intensity of sensory stimuli in self-generated and externally generated conditions [15,81,82]. A frequent criticism is that tactile input may be "predicted" in self-generated conditions through action and "unpredicted" in externally generated (passive) conditions that do not involve action. This concern can be ruled out since in all three experiments, the stimulus in the *baseline (i.e.,* externally generated touch) condition was delivered at a fixed timepoint (800 ms after the cue); therefore, participants could predict it in the absence of motor-based predictions. Secondly, it has been argued that somatosensory attenuation findings may be driven by dual-task requirements [15] present only in self-generated conditions that could increase the working memory load or result in a shift of attention towards the active hand. A similar criticism has been suggested in which the mere movement of the active hand in self-generated conditions can result in a nonpredictive gating of tactile input. However these explanations can also be ruled out, given that dual-task requirements and movement of the active hand were present in the *contact* and *no-contact* conditions of all three experiments without concomitant attenuation effects. Indeed if dual-task requirements, or the mere presence of movement during the perceptual task, was sufficient to cause attenuation effects, then attenuation would have been observed in the *no-contact* conditions of Experiments 1 and 2. Finally, alternative explanations based on differences in other psychophysical parameters, movement kinematics or timings between experiments can also be ruled out (see **Supplementary Material** for additional analyses).

How can the findings that action prediction attenuates touch be reconciled with those showing that expectations outside the domain of action improve sensory perception [15]? While it is difficult to directly compare these lines of research because of differences in

the sensory modality investigated, the task designs used and the perceptual measures employed, we speculate that there are numerous possible reasons why action prediction may not have the same effect on perception as prediction mechanisms outside the domain of action. First, research on action-based predictions concerns ubiquitous associations between actions and their sensory consequences that we are continually exposed to throughout the lifespan. For example, we are constantly exposed to associations between our motor behaviours and their tactile consequences during self-touch, even as early as 13 weeks *in utero* [83]. In contrast, research on sensory expectations outside the domain of action primarily concerns arbitrary associations between stimuli that are typically learned only during the time course of a given task. It is therefore conceivable that separable mechanisms may operate to predict action effects versus stimulus-stimulus associations (see [84,85] for discussion). Second, higher-level expectations, such as explicit prior knowledge that a specific sensory event is likely, might not operate in the same way as lower-level action predictions; for example, it has been proposed that action-based predictions inhibit expected stimuli, while sensory expectations potentiate the expected sensory inputs [86]. Most importantly, from a theoretical perspective, attenuating the predicted sensory consequences of actions does not necessarily mean that the brain forms inaccurate representations of the world but instead indicates a flexible strategy that prioritises more informative externally generated events.

Debates between researchers supporting attenuation or enhancement are useful and fruitful for scientific dialogue and advancement. The present study revisited recent findings on somatosensory enhancement during action and showed that when the requirement to inhibit the action is removed from the baseline, action predictions do not enhance but attenuate the received somatosensory input. Our results are in strong alignment with animal studies showing that action attenuates the predicted sensory consequences (for reviews, see [87–91]). For example, crickets and mice suppress auditory reafferent signals but maintain their sensitivity to external sounds [92–95], the weakly electric fish attenuates its electrosensory reafference to respond to externally generated electrical signals [89,96], and primates attenuate their vestibular reafference and activate vestibular-related reflexes only when the vestibular input is exafferent [97–99]. To this end, the results of the present study prompt a reappraisal of recent experimental findings upon which theoretical frameworks proposing a perceptual enhancement by action prediction are based.

Methods

Participants

Thirty naive adults participated in Experiment 1 (18 female, aged 18-36, 28 right-handed and 2 ambidextrous), thirty naive adults participated in Experiment 2 (17 female, aged 20-37, 29 right-handed and 1 left-handed) and thirty naive adults participated in Experiment 3 (12 female, aged 21-40, 29 right-handed, 1 ambidextrous, 1 left-handed). Current or history of psychological or neurological conditions, as well as the use of any psychoactive drugs or medication to treat such conditions, were criteria for exclusion. The sample size of each Experiment was decided prior to data collection based on our

previous studies using the same task [27,34]. Handedness was assessed using the Edinburgh Handedness Inventory [100]. All experiments were approved by the Swedish Ethical Review Authority (registration no. 2021-03790). Participants provided written informed consent and were compensated for their time.

Pre-registration of experiments

The method and analysis plan for each experiment was pre-registered on the Open Science Framework (OSF) prior to data collection (Experiment 1 https://osf.io/9jkqt, Experiment 2 https://osf.io/hs8au, Experiment 3 https://osf.io/gkwu7).

Modifications/additions to the pre-registered plan

The only modification from the pre-registered plan was a reduction in the number of trials administered in Experiment 3. Following initial piloting, it was deemed appropriate to reduce the overall length of the task by including fewer trials (see *Methods Details)*. Two additional ANOVAs were also conducted to compare the results across the three experiments (**Main Text***)* and the kinematics across experiments (Supplementary Materials).

Experimental setup

Participants sat comfortably on a chair with their arms placed on a table. The left hand rested palm up, with the index finger placed on a molded support. On each trial, a motor (Maxon EC Motor EC 90 flat; Switzerland) delivered two forces (the *test* force and the *comparison* force) on the pulp of the left index finger through a cylindrical probe (25 mm height) with a flat aluminum surface (20 mm diameter) attached to a lever on the motor. A force sensor (FSG15N1A, Honeywell Inc.; diameter, 5 mm; minimum resolution, 0.01 N; response time, 1 ms; measurement range, 0–15 N) within the probe recorded the forces applied on the left index finger. Following the presentation of the two forces, participants were asked to verbally report which of the two forces felt stronger – the first or the second. A second identical force sensor within an identical cylindrical probe ("active force sensor") was placed on top of, but not in contact with, the probe of the left index finger.

Force-discrimination task

Participants judged the intensity of a *test* force and a *comparison* force (100 ms duration each) separated by a random interval between 800ms and 1200ms in a two-alternative forced choice (2AFC) task. The intensity of the test force was 2 N, while the intensity of the comparison force was systematically varied among seven force levels (1, 1.5, 1.75, 2, 2.25, 2.5 or 3 N). In all the conditions, the forces were delivered by the same motor, in order to precisely control their magnitude, however the source of the force was manipulated across conditions such that the force was triggered by the participants contact with a force sensor (*contact* condition), their finger movement (*no-contact* condition) or automatically by the stimulus computer (*baseline* condition).

Experimental design and procedures

Experiment 1

There were three conditions in Experiment 1 presented in three blocks separately: the *baseline* (i.e. externally generated touch) condition, the *contact* condition, and the *no-contact* condition.

In the *baseline* condition, participants did not move their limbs but passively received a *test* and the *comparison* force on the left index finger. This *baseline* condition was used to assess the participants' somatosensory perception in the absence of any movement [26,34,35]. Each trial began with an auditory cue (100 ms duration, 997 Hz) followed by the *test* force delivered to the participants' left index finger 800 ms after the cue by the motor. The *comparison* force was then delivered at a random interval between 800 ms and 1200 ms after the *test* force.

In the *contact* condition, participants started each trial by holding their right index finger approximately 10 cm above their left index finger. The start position was marked with a visual marker placed next to their right index finger while the final position was the probe of the active force sensor. The same auditory cue was presented as in the *baseline* condition, but participants now moved their right index finger downwards towards their left index finger and actively tapped on the active force sensor placed on top of, but not in contact with, the probe. The participant's active tap on the force sensor triggered the motor to apply the *test* force on their left index finger (threshold 0.2 N). The tap of their right index finger triggered the test force on their left index finger with an intrinsic delay of 36 ms. The *comparison* force was then delivered at a random interval between 800ms and 1200ms after the *test* force. As in our previous studies[25,27,39], participants were asked to tap, neither too weakly nor too strongly, with their right index finger, "as if tapping the screen of their smartphone". This instruction was provided to ensure that the relationship between the force they applied with their right index finger on the force sensor and the force they received on their left index finger by the motor (2 N) remained approximately constant throughout the experiment, thereby establishing perceived causality[14,101]. Indeed, it has been previously shown that the magnitude of attenuation remains unaffected by halving or doubling the gain between the force applied by the active finger and the force delivered on the passive hand[14]. This should not be surprising given that when one finger transmits a force through an object to another finger, the resulting force depends also on the object properties (e.g., shape, material, contact area) and the angle at which the finger contacts the object.

In the *no-contact* condition, the participants started each trial by holding their right index finger 10 cm above their left index finger, identically to the *contact* condition. For this block, the active force sensor was removed and replaced by a distance sensor that detected the position of their right index finger. The distance sensor was placed on top of, but not in contact with, the probe. Following the same auditory cue, participants moved their right index finger towards their left index finger. To restrict the participants' right index finger movement within similar movement ranges as in the *contact* condition and

avoid contact with the distance sensor placed underneath, a second visual marker indicated the final position of the right index finger above the distance sensor (5 cm from the marker indicating the initial position). The distance sensor was connected to an Arduino microcontroller that controlled a servo motor. The servo motor and the force sensor were placed 2 meters away from the participants' hands and hidden from view. Once the distance sensor detected that the position of the right index finger became smaller than a preset threshold, it triggered the servo motor that hit the force sensor and triggered the *test* force. We accounted for the additional delay of the distance sensor by setting the position threshold for the distance sensor slightly higher than the lowest position the participants were asked to reach with their right index finger. Therefore, the test force would be delivered at a similar timing to the participants' right index finger endpoint between the *contact* and *no-contact conditions*. This position threshold was set based on significant pilot testing prior all experiments. Indeed, there was a minimal time difference between the two setups across experiments (17 ms average delay) with the *no-contact* condition leading, rather than lagging the *contact* one (see **Supplementary Materials**). The *comparison* force was delivered at a random interval between 800ms and 1200ms after the *test* force. Therefore, in the *no-contact* condition, there was no touch on the right index finger simultaneously with the *test* force on their left index finger.

Before the experiment, participants were trained to make similar movements with their right index finger in the *contact* and *no-contact* conditions, emphasis was placed on restricting their right index finger movements to between the two visual markers in the *no-contact* condition. The 3D position of the right index finger was recorded using a Polhemus Liberty tracker (240 Hz). Kinematic information was used to compare the movements between the *contact* and *no-contact* conditions and to reject any trials in which the participant did not trigger the test stimulus with their movement, or trials in which the participant did not move as instructed in the training (see below). Participants were administered white noise through headphones to mask any sounds made by the motor to serve as a cue for the task. The loudness of the white noise was adjusted such that participants could clearly hear the auditory tones of the trial.

Each block consisted of 70 trials resulting to 210 trials per participant. Thus, the proportion of *contact* and *no-contact* trials was the same (50-50%). The order of the conditions was fully counterbalanced across participants.

Experiment 2

The 2-AFC task was identical to that of Experiment 1. In Experiment 2, there were five conditions in total: the *baseline* condition, the *contact* condition, the *no-contact* condition, and two additional NOGO conditions (*NOGO contact* and *NOGO no-contact*). Trials of the NOGO conditions were pseudo-randomly intermixed with trials of the *contact* and *no-contact* conditions respectively (GO trials). The five conditions were presented in three separate blocks: the *baseline*, *contact* (GO and NOGO trials) and *no-contact* (GO and NOGO trials) blocks.

In the *contact* block, 50% of trials began with an auditory "GO" cue (100 ms duration
high tone of 2458 Hz) instructing participants to tap the active force sensor to trigger the
test force (identically to the *contact* condition of Experiment 1). On the remaining 50% of
trials an auditory "NOGO" cue (100ms duration low tone of 222 Hz) instructed
participants to withhold their movement and the *test* force was then delivered 800ms
following the NOGO cue. The *no-contact* block was identical to the *contact* block, except
that the *test* force was triggered by the position of the right index finger without contact
with the force sensor (identically to the *no-contact* condition of Experiment 1). The cue
tone in the *baseline* block was the same as the cue tone in the NOGO trials (100 ms
duration low tone of 222 Hz).

Therefore, this design replicated the experimental design of Thomas et al. (Thomas et al.,
2021) with the additional inclusion of the *baseline* (i.e. externally generated touch)
condition. As in Experiment 1, we recorded the 3D position of the right index finger and
the registered kinematic information was used to compare the movements between the
contact and *no-contact* conditions and reject any trials in which the participant did not
trigger the *test* stimulus with their movement, trials in which the participant did not move
as instructed in the training (see below), and trials in which the participants moved while
instructed not to do so by the auditory cues (NOGO trials). As in Experiment 1,
participants were administered white noise and the loudness of the white noise was
adjusted such that participants could clearly hear the GO and NOGO auditory cues of the
trial.

The *baseline* block consisted of 70 trials, and the *contact* and *no-contact* blocks consisted
of 70 GO trials and 70 NOGO trials each, resulting to 350 trials in total per participant.
The order of the blocks was fully counterbalanced across participants.

<u>Experiment 3</u>

The 2-AFC task and the experimental conditions (*baseline*, *contact*, and *no-contact*) were
identical to those of Experiment 1, except for the following. *Contact* trials (80%) were
now pseudo-randomly intermixed with *no-contact* trials (20%) within the same block.
The force sensor was now attached to a plastic platform that could be automatically
retracted by a servo motor, depending on the trial type. Upon retraction, a distance sensor
placed underneath the platform was revealed. In the *contact* trials, participants tapped the
force sensor to trigger the *test* force identically to Experiments 1 and 2. In the *no-contact*
trials, the platform was automatically retracted before trial onset, unbeknownst to the
participant. This led participants to unexpectedly miss the active force sensor and instead
trigger the *test* force only by the position of their right index finger. In all conditions, the
participants' vision was occluded, and white noise was administered via headphones to
prevent any visual or auditory cues indicating that the force sensor had been retracted in
no-contact trials. The *baseline* block was identical to that of Experiments 1and 2.

In Experiment 3, the number of trials was reduced from 70 to 56 to shorten the total
experiment time to less than 90 minutes, similar in the Experiment 2. Thus, there were 56
no-contact trials (20%) and 224 contact trials (80%). Similarly, there were 56 trials in the

baseline condition. This resulted to 336 trials per participant. The order of the two blocks
was fully counterbalanced across participants.

Preprocessing of psychophysical trials.

Experiment 1

Experiment 1 included 6300 trials in total (30 participants * 70 trials * 3 conditions).
Twenty-nine (29) trials were excluded (0.5%) because of 1 missing response, 4 trials in
which the force was not applied correctly (1.85 N < *test* force < 2.15 N), and 24 trials
because the *test* force was not triggered when moving towards the distance sensor. All
trials excluded by the psychophysical fits were also excluded in the kinematic analysis.

Experiment 2

Experiment 2 included 10500 trials in total (30 participants * 70 trials * 5 conditions).
One-hundred fourteen (114) trials were excluded (1.1%) because of 2 missing responses,
6 trials in which the force was not applied correctly (1.85 N < *test* force < 2.15 N), 44
trials because the *test* force was not triggered when moving towards the distance sensor
and 62 trials because the finger moved on a NOGO trial. All trials excluded by the
psychophysical fits were also excluded in the kinematic analysis.

Experiment 3

Experiment 3 included 10080 trials in total (30 participants * 224 contact trials + 56 no-
contact trials + 56 baseline trials). Two-hundred twenty-four (29) trials were excluded
(2.42%) because of 29 missing responses, 86 trials in which the force was not applied
correctly (1.85 N < *test* force < 2.15 N), 81 because the test force was not triggered when
moving towards the distance sensor and 48 because the finger contacted the distance
sensor on *no-contact* trials. All trials excluded by the psychophysical fits were also
excluded in the kinematic analysis.

Preprocessing of kinematic recordings.

All kinematic trials were co-registered with the force trials through Transistor-Transistor
Logic (TTL) signals sent by the motor to both file outputs. The kinematic recordings
were corrected for any distortion in the Polhemus sensor from the force sensor and
distance sensor based on measurements made with and without the force/distance sensors
within the same space. Exclusion of trials based on the kinematics was done by assessing
whether the test force was delivered after the position of the active finger reached its
minimum value on the vertical plane. Trials in which the force was delivered after the
minimum value had been reached were rejected. For experiment 2, trials in which the
active finger moved more than 1 cm following a NOGO cue were also rejected.

Fitting of psychophysical responses

723 For each experiment and each condition, the participants' responses were fitted with a
724 generalized linear model using a *logit* link function (Equation 1):

$$p = \frac{e^{\beta_0 + \beta_1 x}}{1 + e^{\beta_0 + \beta_1 x}} \text{ (Equation 1)}$$

We extracted two parameters of interest: the Point of Subjective Equality (PSE) ($PSE = -\frac{\beta_0}{\beta_1}$), which represents the intensity at which the *test* force felt as strong as the *comparison* force ($p = 0.5$) and quantifies the perceived intensity, and the JND ($JND = \frac{\log(3)}{\beta_1}$), which reflects the participants' discrimination ability. Before fitting the responses, the values of the applied comparison *forces* were binned to the closest value with respect to their theoretical values (1, 1.5, 1.75, 2, 2.25, 2.5 or 3 N).

For all participants and all conditions, the fitted logistic models were very good, with McFadden's R squared measures ranging between 0.602 and 0.979.

Normality of data and statistical comparisons.

We used R [102], JASP [103] and MATLAB (2020b) to analyze our data. The normality of the PSE and the JND data distributions, as well as the kinematic information data distributions were assessed using Shapiro-Wilk tests. Depending on the data normality, pairwise comparisons between conditions were performed by using either a paired t-test or a Wilcoxon signed-rank test. We report 95% confidence intervals (CI^{95}) for each statistical test. Effect sizes are reported as the Cohen's *d* for t-tests or the matched rank biserial correlation *rrb* for the Wilcoxon signed-rank tests. In addition, a Bayesian factor analysis using default Cauchy priors with a scale of 0.707 was performed for all statistical tests that led to not statistically significant effects, to provide information about the level of support for the null hypothesis compared to the alternative hypothesis (BF_{01}) based on the data. We interpret a factor between 1/3 and 3 as "anecdotal evidence"[104,105], indicating that support for either the preferred or null hypotheses is insufficient. For the Analysis of Variance (ANOVA, see **Supplementary Material**), homogeneity of variance was assessed using Levene's test for Equality of Variances, which did not reach significance, and the Q-Q plot of the standardized residuals indicated approximately normally distributed residuals. Post-hoc tests were made using Bonferroni corrections. All tests were two-tailed.

Acknowledgments

We thank Evridiki Asimakidou and Lili Timar for their assistance during data collection. We also thank Henrik Ehrsson for his helpful comments on an earlier version of the manuscript. X.J. and K.K. were supported by the Swedish Research Council (VR Starting Grant 2019-01909 granted to K.K.). Experimental costs were covered by the same grant. X.J. is supported by the European Union's Horizon 2021 research and innovation programme under the Marie Skłodowska-Curie grant agreement No. 101059348.

Author Contributions

K.K. and X.J. designed the experiments; X.J. collected the data; X.J. analysed the data; X.J. and K.K. wrote the manuscript.

Declaration of interests

The authors declare no competing interests.

Data availability

Data (PSEs and JNDs) are publicly available along with preregistered analyses plans on the OSF (Experiment 1: https://osf.io/9jkqt, Experiment 2: https://osf.io/hs8au & Experiment 3: https://osf.io/gk).

Code availability

This paper does not report original code. Any additional information is available from the lead contact upon request.

References

1. Shadmehr, R., Krakauer, J. W., Neuroanatomy, A. C. & Motor, F. O. R. A computational neuroanatomy for motor control. *Exp. Brain Res.* **185**, 359–381 (2008). doi: 10.1007/s00221-008-1280-5.

2. McNamee, D. & Wolpert, D. M. Internal Models in Biological Control. *Annu. Rev. Control. Robot. Auton. Syst.* **2**, 339–364 (2019). doi: 10.1146/annurev-control-060117-105206.

3. Wolpert, D. M. & Flanagan, J. R. Motor prediction. *Curr. Biol.* **11**, R729–R732 (2001). doi: 10.1016/S0960-9822(01)00432-8.

4. Shadmehr, R., Smith, M. a & Krakauer, J. W. Error correction, sensory prediction, and adaptation in motor control. *Annu. Rev. Neurosci.* **33**, 89–108 (2010). doi: 10.1146/annurev-neuro-060909-153135.

5. Todorov, E. & Jordan, M. I. Optimal feedback control as a theory of motor coordination. *Nat. Neurosci.* **5**, 1226–1235 (2002). doi: 10.1038/nn963.

6. Scott, S. H. Optimal feedback control and the neural basis of volitional motor control. *Nat. Rev. Neurosci.* **5**, 532–544 (2004). doi: 10.1038/nrn1427.

7. Davidson, P. R. & Wolpert, D. M. Widespread access to predictive models in the motor system: A short review. *Journal of Neural Engineering* vol. 2 at https://doi.org/10.1088/1741-2560/2/3/S11 (2005). doi: 10.1088/1741-2560/2/3/S11.

8. Franklin, D. W. & Wolpert, D. M. Computational mechanisms of sensorimotor control. *Neuron* vol. 72 at https://doi.org/10.1016/j.neuron.2011.10.006 (2011).

doi: 10.1016/j.neuron.2011.10.006.

9.   Wolpert, D. M. & Kawato, M. Multiple paired forward and inverse models for
motor control. *Neural Networks* **11**, (1998). doi: 10.1016/S0893-6080(98)00066-5.

10. Brown, H., Adams, R. a., Parees, I., Edwards, M. & Friston, K. Active inference,
sensory attenuation and illusions. *Cogn. Process.* **14**, 411–427 (2013). doi:
10.1007/s10339-013-0571-3.

11. Corlett, P. Predicting to Perceive and Learning When to Learn. *Trends Cogn. Sci.*
24, 259–260 (2020). doi: 10.1016/j.tics.2019.12.005.

12. Idei, H., Ohata, W., Yamashita, Y., Ogata, T. & Tani, J. Emergence of sensory
attenuation based upon the free-energy principle. *Sci. Rep.* **12**, 14542 (2022). doi:
10.1038/s41598-022-18207-7.

13. Blakemore, S.-J., Wolpert, D. & Frith, C. Why can☐t you tickle yourself?
Neuroreport **11**, R11–R16 (2000). doi: 10.1097/00001756-200008030-00002.

14. Bays, P. M. & Wolpert, D. M. Predictive attenuation in the perception of touch. in
Sensorimotor Foundations of Higher Cognition (2012).
doi:10.1093/acprof:oso/9780199231447.003.0016. doi:
10.1093/acprof:oso/9780199231447.003.0016.

15. Press, C., Kok, P. & Yon, D. The Perceptual Prediction Paradox. *Trends in*
Cognitive Sciences vol. 24 at https://doi.org/10.1016/j.tics.2019.11.003 (2020).
doi: 10.1016/j.tics.2019.11.003.

16. Yon, D., Zainzinger, V., de Lange, F., Eimer, M. & Press, C. Action biases
perceptual decisions towards expected outcomes. *J. Exp. Psychol. Gen.* (2020)
doi:10.31234/osf.io/3zp8n. doi: 10.31234/osf.io/3zp8n.

17. Press, C. & Yon, D. Perceptual Prediction: Rapidly Making Sense of a Noisy
World. *Current Biology* vol. 29 at https://doi.org/10.1016/j.cub.2019.06.054
(2019). doi: 10.1016/j.cub.2019.06.054.

18. Wyart, V., Nobre, A. C. & Summerfield, C. Dissociable prior influences of signal
probability and relevance on visual contrast sensitivity. *Proc. Natl. Acad. Sci. U. S.*
A. **109**, (2012). doi: 10.1073/pnas.1120118109.

19. Chalk, M., Seitz, A. R. & Seriès, P. Rapidly learned stimulus expectations alter
perception of motion. *J. Vis.* **10**, (2010). doi: 10.1167/10.8.2.

20. Stein, T. & Peelen, M. V. Content-specific expectations enhance stimulus
detectability by increasing perceptual sensitivity. *J. Exp. Psychol. Gen.* **144**,
(2015). doi: 10.1037/xge0000109.

21. Teufel, C., Dakin, S. C. & Fletcher, P. C. Prior object-knowledge sharpens
properties of early visual feature-detectors. *Sci. Rep.* **8**, (2018). doi:

830   10.1038/s41598-018-28845-5.

831 22. Hudson, M., Nicholson, T., Simpson, W. A., Ellis, R. & Bach, P. One step ahead:
832   The perceived kinematics of others' actions are biased toward expected goals. *J.*
833   *Exp. Psychol. Gen.* **145**, (2016). doi: 10.1037/xge0000126.

834 23. Hudson, M., Nicholson, T., Ellis, R. & Bach, P. I see what you say: Prior
835   knowledge of other's goals automatically biases the perception of their actions.
836   *Cognition* **146**, (2016). doi: 10.1016/j.cognition.2015.09.021.

837 24. Summerfield, C. & de Lange, F. P. Expectation in perceptual decision making:
838   Neural and computational mechanisms. *Nature Reviews Neuroscience* vol. 15 at
839   https://doi.org/10.1038/nrn3838 (2014). doi: 10.1038/nrn3838.

840 25. Asimakidou, E., Job, X. & Kilteni, K. The positive dimension of schizotypy is
841   associated with a reduced attenuation and precision of self-generated touch. *npj*
842   *Schizophr.* (2022).

843 26. Bays, P. M., Wolpert, D. M. & Flanagan, J. R. Perception of the consequences of
844   self-action is temporally tuned and event driven. *Curr. Biol.* **15**, (2005). doi:
845   10.1016/j.cub.2005.05.023.

846 27. Kilteni, K. & Ehrsson, H. H. Predictive attenuation of touch and tactile gating are
847   distinct perceptual phenomena. *iScience* **25**, (2022). doi:
848   10.1016/j.isci.2022.104077.

849 28. Knoetsch, F. & Zimmermann, E. The spatial specificity of sensory attenuation for
850   self-touch. *Conscious. Cogn.* **92**, 103135 (2021). doi:
851   10.1016/j.concog.2021.103135.

852 29. Wolpe, N. *et al.* Ageing increases reliance on sensorimotor prediction through
853   structural and functional differences in frontostriatal circuits. *Nat. Commun.* **7**,
854   13034 (2016). doi: 10.1038/ncomms13034.

855 30. Weiskrantz, L., Elliott, J. & Darlington, C. Preliminary observations on tickling
856   oneself. *Nature* **230**, 598–599 (1971). doi: 10.1038/230598a0.

857 31. McNaughton, D. *et al.* Methodological considerations for the force-matching task.
858   *Behav. Res. Methods* (2022) doi:10.3758/s13428-022-01954-w. doi:
859   10.3758/s13428-022-01954-w.

860 32. Bays, P. M., Flanagan, J. R. & Wolpert, D. M. Attenuation of self-generated tactile
861   sensations is predictive, not postdictive. *PLoS Biol.* **4**, (2006). doi:
862   10.1371/journal.pbio.0040028.

863 33. Blakemore, S. J., Frith, C. D. & Wolpert, D. M. Spatio-temporal prediction
864   modulates the perception of self-produced stimuli. *J. Cogn. Neurosci.* **11**, (1999).
865   doi: 10.1162/089892999563607.

34. Kilteni, K., Houborg, C. & Ehrsson, H. H. Rapid learning and unlearning of
predicted sensory delays in self-generated touch. *Elife* **8**, (2019). doi:
10.7554/eLife.42888.

35. Kilteni, K., Engeler, P. & Ehrsson, H. H. Efference Copy Is Necessary for the
Attenuation of Self-Generated Touch. *iScience* **23**, (2020). doi:
10.1016/j.isci.2020.100843.

36. Kilteni, K. & Ehrsson, H. H. Sensorimotor predictions and tool use: Hand-held
tools attenuate self-touch. *Cognition* **165**, 1–9 (2017). doi:
10.1016/j.cognition.2017.04.005.

37. Kilteni, K. & Ehrsson, H. H. Body ownership determines the attenuation of self-
generated tactile sensations. *Proc. Natl. Acad. Sci.* **114**, 8426–8431 (2017). doi:
10.1073/pnas.1703347114.

38. Kilteni, K., Andersson, B. J., Houborg, C. & Ehrsson, H. H. Motor imagery
involves predicting the sensory consequences of the imagined movement. *Nat.*
Commun. **9**, 1617 (2018). doi: 10.1038/s41467-018-03989-0.

39. Kilteni, K., Engeler, P., Boberg, I., Maurex, L. & Ehrsson, H. H. No evidence for
somatosensory attenuation during action observation of self-touch. *Eur. J.*
Neurosci. **54**, 6422–6444 (2021). doi: 10.1111/ejn.15436.

40. Blakemore, S. J., Wolpert, D. M. & Frith, C. D. Central cancellation of self-
produced tickle sensation. *Nat. Neurosci.* **1**, (1998). doi: 10.1038/2870.

41. Hesse, M. D., Nishitani, N., Fink, G. R., Jousmäki, V. & Hari, R. Attenuation of
somatosensory responses to self-produced tactile stimulation. *Cereb. Cortex* **20**,
425–432 (2010). doi: 10.1093/cercor/bhp110.

42. Kilteni, K. & Ehrsson, H. H. Functional Connectivity between the Cerebellum and
Somatosensory Areas Implements the Attenuation of Self-Generated Touch. *J.*
Neurosci. **40**, 894–906 (2020). doi: 10.1523/JNEUROSCI.1732-19.2019.

43. Shergill, S. S. *et al.* Modulation of somatosensory processing by action.
Neuroimage **70**, 356–362 (2013). doi: 10.1016/j.neuroimage.2012.12.043.

44. Shergill, S. S. *et al.* Functional magnetic resonance imaging of impaired sensory
prediction in schizophrenia. *JAMA Psychiatry* **71**, (2014). doi:
10.1001/jamapsychiatry.2013.2974.

45. Thomas, E. R., Yon, D., de Lange, F. P. & Press, C. Action Enhances Predicted
Touch. *Psychol. Sci.* **33**, (2022). doi: 10.1177/09567976211017505.

46. Press, C., Thomas, E. & Yon, D. Cancelling cancellation? Sensorimotor control,
agency, and prediction. *PsyArXiv* **195**, 2459–2482 (2022).

47. Führer, E., Voudouris, D., Lezkan, A., Drewing, K. & Fiehler, K. Tactile

suppression stems from sensation-specific sensorimotor predictions.
Proc.Natl.Acad.Sci U.S.A 2021.07.04.451060 (2022)
doi:10.1073/pnas.2118445119/-/DCSupplemental.Published. doi:
10.1073/pnas.2118445119/-/DCSupplemental.Published.

48.   Blakemore, S.-J., Smith, J., Steel, R., Johnstone, E. C. & Frith, C. D. The
perception of self-produced sensory stimuli in patients with auditory hallucinations
and passivity experiences: evidence for a breakdown in self-monitoring. *Psychol.*
Med. **30**, 1131–1139 (2000).

49.   Corlett, P. R. *et al.* Hallucinations and Strong Priors. *Trends in Cognitive Sciences*
vol. 23 at https://doi.org/10.1016/j.tics.2018.12.001 (2019). doi:
10.1016/j.tics.2018.12.001.

50.   Frith, C. The neural basis of hallucinations and delusions. *C. R. Biol.* **328**, 169–75
(2005). doi: 10.1016/j.crvi.2004.10.012.

51.   Frith, C. The self in action: Lessons from delusions of control. *Conscious. Cogn.*
14, (2005). doi: 10.1016/j.concog.2005.04.002.

52.   Frith, C. D., Blakemore, S. J. & Wolpert, D. M. Explaining the symptoms of
schizophrenia: Abnormalities in the awareness of action. *Brain Res. Rev.* **31**, 357–
363 (2000). doi: 10.1016/S0165-0173(99)00052-1.

53.   Frith, C. D. Can a Problem With Corollary Discharge Explain the Symptoms of
Schizophrenia? *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging*
vol. 4 at https://doi.org/10.1016/j.bpsc.2019.07.003 (2019). doi:
10.1016/j.bpsc.2019.07.003.

54.   Shergill, S. S., Samson, G., Bays, P. M., Frith, C. D. & Wolpert, D. M. Evidence
for sensory prediction deficits in schizophrenia. *Am. J. Psychiatry* **162**, 2384–2386
(2005). doi: 10.1176/appi.ajp.162.12.2384.

55.   Parees, I. *et al.* Loss of sensory attenuation in patients with functional
(psychogenic) movement disorders. *Brain* **137**, 2916–2921 (2014). doi:
10.1093/brain/awu237.

56.   Wolpe, N. *et al.* Sensory attenuation in Parkinson's disease is related to disease
severity and dopamine dose. *Sci. Rep.* **8**, 15643 (2018). doi: 10.1038/s41598-018-
33678-3.

57.   Kilteni, K., Houborg, C. & Ehrsson, H. H. Brief temporal perturbations in
somatosensory reafference disrupt perceptual and neural attenuation and increase
supplementary motor-cerebellar connectivity. *bioRxiv* 2022.11.25.517892 (2022)
doi:10.1101/2022.11.25.517892. doi: 10.1101/2022.11.25.517892.

58.   Shergill, S. S., Bays, P. M., Frith, C. D. & Wolpert, D. M. Two eyes for an eye: the
neuroscience of force escalation. *Science* **301**, 187 (2003). doi:

10.1126/science.1085327.

59.   Blakemore, S. J., Goodbody, S. J. & Wolpert, D. M. Predicting the consequences
of our own actions: The role of sensorimotor context estimation. *J. Neurosci.* **18**,
7511–7518 (1998). doi: 10.1523/jneurosci.18-18-07511.1998.

60.   Voss, M., Ingram, J. N., Wolpert, D. M. & Haggard, P. Mere expectation to move
causes attenuation of sensory signals. *PLoS One* **3**, (2008). doi:
10.1371/journal.pone.0002866.

61.   Walsh, E. & Haggard, P. The internal structure of stopping as revealed by a
sensory detection task. *Exp. Brain Res.* **183**, 405–410 (2007). doi: 10.1007/s00221-
007-1128-4.

62.   Hoshiyama, M. & Sheean, G. Changes of somatosensory evoked potentials
preceding rapid voluntary movement in Go/No-go choice reaction time task. *Cogn.*
Brain Res. **7**, (1998). doi: 10.1016/S0926-6410(98)00018-4.

63.   Chiu, Y. C. & Egner, T. Inhibition-induced forgetting results from resource
competition between response inhibition and memory encoding processes. *J.*
Neurosci. **35**, (2015). doi: 10.1523/JNEUROSCI.0519-15.2015.

64.   Chiu, Y. C. & Egner, T. Inhibition-Induced Forgetting: When More Control Leads
to Less Memory. *Psychol. Sci.* **26**, (2015). doi: 10.1177/0956797614553945.

65.   Yebra, M. *et al.* Action boosts episodic memory encoding in humans via
engagement of a noradrenergic system. *Nat. Commun.* **10**, (2019). doi:
10.1038/s41467-019-11358-8.

66.   Voss, M., Ingram, J. N., Haggard, P. & Wolpert, D. M. Sensorimotor attenuation
by central motor command signals in the absence of movement. *Nat. Neurosci.* **9**,
(2006). doi: 10.1038/nn1592.

67.   Chapman, C. E., Bushnell, M. C., Miron, D., Duncan, G. H. & Lund, J. P. Sensory
perception during movement in man. *Exp. Brain Res.* **68**, 516–524 (1987). doi:
10.1007/BF00249795.

68.   Cohen, L. G. & Starr, A. Localization, timing and specificity of gating of
somatosensory evoked potentials during active movement in man. *Brain* (1987)
doi:10.1093/brain/110.2.451. doi: 10.1093/brain/110.2.451.

69.   Colino, F. L., Buckingham, G., Cheng, D. T., van Donkelaar, P. & Binsted, G.
Tactile gating in a reaching and grasping task. *Physiol. Rep.* **2**, 1–11 (2014). doi:
10.1002/phy2.267.

70.   Papakostopoulos, D., Cooper, R. & Crow, H. J. Inhibition of cortical evoked
potentials and sensation by self-initiated movement in man. *Nature* **258**, 321–324
(1975). doi: 10.1038/258321a0.

71. Pertovaara, A., Kemppainen, P. & Leppänen, H. Lowered cutaneous sensitivity to nonpainful electrical stimulation during isometric exercise in humans. *Exp. Brain Res.* **89**, 447–452 (1992). doi: 10.1007/BF00228261.

72. Rushton, D. N., Roghwell, J. C. & Craggs, M. D. Gating of somatosensory evoked potentials during different kinds of movement in man. *Brain* **104**, 465–491 (1981). doi: 10.1093/brain/104.3.465.

73. Schütz-Bosbach, S., Musil, J. J. & Haggard, P. Touchant-touché: The role of self-touch in the representation of body structure. *Conscious. Cogn.* **18**, 2–11 (2009). doi: 10.1016/j.concog.2008.08.003.

74. Busse, L. & Woldorff, M. G. The ERP omitted stimulus response to 'no-stim' events and its implications for fast-rate event-related fMRI designs. *Neuroimage* **18**, (2003). doi: 10.1016/S1053-8119(03)00012-0.

75. Karamürsel, S. Human auditory fast and slow omitted stimulus potentials and steady-state responses. *Int. J. Neurosci.* **100**, (2000). doi: 10.3109/00207450008999674.

76. SanMiguel, I., Saupe, K. & Schröger, E. I know what is missing here: Electrophysiological prediction error signals elicited by omissions of predicted 'what' but not 'when'. *Front. Hum. Neurosci.* (2013) doi:10.3389/fnhum.2013.00407. doi: 10.3389/fnhum.2013.00407.

77. SanMiguel, I., Widmann, A., Bendixen, A., Trujillo-Barreto, N. & Schröger, E. Hearing silences: Human auditory processing relies on preactivation of sound-specific brain activity patterns. *J. Neurosci.* **33**, (2013). doi: 10.1523/JNEUROSCI.5821-12.2013.

78. Schomaker, J. & Meeter, M. Novelty enhances visual perception. *PLoS One* **7**, e50599 (2012).

79. Mayer, J. S., Kim, J. & Park, S. Enhancing visual working memory encoding: The role of target novelty. *Vis. cogn.* **19**, (2011). doi: 10.1080/13506285.2011.594459.

80. Schomaker, J. & Meeter, M. Short- and long-lasting consequences of novelty, deviance and surprise on brain and cognition. *Neuroscience and Biobehavioral Reviews* vol. 55 at https://doi.org/10.1016/j.neubiorev.2015.05.002 (2015). doi: 10.1016/j.neubiorev.2015.05.002.

81. Yon, D., Zainzinger, V., de Lange, F. P., Eimer, M. & Press, C. Action biases perceptual decisions toward expected outcomes. *J. Exp. Psychol. Gen.* (2020) doi:10.1037/xge0000826. doi: 10.1037/xge0000826.

82. Yon, D., Gilbert, S. J., de Lange, F. P. & Press, C. Action sharpens sensory representations of expected outcomes. *Nat. Commun.* **9**, (2018). doi: 10.1038/s41467-018-06752-7.

83.   Kurjak, A. *et al.* Fetal hand movements and facial expression in normal pregnancy
studied by four-dimensional sonography. *J. Perinat. Med.* **31**, (2003). doi:
10.1515/JPM.2003.076.

84.   Dogge, M., Custers, R. & Aarts, H. Moving Forward: On the Limits of Motor-
Based Forward Models. *Trends in Cognitive Sciences* vol. 23 at
https://doi.org/10.1016/j.tics.2019.06.008 (2019). doi: 10.1016/j.tics.2019.06.008.

85.   Dogge, M., Custers, R., Gayet, S., Hoijtink, H. & Aarts, H. Perception of action-
outcomes is shaped by life-long and contextual expectations. *Sci. Rep.* **9**, (2019).
doi: 10.1038/s41598-019-41090-8.

86.   de Lange, F. P., Heilbron, M. & Kok, P. How Do Expectations Shape Perception?
Trends in Cognitive Sciences vol. 22 at https://doi.org/10.1016/j.tics.2018.06.002
(2018). doi: 10.1016/j.tics.2018.06.002.

87.   Brooks, J. X. & Cullen, K. E. Predictive Sensing: The Role of Motor Signals in
Sensory Processing. *Biological Psychiatry: Cognitive Neuroscience and*
Neuroimaging vol. 4 at https://doi.org/10.1016/j.bpsc.2019.06.003 (2019). doi:
10.1016/j.bpsc.2019.06.003.

88.   Crapse, T. B. & Sommer, M. A. Corollary discharge across the animal kingdom.
Nat. Rev. Neurosci. **9**, 587–600 (2008). doi: 10.1038/nrn2457.

89.   Cullen, K. E. Sensory signals during active versus passive movement. *Curr. Opin.*
Neurobiol. **14**, 698–706 (2004). doi: 10.1016/j.conb.2004.10.002.

90.   Schneider, D. M. & Mooney, R. How Movement Modulates Hearing. *Annu. Rev.*
Neurosci. **41**, 553–572 (2018). doi: 10.1146/annurev-neuro-072116-031215.

91.   Straka, H., Simmers, J. & Chagnaud, B. P. A New Perspective on Predictive Motor
Signaling. *Curr. Biol.* **28**, R232–R243 (2018). doi: 10.1016/j.cub.2018.01.033.

92.   Audette, N. J., Zhou, W. & Schneider, D. M. Temporally precise movement-based
predictions in the mouse auditory cortex. *bioRxiv* 2021.12.13.472457 (2021)
doi:10.1101/2021.12.13.472457. doi: 10.1101/2021.12.13.472457.

93.   Poulet, J. F. A. A. & Hedwig, B. The Cellular Basis of a Corollary Discharge.
Science (80-.). **311**, 518–522 (2006). doi: 10.1126/science.1120847.

94.   Poulet, J. F. A. & Hedwig, B. Corollary Discharge Inhibition of Ascending
Auditory Neurons in the Stridulating Cricket. *J. Neurosci.* **23**, 4717–4725 (2003).
doi: 10.1523/JNEUROSCI.23-11-04717.2003.

95.   Schneider, D. M., Sundararajan, J. & Mooney, R. A cortical filter that learns to
suppress the acoustic consequences of movement. *Nature* **561**, 391–395 (2018).
doi: 10.1038/s41586-018-0520-5.

96.   Sawtell, N. B. Neural Mechanisms for Predicting the Sensory Consequences of

Behavior: Insights from Electrosensory Systems. *Annu. Rev. Physiol.* **79**, 381–399
(2017). doi: 10.1146/annurev-physiol-021115-105003.

97.   Brooks, J. X., Carriot, J. & Cullen, K. E. Learning to expect the unexpected: rapid
updating in primate cerebellum during voluntary self-motion. *Nat. Neurosci.* **18**,
1310–1317 (2015). doi: 10.1038/nn.4077.

98.   Cullen, K. E. The vestibular system: multimodal integration and encoding of self-
motion for motor control. *Trends Neurosci.* **35**, 185–196 (2012). doi:
10.1016/j.tins.2011.12.001.

99.   Roy, J. E. Dissociating Self-Generated from Passively Applied Head Motion:
Neural Mechanisms in the Vestibular Nuclei. *J. Neurosci.* **24**, 2102–2111 (2004).
doi: 10.1523/JNEUROSCI.3988-03.2004.

100.   Oldfield, R. C. The assessment and analysis of handedness: The Edinburgh
inventory. *Neuropsychologia* **9**, (1971). doi: 10.1016/0028-3932(71)90067-4.

101.   Kilteni, K. Methods of Somatosensory Attenuation. in *Somatosensory Research*
Methods 35–53 (2023).

102.   R Core Team. R: A Language and Environment for Statistical Computing. at
(2022).

103.   JASP Team. JASP (Version 0.16.3)[Computer software]. at (2022).

104.   Quintana, D. S. & Williams, D. R. Bayesian alternatives for common null-
hypothesis significance tests in psychiatry: A non-technical guide using JASP.
BMC Psychiatry **18**, (2018). doi: 10.1186/s12888-018-1761-4.

105.   van Doorn, J. *et al.* The JASP guidelines for conducting and reporting a Bayesian
analysis. *Psychonomic Bulletin and Review* vol. 28 at
https://doi.org/10.3758/s13423-020-01798-5 (2021). doi: 10.3758/s13423-020-
01798-5.

1073