Fatty acid amide hydrolase drives luminal cell differentiation in the adult mammary gland

Isabel Tundidor[1,2], Marta Seijo-Vila[1,2], Sandra Blasco-Benito[1,2], María Rubert-Hernández[1,2], Gema Moreno-Bueno[3], Laura Bindila[4], Manuel Guzmán[1,5], Cristina Sánchez[1,2*], and Eduardo Pérez-Gómez[1,2*].

[1]Dept. Biochemistry and Molecular Biology, Complutense University, Madrid, Spain.

[2]Instituto de Investigación Sanitaria Hospital 12 de Octubre, Madrid, Spain.

[3]MD Anderson International Foundation; Department of Biochemistry, Autonomous University of Madrid; Instituto de Investigaciones Biomédicas 'Alberto Sols' (CSIC-UAM); Instituto de Investigación Hospital Universitario La Paz (IdiPaz); Centro de Investigación Biomédica en Red de Cáncer (CIBERONC), Madrid, Spain.

[4]Clinical Lipidomics Unit, Institute of Physiological Chemistry, University Medical Center, Mainz, Germany.

[5]Instituto Ramón y Cajal de Investigación Sanitaria, Madrid, Spain.

* emails: eduperez@ucm.es; macsanch@ucm.es,

Summary

Mammary gland development occurs primarily in the adult, undergoing extensive expansion during puberty followed by cycles of functional specialization and regression with every round of pregnancy/lactation/involution. This process is ultimately driven by the coordinated proliferation and differentiation of mammary epithelial cells, but the endogenous factors regulating these developmental dynamics are still poorly defined. Endocannabinoid signaling is known to determine cell fate-related events involved in the development of different organs in the central nervous system and the periphery. Here, we report that the endocannabinoid-degrading enzyme fatty acid amide hydrolase (FAAH) plays a pivotal role in adult mammary gland development. Specifically, it is required for luminal lineage specification in the mammary gland, and it promotes hormone-driven secretory differentiation of mammary epithelial cells by controlling the endogenous levels of anandamide and the subsequent activation of CB_1R. Together, our results shed some light on the role of the endocannabinoid system in breast development and introduce FAAH-signaling as a new therapeutic target in milk production deficits.

Introduction

The mammary gland is a unique glandular organ that distinguishes mammals from all other animals and whose function is to produce and secrete milk to nourish offspring. Architecturally, the virgin mammary gland consists of a bilayered ductal tree with a luminal layer that contains hormone receptor-positive (HR+) cells and HR- cells primed for milk production (alveolar progenitors), and an outer basal layer that contains mostly contractile myoepithelial cells, but also bipotent mammary stem cells (MaSCs) (Visvader & Stingl, 2014). During puberty, steroid hormones drive extensive expansion of the ductal system to fill the mammary fat pad and, when pregnancy occurs, the influence of prolactin orchestrates the secretory differentiation of alveolar progenitors and the development of the lobuloalveolar units specialized in milk production. After weaning, the epithelial compartment undergoes massive cell death and returns to a morphological pre-pregnancy state in a process named post-lactational regression (Watson & Khaled, 2020). Determining the differentiation dynamics driving these postnatal developmental stages is not only essential to understanding normal mammary gland function but also the origins of breast cancer.

The endocannabinoid system (ECS) is a cell signaling system that plays multiple roles in the regulation of several aspects of cell physiology. This system is principally composed of (i) the G protein-coupled cannabinoid-selective receptors CB_1R and CB_2R, (ii) their endogenous ligands (the endocannabinoids) anandamide (N-arachidonoylethanolamine, AEA) and 2-arachidonoylglycerol (2-AG), and (iii) the enzymes that synthesize and metabolize these ligands (Fowler *et al.*, 2021). Previous studies have shown that endocannabinoid-associated signaling pathways regulate cell fate decisions such as proliferation, differentiation, and survival (Galve-Roperh *et al.*, 2013). Here, we tried to unravel the role of the ECS in adult mammary gland development by describing and

modulating the expression of one of the main responsible for endogenous AEA tone: the AEA-degrading enzyme fatty acid amide hydrolase (FAAH). We found that the expression of this enzyme is deeply associated to the luminal lineage in the adult mammary gland, and that it acts as a driver of lactogenic differentiation *in vitro* (by downregulating the endocannabinoid tone and its activity at CB_1R) and *in vivo* (it is necessary for the generation of hormone-sensing luminal cells in the mammary gland). Taken together, our data offer a window into the role of the ECS in the cellular dynamics that occur during human lactation and may provide further insights into the interplay between mammary gland development and breast cancer.

Materials & Methods

Reagents

The FAAH inhibitor URB 597 and the non-hydrolysable anandamide analog R-1 Methanandamide (metAEA) were purchased from Cayman Chemical. The FAAH inhibitor BIA 10-2474 was purchased from Medchem Express. Selective antagonists of CB_1R (SR141716, SR1), CB_2R (SR144528, SR2), PPARα (GW 6471) and PPARγ (GW 9662) were purchased from Tocris Bioscience.

HC11 lactogenic differentiation

The murine cell line HC11 was a kind donation by Dr. Castillo-Lluva (Complutense University, Madrid, Spain) and was authenticated by short tandem repeat profiling at the Genomics Core Facility of Instituto de Investigaciones Biomédicas 'Alberto Sols' (Madrid, Spain). Cells were maintained in subconfluence in RPMI medium (Sigma-Aldrich) with 10 % fetal bovine serum (FBS, Gibco), 10 µg/mL insulin (Sigma-Aldrich), 20 ng/mL epidermal growth factor (EGF, Gibco), 2 mM L-glutamine (Lonza), 10 mM HEPES (Lonza) and 1 % penicillin/streptomycin (Lonza). To induce their differentiation, subconfluent HC11 cells were trypsinized and seeded in complete medium at a density of 8.5×10^5 cells per well in 6-well plates. When they reached confluence (usually, overnight), complete medium was changed to EGF-free medium and predifferentiated cells were obtained after 48 h of growth in this medium. To obtain completely differentiated cells, predifferentiated cells were then treated for additional 5 days with differentiation medium: EGF-free medium containing 1 µg/mL prolactin (PRL) (Sigma-Aldrich) and 100 nM dexamethasone (DXM) (Sigma-Aldrich). Differentiation medium was changed and added fresh on day 3.

Lentiviral transduction

For lentiviral transduction of HC11 cells, a second-generation packaging system was used. Commercial packaging (psPAX2) and envelope (pMD2.G) plasmids (Addgene), as well as the pReceiver-Lv225 (Genecopoeia), were kindly donated by Dr. Fernández-Piqueras ("Severo Ochoa" Centre for Molecular Biology, Madrid, Spain). EX-hFAAH-Lv225 transfer plasmid was home generated by subcloning human *FAAH* into the XhoI and BstbI sites of pReceiver-Lv225. The empty (Ø) vector EX-NEG-Lv225 was used as negative control. Briefly, lentiviruses were generated by transfecting 3×10^6 HEK-293T cells in antibiotic-free medium with 6 µg of total DNA in a 1:2:3 ratio of pMD2.G:psPAX2:transfer plasmid using polyethylenimine (PEI) (Sigma-Aldrich). 48 h after transfection, conditioned medium containing recombinant lentiviruses was collected, filtered through 0.45 µm sterilization filters and added to HC11 cells together with 8 µg/mL polybrene (Sigma-Aldrich). Transduction efficiency was monitored by checking the expression of green fluorescent protein (GFP), which usually happened 48-72 h later. Antibiotic selection was not performed as the transfection efficiency was usually near 100 % and lasted through the complete HC11 differentiation protocol.

Endocannabinoid measurement

For sample preparation, cells were grown in 6-cm dishes and scraped off in ice-cold PBS. They were centrifuged twice at 2000 g for 10' at 4 °C and pellets were frozen in liquid nitrogen and stored at -80 °C until processed. Lipid extraction and parallel quantification of endocannabinoids was performed at the lipidomics unit of the *Universitätsmedizin* associated to the University of Mainz (Germany) as previously described (Bindila *et al*., 2016).

RNA extraction and real-time quantitative PCR (Q-PCR)

RNA was isolated with Nucleozol™ (Macherey-Nagel) following the extraction protocol suggested by the manufacturer. Reverse transcription (RT) was performed from 2-3 µg of RNA with the Transcriptor First Strand cDNA Synthesis Kit (Roche Life Science) using random hexamer primers. Real-time quantitative RT-PCR (Q-PCR) was performed in 384-well plates using the LightCycler® Multiplex DNA Master (Roche Life Science) and hybridization probes from the Universal ProbeLibrary Set (Roche Life Science). The relative expression ratio of target/reference gene was calculated with the $\triangle\triangle$Ct method (Pfaffl *et al*., 2001). Primer sequences for m*FAAH* were GCA GGT GGG CTG TTC AGT (sense) and AAG CAG GGA TCC ACA AAG TC (antisense); for m*Prlr*, TCC CTG GTA TGG CAG ACT TT (sense) and AAC ATC TGC GAT GCT CAC CT (antisense); for m*Tbp*, GGG GAG CTG TGA TGT GAA GT (sense) and CCA GGA AAT AAT TCT GGC TCA (antisense).

Protein extraction and Western blot (WB)

Cells were scraped off in RIPA buffer (0.1 % SDS, 0.5 % sodium deoxycholate, 1 % NP40, 150 mM NaCl, 50 mM Tris–HCl pH 8.0, in PBS) supplemented with protease and phosphatase inhibitors (Sigma). 25 µg of total protein were boiled for 5' at 95 °C, resolved by SDS-PAGE and transferred to PVDF membranes in a Trans-Blot® SD Semi-Dry Transfer Cell (Bio-Rad). Unspecific binding to membranes was prevented by blocking in 5 % bovine serum albumin (BSA) for 1 h at room temperature (RT). Primary antibodies [anti-milk (Nordic-MUbio #RAM/TM, dilution 1:5000), anti-FAAH (Abcam #ab128917, dilution 1:1000), anti-CB$_1$R (Frontier Institute #CB1-GP-Af530-1, dilution 1:500), anti-PRLR (Abcam #ab214303, dilution 1:1000), and anti-α-Tubulin (Sigma-Aldrich T9026, dilution 1:5000)] were incubated overnight at 4 °C in the blocking solution. Membranes

were then washed three times with TBS-T, incubated for 1 h at RT with the corresponding HRP-conjugated secondary antibody (Cytiva Lifescience) and developed with self-prepared ECL reagent (1.25 mM luminol, 0.2 mM p-coumaric acid, 100 mM Tris–HCl pH 8.5, in dH$_2$O). Densitometric analysis was performed with ImageJ™ software (Schindelin *et al*., 2012).

Human samples

Human mammary gland sections from reduction mammoplasties were kindly given by Dr. Moreno-Bueno (MD Anderson Cancer Center, Madrid, Spain). All patients gave informed consent, and the study was authorized by the respective Hospital Ethics Committees.

Immunofluorescence (IF)

Paraffin-embedded human tissue sections were de-waxed in isoparaffin H (Panreac) and hydrated by quick changes through serial ethanol baths. Antigen retrieval was performed by boiling in sodium citrate buffer (10 mM sodium citrate, 0.05 % Tween 20, pH 6.0, in dH$_2$O). Samples were subjected to several washes with dH$_2$O and washing buffer (NaCl 8.3 g/L, Tris 1.2 g/L; pH 7.4). Blocking was performed with 10 % goat serum in PBS, for 1 h at RT. Primary antibodies [(anti-FAAH (Abcam #ab128917, dilution 1:100), anti-CK8 (DSHB #531826, dilution 1:50), and anti-SMA (DSHB #2289065, dilution 1:50)] were diluted in Antibody Diluent (Dako) and incubated overnight at 4 °C. Samples were rinsed 2 x 5' in PBS + 0.25 % triton and incubated with fluorescent secondary antibodies (Invitrogen, dilution 1:200) and 1 µg/mL DAPI for 1 h at RT. Finally, they were mounted with Mowiol® mounting medium. Fluorescence confocal images were acquired by using an Olympus FV1200 microscope.

Animals

C57BL/6J FAAH $^{-/-}$ mice were kindly donated by Dr. Ben Cravatt's laboratory (Scripps Research Institute, La Jolla, CA, US). All procedures involving animals were performed with the approval of the Complutense University Animal Experimentation Committee and Madrid Regional Government according to the European official regulations. Animals were housed in the animal facility of the UCM School of Biology, under a 12 h light-dark cycle, and were allowed to feed and drink *ad libitum.*

Mammary gland dissection

For studies on the virgin mammary gland, tissue was harvested from 12 to 18-week-old C57BL/6J FAAH $^{+/+}$ or FAAH $^{-/-}$ female mice. For studies during the reproductive phase of mammary gland development (gestation, lactation, and involution), 12 to 18-week-old C57BL/6J FAAH $^{+/+}$ or FAAH $^{-/-}$ female mice were mated with wild type studs and allowed to litter. For gestation studies, tissue was harvested at day 15 after vaginal plug detection. For lactation studies, tissue was harvested at days 1, 5, 10 and 15 postpartum. In this case, pups were removed from the cage 2 h before mammary gland extraction to make sure that the differences between FAAH $^{+/+}$ and FAAH $^{-/-}$ mice were not due to different sucking stimuli at the time of dissection. For involution studies, pups were force-weaned at 10 days of lactation, and tissue was harvested 2 days after. Integrity of pairs #1 and #5 was usually compromised after opening the skin and thus were never dissected. Pairs #2 and #3 were usually snap frozen and then used for RNA/protein extraction. Pair #4 was the preferred option for imaging studies as it was easier to remove integrally.

Mammary gland histology and immunohistochemistry

For histology studies, mammary gland pair #4 was spread over a 6 cm dish, fixed overnight in 10 % formalin, and transferred to 50 % ethanol. Fixed mammary glands were then paraffin-embedded and sectioned at the Department of Medicine and Animal Surgery at UCM School of Veterinary, which also performed hematoxylin and eosin (H&E) staining. Prolactin IHC staining was performed in paraffin-embedded mouse tissue sections by the Histopathology Unit at the Spanish National Cancer (CNIO) (Madrid, Spain).

Flow cytometry

Mammary gland pairs #2, #3 and #4 were pooled and processed to obtain a single cell suspension through mechanical disaggregation followed by 2 h of enzymatic digestion in DMEM + 125 µg/mL collagenase (Sigma-Aldrich), 0.25 µg/mL dispase (Sigma-Aldrich) and 10 µg/mL DNAse (Sigma-Aldrich) at 37 °C. Luminal cells were isolated through the staining with Alexa Fluor® 488 CD24 antibody (Biolegend #101816, dilution 1:1000) and Alexa Fluor® 647 CD49f antibody (Biolegend #313610, dilution 1:20) and subsequently segregated into hormone-sensing and alveolar through the staining with Brilliant Violet 605™ Sca-1 antibody (Biolegend #108133, dilution 1:40) and PE CD49b antibody (Biolegend #108907, dilution 1:80). In addition, cell suspension was stained for biotinylated CD31 (Thermo Fisher Scientific #13-0311-82, diluted 1:200), CD45 (Thermo Fisher Scientific #13-0451-82, diluted 1:400) and TER119 (Thermo Fisher Scientific #13-5921-82, diluted 1:100) + PE Streptavidin (Biolegend #405203, diluted 1:300) to exclude endothelial and hematopoietic lineages (Lin$^+$ cells) from the analysis. Antibody incubations were sequentially performed in agitation for 20' at 4 °C and protected from light. A washing step with FACS buffer (PBS + 1 % FBS + 1 % BSA + 0.02 % sodium azide) was performed after each antibody incubation. Finally, cells were resuspended thoroughly in FACS buffer at a concentration of 1 x 10^6 cells/mL and

transferred to polystyrene round-bottom tubes. 7-aminoactinomycin D (7-AAD, Biolegend) was used as death cell marker and added 15' before the analysis. Data acquisition was performed in a FACSCalibur system (Becton Dickinson). Compensation was applied according to single-stained controls of the same cell type. For data analysis, FlowJo™ software was used. Unstained cells were used to set the blank for surface-marker expression after exclusion of cell debris, nonviable cells ($7\text{-}AAD^-$), and Lin^+ cells.

Statistical analyses

Unpaired, independent groups of two were analyzed by 2-tailed Student's t-test. For multi-group comparison, data were analyzed by 1-way ANOVA with post Tukey's multiple comparison test, or by 2-way-ANOVA when required. A p value of less than 0.05 was considered statistically significant. Unless otherwise stated, data are expressed as mean ± SEM from at least three biological replicates.

Results

FAAH expression is associated with luminal differentiated and hormone receptor-positive mammary epithelial cells. In order to investigate the role of FAAH in the adult mammary gland development, we first assessed its mRNA levels and those of other components of the ECS in human mammary epithelial cell (MEC) populations through publicly available datasets. Traditionally, cell sorting techniques for specific cell surface markers have been used to prospectively isolate MEC subpopulations from mammary tissue. More specifically, the combination of antibodies against epithelial cell adhesion molecule (EpCAM) (or the heat stable antigen CD24 as surrogate)- and integrin α6 (CD49f) allows segregation between the differentiated luminal subset (EpCAM$^+$ CD49f$^-$), the luminal progenitor subset (EpCAM$^+$ CD49f$^+$), and the basal/MaSC subset (EpCAM$^-$ CD49f$^+$), which is thought to contain both differentiated basal cells and bipotent MaSCs (Stingl *et al*., 2006). Microarray analyses on FACS sorted populations purified from reduction mammoplasties (Kannan *et al*., 2013) revealed that *FAAH* mRNA levels were significantly higher in the differentiated luminal subset than in the basal/MaSC subset, while expression levels in the luminal progenitor subset were located between the other two (Fig. 1a). Similar observations were made after the analysis of two additional studies involving gene expression profiling of human MEC subpopulations (Raouf *et al*., 2008; Lim *et al*., 2009) (Figs. 1b, c). One of these studies (Raouf *et al*., 2008) provided further segregation of the basal/MaSC subset into mature basal (CD49f$^-$ MUC1/CD133$^-$ CD10/THY1$^+$) and bipotent progenitor (CD49f$^+$ MUC1/CD133$^-$ CD10/THY1$^+$) subsets. No differences in *FAAH* levels were observed in these two subpopulations (Fig. 1c). In order to validate these results at the protein level, we performed immunofluorescence staining of FAAH in human adult mammary gland samples. Consistent with our previous

observations, FAAH was found to colocalize exclusively with luminal cells in the inner layer of the inactive lactiferous duct [characterized by the expression of cytokeratin 8 (CK8)], but not with basal cells in the outer layer [characterized by the expression of smooth muscle actin (SMA)] (Fig. 1d). Together, these results suggest that FAAH expression is associated with the luminal lineage in the adult mammary gland.

No remarkable differences between MEC populations were observed for the mRNA levels of other endocannabinoid (eCB)-synthesizing (*NAPEPLD*, *DAGLA,* and *DAGLB*) or eCB-degrading enzymes (*MGLL*) (Fig. 1a). Regarding cannabinoid receptors CB_1R and CB_2R (encoded by *CNR1* and *CNR2*, respectively) and other cannabinoid-related receptors that may be potentially engaged and activated by FAAH substrates (such as the peroxisome proliferator-activated receptors PPARα and PPARγ, encoded by *PPARA* and *PPARG*, respectively), mRNA levels of *CNR1*, *PPARA,* and *PPARG* also presented differential expression between cell subsets, with levels of *CNR1* and *PPARA* showing opposite patterns to that of FAAH, *i.e.,* higher in the basal/MaSC subset (Fig. 1a).

 In the recent years, single cell RNA sequencing (scRNA-seq) has allowed to comprehensively address the heterogeneity of the mammary epithelium as well as the identification of functionally relevant subpopulations within the luminal and basal compartments. To better characterize the expression pattern of FAAH in the mammary epithelial cell hierarchy, we next studied its mRNA levels in a dataset containing scRNA-seq profiling of mouse mammary glands from four adult developmental time points: nulliparous, mid gestation, lactation, and post involution (Bach *et al*., 2017) (Fig. 2a). Paralleling our observations in human tissue, *Faah* was predominantly expressed in cell clusters of differentiated luminal cells (C3, C4, C5, C8 and C11) and, to a lesser extent, in cell clusters of luminal progenitors (C1, C2, C6, C7, C9 and C10), while it was absent from basal (C12-C14) and MaSC (*i.e.*, Procr+ cells, C15) clusters (Fig. 2a). Importantly,

Faah's highest expression within luminal differentiated cells was found in C3 and C4 clusters, which showed characteristics of hormone-sensing cells such as high expression levels of genes encoding for hormone (estrogen, progesterone, and prolactin) receptors (*Esr1*, *Pgr*, and *Prlr*, respectively), while its expression was lower in HR- alveolar cells (C8 and C11), characterized by the expression of β-Casein (*Csn2*) (Bach *et al*., 2017) (Fig. 2a). Similar observations were made in human tissue after the analysis of *FAAH* mRNA levels in the UCSC Genome Browser, which integrates 24 K human mammary epithelial cell atlases obtained by scRNA-seq (Saeki *et al*., 2021) (Fig. 2b), and also in the Human Protein Atlas (Thul *et al*., 2017) (Supplementary Fig. 1).

Together, our results show a strong association between high FAAH expression and differentiated, HR+ cell phenotypes within the luminal lineage, and suggest that FAAH may be involved in cell differentiation and lineage specification in the mammary gland.

FAAH drives mammary epithelial cell differentiation *in vitro*. To validate the hypothesis that FAAH is involved in the differentiation of MECs, we next investigated its role in an *in vitro* model of lactogenic differentiation, a crucial process that occurs when pregnancy is advanced to expand the ductal system and differentiate it to generate milk secretion sites. Specifically, we utilized HC11 cells, a murine MEC model that displays progenitor-like features and retains its ability to differentiate into milk-producing cells in response to the addition of lactogenic hormones prolactin (PRL) and dexamethasone (DXM) (Ball *et al*., 1988). First, we differentiated HC11 cells from their progenitor-like (*i.e*., undifferentiated) state to an intermediate or "competent" (*i.e*., predifferentiated) state (by removing EGF from the culture medium), from which the addition of the lactogenic stimuli triggers their final differentiation into milk-producing (*i.e*., differentiated) cells (Fig. 3a). Differentiation was monitored by following the gradual increase in β-Casein expression, which is a widely accepted differentiation

marker in this cell line. In line with our previous results, HC11 cell differentiation (*i.e.*, β-Casein increase) was accompanied by a concomitant increase in FAAH expression (Fig. 3b).

To try to establish a cause-effect link between the increase in FAAH expression and HC11 differentiation, we next blocked FAAH activity by using the selective inhibitor URB 597 during the differentiation process. FAAH inhibition resulted in impaired lactogenic differentiation as determined by the prevention of the increase in β-Casein at the final stage of the process (Fig. 3c). These results were mimicked by using a different FAAH inhibitor (BIA 10-2474) (Fig. 3c), indicating that the catalytic activity of FAAH is required for the differentiation of HC11 cells.

To further confirm that FAAH is involved in the differentiation process associated with the lactogenic switch, we overexpressed the enzyme in HC11 cells via a lentiviral delivery system. As shown in Fig. 3d, FAAH overexpression led to an increase in β-Casein and other milk proteins corresponding to molecular weights previously observed for lactoferrin and γ-Casein (with apparent molecular weights of 80 and 25 kDa, respectively) (Boumahrou *et al*., 2011), even in those stages where lactogenic hormones had not been added yet (*i.e*., undifferentiated and predifferentiated), which further supports the notion that FAAH drives the differentiation of mammary epithelial cells.

FAAH drives MEC differentiation by decreasing anandamide-mediated activation of cannabinoid receptors. The results obtained from the experiments modulating the activity and expression of FAAH support the idea that the enzymatic activity of FAAH participates in the control of lactogenic differentiation of MECs *in vitro*, and therefore, they put forward the possible participation of one or several FAAH substrates in this process. FAAH is the main enzyme in charge of physiologically terminating the action of anandamide (AEA) and other bioactive lipids of the N-acylethanolamine (NAE) family,

such as palmitoylethanolamide (PEA) and oleoylethanolamide (OEA), that lack affinity for cannabinoid receptors (CBRs) but are active at non-canonical CBRs such as PPARα and PPARγ. To unmask the primary molecular target of the inhibition of lactogenic differentiation that may be driven by FAAH substrates, we next blocked a battery of receptors that can be engaged and activated by these compounds. Differentiating HC11 cells were incubated with selective antagonists for CB_1R (SR141716, SR1), CB_2R (SR144528, SR2), PPARα (GW 6471) or PPARγ (GW 9662) in the presence of URB 597. Only SR1 completely prevented the downregulation of β-Casein triggered by URB 597 (Fig. 4a), indicating that CB_1R mediates the FAAH-controlled inhibition of MEC differentiation. A partial prevention of the downregulation of β-Casein by SR2 was also observed, suggesting CB_2R might be partly involved too (Fig. 4a).

Given that only CBRs (and not PPARs) were proven to participate in the molecular mechanism underlying the effect of FAAH on MEC differentiation, we next analyzed the intracellular levels of CBR's endogenous ligands (the endocannabinoids, eCBs) in the HC11 *in vitro* differentiation model. We first observed that, in parallel to the increase in FAAH expression, AEA intracellular levels decreased throughout the differentiation process, while no changes in 2-arachidonoylglycerol (2-AG) levels were found (Fig. 4b). As expected, FAAH inhibition by URB 597 led to an increase in AEA levels in the three different differentiation stages (Fig. 4b), while the complementary approach (*i.e.*, FAAH overexpression) led to a decrease in AEA levels in undifferentiated, but not in predifferentiated and differentiated HC11 cells, probably due to the already high FAAH levels expressed by the cells in those stages (Fig. 4b).

All of the above points to an activation of CBRs by endogenous AEA as the responsible for the block of cell differentiation seen upon FAAH inhibition. In order to further study this causative link, we next challenged differentiating HC11 cells with methanandamide

(metAEA), a non-hydrolysable AEA analogue. MetAEA phenocopied the effect of FAAH pharmacological inhibition (*i.e.*, it impaired HC11 differentiation), and this effect was prevented by both SR1 and SR2 (Fig. 4c), further reinforcing that the AEA tone, as driven by FAAH, is involved in the control of MEC differentiation through the activation of CBRs. Of interest, SR1 not only prevented URB 597 (Fig. 4a) and metAEA (Fig. 4c) effect on β-Casein levels but increased them above those of the vehicle-treated cells. To better understand the importance of this observation, we treated HC11 cells during differentiation with SR1 alone and detected a concentration-dependent increase in β-Casein (Fig. 4d), this is, an induction of cell differentiation. All the above indicates that a high AEA tone, produced by low levels of FAAH, keeps MECs in an undifferentiated state through CB_1R activation. Following this rationale, a poorly active ECS would be expected in differentiated cells. In line with this idea, and in addition to an increase in FAAH expression and a decrease in AEA intracellular levels, HC11 differentiation was accompanied by a decrease in CB_1R (Fig. 4e).

Together, these data demonstrate that FAAH drives MEC differentiation by decreasing AEA-mediated activation of CB_1R.

FAAH expression is regulated by estradiol in mammary epithelial cells. Next, we aimed to find upstream regulators of FAAH expression that help clarify the biological significance of its role in MEC differentiation. Hormones are the main regulators of the adult phase mammary gland development. While estrogen and progesterone are key for ductal growth and tertiary branching during puberty and early pregnancy, PRL and glucocorticoids mostly drive alveolar morphogenesis that culminates at lactation (Neville *et al.*, 2002). Of interest, results presented in Fig. 2 and Supplementary Fig. 1 showed high FAAH expression in hormone receptor-positive luminal cells. To determine whether there is a functional link between these hormones and FAAH activity, we treated

undifferentiated HC11 cells with increasing concentrations of estradiol (E$_2$), progesterone (P$_4$), PRL and DXM. Among all treatments, only E$_2$ was able to stimulate FAAH expression (Fig. 5a). This effect was maximum 2 h after E$_2$ addition and quicky dissipated afterwards (Fig. 5b). The connection between estrogen signaling and FAAH was further strengthen by the analysis of mouse mammary gland *Faah* mRNA levels during pregnancy, lactation, and involution (Clarkson *et al*., 2004), that showed a parallel variation with that of the estrogen receptor (*Esr1*) (Fig. 5c). Together, these results raise the possibility that the regulation of FAAH expression in MECs by E$_2$ may have physiological relevance for the success of mammary gland development during the adult phase.

FAAH is necessary for the generation of hormone-sensing luminal cells and a functional PRLR-β-Casein axis in the mouse mammary gland. Since FAAH was found to be a relevant driver of MEC differentiation *in vitro*, we hypothesized that its genetic inactivation would have deleterious effects over mammary gland development in adult mice. Histological analyses of mouse tissue showed that the structure of FAAH $^{-/-}$ mammary glands at different adult developmental points (virgin, gestation, lactation, and involution) was essentially normal compared to that of FAAH $^{+/+}$ littermates, and that FAAH knockout did not affect the architecture or the number of mammary ducts or alveoli (Fig. 6a). However, flow cytometry-based analysis of Sca1 and CD49b markers (which allows to segregate the CD24$^+$ CD49f$^-$ luminal compartment into hormone-sensing and alveolar cells) revealed that virgin FAAH $^{-/-}$ mammary glands contained a significantly lower proportion of hormone-sensing cells (and a consequent higher proportion of alveolar cells) in their luminal layer compared to FAAH $^{+/+}$ controls (Fig. 6b), suggesting that FAAH knockout may have a negative impact on how MECs sense and integrate hormonal cues in order to drive mammary gland development. Given that

the developmental features mainly orchestrated by E_2 and P_4 (*i.e.*, duct length and tertiary branching during puberty and gestation) seemed unaffected in FAAH [-/-] mice (Fig. 6a), and based on the role that FAAH seems to play in PRL-dependent secretory differentiation *in vitro* (Fig. 4), we hypothesized that FAAH [-/-] mammary glands may have functional defects during lactation, which is mostly dependent on PRL signaling. In line with this, Western blot analysis demonstrated that FAAH [-/-] mammary glands had lower expression levels of β-Casein at different time points of lactation, and this was accompanied by a downregulation of PRL receptor (PRLR) at the same time points (Fig. 6c). Interestingly, treatment of HC11 cells with the FAAH inhibitor URB 597 also downregulated mRNA levels of *Prlr* (Fig. 6d). All of this suggests that FAAH [-/-] mammary glands may not undergo complete lactogenic differentiation because they exhibit partial resistance to PRL action. In line with this, no differences regarding PRL expression levels were found between FAAH [+/+] and FAAH [-/-] mammary glands during lactation.

Discussion

The reproductive phase of mammary gland development is a hormone-dependent morphogenetic process where the different epithelial cell types collaborate to generate lobuloalveolar units specialized in milk production. HR+ (or hormone-sensing) luminal cells play a key role in this process as they sense systemic hormonal changes and translate these into paracrine instructions for neighboring HR- (or alveolar progenitors) cells which, in response, will differentiate into alveolar cells able to synthesize and secrete milk (Visvader & Stingl, 2014). In this study, we demonstrate that the AEA-degrading enzyme FAAH is highly expressed by the hormone-sensing lineage of the adult mammary gland, and that it is involved in the acquisition of the hormone-sensing phenotype that integrates hormonal signaling in order to drive MEC differentiation.

Here, we used the HC11 cell line as a model of progenitor-like cells with the capability to differentiate into milk-producing cells in response to the addition of lactogenic stimuli. Treatment of these cells with the FAAH inhibitor URB 597 blocked lactogenic differentiation in a cannabinoid receptor (CBR)-manner, and this was mimicked by the AEA analogue metAEA, pointing to an elevated endogenous AEA tone triggered by FAAH inhibition as the main responsible for the phenotype in the former case. Over the years, attention has been focused on the possible role of AEA and other eCBs in regulating cell differentiation, which might account for some physio-pathological effects of these lipids (Galve-Roperh *et al*., 2013). For example, AEA treatment was shown to inhibit the differentiation of human keratinocytes (Paradisi *et al*., 2008) and neurons (Rueda *et al*., 2002) through CB_1R-mediated attenuation of cell type-specific differentiation pathways. Lactogenic failure in our FAAH-deficient models (both *in vitro* and *in vivo*) was accompanied by a downregulation of PRLR expression levels, pointing to an involvement of the PRLR pathway in the phenotype; however, metAEA treatment

of HC11 cells was unable to do so (data not shown). In spite of this, AEA has been previously reported to mediate CB_1R-dependent downregulation of PRLR in breast cancer cells -thus rendering them insensitive to the mitogenic action of PRL- (De Petrocellis *et al*., 1998), and therefore, a possible regulation by AEA of the PRLR pathway also in MECs should not be discarded. Future studies may inquire further into the relationship between the signaling pathways downstream CBR activation and those involved in MEC differentiation in order to shed some light on the role of FAAH in this process.

Importantly, treatment of differentiating HC11 cells with the CB_1R selective antagonist SR141716 (SR1) was sufficient to enhance cell-fate specification in a concentration-dependent fashion, pointing to either a ligand-independent constitutive activation of CB_1R or a basal eCB tone acting on CB_1R and actively inhibiting cell differentiation in MECs. Although the former cannot be ruled out, our results support the later hypothesis, given that constitutive activation of FAAH (accompanied by downregulation of its mains substrates) mimicked CB_1R antagonism, *i.e*., it promoted HC11 differentiation. No concentration-dependent effects on cell-fate specification were observed after treating HC11 cells with the CB2R selective antagonist SR144528 alone (data not shown), which is consistent with the scarce expression of CB_2R in the healthy mammary gland (Caffarel *et al*., 2006; Qamri *et al*., 2009; Pérez-Gómez *et al*., 2015). Together, these observations support the existence of an AEA-driven, basal CB_1R activity that prevents differentiation in luminal cells and is ultimately controlled by FAAH activity.

However, the contribution to inhibition of lactogenic differentiation of (i) FAAH substrates other than AEA that lack affinity for CB_1R as well as of (ii) other CB_1R agonists whose endogenous levels are not directly regulated by FAAH cannot be ruled out. Regarding the first scenario, it is well-known that palmitoylethanolamide (PEA) and

oleoylethanolamide (OEA) (also FAAH substrates) can downregulate the expression and activity of FAAH in breast cancer cells (Bisogno *et al*., 1998; Di Marzo *et al*., 2001). It could be argued that, by downregulating FAAH, these could be further enhancing the anti-differentiation effects of AEA in MECs. Regarding the second, it is worth noticing that the levels of 2-AG in HC11 cells also suffered a notable (although not statistically significant) increase in response to URB 597. Although monoacylglycerol lipase (MAGL) is responsible for 85 % of 2-AG hydrolysis in the mouse brain, it can also be cleaved into glycerol and arachidonic acid by FAAH (Maccarrone *et al*., 2015). However, 2-AG levels did not decrease upon FAAH constitutive activation, so we concluded that it is probably not contributing to the FAAH-controlled eCB tone acting at CB_1R in HC11 cells.

Our data also demonstrate that transgenic mice lacking FAAH show an aberrant development of the mammary gland during lactation, suggesting that FAAH may be necessary for this process to take place *in vivo*. In this regard, several hormones involved in the development of the mammary gland have been found to regulate FAAH expression through direct or indirect interaction with its promoter. Research in human neuroblastoma cells first identified estrogen and glucocorticoid response elements (EREs and GREs, respectively) in the *FAAH* promoter that regulated transcriptional activity independent of their ligand (Waleh *et al*., 2002). In the uterus, *FAAH* expression is downregulated by sex hormones (Maccarrone *et al*., 2000), and the opposite has been reported in mouse Sertoli cells, where E_2 induces ER binding to ERE2/3 sites in the *Faah* promoter and activates its transcription (Grimaldi *et al*., 2012). Likewise, there is evidence that progesterone binds and regulates transcriptional activity of the *FAAH* gene in human T lymphocytes through the transcription factor Ikaros (Maccarrone *et al*., 2003). Here, we demonstrate

that *FAAH* expression is induced by E_2 in MECs, and put forward the biological importance of ER-mediated regulation of FAAH in mammary gland development.

The evidence gathered in this study reveals the profound importance of FAAH in the differentiation dynamics of MECs in the adult mammary gland and raises important implications for the pathogenesis of breast cancer. Recent work from our lab established that FAAH genetic inactivation in the MMTV-neu transgenic mouse preceded the formation of highly proliferative, more aggressive tumors (manuscript under review). Here, we demonstrated that mammary glands from FAAH $^{-/-}$ mice show an expanded population of alveolar progenitors, which are the likely target for transformation in the MMTV-neu model (Henry *et al*., 2004). Therefore, it could be argued that, by regulating MEC plasticity in the normal mammary gland, FAAH could be involved in the early stages of HER2/neu-induced tumorigenesis, and probably other breast cancer subtypes.

Setting aside the insight provided into the potential role of FAAH in breast cancer, these results have translational relevance on their own. Inadequate milk production affects many women after giving birth, but the molecular determinants underlying lactational failure are still poorly understood (Olsen & Gordon, 1990; Kent *et al*., 2012). Recent studies demonstrated that treatment of HC11 cells with the plant-derived cannabinoids delta-9-tetrahydrocannabinol (THC) and cannabidiol (CBD) reduced the levels of the main nutritional components of the milk, thus reiterating the risks of cannabis consumption during pregnancy and lactation (Josan *et al*., 2022). In the same line, we demonstrate that FAAH KO animals (which are subjected to a higher eCB tone) show a significant decrease in β-Casein levels, which is an extremely important milk protein for infant health due to its ability to improve nutrient bioavailability and absorption at the intestinal level (Cosentino *et al*., 2010).

Together, these findings put forward the implication of the ECS in adult mammary gland development, and therefore contribute to the identification of physiological barriers for appropriate milk production, which are necessary to develop novel interventions to support breastfeeding. In addition, they represent a note of caution for the use of cannabis during pregnancy until further research unveils the physio-pathological relevance of these observations.

Figure legends

Figure 1. Expression of FAAH and related components of the ECS in human mammary gland cell populations. a-c Data were obtained from the microarray dataset published by Kannan *et al.*, 2013 (a), Lim *et al.*, 2009 (b), and Raouf *et al.*, 2008 (c). *FAAH*: fatty acid amide hydrolase; *MGL*: monoacylglycerol lipase; *NAPEPLD*: N-acyl phosphatidylethanolamine-specific phospholipase D; *DAGLA*, *DAGLB*: diacylglycerol lipase A and B; *CNR1*: cannabinoid receptor 1; *CNR2*: cannabinoid receptor 2; *PPARA*, *PPARG*: peroxisome proliferator activated receptor alpha and gamma. ANOVA: * p < 0.05, ** p < 0.01, *** p < 0.001. **d** Representative immunofluorescence staining of FAAH with cytokeratin 8 (CK8) as a luminal marker (upper panels) and with smooth muscle actin (SMA) as a basal marker (lower panels) in human mammary samples obtained from breast reductions. Cell nuclei are stained in blue. Scale bar = 50 μm.

Figure 2. FAAH is expressed in luminal hormone-sensing cell populations in all stages of adult mammary gland development. a,b t-SNEs plots representing mRNA expression of *FAAH* and cell population-specific genes in putative cell clusters of mammary cell populations defined by scRNA-seq analysis of developing mouse mammary glands as published by Bach *et al.*, 2017 (a) and Saeki *et al.*, 2021 (b). t-SNEs plots are colored by the normalized log-transformed expression of each of the genes. *ESR1*: estrogen receptor alpha; *PGR*: progesterone receptor; *PRLR*: prolactin receptor; *CSN2*: β-Casein.

Figure 3. FAAH drives lactogenic differentiation of MECs *in vitro*. a Schematic representation of the differentiation protocol of HC11 cells. **b** Representative WB analysis of β-Casein and FAAH at the three differentiation points. The specific band for FAAH in HC11 cells (pointed with an arrow) was determined by WB analysis of FAAH KO mouse tissue (data not shown). **c** Representative WB analysis of β-Casein in HC11

cells that have completed the differentiation protocol in the presence of the FAAH inhibitors URB 597 or BIA 10-2494 at 5 µM. **d** Representative WB analysis of β-Casein and other milk proteins during the complete differentiation of HC11 cells harboring an empty (Ø) or a FAAH overexpression vector. Densitometric values for all WBs after normalization against α-Tubulin and control condition are depicted in purple.

Figure 4. FAAH triggers HC11 cell differentiation by decreasing AEA-mediated cannabinoid receptor activity. a Representative WB analysis of β-Casein in HC11 cells after being differentiated in the presence of URB 597 at 5 µM and selective receptor antagonists at 1 µM. **b** Levels of the eCBs anandamide (AEA) and 2-arachidonoylglycerol (2-AG) in FAAH-modulated HC11 cells normalized to protein content. Values represent mean ± SEM of three independent samples. Student's t-test: * $p < 0.05$. **c** Representative WB analysis of β-Casein in HC11 cells after being differentiated in the presence of metAEA at 0.5 µM and selective antagonists to CB_1R and CB_2R at 1 µM. **d** Representative WB analysis of β-Casein in HC11 cells after being differentiated in the presence of increasing concentrations of the CB_1R selective antagonist SR1. **e** Representative WB analysis of CB_1R during HC11 differentiation with or without URB 597 at 5 µM. Densitometric values after normalization against α-Tubulin and control condition are depicted in purple. Dashed lines represent non-consecutive lanes.

Figure 5. FAAH expression is regulated by estradiol in undifferentiated HC11 cells. a, b qPCR analysis of *Faah* mRNA expression in undifferentiated HC11 cells after treatment with increasing concentrations of estradiol (E_2), progesterone (P_4), prolactin (PRL) and dexamethasone (DXM) for 4 h (a) or with 10 nM E_2 for the indicated times (b). Expression levels were normalized against *Tbp*. **c** *Faah* and *Esr1* mRNA expression at different time points of adult mammary gland development, obtained from the microarray dataset published by Clarkson *et al.*, 2004.

Figure 6. FAAH is necessary for the generation of hormone-sensing luminal cells and a functional PRLR-β-Casein axis in the mouse mammary gland. a Representative H&E-stained sections of FAAH $^{+/+}$ and FAAH $^{-/-}$ mouse mammary glands harvested at different time points of adult development (virgin, gestation day 15, lactation day 15, and involution day 2). n = 2 mice per time point. Scale bar = 200 μm (H&E). **b** Representative flow cytometric analysis of MEC populations of FAAH $^{+/+}$ and FAAH $^{-/-}$ virgin mammary glands. CD24$^+$ CD49f$^-$ luminal cells were isolated and then separated into hormone-sensing or alveolar lineages based on their expression of Sca-1 and CD49b (see Methods). Bar graph represents the mean ± SEM of n = 7 mice per group. **c** Representative WB analysis of β-Casein and PRLR at L1, L5, L10 and L15. Densitometric values after normalization against α-Tubulin and WT condition for every time point are depicted in purple. **d** qPCR analysis of *Prlr* mRNA expression in HC11 cells after being differentiated in the presence of the FAAH inhibitor URB 597 at 5 μM. Expression levels were normalized against *Tbp*. Data are shown as mean ± SEM from at least 3 independent experiments. Student's t-test: * p < 0.05. **e** IHC detection of PRL in samples at L10. Scale bar = 200 μm.

Supplementary Figure 1. FAAH is expressed in luminal hormone-sensing cell populations. t-SNEs plots representing mRNA expression of *FAAH* and cell population-specific genes in putative cell clusters of mammary cell populations defined by scRNA-seq analysis of developing mouse mammary glands as published by Thul *et al.*, 2017. t-SNEs plots are colored by the normalized log-transformed expression of each of the genes. *ESR1*: estrogen receptor alpha; *PGR*: progesterone receptor; *PRLR*: prolactin receptor; *CSN2*: β-Casein.

References

● Bach, K., Pensa, S., Grzelak, M., Hadfield, J., Adams, D. J., Marioni, J. C., & Khaled, W. T. (2017). Differentiation dynamics of mammary epithelial cells revealed by single-cell RNA sequencing. Nature communications, 8(1), 1-11.

● Ball, R. K., Friis, R. R., Schoenenberger, C. A., Doppler, W., & Groner, B. (1988). Prolactin regulation of beta-casein gene expression and of a cytosolic 120-kd protein in a cloned mouse mammary epithelial cell line. The EMBO journal, 7(7), 2089-2095.

● Bindila, L., & Lutz, B. (2016). Extraction and simultaneous quantification of endocannabinoids and endocannabinoid-like lipids in biological tissues. Endocannabinoid Signaling: Methods and Protocols, 9-18.

● Bisogno, T., Katayama, K., Melck, D., Ueda, N., De Petrocellis, L., Yamamoto, S., & Di Marzo, V. (1998). Biosynthesis and degradation of bioactive fatty acid amides in human breast cancer and rat pheochromocytoma cells: implications for cell proliferation and differentiation. European journal of Biochemistry, 254(3), 634-642.

● Boumahrou, N., Bevilacqua, C., Beauvallet, C., Miranda, G., Andrei, S., Rebours, E., … & Martin, P. (2011). Evolution of major milk proteins in Mus musculus and Mus spretus mouse species: a genoproteomic analysis. BMC genomics, 12(1), 1-13.

● Brisken, C., Kaur, S., Chavarria, T. E., Binart, N., Sutherland, R. L., Weinberg, R. A., ... & Ormandy, C. J. (1999). Prolactin controls mammary gland development via direct and indirect mechanisms. Developmental biology, 210(1), 96-106.

● Caffarel, M. M., Sarrió, D., Palacios, J., Guzmán, M., & Sánchez, C. (2006). Δ9-tetrahydrocannabinol inhibits cell cycle progression in human breast cancer cells through Cdc2 regulation. Cancer research, 66(13), 6615-6621.

● Clarkson, R. W., Wayland, M. T., Lee, J., Freeman, T., & Watson, C. J. (2004). Gene expression profiling of mammary gland development reveals putative roles for

death receptors and immune mediators in post-lactational regression. Breast Cancer Research, 6(2), 1-18.

● Cosentino, S., Gravaghi, C., Donetti, E., Donida, B. M., Lombardi, G., Bedoni, M., ... & Ferraretto, A. (2010). Caseinphosphopeptide-induced calcium uptake in human intestinal cell lines HT-29 and Caco2 is correlated to cellular differentiation. *The Journal of Nutritional Biochemistry*, *21*(3), 247-254.

● De Petrocellis, L., Melck, D., Palmisano, A., Bisogno, T., Laezza, C., Bifulco, M., & Di Marzo, V. (1998). The endogenous cannabinoid anandamide inhibits human breast cancer cell proliferation. Proceedings of the National Academy of Sciences, 95(14), 8375-8380.

● Di Marzo, V., Melck, D., Orlando, P., Bisogno, T., Zagoory, O., Bifulco, M., ... & de Petrocellis, L. (2001). Palmitoylethanolamide inhibits the expression of fatty acid amide hydrolase and enhances the anti-proliferative effect of anandamide in human breast cancer cells. Biochemical Journal, 358(1), 249-255.

● Fowler, C. J. (2021). The endocannabinoid system–current implications for drug development. Journal of Internal Medicine, 290(1), 2-26.

● Galve-Roperh, I., Chiurchiù, V., Díaz-Alonso, J., Bari, M., Guzmán, M., & Maccarrone, M. (2013). Cannabinoid receptor signaling in progenitor/stem cell proliferation and differentiation. Progress in lipid research, 52(4), 633-650.

● Grimaldi, P., Pucci, M., Di Siena, S., Di Giacomo, D., Pirazzi, V., Geremia, R., & Maccarrone, M. (2012). The faah gene is the first direct target of estrogen in the testis: role of histone demethylase LSD1. Cellular and molecular life sciences, 69(24), 4177-4190.

● Henry, M. D., Triplett, A. A., Oh, K. B., Smith, G. H., & Wagner, K. U. (2004). Parity-induced mammary epithelial cells facilitate tumorigenesis in MMTV-neu transgenic mice. Oncogene, 23(41), 6980-6985.

● Josan, C., Podinic, T., Pfaff, N., & Raha, S. (2022). Effect of Delta-9-tetrahydrocannabinol and cannabidiol on milk proteins and lipid levels in HC11 cells. *Plos one*, *17*(8), e0272819.

● Kannan, N., Huda, N., Tu, L., Droumeva, R., Aubert, G., Chavez, E., ... & Gilley, D. (2013). The luminal progenitor compartment of the normal human mammary gland constitutes a unique site of telomere dysfunction. Stem Cell Reports, 1(1), 28-37.

● Kent, J. C., Prime, D. K., & Garbin, C. P. (2012). Principles for maintaining or increasing breast milk production. Journal of Obstetric, Gynecologic & Neonatal Nursing, 41(1), 114-121.

● Lim, E., Wu, D., Pal, B., Bouras, T., Asselin-Labat, M. L., Vaillant, F., ... & Visvader, J. E. (2010). Transcriptome analyses of mouse and human mammary cell subpopulations reveal multiple conserved genes and pathways. Breast Cancer Research, 12(2), 1-14.

● Maccarrone, M., Bari, M., Di Rienzo, M., Finazzi-Agrò, A., & Rossi, A. (2003). Progesterone activates fatty acid amide hydrolase (FAAH) promoter in human T lymphocytes through the transcription factor Ikaros: evidence for a synergistic effect of leptin. Journal of Biological Chemistry, 278(35), 32726-32732.

● Maccarrone, M., De Felici, M., Bari, M., Klinger, F., Siracusa, G., & Finazzi-Agrò, A. (2000). Down-regulation of anandamide hydrolase in mouse uterus by sex hormones. European journal of Biochemistry, 267(10), 2991-2997.

●   Neville, M. C., McFadden, T. B., & Forsyth, I. (2002). Hormonal regulation of mammary differentiation and milk secretion. Journal of mammary gland biology and neoplasia, 7(1), 49-66.

●   Olsen, C. G., & Gordon Jr, R. E. (1990). Breast disorders in nursing mothers. American family physician, 41(5), 1509-1516.

●   Paradisi, A., Pasquariello, N., Barcaroli, D., & Maccarrone, M. (2008). Anandamide regulates keratinocyte differentiation by inducing DNA methylation in a CB1 receptor-dependent manner. Journal of Biological Chemistry, 283(10), 6005-6012.

●   Pérez-Gómez, E., Andradas, C., Blasco-Benito, S., Caffarel, M. M., García-Taboada, E., Villa-Morales, M., ... & Sánchez, C. (2015). Role of cannabinoid receptor CB2 in HER2 pro-oncogenic signaling in breast cancer. Journal of the National Cancer Institute, 107(6), djv077.

●   Pfaffl, M. W. (2001). A new mathematical model for relative quantification in real-time RT-PCR Nucleic Acids Res 29: e45. Find this article online.

●   Qamri, Z., Preet, A., Nasser, M. W., Bass, C. E., Leone, G., Barsky, S. H., & Ganju, R. K. (2009). Synthetic cannabinoid receptor agonists inhibit tumor growth and metastasis of breast cancer. Molecular cancer therapeutics, 8(11), 3117-3129.

●   Raouf, A., Zhao, Y., To, K., Stingl, J., Delaney, A., Barbara, M., ... & Eaves, C. (2008). Transcriptome analysis of the normal human mammary cell commitment and differentiation process. Cell stem cell, 3(1), 109-118.

●   Rueda, D., Navarro, B., Martínez-Serrano, A., Guzmán, M., & Galve-Roperh, I. (2002). The endocannabinoid anandamide inhibits neuronal progenitor cell differentiation through attenuation of the Rap1/B-Raf/ERK pathway. Journal of Biological Chemistry, 277(48), 46645-46650.

● Saeki, K., Chang, G., Kanaya, N., Wu, X., Wang, J., Bernal, L., ... & Chen, S. (2021). Mammary cell gene expression atlas links epithelial cell remodeling events to breast carcinogenesis. Communications biology, 4(1), 1-16.

● Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., ... & Cardona, A. (2012). Fiji: an open-source platform for biological-image analysis. Nature methods, 9(7), 676-682.

● Stingl, J., Raouf, A., Eirew, P., & Eaves, C. J. (2006). Deciphering the mammary epithelial cell hierarchy. Cell cycle, 5(14), 1519-1522.

● Thul, P. J., Åkesson, L., Wiking, M., Mahdessian, D., Geladaki, A., Ait Blal, H., ... & Lundberg, E. (2017). A subcellular map of the human proteome. Science, 356(6340), eaal3321.

● Visvader, J. E., & Stingl, J. (2014). Mammary stem cells and the differentiation hierarchy: current status and perspectives. Genes & development, 28(11), 1143-1158.

● Waleh, N. S., Cravatt, B. F., Apte-Deshpande, A., Terao, A., & Kilduff, T. S. (2002). Transcriptional regulation of the mouse fatty acid amide hydrolase gene. Gene, 291(1-2), 203-210.

● Watson, C. J., & Khaled, W. T. (2020). Mammary development in the embryo and adult: new insights into the journey of morphogenesis and commitment. Development, 147(22), dev169862.

Figure 1

Figure 2

a

HC11 Diff

URB 597 +

Veh | URB 597 | SR1 | SR2 | GW 6471 | GW 9662

β-Casein   1   0.3   1.6   0.7   0.3   0.4 — 37 kDa

α-Tubulin — 50 kDa

b

AEA

ng/g protein

Undiff | Prediff | Diff

2-AG

µg/g protein

Undiff | Prediff | Diff

Vehicle/∅   URB 597   FAAH-OE

c

HC11 Diff

Veh | metAEA | metAEA + SR1 | metAEA + SR2

β-Casein   1   0.4   1.5   0.9 — 37 kDa

α-Tubulin — 50 kDa

d

HC11 Diff

Veh | SR1 2 µM | SR1 4 µM | SR1 6 µM | SR1 8 µM

β-Casein   1   2.7   4.8   5.4   9.3 — 37 kDa

α-Tubulin — 50 kDa

e

Undiff | Prediff | Diff

CB$_1$R   1   0.8   0.4 — 50 kDa / 37 kDa

α-Tubulin — 50 kDa

a

E$_2$

Faah mRNA levels (Fold change vs. Veh)

Veh   10 nM   100 nM   1 μM

P$_4$

Veh   10 nM   100 nM   1 μM

PRL

Veh   10 nM   100 nM   1 μM

DXM

Veh   10 nM   100 nM   1 μM

b

HC11 Undiff

Faah mRNA levels (Fold change vs. Veh)

Vehicle
E$_2$ 10 nM

T0   1 h   2 h   4 h   8 h   24 h

c

mRNA levels (Fold change vs. T0)

Esr1
Faah

Virgin   G5   G10   G15   L0   L5   L10   I0.5   I1   I2   I3   I4

Gestation   Lactation   Involution

a

	Virgin	Gestation	Lactation	Involution
FAAH +/+				
FAAH -/-				

b

FAAH +/+

Sca-1 / CD49b

hormone sensing 67.0%

alveolar 28.8%

FAAH -/-

Sca-1 / CD49b

hormone sensing 32.5%

alveolar 60.4%

% of luminal cells

Hormone-sensing   Alveolar

FAAH +/+   FAAH -/-

c

Mammary glands

	L1		L5		L10		L15	
	WT	KO	WT	KO	WT	KO	WT	KO
PRLR	1	0.4	1	0.5	1	1.4	1	0.2
β-Casein	1	0.9	1	0.6	1	1.9	1	0.4
α-Tubulin								

50 kDa
37 kDa
37 kDa
50 kDa

d

HC11 Diff

Prlr/Tbp

mRNA levels (Fold change vs. Veh)

Veh   URB

e

FAAH +/+   FAAH -/-

PRL