Pelvic spine reduction affects diet but not gill raker morphology in two polymorphic brook

stickleback (*Culaea inconstans*) populations

Running title: Brook stickleback diet and gill rakers

Jonathan A. Mee[1*], Emily Yap[1], and Daniel M. Wuitchik[2]

[1] Department of Biology, Mount Royal University, 4848 Mount Royal Gate, Calgary, Alberta,

T3E 6K6, Canada

[2] Department of Biology, Boston University, 5 Cummington Mall, Boston, Massachusetts,

02215, United States

[*] Corresponding author: jmee@mtroyal.ca

Author contributions:

Jonathan A. Mee conceived of the study, performed the statistical analyses, and contributed to

writing the manuscript.

Emily Yap collected the gill raker data and contributed to writing of manuscript.

Daniel Wuitchik performed the metabarcoding bioinformatic analyses and contributed to writing

the manuscript.

Abstract

Pelvic spine polymorphism occurs in several species in the stickleback family

(Gasterosteidae). Given the similar phenotypic polymorphisms in multiple stickleback species,

we sought to determine the extent of parallelism in the ecological correlates of pelvic spine

reduction. Based on a metabarcoding analysis of brook stickleback gut contents in two

polymorphic populations, we found significant diet differences were associated with pelvic spine

reduction, but we found no clear or consistent trend supporting a tendency for benthic feeding in

pelvic-reduced brook stickleback. These results contrast with those found in threespine

stickleback where pelvic spine reduction is often associated with a benthic diet. Hence, we found

non-parallel consequences of spine polymorphism across species. Furthermore, a difference in

gill raker morphology has been frequently observed between ecomorphs with differen diets in

many fish species. But, we found no evidence of any difference in gill raker morphology

associated with pelvic spine polymorphism in brook stickleback.

Introduction

The maintenance of polymorphisms in natural populations is a central concept in

evolutionary biology. Polymorphisms are of particular interest when they are maintained at a

frequency that is too high to be explained by mutation-selection balance (Ford 1945, Lande

1975). In cases where a polymorphism persists in a population at high and stable frequency, the

maintenance of the polymorphism is typically explained by some form of balancing selection or

by heterosis (Dobzhansky 1971, Takahata et al. 1992, Rainey et al. 2000, Gigord et al. 2001,

Punzalan et al. 2005). These cases are of interest because they afford the opportunity to study the

genetic and ecological mechanisms that result in the balance of selective forces that maintain

diversity. For example, balancing selection, in the form of pollinator-mediated negative

frequency dependent selection, maintains a stable polymorphism in flower colour in the

rewardless orchid *Dactylorhiza sambucina* (Gigord et al. 2001). Regardless of whether or not a

given polymorphism is maintained by balancing selection (e.g. by frequency-dependent selection

or temporally-fluctuating selection), a polymorphism has the potential to affect other traits via

pleiotropy, and a polymorphism may cause divergence in a suite of traits via plastic or

evolutionary responses to the ecological or behavioural consequences of the polymorphism.

Studies of polymorphisms in threespine sticklebacks have provided foundational

contributions to our understanding of evolutionary processes (Schluter and McPhail 1992,

Reimchen 1994, Schluter 2000, Bolnick 2004, Colosimo et al. 2005, Chan et al. 2010, Hendry et

al. 2013, Schluter et al. 2022). The establishment of threespine stickleback as a model system in

evolutionary biology (Reid et al. 2021) has also increased interest in the study of adaptive

polymorphisms within other species in the stickleback family, Gasterosteidae (Blouw and Boyd

1992, Shapiro et al. 2009, Klepaker et al. 2013, Shikano et al. 2013, Lowey et al. 2020, Jeffries

et al. 2022, Willerth et al. 2022). For example, spine reduction or loss has evolved independently

in four stickleback genera that share a common ancestor approximately 27 million years ago:

Gasterosteus, *Pungitius*, *Apeltes*, and *Culaea* (Nelson 1969, Nelson 1977, Hagen and Blouw

1983, Blouw and Hagen 1984a, Chan *et al.* 2010, Varadharajan et al. 2019). The involvement of

predation by birds, fish, mammals, and insects as a selective agent on spines has been established

by many studies involving multiple stickleback species (Hoogland et al. 1957, Hall 1956,

Reisman and Cade 1967, Reimchen 1980, Reist 1980*a*, Reist 1980*b*, Reist 1983, Blouw and

Hagen 1984b, Foster et al. 1988, Reimchen 1992, Reimchen 1994, Vamosi 2002, Rundle et al.

2003, Vamosi and Schluter 2004, Marchinko 2009, Lescak and von Hippel 2011, Lescak et al.

2012, Miller et al. 2017). These studies typically take advantage of the existence of variation

among individuals within a population (or within sympatric or parapatric populations) to

determine which pelvic phenotype (e.g. spined or unspined) is favoured by a given predator. For

example, several studies have shown that gape-limited predators, such as trout (*Oncorhynchus*

mykiss) or juvenile pike (*Esox lucius*), select for increased spine length or against spine loss,

whereas predators that are not gape-limited, such as adult pike or larvae of *Dytiscus* spp. (water

beetles), select for reduced spines or spine loss (Reist 1980a, 1980b). Given that multiple

predators occur in an environment (e.g. a lake may contain insects, fish, and birds that prey on

stickleback; see Reimchen 1994), and that predators are not uniformly distributed within an

environment across space, seasonsm, or years (Reimchen 1994), spine polymorphism within

stickleback populations may be maintained by a balance of selection driven by multiple

predators (e.g. gape-limited and non-gape-limited) across multiple habitats or multiple points in

time. There is also evidence that spines may be selected against in calcium-deficient

environments, presumably due to the resource cost of gowing these bony structures (Giles 1983,

Bell et al. 1993, Spence et al. 2013, Haines et al. 2023). So, spine polymorphism may be

maintained by a balance of selection for spines by gape-limited predators and against spines due

to physiological costs in low-calcium environments (e.g. in some freshwater lakes).

The ecological correlates of spine polymorphism have been well-studied and

characterized in threespine stickleback (Bell and Foster 1994, Klepaker et al. 2013, Haines et al.

2023). These ecological correlates include differences between spine morphs in predation,

habitat preference, and diet, which, collectively or invidually, give the opportunity for additional

adaptive or plastic morphological or behavioural differences between spine morphs. Pelvic spine

divergence in some lakes is known to be associated with morphological and ecological

divergence between reproductively isolated limnetic and benthic threespine stickleback, wherein

the benthic species often evolves reduced spines (Schluter and McPhail 1992, Hatfield and

Schluter 1999, Schluter 1993, Hatfield 1997, Boughman 2001, Peichel et al. 2001, Rundle and

Schluter 2004). Even without reproductive isolation, a well-established difference associated

with spine reduction in threespine stickleback is that spined stickleback consume more

planktonic food sources and unspined stickleback consume more benthic food items, which is

associated with a difference in gill raker morphology where fewer, shorter, and thicker gill rakers

are found in benthic feeding fish (Reimchen 1980, McPhail 1984, Ridgeway and McPhail 1984,

Schluter 1993, Nagel and Schluter 1998, Rundle et al. 2003, Matthews et al. 2010). It is unclear,

however, whether the reduction in spines that has occurred in other stickleback species is

associated with parallel ecological and morphological changes, such as parallel changes in diet or

gill raker morphology.

In brook stickleback (*Culaea inconstans* Kirtland, 1840), within-population pelvic spine

polymorphism does not involve reproductively isolated morphs (Lowey et al. 2020) and the

ecological and morphological changes associated with pelvic spine reduction are not necessarily

consistent among populations (Willerth et al. 2022). To date, however, no study has directly

addressed whether pelvic spine reduction is associated with a difference in diet in brook

stickleback in a parallel manner to the difference in diet in pelvic-reduced relative to spined

threespine stickleback. Hence, the extent of ecological or phenotypic parallelism among

stickleback species associated with the reduction or loss of pelvic spines is unknown. Resolving

this question about the extent of parallelism among stickleback species will help address more

general questions about the extent of phenotypic parallelism across taxa (Bolnick et al. 2018, De

Lisle and Bolnick 2020).

In this study, we sought to understand whether pelvic spine reduction is correlated with

ecological and morphological changes in brook stickleback. We addressed the hypothesis that

pelvic spine polymorphism is associated with a difference in diet in brook stickleback, and we

predicted that, as in threespine stickleback, pelvic-reduced fish eat a more benthic diet. To

address our hypothesis, we used metabarcoding of gut contents to compare diet among spined

and pelvic-reduced brook stickleback in two polymorphic populations. Because gill raker

morphology differs between ecomorphs of threespine stickleback and many other fish species

(see, for example, Mee et al. 2015), we also hypothesized that gill raker morphology would

differ between spined and pelvic-reduced brook stickleback.

Methods

Sample collection and study site

Brook stickleback were collected in Alberta, Canada, from Muir Lake (UTF-8 encoded

WGS84 latitude and longitude: 53.627659, -113.957524) and Shunda Lake (52.453899, -

116.146192) in the summer of 2017 and 2019 using unbaited minnow traps. To sample fish from

the littoral zone, where they were more likely to have a benthic diet, traps were set within

throwing distance from the shore (maximum 4m to 5m from shore) at a depth of 0.5m to 2m. To

sample fish from the limnetic zone, where they were more likely to have a planktonic diet, traps

were set at least 50m from shore and suspended from floats at a depth of 1-2m. All traps were

retrieved within one to twelve hours after being set. In each year, all brook stickleback were

retained until we had collected 40 (in 2017) or 30 (in 2019) spined individuals. Then, once we

had captured 30 or 40 spined fish, we retained only pelvic-reduced individuals until we had

achieved a balanced sample of spined and pelvic-reduced morphotypes (Table 1). Pelvic

morphology was assessed at the time of capture via close inspection of the pelvic region of each

fish with forceps, and sex was identified at the time of capture based on the presence of male

nuptial colouration or the presence of eggs. Each fish included in this study was euthanized by

immersion in an overdose mixture of lake water and eugenol. In 2019, the entire gut of each fish

was removed and preserved in 95% ethanol. The body of each fish was preserved in 70%

ethanol. As described in Willerth et al. (2022), we stained each fish with Alizarin red to observe

calcium-containing osteocytes more clearly (such as in pelvic bones, spines, branchial arches,

and gill rakers) and we confirmed pelvic morphology in the lab using magnified digital images.

The fish collected from Muir Lake and Shunda Lake were used in previous comparisons

of stable isotope signatures and morphology between spined and unspined brook stickleback

(Willerth et al. 2022). These lakes are well-suited for studies of brook stickleback pelvic spine

polymorphism because they contain relatively high abundances of spined and pelvic-reduced

morphs (Nelson 1977, Lowey et al. 2020). The pelvic-reduced morphs include unspined

individuals that completely lack spines or any vestige of a pelvic girdle, but also individuals with

vestigial (a.k.a. intermediate) pelvic structures that lack some pelvic element (e.g. one spine

missing, half the pelvic girdle missing, etc.; see Klapaker et al. 2013 for a detailed description of

variation in pelvic morphology across stickleback species). In Muir Lake, approximately 40% of

the brook stickleback are unspined, with very few (less than 2%) intermediate morphs (Lowey et

al. 2020, and personal observation). In Shunda Lake, approximately 20% of the brook

stickleback have some sort of pelvic reduction (unspined or intermediate morphs), and

approximately 25% of these pelvic-reduced morphs are completely unspined (Lowey et al. 2020,

and personal observation). Recent surveys and stocking records (Alberta Environment and Parks

2021) suggest that salmonids, such as brown trout (*Salmo trutta* Linnaeus 1758) and rainbow

trout [*Ocorhychus mykiss* (Walbaum 1792)], are present in both lakes, although we also observed

other predators, such as loons [*Gavia immer* (Brunnich 1764)], dragonfly nymphs (Gomphidae

and other unidentified families), giant water bugs [*Lethocerus americanus* (Leidy 1847)], and

backswimmers (Notonectidae) in both lakes (Willerth et al. 2022). Both lakes contain abundant

Daphniid zooplankton, and littoral macroinvertebrate sampling revealed Trichoptera larvae,

amphipods, leeches, and Chironomidae larvae (Willerth et al. 2022). Sampling permits were

issued by the Government of Alberta, and fish handling protocols were approved by the Animal

Care Committee at Mount Royal University (Animal Care Protocol ID 101029 and 101795).

Diet analysis

Fish guts collected from 95 fish in 2019 (Table 1) were shipped to the Canadian Centre

for DNA Barcoding at the University of Guelph for membrane-based DNA extraction and

amplicon sequencing (Ivanova et al. 2006). This metabarcoding approach used five PCR primer

sets designed to amplify a barcode region of the cytochrome c oxidase subunit I (COI) gene in

arthropods, mollusks, annelids, amphipods, and microalgae (details of COI amplification can be

found at http://ccdb.ca/site/wp-content/uploads/2016/09/CCDB_Amplification.pdf). A second

round of PCR amplification added Ion Torrent sequencing adapters and a unique multiplex

identifier (MID) sequence to the 5' end of amplicons for each sample. Amplified COI fragments

were then pooled and single-end sequenced on an Ion Torrent PGM sequencing platform

(Thermo Fisher, Waltham, MA, USA). The resulting sequence data was automatically de-

multiplexed by the PGM Torrent Browser (Thermo Fisher, Waltham, MA, USA). We removed

primer and adapter sequences from the de-multiplexed reads, and discarded reads that consisted

of only primer or adapter sequence using CUTADAPT v2.3 (Martin 2011). We used DADA2

(Callahan et al. 2016) in R v4.2.2 (R Core Team 2022) to trim and filter reads, estimate error

rates for the filtered and trimmed amplicon dataset, combine identical reads into unique

sequences, and infer biologically meaningful (as opposed to spurious) amplified sequence

variants (ASVs). In the filtering and trimming step, we removed reads containing any unknown

basepairs, truncated reads to 200 bp, discarded reads less than 200 bp, truncated reads at the first

instance of a quality score less than 2, discarded reads that match the phiX genome, and

discarded reads with greater than 2 "expected errors" after truncation (where the number of

"expected errors" is calculated from the per-basepair quality scores for each read). In the

inference of ASVs, because Ion Torrent sequencing tends to make homopolymer errors and

produces higher numbers of indels (relative to Illumina sequencing), we applied a homopolymer

gap penalty of -1 and applied a more restrictive threshold for the number of insertions in one

sequence relative to another (BAND_SIZE = 32). We then created an ASV abundance table

(with the abundance of each ASV for each fish) and removed chimeric sequences using the

"consensus" method. We exported the list of ASVs across all samples to a fasta file and used the

JAMP pipeline in BOLDigger (Buchner & Leese 2020) to identify the best-fitting taxonomic

identification for each ASV.

We used PHYLOSEQ (McMurdie et al. 2013) to create a data object in R that allowed us

to associate the abundance of each taxon identified in each fish's gut with other sample data (i.e.

lake, sex, size, and pelvic phenotype). We removed any taxa identified as brook stickleback or

bacteria (the latter being more indicative of gut microbiome than diet). Because many taxa were

not identified to genus or species, we agglomerated taxa at the rank of family, while also

removing taxa that were not identified to the level of family, using the tax_glom function in

phyloseq. We analyzed individual-level variation in gut contents by performing an ordination

using non-metric multidimensional scaling (NMDS) and Bray-Curtis distances based on family-

level taxonomic assignment.

To test the hypothesis that brook stickleback pelvic spine reduction is associated with a

difference in diet, we used two analyses. First, to investigate an overall difference in diet, we

conducted a PERMANOVA using the adonis2 function in the vegan package (Oksanen et al,

2022) with 10000 permutations to evaluate whether centroids of the Bray-Curtis distances

(calculated using phyloseq, as described above) differed among groups. Second, to investigate

differences between groups for specific diet items, we used DESeq2 (Love et al. 2014) to

conduct Wald tests of the Log_2 fold-differences in the normalized abundances of each eukaryote

family in the diets of brook stickleback (normalized by individual fish) between lakes and

between pelvic phenotypes (with p-values corrected for multiple testing with the Benjamini-

Hochberg method; Benjamini and Hochberg 1995). Because we had previously found a

significant difference in length between fish collected in near-shore "littoral" traps and fish

collected in "limnetic" traps suspended >50m from shore (Willerth et al. 2022), which would

lead to collinearity in statistical models investigating the effects of fish length and trap location

on diet, we looked for correlations among predictor variables using generalised linear models

with a Gaussian error distribution (for length – a continuous numerical variable) or a binomial

error distribution (for factor variables – pelvic phenotype, sex, and trap location). Given the

results of these tests (details below in the results section) showing significant correlation between

length and habitat (but between no other predictor variables), we used only one of either habitat

or length in our analyses. In the comparison between lakes (for the PERMANOVA and Wald

tests), we specified a model that included lake, pelvic morphology, sex, and scaled total length

(i.e. scaled to mean = 0 and variance = 1), but not habitat because no fish were captured in

"limnetic" traps in Shunda Lake. Because we expect diet might be affected by allometry

(Aguirre et al. 2008, Reimchen et al. 2016), we also included interactions with length in all our

diet models. Because of concerns with over-fitting given our sample size, we did not include

other interaction terms in our diet models. For the Wald tests, we subsequently conducted

separate analyses for each lake, wherein we specified models (one per lake) that included pelvic

phenotype (spined or pelvic-reduced), sex, scaled total length (or habitat in the case of Muir

Lake), and two-way interactions with length.

Gill rakers

Fish that had been bleached and stained with alizarin red for morphological analyses

(Willerth et al. 2022) were used to investigate variation in gill raker morphology associated with

different pelvic phenotypes. We removed the first branchial arches from both sides of each fish

with forceps and scissors, cleared away all the gill filaments, and photographed the gill rakers in

a plastic weighing boat using a Nikon SMZ-75T Greenough-type stereo microscope with a

mounted Imaging Source camera and NIS-Elements D Software (©2021 Nikon Corporation).

Following Kaeuffer et al. (2011), we counted the number of gill rakers on the left and right gill

arches, and we measured the length of the second through fourth gill raker from the epibranchial-

ceratobranchial joint on the ceratobranchial, which were consistently the three longest gill rakers

on each arch. We used generalized linear models in R with a gaussian error distribution to model

the effects of fish length, sex, year of capture, lake, and pelvic phenotype on variation in gill

raker length or number. As described above, we used length as a proxy for habitat to avoid

collinearity between the length and habitat variables. We set contrasts among factors using the

contr.sum function, and we used the CAR package (Fox and Weisberg 2019) to generate an

ANOVA table with type III sums of squares. Given the larger sample size for our gill raker

dataset, and the reasonable possibility that the effect of pelvic reduction on gill raker morphology

differs between sexes, between lakes, or between years, we included all two-way interactions in

our initial models, and then used a reverse-stepwise approach for model selection, removing any

interaction terms that were not significant (p > 0.05).

Results

A list of all taxa identified via metabarcoding in Muir Lake and Shunda Lake is available

in the supplementary material (Table S1 and Table S2). At the population level, brook

stickleback in Muir Lake and Shunda Lake had taxonomically diverse diets. We identified

thirteen Eukaryote phyla among the Amoebozoa, Archaeplastida, Chromalveolata, Fungi, and

Metazoa. The Metazoa was by far the most diverse taxon in both lakes where we identified six

different phyla, the most abundant and diverse of which were the arthropods. We identified six

arthropod classes: Arachnida, Branchiopoda, Copepoda, Insecta, Malacostraca, and Ostracoda.

Insecta was by far the most abundant and diverse class (among all phyla) in the diets of brook

stickleback in both lakes, with six orders and sixteen families. We focused on family-level

classifications because many taxa were not resolved to genus or species. Prior to statistically

evaluating differences in diet between lakes and between pelvic phenotypes, we evaluated

correlations among predictor variables. Using a type II sums of squares ANOVA (i.e. evaluating

a main effect conditional on all other main effects without significant interaction effects), we

found a significant effect of habitat (limnetic or littoral) on fish length in Muir Lake (df = 1,

Wald's X^2 = 4.7997, p = 0.028), and a significant effect of fish length on habitat in Muir Lake (df

= 1, Wald's X^2 = 4.8788, p = 0.027). No other predictor variables were correlated.

Large individual-level differences in diet among brook stickleback obscured any obvious

group-level differences in diet between lakes or between pelvic morphologies in an ordination

plot (Figure 1). A PERMANOVA test of overall differences in diet did, however, reveal that we

can reject the null hypothesis that pelvic spine morphs have the same centroid, indicating that

diet does differ significantly between pelvic spine morphs (Table 2). Wald tests of the Log_2 fold-

differences in the abundances of each eukaryote family in the diets of brook stickleback between

lakes suggested that there were significant differences in the diets of brook stickleback in Muir

Lake compared to Shunda Lake attributable to several taxa (Figure 2). Several families found in

Muir Lake were not found in Shunda Lake and vice versa: Unionicolidae, Culicidae, Muscidae,

Chaoboridae, Hydrozetidae, Ascarididae, Succineidae, Planorbidae, Crambidae, Candonidae,

Naididae, Gelechiidae, Chlorellaceae, Scirtidae, and Cyclopidae were not found in Shunda Lake,

whereas Simuliidae, Baetidae, Tephritidae, Sellaphoraceae, Saprolegniaceae, Leptolegniaceae,

Spionidae, Cyprididae, and Bucculatricidae were not found in Muir Lake. The diet differences

between lakes were not driven entirely by the absence of taxa in one lake – several taxa that were

present in both lakes (e.g. Caenidae) were consumed in greater abundance in one lake relative to

the other (Figure 2).

To further address the hypothesis that brook stickleback pelvic spine reduction (or loss) is

associated with a difference in diet, we evaluated the differential abundance of each taxon

identified in the stomachs of spined and pelvic-reduced brook stickleback in Muir Lake and

Shunda Lake. When fish length was included as a factor in analyses of diets in Muir Lake,

Daphniidae (daphnia, or water fleas) were significantly more abundant in the diets of spined fish,

whereas Caenidae (squaregill mayflies), Gammaridae (amphipods commonly known as "scuds"),

and Hyalelidae (amphipods) were significantly more abundant in the diets of pelvic-reduced fish

(Figure 3). When habitat (instead of length) was included as a factor in analyses of diets in Muir

Lake, only Caenidae were significantly more abundant in the diets of pelvic-reduced fish (Figure

4). In Shunda Lake, Lumbricidae (earthworms) were significantly more abundant in the diets of

spined fish (Figure 3). Diets also differed significantly based on fish length and between sexes

(Figure 3, Figure 4).

To investigate a potential change in gill raker morphology between spined and unspined

brook stickleback, we compared gill raker morphology between individuals with spined and

reduced pelvic phenotypes in Muir Lake and Shunda Lake. We conducted separate analyses

where we looked for an effect on maximum gill raker length (i.e. the length of the longest gill

raker in each individual), mean gill raker length, the length of a specific gill raker (e.g. the

second gill raker from the epibranchial-ceratobranchial joint on the ceratobranchial), the number

of gill rakers on the left or right gill arch, or the mean number of gill rakers on both arches. We

also conducted analyses involving the effect of pelvic morphology with three categories (i.e.

spined, vestigial, and unspined) instead of two categories (i.e. spined and pelvic-reduced).

Whereas the effects of some factors varied in magnitude among these analyses, no analysis

revealed a significant effect of pelvic morphology on gill raker size or number. For clarity, we

report only the details of our analyses involving the effect of pelvic reduction (with two

categories of pelvic morphology) on maximum gill raker length (Table 3, Figure 5) and on the

mean number of gill rakers (Table 4, Figure 6). Gill raker length and number did not differ

significantly between spined and pelvic-reduced brook stickleback. Gill raker length and number

were significantly affected by fish size, with larger fish having more and longer gill rakers.

Brook stickleback in Muir Lake had more gill rakers and longer gill rakers than those in Shunda

Lake. Male brook stickleback had significantly longer gill rakers than female brook stickleback,

especially at larger sizes. The effect of fish size on gill raker length was weaker for samples

collected in 2019 than for samples collected in 2017 (see Tables 3 and 4 for details of statistical

results).

Discussion

Whereas the factors maintaining the pelvic spine polymorphism within brook stickleback

populations are unclear, it is possible, if not likely, that spined and unspined fish differ

ecologically. A previous study of stable isotope signatures revealed that spined and pelvic

reduced brook stickleback are associated with different habitat types (Willerth et al. 2022),

suggesting that the pelvic spine polymorphism may have ecological correlates. In this study, we

tested the hypothesis that spined and pelvic-reduced brook stickleback have different diets.

Consistent with our hypothesis, we found significant differences in the abundance of some taxa

in the diets of spined and pelvic-reduced brook stickleback.

Pelvic spine reduction in brook stickleback and threespine stickleback populations has

evolved in parallel, likely driven by similar selective forces. The extent to which we should

expect parallelism to extend beyond a particular phenotype of interest (in this case, pelvic spine

reduction), resulting in parallel ecological, behavioural, and morphological consequences, is

unknown. We predicted that, like threespine stickleback, pelvic spine reduction in brook

stickleback should be associated with a more benthic diet. Our observations did not align with

this prediction, which implies non-parallelism in the ecological correlates of spine reduction.

Diet differences between spined and pelvic-reduced brook stickleback did not indicate a clear or

consistent trend towards a benthic or limnetic diet in either morph. The taxon-specific

differences in diet between spined and pelvic-reduced brook stickleback that we report here

should not, however, be over-interpreted. The specific diet items that differed significantly

between pelvic morphs were highly dependent on choices in our statistical analyses. For

example, we note that our choice whether to used fish size or habitat as a variable in our model

for Muir Lake influenced our inferences, although this was likely partly due to the reduction in

power associated with replacing a continuous variable (length) with a factor variable (habitat). In

addition, whether we included more or fewer two-way interactions in our analyses also had a

strong effect on which specific taxa differed significantly between pelvic morphs (these

alternative analyses are not presented here). Furthermore, our metabarcoding approach did not

allow us to differentiate taxa that were actually targeted by stickleback from taxa that happen to

be in the guts of the stickleback's prey. For example, Pythiaceae, known as water moulds, and

which we identified in the diets of brook stickleback, are parasites of plants and animals, and

may have been found in the guts of mayflies (or other herbivorous prey), or may have been

ingested as part of some other diet item, such as rotting vegetation. Our metabarcoding approach

also did not allow us to distinguish different prey life stages, which may be associated with

different habitats, making it difficult to make inferences regading stickleback foraging habitat

based on diet. Nonetheless, all of our analyses indicate some significant difference in diet

between spined and pelvic-reduced brook stickleback, providing support to the possibility that

pelvic reduction has ecological consequences, such as a difference in foraging habitat.

Differences in diet associated with pelvic spine reduction in threespine stickleback are

frequently accompanied by differences in gill raker morphology (McPhail 1984, Ridgeway and

McPhail 1984, Schluter 1993, Nagel and Schluter 1998, Rundle et al. 2003, Matthews et al.

2010). Longer, more slender, more numerous gill rakers are better for planktonic feeding,

whereas shorter, thicker, less numerous gill rakers are better for benthic feeding. In brook

stickleback, we did not observe a significant difference in gill raker morphology between spined

and pelvic-reduced fish. There are several possible reasons why the difference in diet between

spined and pelvic-reduced brook stickleback is not associated with a difference in gill raker

morphology. First, the diet difference associated with pelvic reduction in brook stickleback does

not necessarily involve a decrease in planktonic feeding. Second, spined and pelvic-reduced

brook stickleback are not reproductively isolated, and each lake likely contains a panmictic

population (Lowey et al. 2020). So, in the absence of pleiotropic effects of the spine

polymorphism, divergence (or polymorphism) in other traits that are not directly affected by

selection is unlikely to evolve. Lastly, the diet differences between spined and pelvic-reduced

brook stickleback may not induce any developmental plasticity in gill raker morphology.

It is unclear whether the diet differences between spined and pelvic-reduced brook

stickleback are, in fact, associated with a difference in habitat. It is possible that different pelvic

morphs use different resources in the same habitat. If this were the case, then maintenance of

spine polymorphism must depend on predators in the same habitat exerting balancing selection

pressure simultaneously or in a temporally segregated manner, or on a balance between predation

pressure and selection based on physiological costs in calcium-deficient environments. A fruitful

avenue of future research would likely be an investigation of temporally varying predation on

brook stickleback between seasons, between years, and across multiple generations.

Observations of parallel evolution, such the pelvic spine reduction in multiple stickleback

species, have long been interpreted as evidence for the involvement of natural selection as a

mechanism of evolutionary change. Evolutionary biologists are now trying to understand when,

and when not, to expect the evolution of parallel phenotypes to be associated with parallel

ecological consequences (e.g. the same behavioural or habitat divergence) and parallel biological

pathways (e.g. the same genes or the same mutations). In the case of threespine stickleback

pelvic spine reduction, there seems to be parallelism among populations in the genetic basis of

spine reduction, and there appears to be parallelism in the ecological consequences and correlates

of spine reduction (Miller et al. 2015). Our comparison with instances of pelvic spine reduction

in brook stickleback suggests little parallelism between species. Whether, and why, the extent of

parallelism might generally be lower in between-species versus between-population

comparisons, despite the likelihood of very similar selective scenarios (e.g. based on balancing

or divergent selection due to predation in freshwater environments), will undoubtedly be the

focus of future research.

Acknowledgments

We thank two anonymous reviewers and the associate editor who reviewed a previous version of

this manuscript for constructive feedback. Samples were collected with the assistance of Alex

Farmer, Carolyn Ly, Alyce Straub, Bryce Carson, Moroni Lopez Vasquez, and Kaylee

Olszewski. A Natural Sciences and Engineering Research Council of Canada Discovery Grant

awarded to J.A.M. supported this work, and E.Y. was supported by an NSERC URSA.

Data Availability

Private URL for DRYAD archive during peer review (Clicking the link immediately launches a

download of the data files):

https://datadryad.org/stash/share/CJuLVp4HYMFaQMBDsCUnMMhK_jh4j4V-eSz9Qp6T21E

Literature Cited

Aguirre, W. E., K. E. Ellis, M. Kusenda, and M. A. Bell. 2008. Phenotypic variation and sexual

413   dimorphism in anadromous threespine stickleback: implications for postglacial adaptive

414   radiation. *Biological Journal of the Linnean Society*, 95:465-478.

415   Alberta Environment and Parks. (Accessed 2021, June 1). Fish and Wildlife Information

416   Management System. https://www.alberta.ca/fisheries-and-wildlife-management-

417   information-system-overview.aspx

418   Bell, M. A., Ortí, G., Walker, J. A., & Koenings, J. P. (1993). Evolution of pelvic reduction in

419   threespine stickleback fish: a test of competing hypotheses. *Evolution*, 47(3), 906-914.

420   Bell, M.A., and S.A. Foster. 1994. The evolutionary biology of the threespine stickleback.

421   Oxford University Press, New York, NY.

422   Benjamini, Y., & Hochberg, Y. (1995). Controlling the false discovery rate: a practical and

423   powerful approach to multiple testing. *Journal of the Royal Statistical Society: Series B*

424   *(Methodological)*, 57(1), 289-300.

425   Blouw, D. M., and G. J. Boyd. 1992. Inheritance of reduction, loss, and asymmetry of the pelvis

426   in Pungitius pungitius (ninespine stickleback). *Heredity*, *68*(1), 33-42.

427   Blouw, D. M., and D. W. Hagen. 1984a. The adaptive significance of dorsal spine variation in

428   the fourspine stickleback, *Apeltes quadracus*, I. Geographic variation in spine number.

429   *Canadian Journal of Zoology*, 62(7):1329-1339.

430   Blouw, D. M., and D. W. Hagen. 1984b. The adaptive significance of dorsal spine variation in

431   the fourspine stickleback, *Apeltes quadracus*. III. Correlated traits and experimental

432   evidence on predation. *Heredity*, 53(2):371-382.

433   Bolnick, D. I. 2004. Can intraspecific competition drive disruptive selection? An experimental

434   test in natural populations of sticklebacks. *Evolution*, *58*(3), 608-618.

Bolnick, D. I., R. D. Barrett, K. B. Oke, D. J. Rennison, and Y. E. Stuart. 2018. (Non) parallel

evolution. *Annual Review of Ecology, Evolution, and Systematics, 49*(1):303-330.

Boughman, J. W. 2001. Divergent sexual selection enhances reproductive isolation in

sticklebacks. *Nature*, 411(6840):944-948.

Buchner, D., and F. Leese. 2020. BOLDigger–a Python package to identify and organise

sequences with the Barcode of Life Data systems. *Metabarcoding and Metagenomics*,

4:e53535.

Callahan B. J., P. J. McMurdie, M. J. Rosen, A. W. Han, A. J. A. Johnson, and S. P. Holmes

(2016). DADA2: High-resolution sample inference from Illumina amplicon data. *Nature*

Methods, 13:581-583.

Chan, Y. F., M. E. Marks, F. C. Jones, G. Villarreal Jr, M. D. Shapiro, S. D. Brady, A. M.

Southwick, D. M. Absher, J. Grimwood, J. Schmutz, R. M. Myers, D. Petrov, B. Jónsson,

D. Schluter, M. A. Bell, and D. M. Kingsley. 2010. Adaptive evolution of pelvic

reduction in sticklebacks by recurrent deletion of a Pitx1 enhancer. *Science*,

327(5963):302–305.

Colosimo, P. F., K. E. Hosemann, S. Balabhadra, G. Villarreal Jr, M. Dickson, J. Grimwood, J.

Schmutz, R. M. Myers, D. Schluter, and D. M. Kingsley. 2005. Widespread parallel

evolution in sticklebacks by repeated fixation of ectodysplasin alleles. *Science*,

307(5717):1928-1933.

De Lisle, S. P., and D. I. Bolnick. 2020. A multivariate view of parallel evolution. *Evolution*,

74(7):1466-1481.

Dobzhansky, T. 1971. *Genetics of the Evolutionary Process*. Columbia University Press, New

York, NY.

Ford, E. B. 1945. Polymorphism. *Biological Reviews* 20:73–88.

Foster, S.A., V.B. Garcia, and M.Y. Town. 1988. Cannibalism as the cause of an ontogenetic

460   shift in habitat use by fry of the threespine stickleback. *Oecologia*, 74(4):577-585.

461   Fox, J., and S. Weisberg. 2019. An {R} Companion to Applied Regression, Third Edition.

462   Thousand Oaks CA: Sage. URL:

463   https://socialsciences.mcmaster.ca/jfox/Books/Companion/

464   Gigord, L. D., M. R. Macnair, and A. Smithson. 2001. Negative frequency-dependent selection

465   maintains a dramatic flower color polymorphism in the rewardless orchid *Dactylorhiza*

466   *sambucina* (L.) Soò. *Proceedings of the National Academy of Sciences of the United*

467   *States of America*, 98(11):6253-6255.

468   Giles, N. (1983). The possible role of environmental calcum levels during the evolution of

469   phenotypic diversity in Outer Hebridean populations of the three‐spined stickleback,

470   Gasterosteus aculeatus. Journal of zoology, 199(4), 535-544.

471   Hagen, D. W., and D. M. Blouw. 1983. Heritability of dorsal spines in the fourspine stickleback

472   (*Apeltes quadracus*). *Heredity*, 50(3):275-281.

473   Haines, G. E., Moisan, L., Derry, A. M., & Hendry, A. P. (2023). Dimensionality and modularity

474   of adaptive variation: Divergence in threespine stickleback from diverse environments.

475   *The American Naturalist*, 201(2), 175-199.

476   Hall, M. F. 1956. A comparative study of the reproductive behaviour of the sticklebacks

477   (Gasterosteidae). Ph.D. Thesis. St. Hugh's College, Oxford. 370 p.

478   Hatfield, T. 1997. Genetic divergence in adaptive characters between sympatric species of

479   stickleback. *The American Naturalist*, 149(6):1009-1029.

Hatfield, T., and D. Schluter. 1999. Ecological speciation in sticklebacks:

environment☐dependent hybrid fitness. *Evolution*, *53*(3):866-873.

Hendry, A. P., C. L. Peichel, B. Matthews, J. W. Boughman, and P. Nosil. 2013. Stickleback

research: the now and the next. *Evolutionary Ecology Research*, *15*(2):111-141.

Hoogland, R., D. Morris, and N. Tinbergen. 1957. The spines of sticklebacks (*Gasterosteus* and

Pygosteus) as means of defence against predators (*Perca* and *Esox*). *Behaviour*, 10:205-

236.

Ivanova N. V., J. R. deWaard, and P. D. N. Hebert. 2006. An inexpensive, automation-friendly

protocol for recovering high-quality DNA. *Molecular Ecology Notes*, 6:998-1002.

Jeffries, D. L., J. A. Mee, and C. L. Peichel. 2022. Identification of a candidate sex determination

gene in Culaea inconstans suggests convergent recruitment of an Amh duplicate in two

lineages of stickleback. *Journal of Evolutionary Biology*, 35(12):1683-1695.

Kaeuffer, R., C. L. Peichel, D. I. Bolnick, and A. P. Hendry. 2012. Parallel and nonparallel

aspects of ecological, phenotypic, and genetic divergence across replicate population

pairs of lake and stream stickleback. *Evolution*, *66*(2):402-418.

Klepaker, T., K. Østbye, and M. A. Bell. 2013. Regressive evolution of the pelvic complex in

stickleback fishes: a study of convergent evolution. *Evolutionary Ecology Research,*

15(4):413-435.

Lande, R. 1975. The maintenance of genetic variability by mutation in a polygenic character

with linked loci. *Genetics Research*, *26*(3):221-235.

Lescak, E. A., and F. A. von Hippel. 2011. Selective predation of threespine stickleback by

rainbow trout. *Ecology of Freshwater Fish*, 20(2):308-314.

Lescak, E. A., F. A. von Hippel, B. K. Lohman, and M. L. Sherbick. 2012. Predation of

threespines stickleback by dragonfly naiads. *Ecology of Freshwater Fish*, 21(4):581-587.

Love, M. I., W. Huber, and S. Anders. 2014. Moderated estimation of fold change and dispersion

for RNA-seq data with DESeq2. *Genome Biology*, 15(12):550.

Lowey, J., Q. Cheng, S. M. Rogers, and J. A. Mee. 2020. Persistence of pelvic spine

polymorphism in a panmictic population of brook stickleback (*Culaea inconstans*) in

Alberta, Canada. *Canadian Journal of Zoology*, 98(10):643-649.

Marchinko, K. B. 2009. Predation's role in repeated phenotypic and genetic divergence of armor

in threespine stickleback. *Evolution*, 63(1):127-138.

Martin, M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads.

EMBnet.Journal, 17(1):10-12.

Matthews, B., K. B. Marchinko, D. I. Bolnick, and A. Mazumder. 2010. Specialization of trophic

position and habitat use by sticklebacks in an adaptive radiation. *Ecology*, *91*(4):1025-

1034.

McMurdie, P. J., and S. Holmes. 2013. phyloseq: an R package for reproducible interactive

analysis and graphics of microbiome census data. *PloS One*, *8*(4):e61217.

McPhail, J. D. 1984. Ecology and evolution of sympatric sticklebacks (Gasterosteus):

morphological and genetic evidence for a species pair in Enos Lake, British Columbia.

Canadian Journal of Zoology, *62*(7):1402-1408.

Mee, J. A., Bernatchez, L., Reist, J. D., Rogers, S. M., & Taylor, E. B. (2015). Identifying

designatable units for intraspecific conservation prioritization: a hierarchical approach

applied to the lake whitefish species complex (*Coregonus* spp.). *Evolutionary*

Applications, *8*(5), 423-441.

Menon, P. S. (1969). Population ecology of Gammarus lacustris sars in Big Island Lake. I

Habitat preference and relative abundance. *Hydrobiologia*, 33, 14-32.

Miller, S. E., D. Metcalf, and D. Schluter. 2015. Intraguild predation leads to genetically based

character shifts in the threespine stickleback. *Evolution*, *69*(12):3194-3203.

Miller, S.E., M. Barrueto, and D. Schluter. 2017. A comparative analysis of experimental

selection on the stickleback pelvis. *Journal of Evolutionary Biology*, 30(6):1165-1176.

Nagel, L., and D. Schluter. 1998. Body size, natural selection, and speciation in sticklebacks.

Evolution, *52*(1):209-218.

Nelson, J. S. 1969. Geographic variation in the brook stickleback, *Culaea inconstans*, and notes

on nomenclature and distribution. *Journal of the Fisheries Board of Canada*, 26(9):2431-

2447.

Nelson, J. S. 1977. Evidence of a genetic basis for absence of the pelvic skeleton in brook

stickleback, *Culaea inconstans*, and notes on the geographical distribution and origin of

the loss. *Journal of the Fisheries Board of Canada*, 34(9):1314-1320.

Nelson, J. S., and M. J. Paetz. 1992. *The fishes of Alberta* (pp. 319–323). The University of

Alberta Press, Edmonton, AB, Canada.

Oksanen J, Simpson G, Blanchet F, Kindt R, Legendre P, Minchin P, O'Hara R, Solymos P,

Stevens M, Szoecs E, Wagner H, Barbour M, Bedward M, Bolker B, Borcard D,

Carvalho G, Chirico M, De Caceres M, Durand S, Evangelista H, FitzJohn R, Friendly

M, Furneaux B, Hannigan G, Hill M, Lahti L, McGlinn D, Ouellette M, Ribeiro Cunha E,

Smith T, Stier A, Ter Braak C, Weedon J (2022). _vegan: Community Ecology

Package_. R package version 2.6-4, <https://CRAN.R-project.org/package=vegan>.

Peichel, C. L., K. S. Nereng, K. A. Ohgi, B. L. Cole, P. F. Colosimo, C. A. Buerkle, D. Schluter,

and D. M. Kingsley. 2001. The genetic architecture of divergence between threespine

stickleback species. *Nature*, *414*(6866):901-905.

Punzalan, D., F.H. Rodd, and K. A. Hughes. 2005. Perceptual processes and the maintenance of

polymorphism through frequency-dependent predation. *Evolutionary Ecology*, 19(3):303-

320.

R Core Team. 2022. R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Rainey, P. B., A. Buckling, R. Kassen, and M. Travisano. 2000. The emergence and maintenance

of diversity: insights from experimental bacterial populations. *Trends in Ecology and*

Evolution, 15(6):243-247.

Reid, K., M. A. Bell, and K. R. Veeramah. 2021. Threespine stickleback: a model system for

evolutionary genomics. *Annual Review of Genomics and Human Genetics*, *22*:357-383.

Reimchen, T. E. 1980. Spine deficiency and polymorphism in a population of *Gasterosteus*

aculeatus: an adaptation to predators? *Canadian Journal of Zoology*, 58(7): 1232-1244.

Reimchen. T. E. 1992. Injuries on stickleback from attacks by a toothed predator

(*Onchorhynchus*) and implications for the evolution of lateral plates. *Evolution*,

46(4):1224-1230.

Reimchen, T. E. 1994. Predators and morphological evolution in threespine stickleback. *in* The

evolutionary biology of the threespine stickleback. *Edited by* Bell, M.A., and S.A. Foster.

Oxford University Press, New York, NY, pp. 240-276.

Reimchen, T. E., D. Steeves, and C. A. Bergstrom. 2016. Sex matters for defence and trophic

traits of threespine stickleback. *Evolutionary Ecology Research*, 17(4):459-485.

Reisman, H. M., and T. J. Cade. 1967. Physiological and behavioral aspects of reproduction in

571   the brook stickleback, *Culaea inconstans*. *American Midland Naturalist*, 77:275-295.

572   Reist, J. D. 1980a. Selective predation upon pelvic phenotypes of brook stickleback, *Culaea*

573   *inconstans*, by northern pike, *Esox lucius*. *Canadian Journal of Zoology*, 58(7):1245-

574   1252.

575   Reist, J. D. 1980b. Predation upon pelvic phenotypes of brook stickleback, *Culaea inconstans*,

576   by selected invertebrates. *Canadian Journal of Zoology*, 58(7):1253-1258.

577   Reist, J. D. 1983. Behavioural variation in pelvic phenotypes of brook stickleback, *Culaea*

578   *inconstans*, in response to predation by northern pike, *Esox lucius*. *Environmental*

579   *Biology of Fishes*, 8(3-4):255-267.

580   Rennison, D. J., S. M. Rudman, and D. Schluter. 2019a. Parallel changes in gut microbiome

581   composition and function during colonization, local adaptation and ecological speciation.

582   *Proceedings of the Royal Society B*, 286(1916):20191911.

583   Rennison, D. J., Y. E. Stuart, D. I. Bolnick, and C. L. Peichel. 2019b. Ecological factors and

584   morphological traits are associated with repeated genomic differentiation between lake

585   and stream stickleback. *Philosophical Transactions of the Royal Society*

586   *B*, *374*(1777):20180241.

587   Ridgway, M. S., and J. D. McPhail. 1984. Ecology and evolution of sympatric sticklebacks

588   (*Gasterosteus*): mate choice and reproductive isolation in the Enos Lake species pair.

589   *Canadian Journal of Zoology*, *62*(9):1813-1818.

590   Rundle, H. D., S. M. Vamosi, and D. Schluter. 2003. Experimental test of predation's effect on

591   divergent selection during character displacement in sticklebacks. *Proceedings of the*

592   *National Academy of Sciences*, 100(25):14943-14948.

Rundle, H. D., and D. Schluter. 2004. Natural selection and ecological speciation in sticklebacks.

Adaptive Speciation, *19*(3):192-209.

Schluter, D. 1993. Adaptive radiation in sticklebacks: size, shape, and habitat use efficiency.

Ecology, 74(3):699-709.

Schluter, D. 2000. *The Ecology of Adaptive Radiation*. Oxford University Press, Oxford.

Schluter, D., and J. D. McPhail. 1992. Ecological character displacement and speciation in

sticklebacks. *The American Naturalist*, 140(1):85-108.

Schluter, D., K. B. Marchinko, M. E. Arnegard, H. Zhang, S. D. Brady, F. C. Jones, M. A. Bell,

and D. M. Kingsley. 2021. Fitness maps to a large-effect locus in introduced stickleback

populations. *Proceedings of the National Academy of Sciences*, 118(3):e1914889118.

Shapiro, M. D., B. R. Summers, S. Balabhadra, J. T. Aldenhoven, A. L. Miller, C. B.

Cunningham, M. A. Bell, and D. M. Kingsley. 2009. The genetic architecture of skeletal

convergence and sex determination in ninespine sticklebacks. *Current Biology*,

19(13):1140-1145.

Shikano, T., V. N. Laine, G. Herczeg, J. Vilkki, and J. Merilä. 2013. Genetic architecture of

parallel pelvic reduction in ninespine sticklebacks. *G3: Genes, Genomes, Genetics*,

3(10):1833-1842.

Spence, R., Wootton, R. J., Barber, I., Przybylski, M., & Smith, C. (2013). Ecological causes of

morphological evolution in the three□spined stickleback. *Ecology and Evolution*, 3(6),

1717-1726.

Takahata, N., Y. Satta, and J. Klein. 1992. Polymorphism and balancing selection at major

histocompatibility complex loci. *Genetics* 130(4):925-938.

Vamosi, S. M. 2002. Predation sharpens the adaptive peaks: survival trade-offs in sympatric

sticklebacks. *Annales Zoologici Fennici*, 39(3):237-248.

Vamosi, S. M., and D. Schluter. 2004. Character shifts in the defensive armor of sympatric

sticklebacks. *Evolution*, 58(2):376-385.

Varadharajan, S., P. Rastas, A. Löytynoja, M. Matschiner, F. C. F. Calboli, B. Guo, A. J.

Nederbragt, K. S. Jakobsen, and J. Merilä. 2019. A high-quality assembly of the nine-

spined stickleback (*Pungitius pungitius*) genome. *Genome Biology and Evolution*,

11:3291–3308.

Willerth, K., E. Franks, and J. A. Mee. 2022. Parallel and non-parallel divergence within

polymorphic populations of brook stickleback, *Culaea inconstans* (Actinopterygii:

Gasterosteidae). *Biological Journal of the Linnean Society*, 135(1):159-173.

Table 1. Summary of samples analyzed in this study. Numbers in parentheses indicate the

samples used in the gut metabarcoding analysis.

	Pelvic Phenotype		
	Spined	Vestigial	Unspined
2017			
Muir Lake			
female	19	1	25
male	22	1	18
Shunda Lake			
female	20	22	6
male	25	8	3
2019			
Muir Lake			
female	3 (3)	0	7 (7)
male	25 (18)	1	27 (18)
Shunda Lake			
female	13 (13)	12 (10)	3 (3)
male	17 (12)	10 (9)	2 (2)

631   **Table 2**. Results of a PERMANOVA test of diet differences

	df	Sum of Squares	F	p-value
Main effects:				
Lake	1	0.648	1.549	0.074
Pelvic reduction	1	0.902	2.154	0.008
Sex	1	0.614	1.466	0.094
Body length	1	0.923	2.205	0.008
Interaction effects:				
Lake x Body length	1	0.49	1.172	0.256
Pelvic reduction x Body length	1	0.649	1.550	0.075
Sex x Body length	1	0.271	0.647	0.877
Residual	86	35.998		
Total	93	40.495		

Table 3. ANOVA table for the generalized linear model of the effects of pelvic reduction (spined

or pelvic-reduced), lake (Muir Lake or Shunda Lake), sex, sampling year (2017 or 2018), and

body length (total length in mm) on maximum gill raker length in brook stickleback. We

included all two-way interactions in our initial models, and then used a reverse-stepwise

approach for model selection, removing any interaction terms that were not significant. Two-way

interaction terms that were not significant ($p > 0.05$) were not included in the final model shown

here.

	Sum of Squares	df	*F*	p-value
Main effects:				
Pelvic reduction	5109	1	0.9238	0.337
Lake	112007	1	20.2544	< 0.001
Sex	84370	1	15.2569	< 0.001
Year	16909	1	3.0578	0.081
Body length	417938	1	75.5767	< 0.001
Interaction effects:				
Lake x body length	55977	1	10.1224	0.002
Sex x body length	109862	1	19.8665	< 0.001
Year x body length	28198	1	5.099	0.025
Residual	1548394	280		
Total	2378764			

Table 4. ANOVA table for the generalized linear model of the effects of pelvic reduction (spined

or pelvic-reduced), lake (Muir Lake or Shunda Lake), sex, sampling year (2017 or 2018), and

body length (total length in mm) on mean number of gill rakers in brook stickleback. We

included all two-way interactions in our initial models, and then used a reverse-stepwise

approach for model selection, removing any interaction terms that were not significant.

Interaction terms were not significant ($p > 0.05$) and were not included in the final model shown

here.

	Sum of Squares	df	F	p-value
Pelvic reduction	0.64	1	0.5474	0.460
Lake	5.65	1	4.8253	0.029
Sex	1.52	1	1.3004	0.255
Year	2.05	1	1.7528	0.187
Body length	29.06	1	24.8229	< 0.001
Residual	331.28	283		
Total	370.20			

Figure 1. Ordination plot of Non-metric Multidimensional Scaling (NMDS) axes based on

sample-wise Bray-Curtis distances in brook stickleback diet (with diet items classified to the

level of family). Ellipses show the 95% probability interval for each group, assuming a normal

distribution of points. The larger points indicate the center of each ellipse.

Figure 2. Log$_2$ fold change between Shunda Lake and Muir Lake in the abundance of eukaryotic

families in the diets of brook stickleback. Zero indicates no difference in abundance between

lakes, positive numbers indicate higher abundance in Shunda Lake, and negative numbers

indicate higher abundance in Muir Lake. Red symbols indicate a significant difference in

abundance with an adjusted p-value < 0.05 corrected for multiple testing. Families are arranged

(left to right on the x-axis) in order of highest to lowest fold change (in an absolute sense)

between lakes.

Figure 3. Log$_2$ fold change in the abundance of eukaryotic families in the diets of brook

stickleback in Muir Lake and Shunda Lake with length as a co-factor. Points show the change in

abundance between spined and pelvic-reduced individuals, "S" symbols indicate change between

sexes, and "L" symbols indicate change attributed to fish length. A log$_2$ fold change of zero

indicates no difference in abundance, whereas positive values indicate higher abundance in

spined individuals, in males, or in longer individuals. Red symbols indicate a significant

difference in abundance with an adjusted p-value < 0.05 corrected for multiple testing. Two-way

interactions with length were included in the statistical model but, for clarity, are not included in

this visualization. Families are arranged (left to right on the x-axis) in order of highest to lowest

fold change (in an absolute sense) between pelvic morphotypes in each lake. Histograms of

Log_{10} abundances across all sampled individuals for each diet item are shown above the fold-

change points.

Figure 4. Log_2 fold change in the abundance of eukaryotic families in the diets of brook

stickleback in Muir Lake with habitat as a co-factor. Points show the change in abundance

between spined and pelvic-reduced individuals, "S" symbols indicate change between sexes, and

"H" symbols indicate change attributed to habitat. A log_2 fold change of zero indicates no

difference in abundance, whereas positive values indicate higher abundance in spined

individuals, in males, or in the littoral habitat. Red symbols indicate a significant difference in

abundance with an adjusted p-value < 0.05 corrected for multiple testing. Families are arranged

(left to right on the x-axis) in order of highest to lowest fold change (in an absolute sense)

between pelvic morphotypes in Muir lake. Histograms of Log_{10} abundances across all sampled

individuals for each diet item are shown above the fold-change points.

Figure 5. Comparison of maximum gill raker lengths between spined and pelvic-reduced brook

stickleback in Muir Lake and Shunda Lake. The horizontal span of each line indicates the range

of fish lengths for each group in the comparison (i.e. male spined, female spined, male reduced,

and female reduced). For this visualization, only the effects included in the final statistical model

(Table 3) are depicted (although, for clarity, the effect of year of sampling is not shown).

Figure 6. Comparison of mean number of gill rakers between spined and pelvic-reduced brook

stickleback in Muir Lake and Shunda Lake. The horizontal span of each line indicates the range

of fish lengths for each group in the comparison (i.e. male spined, female spined, male reduced,

and female reduced). For this visualization, only the effects included in the final statistical model

(Table 4) are depicted (although, for clarity, the effect of year of sampling is not shown).

706