Chemical-guided SHAPE sequencing (cgSHAPE-seq) informs the binding site of RNA-degrading chimeras targeting SARS-CoV-2 5' untranslated region

Zhichao Tang[1,3], Shalakha Hegde[1,3], Siyuan Hao[2], Manikandan Selvaraju[1], Jianming Qiu[2], Jingxin Wang*[1]

[1]Department of Medicinal Chemistry, University of Kansas, Lawrence, KS, USA.

[2]Department of Microbiology, Molecular Genetics and Immunology, University of Kansas Medical Center, Kansas City, KS, USA.

[3]These authors contributed equally.

*Email: wang.jingxin@ku.edu

One of the hallmarks of RNA viruses is highly structured untranslated regions (UTRs) in their genomes. These conserved RNA structures are often essential for viral replication, transcription, or translation. In this report, we discovered and optimized a new coumarin derivative **C30** that binds to a four-way RNA helix called SL5 in the 5' UTR of the SARS-CoV-2 RNA genome. To locate the binding site, we developed a novel sequencing-based method namely cgSHAPE-seq, in which the acylating chemical probe was directed to crosslink with the 2'-OH groups of ribose at the ligand binding site. This crosslinked RNA could then create read-through mutations during reverse transcription (i.e., primer extension) at single-nucleotide resolution to uncover the acylation locations. cgSHAPE-seq unambiguously determined that a bulged G in SL5 was the primary binding site of **C30** in the SARS-CoV-2 5' UTR, which was validated through mutagenesis and in vitro binding experiments. **C30** was further used as a warhead in RNA-degrading chimeras (RIBOTACs) to reduce viral RNA expression levels. We demonstrated that replacing the acylating moiety in the cgSHAPE probe with ribonuclease L recruiter (RLR) moieties yielded RNA degraders active in the in vitro RNase L degradation assay and SARS-CoV-2 5' UTR expressing cells. We further explored another RLR conjugation site on the E ring of **C30** and discovered potent activity in vitro and in cells. The optimized RIBOTAC **C64** inhibited live virus replication in lung epithelial carcinoma cells.

RNA viruses usually have highly structured 5' and 3' UTRs in their RNA genome, which can potentially serve as therapeutic targets[1]. In this report, we used SARS-CoV-2 as a specific test-case example and explore to use RNA-degrading chimeras to inhibit virus replication. SARS-CoV-2 is an enveloped ssRNA(+) virus. The whole genome of SARS-CoV-2 (~30,000 nucleotides) is encoded in a single RNA molecule[2]. The viral RNA in transmitted virions is 5' capped and 3' polyadenylated, so that it is recognized and treated as an mRNA[3]. In this step, the 5' UTR is used to hijack the host ribosome to translate viral proteins[4]. Furthermore, the 5'

UTR plays an essential role in RNA transcription for each coronavirus structural protein, which
is accomplished through a "discontinuous" transcription mechanism. Specifically, the replication
transcription complex binds to the 5' UTR leader transcriptional regulatory sequences (TRS-L),
and then "hops" onto the body TRS (TRS-B) sequence located at the 5'-end of each structural
gene[5,6]. That said, all SARS-CoV-2 transcripts share the same 5' UTR leader sequence. In
addition, the SARS-CoV-2 5' UTR was reported essential for viral RNA packaging[7].

Given the importance of the UTRs in SARS-CoV-2, we and others elucidated the RNA
structures in SARS-CoV-2 UTRs[8–12]. The 5' UTR RNA structures in cell-free buffers, in virus-
infected cells, and in our reporter cell model are highly consistent [8–13], suggesting superior
stability and suitability serving as drug targets. The 5' UTR of SARS-CoV-2 contains five stem-
loops, namely SL1–5. The start codon resides in SL5, a unique four-way helix[8–12] (Fig. 1a). SL5
exists in all betacoronavirus species, including MERS and SARS-CoV, and the shapes of this
RNA structure are similar[9]. We aligned the SARS-CoV-2 RefSeq and different lineages and
demonstrated that the SL5 is highly conserved among all strains[1]. Although a predominant
mutation was found in SL5B loop region from recent lineages (C241T), this mutation is unlikely
to change the overall structure of SL5[1]. In SARS-CoV-2, several structures in the 5' UTR,
including SL4, SL5A, and SL6, were found binding to amilorides[14] (Fig. 1a). Amilorides
demonstrated antiviral activity in SARS-CoV-2 infected cells.

Here, we report a pipeline in antiviral discovery and optimization of RNA-degrading chimeras
targeting SL5, highlighting a novel sequencing-based method namely chemical-guided (cg)
selective 2′-hydroxyl acylation analyzed by primer extension (SHAPE) sequencing (seq), or
cgSHAPE-seq, to rapidly locate the RNA ligand binding site (Fig. 1a). First, we screened a small
coumarin derivative library in SL5 RNA binding assay and optimized the hit through structure-
activity-relationship studies. To elucidate the RNA ligand binding site, we synthesized and
applied a new type of chemical probe that can selectively acylate the 2'-OH on the ribose at the
location of binding (Fig. 1b)[26,27]. The 2'-OH acylation locations were "recorded" onto RNA
molecules by reverse transcriptase as single-point mutations at the modification sites during
primer extension. The mutation sites were then captured and deconvoluted by next-generation
sequencing[28,29]. Mutational profiling analysis in cgSHAPE-seq unambiguously identified a
bulged G in SL5 as the primary binding site in the SARS-CoV-2 5' UTR. In the literature, other
sequencing-based methods were reported using affinity probes bearing nitrogen mustard or
diazirine moiety (e.g., ChemCLIP-seq[15–19] and PEARL-seq[20]). However, a major limitation of
these methods is a strong labeling bias toward guanosines[21]. Similar to the SHAPE, cgSHAPE-
seq reacts with 2'-OH of A, U, G, or C at similar rates. This can potentially increase the scope
and accuracy of proximity-induced chemical reactions on RNAs for mapping purposes.

We further developed RNA-degrading chimeras by replacing the 2'-OH acylating moiety with
RNase L recruiter (RLR) moieties on the cgSHAPE probe, as well as by conjugating the RLR
moieties on other putative solvent accessible site on the RNA ligand. RNA-degrading chimera
utilizing endogenous ribonuclease (RNase) L was first reported by the Silverman group in
1993[22]. Recently, the modality of RNA-degrading chimera was further developed by the Disney
group and was demonstrated to be active using small-molecule RNA ligands[23–30] (Fig. 1c). The
Disney group also coined the name ribonuclease targeting chimera or RIBOTAC for this type of

RNA degraders. RIBOTACs were shown efficacious to degrade microRNA in cells[24,25] and
mouse models[26]. Importantly, a small-molecule RIBOTAC was recently used to degrade the
SARS-CoV-2 RNA genome by targeting an RNA structure named attenuator hairpin near the
programmed frameshift (PFS) regulatory element[23]. The viral RNA transcript level was shown to
be reduced in a model cell system by ~50% at 8 μM RIBOTAC[23]. Antisense-based RNA-
degrading chimeras targeting spike-protein or envelope-protein encoding RNAs were also
demonstrated efficacy in SARS-CoV-2 infected cells[28]. Our optimized RIBOTAC robustly
degraded SARS-CoV-2 RNA in cellular models at 1 μM and inhibited virus replication at 20 μM
in lung epithelial cells. No significant toxicity was observed. Interestingly, we discovered the
natural RNase L binding moiety is similar or less active than the synthetic RLR in the RIBOTAC
modality.

91
Fig. 1 | Development of cgSHAPE-seq and anti-viral RIBOTAC. a, RNA secondary structures in SARS-CoV-2 5'
UTR and the pipeline in identification of the ligand binding site and the development of RNA-degrading chimeras. **b**,
Principle of cgSHAPE-seq for identifying small molecule binding sites in four steps. Step 1: Synthesis of FAI conjugated
chemical probe. Step 2: Chemical-guided acylation at the 2'-OH of ribose at the binding site. Step 3: Reverse
transcription in the presence of Mn2+ creates single-point mutations at the acylation site. Step 4: Mutational profiling
and quantification identify the putative binding sites (figure adapted from Figdraw). **c**, RNA-degrading chimeras
(RIBOTACs) recruit RNase L at the target RNA to degrade viral RNAs.

Results

Chemical optimization of the coumarin derivatives for SL5 RNA binding

We previously synthesized a collection of coumarin derivatives that are known to bind to
RNAs[31]. Each of these coumarin derivatives is fluorescent (excitation/emission = 400/480 nm),
enabling us to use fluorescence polarization (FP) to determine the in vitro binding affinity with
the RNA receptors. We in vitro transcribed SL5 RNA (144–303, RefSeq NC_045512) and
screened the compound library using the FP assay (Extended Data Fig. 1). **C2** binds to SL5 at a
dissociation constant (K_d) of 1.45 µM (Table 1). Elimination of the ethyl substituent on the A ring
(**C4**) did not improve the binding affinity to SL5 RNA (Table 1). In contrast, a bulky substituent
on the A ring (**C6**) impeded the interaction with SL5 (Table 1). This indicated that a non-
sterically hindered linker might be suitable for conjugation on ring A without affecting the ligand
binding affinity to SL5 RNA. We found **C29** the best ligand in our compound collection, which
binds to SL5 RNA at a K_d of 0.47 µM. Comparing to **C4**, **C29** has a different set of substituents
on ring E, and therefore, we further investigated ring E while keep ring A as unsubstituted
piperazine (Table 1). In vitro FP assay showed that a fluorinated analog, **C30**, further improves
the binding affinity (K_d = 0.22 µM). Changing the F group into Cl (**C32**) or CF_3 (**C36**) groups or
alternating the fluorinated site (**C31**) all reduced the binding affinities (Table 1). To our
knowledge, **C30** is the strongest small-molecule ligand to the SARS-CoV-2 SL5 RNA.

Table 1. Structure-activity-relationship study of coumarin derivatives for in vitro binding affinities with SL5
RNA.

Name	R^1	R^2	R^3	SL5 K_d (µM)
C2	CH$_2$CH$_3$	NA	NA	1.45 ± 0.38
C4	H	NA	NA	1.24 ± 0.29
C6	CH$_2$CH$_2$NHBoc	NA	NA	5.67 ± 4.78
C29	NA	H	H	0.47 ± 0.09
C30	NA	H	F	0.22 ± 0.09
C31	NA	F	H	0.69 ± 0.07
C32	NA	H	Cl	0.42 ± 0.09
C36	NA	H	CF$_3$	1.41 ± 0.25

cgSHAPE-seq uncovered the bulged G in SL5 as the C30 binding site
We were inspired by a commonly used method for RNA structure elucidation called SHAPE and
sought to develop a new chemical-guided sequencing-based method to identify the binding site
of **C30** in SL5 RNA. We named the new chemical probing method chemical-guided SHAPE
sequencing, or cgSHAPE-seq. Conventional SHAPE uses electrophilic reagents that can form
ester adducts on the 2'-OH of the ribose. The unpaired nucleotides have higher accessibility for
acylation reactions, which is the basis of structure-based differential acylation activity.
Therefore, identification of the acylation site would provide information in RNA base-pairing in

the conventional SHAPE. Importantly, the electrophile-ribose adduct can create a mutation
during reverse transcription (i.e., primer extension). As a result, conventional SHAPE coupled
with quantitative mutational profiling has become a gold standard method to explore RNA
topology in vitro and in vivo[32–36]. The key advantage of SHAPE is that the electrophile can
usually react with all four nucleotides (A, U, G, or C) with similar activity. We wondered if the
ribose acylation could be repurposed for identification of small-molecule binding sites by
covalently linking electrophile moieties to RNA-binding chemical ligands.

First, we selected furoyl acylimidazole (FAI) as the electrophile for synthesizing the chemical
probe. Compared to other electrophile moieties such as anhydride and acyl cyanide, FAI is
more resistant to hydrolysis with a half-life of 73 min in H_2O[37,38]. The hydrolysis-resistant probe
design renders the synthesis and storage less demanding for anhydrous experimental facilities.
An azide group on the furoyl moiety was used to provide a click-chemistry handle to conjugate
with **C30** (Fig. 2a). After the [3+2] cycloaddition reaction, the carboxylic acid was then converted
into an acyl imidazole moiety under a mild condition, finalizing the synthesis of **C30-FAI** probe
(Fig. 2a). The acyl imidazole probe was prepared freshly and used directly without further
purification (see Method). We tested the reactivity of FAI-N_3 with a denatured RNA and
confirmed that FAI-N_3 can react with all four nucleotides and generate SHAPE signal at a high
concentration (Extended Data Fig. 2).

Next, we applied **C30-FAI** in SARS-CoV-2 5' UTR RNA to elucidate the binding site of **C30**. It is
known that conventional SHAPE experiments usually require high concentration (10–100 mM)
of acyl imidazole (e.g., FAI) for ribose acylation. To avoid obtaining structure-based differential
acylation activity caused by FAI moiety alone, we chose to use a much lower dose of the
chemical probe for cgSHAPE-seq. We reasoned that at low concentration of the probe (0.02–1
mM), the differential acylation activity would be predominantly caused by ligand binding (i.e.,
proximity-promoted acylation). Briefly, the total RNA was extracted from SARS-CoV-2 5' UTR
expressing cells (see Method) and refolded in buffer. **C30-FAI**, FAI-N_3, or DMSO was
individually reacted with the folded RNA for 15 min at 37 °C. After the reaction, we used
ProtoScript II reverse transcriptase (New England Biolabs) in the presence of $MnCl_2$ (3 mM) for
primer extension using a protocol modified from the literature report (see Method)[32]. The cDNA
was then amplified by PCR in the SARS-CoV-2 5' UTR region and the resulting amplicon was
subsequently sequenced. We applied an existing software package ShapeMapper2 developed
by the Weeks group for mutational profiling analysis[32,39]. We calculated the background
Δmutation rate (FAI-N_3 – DMSO) for each nucleotide and pleasingly observed a low background
signal at 0.02 and 0.1 mM of FAI-N_3, indicating that the structure-based differential acylation
activity is negligible in cgSHAPE-seq at these concentrations (Extended Data Fig. 3). We then
calculated the RNA ligand-induced ΔΔmutation rate [(C30-FAI – DMSO) – (FAI-N_3 – DMSO)] for
each nucleotide and identified G174 as the only significantly mutated nucleotides in 0.1 mM
probe-treated samples (Fig. 2b,c). In 1 mM probe treated samples, a lower signal-to-noise ratio
was observed even though the mutational result also implied the G174 as the primary **C30**
binding site (Fig. 2d). At 1 mM, FAI-N3 also significantly increased the Δmutation rate at G174,
implying the contribution of structure-based SHAPE activities started to emerge (Fig. 2d).
Mapping the nucleotide with previously identified secondary structures uncovered that G174 is a
single-nucleotide bulge in the SL5 stem region[8–12].

We then validated the **C30** binding site in SL5 by testing individual substructures of the SL5
RNA. The loop region of SL5A, SL5B, and a minimized four-helix junction, named SL5^M
(containing shorter stems) were synthesized chemically or enzymatically (Fig. 3a). The in vitro
binding results demonstrated that only SL5^M retained similar binding affinity to **C30** (Fig. 3a,
Extended Data Fig. 4). To further validate the putative binding site G174 in SL5^M, we designed
and synthesized SL5^M RNAs with different mutations that disrupt the bulged G or other RNA
structures (Fig. 3b). As expected, deletion of G174 or base-pairing G174 with an additional C
both resulted in a 7-fold decrease in binding affinity to **C30**. Replacement of G174 with A, C or U
also significantly reduced **C30** binding. We also expanded the bulged G by inserting different
nucleotides between C173 and U175 or between A270 and G271, all these mutated RNAs were
demonstrated 5–6-fold reduced binding affinity to **C30**. These results suggested the importance
of a single bulged G in accommodating C30's binding. Changing the closing U-A base pair into
C-G (3'-end of G174) or C-G base pair into U-A (5'-end of G174) also resulted in a 4-fold
decreased binding. Mutations on other parts of the RNA has less impact (i.e., within 2-fold) in
changing the binding affinity to **C30** (Fig. 3b). Altogether, these observations validated that the
bulged G region is the primarily binding site in SL5 RNA for **C30**. We concluded that cgSHAPE-
seq is a validated method for identifying the binding site of RNA-binding small molecules.

Fig. 3 | Validation of the binding site. a, Structural fragments of SL5 and their binding affinities to C30. b, SL5^M
mutants and their binding affinities to C30.

Comparison of two RNase L recruiting moieties in RIBOTACs

We then conjugated **C30** with RLRs to synthesize SL5-targeting RNA degraders (RIBOTACs).
Two different conjugation sites were used on rings A and E of **C30**, respectively (Fig. 4a). For
RLR conjugation on the A ring, the acylating moiety in the above cgSHAPE probe (**C30-FAI**)
was replaced with RLR moieties. We notice that in most reported co-NMR structures of RNA
bulges and small molecules that are similar to **C30**, both ends of the small molecules are
solvent accessible[40–42]. Therefore, ring E was also explored for conjugation to RLRs. We
replaced the electronegative fluorine atom on ring E into an oxygen, which was further used for
RLR attachment (Fig. 4a).

For RLR moieties, the natural RNase L ligand 2'-5'-lined oligoadenylate (2-5A) and its synthetic
mimic **D1** were both previously reported to be used in RNA-degrading chimeras (Extended Data
Fig. 5)[23,24]. Combining the two conjugation sites and two RLR structures, we obtained four
RIBOTAC candidates, **C47**, **C48**, **C64**, and **C65**, for SL5 RNA degradation (Fig. 4a). We
validated that the polyethylene glycol (PEG) linker on 2-5A does not affect the activity in a
reported RNase L degradation assay with a 5' 6-fluorescein-tagged model RNA containing
multiple RNase L cleavage sites[43,44] (Extended Data Fig. 5a). It was demonstrated that the
binding affinity between RNase L and **D1** ($K_d ≈ 18$ µM) is 80,000-fold weaker than that observed
for 2-5A[43]. Consistent with this reported in vitro binding data, the synthetic **D1** alone is > 10,000
times weaker than 2-5A in the in vitro RNase L degradation assay (Extended Data Fig. 5b,c).

Next, we tested the four RIBOTACs in the RNase L degradation assay with purified SL5 RNA
and observed their activities in order: **C64** > **C47** ≈ **C48** > **C65** (Fig. 4b, Extended Data Fig. 6a).
To our surprise, the RIBOTAC **C64** with **D1** as the RLR moiety is much stronger than **C65** with
2-5A at 50 µM (Fig. 4b). This result is contrary to what we would have predicted based on the
activities of the RLR moieties per se. We validated these in vitro findings in SARS-CoV-2 5'

UTR expressing 293T cells. In this cell model, the SARS-CoV-2 5' UTR sequence was fused to
a CMV promoter-controlled Gaussia luciferase expression cassette (Fig. 4c; for sequences, see
Method). Consistent with the RNase L degradation assay result, the maximum potency of **C64**
(i.e., RNA reduction level) was significantly better than **C65** (Fig. 4c). The activities of **C47** and
C48 in this cell model are similar, between those of **C64** and **C65** (Extended Data Fig. 6b).

Fig. 4 | RNA degrading activity and anti-viral activity of C30-based RIBOTACs. a, Synthesis of **C30**-based
RIBOTACs using conjugation sites on rings A or E of **C30**. **b**, Comparison of two RLR moieties in the RIBOTAC
modality using the in vitro RNase L degradation assay with purified SL5 RNA. **c**, Cellular activity of RIBOTACs in
SARS-CoV-2 5' UTR expressing cells. **d**, Inhibitory effect of RIBOTAC **C64** in SARS-CoV-2 infected A549 cells. The
cytotoxicity of the compound was also evaluated. The dose-response curves are representative of three independent
measurements (N = 3).

Efficacy of RIBOTAC in live virus infection assay

Finally, we tested the activity of **C64** in SARS-CoV-2 infected cells. The SARS-CoV-2 virus was
engineered to include a Nano Luciferase (NLuc) reporter by fusing NLuc onto ORF7 of the
SARS-CoV-2 genome[45]. In this way, the NLuc signal is proportional to the viral protein copy
number in cells. We applied a human lung epithelial carcinoma cell line A549 expressing high
level of ACE2 as the host cell[45]. The cells were infected with the SARS-CoV-2-NLuc virus at a
multiplicity of infection (MOI) of 2.0 at 1 h before the treatment with RIBOTACs **C64** for 3 d. To
our satisfactory, **C64** showed > 95% inhibition at 20 µM (Fig. 4d). At the same concentration, no
major toxicity is observed in A549 cells (Fig. 4d).

Discussion

cgSHAPE-seq has several potential limitations in application. For example, the FAI-based
probes used in this report would not be compatible with strong nucleophilic RNA ligands[14] due
to self-reaction. In addition, as shown in conventional SHAPE, FAI moiety has a higher reactivity

towards unpaired RNA nucleotides. Although most of reported RNA ligands targets the unpaired
region[46,47], cgSHAPE-seq may be less reactive for ligands that binds to the double-stranded
RNA grooves. Finally, in our model denatured RNA, FAI-N_3 react with all four nucleotides.
However, the reactivity of FAI-N_3 and A, U, G, and C are not equal. In this model, we found FAI-
N_3 has a higher reactivity bias towards G and against U (Extended Data Fig. 2). For this reason,
the cgSHAPE activities on G and U might be over- and under-estimated, respectively.

In our cgSHAPE-seq result, apart from G174, we also observed a cluster of nucleotides from
A131 to G149 showing slightly higher mutation rate than others (Fig. 2c). This can be potentially
caused by the nonspecific binding of **C30** with flexible sequence[31]. RNA targeting strategies are
known to have off-target effects due to shallow binding sites on RNAs and relatively weak
binding affinity for small molecules. **C64** at 3 µM can cause 13 gene down regulation
(log2FoldChange < −2) and > 23 gene upregulation (log2FoldChange > 2) in the transcriptome
(Extended Data Fig. 7, Supplementary Table S1). The activity of the RIBOTAC might be
improved if a more potent and selective RNase L recruiter is used[48]. Specifically, we showed
that the natural RNase L recruiter/activator 2-5A, which is negatively charged, is sometimes not
compatible with the positively charged RNA binder **C30** (Fig. 4b). For this reason, new synthetic
RNase L recruiter should probably be considered to be neutral or positively charged as most of
the reported RNA ligands are also positively charged[49–51].

In summary, we developed a new generalizable chemical probing method called cgSHAPE-seq
for quickly identifying small molecule-RNA binding sites by sequencing. cgSHAPE probes react
with the 2'-OH groups on the ribose close to the binding sites with a mitigated dependency of
the nucleobase identity observed in other reported methods. We used cgSHAPE-seq to identify
a bulged G on SL5 as the primary binding site on the SARS-CoV-2 5' UTR targeted by the
newly discovered coumarin derivative **C30**. Finally, we developed a novel **C30**-based RNA
degrader (RIBOTAC) capable of degrading viral RNA transcripts in cells and inhibiting virus
replication in SARS-CoV-2 infected cells, offering crucial insights into RNA degrading chimeras'
design.

References
1.   Hegde, S., Tang, Z., Zhao, J. & Wang, J. Inhibition of SARS-CoV-2 by Targeting
Conserved Viral RNA Structures and Sequences. *Front. Chem.* **9**, 802766 (2021).

2.   Wu, F. *et al.* A new coronavirus associated with human respiratory disease in China.
Nature **579**, 265–269 (2020).

3.   Malone, B., Urakova, N., Snijder, E. J. & Campbell, E. A. Structures and functions of
coronavirus replication-transcription complexes and their relevance for SARS-CoV-2 drug
design. *Nat. Rev. Mol. Cell Biol.* **23**, 21–39 (2022).

4.   de Breyne, S. *et al.* Translational control of coronaviruses. *Nucleic Acids Research* vol.
48 12502–12522 (2020).

5.   Zúñiga, S., Sola, I., Alonso, S. & Enjuanes, L. Sequence motifs involved in the regulation
of discontinuous coronavirus subgenomic RNA synthesis. *J. Virol.* **78**, 980–94 (2004).

6.   Sola, I., Almazán, F., Zúñiga, S. & Enjuanes, L. Continuous and Discontinuous RNA
Synthesis in Coronaviruses. *http://dx.doi.org/10.1146/annurev-virology-100114-055218* **2**,
265–288 (2015).

7.   Escors, D., Izeta, A., Capiscol, C. & Enjuanes, L. Transmissible Gastroenteritis
Coronavirus Packaging Signal Is Located at the 5′ End of the Virus Genome. *J. Virol.* **77**,
7890–7902 (2003).

8.   Zhao, J., Qiu, J., Aryal, S., Hackett, J. L. & Wang, J. The RNA Architecture of the SARS-
CoV-2 3′-Untranslated Region. *Viruses* **12**, 1473 (2020).

9.   Sun, L. *et al.* In vivo structural characterization of the SARS-CoV-2 RNA genome
identifies host proteins vulnerable to repurposed drugs. *Cell* **184**, 1865-1883.e20 (2021).

10.   Huston, N. C. *et al.* Comprehensive in vivo secondary structure of the SARS-CoV-2
genome reveals novel regulatory motifs and mechanisms. *Mol. Cell* **81**, 584-598.e5
(2021).

11.   Manfredonia, I. *et al.* Genome-wide mapping of SARS-CoV-2 RNA structures identifies
therapeutically-relevant elements. *Nucleic Acids Res.* (2020) doi:10.1093/nar/gkaa1053.

12.   Sanders, W. *et al.* Comparative analysis of coronavirus genomic RNA structure reveals
conservation in SARS-like coronaviruses. *bioRxiv Prepr. Serv. Biol.* (2020)
doi:10.1101/2020.06.15.153197.

13.   Lan, T. C. T. *et al.* Structure of the full SARS-CoV-2 RNA genome in infected cells.
bioRxiv (2020) doi:10.1101/2020.06.29.178343.

14.   Zafferani, M. *et al.* Amilorides inhibit SARS-CoV-2 replication in vitro by targeting RNA
structures. *Sci. Adv.* **7**, eabl6096 (2021).

15.   Velagapudi, S. P. *et al.* Defining RNA-Small Molecule Affinity Landscapes Enables
Design of a Small Molecule Inhibitor of an Oncogenic Noncoding RNA. *ACS Cent. Sci.* **3**,
205–216 (2017).

16.   Suresh, B. M. *et al.* A general fragment-based approach to identify and optimize bioactive
ligands targeting RNA. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 33197–33203 (2020).

17.   Velagapudi, S. P. *et al.* Design of a small molecule against an oncogenic noncoding
RNA. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 5898–903 (2016).

18.   Velagapudi, S. P., Li, Y. & Disney, M. D. A cross-linking approach to map small molecule-
RNA binding sites in cells. *Bioorganic Med. Chem. Lett.* **29**, 1532–1536 (2019).

19.   Guan, L. & Disney, M. D. Covalent Small-Molecule-RNA Complex Formation Enables
Cellular Profiling of Small-Molecule-RNA Interactions. *Angew. Chemie Int. Ed.* **52**,
10010–10013 (2013).

20.   Mukherjee, H. *et al.* PEARL-seq: A Photoaffinity Platform for the Analysis of Small

Molecule-RNA Interactions. *ACS Chem. Biol.* **15**, 2374–2381 (2020).

21. Balaratnam, S. *et al.* A chemical probe based on the PreQ1 metabolite enables
transcriptome-wide mapping of binding sites. *Nat. Commun.* **12**, 5856 (2021).

22. Torrence, P. F. *et al.* Targeting RNA for degradation with a (2'-5')oligoadenylate-
antisense chimera. *Proc. Natl. Acad. Sci. U. S. A.* **90**, 1300–4 (1993).

23. Haniff, H. S. *et al.* Targeting the SARS-CoV-2 RNA Genome with Small Molecule Binders
and Ribonuclease Targeting Chimera (RIBOTAC) Degraders. *ACS Cent. Sci.* **6**, 1713–
1721 (2020).

24. Costales, M. G., Matsumoto, Y., Velagapudi, S. P. & Disney, M. D. Small Molecule
Targeted Recruitment of a Nuclease to RNA. *J. Am. Chem. Soc.* **140**, 6741–6744 (2018).

25. Costales, M. G., Suresh, B., Vishnu, K. & Disney, M. D. Targeted Degradation of a
Hypoxia-Associated Non-coding RNA Enhances the Selectivity of a Small Molecule
Interacting with RNA. *Cell Chem. Biol.* **26**, 1180-1186.e5 (2019).

26. Costales, M. G. *et al.* Small-molecule targeted recruitment of a nuclease to cleave an
oncogenic RNA in a mouse model of metastatic cancer. *Proc. Natl. Acad. Sci.* **117**, 2406–
2411 (2020).

27. Liu, X. *et al.* Targeted Degradation of the Oncogenic MicroRNA 17-92 Cluster by
Structure-Targeting Ligands. *J. Am. Chem. Soc.* **142**, 6970–6982 (2020).

28. Su, X. *et al.* Efficient Inhibition of SARS-CoV-2 Using Chimeric Antisense
Oligonucleotides through RNase L Activation*. *Angew. Chem. Int. Ed. Engl.* **60**, 21662–
21667 (2021).

29. Zhang, P. *et al.* Reprogramming of Protein-Targeted Small-Molecule Medicines to RNA
by Ribonuclease Recruitment. *J. Am. Chem. Soc.* **143**, 13044–13055 (2021).

30. Bush, J. A. *et al.* Ribonuclease recruitment using a small molecule reduced c9ALS/FTD
r(G4C2) repeat expansion in vitro and in vivo ALS models. *Sci. Transl. Med.* **13**,
eabd5991 (2021).

31. Tang, Z. *et al.* Recognition of single-stranded nucleic acids by small-molecule splicing
modulators. *Nucleic Acids Res.* **49**, 7870-7883. PMC8373063 (2021).

32. Smola, M. J., Rice, G. M., Busan, S., Siegfried, N. A. & Weeks, K. M. Selective 2 ' -
hydroxyl acylation analyzed by primer extension and mutational profiling (SHAPE-MaP)
for direct , versatile and accurate RNA structure analysis. *Nat. Protoc.* **10**, 1643–1669
(2015).

33. Weeks, K. M. & Mauger, D. M. Exploring RNA Structural Codes with SHAPE Chemistry.
Acc. Chem. Res. **44**, 1280–1291 (2011).

34. Siegfried, N. A., Busan, S., Rice, G. M., Nelson, J. A. E. & Weeks, K. M. RNA motif

discovery by SHAPE and mutational profiling (SHAPE-MaP). *Nat. Methods* **11**, 959
(2014).

35. Spitale, R. C. *et al.* Structural imprints in vivo decode RNA regulatory mechanisms.
Nature **519**, 486–490 (2015).

36. Flynn, R. A. *et al.* Transcriptome-wide interrogation of RNA secondary structure in living
cells with icSHAPE. *Nat. Protoc.* **11**, 273–290 (2016).

37. Spitale, R. C. *et al.* RNA SHAPE analysis in living cells. *Nat. Chem. Biol.* **9**, 18–20 (2013).

38. Chan, D., Beasley, S., Zhen, Y. & Spitale, R. C. Facile synthesis and evaluation of a dual-
functioning furoyl probe for in-cell SHAPE. *Bioorg. Med. Chem. Lett.* **28**, 601–605 (2018).

39. Busan, S. & Weeks, K. M. Accurate detection of chemical modifications in RNA by
mutational profiling (MaP) with ShapeMapper 2. *RNA* **24**, 143–148 (2018).

40. Campagne, S. *et al.* Structural basis of a small molecule targeting RNA for a specific
splicing correction. *Nat. Chem. Biol. 2019 1512* **15**, 1191–1198 (2019).

41. Nagano, K., Kamimura, T. & Kawai, G. Interaction between a fluoroquinolone derivative
and RNAs with a single bulge. *J. Biochem.* **171**, 239–244 (2022).

42. Ichijo, R., Kamimura, T. & Kawai, G. Interaction between a fluoroquinolone derivative
KG022 and RNAs: Effect of base pairs 3′ adjacent to the bulged residues. *Front. Mol.*
Biosci. **10**, 168 (2023).

43. Thakur, C. S. *et al.* Small-molecule activators of RNase L with broad-spectrum antiviral
activity. *Proc. Natl. Acad. Sci. U. S. A.* **104**, 9585–90 (2007).

44. Thakur, C. S., Xu, Z., Wang, Z., Novince, Z. & Silverman, R. H. A convenient and
sensitive fluorescence resonance energy transfer assay for RNase L and 2',5'
oligoadenylates. *Methods Mol. Med.* **116**, 103–13 (2005).

45. Adhikary, P. *et al.* Discovery of Small Anti-ACE2 Peptides to Inhibit SARS-CoV-2
Infectivity. *Adv. Ther.* **4**, 2100087 (2021).

46. Hargrove, A. E. Small molecule–RNA targeting: starting with the fundamentals. *Chem.*
Commun. **56**, 14744–14756 (2020).

47. Childs-Disney, J. L. *et al.* Targeting RNA structures with small molecules. *Nat. Rev. Drug*
Discov. **21**, 736–762 (2022).

48. Meyer, S. M. *et al.* DNA-Encoded Library Screening To Inform Design of a Ribonuclease
Targeting Chimera (RiboTAC). *J. Am. Chem. Soc.* **144**, 21096–21102 (2022).

49. Morgan, B. S., Forte, J. E., Culver, R. N., Zhang, Y. & Hargrove, A. E. Discovery of Key
Physicochemical, Structural, and Spatial Properties of RNA-Targeted Bioactive Ligands.
Angew. Chemie - Int. Ed. **56**, 13498–13502 (2017).

50.   Velagapudi, S. P., Gallo, S. M. & Disney, M. D. Sequence-based design of bioactive
small molecules that target precursor microRNAs. *Nat. Chem. Biol.* **10**, 291–297 (2014).

51.   Disney, M. D. *et al.* Informa 2.0: A Platform for the Sequence-Based Design of Small
Molecules Targeting Structured RNAs. *ACS Chem. Biol.* **11**, 1720–1728 (2016).

Methods
Synthesis of C30-FAI (cgSHAPE probe)

Compound **C30-alkyne** (50 mg, 0.08 mmol) in DMSO (1 mL) was added 2-(azidomethyl)furan-
3-carboxylic acid (15 mg, 0.09 mmol), THPTA (9 mg, 0.02 mmol), sodium ascorbate (8 mg, 0.04
mmol) and $CuSO_4$ (3 mg, 0.02 mmol). The reaction vial was sealed, evacuated, and refilled with
N_2 three times and stirred at room temperature for overnight. DMSO was removed under
vacuum and the residue was purified by silica gel column chromatography (0 – 10% CH_3OH in
CH_2Cl_2) to afford **C30-FCA** as a yellow solid (45 mg, 70%). MS-ESI (*m/z*) [M+1]$^+$ 746.28.

^{1}H NMR (500 MHz, DMSO-d_6) δ 8.71 – 8.68 (m, 2H), 8.52 (s, 1H), 8.09 (s, 1H), 7.71 – 7.68 (m,
2H), 7.38 (dd, *J* = 10.1, 2.6 Hz, 1H), 7.02 (dd, *J* = 9.0, 2.3 Hz, 1H), 6.97 (td, *J* = 7.6, 2.6 Hz, 1H),
6.88 (d, *J* = 2.4 Hz, 1H), 6.72 (d, *J* = 1.9 Hz, 1H), 5.92 (s, 2H), 4.52 (s, 2H), 3.58 – 3.51 (m,
14H), 3.38 (t, *J* = 5.1 Hz, 4H), 2.59 (t, *J* = 5.1 Hz, 4H), 2.56 (t, *J* = 5.8 Hz, 2H).

^{13}C NMR (126 MHz, DMSO-d_6) δ 160.2 (d, *J* = 252 Hz), 159.5, 154.9, 153.3, 152.2, 144.4,
144.3, 143.4, 139.4, 138.6, 129.6, 129.2 (d, *J* = 11.8 Hz), 124.2, 114.5, 112.1, 111.7, 111.4,
110.2, 104.2 (d, *J* = 29.8 Hz), 99.6, 99.4, 69.8, 69.7, 69.1, 63.4, 57.0, 54.9, 52.6, 46.7, 44.8.

Compound **C30-FCA** (45 mg, 0.06 mmol) in anhydrous DMSO (0.6 mL) was added
carbonyldiimidazole (CDI, 10 mg, 0.06 mmol) and the reaction mixture was stirred at room
temperature for 1 h. The reaction mixture contains ~75% **C30-FAI** and ~25% unreacted **C30-**
FCA (see Supplementary Information) and was used directly in RNA modification. The stock
solution was used as a **75 mM** and can be stored at –80 °C for long-term storage.

cgSHAPE-seq using Total RNA Extract from Cells

SARS-CoV-2 5' UTR expressing cells were harvested and pelleted. Total RNA was extracted
using TRIzol Reagent (Invitrogen) per the user's manual. An on-column DNA digestion was
performed to remove the residual genomic DNA in total RNA using DNase I (10 U/µL, Roche)
and RDD buffer (Qiagen). Purified total RNA was dissolved in water and stored at –80 °C before
use. For RNA modification, 5 µg total RNA was used for each reaction. **C30-FAI** and FAI-N_3
were prepared at 20 mM, 2 mM, and 0.4 mM in DMSO as 20× working solution. Briefly, total
RNA was added water and 5× folding/reaction buffer (500 mM HEPES pH 7.4, 500 mM KCl, 30
mM $MgCl_2$) to make a 47.5 µL solution. The solution was incubated at 37 °C for 30 min to refold.
2.5 µL **C30-FAI** (cgSHAPE probe), FAI-N_3 (background control) or DMSO was added to the
total RNA and mix well by pipetting. The mixture was incubated at 37 °C for 15 min and then
quenched by adding RLT buffer (Qiagen). The RNA was then extracted using RNeasy kit

(Qiagen). 500 ng total RNA was used for reverse transcription and then PCR as described
below. All reactions were performed in triplicates.

For reverse transcription (10 µL reaction), probe or DMSO treated RNA and reverse
transcription primer (0.5 µM in final reaction buffer) were heated at 70 °C for 5 min and snap-
cooled on ice for 1min. 5× reaction buffer (375 mM Tris-HCl, 500 mM KCl, 15 mM MnCl$_2$, pH
7.4, 2 µL), DTT (100 mM, 1 µL), dNTP (10 mM, 0.5 µL), ProtoScript II (0.5 µL, New England
Biolabs, M0368L) and RNase inhibitor (0.2 µL, ApexBio, K1046) were added. The reaction was
incubated at 42 °C for 1 h and deactivated at 70 °C for 15 min. In each PCR reaction (50 µL),
cDNA (2.5 µL) was mixed with Phire Hot Start II DNA Polymerase (Thermo Fisher, 1 µL), dNTP
(10 mM, 1 µL), 5× Phire Green reaction buffer (10 µL), primers (0.5 µM in final reaction buffer)
and water (35.5 µL). After reaction, the amplicon was purified using a DNA Clean &
Concentrator kit (Zymo Research) following user's manual. The purified DNA was submitted for
next-generation sequencing (Amplicon-EZ, Azenta Life Sciences).

An integrated software package developed by Busan and Weeks, ShapeMapper2 was used to
analyze the fastq files for mutational profiling and the rresult was used to generate Figures 2b,
2c, and Extended Data Fig. 3[39]. The reference sequence (SARS-CoV-2_5_UTR.fa) required for
ShapeMapper2 is listed below.

>SARS-CoV-2_5_UTR

aggtttataccttcccaggtaacAAACCAACCAACTTTCGATCTCTTGTAGATCTGTTCTCTAAACGAAC
TTTAAAATCTGTGTGGCTGTCACTCGGCTGCGTGCTTAGTGCACTCACGCAGTATAATTAAT
AACTAATTACTGTCGTTGACAGGACACGAGTAACTCGTCTATCTTCTGCAGGCTGCTTACG
GTTTCGTCCGTGTTGCAGCCGATCATCAGCACATCTAGGTTTCGTCCGGGTGTGACCGAAA
GGTAAGATGGAGAGCCTTGTCCCTGGTTTCAACGAGGGAGTCAAAGTTCTGTTTGCCCTGA
TCTGCATCGCTGTGGCCGAGGCCAAGCCCACCGAGAACAACGAagacttcaacatcgtggccg.
(lowercase = primer binding sequences).

In Vitro RNase L Degradation Assay

Purified recombinant GST-tagged RNase L was purchased from MyBioSource (MBS1041064).
The buffer of RNase L was exchanged into a buffer containing 50 mM Tris-HCl (pH 7.4) and 100
mM NaCl using Zeba Desalting Column (Thermo Fisher, 8766) using the manufacturer's
protocol. For RNase L degradation of SL5 RNA, T7 transcribed SL5 RNA was first purified by
polyacrylamide gel electrophoresis (PAGE) and recovered using small-RNA PAGE Recovery Kit
(Zymo Research, R1070)**.** RNase L (1.3 µg in 5 µL) was incubated in the presence of **C47**, **C48**,
C64, **C65**, or DMSO control in the cleavage buffer (final reaction volume is 8 µL) at 4 °C for 12
h. The 1X cleavage buffer contains 25 mM Tris-HCl (pH 7.4), 10 mM MgCl$_2$, 100 mM KCl, 50
µM ATP, and 7 mM β-mercaptoethanol. The SL5 RNA (120 ng in 2 µl H$_2$O) was then added into
the reaction mixture and incubated for another 2 h at 22 °C. The reaction was stopped by
adding RNA Gel Loading Dye (Thermo Fisher, R0641) at 1:1 ratio. The samples (4 µl) were
then loaded on TBE-urea polyacrylamide gel (20%) for electrophoresis (180 V for 85 min). The

gel was stained with SYBR safe (1/50,000, ApexBio, A8743) in TBE buffer for 1 min and
visualized on gel imager (Thermo Fisher, iBright FL1500).

The RNA sequence of SL5 used in this assay is: 5'-
UCGUUGACAGGACACGAGUAACUCGUCU
AUCUUCUGCAGGCUGCUUACGGUUUCGUCCGUGUUGCAGCCGAUCAUCAGCACAUCUA
GGUUUCGUCCGGGUGUGACCGAAAGGUAAGAUGGAGAGCCUUGUCCCUGGUUUCAACG
A.

For RNase L degradation of a model 6-FAM-tagged RNA, RIBOTACs in the above protocol
were replaced with **D1** (0.37 µg in 0.75 µl DMSO) or **2-5A-N₃** (0.12 ng in 0.75 µl H_2O)[43,44]. After
electrophoresis, the gel was not stained and was directly visualized on the gel imager at the 6-
FAM fluorescence channel. The 6-FAM RNA (5'-6-FAM-
UUAUCAAAUUCUUAUUUGCCCCAUU
UUUUUGGUUUA-BHQ) was purchased from IDT.

SARS-CoV-2 5' UTR Expressing Stable Cell Line

293T cells (Thermo Fisher, R70007) were cultured in DMEM growth medium (Gibco, 11995040)
supplemented with 10% FBS (Cytiva, SH30910.03) and 1% Antibiotic-Antimycotic (Gibco,
15240062) at 37 °C in 5% CO_2 atmosphere. For producing the lentivirus, 293T cells were
seeded in a 6-well plate (Fisher, FBO12927) at 3×10^5 cells per well and transfected with 1 µg
of SARS-CoV-2 5' UTR expressing lentivirus vector (pLV-SARS-CoV-2-5'UTR-GLuc) along with
the packaging plasmids pMD2.G (0.4 µg) and psPAX2 (0.6 µg) using Lipofectamine 2000
(Invitrogen, 11668019). At 24 h post-transfection, the cell medium was replaced with fresh
growth medium. 48 h after the change of media, the supernatant containing the lentivirus
particles was siphoned and centrifuged at 500 g for 10 min at 4 °C to remove the cell debris.
The virus particles were further concentrated at 10X in volume using Lenti-X-concentrator
(Clontech, PT4421-2) according to the manufacturer's protocol. The lentivirus can be quantified
using literature method[52]. Usually, 10^7–10^8 plaque forming units (pfu)/mL lentivirus was obtained
after the concentrator treatment. For lentiviral transduction, 293T cells were inoculated with the
concentrated viral suspension (multiplicity of infection ~10) using polybrene (Sigma-Aldrich, TR-
1003-G) at a final concentration of 8 µg/mL. At 24 h post-transduction, the culture medium was
replaced with fresh growth media. After recovery for 24 h, the transduced cells were then
selected in blasticidin (10 µg/mL, Invivogen, ant-bl) for 2 weeks. For stable single clone
selection, the cells were diluted in the growth medium containing blasticidin (10 µg/mL) to a final
density of 1 cell per 100 µL. The diluted cell suspension was then dispensed to a 96-well plate
(100 µL per well). The plate was incubated at 37 °C for 4 weeks. A single cell colony from one
of the wells was then selected for experiments.

pLV-SARS-CoV-2-5UTR-Luc was constructed by inserting the SARS-CoV-2 5' UTR and
Gaussian luciferase into the pLV vector, under the control of the CMV promoter. The insert
sequence is as follows:

attaaaggtttataccttcccaggtaacaaaccaaccaactttcgatctcttgtagatctgttctctaaacgaactttaaaatctgtgtggctg

tcactcggctgcgtgcttagtgcactcacgcagtataattaataactaattactgtcgttgacaggacacgagtaactcgtctatcttctgc

aggctgcttacggtttcgtccgtgttgcagccgatcatcagcacatctaggtttcgtccgggtgtgaccgaaaggtaag<u>atg</u>gagagcct

tgtccctggtttcaacgagggagtcaaagttctgtttgccctgatctgcatcgctgtggccgaggccaagcccaccgagaacaacgaa

gacttcaacatcgtggccgtggccagcaacttcgcgaccacggatctcgatgctgaccgcgggaagttgcccggcaagaagctgcc

gctggaggtgctcaaagagatggaagccaatgcccggaaagctggctgcaccaggggctgtctgatctgcctgtcccacatcaagt

gcacgcccaagatgaagaagttcatcccaggacgctgccacacctacgaaggcgacaaagagtccgcacagggcggcataggc

gaggcgatcgtcgacattcctgagattcctgggttcaaggacttggagcccatggagcagttcatcgcacaggtcgatctgtgtgtgga

ctgcacaactggctgcctcaaagggcttgccaacgtgcagtgttctgacctgctcaagaagtggctgccgcaacgctgtgcgacctttg

ccagcaagatccagggccaggtggacaagatcaaggggggccggtggtgactaa (lowercase = SARS-CoV-2 5'

UTR; uppercase = Gaussia luciferase; underline = start codon).

Quantitative Reverse Transcription PCR (RT-qPCR) Assay

The SARS-CoV-2 5' UTR expressing cells were seeded at 3 x 10^5 cells per well in 12-well
plates in 1 mL growth medium at 37 °C for 3 h. The cells were then treated with the compounds
(**C47**, **C48**, **C64**, or **C65**) at various concentrations (1.3 nM–3 µM) for 48 h. After treatment, the
supernatant was aspirated from each well and the total RNA was then extracted from the cells
using RNeasy mini kit (Qiagen, 74104). The total RNAs were quantified by ultraviolet absorption
at 260 nm (Thermo Fisher, NanoDrop 1000). Usually 10–20 µg total RNA was obtained from
each well. cDNAs were synthesized from 500 ng of total RNA for each sample using M-MLV
reverse transcriptase (Promega, M1701) and $(dT)_{25}$ according to the manufacturer's protocol. 1
µl of cDNA mixture was used in a 15 µl RT-qPCR reaction (Apex-Bio, K1070). The human
GAPDH RNA level was used as the reference for normalization. The RT-qPCR primer
sequences used for the PCR are shown below:

SL5-SYBR-FW: 5'-CGTTGACAGGACACGAGTAA

SL5-SYBR-RV: 5'-TTGAAACCAGGGACAAGGCTC

GAPDH-FW: 5'-GACAAGGCTGGGGCTCATTT

GAPDH-RV: 5'-CAGGACGCATTGCTGATGAT

SARS-CoV-2 Inhibition Assay

538   Vero-E6 cells (ATCC® CRL-1586™) and A549 cells (ATCC® CCL-185) were cultured in
539   Dulbecco's modified Eagle's medium (DMEM, Cytiva Life Science, SH30022) with addition of
540   10% fetal bovine serum (FBS, Millipore Sigma, F0926) at 37°C under 5% CO_2 atmosphere.
541   A549 cells were transduced with a human ACE2-expressing lentivirus vector, and the
542   transduced were cultured in the DMEM plus 2 µg/µL puromycin[45].

Virus and titration

SARS-CoV-2-Nluc was created by engineering the nanoluciferase (Nluc) gene into the OFR7 of
the SARS-CoV-2 genome. The insertion site of Nluc at ORF7 was based on previous
mNeonGreen reporter SARS-CoV-2[53]. The virus was propagated in Vero-E6 cells once,
aliquoted in DMEM, and stored at –80°C. A biosafety protocol to work on SARS-CoV-2 in the
BSL3 Lab was approved by the Institutional Biosafety Committee of the University of Kansas
Medical Center.

Plaque assay[54,55]

Vero-E6 cells were seeded in 24-well plates at a density of 0.5 ×10^6 cells per well. A virus stock
was serially diluted at 10-fold in Dulbecco's phosphate-buffered saline, pH7.4 (DPBS). 200 μL of
the diluent were added to each well and incubated for 1 h on a rocking rotator. After removing
the virus diluent, 0.5 mL of overlay media (1% methylcellulose in DMEM with 5% FBS) were
added to each well. The plates were incubated at 37 °C under 5% CO_2 for 4 days. The
methylcellulose overlays were aspirated, and the cells were fixed with 10% formaldehyde
solution for 30 min and stained with 1% crystal violet solution followed by extensive washing.
Plaques in each well were counted and multiplied by the dilution factor to determine the virus
titer at pfu/mL.

Determination of half-maximal inhibitory concentration (IC$_{50}$)[45,56]

ACE2-A549 cells were seeded into 96 well plates. When the cells were confluent, SARS-CoV-2-
NLuc viruses were diluted with cold PBS and added into each well at a multiplicity of infection
(MOI) of 2 (2 pfu/cell). The plates were kept in the CO_2 incubator for 1 hour. Compound **C64**
was diluted at 2 x serials from 20 μM to 0.002 μM. The virus-PBS solution was aspirated. Each
well was washed with cold PBS three times, and was loaded with the diluted compounds. Each
concentration was loaded in triple wells in the plates, and the total volume of each well was 0.2
mL. The plates were kept in the incubator. After 3 days post-infection, the culture media were
aspirated from each well and the wells were washed with PBS for three times. The nano-
luciferase activity assay (Promega, N1110) was carried out by following the manufacturer's
instructions. Briefly, 100 ul of cell lysis buffer were added to each well for 10 minutes to
completely lyse the cells. Then 100 ul of nano-luciferase reaction reagent were add to each well
and the luminescent signal was determined at A490 absorbance on a plate reader (Bio-Tek,
Synergy). The IC$_{50}$ was calculated using GraphPad Prism 8.0 software.

Statistical Analysis

All data shown as means ± s.d. with sample size (N) listed for each experiment. Statistical
analysis was carried out with Prism GraphPad 8.0. Unpaired two-sample t-tests were used to
analyze significant differences between the group means. The P values were calculated by
Prism GraphPad 8.0 or R. For data generated from ShapeMapper2, the standard error (stderr)
associated with the mutation rate at a given nucleotide in the S (probe treated) or U (DMSO

treated) samples was calculated as: stderr = $\sqrt{mutation\ rate}/\sqrt{reads}$. The standard error of the

Δmutation rate at a given nucleotide is: $\sqrt{stderr_S{}^2 + stderr_U{}^2}$.

Data Availability

585 The cgSHAPE-seq data for **C30-FAI**, FAI-N$_3$ or DMSO-treated total RNAs, and RNA-seq data
586 for **C64**-treated cells were deposited in NCBI SRA with accession numbers PRJNA950557 and
587 PRJNA947619, respectively.

References

52.   Barczak, W., Suchorska, W., Rubiś, B. & Kulcenty, K. Universal Real-Time PCR-Based
Assay for Lentiviral Titration. *Mol. Biotechnol.* **57**, 195–200 (2015).

53.   Xie, X. *et al.* A nanoluciferase SARS-CoV-2 for rapid neutralization testing and screening
of anti-infective drugs for COVID-19. *Nat. Commun.* **11**, 5214 (2020).

54.   Hao, S. *et al.* Long-term modeling of SARS-CoV-2 infection of in vitro cultured polarized
human airway epithelium. *MBio* **11**, 1–17 (2020).

55.   Zou, W. *et al.* The sars-cov-2 transcriptome and the dynamics of the s gene furin
cleavage site in primary human airway epithelia. *MBio* **12**, (2021).

56.   Hao, S. *et al.* Establishment of a Replicon Reporter of the Emerging Tick-Borne Bourbon
Virus and Use It for Evaluation of Antivirals. *Front. Microbiol.* **11**, 2229 (2020).

Acknowledgments

Research reported in this article was supported by the National Institute of General Medical
Sciences (NIGMS) of the National Institutes of Health (NIH) under award numbers
R35GM147498 and P20GM113117, W. M. Keck Foundation, and the University of Kansas
General Research Funds. We also thank Dr. Robert Silverman at Cleveland Clinic for providing
the authentic 2-5A samples for comparison. We have obtained SARS-CoV-2-Nluc from Drs. Shi
and Menachery through The University of Texas Medical Branch (UTMB)'s World Reference
Center for Emerging Viruses and Arboviruses. We thank the University of Kansas Medical
Center Genomics Core for doing RNA-seq experiment. (The KUMC Genomics Core was
supported by Kansas Intellectual and Developmental Disabilities Research Center (NIH U54 HD
090216), the Molecular Regulation of Cell Development and Differentiation COBRE (P30
GM122731-03), the NIH S10 High-End Instrumentation Grant (NIH S10OD021743) and the
Frontiers CTSA grant (UL1TR002366)).

Author contributions

J.W. conceived the work. J.W., Z.T., S. Hegde and J.Q. wrote the paper. Z.T and M.S. performed chemical synthesis. Z.T. and J.W. conducted the RNA sequencing and analysis. S. Hegde collected RNA degradation data in vitro and in cells. S. Hao and J.Q. performed the live virus assay.

Competing interests

The authors declare no competing interests.

Extended Data Fig. 1 | Binding affinity of coumarin derivatives to SL5 RNA. a, Dose-response curves of
coumarin derivatives in fluorescence polarization assay with in vitro transcribed SL5 RNA. All compounds were used
at a concentration of 80 nM. Each data point represents the mean fluorescence polarization value of two independent
replicates (N = 2). **b**, Surface plasmon resonance (SPR) binding analysis of **C30** with SL5 RNA. (Curve fitting is
shown as black line. [SL5 RNA] = 0.075, 0.15, 0.3, and 0.6 µM).

**Extended Data Fig. 2 | Background Δmutation rate (FAI-N$_3$ – DMSO) of A, C, G, and U in denatured RNA
treated with FAI-N$_3$.** Denatured RNA was treated with **FAI-N$_3$** (100 mM) at 80 °C for 5 min in the denaturing buffer
containing 90% formamide, 5 mM EDTA. Mutation rate was calculated with ShapeMapper2 software package[39]. RNA
sequence is 5'-
aacuuccuuuauuuuccuuacaggguuuuAGACAAAAUCAAAAAGAAGGAAGGUGCUCACAUUCCUUAAAU
UAAGGAGUAAGUCUGCCAGCAUUAUGAAAGUGAAUCUUACUUUUGUAAAACUUUAUGGUUUGUGGAAAACAA
AUGUUUUUGAACAUUUAAAAAGUUCAGAUGUUAGAAAGUUGAAAGGUUAAUGUAAAACAAUCAAUAUUAAAGA
AUUUUGAUGCCAAAACUAUUAGAUAAAAGGUUAAUCUACAUCCCUACUAGAAUUCUCAUACUUAACUGGUUGG
UUGuguggaagaaacauacuuucacaauaaagagc. (lowercase = primer binding sequences).

Extended Data Fig. 3 | Background Δmutation rate (FAI-N_3 – DMSO) of the SL5 sequence in total RNA extract treated with different concentrations of FAI-N_3. One data point (C75) was removed as an outlier as it had an abnormally high mutation rate (*Z*-score > 4.0) for the DMSO-treated sample.

Extended Data Fig. 4 | Dose-response curves of **C30** (80 nM) in fluorescence polarization assay with SL5 RNA and its substructures. Each data point represents the mean fluorescence polarization value of two independent replicates (N = 2).

Extended Data Fig. 5 | In vitro RNase L degradation assay with a 5' 6-fluorescein-tagged model RNA
containing multiple RNase L cleavage sites. **a**, Comparison of RNA degradation activity of RNase L in the
presence of **2-5A** and **2-5A-N₃** with no significant differential activity observed. **b**, Comparison of synthetic RNase L
recruiter (D1) and 2-5A. The activity of **D1** in RNase L activation is ~ 10,000 times weaker than **2-5A**. **c**, Chemical
structures of **2-5A**, **2-5A-N₃**, and **D1**.

Extended Data Fig. 6 | RNA degrading activity of C47 and C48. **a**, Comparison of two RLR moieties in the
RIBOTAC modality using the in vitro RNase L degradation assay with purified SL5 RNA (red circle is a staining
artifact). **b**, Cellular activity of **C47** and **C48** in SARS-CoV-2 5' UTR expressing cells.

Extended Data Fig. 7 | Volcano plot of differential gene expression in SARS-CoV-2 5' UTR expressing cells treated
with C64 (3 µM). DMSO-treated cells were used as a control. Red spot = SARS-CoV-2 5' UTR transcript. The RNA-
seq analysis was performed with three biological replicates (N = 3).