Automated tumor immunophenotyping predicts clinical benefit from anti-PD-L1 immunotherapy

Xiao Li[1]*, Jeffrey Eastham[1]*, Jennifer M. Giltnane[1], Wei Zou[1], Andries Zijlstra[1], Evgeniy Tabatsky[1], Romain Banchereau[1], Ching-Wei Chang[1], Barzin Nabet[1], Namrata Patil[1], Luciana Molinero[1], Steve Chui[1], Maureen Peterson[1], Shari Lau[1], Linda Rangell[1], Yannick Waumans[2], Mark Kockx[2], Darya Orlova[1]*#, Hartmut Koeppen[1]*#

[1] Genentech, South San Francisco, California, 94080, USA

[2] CellCarta, 2610 Antwerpen, Belgium

To whom correspondence should be addressed: dyorlova@gmail.com ,

koeppen.hartmut@gene.com

* XL and JE are joint first authors

Running Title: Tumor immunophenotype predicts clinical benefit from immunotherapy

List of abbreviations

TME: tumor microenvironment; PD-L1: programmed death ligand-1; NSCLC: non-small cell lung cancer; TNBC: triple-negative breast cancer; IO: immuno-oncology; CI: checkpoint inhibitor; IP: immunophenotype, IHC: immunohistochemistry; H&E: hematoxylin and eosin; IC: immune cells; TILs: tumor-infiltrating lymphocytes; CK-cytokeratin; OS: overall survival; PFS: progression-free survival; WSI: whole slide images; MOCHA: randoM fOrest Classifier witH spAtial statistics; MOCHA-BITE: randoM fOrest Classifier witH spAtial statistics and BInned CD8 T-cell dEnsity

Abstract

Background. Cancer immunotherapy has transformed the clinical approach to patients with malignancies as profound benefits can be seen in a subset of patients. To identify this subset, biomarker analyses increasingly focus on phenotypic and functional evaluation of the tumor microenvironment (TME) to determine if density, spatial distribution, and cellular composition of immune cell infiltrates can provide prognostic and/or predictive information. Attempts have been made to develop standardized methods to evaluate immune infiltrates in the routine assessment of certain tumor types; however, broad adoption of this approach in clinical decision-making is still missing. **Methods.** We developed approaches to categorize solid tumors into "Desert", "Excluded" and "Inflamed" types according to the spatial distribution of CD8+ immune effector cells to determine the prognostic and/or predictive implications of such labels. To overcome the limitations of this subjective approach we incrementally developed four automated analysis pipelines of increasing granularity and complexity for density and pattern assessment of immune effector cells. **Results.** We show that categorization based on "manual" observation is predictive for clinical benefit from anti-programmed cell death ligand-1 (PD-L1) therapy in two large cohorts of patients with non-small cell lung cancer (NSCLC) or triple-negative breast cancer (TNBC). For the automated analysis we demonstrate that a combined approach outperforms individual pipelines and successfully relates spatial features to pathologist-based read-outs and patient response to therapy. **Conclusions.** Our findings suggest tumor immunophenotype (IP) generated by automated analysis pipelines should be evaluated further as potential predictive biomarkers for cancer immunotherapy.

What is already known on this topic

Clinical benefit from checkpoint inhibitor-targeted therapies is realized only in a subset of patients. Robust biomarkers to identify patients who may respond to such therapies are needed.

What this study adds

We have developed manual and automated approaches to categorize tumors into immunophenotypes based on the spatial distribution of CD8+ T effector cells that predict clinical benefit from anti-PD-L1 immunotherapy for patients with advanced non-small cell lung cancer or triple-negative breast cancer.

How this study might affect research, practice or policy

Tumor immunophenotypes should be further validated as predictive biomarker for checkpoint inhibitor-targeted therapies in prospective clinical studies.

Keywords

Immunophenotype, panCK/CD8 immunohistochemistry, CD8+ T cell, non-small cell lung cancer, triple-negative breast cancer, artificial intelligence, anti-PD-L1 immunotherapy, survival analysis

Background

Immuno-oncology (IO) has been practice-changing for the clinical care of patients with a variety of malignancies; long-lasting responses including cures have been observed in patients treated with therapeutics targeting checkpoint inhibitor (CI) molecules such as Cytotoxic T Lymphocyte Antigen 4 and Programmed Death Ligand 1 (PD-L1) [1] [2] [3]. Furthermore, therapeutics targeting other CI molecules or antibodies with immunomodulatory properties such as bi-specific antibodies have shown promise in early stages of clinical evaluation [4]. However, these encouraging results with IO therapeutics are seen only in a subset of patients [5]. This

realization has triggered extensive efforts to identify predictive biomarkers in order to maximize efficacy while identifying patients unlikely to respond but who would still be exposed to potentially life-threatening side effects.

Three assays have received regulatory approval for prospective patient identification for CI-targeted therapy: Immunohistochemistry (IHC) for PD-L1 and assessment of tumor mutational burden or microsatellite instability. PD-L1 IHC is by far the most commonly used methodology; however, the requirement of different assays and scoring algorithms across different indications and the imperfect correlation between IHC results and therapy outcome have caused frustration amongst clinicians and pathologists [6] [7] [8] [9] [10].

Success of IO relies on generating or facilitating an anti-tumor immune response. The vast number of known interactions between immune cells (IC) and an established tumor along with the cellular complexity and plasticity of such an immune response makes it difficult to identify a single parameter with sufficient predictive power [11] [12] [13]. Density and phenotype of IC in the TME have previously been used to identify patients with better clinical outcome and/or response to therapy [14]. The Immunoscore [15] [16] [17], originally developed for primary colorectal cancer, evaluates the spatial distribution and density of Cluster of Differentiation 3-positive (CD3+) and CD8+ T lymphocytes and identifies patients with improved clinical outcome independent of prognostic factors such as age, sex, tumor and lymph node status. This approach has been validated in a large patient cohort across multiple institutions using a digital pathology-based image analysis tool [17]. A scoring system for tumor-infiltrating lymphocytes (TILs) has been developed for ductal breast cancer and other carcinomas. It estimates on hematoxylin-eosin (H&E) stained sections the stromal density of all mononuclear cells including plasma cells but excludes granulocytes as well as intra-epithelial immune cells [18] [19] [20] [21]. High density of stromal TILs tends to correlate with better clinical outcome but is also associated with inter-observer variability [22] [23]. Lately, different approaches to

assess TILs in solid tumors using digital quantification have been published [24] [25] [26] [27]. Heterogeneity of these approaches with respect to tumor indication, methodology to identify TILs and readouts makes it challenging to compare results and emphasizes the need for standardization as well as validation in large and well-annotated clinical cohorts [28].

Recent studies have described approaches using image analysis with or without machine-learning to delineate the spatial distribution of immune effector cells, both stromal and intra-epithelial, correlating the pattern with gene expression signatures [29] and clinical outcome for patients treated with CI-targeted therapies [30]. Here, we categorized tumors into "Desert", "Excluded" and "Inflamed" immunophenotypes (IP) based on the density and spatial distribution of CD8+ T effector cells. We reasoned that such categories would provide prognostic and/or predictive information for patients treated with the anti-PD-L1 agent atezolizumab. We show that manual immunophenotyping of tumors is predictive for clinical benefit by CI-targeted therapy in two independent cohorts of two distinct indications. We describe the development and performance of an automated classification pipeline that yields equivalent or superior results compared with the manual methodology. Our results indicate that IP information generated by an automated classifier could represent an important predictive biomarker for IO clinical studies.

Methods

Patient cohorts and histology slides

We had access to paraffin embedded sections from three clinical trials. POPLAR (NCT01903993) is a randomized phase 2 study of atezolizumab compared with docetaxel in 287 patients with locally advanced or metastatic NSCLC who failed platinum therapy [31]. OAK (NCT02008227) is a randomized phase 3 trial of 1225 patients with identical design as POPLAR [32]. IMpassion130 (NCT02425891) is a phase 3 randomized clinical study of atezolizumab in

combination with nab-paclitaxel (atezolizumab arm) compared with placebo with nab-paclitaxel (placebo arm) for 902 patients with previously untreated TNBC [33]. Tissue sections had to show at least 100 viable invasive tumor cells with associated stroma for immunophenotype evaluation. In the rare case where more than one sample had been submitted for a patient, the sample with the largest tumor area was selected for evaluation. Samples from the same patient with divergent manual tumor IP's were excluded when considering patient response. This resulted in the evaluation of 745 patient samples for OAK, and 800 for IMpassion130 for manual tumor IP versus automated calls. The study was conducted according to the Declaration of Helsinki and all patients had provided written consent.

Immunohistochemistry (IHC)

Sections were stained for CD8 and pan-cytokeratin (CK) in a dual chromogenic IHC assay on a Discovery ULTRA platform (Roche Tissue Diagnostics (RTD), Tucson, AZ). Primary antibodies added to the sections were at 3µg/ml (CD8; C8144B) and at 1.6µg/ml (pan-CK; AE1/AE3), respectively (Agilent, Santa Clara, CA). Antigen retrieval was performed with Cell Conditioning Solution 1 (CC1, Cat.# 950-224; RTD) for 64 minutes at 95°C and sections were exposed to primary antibody for 32 minutes at 37°C. Specifically-bound primary antibody was detected using OmniMap anti-mouse HRP (Cat.# 760-4310; RTD) with diaminobenzidine (DAB; CD8) and DISCOVERY Purple (pan-CK; Cat.# 760-229; RTD) followed by counterstain with hematoxylin.

IP assessment - Manual approach

The tumor area was defined as viable tumor cells with associated stroma. Areas of necrosis or intraluminal aggregates of CD8+ cells were excluded from evaluation. CD8+ cells were detected in essentially all cases; tumors with density of CD8+ cells of "0", i.e. single dispersed cells with a density too sparse to allow identification of a distinct pattern, were classified as "Desert" (Figure

1A). "Excluded" tumors showed distribution of CD8+ cells limited to the CK- stroma (Figure 1B). "Inflamed" tumors showed co-localization of CD8+ cells with CK+ tumor cells either as a diffuse infiltrate with or without involvement of CK- stroma, or with a predominantly stromal distribution of CD8+ cells with "spill-over" into CK+ tumor cell aggregates (Figure 1C & D). We considered these two patterns as respective ends of a spectrum and did not distinguish between them for our downstream analysis. To address intratumoral heterogeneity of density and pattern of the infiltrate (Figure 1 E & F) we implemented a 20% cut-off; patterns occupying less than 20% of the tumor area were not considered for categorization. Cases showing an inflamed phenotype in >20% of the tumor area were labeled "Inflamed", independent of the pattern(s) observed in the remaining areas. Implementation of the cutoff was based on a subjective estimate by the inspecting pathologist (Figure 1G).

IP assessment - Automated approach

Whole slide image (WSI) segmentation and tiling

Slides were scanned on a NanoZoomer XR whole slide imager (Hamamatsu, Bridgewater NJ) with a 20x lens, and the images were used to quantify the tumor and IHC positive areas. Annotation of the tumor area was performed manually, excluding areas of assay or scanning artifact and large areas of necrosis. Within these regions, segmentation of CK+ tumor regions and CD8+ DAB pixels were performed by a custom algorithm using Hue, Saturation, and Value (HSV) color thresholding and standard morphological operations in MatLab 2020b (Mathworks, Natick, MA). Images were analyzed using a sliding window of 280x280 pixels and a stride of 200 pixels (0.46 um per pixel) for every analysis except Pipeline #1. CD8 density was quantified at the pixel level in both the CK+ and CK- compartments in each tile, and recorded separately as tile specific measurements, including tile coordinates to enable spatial distribution analysis.

Pipeline #1 - Density cutoff

Whole-slide level algorithmic readouts of CD8 density in the CK+ compartment were used to rank order the POPLAR dataset. Using the relative proportions of manual tumor IP calls as a guide, the lowest 20% of these samples were designated as Desert, the next 40% as Excluded, and the highest 40% as Inflamed. The ratio of CD8+ area to CK+ area cutoffs between the classes in POPLAR, 0.00059 CD8+ area / CK+ area between Desert and Excluded, and 0.005 CD8+ area / CK+ area between Excluded and Inflamed, was then used to classify samples in OAK and IMpassion130.

Pipeline #2 - Binned CD8 density

For each slide, 10 bins for the density of CD8+ pixels within the CK+ compartment in each tile were empirically defined (with upper bounds of: 0.005, 0.01, 0.02, 0.04, 0.06, 0.08, 0.12, 0.16, 0.2, 1) and the number of tiles in each bin was recorded separately for both compartments. Tiles with less than 25% of their area containing CK+ area were excluded from this count. The same process was performed using the CK- compartment, resulting in 10 additional bins. These two sets of bins are considered entirely separately, and as such it is possible for one tile to contribute to both sets of bin distributions. This resulted in a 20-element vector for each slide, with 10 measurements from each compartment. The vectors from the entire POPLAR dataset were used to train a three class Support Vector Machine (SVM) in Matlab using built-in hyperparameter optimization. This trained SVM was then validated by predicting the phenotype classification for OAK, and tested without further optimization on IMpassion130.

Pipeline #3 - randoM fOrest Classifier witH spAtial statistics (MOCHA)

A collection of spatial analytics approaches based on the lattice system was utilized. Firstly, we extracted the center coordinates of the tiles and treated each tile as a spatial unit. Analogous to spatial epidemiology maps, a CK/CD8 prevalence map was created [34], [35]. Specifically, for

each tile, the CK+/CD8+ area was calculated, i.e. number of CK+/CD8+ pixels. We further

normalized the areas for each tile divided by the sum of CK/CD8 areas across all tiles for the

same slide. In addition, we define CK+ tiles as tiles with ratio ≥0.25, where the ratio is

calculated as CK+ pixels/(CK+ pixels + CK- pixels) and CK- tiles otherwise. Secondly, based on

the prevalence maps, we derived 50 spatial features that either captured the co-localization of

CK and CD8 in each slide or spatial distribution of CK or CD8 in each slide, respectively. Some

of the features, such as the CD8-CK ratio and Bhattacharyya coefficient in CK+/CK- regions,

allow to quantify the CD8 cell density and their colocalization with tumor/stroma cells, and thus

could serve as rough proxy for the manual IP (Figure S1A). Collectively, the following set of

features was derived:

 a) spatial co-localization features of CK and CD8 in CK+ and CK- tiles, respectively:

Bhattacharyya coefficient, Morisita-Horn index, Jaccard index, Sorensen index and bivariate

Moran's Index, co-localized Getis-Ord hotspot [36] [37] [38] [39] and CD8-CK area ratio ;

b) spatial distribution features of CK and CD8 in CK+ and CK- tiles, respectively: local and

global Moran's and Geary's C indices [40] and Getis-Ord hotspot [41];

A three-class random forest classifier was then trained based on the entire POPLAR dataset

with these 50 features per slide as input. This trained model was then tested by predicting the

phenotype classification for OAK and IMpassion130 without further model optimization.

Pipeline #4 - randoM fOrest Classifier witH spAtial statistics and BInned CD8 T-cell dEnsity
(MOCHA-BITE)

In order to capture the density information as well as the spatial distribution of CK and CD8,

respectively, for each slide, we concatenated the features described in Pipeline #2 and Pipeline

#3 for each slide. A three-class random forest classifier was then trained based on the entire

POPLAR dataset with the combined feature vector per slide as input. This trained model was

then tested by predicting the phenotype classification for OAK and IMpassion130 without further model optimization. The five top performing features are shown in Figure S1B.

Results

Manual tumor IP classification

We established a methodology to assign tumor IP based on the spatial distribution and density of CD8+ T effector cells on sections stained for CD8 and CK and assigned each case a label of Desert, Excluded or Inflamed. The impact of this categorization on patient outcome was evaluated in the atezolizumab-treatment arms of the OAK and IMpassion130 clinical studies. Both studies showed longer patient overall survival (OS) when tumors displayed an Inflamed phenotype; patients with Desert tumors had the lowest OS, while patients with Excluded tumors showed intermediate outcome (Figure 1H). Median OS for Inflamed, Excluded and Desert tumors were 15.4, 14.1 and 9.5 months for OAK and 27.3, 22.6 and 15.5 months for IMpassion130, respectively. Patients in the control arm for either trial did not show differences in clinical outcome based on tumor IP (Figure S2).

Automated tumor IP classification

In order to address subjectivity and lack of scalability of manual tumor IP assignment, we developed an integrated automated classification approach consisting of four distinct pipelines; all pipelines require pre-processed WSI of CD8/CK-stained slides with annotated tumor areas as input (Figure 2A; also see Materials & Methods for details).

Pipeline#1 is based on the averaged CD8+ cell density score at the slide level as the simplest representation of manually derived IP. In this analysis, the average density of CD8 in the CK+ compartment of the annotated tumor region is determined. We observed a unimodal distribution

of CD8 density across all samples with an enrichment of manually classified Desert samples at the low and Inflamed at the high end (Figure 3A). Unimodality of this distribution suggested that fully unsupervised classification of samples into three IP groups would lead to a high misclassification error. IP classification was achieved in a supervised fashion by application of CD8 density cutoffs established using the manual IP calls from POPLAR. The prevalence of manual IP calls in this trial was used to determine these cutoffs, with the 20th percentile of CD8 densities in CK+ regions corresponding to the percentage of Desert samples, resulting in a CD8 density ratio cutoff of 0.059, and the 60th percentile and a CD8 density cutoff of 0.5 for Excluded samples, with the remaining cases assigned to Inflamed samples. These CD8 density cutoffs were then applied to OAK and IMpassion130. A surprisingly high accuracy and linear weighted Cohen's kappa (0.558 in OAK, and 0.443 in IMpassion130) were observed (Figure 4A, 4C).

This same cutoff methodology of using the 20th percentile as the cutoff between Desert and Excluded, and the 60th percentile between Excluded and Inflamed was applied to CD8 density calculated across the entire tumor area without regard to CK+ regions. This resulted in a CD8 density ratio cutoff of 0.003 for Desert samples, and a cutoff of 0.0014 for Excluded samples. This resulted in a decrease in overall accuracy and Cohen's kappa by approximately 4% for each measure compared to the analysis of the CK+ compartment (data not shown).

Pipeline #2 incorporates the concept of local density used in the manual process. Each image was analyzed using small tiles of 16,000 um^2, providing localized measurements of CD8 density with respect to CK staining (Figure 2B). Within each tile, the CD8+ density was measured separately within CK+ and CK- compartments. The readouts for each tile were divided into 10 bins based on CD8 density within their respective CK+ or CK- compartments, resulting in 20 readouts per sample. These readouts alone, used in the fully unsupervised manner, could not clearly segregate tumors into three clean clusters (Figure 3B); however, there appeared to be a trend for Desert, Excluded and Inflamed enriched regions in that 20

dimensional binned IC density feature space projected into the two dimensional Uniform Manifold Approximation and Projection (UMAP) space. To arrive at a cleaner IP assignment we adopted a supervised classification approach. Each of the twenty CD8+ density bins was used to define a multidimensional feature space, which was separated by a multiclass Support Vector Machine (SVM) trained with the corresponding manual phenotype classifications in POPLAR. The resulting SVM was then used to classify OAK and IMpassion130 (Figure 4A). The weighted Cohen's kappa for the Binned CD8 density pipeline was 0.562 in OAK, and 0.628 in IMpassion130 compared to the manual immunophenotype classification (Figure 4C).

Pipeline #3 (MOCHA) considers the more granular spatial pattern and density of immune infiltrates (Figure 2A). CK and CD8+ densities calculated in each tile allowed us to obtain a two dimensional density distribution (Figure 2C) of tumor cells and CD8+ T cells. In order to quantitate spatial patterns of tumor cells and CD8 T cells such as cell colocalization, presence of cell "hot spots" etc., we used a collection of lattice-based spatial analytics approaches (see Methods for details). As a result, a set of fifty spatial features was extracted for each tiled WSI from POPLAR, OAK and IMpassion130 datasets. Since these spatial features alone do not clearly differentiate samples into the three IP categories (Figure 3C), we used manual IP calls from the entire POPLAR dataset to train the random forest classifiers and then tested them on OAK and IMpassion130 datasets. This pipeline successfully relates spatial features to manual IP calls (Figure 4A) with the weighted Cohen's kappa of 0.573 for OAK, and 0.456 for IMpassion130 (Figure 4C).

Pipeline #4 (MOCHA-BITE) combines the 50 spatial features of pipeline #3 with the binned CD8+ cell densities of pipeline #2 and a random forest classifier with manual IP calls is trained in a supervised learning framework. We used IP classification agreement and OS/PFS log-rank test results as criteria to assess pipeline performance.

Based on Cohen's kappa MOCHA-BITE outperforms the other three pipelines (Figure 4, Figures S1-7) and recapitulates the results observed by manual IP assignment; the only exception is the analysis using pipeline #2 (binned CD8+ cell density) for IMpassion130, which on the other hand performs inferiorly in the PFS analysis (Figure 4C). MOCHA-BITE yielded significant log-rank test results when comparing the three predicted IP groups for OS and PFS using OAK and IMpassion130 as test sets (Figure 4C).

In order to determine if the treatment effect differs between IP patient subgroups (identified by manual or MOCHA-BITE analysis) we performed an interaction test as illustrated in the forest plot analyses for both clinical trials; we focused on OS and "condensed" Desert and Excluded tumors into a "Non-Inflamed" group. We observed that atezolizumab is linked to improved OS clinical benefit in Inflamed vs Non-Inflamed tumors in both, OAK (HR 0.65 [CI 95% 0.48-0.97] vs 0.88 [CI 95% 0.72-1.08], respectively) and IMpassion130 (HR 0.53 [CI 95% 0.38-0.75] vs 0.93 [CI 95% 0.77-1.11], respectively). The difference in HR between Inflamed and Non-Inflamed tumors for either IP classification methodology is statistically significant for IMpassion130 (Figure 5A, 5B). This is also shown in the separation of the accompanying OS KM curves for Inflamed and Non-Inflamed tumors in the atezolizumab vs control arm of each trial.

Discussion

Analysis of digital pathology images based on spatial statistics has previously shown value in different clinical use cases for solid and liquid tumors [36] [37] [42] [43] [44] [27]. In this study, we show that tumor IP assignment using different methodologies is predictive for outcome of anti-PD-L1 targeted therapy. We developed our own approach instead of applying previously published algorithms such as Immunoscore [45] or traditional TIL scoring [19] for several reasons. We wanted to pursue the task in an indication-agnostic fashion. Unambiguous

identification of epithelial and stromal compartments as well as labeling of the relevant immune cell type allowed us to implement our analyses in two different epithelial tumor types. Furthermore, diverse specimen types are typically encountered in the clinical trial setting (resections, excisional and needle core biopsies); while our methodology has a minimum requirement of viable, invasive tumor and associated stroma, we did not want to solely rely on resection specimens with representation of tumor center and invasive edge. This would severely limit the number of available specimens and may introduce unwanted bias toward better clinical outcome as resections in NSCLC patients are primarily performed with curative intent. Lastly, we felt strongly about the value of our focus on intra-epithelial CD8+ T-cells in our assessment.

In order to address the limitations of the manual approach of scalability and inter-/intra-observer variability, we developed individual automated pipelines for IP calls and show that using a combined approach (pipeline #4) outperforms individual pipelines in two distinct tumor types. While the training set was limited to cases of NSCLC, implementation of the analysis was successful in TNBC without the need for additional machine-learning efforts. Extension to other tumor types should be straightforward as long as the malignant cells can be unambiguously identified. Furthermore, our approach lends itself to evaluate the distribution of immune cells other than CD8+ T-effector cells as long as they can be identified by immunohistochemical means. This could be relevant for IO therapies targeting molecules other than PD-1 and PD-L1. Also, expansion into a multiplexed methodology with identification of more than one immune cell phenotype should be rather seamless.

An end-to-end deep learning–based AI model that segments cancer epithelium, cancer stroma, as well as TILs to categorize tumors into IP similar to our classification has recently been developed for NSCLC [46]. Similar to our findings, the authors describe improved clinical outcomes for patients with inflamed tumors when treated with anti-PD-1 or anti-PD-L1 agents .

The proportion of the three IP categories are similar but far from identical between their and our methodology. There are several reasons for this "discrepancy": The composition of the respective patient cohorts is most likely different with respect to clinical parameters, cut-offs to define IP categories are different between the two methodologies, and identification of the two tumor compartments and immune cells relies on machine-learning (and thus requires training data) in one and presence/absence of an immunohistochemical stain in the other (our) approach. The latter is an important distinction as it offers the opportunity to expand this approach to other epithelial tumor types without the need to train a new algorithm for recognition of tumor cells in a separate indication. Importantly, two distinct approaches on different patient cohorts show that the spatial distribution of immune cells has predictive value for CI-based therapies suggesting biologically relevant mechanisms that deserve further exploration.

While our results are encouraging in terms of the overall correlation between the immune status of the tumor and patient outcome, there are still the outliers; we observed a certain degree of tumor IP heterogeneity among the patients who responded well to immunotherapy treatment. Incorporating clinical response data in the training of the model may optimize performance and identify features that could serve as additional biomarkers. However, we are also keenly aware of the inherent caveat of the approach of interrogating an environment of high plasticity such as the TME in archival tissues to predict therapy outcome. There is ample clinical and experimental evidence that tumor IP can change during immunotherapy [47] [48] [49] [50]; in vivo imaging approaches could circumvent the need for on-treatment biopsies for immune profiling [51].

We realize that our study – albeit on rather large patient cohorts – is based on retrospective analyses. Further optimization and validation in a prospective setting are necessary prior to routine clinical implementation.

Declarations

Ethics approval and consent to participate

All patients provided informed consent to participate in the respective clinical trials.

Consent for publication

All authors agree to the publication of this manuscript.

Availability of data and material

Data and materials are available upon reasonable request

The source code is available at https://github.com/geofly1985/Automated_IP

Competing interests

X.L., J.E., J.M.G., W.Z., A.Z., R.B., C.W.C., B.N., N.P., L.M., S.C., M.P., S.L., L.R., H.K. are employees of Genentech, Inc. X.L., J.E., D.O., H.K. are inventors on a patent related to automated immunophenotyping of tumors.

Funding

This work was fully funded by Genentech, Inc.

Contributors

X.L., J.E., D.O. and H.K contributed to the study design and wrote the paper. X.L., J.E., R.B., C.W.C., L.M., B.N., M.K., S.L., L.R., D.O., H.K. made critical contributions to the acquisition and/or analysis of data. All authors were involved in data interpretation and review of the manuscript.

Acknowledgements
We thank the patients, their families, and all of the investigators and their staff involved in the POPLAR, OAK and IMpassion130 studies. The authors would like to thank the Digital Pathology and Image Acquisition staff and the Histology Archive (both Genentech, Inc.) for scanning of slides, image acquisition and slide retrieval and management, respectively. We would like to

thank Robin Lorenz, Lisa McGinnis and Frank Peale for critical review of the manuscript and Meredith Triplett for excellent administrative support.

Figure 1. Manually assigned tumor IP and association with treatment outcome. Representative examples of Desert (**A**), Excluded (**B**) and Inflamed (**C** & **D**) immunophenotypes; areas from the same specimen demonstrating areas of Desert (**E**) and Excluded (**F**; intratumoral heterogeneity). Shown are five cases of NSCLC stained for CD8 and panCK. 200x magnification, scale bar indicates 100μm. Cut-off criteria for density and distribution of CD8+ cells for manual IP assignment; % indicate proportion of tumor area involved; 0, 1+, 2+ and 3+ indicate observed densities of CD8+ cells (**G**). Kaplan-Meier curves for OS for OAK and IMpassion130 according to manually assigned IP for patients receiving atezolizumab (**H**).

Figure 2. **Workflow and output of automated tumor immunophenotype classification pipelines.** Input for all four pipelines are whole slide images with manually annotated tumor areas, excluding artifacts (green rectangle; **A**) and manual IP calls from POPLAR. CD8+ and CK+ pixels are automatically identified in the manually annotated tumor area. CD8 density measurements within CK+ regions are the only input for the slide level density cutoff pipeline (#1). CD8 and CK regions are then incorporated in a tile-based analysis, with individual tile densities resulting in a multidimensional readout classified by a SVM into one of three immunophenotype classes, pipeline #2 (**B**). **C**. Data analysis workflow in MOCHA pipeline (#3): based on 50 spatial features (e.g., co-localization of CD8 T cells and tumor cells) extracted for

each tiled WSI, a random forest classifier with manual immunophenotype call is trained in a supervised learning framework.

Figure 3. Overlay of the manual tumor IP calls with the features generated by the respective automated analysis pipelines for POPLAR, OAK and IMpassion130. Distribution of patient samples based on the averaged slide level CD8 density readout in CK+ regions (pipeline #1); counts (y-axis) refer to the number of unique patient samples (**A**). UMAP plots based on: Binned CD8+ cell density in CK+ and CK- tiles (pipeline #2; **B**); a set of spatial features derived from tile based measurements (pipeline #3; **C**); a combination of spatial

features with binned tile-based CD8 density (pipeline #4; **D**). Cases are color-coded based on manually assigned tumor IP categories.

Figure 4. Concordance between manually and algorithmically assigned tumor IP calls and association with treatment outcome. Confusion matrices showing the distribution of manually assigned tumor IP calls (rows) among the automated IP calls (columns) for the four automated pipelines for OAK and IMpassion130 (**A**). Kaplan-Meier curves for OS for atezolizumab-containing treatment arms for OAK and IMpassion130 according to IP categories determined by MOCHA-BITE or manually (see suppl. Fig for PFS curves) **(B)**. Performance of the four automated pipelines based on agreement of IP classification using manual IP calls as ground truth; listed for each pipeline are the Cohen's kappa coefficient as well as median OS and PFS times for Desert (D), Excluded (E) and Inflamed (I) IP in descending order for the atezolizumab-containing (blue) and control arms (red) (**C**). Blue and red values indicate

statistically significant and non-significant differences, respectively according to the log-rank test, separation of OS (or PFS) Kaplan-Meier curves with the given IP categorization.

Figure 5. Association of IP with atezolizumab outcome in NSCLC and TNBC patients. Forest plot analysis and Kaplan-Meier curves for patients with "Inflamed" vs "Non-Inflamed"

(Excluded+Desert) tumors for OAK (**A**) and IMpassion130 (**B**). The vertical line of no effect in the forest plots indicates a hazard ratio (HR) of 1, shown are the number of patients in control and treatment group for each IP category, median survival times (MST) with confidence intervals (CI) and p values.

References

Korman AJ, Garrett-Thomson SC, Lonberg N. The foundations of immune checkpoint blockade and the ipilimumab approval decennial. *Nat Rev Drug Discov* 2021: 1-20

Robert C. A decade of immune-checkpoint inhibitors in cancer therapy. *Nat Commun* 2020; **11**: 3801

Marin-Acevedo JA, Kimbrough EO, Lou Y. Next generation of immune checkpoint inhibitors and beyond. *J Hematol Oncol* 2021; **14**: 45

Labrijn AF, Janmaat ML, Reichert JM, *et al.* Bispecific antibodies: a mechanistic review of the pipeline. *Nat Rev Drug Discov* 2019; **18**: 585-608

Bagchi S, Yuan R, Engleman EG. Immune Checkpoint Inhibitors for the Treatment of Cancer: Clinical Impact and Mechanisms of Response and Resistance. *Annu Rev Pathology Mech Dis* 2020; **16**: 1-27

Taube JM, Klein A, Brahmer JR, et al. Association of PD-1, PD-1 Ligands, and Other Features of the Tumor Immune Microenvironment with Response to Anti–PD-1 Therapy. *Clin Cancer Res* **20**14; 20: 5064-5074

Tsao MS, Kerr KM, Kockx M, et al. PD-L1 Immunohistochemistry Comparability Study in Real-Life Clinical Samples: Results of Blueprint Phase 2 Project. *J Thorac Oncol* 2018; **13**: 1302-1311

Rimm DL, Han G, Taube JM, et al. A Prospective, Multi-institutional, Pathologist-Based Assessment of 4 Immunohistochemistry Assays for PD-L1 Expression in Non–Small Cell *Lung Cancer*. *Jama Oncol* 2017; **3**: 1051

Doroshow DB, Bhalla S, Beasley MB, et al. PD-L1 as a biomarker of response to immune-checkpoint inhibitors. *Nat Rev Clin Oncol* 2021; **18**: 345-362

Kefford R, Ribas A, Hamid O, et al. Clinical efficacy and correlation with tumor PD-L1 expression in patients (pts) with melanoma (MEL) treated with the anti-PD-1 monoclonal antibody MK-3475. *J Clin Oncol* 2014; **32**: 3005-3005

Kockx MM, McCleland M, Koeppen H. Microenvironmental regulation of tumour immunity and response to immunotherapy. *J Pathology* 2021; **254**: **37**4-383

Bai R, Lv Z, Xu D, et al. Predictive biomarkers for cancer immunotherapy with immune checkpoint inhibitors. *Biomark Res* 2020; **8**: 34

13 Li N, Hou X, Huang S, et al. Biomarkers related to immune checkpoint inhibitors therapy. *Biomed Pharmacother* 2022; **147**: 112470

Johnson SK, Kerr KM, Chapman AD, et al. Immune cell infiltrates and prognosis in primary carcinoma of the lung. Lung Cancer 2000; **27**: 27-35

Galon J, Costes A, Sanchez-Cabo F, et al. Type, Density, and Location of Immune Cells Within Human Colorectal Tumors Predict Clinical Outcome. *Science* 2006; **313**: 1960-1964

Galon J, Angell HK, Bedognetti D, et al. The Continuum of Cancer Immunosurveillance: Prognostic, Predictive, and Mechanistic Signatures. *Immunity* 2013; **39**: 11-**26**

Pagès F, Mlecnik B, Marliot F, et al. International validation of the consensus Immunoscore for the classification of colon cancer: a prognostic and accuracy study. *Lancet* 2018; **391**: 2128-2139

Savas P, Salgado R, Denkert C, et al. Clinical relevance of host immunity in breast cancer: from TILs to the clinic. Nat Rev Clin Oncol 2016; 13: 228-241

Bairi KE, Haynes HR, Blackley E, et al. The tale of TILs in breast cancer: A report from The International Immuno-Oncology Biomarker Working Group. *Npj Breast Cancer* 2021; **7**: 150

Hendry S, Salgado R, Gevaert T, et al. Assessing Tumor-Infiltrating Lymphocytes in Solid Tumors. *Adv Anat Pathol* 2017; **24**: 311-335

Fuchs TL, Sioson L, Sheen A, et al. Assessment of Tumor-infiltrating Lymphocytes Using International TILs Working Group (ITWG) System Is a Strong Predictor of Overall Survival in Colorectal Carcinoma. *Am J Surg Pathology* 2020; **44**: 53**6**-544

Jong VMT de, Wang Y, Hoeve ND ter, et al. Prognostic Value of Stromal Tumor-Infiltrating Lymphocytes in Young, Node-Negative, Triple-Negative Breast Cancer Patients Who Did Not Receive (neo)Adjuvant Systemic Therapy. J Clin Oncol 2022: JCO2101536

Kos Z, Roblin E, Kim RS, et al. Pitfalls in assessing stromal tumor infiltrating lymphocytes (sTILs) in breast cancer. Npj Breast Cancer 2020; 6: 17

Amgad M, Stovgaard ES, Balslev E, et al. Report on computational assessment of Tumor Infiltrating Lymphocytes from the International Immuno-Oncology Biomarker Working Group. Npj Breast Cancer 2020; 6: 16

Ruiter EJ de, Roest RH de, Brakenhoff RH, et al. Digital pathology-aided assessment of tumor-infiltrating T lymphocytes in advanced stage, HPV-negative head and neck tumors. *Cancer Immunol Immunother* 2020; **69**: 581-591

Sun P, He J, Chao X, et al. A Computational Tumor-Infiltrating Lymphocyte Assessment Method Comparable with Visual Reporting Guidelines for Triple-Negative Breast Cancer. *Ebiomedicine* 2021; **70**: 103492

Albusayli R, Graham JD, Pathmanathan N, et al. Artificial intelligence based digital scores of stromal tumour-infiltrating lymphocytes and tumour-associated stroma predict disease specific survival in triple-negative breast cancer. J Pathol 2023

Acs B, Salgado R, Hartman J. What do we still need to learn on digitally assessed biomarkers? Ebiomedicine 2021; 70: 103520

Szabo PM, Lee G, Ely S, et al. CD8+ T cells in tumor parenchyma and stroma by image analysis (IA) and gene expression profiling (GEP): Potential biomarkers for immuno-oncology (I-O) therapy. J Clin Oncol 2019; 37: 2594-2594

Lee G, Desai K, Tang H, et al. 387 The utility of AI-powered spatial classification of intratumoral CD8+ immune-cell distribution in predicting overall survival in patients with melanoma as part of the checkMate 067 clinical trial. *J Immunother Cancer* 2021; **9**: A420-A420

Fehrenbacher L, Spira A, Ballinger M, et al. Atezolizumab versus docetaxel for patients with previously treated non-small-cell lung cancer (POPLAR): a multicentre, open-label, phase 2 randomised controlled trial. *Lancet Lond Engl* 2016; **387**: 1837-1846

Rittmeyer A, Barlesi F, Waterkamp D, et al. Atezolizumab versus docetaxel in patients with previously treated non-small-cell lung cancer (OAK): a phase 3, open-label, multicentre randomised controlled trial. Lancet Lond Engl 2016; **389**: 255-265

Schmid P, Adams S, Rugo HS, et al. Atezolizumab and Nab-Paclitaxel in Advanced Triple-Negative Breast Cancer. *New Engl J Med* 2018; **379**: 2108-2121

Chien L-C, Li X, Staudt A. Physical inactivity displays a mediator role in the association of diabetes and poverty: A spatiotemporal analysis. *Geospatial Health* 2016; **12**: 528

Li X, Staudt A, Chien L-C. Identifying counties vulnerable to diabetes from obesity prevalence in the United States: a spatiotemporal analysis. Geospatial Health 2015; **11**: 439

Li X, Gaire F, Jansen G, et al. 1381P Spatial-statistics-based modeling for predicting treatment response in non-small cell lung cancer (NSCLC) patients using H&E pathology images. *Ann Oncol* 2020; **31**: S879

Nawaz S, Heindl A, Koelble K, et al. Beyond immune density: critical role of spatial heterogeneity in estrogen receptor-negative breast cancer. *Modern Pathol* 2015; **28**: 766-777

Rempala GA, Seweryn M. Methods for diversity and overlap analysis in T-cell receptor populations. *J Math Biol* 2013; **67**: 1339-1368

Lee S-I. Developing a bivariate spatial association measure: An integration of Pearson's r and Moran's I. *J Geogr Syst* 2001; **3**: 369-385

Geary RC. The Contiguity Ratio and Statistical Mapping. *Incorporated Statistician* 19**5**4; 5: 115-141

Anselin L. Local Indicators of Spatial Association—LISA. *Geogr Anal* 1995; **27**: 93-115

Karrer TM, Garmhausen M, Li X, et al. Combining hematoxylin and eosin (H&E) stained images and RNA sequencing (RNA-Seq) data to predict overall survival (OS) in patients with non-small cell lung cancer (NSCLC). J Clin Oncol 2021; 39: 1547-1547

Hussein SE, Chen P, Medeiros LJ, et al. Artificial intelligence strategy integrating morphologic and architectural biomarkers provides robust diagnostic accuracy for disease progression in chronic lymphocytic leukemia. J Pathol 2022; **256**: 4-14

Allam M, Hu T, Lee J, et al. Spatially variant immune infiltration scoring in human cancer tissues. *Npj Precis Oncol* 2022; **6**: 60

Angell HK, Bruni D, Barrett JC, et al. The Immunoscore: Colon Cancer and Beyond. Clin Cancer Res 2020; 26: 332-339

Park S, Ock C-Y, Kim H, et al. Artificial Intelligence-Powered Spatial Analysis of Tumor-Infiltrating Lymphocytes as Complementary Biomarker for Immune Checkpoint Inhibition in Non-Small-Cell Lung Cancer. *J Clin Oncol Official J Am Soc Clin Oncol* 2022: JCO2102010

Tumeh PC, Harview CL, Yearley JH, et al. PD-1 blockade induces responses by inhibiting adaptive immune resistance. *Nature* 2014; **515**: 568-571

Westergaard MCW, Milne K, Pedersen M, et al. Changes in the Tumor Immune Microenvironment during Disease Progression in Patients with Ovarian Cancer. *Cancers* 2020; **12**: 3828

Yang J, Zhang Q, Wang J, et al. Dynamic profiling of immune microenvironment during pancreatic cancer development suggests early intervention and combination strategy of immunotherapy. Ebiomedicine 2022; **78**: 103958

Murciano-Goroff YR, Warner AB, Wolchok JD. The future of cancer immunotherapy: microenvironment-targeting combinations. *Cell Res* 2020; **30**: 507-519

Ruijter LK de, Donk PP van de, Hooiveld-Noeken JS, et al. Whole-body CD8+ T cell visualization before and during cancer immunotherapy: a phase 1/2 trial. *Nat Med* 2022; **28**: 2601-2610