Sex-dependent contributions of ventrolateral orbitofrontal cortex and basolateral amygdala to learning under uncertainty

Running Title: stimulus and action-based reversal learning

C.G. Aguirre[1]#, J.H. Woo[2]#, J.L.R Sosa[1], J. J. Munier[3], J. Perez[1], M. Goldfarb[1], K. Das[1], M. Gomez[1], T. Ye[1], J. Pannu[3], K. Evans[1], P.R. O'Neill[4], I. Spigelman[3], A. Soltani[2], and A. Izquierdo[1]

[1]Department of Psychology, University of California, Los Angeles, Los Angeles, CA 90095

[2]Department of Psychological and Brain Sciences, Dartmouth College, Hanover, NH 03755

[3]Section of Biosystems and Function, School of Dentistry, University of California, Los Angeles, Los Angeles, CA 90095

[4]Shirley and Stefan Hatos Center for Neuropharmacology, Department of Psychiatry and Biobehavioral Sciences, University of California Los Angeles, Los Angeles, CA 90095

#**co-first authors**

Author contributions: CGA and AI designed the research; CGA, JLRS, JM, JP, MG, KD, MG, TY, KE, and JP performed the research; CGA, JHW, JM, AS and AI analyzed the data; CGA, JHW, PRO, IS, AS, and AI interpreted the data; AI, AS, PRO, and IS acquired funding for the project; CGA and AI wrote the paper; all authors edited the final version.

Acknowledgements: This work was supported by UCLA's Division of Life Sciences Retention fund (Izquierdo), National Institutes of Health grants R01 DA047870 (Izquierdo and Soltani), R21 MH122800 (Izquierdo and Blair), R01AA024527 (Spigelman), K01 DA042219 (O'Neill), and F31 AA028183 (Munier), the NSF GRFP, Cota-Robles Fellowship, and Charles E. and Sue K. Young Fellowship (Aguirre), Ursula Mandel Fellowship and Graduate Research Mentorship award (Romero-Sosa), and the Training program in Neurotechnology Translation T32 NS115753 (Ye). We acknowledge the Staglin Center for Brain and Behavioral Health for additional support related to fluorescence microscopy. We also thank the NIDA Drug Supply program for the supply of clozapine-N-oxide.

Correspondence: *Alicia Izquierdo, Ph.D. email: aizquie@psych.ucla.edu Ph: +1 310 825 3459

Manuscript Information:
Pages   47
Figures   5   Extended Data Figures 5

Abstract

Reversal learning measures the ability to form flexible associations between choice outcomes with stimuli and actions that precede them. This type of learning is thought to rely on several cortical and subcortical areas, including highly interconnected orbitofrontal cortex (OFC) and basolateral amygdala (BLA), and is often impaired in various neuropsychiatric and substance use disorders. However, unique contributions of these regions to stimulus- and action-based reversal learning have not been systematically compared using a chemogenetic approach and particularly before and after the first reversal that introduces new uncertainty. Here, we examined the roles of ventrolateral OFC (vlOFC) and BLA during reversal learning. Male and female rats were prepared with inhibitory DREADDs targeting these regions and tested on a series of deterministic and probabilistic reversals during which they learned about stimulus identity or side (left or right) associated with different reward probabilities. Using a counterbalanced within-subject design, we inhibited these regions prior to reversal sessions. We measured initial and pre-post reversal changes in accuracy to measure first detection and further adjustment to reversals, respectively. We found that inhibition of vlOFC, but not BLA, eliminated detection of stimulus-based reversals. Conversely, both BLA and vlOFC inhibition resulted in significantly slower action-based reversal learning in females, not males, indicating a sex-dependent role for these regions in this type of learning. Further, learning in females was more impacted in first reversal by vlOFC inhibition than inhibition of BLA, the latter more involved in probabilistic reversal learning. These findings add to mounting evidence of sex-dependent learning flexibility.

Keywords: **stimulus learning, action learning, deterministic, probabilistic, reward learning, DREADDs**

Significance Statement

Inflexible learning is a feature of several neuropsychiatric disorders. We investigated how the ventrolateral orbitofrontal cortex (vlOFC) and basolateral amygdala (BLA) are involved in learning of stimuli or actions under different forms of uncertainty. Following chemogenetic inhibition of these regions in both male and females, we measured detection and adjustment to fully-predictive (i.e., deterministic) and subsequent probabilistic reversals. For action learning, vlOFC and BLA exhibited a sex-dependent role in deterministic and probabilistic learning with females exhibiting the slowest learning following inhibition. For stimulus learning, vlOFC, but not BLA, was required for reversal detection and adjustment. These findings provide insight into the mechanisms of learning under different forms of uncertainty and the sex-dependency of these mechanisms.

Introduction

Reversal learning, impacted in various neuropsychiatric conditions, measures subjects' ability to form flexible associations between stimuli and actions with outcomes (Schoenbaum et al., 2003; Izquierdo et al., 2013; Dalton et al., 2016). Reversal learning tasks can also be used to probe learning following expected and unexpected uncertainty in the reward environment (Behrens et al., 2007; Jang et al., 2015; Winstanley and Floresco, 2016; Soltani and Izquierdo, 2019). For example, after the experience of the first reversal, all others are expected to some extent (Jang et al., 2015). Additionally, unexpected uncertainty can be introduced by changes in reward probabilities, after taking the baseline, expected uncertainty into account.

The basolateral amygdala (BLA) is an area of interest in reversal learning due its involvement in value updating (Tye and Janak, 2007; Janak and Tye, 2015; Wassum and Izquierdo, 2015; Groman et al., 2019) and the encoding of both stimulus-outcome and action-outcome associations typically probed in Pavlovian-to-Instrumental tasks (Corbit and Balleine, 2005; Lichtenberg et al., 2017; Malvaez et al., 2019; Sias et al., 2021). Manipulations of amygdala and specifically BLA have resulted in reversal learning impairments (Schoenbaum et al., 2003; Churchwell et al., 2009; Groman et al., 2019), impaired learning from positive feedback (Costa et al., 2016; Groman et al., 2019), enhanced learning from negative feedback (Rudebeck and Murray, 2008; Izquierdo et al., 2013; Taswell et al., 2021), and even improvements of deficits produced by OFC lesions (Stalnaker et al., 2007). Yet BLA has not been extensively studied in the context of flexible reversal learning of stimuli vs. actions with the exception of a recent lesion study in rhesus macaques (Taswell et al., 2021). BLA has also not been systematically evaluated for its contributions to deterministic vs. probabilistic schedules, with the exception of another lesion study in monkeys (Costa et al., 2016). The idea that BLA encodes changes in the environment in terms of salience and

associability (Roesch et al., 2010) suggests this region may facilitate rapid updating to incorporate

new information. The contribution of BLA to reversal learning and its dependence on the nature

of the association (i.e., stimulus- vs. action-based), sensory modality (i.e., visual), and type of

uncertainty introduced by the task design (i.e., deterministic vs. probabilistic, but also first reversal

versus all subsequent reversals) has not been extensively studied using a chemogenetic approach

in rats.

In parallel, studies with manipulations in rat OFC in reversal learning have included targeting

of the entire ventral surface (Izquierdo, 2017), or systematic comparisons of medial vs. lateral OFC

(Hervig et al., 2020; Verharen et al., 2020). Here we examined the role of vlOFC, a subregion not

as often probed in reward learning as medial and more (dorso)lateral OFC (cf. Zimmermann et al.

(2018)) but also densely-interconnected with BLA (Barreiros et al., 2021a; Barreiros et al., 2021b).

Additionally, unlike almost all previous studies on reversal learning, we included both male and

female subjects.

Using a within-subject counterbalanced design, we inactivated these regions prior to reversal

sessions and measured both learning and detection/adjustment to reversals. We found that vlOFC,

but not BLA, inhibition impaired detection of deterministic and probabilistic stimulus-based,

reversals. Conversely, BLA and vlOFC inhibition resulted in significantly slower action-based

reversal learning in females, but not males, suggesting a sex-dependent role for these regions.

Learning in females was more impacted in first (deterministic) reversal by vlOFC inhibition, and

more robustly affected by BLA inhibition in the subsequent probabilistic reversal. These results

suggest similar roles for vlOFC and BLA in flexible action-based learning, but a more specialized

role for vlOFC in setting adjustments in stimulus-based learning. Further, fitting choice data with

reinforcement learning models indicated the attenuated probabilistic action-based reversal learning

deficits were mediated by a larger memory decay for the unchosen option, especially following

vlOFC inhibition. Finally, our findings underscore the importance of including male and female

animals in neuroscience studies, adding to mounting evidence of sex-modulated learning (Chen et

al., 2021a; Chen et al., 2021b).

Materials and Methods

Subjects

Animals for behavioral experiments were adult (N=56, 25 females; 52 used for behavioral

study and 4 for *ex vivo* imaging) Long-Evans rats (Charles River Laboratories) average age post-

natal-day (PND) 65 at the start of experiments, with a 280g body weight minimum for males and

240g body weight minimum for females at the time of surgery and the start of the experiment. Rats

were approximately PND 100 (emerging adulthood; Ghasemi et al. (2021)) when behavioral

testing commenced. Before any treatment, all rats underwent a 3-day acclimation period during

which they were pair-housed and given food and water *ad libitum*. During that time, they remained

in their home cage with no experimenter interference. Following this 3-day acclimation period,

animals were handled for 10 min per animal for 5 consecutive days. During the handling period,

the animals were also provided food and water *ad libitum*. After the handling period, animals were

individually-housed under standard housing conditions (room temperature 22–24° C) with a

standard 12 h light/dark cycle (lights on at 6am). Animals were then surgerized and tested on

discrimination and reversal learning 1-week post-surgery. At the point of reversal, they were

beyond the 3-week expression time for Designer Receptors Exclusively Activated by Designer

Drugs (DREADDs).

A separate group of Long-Evans rats (N=4, all males) were used for validation of

effectiveness of DREADDs in slides of BLA and vlOFC, using *ex vivo* calcium imaging

procedures. All procedures were conducted in accordance to the recommendations in the Guide

for the Care and Use of Laboratory Animals of the National Institutes of Health and with the

approval of the Chancellor's Animal Research Committee at the University of California, Los

Angeles.

Surgery

Viral Constructs

Rats were singly-housed and remained in home cages for 4 weeks prior to testing while the

inhibitory hM4Di DREADDs expressed in BLA (n=16, 9 females), vlOFC (n=19, 9 females), or

eGFP control virus (n=17, 7 females) in these regions. In rats tested on behavior, an adeno-

associated virus AAV8 driving the hM4Di-mCherry sequence under the CaMKIIa promoter was

used to express DREADDs bilaterally in BLA neurons (0.1 μl, AP = -2.5; ML= ±5; DV = −7.8

and 0.2 μl, AP= -2.5; ML= ±5; DV= −8.1, from bregma at a rate of 0.1 μl/min; AAV8-CaMKIIa-

hM4D(Gi)-mCherry, Addgene, viral prep #50477-AAV8). In other animals, this same virus

(AAV8-CaMKIIa-hM4Di-mCherry, Addgene) was bilaterally infused into two sites in vlOFC (0.2

μl, AP = +3.7; ML= ±2.5; DV = −4.6 and 0.15 μl, AP= 4; ML= ±2.5; DV= −4.4, from bregma at

a rate of 0.1 μl/min). A virus lacking the hM4Di DREADD gene and only containing the green

fluorescent tag eGFP (AAV8-CaMKIIa-eGFP, Addgene) was also infused bilaterally into either

BLA (n=7), vlOFC (n=5), or anterior cingulate cortex [(n=5); 0.3 μl, AP = +3.7; ML= ±2.5; DV =

−4.6, rate of 0.1 μl/min] as null virus controls. Our vlOFC targeting is most similar to infusion

sites reported previously by Dalton et al. (2016), and 0.7 mm more medial than others (Costa et

al., 2023). In rats used for *ex vivo* calcium imaging, the same target regions were infused with

either GCaMP6f (AAV9-CaMKIIa-GCaMP6f, Addgene), a 1:1 combination of GCaMP6f+mCherry

(AAV8-CamKIIa-mCherry, Vector BioLabs, #VB1947), or a 1:1 combination of

GCaMP6f+hM4Di-mCherry (same as used for behavior, AAV8-CaMKIIa-hM4Di-mCherry,

Addgene).

Figure 1. Bilateral targeting of basolateral amygdala (BLA) and ventrolateral orbitofrontal cortex (vlOFC) and confirmation of effective DREADDs inhibition using ex vivo Ca²⁺ imaging in slices. *(A) Photomicrograph of hM4Di-mCherry DREADDs expression in BLA. Numerals indicate mm anterior to Bregma. (B) Reconstructions of viral expression of hM4Di (magenta) and enhanced green fluorescent protein, eGFP (green) in BLA. The more intense colors represent regions where expression overlapped the most across animals. (C) In BLA neurons expressing GCaMP6f and GCaMP6f+mCherry, application of CNO (10μM) in the presence of picrotoxin (50 µM) had no effect on the frequency of elicited Ca²⁺ events (top left and right: example of single cell traces, bottom left: Ca²⁺ event changes, each line is a cell). In BLA neurons that expressed hM4Di, there was a reduction in the frequency of elicited Ca²⁺ events during CNO application (bottom right). (D) Photomicrograph of hM4Di-mCherry DREADDs expression in vlOFC. Numerals indicate mm anterior to Bregma. (E) Reconstruction of viral expression of hM4Di (magenta) and enhanced green fluorescent protein, eGFP (green). The more intense colors represent regions where expression overlapped the most across animals. (F) In vlOFC neurons expressing GCaMP6f and GCaMP6f+mCherry, application of CNO (10μM) had no effect on the frequency of elicited Ca²⁺ events (top left and right: example of single cell traces, bottom left: Ca²⁺ event changes, each line is a cell). In vlOFC neurons that expressed hM4Di, there was a reduction in the frequency of Ca²⁺ events during CNO application (bottom right). n=2-5 slices/rat, 2-way ANOVA and multiple comparison tests ***p<0.001, *p<0.05.*

Surgical Procedure

Infusions of DREADD or eGFP control virus were performed using aseptic stereotaxic

techniques under isoflurane gas (1-5% in O_2) anesthesia prior to any behavioral testing experience.

Before surgeries were completed, all animals were administered 5 mg/kg s.c. carprofen (NADA

#141–199, Pfizer, Inc., Drug Labeler Code: 000069) and 1cc saline. After being placed in the

stereotaxic apparatus (David Kopf; model 306041), the scalp was incised and retracted. The skull

was leveled with a +/- 0.3 mm tolerance on the A-P to ensure that bregma and lambda were in the

same horizontal plane. Small burr holes were drilled in the skull above the infusion target. Virus

was bilaterally infused at a rate of 0.01 µl per minute in target regions (coordinates above). After

each infusion, 5 min elapsed before exiting the brain.

Histology

At the end of the experiment, rats were euthanized with an overdose of Euthasol (Euthasol,

0.8 mL, 390 mg/mL pentobarbital, 50 mg/mL phenytoin; Virbac, Fort Worth, TX), were

transcardially perfused, and their brains removed for histological processing. Brains were fixed in

10% buffered formalin acetate for 24 h followed by 30% sucrose for 5 days. To visualize hM4Di-

mCherry and eGFP expression in BLA or vlOFC cell bodies, free-floating 40-µm coronal sections

were mounted onto slides and cover-slipped with mounting medium for DAPI. Slices were

visualized using a BZ-X710 microscope (Keyence, Itasca, IL), and analyzed with BZ-X Viewer

and analysis software.

Reconstructions of viral expressions of hM4Di (magenta) and green fluorescent protein,

eGFP (green) across the AP plane (**Fig. 1BE**) were conducted using Photoshop and Illustrator

(Adobe, Inc., San Jose, CA) by individuals blind to condition. Two independent raters then used

ImageJ (U. S. National Institutes of Health, Bethesda, Maryland, USA) to trace and quantify pixels

at AP +3.7 (vlOFC) and AP -2.8 (BLA) for each animal. Three measures were obtained per

hemisphere and were highly correlated (Spearman rank correlation: $r = 0.93$, $p < 0.01$). Only

subjects with bilateral expression were included in behavioral analyses (3 BLA rats excluded due

to unilateral expression). There were no differences in expression level between males and females

for pixel count reconstructions [$F_{(1,12)} = 0.32$, $p = 0.58$], but there was a significant difference

between target brain region (vlOFC, BLA) in the hM4Di virus group [$F_{(1,12)} = 9.71$, $p = 0.009$]; the

latter was expected given the larger infusion volumes in vlOFC.

Food Restriction

Five days prior to any behavioral testing, rats were placed on food restriction with females

on average maintained on 10-12 grams/ day and males given 12-14 grams/ day of chow. Food

restriction level remained unchanged throughout behavioral testing, provided animals completed

testing sessions. Water remained freely available in the home cage. Animals were weighed every

other day and monitored closely to not fall below 85% of their maximum, free-feeding weight.

Drug administration

Inhibition of vlOFC or BLA was achieved by systemic administration of clozapine-N-

oxide, CNO (i.p., 3mg/kg in 95% saline, 5% DMSO) in animals with DREADDs. Rats with eGFP

in these regions underwent identical drug treatment. Rats were randomly assigned to drug

treatment groups, irrespective of performance in pretraining. CNO was administered only during

reversal learning, 30 min prior to behavioral testing. We followed previous work on timing and

dose of systemic CNO (Stolyarova, Rakhshan et al., 2019; Hart et al., 2020) and considering the

long duration of test sessions. To control for nonspecific effects of injections and handling stress,

we also injected animals with saline vehicle (VEH). To increase power and decrease the number

of animals used in experiments, we used a within-subject design for assessing the effects of CNO,

with all rats receiving CNO and VEH injections in a counterbalanced order. Thus, for drug

administration if a rat received CNO on the first reversal (R1), it was administered VEH on the

second reversal (R2), CNO on the third reversal (R3), and VEH on the fourth reversal (R4), or vice

versa: VEH on R1, CNO on R2, VEH on R3, and CNO on R4.

Behavioral Testing

Pretraining. Behavioral testing was conducted in operant conditioning chambers outfitted

with an LCD touchscreen opposing the sugar pellet dispenser. All chamber equipment was

controlled by customized ABET II TOUCH software (Lafayette Instrument Co., Lafayette, IN).

The pretraining protocol, adapted from established procedures (Stolyarova and Izquierdo,

2017), consisted of a series of phases: Habituation, Initiation Touch to Center Training (ITCT),

Immediate Reward Training (IMT), designed to train rats to nose poke, initiate a trial, and select a

stimulus to obtain a reward (i.e., sucrose pellet). Pretraining stages have been reported in detail

elsewhere (Stolyarova, Rakhshan et al., 2019). For habituation pretraining, the criterion for

advancement was collection of all 5 sucrose pellets. For ITCT, the criterion to the next stage was

set to 60 rewards consumed in 45 min. The criterion for IMT was set to 60 rewards consumed in

45 min across two consecutive days. After completion of all pretraining schedules, rats were

advanced to the discrimination (initial) phase of either the action- or stimulus-based reversal

learning task, with the task order counterbalanced (**Fig. 2AB** or **Fig. 4AB**). A subset of animals

was tested first on the action-based task (10 vlOFC hM4Di, 9 BLA hM4Di), while others were

tested first on the stimulus-based task (9 vlOFC hM4Di, 7 BLA hM4Di, 17 eGFP). Three vlOFC

rats completed only the stimulus-based task.

Action-based deterministic discrimination learning. After completion of either all

pretraining schedules or all four reversals of the stimulus-based task, rats were advanced to the

discrimination (initial) phase of the action-based task (**Fig. 2A**). Rats were required to initiate a

trial by touching the white graphic stimulus in the center screen (displayed for 40 s), and after

initiation rats would be presented with two stimuli (i.e., fan or marble) on the left and right side of

the screen (displayed for 60 s). Rats could nosepoke either the spatial side rewarded with one

sucrose pellet (the better option, $p_R(B)=1.0;$) or the spatial side that went unrewarded (the worse

option, $p_R(W)=0.0$). Thus, rats were required to ignore the properties of the stimuli and determine

the better rewarded side. If a side was not selected, it was scored as a choice omission, and a 10 s

inter-trial interval (ITI) ensued. If a trial was not rewarded, a 5 s time-out would occur, followed

by a 10 s ITI. If a trial was rewarded, a 10 s ITI would occur after the reward was collected. The

criterion was set to 60 or more rewards consumed and selection of the correct option in 75% of the

trials or higher during a 60 min session across two consecutive days. After reaching the criterion

for the discrimination phase, the rats were advanced to the reversal phase beginning on the next

session. Animals were not administered either CNO or VEH injections during discrimination

learning.

Action-based reversal learning. After the discrimination phase, the rats advanced to the

reversal phase. Before a reversal learning session, rats were injected intraperitoneally with either

3 mg/kg of CNO or VEH 30 min prior to each reversal testing session. The side previously

associated with the $p_R(B)=1.0$ probability was now associated with a $p_R(W)=0.0$ probability of

being rewarded, and vice versa. The criterion was the same as the deterministic discrimination

phase. After reaching the criterion for the first deterministic reversal phase (i.e., R1), the rats

advanced to the second deterministic reversal phase (i.e., R2) beginning on the next session. Rats

that had previously received VEH during the first reversal would now receive CNO injections, and

vice versa.

268   After completing both deterministic reversal learning phases, rats advanced to the first

probabilistic reversal learning phase (i.e., reversal 3, R3). Rats underwent the same injection

procedure as the prior reversals. However, the spatial side (i.e., left or right) previously associated

with $p_R(B)=1.0$, was now associated with a $p_R(W)=0.1$ probability of being rewarded, whereas the

spatial side previously associated with $p_R(W)=0.0$ probability, was now associated with $p_R(B)=0.9$.

The criterion was the same as the previous deterministic reversal learning phases. After reaching

the criterion for the first probabilistic reversal learning phase (i.e., reversal 3, R3), rats were

advanced to the second probabilistic reversal phase (i.e., reversal 4, R4) beginning on the next

testing day, where the probabilities would be reversed once again. Rats that had previously

received VEH during the first probabilistic reversal now received CNO injections, and vice versa.

278   *Stimulus-based deterministic discrimination learning.* After completion of all pretraining

schedules (or all reversals of the action-based task), rats were advanced to the discrimination

(initial) phase of learning in which they would initiate a trial by touching a white graphic stimulus

in the center screen (displayed for 40 s), and choose between two different visual stimuli

pseudorandomly presented on the left and right side of the screen. Stimuli were displayed for 60 s

each, randomly assigned as the better or worse stimulus: $p_R(B)=1.0$ or $p_R(W)=0.0$. If a trial was

not initiated within 40 s, it was scored as an initiation omission. If a stimulus was not selected, it

was scored as a choice omission, and a 10 s ITI ensued. If a trial was not rewarded, a 5 s time-out

would occur, followed by a 10 s ITI. Finally, if a trial was rewarded, a 10 s ITI would follow after

the reward was collected. A criterion was set to 60 or more rewards consumed and selection of the

correct option in 75% of the trials or higher during a 60 min session across two consecutive days.

After reaching the criterion for the discrimination phase or if rats were unable to achieve criterion

after 10 days, rats were advanced to the reversal phase beginning on the next session. Animals

were not administered either CNO or VEH injections during discrimination learning.

Stimulus-based reversal learning. After the discrimination phase, rats advanced to the first

deterministic reversal learning phase (i.e., reversal 1, R1) where they were required to remap

stimulus-reward contingencies. As above, before a reversal learning session, rats were injected

intraperitoneally with either 3 mg/kg of CNO or VEH 30 min prior to each reversal testing session.

The criterion was the same as discrimination learning. After reaching the criterion for the first

reversal phase or if they were unable to achieve criterion after 10 days, the rats were advanced to

the second deterministic reversal phase (i.e., reversal 2, R2) beginning on the next testing day,

where the reward contingencies were reversed once again. Rats that had previously received VEH

during the first reversal now received CNO injections, and vice versa.

After completing both deterministic reversal learning phases, rats advanced to the first

probabilistic reversal learning phase (i.e., reversal 3, R3). The injection procedure remained the

same as prior reversals. However, the visual stimulus previously associated with $p_R(B)=1.0$, would

now be associated with $p_R(W)=0.1$, whereas the stimulus previously associated with $p_R(W)=0.0$,

would now be associated with $p_R(B)=0.9$. The criterion remained the same as prior reversals. After

reaching the criterion for the first probabilistic reversal learning phase or if rats were unable to

achieve criterion after 10 days, the rats were advanced to the second probabilistic reversal phase

(i.e., reversal 4, R4) beginning on the next testing day, where the probabilities would be reversed

once again. As above for action-based reversal learning, rats that had previously received VEH

during the first probabilistic reversal now received CNO injections, and vice versa.

Ex vivo **calcium imaging**

In N=4 animals (all males), following >3 weeks following stereotaxic viral injections, rats

(n=1 rat/brain region/virus combination; n=2-5 slices/rat) were deeply anesthetized with isoflurane

(Patterson Veterinary, MA, USA), decapitated and brains submerged in ice cold oxygenated

(95/5% O_2/CO_2) slicing artificial cerebrospinal fluid (ACSF) containing (in mM): 62 NaCl, 3.5

KCl, 1.25 NaH_2PO_4, 62 choline chloride, 0.5 $CaCl_2$, 3.5 $MgCl_2$, 26 $NaHCO_3$, 5 N-acetyl L-

cysteine, and 5 glucose, pH adjusted to 7.3 with KOH. Acutely microdissected vlOFC or BLA

slices (300 μM thick) were obtained (VT1200s, Leica, Buffalo Grove, IL) and transferred to room

temperature normal ACSF containing (in mM): 125 NaCl, 2.5 KCl, 1.25 NaH_2PO_4, 2 $CaCl_2$, 2

$MgCl_2$, 26 $NaHCO_3$, 10 glucose, pH adjusted to 7.3 with KOH, and allowed equilibrate for >1 hr

prior to transfer into a perfusion chamber for imaging.

Imaging was performed on a Scientifica SliceScope, with imaging components built on an

Olympus BX51 upright fluorescence microscope equipped with an sCMOS camera (Hamamatsu

Orca Flash 4.0v3). Anatomical regions in brain sections for Ca^{2+} imaging were first identified by

brightfield imaging with 780nm LED (Scientifica) illumination. Ca^{2+} imaging was performed

using a 40x, 0.80NA water immersion objective (Olympus), continuous 470nm LED illumination

(ThorLabs), and a filter cube suitable for GCaMP6f imaging: Excitation: Brightline 466/40,

Dichroic: Semrock FF495-Di03, Emission: Brightline 525/50. Slices were housed on poly-D

lysine cover slips attached to RC-26G chamber (Warner Instruments, Holliston, MA), which was

modified with platinum wires to apply electric field stimulation. Images were acquired continually

with 20 ms exposure time. Electric field stimulation was applied at 110 mV (twin pulse every 5s).

Temperature of ACSF during the recorded sessions was held at 28° C to minimize bubble

formation.

Calcium data extraction

Prior to imaging sessions, 40x images of red and green fluorescence were captured and

subsequently overlaid for post-hoc genotyping of individual cells (GCaMP6f[+],

GCaMP6f[+]/hM4Di-mCherry[+], or GCaMP6f/mCherry[+]). Blinded scorers semi-manually curated

regions of interest (ROIs) using Python-based Suite2P software (Pachitariu et al., 2017). ROI

fluorescence was subtracted from the annular surround fluorescence, low-pass filtered, and

transformed to dF/F_0 as previously described (Asrican and Song, 2021) where F_0 is calculated with

a boxcar filter with a 200-frame lookback window. dF/F_0 values were clipped between 0 and 9000

to eliminate negative changes. Area under the curve and event frequency of each cell was

calculated for each drug treatment. A threshold of 0.15 dF/F was used to determine significant

events, which is lower than the dF/F of a single *ex vivo* action potential, but significantly above

signal to noise in our recorded traces (Tada et al., 2014), **Fig. 1CF**.

Data Analyses

MATLAB (MathWorks, Natick, Massachusetts; Version R2021a) was used for all

statistical analyses and figure preparation. Data were analyzed with a series of mixed-effects

General Linear Models (GLM) for the discrimination learning phase to establish there were no

baseline differences in learning measures between the hM4Di and eGFP animals (i.e., virus group)

prior to any drug treatment for each task separately. Mixed-effects GLMs were also conducted on

reversal phases, with all fixed factors included in the model [i.e., reversal number (1-4), virus

group (hM4Di, eGFP), drug (CNO, VEH), sex (female, male), drug order (CNO1, VEH1)] and

individual rat as a random factor. These GLMs were run for each task type (stimulus- and action-

based tasks) separately. Since learning reached asymptote at 5-days for stimulus-based reversal

learning, only the first 5 days were included in the GLM. Similarly, since rats typically reached a

plateau (and criterion) at 150 trials for action-based reversal learning, we included only the first

150 trials in the GLM. Significant reversal number and/or drug order interactions were further

analyzed with a narrower set of fixed factors and Bonferroni-corrected post-hoc comparisons. In

the instance where sex was found a significant predictor (moderator), sex was entered as a

covariate factor in subsequent reversals. Accuracy (probability correct) before and after a reversal

(-3 and +3 sessions surrounding a reversal) was analyzed using ANOVA with virus group (hM4Di,

eGFP) and drug order (CNO1, VEH1) as fixed factors on the average change pre-post reversal.

Virus expression level was analyzed with ANOVA by sex (male, female) and region (vlOFC,

BLA) on pixel counts.

Dependent measures for learning included probability of choosing the correct or better

option, initiation latencies and omissions (failure to initiate a trial, latency to initiate a trial,

respectively), correct and incorrect choice latencies (latency to select the correct or better stimulus

or spatial side and latency to select the incorrect or wrong stimulus or spatial side, respectively),

reward latencies (latency to collect the reward), probability of win-stay, and probability of lose-

shift. Probability of win-stay and lose-shift adaptive strategies were calculated for the stimulus-

based task such that each trial was classified as a *win* if an animal received a sucrose pellet, and as

a *loss* if no reward was delivered. Statistical significance was noted when p-values were less than

0.05. All Bonferroni post-hoc tests were corrected for number of comparisons.

To analyze *ex vivo* calcium imaging data, 2-way ANOVAs with drug and virus as factors

were conducted to compare calcium event changes in GCaMP6f and GCaMP6f+mCherry in each

brain region for control experiments, and for GCaMP6f and GCaMP6f+hM4Di-mCherry in each

brain region for the experimental group. Tests corrected for number of comparisons were

conducted for interactions.

Reinforcement Learning Models

To capture the differences in learning and choice behavior during the action-based task, we

utilized two conventional reinforcement learning (RL) models. Specifically, the subjective

estimate of reward (V) for each choice option was updated on a trial-by-trial basis using reward

prediction error (RPE), the discrepancy between actual and expected reward value. In the first

model, which we refer to as *RL*, the value estimate of the chosen option (V_C) for a trial t was

updated using the following equations:

$$V_C(t + 1) = V_C(t) + \alpha\big(R(t) - V_C(t)\big), \qquad (1)$$

where $R(t)$ indicates the presence (1) or absence (0) of a reward for the given trial, and α is the

learning rate dictating the amount of update in the value estimate by RPE. In this model, the value

of unchosen option was not updated.

The second model, referred to as RL_{decay}, used the same learning rule as Equation (1) for

updating the value of the chosen option, and additionally updated the value of the unchosen option

(V_U) as follows:

$$V_U(t + 1) = V_U(t) - \gamma_d(V_U(t)), \qquad (2)$$

where γ_d is a decay rate controlling the amount of passive decay in value of the unchosen option.

In both models above, the probability of choosing a particular option was computed using the

following decision rule:

$$P_i(t) = (1 + e^{-\beta(V_i(t) - V_j(t))})^{-1}, \qquad (3)$$

where i and j corresponds to two alternative options (i.e., left and right for action-based task), and

β is the inverse temperature or sensitivity governing the extent to which higher-valued options are

consistently selected.

We used the standard maximum likelihood estimation method to fit choice data and

estimate the parameters for each session of the experiment. The values of the learning rate α and

decay rate γ_d were bounded between 0 and 1, and β was bounded between 1 and 100. Initial

parameter values were selected from this range, and fitting was performed using the MATLAB

function *fmincon*. For each set of parameters fitted to each session, we repeated 30 different initial

conditions selected from evenly-spaced search space to avoid local minima. The best fit was

selected from the iteration with the minimum negative log-likelihood (*LL*). For the first model

(*RL*), we treated the uninitiated or uncommitted trials with no choice data as if they had not

occurred. In contrast, for the second model (*RL_decay*), both choice options were considered

unchosen for those trials and both of the value estimates decayed passively according to Equation

(2).

To quantify goodness of fit, we computed both Akaike Information Criterion (AIC) and

Bayesian Information Criterion (BIC) for each session as follows:

$$AIC = -2 * LL + 2 * k, \qquad (4)$$

$$BIC = -2 * LL + \ln(n) * k, \qquad (5)$$

where k is the number of parameters in the model (two for RL and three for RL_{decay}), and n is the

number of choice trials in the session.

Results

Ex vivo calcium imaging in slices

We performed ex *vivo* Ca^{2+} imaging to confirm the selective action on CaMKII$^+$ neuronal

excitability in vlOFC and BLA in rats expressing hM4Di DREADD vs. controls expressing

mCherry. In BLA, there was no significant effect of CNO (10μM) on Ca^{2+} events for neurons

expressing GCaMP6f or GCaMP6f+mCherry (**Fig. 1C**). A 2-way ANOVA resulted in a significant

drug × virus interaction [$F_{(2,324)} = 3.367$, $p = 0.036$], with a selective reduction in the frequency of

elicited Ca^{2+} events during CNO only in neurons expressing GCaMP6f+hM4Di (multiple

comparison test, p=0.049).

In vlOFC, there was also no significant effect of CNO (10μM) on Ca^{2+} events for neurons

expressing GCaMP6f or GCaMP6f+mCherry (**Fig. 1F**). However, in CaMKII$^+$ vlOFC neurons

expressing GCaMP6f+hM4Di there was a decrease in the frequency of Ca^{2+} events during CNO

application. A 2-way ANOVA revealed a significant drug x virus interaction [$F_{(2,400)} = 8.349$, $p<$

0.001], with multiple comparisons test resulting in decreased Ca^{2+} events in GCaMP6f+hM4Di

following CNO ($p = 0.02$), and increased activity in GCaMP6f expressing neurons after CNO (p

$= 0.02$).

Discrimination learning: eGFP controls

Mixed-effects GLMs for the discrimination learning phase were conducted for each task

separately to establish if there were baseline differences in learning measures between animals

infused with eGFP virus in different brain regions. There were no differences between the eGFP

groups by target region on learning (i.e., the probability of choosing the correct side) across trials

in the action-based task (β_{region} = -0.13, t(2392) = -0.72, p = 0.47), as well as no differences in

learning (i.e., probability of choosing the correct visual stimulus) across sessions in the stimulus-

based task (β_{region} = 0.10, t(77) = 1.70, p = 0.09). Thus, animals' data were collapsed into a single

eGFP virus group for subsequent analyses.

Discrimination learning: hM4Di vs. eGFP

For the action-based task, there were no significant effects of virus or virus interactions for

vlOFC vs. eGFP on *probability correct* (β_{virus} = -0.13, t(4792) = -1.02, p = 0.31), with similar

findings for the comparison of BLA vs. eGFP (β_{virus} = -0.14, t(4792) =- 1.06, p= 0.29; **Fig. 2C**).

All animals met criterion very quickly (~2 days), thus, we compared trials to reach 75% criterion

(i.e., probability of choosing the correct side). Both hM4Di virus groups performed comparably

[M±SEM: vlOFC hM4Di (81.1±23.0), BLA hM4Di (84.1±21.7)], whereas the eGFP group met

criterion within fewer trials (59.5±18.6), but the difference was not statistically significant [vlOFC

hM4Di vs. eGFP: β_{virus} = 54.5, t(25) = 1.34, p = 0.19; BLA hM4Di vs. eGFP: β_{virus} = 54.8, t(25) =

1.38, p = 0.18].

For the stimulus-based task, there were also no significant effects of virus or virus

interactions for either vlOFC vs. eGFP on *probability correct* (β_{virus} = -0.06, t(141) = -1.26, p =

0.21), or for BLA vs. eGFP (β_{virus} = -0.05, t(152) = -1.32, p = 0.19; **Fig. 4C**). The animals on

average took approximately ~6 days to meet criterion regardless of virus group [M±SEM: vlOFC

hM4Di (6.1±0.7), BLA hM4Di (6.5±1.2), eGFP (6.9±0.7)]. However, many animals did not meet

criterion after a maximum of 10 days of testing (61% of rats).

Given the poorer learning in the stimulus-based task, we evaluated whether this was due to

the order of task administered [i.e., Stimulus → Action or Action → Stimulus]. To test whether

learning was influenced by task order, we analyzed *probability correct* during initial

discrimination learning for the stimulus-based task, which resulted in no effect of task order (β_{order}

= 0.03, t(220) = 0.97, *p* = 0.33), but a significant task order x session interaction ($\beta_{order \, x \, session}$ =-

0.03, t(220) = -3.06, *p* = 0.002). Thus, subsequent analyses were conducted with task order

analyzed separately by session, which revealed that animals administered the Action → Stimulus

task order exhibited poorer learning across sessions ($\beta_{session}$ = 0.01, t(58) = 2.03, *p* = 0.05),

compared to those administered the Stimulus→ Action task order ($\beta_{session}$ = 0.04, t(162) = 6.14, *p*

< 0.0001). Notably, only 1 BLA hM4Di animal successfully met criterion on stimulus-based

learning for the Action → Stimulus order, while no OFC hM4Di animals achieved this.

Accuracy in reversal learning across session and trials

Action-based reversal learning. Mixed-effects GLMs were used to analyze *probability*

correct, with trial number, reversal number, drug order, drug, virus, and sex as between-subject

factors, trial number, reversal number, and drug as within-subject factors, and individual rat as

random factor. GLMs were conducted separately by target region (vlOFC vs. eGFP and BLA vs.

eGFP), using the following formula for the full model: *γ ~ [1 + trial number * reversal number **

virus *drug * drug order * sex + (1 + trial number * reversal number * drug| rat)].

For the comparison of vlOFC with eGFP, several interactions were found: interaction of

sex, virus, drug, and trial number ($\beta_{sex \, x \, virus \, x \, drug \, x \, trial \, number}$ = 0.035, t(19136) = 2.16, *p* = 0.03), an

interaction of sex, virus, drug, drug order, and trial number ($\beta_{sex \, x \, virus \, x \, drug \, order \, x \, trial \, number}$ = -0.006,

t(19136) = -2.32, *p* = 0.02), as well as an interaction of virus, drug, drug order, reversal number,

and trial number ($\beta_{\text{virus x drug x drug order x reversal number x trial number}}$ = -0.001, t(19136) = -2.19, p = 0.03).

Due to these interactions, we were justified to look further at individual reversals. For the comparison of vlOFC hM4Di vs eGFP, sex emerged as a significant predictor of R1 probability correct (β_{sex} = -0.279, t(4784) = -2.03, p = 0.04), and thus sex was entered in the model as a covariate: *γ ~ [1 + trial number * reversal number * virus * drug order + sex + (1 + trial number * drug| rat)]*. With sex as a covariate, there was a significant effect of trial number ($\beta_{\text{trial number}}$ = 0.003, t(4791) = 7.08, p = 1.63e^{-12}) and a nonsignificant but trending interaction of virus, drug order, and trial number for R1: ($\beta_{\text{virus x drug order x trial number}}$ = -0.002, t(4791) = -1.80, p = 0.07; **Fig. 2D**). For probability correct in R3, we also found sex a significant moderator, interacting with drug ($\beta_{\text{sex x drug}}$ = 0.334, t(4784) = 2.18, p = 0.029), and drug x trial number ($\beta_{\text{sex x drug x trial number}}$ = -0.003, t(4784) = -2.55, p = 0.01; **Fig. 2D**). When sex was entered as a covariate in the model, we found a

Figure 2. Either BLA or vlOFC inhibition attenuates action-based reversal learning. (A-B) *Trial structure (A) and timeline (B) of the action-based task. Rats were first surgerized with either hM4Di DREADDs on a CaMKII promoter or eGFP null virus, also on the same promoter. Rats were allowed to recover for 1 week before testing on a stimulus-or action-based reversal learning task. (C) Initial learning of a rewarded side. (D) Learning during subsequent deterministic (100/0) and probabilistic (90/10) reversals. Plots show cumulative P(Correct) for first 100 trials with a sliding window of 10 trials is shown. Drug order was counterbalanced such that on R2 and R4 animals received VEH if they were administered CNO first on R1 and R3, and vice versa. There was no effect of CNO on learning in the eGFP group. *p<0.05 sex as a significant predictor (see* **Fig. 3** *for learning curves plotted by sex). Bonferroni-corrected post-hocs following mixed-effects GLM with sex as a covariate fixed factor wherein a drug x virus interaction was found resulted in **p<0.01 effect of drug only in BLA hM4Di, not in eGFP.*

significant effect of trial number as all animals exhibited improvements in probability correct

across trials ($\beta_{\text{trial number}} = 0.003$, t(4791) = 7.73, p = 1.34e^{-14}), with males performing significantly

better than females (β_{sex} 0.0841, t(4791) = 2.088, p = 0.04; **Fig. 3B**), but no other significant

interactions with any other factor for action-based reversal learning for vlOFC hM4Di compared

to eGFP.

Starting with the full model as above for the comparison of BLA with eGFP, several

interactions of virus x drug were observed, including: virus x drug order x trial number, ($\beta_{\text{virus x drug}}$

$_{\text{order x trial number}}$ = -0.008, t(18986) = -2.58, p = 0.01), virus x reversal number x drug order x trial

number ($\beta_{\text{virus x reversal number x drug order x trial number}}$ = 0.002, t(18986) = 2.59, p = 0.01), and virus x drug

x reversal number x drug order x trial number ($\beta_{\text{virus x drug x reversal number x drug order x trial number}}$ = -0.002,

Figure 3. Female learning was more adversely affected by vlOFC and BLA inhibition than male learning during deterministic and probabilistic action-based reversals. *Cumulative P(Correct) for first 150 trials with a sliding window of 10 trials is shown. Learning of deterministic (100/0) (A) and probabilistic (90/10) (B) reversals as measured by probability correct (P(Correct)). Drug order was counterbalanced such that on R2 and R4 (not shown) animals received VEH if they were administered CNO first on R1 and R3, and vice versa. Chemogenetic inhibition of vlOFC lowered P(Correct) on first deterministic R1 and first probabilistic reversal R3, whereas BLA inhibition attenuated probabilistic learning. Bonferroni-corrected post-hocs following GLM following a sex x drug interaction resulted in effect of drug only in vlOFC or BLA hM4Di, not in eGFP. There were no significant sex differences and no effect of CNO on learning in the eGFP group. **p<0.01, *p<0.05.*

$t(18986) = -2.00$, $p = 0.046$). There were also interactions with sex, including: sex x drug x drug

order ($\beta_{sex\ x\ drug\ x\ drug\ order} = -1.07$, $t(18986) = -2.72$, $p = 0.01$), and sex x drug x reversal number x

drug order ($\beta_{sex\ x\ drug\ x\ reversal\ number\ x\ drug\ order} = 0.335$, $t(18986) = 2.19$, $p = 0.03$). Due to these

interactions, we were justified to look further at individual reversals. For BLA hM4Di compared

to eGFP in R1 there was also a sex difference (females > males) and an effect of trial number on

probability correct ($\beta_{sex} = -0.279$, $t(4784) = -2.38$, $p = 0.02$; $\beta_{trial\ number} = 0.002$, $t(4784) = 3.87$, $p =$

$1.0e^{-04}$). With sex as a covariate, there was only a significant effect of trial number for R1 (β_{trial}

$_{number} = 0.003$, $t(4791) = 7.17$, $p = 8.93e^{-13}$). Sex was also a significant moderator of probability

correct in R3, with significant interactions of sex x trial number ($\beta_{sex\ x\ trial\ number} = 0.002$, $t(4784) =$

2.25, $p = 0.03$) and sex x trial number x drug ($\beta_{sex\ x\ drug\ x\ trial\ number} = -0.003$, $t(4784) = -2.42$, $p =$

0.02). When sex was included as a covariate, there was a significant effect of BLA inhibition on

probability correct in probabilistic R3: (GLM: $\beta_{drug*virus} = -0.31$, $t(4791) = -2.08$, $p = 0.038$).

Bonferroni-corrected post-hoc comparisons revealed an effect of CNO in hM4Di ($p < 0.01$), not

in eGFP ($p = 1.0$), with both males and females exhibit attenuated learning of probabilistic R3

following BLA inhibition.

Given the sex x drug interactions observed in both hM4Di groups, *probability correct* was

also analyzed separately for hM4Di males, eGFP males, hM4Di females, and eGFP females using

the following formula: $\gamma \sim [1 + drug + (1 + drug|\ rat)]$. We found a significant effect of drug for

R1 ($\beta_{drug} = -0.24$, $t(1348) = -2.49$, Bonferroni corrected $p = 0.03$) and R3 ($\beta_{drug} = -0.11$, $t(1348) =$

-3.09, Bonferroni corrected $p = 0.004$) for vlOFC hM4Di females and significant effect of drug

for R3 for BLA hM4Di females ($\beta_{drug} = -0.27$, $t(1348) = -2.97$, Bonferroni corrected $p = 0.006$).

There was no significant effect of drug in eGFP females, eGFP males, or hM4Di males across

regions and reversals (**Fig. 3**).

To gain more insight into this sex difference and its potential underlying mechanism, we

next compared the estimated model parameters from the reinforcement learning models (*RL*,

RL$_{decay}$). Comparing the goodness of fit between the two models, we found that the second model

with the decay parameter (*RL$_{decay}$*) better accounted for the animals' choice behavior as indicated

by significantly lower AIC (paired sample t-test; t(1554) = 6.792, *p* = 1.56e^{-11}). Overall mean BIC

value was also lower for the second model, although the values did not significantly differ from

the first model (t(1554) = 0.93806, *p* = 0.348). Therefore, we focused on the estimated parameters

from the *RL1$_{decay}$* model only.

Comparison of the estimated parameters across groups revealed that female and male rats

differed mainly in the decay parameter γ_d, which governs the amount of passive decay or

forgetting in the value estimate of the unchosen option (**Fig. 3-1**). During the first deterministic

reversal (R1, 100/0), eGFP females showed overall significantly lower values of γ_d than eGFP

males (mean difference in γ_d = -0.098; Wilcoxon rank sum test, p=0.00537), suggesting a different

mechanism of adjustment to the reversal between females and males. While there was no clear

evidence for such sex difference in γ_d for the vlOFC or BLA hM4Di groups during the first

deterministic reversal (R1), the first probabilistic reversal (R3, 90/10) instead revealed a sex-

specific effect between CNO and VEH groups: vlOFC hM4Di females who were administered

CNO to inhibit vlOFC had significantly higher values of γ_d compared to those who received VEH

(mean difference in γ_d = 0.150; *p* = 0.00757). In contrast, vlOFC hM4Di males did not show a

significant difference in γ_d between CNO and VEH groups (mean difference in γ_d = -0.0753; *p* =

0.228). BLA hM4Di groups showed a similar trend in γ_d, with the females exhibiting a larger

difference between CNO and VEH groups (mean difference in γ_d = 0.137; *p* = 0.0673) than males

(mean difference in γ_d = 0.052; *p* = 0.961). These results based on RL model fitting suggest that

the attenuated probabilistic learning for the female CNO groups is mediated by larger γ_d or

decreased memory for the unchosen options after vlOFC and BLA inhibition. Importantly, this

significant difference emerged due to hM4Di VEH females exhibiting *enhanced* memory in R3.

This is because in this condition, rats had previously received CNO in R2 and thus did not encode

the reversal very well, making it easier to return to the R1 contingency (which was the same as in

R3). Collectively, this shows that vlOFC, and perhaps secondarily BLA, is necessary for the

encoding and retrieval of action-based values.

Stimulus-based reversal learning. In contrast to the acquisition curves that demonstrated

learning of the initial visual discrimination (**Fig. 4C**), all animals exhibited difficulty with

stimulus-based reversal learning, rarely achieving above 60% after 10 sessions (**Fig. 4-1**), similar

to recent reports (Harris, Aguirre et al., 2021; Ye et al., 2023). Here, due to several non-learners,

we adhered to the criterion of rats reaching greater than a 50% running window average for the

last 100 trials in discrimination, for inclusion in subsequent reversal learning analyses. The

following numbers did not meet this criterion and were excluded from these groups: 0 of 13 vlOFC

hM4Di, 6 of 15 BLA hM4Di, and 6 of 17 eGFP. As above, GLMs were conducted separately for

accuracy (*probability correct*) by target region comparison, but with session instead of trial

number as a within-subject factor. Thus, the GLM formula was as follows: *γ ~ [1 + session*

*reversal number * virus * drug * drug order * sex + (1 + session * reversal number * drug| rat)].

For vlOFC hM4Di comparison with eGFP, we observed several interactions including:

session, drug, and virus ($\beta_{\text{session x drug x virus}}$ = -0.096, t(536) = -2.83, p = 4.81e^{-03}), session, drug, drug

order, and virus ($\beta_{\text{session x drug x drug order x virus}}$ = 0.146, t(536) = 2.62, p = 9.17e^{-03}) and drug, drug

order, and reversal number ($\beta_{\text{drug x drug order x reversal number}}$ = 0.107, t(536) = 2.18, p = 0.03). We also

observed a sex x reversal number interaction ($\beta_{\text{sex x reversal number}}$ = -0.084, t(536) = -2.50, p = 0.01).

Aside from a significant effect of session in R1 ($\beta_{session}$ = 0.03, t(134) = 2.67, p = 8.55e^{-03}) there

were no significant predictors of learning with follow-up GLM analyses with sex in the model.

With sex as a covariate in the model there was a significant effect of session for R1 and R3 (R1:

$\beta_{session}$ = 0.026, t(141) = 3.80, p = 2.16e^{-04}; R3: $\beta_{session}$ = 0.018, t(141) = 2.92, p = 4.11e^{-03}), and a

session x virus interaction in R3 ($\beta_{session \, x \, virus}$ = -0.020, t(141) = -2.07, p = 0.04). Bonferroni-

corrected post-hoc comparisons revealed an effect of session in eGFP (p < 0.01), but not in hM4Di

(p = 0.45), indicating that only the eGFP group improved across session in R3 (**Fig. 4-1**).

For BLA hM4Di compared to eGFP, several interactions were observed including

interactions of session, drug, and virus ($\beta_{session \, x \, drug \, x \, virus}$ = -0.087, t(575) = -2.25, p = 0.02), drug,

drug order, and reversal number ($\beta_{drug \, x \, drug \, order \, x \, reversal \, number}$ = 0.107, t(575) = 2.29, p = 0.02), and

sex and reversal number ($\beta_{sex \, x \, reversal \, number}$ = -0.084, t(575) = -2.15, p = 0.03). However, none of

the post-hoc GLM analyses resulted in significant predictors of learning with the exception of

session for R1 and R3 (R1: $\beta_{session}$ = 0.026, t(151) = 2.35, p = 0.02; R3: $\beta_{session}$ = 0.018, t(151) =

2.67, p = 8.43e^{-03}) when sex was included as a covariate in the model.

Due to the slow stimulus-based reversal learning, we next assessed *probability correct*

around reversals (three sessions before and after reversals for stimulus-based learning, and one

session before and after reversals for action-based learning) to test for detection and adjustments

to reversals.

Accuracy around reversals: Reversal detection

Stimulus-based reversal learning. We analyzed accuracy (*probability correct*) around

reversals, as overall stimulus-based reversal learning was modest. ANOVAs with virus and drug

order as between subject factors were conducted on the mean change in accuracy between one

reversal and the next. vlOFC hM4Di was significantly different from eGFP for R1-to-R2 [$F_{(1,24)}$=

9.49, $p < 0.01$] and R2-to-R3 [$F_{(1,24)}$= 10.1, $p < 0.01$], but not R3-to-R4 [$F_{(1,24)}$= 2.61, $p = 0.12$],

Fig. 4D. In contrast, BLA hM4Di was not significantly different from eGFP on changes in

accuracy around any of the reversals.

Figure 4. Inhibition of vlOFC, but not BLA, impairs the detection of stimulus-based reversals as measured by probability correct adjustments. (A-B) *Trial structure (A) and timeline (B) of the stimulus-based task. Rats were first surgerized with either hM4Di DREADDs or eGFP null virus on a CaMKII promoter. Rats were allowed to recover for 1 week before testing on a stimulus-or action-based reversal learning task. (**C**) Initial learning of a rewarded stimulus, presented pseudorandomly on the left or right side of the touchscreen for eGFP (top), vlOFC hM4Di (middle), and BLA hM4Di (bottom). (**D**) Rats were always tested on a deterministic schedule before a probabilistic one. Shown are subsequent deterministic (100/0) and probabilistic (90/10) reversal "transitions," 3 sessions before and after each reversal. Drug order was counterbalanced such that on R2 and R4 animals received VEH if they were administered CNO first on R1 and R3, and vice versa. Chemogenetic inhibition of vlOFC abolishes changes in probability correct over the last 3 (pre) and first 3 (post)-reversal sessions, indicating impaired detection of reversal. In contrast, BLA inhibition had no impact on this detection. There was also no effect of CNO in eGFP group learning. **p<0.01 different than eGFP following ANOVA of pre-post difference.*

Action-based reversal learning. We also assessed accuracy (*probability correct*) around

reversals for action-based reversal learning. As above, ANOVAs with virus and drug order as

between subject factors were conducted on the mean accuracy change between one reversal and

the next. Other than confirming the probabilistic reversal learning (R3) impairment for BLA

hM4Di (**Fig. 2**), there were no significant effects of virus groups on accuracy changes on any other

reversal transition in action-based reversal learning (**Fig. 2-1).**

In summary, our results indicate that both BLA and vlOFC are required for learning action-

based reversals in females. Conversely, vlOFC, but not BLA, is necessary for detecting stimulus-

based, not action-based, reversals.

Win-Stay, Lose-Switch Strategies around reversals

Stimulus-based reversal learning. We also analyzed adaptive response strategies (*Win-Stay*

and *Lose-Switch*) around reversals. ANOVAs with virus and drug order as between subject factors

were conducted on mean Win-Stay or Lose-Switch between one reversal and the next. vlOFC

hM4Di was significantly different from eGFP for *Win-Stay* R1-to-R2 [$F_{(1,24)} = 9.91$, $p < 0.01$] and

R2-to-R3 [$F_{(1,24)} = 11.61$, $p < 0.01$], but not R3-to-R4 [$F_{(1,24)} = 1.49$, $p = 0.24$]. Similarly, vlOFC

Figure 5. Inhibition of vlOFC, but not BLA, impairs the detection of deterministic stimulus-based reversals as measured by changes in Win-Stay, Lose-Switch strategies. (A) *Inhibition of vlOFC abolishes changes in Win-Stay over the last 3 (pre) and first 3 (post)-reversal sessions, indicating impaired detection of R1-R2, R2-R3. In contrast, BLA inhibition had no impact on this detection.* **(B)** *Inhibition of vlOFC abolishes changes in Lose-Shift over the last 3 (pre) and first 3 (post)-reversal sessions, indicating impaired detection of R1-R2, R2-R3. In contrast, BLA inhibition had no impact on this detection. There was also no effect of CNO in eGFP group learning. **p<0.01 different than eGFP following ANOVA of pre-post difference.*

hM4Di differed from eGFP for *Lose-Switch* in R1-to-R2 [$F_{(1,24)}$=6.00, *p*=0.023] and R2-to-R3

[$F_{(1,24)}$=5.00, *p*=0.036], but not R3-to-R4 [$F_{(1,24)}$=0.08, *p*=0.78], **Fig. 5**.

In contrast, BLA hM4Di was not significantly different from eGFP on changes in *Win-Stay*

or *Lose-Switch* strategies around any of the reversals. Therefore, the results for these adaptive

strategies reflect an identical pattern to that observed for *probability correct* for both vlOFC and

BLA hM4Di, above.

Performance measures in reversal learning

We analyzed other performance measures including *initiation, choice,* and *reward*

latencies, as proxies for attention, deliberation, and motivation, respectively (Harris, Aguirre et al.,

2021; Ye et al., 2023). Given that the significant effects in accuracy were observed only in R1 and

R3, we elected to analyze latency measures during those reversals only, with sex added as a

covariate in the GLM models. In this case, a single measure per animal per reversal number was

obtained: the GLM formula was as follows for R1 and R3: *γ ~ [1 + virus * drug + sex + (1 +*

drug| rat)].

Action-based reversal learning. vlOFC comparisons with eGFP controls did not yield any

significant effects for R1; however, a significant drug effect emerged in R3, such that VEH-treated

animals exhibited longer *initiation latencies* than when treated with CNO, irrespective of virus

group (β_{virus} = -0.524, t(27) = -2.06, *p* = 0.049). Females tended to exhibit longer latencies than

males, although this was only trending toward significance (β_{sex} = -0.353, t(27) = -1.96, *p* = 0.06).

When comparing BLA hM4Di vs. eGFP, we found a significant effect of virus in the full

model (β_{virus} = 1.44, t(93) = 2.07, *p* = 0.039), with BLA hM4Di animals exhibiting longer initiation

latencies than eGFP controls. We also observed a significant effect of virus for both *incorrect*

choice (R1: β_{virus} = 0.38, t(27) = 4.68 p < 0.0001; R3: β_{virus} = 0.28, t(28) = 3.40, p = 0.002) and

reward latencies (R1: β_{virus} = 0.34, t(30) = 3.78 p = 0.0007; R3: β_{virus} = 0.22, t(30) = 2.31, p = 0.03),

such that BLA hM4Di exhibited longer latencies compared to eGFP (**Fig. 2-2B**). This effect was

not observed when comparing vlOFC hM4DI vs. eGFP (**Fig. 2-2A**).

652   *Stimulus-based reversal learning.* We similarly analyzed performance measures in

stimulus-based reversal learning during R1 and R3, with sex added as a covariate to the model.

We found a robust sex difference, with females committing more *initiation omissions* than males,

irrespective of virus or drug, across reversals (vlOFC vs. eGFP: β_{sex} = -88.52, t(80) = -2.40 p =

0.02; BLA vs. eGFP: β_{sex} = -80.38, t(89) = -2.14, p = 0.04; **Fig. 4-2AB**). For *initiation latencies,*

vlOFC hM4Di vs. eGFP analysis yielded an overall significant effect of sex as well (β_{sex} = -3.23,

t(81) = -3.72, p = 0.0004), such that females exhibited longer initiation latencies than males across

reversals (**Fig. 4-2CD**). BLA hM4Di vs. eGFP comparisons also yielded a significant effect of sex

(β_{sex} = -3.26, t(88) = -3.88, p = 0.0002), however, there was also a significant interaction between

reversal number, virus, drug, drug order, and sex ($\beta_{reversal\ number\ x\ virus\ x\ drug\ x\ drug\ order\ x\ sex}$ = 1.46, t(88)

= 2.13, p = 0.036), which justified analysis of each reversal number separately. Subsequent

analyses revealed a significant effect of sex for both R1(β_{sex} = -1.58, t(24) = -3.40, p = 0.002), and

R3 (β_{sex} = -0.67, t(24) = -2.42, p = 0.024), with females exhibiting longer initiation latencies than

males, irrespective of virus or drug (**Fig. 4-2D**).

666   In summary, across all reversals BLA hM4Di animals exhibited slower deliberation speed

667 during incorrect choices, and took longer to collect reward compared to eGFP controls in the

668 action-based task. In contrast, sex emerged as a strong predictor of performance measures, but not

669 learning, in the stimulus-based task, as it accounted for much of the variance in initiation omissions

670 and latencies.

Discussion

We used a chemogenetic approach to transiently inactivate neurons in either vlOFC or BLA to assess how these regions are involved in different aspects of reversal learning. Although the role of OFC in reversal learning has been instantiated in different paradigms using visual stimuli and cues (Izquierdo et al., 2013; Piantadosi et al., 2018; Hervig et al., 2020; Alsio et al., 2021) as well as olfactory ones (Schoenbaum et al., 2003; Kim and Ragozzino, 2005), several groups also report a strong role for OFC in action (spatial)-based reversal learning (Dalton et al., 2016; Groman et al., 2019; Verharen et al., 2020). Almost all of these reversal learning investigations have involved irreversible lesions or baclofen/muscimol inactivations of OFC. Testing both types with a chemogenetic approach, here we found that different aspects of both action- and stimulus-based reversal learning rely on vlOFC.

In parallel, the specific role of BLA in stimulus- vs. action-based reversal learning is poorly understood given mixed results (Schoenbaum et al., 2003; Izquierdo and Murray, 2004; Churchwell et al., 2009; Hervig et al., 2020). Recent studies suggest amygdala may be involved in both types of learning (Taswell et al., 2021; Keefer and Petrovich, 2022) as BLA activity is modulated by violations in reward expectations generally, which are not association-specific (Roesch et al., 2012). To probe this, we tested animals on both stimulus- and action-based tasks and found that BLA is more selectively involved in action-based reversal learning.

As additional motivations for the present study, several reports suggest that neural recruitment in reversal learning may depend on certainty of rewards (Boulougouris et al., 2007; Boulougouris and Robbins, 2009; Ward et al., 2015; Costa et al., 2016; Dalton et al., 2016; Piantadosi et al., 2018; Verharen et al., 2020). To further understand this, we tested animals on both deterministic (100/0) and probabilistic reversals (90/10). We found vlOFC to be involved in

the detection of stimulus-based reversals and initial learning of both deterministic and probabilistic

learning across stimuli and actions, whereas BLA was more selectively involved in probabilistic

reversal learning of actions.

Finally, due to the sparsity of research probing sex differences in flexible learning and

decision making (Orsini and Setlow, 2017; Grissom and Reyes, 2019; Orsini et al., 2022; Cox et

al., 2023), where an overwhelming number of reversal learning studies include only males

(Schoenbaum et al., 2003; Izquierdo et al., 2013; Dalton et al., 2016; Groman et al., 2019; Hervig

et al., 2020; Verharen et al., 2020), we included both male and female rats here. We found sex-

dependent contributions of both vlOFC and BLA in action-based reversal learning. We elaborate

on these findings within the context of the existing literature below.

Similar recruitment of BLA and vlOFC during action-based reversal learning

All animals learned to flexibly adjust their responses following deterministic and

probabilistic reversals, indicating successful remapping of reward contingencies as accuracy

increased across trials. Importantly, we found no effect of CNO in eGFP animals, suggesting that

it was activation of hM4Di receptors in BLA and vlOFC that were crucial to any impairments

observed. After considering sex as a covariate in our analyses, we determined that vlOFC was

necessary for learning of first deterministic (R1) and probabilistic reversal (R3). This is consistent

with findings following pharmacological inactivations or lesions of OFC (Boulougouris et al.,

2007; Boulougouris and Robbins, 2009; Dalton et al., 2016; Piantadosi et al., 2018; Verharen et

al., 2020).

BLA inhibition was not expected to impair deterministic reversal learning as it is thought

to be mostly recruited when there is some level of uncertainty, e.g., probabilistic outcomes (Roesch

et al., 2012). Amygdala-lesioned monkeys are also impaired on action(spatial)-based probabilistic

reversal learning, exhibiting decreased probability of choosing the better option, and increased

switching behavior following negative outcomes (Taswell et al., 2021). BLA may indeed be

critical in generating prediction error signals following changes in reward associations (Esber et

al., 2012; Roesch et al., 2012; Iordanova et al., 2021), with particular involvement in detecting

unexpected upshifts or downshifts in value (Roesch et al., 2010; Stolyarova and Izquierdo, 2017).

Our finding of attenuated learning of probabilistic reversal R3 suggests it is the misleading

feedback that most engages BLA. Future investigations should probe the role of BLA in initial

learning of probabilistic outcomes, without reversal experience.

vlOFC, but not BLA, is necessary for detection of reversals in stimulus-based learning

As described above, unlike the ease of action-based reversal learning, rats exhibited

difficulty learning reversals of stimulus-reward contingencies, as previously reported (Harris,

Aguirre et al., 2021). Thus, instead of examining acquisition curves which reached asymptote

slightly above chance, we elected to study detection and adjustment to reversals by comparing

accuracy and strategy prior to and after a reversal occurred. Furthermore, this enabled assessment

about whether prior inhibition affected future detection of reversals and whether this varied by

transition type [i.e., between deterministic reversals (R1→R2), deterministic and probabilistic

reversal (R2→R3), or probabilistic reversals (R3→R4)]. We found that vlOFC, but not BLA,

inhibition produced a failure in detecting first deterministic and first probabilistic reversal. This

pattern was not observed in animals that received VEH during the first deterministic reversal,

suggesting vlOFC needs to be "online" when experiencing a reversal for the very first time as this

determines how flexibly animals respond to future reversals. Employment of adaptive strategies

matched this effect, such that vlOFC-inhibited animals did not employ either win-stay or lose-

switch strategies after the first reversal. That vlOFC inhibition did not impair the ability to detect

transitions between probabilistic reversals (R3→R4) supports the idea that other brain regions may

be recruited when the probabilistic reward contingencies have already been established. The role

of OFC in establishing an "expected uncertainty" (Soltani and Izquierdo, 2019) has been

instantiated experimentally in several recent studies using different methodologies (Namboodiri et

al., 2019; Namboodiri et al., 2021; Jenni et al., 2022), and we add detection and adjustment to

stimulus-based reversals to this evidence. In contrast, BLA inhibition did not result in any

impairment in the ability to detect and flexibly adjust to reversals, regardless of whether they were

deterministic or probabilistic.

Females exhibited poorer action-based reversal learning following vlOFC and BLA inhibition

We found a significant sex-dependent effect of vlOFC inhibition for both the first

deterministic (R1) and the first probabilistic reversal (R3), and a similar effect in females following

BLA inhibition for R3. Importantly, although BLA hM4Di males also exhibited the general pattern

of attenuated learning of R3, the effect of BLA inhibition was largely driven by females. These

findings were unexpected as we did not anticipate sex differences in the recruitment of brain

regions involved in action-based reversal learning, mostly due to the lack of studies to date

investigating the effect of sex. Nonetheless, there is evidence that OFC is differentially activated

in males and females during risky decisions such that activity in the lateral OFC is inversely

correlated with the proportion of advantageous choices in female, but not in male, rats (van Hasselt

et al., 2012), with similar effects found in humans (Bolla et al., 2004) and in amygdala (Dreher et

al., 2007).

Unfortunately, the potential of estrous-driven behavioral variation in females has

commonly been used as a rationale for excluding female rodents in behavioral neuroscience

research (Beery and Zucker, 2011; McCarthy et al., 2012; Shansky and Murphy, 2021). However,

our findings, and other recent studies of cortical circuits exhibiting sex-mediated influences on

reward-motivated behavior (Cox et al., 2023) should instead encourage the inclusion of both sexes

in experiments as clear patterns emerge. Additionally, the learning impairment observed in females

following vlOFC and BLA inhibition may not be primarily due to fluctuations in hormone levels,

but rather may reflect differential adoption of strategies between sexes (Grissom and Reyes, 2019;

Chen et al., 2021a; Chen et al., 2021b). Consistent with this view, our results based on estimated

RL model parameters suggests differential mechanisms for adjustment to reversals between males

and females. Specifically, we found some evidence for differential effects on γ_d, where the decay

rate for unchosen action values was greater for females than males. This is consistent with a

previous study which also observed similar disruption in retention of action values after ablating

OFC neurons projecting to BLA (Groman et al., 2019). Given that study involved only male rats

and involved a stronger manipulation than our chemogenetic approach (i.e., one that caused

pathway-specific neuronal apoptosis), it is plausible their observed effect on γ_d would have been

different between females and males.

Stimulus-based vs. action-based learning

Interestingly, we discovered task order to be significant in rats' ability to learn to

discriminate stimuli: stimulus→action was learned much more readily than action→stimulus. This

can be explained by noting that rats are heavily biased to acquire spatial associations (Wright et

al., 2019), and reinforcing this already-strong learning likely inhibits the ability to learn

associations where spatial information should be ignored. In contrast, nonhuman primates are able

to quickly transition between "what" vs. "where" blocks of trials (Rothenhoefer et al., 2017;

Taswell et al., 2021). Nonetheless, learning both types of associations is crucial for flexibility

required in naturalistic environments and thus, it is important to examine how stimulus-based and

action-based learning systems interact with each other (Soltani & Koechlin, 2022). Moreover,

although the role of OFC in stimulus- or cue-based reversal learning has been probed using

olfactory and visual stimuli, more viral-mediated approaches employing targeted chemogenetic

and optogenetic manipulations across sensory modalities in both males and females are warranted.

Conclusion

The present results suggest similar roles for vlOFC and BLA in flexible action-based

learning, but a more specialized role for vlOFC in setting adjustments in stimulus-based learning.

Additionally, our findings underscore the importance of including both male and female animals

in behavioral neuroscience studies, adding to the mounting evidence of sex-modulated learning

flexibility (Chen et al., 2021a; Chen et al., 2021b).

References

References

Alsio J, Lehmann O, McKenzie C, Theobald DE, Searle L, Xia J, Dalley JW, Robbins TW
(2021) Serotonergic Innervations of the Orbitofrontal and Medial-prefrontal Cortices are
Differentially Involved in Visual Discrimination and Reversal Learning in Rats. Cereb
Cortex 31:1090-1105.
Asrican B, Song J (2021) Extracting meaningful circuit-based calcium dynamics in astrocytes
and neurons from adult mouse brain slices using single-photon GCaMP imaging. STAR
Protoc 2:100306.
Barreiros IV, Panayi MC, Walton ME (2021a) Organization of Afferents along the Anterior-
posterior and Medial-lateral Axes of the Rat Orbitofrontal Cortex. Neuroscience 460:53-
68.
Barreiros IV, Ishii H, Walton ME, Panayi MC (2021b) Defining an orbitofrontal compass:
Functional and anatomical heterogeneity across anterior-posterior and medial-lateral
axes. Behav Neurosci 135:165-173.
Beery AK, Zucker I (2011) Sex bias in neuroscience and biomedical research. Neurosci
Biobehav Rev 35:565-572.
Behrens TE, Woolrich MW, Walton ME, Rushworth MF (2007) Learning the value of
information in an uncertain world. Nat Neurosci 10:1214-1221.
Bolla KI, Eldreth DA, Matochik JA, Cadet JL (2004) Sex-related Differences in a Gambling
Task and Its Neurological Correlates. Cerebral Cortex 14:1226-1232.
Boulougouris V, Robbins TW (2009) Pre-surgical training ameliorates orbitofrontal-mediated
impairments in spatial reversal learning. Behavioural Brain Research 197:469-475.
Boulougouris V, Dalley JW, Robbins TW (2007) Effects of orbitofrontal, infralimbic and
prelimbic cortical lesions on serial spatial reversal learning in the rat. Behavioural Brain
Research 179:219-228.
Chen CS, Knep E, Han A, Ebitz RB, Grissom NM (2021a) Sex differences in learning from
exploration. Elife 10.
Chen CS, Ebitz RB, Bindas SR, Redish AD, Hayden BY, Grissom NM (2021b) Divergent
Strategies for Learning in Males and Females. Curr Biol 31:39-50 e34.
Churchwell JC, Morris AM, Heurtelou NM, Kesner RP (2009) Interactions between the
prefrontal cortex and amygdala during delay discounting and reversal. Behav Neurosci
123:1185-1196.
Corbit LH, Balleine BW (2005) Double dissociation of basolateral and central amygdala lesions
on the general and outcome-specific forms of pavlovian-instrumental transfer. J Neurosci
25:962-970.
Costa KM, Scholz R, Lloyd K, Moreno-Castilla P, Gardner MPH, Dayan P, Schoenbaum G
(2023) The role of the lateral orbitofrontal cortex in creating cognitive maps. Nat
Neurosci 26:107-115.
Costa VD, Dal Monte O, Lucas DR, Murray EA, Averbeck BB (2016) Amygdala and Ventral
Striatum Make Distinct Contributions to Reinforcement Learning. Neuron 92:505-517.
Cox J, Minerva AR, Fleming WT, Zimmerman CA, Hayes C, Zorowitz S, Bandi A, Ornelas S,
McMannon B, Parker NF, Witten IB (2023) A neural substrate of sex-dependent
modulation of motivation. Nature Neuroscience 26:274-284.
Dalton GL, Wang NY, Phillips AG, Floresco SB (2016) Multifaceted Contributions by Different
Regions of the Orbitofrontal and Medial Prefrontal Cortex to Probabilistic Reversal
Learning. J Neurosci 36:1996-2006.

Dreher J-C, Schmidt PJ, Kohn P, Furman D, Rubinow D, Berman KF (2007) Menstrual cycle
850   phase modulates reward-related neural function in women. Proc Natl Acad Sci U S A
851   104:2465-2470.
852   Esber GR, Roesch MR, Bali S, Trageser J, Bissonette GB, Puche AC, Holland PC, Schoenbaum
853   G (2012) Attention-related Pearce-Kaye-Hall signals in basolateral amygdala require the
854   midbrain dopaminergic system. Biol Psychiatry 72:1012-1019.
855   Ghasemi A, Jeddi S, Kashfi K (2021) The laboratory rat: Age and body weight matter. EXCLI J
856   20:1431-1445.
857   Grissom NM, Reyes TM (2019) Let's call the whole thing off: evaluating gender and sex
858   differences in executive function. Neuropsychopharmacology 44:86-96.
859   Groman SM, Keistler C, Keip AJ, Hammarlund E, DiLeone RJ, Pittenger C, Lee D, Taylor JR
860   (2019) Orbitofrontal Circuits Control Multiple Reinforcement-Learning Processes.
861   Neuron 103:734-746 e733.
862   Harris C, Aguirre C, Kolli S, Das K, Izquierdo A, Soltani A (2021) Unique features of stimulus-
863   based probabilistic reversal learning. Behav Neurosci 135:550-570.
864   Hart EE, Blair GJ, O'Dell TJ, Blair HT, Izquierdo A (2020) Chemogenetic Modulation and
865   Single-Photon Calcium Imaging in Anterior Cingulate Cortex Reveal a Mechanism for
866   Effort-Based Decisions. J Neurosci 40:5628-5643.
867   Hervig ME, Fiddian L, Piilgaard L, Bozic T, Blanco-Pozo M, Knudsen C, Olesen SF, Alsio J,
868   Robbins TW (2020) Dissociable and Paradoxical Roles of Rat Medial and Lateral
869   Orbitofrontal Cortex in Visual Serial Reversal Learning. Cereb Cortex 30:1016-1029.
870   Iordanova MD, Yau JO-Y, McDannald MA, Corbit LH (2021) Neural substrates of appetitive
871   and aversive prediction error. Neurosci Biobehav Rev 123:337-351.
872   Izquierdo A (2017) Functional Heterogeneity within Rat Orbitofrontal Cortex in Reward
873   Learning and Decision Making. J Neurosci 37:10529-10540.
874   Izquierdo A, Murray EA (2004) Combined Unilateral Lesions of the Amygdala and Orbital
875   Prefrontal Cortex Impair Affective Processing in Rhesus Monkeys. Journal of
876   Neurophysiology 91:2023-2039.
877   Izquierdo A, Darling C, Manos N, Pozos H, Kim C, Ostrander S, Cazares V, Stepp H, Rudebeck
878   PH (2013) Basolateral amygdala lesions facilitate reward choices after negative feedback
879   in rats. J Neurosci 33:4105-4109.
880   Janak PH, Tye KM (2015) From circuits to behaviour in the amygdala. Nature 517:284-292.
881   Jang AI, Costa VD, Rudebeck PH, Chudasama Y, Murray EA, Averbeck BB (2015) The Role of
882   Frontal Cortical and Medial-Temporal Lobe Brain Areas in Learning a Bayesian Prior
883   Belief on Reversals. J Neurosci 35:11751-11760.
884   Jenni NL, Rutledge G, Floresco SB (2022) Distinct Medial Orbitofrontal-Striatal Circuits
885   Support Dissociable Component Processes of Risk/Reward Decision-Making. J Neurosci
886   42:2743-2755.
887   Keefer SE, Petrovich GD (2022) Necessity and recruitment of cue-specific neuronal ensembles
888   within the basolateral amygdala during appetitive reversal learning. Neurobiol Learn
889   Mem 194:107663.
890   Kim J, Ragozzino ME (2005) The involvement of the orbitofrontal cortex in learning under
891   changing task contingencies. Neurobiol Learn Mem 83:125-133.
892   Lichtenberg NT, Pennington ZT, Holley SM, Greenfield VY, Cepeda C, Levine MS, Wassum
893   KM (2017) Basolateral Amygdala to Orbitofrontal Cortex Projections Enable Cue-
894   Triggered Reward Expectations. J Neurosci 37:8374-8384.

Malvaez M, Shieh C, Murphy MD, Greenfield VY, Wassum KM (2019) Distinct cortical-
amygdala projections drive reward value encoding and retrieval. Nat Neurosci 22:762-
769.
McCarthy MM, Arnold AP, Ball GF, Blaustein JD, De Vries GJ (2012) Sex differences in the
brain: the not so inconvenient truth. J Neurosci 32:2241-2247.
Namboodiri VMK, Hobbs T, Trujillo-Pisanty I, Simon RC, Gray MM, Stuber GD (2021)
Relative salience signaling within a thalamo-orbitofrontal circuit governs learning rate.
Curr Biol 31:5176-5191.e5175.
Namboodiri VMK, Otis JM, van Heeswijk K, Voets ES, Alghorazi RA, Rodriguez-Romaguera J,
Mihalas S, Stuber GD (2019) Single-cell activity tracking reveals that orbitofrontal
neurons acquire and maintain a long-term memory to guide behavioral adaptation. Nat
Neurosci 22:1110-1121.
Orsini CA, Setlow B (2017) Sex differences in animal models of decision making. J Neurosci
Res 95:260-269.
Orsini CA, Truckenbrod LM, Wheeler A-R (2022) Regulation of sex differences in risk-based
decision making by gonadal hormones: Insights from rodent models. Behav Processes
200:104663-104663.
Pachitariu M, Stringer C, Dipoppa M, Schröder S, Rossi LF, Dalgleish H, Carandini M, Harris
KD (2017) Suite2p: beyond 10,000 neurons with standard two-photon microscopy.
bioRxiv:061507.
Piantadosi PT, Lieberman AG, Pickens CL, Bergstrom HC, Holmes A (2018) A novel
multichoice touchscreen paradigm for assessing cognitive flexibility in mice. Learn Mem
26:24-30.
Roesch MR, Calu DJ, Esber GR, Schoenbaum G (2010) Neural correlates of variations in event
processing during learning in basolateral amygdala. J Neurosci 30:2464-2471.
Roesch MR, Esber GR, Bryden DW, Cerri DH, Haney ZR, Schoenbaum G (2012) Normal aging
alters learning and attention-related teaching signals in basolateral amygdala. J Neurosci
32:13137-13144.
Rothenhoefer KM, Costa VD, Bartolo R, Vicario-Feliciano R, Murray EA, Averbeck BB (2017)
Effects of ventral striatum lesions on stimulus-based versus action-based reinforcement
learning. J Neurosci 37: 6902-6914.
Rudebeck PH, Murray EA (2008) Amygdala and orbitofrontal cortex lesions differentially
influence choices during object reversal learning. J Neurosci 28:8338-8343.
Schoenbaum G, Setlow B, Nugent SL, Saddoris MP, Gallagher M (2003) Lesions of
orbitofrontal cortex and basolateral amygdala complex disrupt acquisition of odor-guided
discriminations and reversals. Learn Mem 10:129-140.
Shansky RM, Murphy AZ (2021) Considering sex as a biological variable will require a global
shift in science culture. Nature Neuroscience 24:457-464.
Sias AC, Morse AK, Wang S, Greenfield VY, Goodpaster CM, Wrenn TM, Wikenheiser AM,
Holley SM, Cepeda C, Levine MS, Wassum KM (2021) A bidirectional corticoamygdala
circuit for the encoding and retrieval of detailed reward memories. Elife 10.
Soltani A, Izquierdo A (2019) Adaptive learning under expected and unexpected uncertainty. Nat
Rev Neurosci 20:635-644.
Soltani A, Koechlin E (2022) Computational models of adaptive behavior and prefrontal cortex.
Neuropsychopharmacology 47: 58-71.

Stalnaker TA, Roesch MR, Calu DJ, Burke KA, Singh T, Schoenbaum G (2007) Neural
correlates of inflexible behavior in the orbitofrontal-amygdalar circuit after cocaine
exposure. Ann N Y Acad Sci 1121:598-609.
Stolyarova A, Izquierdo A (2017) Complementary contributions of basolateral amygdala and
orbitofrontal cortex to value learning under uncertainty. Elife 6.
Stolyarova A, Rakhshan M, Hart EE, O'Dell TJ, Peters MAK, Lau H, Soltani A, Izquierdo A
(2019) Contributions of anterior cingulate cortex and basolateral amygdala to decision
confidence and learning under uncertainty. Nat Commun 10:4704.
Tada M, Takeuchi A, Hashizume M, Kitamura K, Kano M (2014) A highly sensitive fluorescent
indicator dye for calcium imaging of neural activity in vitro and in vivo. Eur J Neurosci
39:1720-1728.
Taswell CA, Costa VD, Basile BM, Pujara MS, Jones B, Manem N, Murray EA, Averbeck BB
(2021) Effects of Amygdala Lesions on Object-Based Versus Action-Based Learning in
Macaques. Cereb Cortex 31:529-546.
Tye KM, Janak PH (2007) Amygdala neurons differentially encode motivation and
reinforcement. J Neurosci 27:3937-3945.
van Hasselt FN, de Visser L, Tieskens JM, Cornelisse S, Baars AM, Lavrijsen M, Krugers HJ,
van den Bos R, Joëls M (2012) Individual variations in maternal care early in life
correlate with later life decision-making and c-fos expression in prefrontal subregions of
rats. PLoS One 7:e37820-e37820.
Verharen JPH, den Ouden HEM, Adan RAH, Vanderschuren L (2020) Modulation of value-
based decision making behavior by subregions of the rat prefrontal cortex.
Psychopharmacology (Berl) 237:1267-1280.
Ward RD, Winiger V, Kandel ER, Balsam PD, Simpson EH (2015) Orbitofrontal cortex
mediates the differential impact of signaled-reward probability on discrimination
accuracy. Front Neurosci 9:230.
Wassum KM, Izquierdo A (2015) The basolateral amygdala in reward learning and addiction.
Neurosci Biobehav Rev 57:271-283.
Winstanley CA, Floresco SB (2016) Deciphering Decision Making: Variation in Animal Models
of Effort- and Uncertainty-Based Choice Reveals Distinct Neural Circuitries Underlying
Core Cognitive Processes. J Neurosci 36:12069-12079.
Wright SL, Martin GM, Thorpe CM, Haley K, Skinner DM (2019) Distance and direction, but
not light cues, support response reversal learning. Learning & Behavior 47:38-46.
Ye T, Romero-Sosa JL, Rickard A, Aguirre CG, Wikenheiser AM, Blair HT, Izquierdo A (2023)
Theta oscillations in anterior cingulate cortex and orbitofrontal cortex differentially
modulate accuracy and speed in flexible reward learning. Oxford Open Neuroscience
kvad005.
Zimmermann KS, Li CC, Rainnie DG, Ressler KJ, Gourley SL (2018) Memory Retention
Involves the Ventrolateral Orbitofrontal Cortex: Comparison with the Basolateral
Amygdala. Neuropsychopharmacology 43:373-383.
980

Extended Data

Figure 2-1. Neither BLA or vlOFC inhibition affected the detection of action-based reversals as measured by probability correct adjustments. Rats were always tested on a deterministic schedule before a probabilistic one. Shown are deterministic (100/0) and probabilistic (90/10) reversal "transitions," 100 trials before and after each reversal. Drug order was counterbalanced such that on R2 and R4 animals received VEH if they were administered CNO first on R1 and R3, and vice versa. Inhibition of BLA impaired probabilistic reversal learning as indicated by selectively poor performance in 90/10 at the beginning and end of that reversal. OFC inhibition resulted in no impairment around these reversals. There was also no effect of CNO in the eGFP group.

983
984

Action

Figure 2-2. BLA, but not vlOFC, hM4Di exhibited longer incorrect choice and reward latencies than eGFP during action-based reversals. (A) *There were no significant differences in incorrect choice and reward latencies when comparing vlOFC hM4Di vs. eGFP controls for any reversals.* ***(B)*** *BLA hM4Di animals exhibited longer incorrect choice and reward latencies than eGFP controls during reversals. ***p<0.0001, **p<0.001, *p<0.05*

Figure 3-1. Reinforcement learning model fit to choice behavior indicates differential effect on the decay parameter between male and female rats during deterministic and probabilistic action-based reversals. *Plotted are estimated model parameters for RL1$_{decay}$ fitted to each session during deterministic (100/0) **(A)** and probabilistic (90/10) **(B)** reversals. α = learning rate, β = inverse temperature, γ_d = decay rate for unchosen option. Drug order was counterbalanced such that on R2 and R4 (not shown) animals received VEH if they were administered CNO first on R1 and R3, and vice versa. Female eGFP group exhibited overall lower γ_d than male eGFP rats during the first deterministic reversal R1. Chemogenetic inhibition of vlOFC or BLA increased γ_d on the first probabilistic reversal R3 for female rats. Only comparisons with p<0.075 are shown. **p<0.01, *p<0.05.*

Figure 4-1. Probability correct during stimulus-based reversals. *Cumulative P(correct) for first 5 sessions of each deterministic (100/0) and probabilistic (90/10) reversal. Drug order was counterbalanced such that on R2 and R4 animals received VEH if they were administered CNO first on R1 and R3, and vice versa. Despite most animals reaching criterion on initial learning (Fig. 2C), animals exhibited poor reversal learning. Performance around reversals (3 sessions before and after each reversal) is shown in Fig.2. Reversal learning was particularly flat for R3 (first probabilistic reversal) for vlOFC hM4Di group compared to eGFP. Bonferroni-corrected post-hocs following mixed-effects GLM with sex as a covariate fixed factor wherein a session x virus interaction was found resulted in **p<0.01 effect of session in eGFP, not in hM4Di.*

Figure 4-2. Females commit more initiation omissions and take longer to initiate trials than males in stimulus-based reversals. (A-B) *Females committed more initiation omissions than males irrespective of virus group or drug condition across reversals. **(C-D)** Females take longer to initiate trials than males irrespective of virus group or drug condition across reversals.*