Identification of overlapping and distinct mural cell populations during early

embryonic development

Sarah Colijn[1#], Miku Nambara[1#], and Amber N. Stratman[1*]

[1]Department of Cell Biology and Physiology, Washington University in St. Louis, School

of Medicine, St. Louis, MO 63110

[#]Contributed equally

[*]Correspondence: a.stratman@wustl.edu (ANS), 1-314-273-7928

Running Title: *pdgfrb* and *tagln* in zebrafish embryos

Keywords**:** mural cell progenitor, vSMC, pericyte, perivascular cells, hypochord,

sclerotome, zebrafish

Summary Statement:

Detailed analysis of *pdgfrb/tagln* vascular expression patterns in embryonic zebrafish

reveals novel regions of *tagln*-positive populations such as the hypochord and a mural

cell progenitor population adjacent to the midline.

Abstract:

Mural cells are an essential perivascular cell population that associate with blood vessels and contribute to vascular stabilization and tone. In the embryonic zebrafish vasculature, *pdgfrb* and *tagln* are commonly used as markers for identifying pericytes and vascular smooth muscle cells (vSMCs). However, the expression patterns of these markers used in tandem have not been fully described. Here, we used the *Tg(pdgfrb:Gal4FF; UAS:RFP)* and *Tg(tagln:NLS-EGFP)* transgenic lines to identify single- and double-positive perivascular populations in the cranial, axial, and intersegmental vessels between 1 and 5 days post-fertilization. From this comparative analysis, we discovered two novel regions of *tagln*-positive cell populations that have the potential to function as mural cell precursors. Specifically, we found that the hypochord—a reportedly transient structure—contributes to *tagln*-positive cells along the dorsal aorta. We also identified a unique sclerotome-derived mural cell progenitor population that resides along the midline between the neural tube and notochord and contributes to intersegmental vessel mural cell coverage. Together, our findings highlight the variability and versatility of tracking *pdgfrb* and *tagln* expression in mural cells of the developing zebrafish embryo.

Introduction:

In vertebrate development, early blood vessels are initially comprised of a single layer of endothelial cells—which then recruit mural cells to assist in basement membrane deposition, vessel contractility, and cellular communication (Armulik et al., 2005). Mural cells can vary widely in morphology and gene expression patterns depending on vessel type, vascular bed, tissue origin, differentiation status, developmental stage, and more (Donadon and Santoro, 2021, Owens et al., 2004, Holm et al., 2018). Types of mural cells include pericytes—which are associated with smaller vessels such as capillaries—and vascular smooth muscle cells (vSMCs) on larger vessels. vSMCs will accumulate on arteries and—to a lesser extent—on veins during vascular development to provide tensile strength and promote basement membrane deposition (Stratman et al., 2017). Due to the diverse properties of mural cells, different gene expression profiles can be used to identify mural cell types along various regions of the vasculature. Therefore, identification of appropriate cellular markers for distinct developmental stages is necessary for the study of vessel stabilization in different vascular beds.

In the embryonic zebrafish, platelet-derived growth factor receptor beta (*pdgfrb*) and transgelin (*tagln*) are commonly used as markers to identify pericytes and vSMCs. However, the temporal expression patterns of these markers used in tandem has not been fully described. Pdgfrb—a surface receptor for the chemoattractant ligands of the Pdgf family—is used as a marker for pericytes and immature vSMCs (Hellstrom et al., 1999, Armulik et al., 2005, Ando et al., 2021b). Pdgfrb has been shown to be required for mural cell expansion both embryonically and postnatally (Leonard et al., 2022, Ando et al., 2021b, Stratman et al., 2020). Cranial and cardiac neural crest cells also express Pdgfrb and have the capacity to become mural cells in addition to giving rise to other mesenchymal cell populations (Smith and Tallquist, 2010, Majesky, 2007, Ando et al., 2016). Some mature mural cells—such as pericytes—continue to express Pdgfrb, though less than 1% of Pdgfrb-positive cells in a zebrafish larvae at 5 days post-fertilization (dpf) are pericytes (Shih et al., 2021). This suggests that most Pdgfrb-positive cells in early development are progenitors for diverse cell populations.

Tagln (or SM22)—an actin-binding protein from the calponin family—is commonly used to identify mature, contractile vSMCs (Chakraborty et al., 2019, Santoro et al.,

2009). Tagln and other actin-binding proteins confer contractile function to mature mural

cells, allowing them to regulate vascular tone and blood pressure (Rensen et al., 2007).

During development, Tagln is also expressed in intestinal smooth muscle, cardiac

muscle, and skeletal muscle cells (Seiler et al., 2010, Li et al., 1996), though these muscle

cells can be distinguished from vSMCs by observing cellular morphology and proximity

to the vasculature. Altogether, this suggests that the *pdgfrb* and *tagln* reporters can be

used to identify mural cells and their differentiation status by tracking single- versus

double-positive cells in discrete vascular beds over the course of early development.

In this report, we use *Tg(pdgfrb:Gal4FF; UAS:RFP)* and *Tg(tagln:NLS-EGFP)*

transgenic lines to identify single- and double-positive pericytes/vSMCs in the cranial,

axial, and intersegmental vessels (ISVs) at 1 to 5 dpf. Through this study, we have

uncovered two novel sites of *tagln*-positive cell populations that have the potential to

function as mural cell precursors in the zebrafish trunk. First, we found that the

hypochord—a reportedly transient structure present in fish and amphibians—contributes

to *tagln*-positive cells along the dorsal aorta. Second, we identified a sclerotome-derived

mural cell progenitor population that resides along the midline at the notochord-neural

tube interface and contributes to ISV mural cell coverage. These findings highlight

unexpected and nuanced gene expression patterns across perivascular cell maturation.

Results:

***pdgfrb* and *tagln* transgenic reporter lines**

To visualize mural cells in the zebrafish embryo, we crossed the established

transgenic lines *TgBAC(pdgfrb:Gal4FF)[ncv24]*, *Tg(5xUAS:RFP)[zf83]*, and *Tg(tagln:NLS-*

EGFP-2A-CFP-FTASE)[y450] to generate embryos that were heterozygous carriers for each

transgene. Expression of both *pdgfrb* and *tagln* were notable at all embryonic stages

analyzed—with the brightest signal for *pdgfrb* visible in the anterior and cranial regions,

and *tagln* in the skeletal muscle of the trunk and later in the intestinal compartment (Fig.

S1).

The *pdgfrb* reporter line utilizes the Gal4/UAS system (Ando et al., 2016)—

designated as *Tg(pdgfrb:Gal4FF; UAS:RFP)* in this report—and mosaic fluorescence was

occasionally observed during screening; embryos with the least mosaicism were selected

for use. The *tagln* reporter line—designated as *Tg(tagln:NLS-EGFP)*—contains a nuclear

localization signal that allows for GFP to be concentrated in the nucleus (Stratman et al.,

2017), a modification that facilitates the identification of individual *tagln*-positive cells in

regions with dense *tagln* expression. With this tool, our study provides a unique approach

to tracking *pdgfrb* and *tagln* localization in the zebrafish.

Imaging the mural cells of the cranial vasculature

Development of the zebrafish cranial vasculature has been thoroughly mapped

through numerous studies. This process involves the formation of the primordial hind- and

midbrain channels (PHBC/PMBC) and the lateral dorsal aorta (LDA) by 1 dpf, the growth

of the middle cerebral vein (MCeV), metencephalic artery (MtA), basilar artery (BA), and

Circle of Willis (CoW) by 1.5 dpf, and the sprouting and fusion of various central arteries

(CtAs) from the PHBC to the BA by 2 dpf (Ulrich et al., 2011, Fujita et al., 2011, Isogai et

al., 2001). These vessels continue to remodel, grow, and stabilize, though the patterning

remains mostly intact from 6 dpf and beyond (Isogai et al., 2001).

Mural cell coverage of the cranial vessels has also been frequently studied, with

the common understanding being that neural crest-derived *pdgfrb*-positive pericytes are

the predominant mural cell in the mid- and hindbrain prior to 5 dpf (Ando et al., 2019,

Ando et al., 2016, Ando et al., 2021a, Wang et al., 2014). Interestingly, Ando et al have

shown that some mural cells in the hindbrain are derived from mesoderm. Specifically,

pdgfrb-positive cells surrounding the PHBC at 1 dpf are purportedly mesoderm-derived

mesenchyme; these cells then increase *pdgfrb* expression, migrate to CtAs, and

eventually acquire vSMC markers—such as *tagln*—at 5 dpf (Ando et al., 2016, Ando et

al., 2021a). Prior to 5 dpf, vSMC markers are sparse in the cranial vessels, with less than

10% of mural cells on the CtAs expressing *tagln* by 5 dpf (Ando et al., 2019).

In our *Tg(tagln:NLS-EGFP); Tg(pdgfrb:Gal4FF; UAS:RFP)* embryos, as expected,

pdgfrb was the predominant marker of cranial mural cells between 1 and 5 dpf (Figs 1A-

J; S2A-K). At 1 dpf, *pdgfrb*-positive cells were loosely associated with the PHBC (Fig.

1B)—corroborating previous data (Ando et al., 2021a, Ando et al., 2016)—and with the

MtA and MCeV (Fig. S2B,C). Based on the current understanding of the literature, these

early *pdgfrb*-positive cells are likely a combination of neural crest and mesoderm-derived

mesenchyme that have the potential to migrate to the vessels and continue to mature into
mural cells throughout development.

After 1 dpf, *pdgfrb*-positive cells slowly accumulate on the CoW and CtAs until 5
dpf, the latest stage analyzed (Figs 1C-J; S2D-K). A single *pdgfrb*/*tagln* double-positive
cell can be seen at 4 dpf on the middle mesencephalic central artery (MMCtA; Fig. 1H,
inset). However, throughout this region, *tagln*-positive cells were extremely sparse at all
developmental stages assessed, consistent with previous reports that vSMC markers
such as *tagln* are not prevalent in cranial mural cells prior to 4 dpf. These results indicate
that *pdgfrb* is the predominant mural cell marker along embryonic cranial vessels, though
a small proportion of these cells—particularly on the arteries—can mature into
pdgfrb/*tagln* double-positive vSMCs by 5 dpf.

Imaging the mural cells of the axial vasculature
The main axial vessels—i.e. dorsal aorta (DA) and posterior cardinal vein (PCV)
in the anterior trunk and the caudal artery (CA) and caudal vein (CV) in the tail—are some
of the first vessels to form prior to the onset of circulation at 1 dpf (Isogai et al., 2001).
The mural cells that accumulate on the ventral side of the DA have been well-described;
pdgfrb-positive cells begin to accumulate at 1.5 dpf and then acquire the vSMC marker
tagln at 3 dpf (Ando et al., 2021a, Stratman et al., 2017, Ando et al., 2016). Approximately
25% of vSMCs on the ventral side of the DA are reported to be *pdgfrb*/*tagln* double-
positive by 5 dpf (Ando et al., 2019).

The mural cells that cover the dorsal side of the DA are not as well-defined.
Intriguingly, the mural cells that accumulate on the dorsal and ventral sides of the DA are
initially derived from different mesodermal origins in mice (Wasteson et al., 2008). This
lineage difference has not been definitively shown in zebrafish. However, there are clear
distinctions between the cells that initially populate the dorsal versus ventral sides of the
DA in zebrafish embryos—with the mural cells on the dorsal side of the DA reported to
express *tagln* but not *pdgfrb* (Ando et al., 2016).

By tracking the expression of *pdgfrb* and *tagln* in vSMCs on the DA, PCV, CA, and
CV, we identify the hypochord as a novel source of *tagln*-positive cells on the dorsal side
of the DA and CA (Fig. 2). We also corroborate the previous findings regarding vSMC

expression patterns on the ventral side of the DA, and observe a slow accumulation of
pdgfrb single-positive cells on the PCV and CV between 1 to 5 dpf (Fig. 2). Finally, we
find that accumulation and maturation of these mural cells occurs in an anterior to
posterior fashion, with mural cells on the anterior DA and PCV maturing one to two days
earlier than the mural cells of the posterior CA and CV (Fig. 2).

To begin, we analyzed *pdgfrb* and *tagln* expression in mural cells of the anterior
DA and PCV in *Tg(tagln:NLS-EGFP); Tg(pdgfrb:Gal4FF; UAS:RFP)* embryos between 1
to 5 dpf (Fig. 3A-I). We identified the hypochord at 1 dpf as the *pdgfrb*-positive structure
immediately dorsal to the DA that runs along the midline (Fig. 3A). The hypochord is a
reportedly transient, rod-like structure—only one cell in width and height—that resides
beneath the notochord and assists in the formation and patterning of the DA (Eriksson
and Löfberg, 2000, Cleaver et al., 2000, Löfberg and Collazo, 1997). Once the DA is
formed, the hypochord cells are reported to delaminate and degenerate—in part due to
apoptosis (Löfberg and Collazo, 1997, Cleaver et al., 2000)—then by 2.5 dpf either
disappear or become indistinguishable from the cells that surround the DA in the zebrafish
embryo (Eriksson and Löfberg, 2000).

pdgfrb expression in the hypochord has been previously reported, in addition to
pdgfrb expression in the floor plate—the structure that sits on the ventral side of the neural
tube (Ando et al., 2019, Ando et al., 2016, Wiens et al., 2010). We found that in addition
to *pdgfrb*, the hypochord also expresses *tagln* at the 1 dpf stage (Fig. 3A,F). At 2 dpf, the
hypochord cells began to delaminate, as evidenced by the elongating nuclei (Fig. 3B). At
3 dpf, when the hypochord is allegedly absent from the anterior trunk (Eriksson and
Löfberg, 2000), we observed that the *tagln*-positive cells persist and begin to decrease
pdgfrb expression (Fig. 3C,F). By 4 dpf, these cells were *tagln* single-positive, with
noticeably fewer nuclei visible per vessel length (Fig. 3D,F; Movie 1), indicating that a
portion of hypochord cells do degenerate after delamination. However, a portion of
elongated *tagln*-positive cells were still present at 5 dpf (Fig. 3E,F), and Ando et al reports
observing these *tagln* single-positive cells above the DA as late as 7 dpf (Ando et al.,
2016).

To confirm that these hypochord-derived cells are maintained—and that they are
not ventral vSMCs that have migrated to the dorsal side of the DA—we used a

sclerotome-marker twist family bHLH transcription factor 1a (*twist1a*) transgenic line to
distinguish their origin (Stratman et al., 2017). The hypochord—in addition to the
notochord and floor plate—has been shown to be derived from the multipotent midline
progenitor cells in the dorsal organizer or the tailbud (Row et al., 2016, Latimer and Appel,
2006). Thus, hypochord cells maintain distinct origins from sclerotome cells, which
emerge from the somites (Tani et al., 2020). Indeed, we found that the *tagln* single-
positive cells on the dorsal side of the DA are not sclerotome-derived, unlike the vSMCs
on the ventral side of the DA (Fig. S3A-F). Altogether, these results suggest that the
hypochord is the source of the *tagln* single-positive cells that reside on the dorsal side of
the DA. This also indicates that the cells of the hypochord are not entirely transient as
previously assumed, but instead a portion reprogram their cellular function and
morphology after the DA has been appropriately oriented.

We next observed the development of the mural cells on the ventral side of the
DA. While the hypochord-derived cells on the dorsal side of the DA flattened and
elongated over time, the vSMCs on the ventral side were more globular (Fig. 3A-E).
pdgfrb single-positive vSMCs were observed beginning at 2 dpf, where they then
expanded in number and increased expression of *tagln* (Fig. 3A-E,G). By 5 dpf,
pdgfrb/*tagln* double-positive cells accounted for approximately 60% of ventral vSMCs,
and no *tagln* single-positive cells were detected. These data are consistent with previous
reports of vSMCs on the ventral side of the DA, though we observe a higher percentage
of double-positive vSMCs (60% versus 25%) by 5 dpf (Ando et al., 2019, Ando et al.,
2016, Ando et al., 2021a).

Furthermore, we detected *pdgfrb* single-positive cells associated with the PCV as
early as 1 dpf—though these cells were sparse—followed by the slow and sporadic
accumulation of *pdgfrb*-positive cells until 5 dpf (Fig. 3A-E,H). This timing is earlier than
suggested in previous reports where *pdgfrb* expression was undetectable in cells around
the PCV prior to 7 dpf (Ando et al., 2016). Stratman et al describes a small number of
tagln-positive cells on the PCV around 4 dpf (Stratman et al., 2017). Similarly, we saw
low numbers of weak *tagln*-positive cells beginning at 3 dpf, though these cells were less
common than *pdgfrb* single-positive cells (Fig. 3H). These data indicate that mural cells
of the PCV associate and mature at much later developmental stages than aortic vSMCs.

In the posterior trunk region, mural cells of the CA and CV associate with vessels
and pattern similarly to the anterior mural cells but on a delayed time course (Fig. 4A-I).
At 1 dpf, the hypochord cells were block-like and were still distinguishable as hypochord
cells by 2 dpf (Fig. 4A,B,F). This was expected, as the hypochord "disappears" in an
anterior to posterior fashion (Eriksson and Löfberg, 2000), and is evident at 2 dpf in the
posterior region even when it is no longer detectable in the anterior region. We saw the
hypochord cells had begun to delaminate by 3 dpf (Fig. 4C), at which point they appeared
comparable to the 2 dpf hypochord cells in the anterior region (Fig. 3B). While the cells
then began turning off *pdgfrb*, only about a third of hypochord-derived cells in the posterior
trunk were *tagln* single-positive by 5 dpf (Fig. 4C-F), indicating a delay in hypochord
reprogramming compared to the anterior hypochord.

Furthermore, the ventral side of the CA began acquiring *pdgfrb* single-positive cells
at 2 dpf—similar to the DA—but they did not begin co-expressing *tagln* until 5 dpf (Fig.
4B-E,G). This amounted to a 2-day delay in mural cell development compared to the
anterior region (Fig. 3G). Finally, mural cell coverage of the CV was particularly sparse
(Fig. 4A-E,H), with half as many cells associated with the CV compared to the PCV at the
same developmental stage (Fig. 3H). These data indicate that association and maturation
of vSMCs on the main axial vessels occurs in an anterior to posterior fashion across time.

Of note, weak *tagln* expression was detected in endothelial cells of the axial
vasculature (Fig. S4A,B), which is consistent with previous reports that the endothelium
can express low levels of *tagln* (Chakraborty et al., 2019, Tsuji-Tamura et al., 2021).
Despite this, *tagln* expression in the endothelial cells was easily distinguishable from
vSMCs; therefore, it did not confound the mural cell counts of the axial vessels.

Imaging the mural cells of the intersegmental vessels (ISVs)

We next analyzed the mural cell populations surrounding the ISVs of the trunk
region. At 1 dpf, arterial ISVs (aISVs) begin to sprout from the DA and grow dorsally
between the somites and the notochord/neural tube. At 1.5 dpf, venous ISVs (vISVs)
sprout from the PCV, grow dorsally, and then with aISVs branch laterally to establish the
dorsal longitudinal anastomotic vessel (DLAV) that resides above the neural tube (Isogai

et al., 2003). By 2 dpf, the arterial and venous ISVs are mostly lumenized and reach from

the DA/PCV below the notochord to the DLAV above the neural tube (Isogai et al., 2001).

Mural cells of the ISVs have been reported to emerge by either *de novo*

differentiation of nearby mesenchyme followed by subsequent proliferation or through

migration from the ventral side of the DA (Ando et al., 2016). Studies identifying the origin

of ISV mural cells show that they are primarily derived from the sclerotome (Stratman et

al., 2017, Rajan et al., 2020)—a sub-compartment of the somites and a derivative of

paraxial mesoderm (Tani et al., 2020). Zebrafish have recently been shown to have a

dorsal and ventral sclerotome, both of which contribute to the cells that associate with the

ISVs: dorsal sclerotome populates the ISV region adjacent to the neural tube while ventral

sclerotome populates the full ISV (Rajan et al., 2020, Ma et al., 2018). However, while the

dorsal and ventral sclerotome cells can contribute to the development of different regions

of the embryo (Ma et al., 2018), it is currently unclear whether they differ in gene

expression.

Rajan et al recently described a population of sclerotome-derived cells called

perivascular fibroblasts that associate with ISVs by 2 dpf and produce collagen matrix,

thus providing structural support prior to mural cell migration or differentiation (Rajan et

al., 2020). These cells are primarily derived from ventral sclerotome, are evenly

distributed between arterial and venous ISVs, remain constant in number on the ISVs

from 2 to 4 dpf, and express low levels of *pdgfrb* (Rajan et al., 2020). At 3 dpf, a subset—

but not all—of perivascular fibroblasts will increase *pdgfrb* expression to become ISV

mural cells (Rajan et al., 2020). Ando et al also describes a naïve *pdgfrb*-low

mesenchymal population—likely the same cell type or similar to perivascular fibroblasts—

that can mature into *pdgfrb*-high mural cells on the ISVs starting at approximately 2.5 dpf

(Ando et al., 2016, Ando et al., 2021a).

Both studies observe that mural cells with *pdgfrb*-high expression are more

apparent on aISVs than vISVs (Ando et al., 2016, Rajan et al., 2020), which is in part due

to maturation of *pdgfrb*-low mesenchyme to *pdgfrb*-high mural cells through Notch

signaling in arterial endothelium (Ando et al., 2019). Ando et al also observes that

approximately 15% of mural cells are *tagln*-positive on ISVs by 5 dpf (Ando et al., 2019),

potentially indicating further maturation of these *pdgfrb*-positive cells.

In support of these previous studies, we observed *pdgfrb*-low and -high cell
populations surrounding the ISVs up to 5 dpf in *Tg(tagln:NLS-EGFP); Tg(pdgfrb:Gal4FF;*
UAS:RFP) embryos (Figs 5A-H; S5A-I). Unexpectedly, we also detected *tagln*-low and -
high populations as early as 2 dpf (Figs 5C,D; S5A,B), prior to the developmental stages
in which ISV mural cells reportedly mature. While a subset of *tagln*-low cells in this region
were endothelial cells, we confirmed that many of the *tagln*-low and -high cells associated
with the early ISVs were perivascular (Fig. S4C,D), indicating that *tagln* may not
exclusively label mature mural cells, but instead also labels a subset of mural cell
precursors.

From our ISV data, we observed findings similar to that of the perivascular
fibroblasts (Rajan et al., 2020): the number of perivascular cells associated with the ISVs
remained constant between 2 to 5 dpf, the cells were equally distributed on aISVs and
vISVs, and *pdgfrb*-high cells became more apparent on aISVs than vISVs over time (Figs
5A-D; S5A-H). Furthermore, we were able to identify *pdgfrb*/*tagln* double-positive cells
with all combinations of high and low expression that increased in proportion up to 4 dpf
(Fig. 5E-G). Our data showed that double-positive cells were more common on aISVs
than vISVs and, as expected, the double-positive cells on the aISVs were more likely to
be *pdgfrb*-high (Fig. 5F,G). *tagln*-high mural cells were constant on all ISVs between 2-5
dpf (Fig. 5C,D), suggesting that on ISVs, mural cells may diverge from the typical
expression patterns of differentiation that would predict upregulated *tagln* at late stages
in the process of maturation.

tagln-low cells were the most common perivascular cell associated with the ISVs
at early timepoints (Fig. 5C,D). We speculated that the identity of these cells would
overlap with the previously described sclerotome-derived perivascular fibroblast
population. Indeed, when using the *twist1a* transgenic line as a sclerotome marker, we
identified *twist1a*-positive cells that reside at the notochord-neural tube interface and
notochord region as *tagln*-low, even prior to their association with the ISVs (Fig. S6A,D,E).
Conversely, a subset of cells that likely originated from the dorsal sclerotome—based on
their localization adjacent to the neural tube (Rajan et al., 2020)—were *tagln*-negative
(Fig. S6B,C). These data indicate that the ventral sclerotome is likely distinct from dorsal

sclerotome by the early and low expression of *tagln*. Additionally, these data reveal that

ISV mural cells can express *tagln* as precursors prior to upregulating *pdgfrb*.

We also observed consistent *tagln*/*pdgfrb* expression patterns in mural cells based

on their localization on the ISVs. *tagln* single-positive cells were more apparent on the

ISVs in the region adjacent to the notochord (Fig. 5H), consistent with previous reports of

ventral sclerotome migration to this region (Rajan et al., 2020, Ma et al., 2018). *pdgfrb*

single-positive cells were highly associated with the ISVs near to the neural tube (Fig.

5H), where the ISVs are primarily populated by dorsal sclerotome (Rajan et al., 2020).

Finally, *pdgfrb*/*tagln* double-positive cells were evident at two sites by 5 dpf—closest to

the notochord-neural tube interface (Fig. 5H), which could indicate a site-specific

maturation event, and near DA/aISV boundaries, which would suggest that these are

vSMCs from the ventral side of the DA that have migrated to the aISVs.

A unique sclerotome-derived mural cell progenitor population resides along the

zebrafish midline

From imaging *tagln*- and *pdgfrb*-positive cells near the ISVs in *Tg(tagln:NLS-*

EGFP); Tg(pdgfrb:Gal4FF; UAS:RFP) embryos, we discovered several cell populations

with unexpected *pdgfrb*/*tagln* expression near the midline. First, we note the medial floor

plate cells are *pdgfrb*/*tagln* double-positive (Fig. 6, MFP cells). Additionally, we observe

a *tagln* single-positive population of cells along the midline that accumulates below the

medial floor plate cells and above the notochord beginning at 3 dpf (Fig. 6, *tagln*+ midline

cells). Finally, we describe a mural cell progenitor population near the notochord-neural

tube interface that borders either side of the midline. This population—which begins as

tagln single-positive cells before acquiring *pdgfrb* expression—can be observed migrating

to the nearby ISVs from 2 to 5 dpf (Fig. 6, MC progenitor population).

Of the structures that reside at or near the midline, the medial floor plate is one of

the most recognizable due to its hourglass-like shape (Zheng et al., 2022). Medial floor

plate cells—which reside at the ventral side of the neural tube and contribute to neuronal

differentiation (Strähle et al., 2004)—have been previously identified as *pdgfrb*-positive

(Ando et al., 2016, Wiens et al., 2010). We observed intense *tagln* co-expression in the

medial floor plate cells from 1 to 5 dpf (Fig. 7A-K), indicating that *tagln* may have as-of-
yet unidentified developmental functions in primitive, non-muscular tissues.

Furthermore, we note a *tagln* single-positive population of elongated cells along
the midline beginning at 3 dpf that accumulates between the notochord and neural tube—
immediately ventral to the medial floor plate cells (Fig. 7A-K). This population appears to
organize as a continuous, thin, rod-like structure that is primarily one cell in width and
height. We confirmed that this *tagln*-positive midline cell population originates from the
sclerotome—unlike the adjacent medial floor plate—as evidenced by *twist1a* expression
(Fig. 7L-O). The function of this previously unidentified structure is currently unknown.

Finally, we observed a cell population—which we will term mural cell progenitors—
near the notochord-neural tube interface that borders either side of the midline and
migrates to the ISVs (Fig. 7A-W; Movies 2,3). At 1 dpf, these cells expressed *tagln*, but
not *pdgfrb* (Fig. 7A'), and at 2 dpf, a fraction of these cells began to express *pdgfrb* (Fig.
7C'). Over the next 3 days these cells elongated and increased *pdgfrb* expression until
approximately half were *pdgfrb*/*tagln* double-positive at 5 dpf (Fig. 7E',G',I').

Due to this mural cell progenitor population's proximity to the notochord and floor
plate, we questioned whether they originated from or resided within either of these
structures. Using a sonic hedgehog signaling molecule a (*shha*) transgenic line to label
the floor plate (Shkumatava et al., 2004, Ertzer et al., 2007), we were able to visualize the
mural cell progenitor population as sitting outside of the neural tube (Fig. S7A,B). Next,
we used a procollagen, type IX, alpha 2 (*col9a2*) transgenic line to visualize notochord
sheath cells in relation to the progenitor population (Garcia et al., 2017). Notochord
sheath cells are reported to express *tagln* in the notochord bud during regeneration
(Sinclair et al., 2021); therefore, we wanted to determine if the sheath cells may be the
source of the mural cell progenitor population. However, the progenitor population did not
overlap with notochord sheath cells (Fig. S7C,D), though we confirm that sheath cells do
express low levels of *tagln* during development (Fig. S7E). Jointly, these data showed
that the mural cell progenitor population abuts the neural tube and notochord, but does
not overlap with these structures.

We next used the sclerotome-marker *twist1a* and the neural crest-marker SRY-
box transcription factor 10 (*sox10*) transgenic lines (Lee et al., 2013, Kucenas et al., 2008)

to test the origins of this novel mural cell progenitor population. We identified this
progenitor population as exclusively derived from *twist1a*-positive sclerotome and not
sox10-positive neural crest (Figs 7L-O; S7F,G; Movie 3). Since Notch signaling from
arterial endothelium has been shown to be involved in maturation of mural cell progenitors
to *pdgfrb*-high mural cells (Ando et al., 2019), we speculated that the progenitor
population could be in contact with a blood vessel along the midline that would induce an
increase in *pdgfrb* expression. However, there is no blood vessel present along the
midline at this time point (Fig. S7H), nor do cells that increase *pdgfrb* expression in the
progenitor population reside near blood vessels (Fig. S7I,J). This suggests that the
induction of *pdgfrb* expression in this mural cell progenitor population is not controlled by
contact with the arterial vasculature.

Using *tagln*, *pdgfrb*, and *twist1a* as cellular markers, we observed cells from this
progenitor population migrate to the ISVs along the region adjacent to the notochord-
neural tube interface (Fig. 7P-W; Movies 2,3). Progenitor cells migrated to both aISVs
and vISVs from 2 to 5 dpf. Due to the location of this mural cell progenitor population in
relation to the ISVs, a proportion of the *pdgfrb*/*tagln* double-positive cells that we observe
on ISVs in the region closest to the notochord-neural tube interface (Fig. 5H) likely
originate from this progenitor source.

Discussion:
pdgfrb **and** ***tagln*** **transgenic zebrafish reporters**
In this study, we use *Tg(pdgfrb:Gal4FF; UAS:RFP)* and *Tg(tagln:NLS-EGFP)*
transgenic lines to identify single- and double-positive pericytes/vSMCs in the cranial,
axial, and ISVs at 1 to 5 dpf. Our nuclear-GFP *tagln* reporter facilitated identification of
novel *tagln*-positive cells in the zebrafish trunk. High *tagln* expression is evident in skeletal
muscle, which has made it difficult in the past to explore *tagln* expression patterns in cells
near to the skeletal muscle with cytosolic-GFP *tagln* reporters. With the nuclear-GFP *tagln*
reporter, we have revealed *tagln* expression in the hypochord, the medial floor plate, and
in sclerotome-derived mesenchyme in the trunk region.

To rule out whether these expression patterns were an artifact of the nuclear-GFP
tagln reporter, we sought to confirm *tagln* expression in these structures using an

independently generated cytosolic-GFP *tagln* reporter (*Tg(tagln:EGFP)*; Fig. S8A-H). As

expected, GFP expression in the skeletal muscle decreased the efficiency and resolution

of identifying *tagln*-positive cells in the trunk, though we were able to visualize GFP

expression in the hypochord, ISV mural cells, the novel mural cell progenitor population

we describe, the medial floor plate, and *tagln*-positive midline cells. Thus, despite high

background GFP signal, we validate *tagln* expression in these structures using the

cytosolic-GFP *tagln* reporter.

The *pdgfrb* reporter line—which utilizes the Gal4/UAS system (Ando et al., 2016)—

presented with occasional RFP expression mosaicism. This mosaicism, when present,

was most apparent in the medial floor plate cells (Fig. 7A-J), though it is difficult to assess

the penetrance at which other structures were affected. Due to this, there may be *pdgfrb*-

positive cells that are underrepresented in this study, particularly cells that express low

levels of *pdgfrb*. However, despite potential mosaicism issues, the expression patterns

that we visualized over time and in various regions were consistent across all samples.

Cranial Mural Cells

Previous reports show that *pdgfrb*-positive pericytes are the predominant mural

cell in the mid- and hindbrain prior to 5 dpf, at which point arterial-associated mural cells

begin to express *tagln* and other vSMC markers (Ando et al., 2019, Ando et al., 2016,

Ando et al., 2021a, Whitesell et al., 2014, Wang et al., 2014). We observed similar

findings, in that *pdgfrb* single-positive pericytes were the most prevalent type of mural cell

along embryonic cranial vessels, with eventual *tagln* co-expression evident—yet sparsely

represented—in a subset of *pdgfrb*-positive arterial mural cells at 4 dpf. No *tagln* single-

positive cells were detected at the time points assessed. Based on these data, and

supported by the findings of previous studies, we conclude that *pdgfrb* is an ideal reporter

to visualize cranial mural cell association with blood vessels up to 5 dpf, with the caveat

that a large proportion of early *pdgfrb*-positive cranial cells are neural crest or

mesenchyme progenitors that are not associated with vessels.

Mural Cells of the Axial Vessels

In the axial vessels (i.e. DA, PCV, CA, and CV), we corroborate the previous findings regarding vSMC expression patterns on the ventral side of the DA (Ando et al., 2021a, Stratman et al., 2017, Ando et al., 2016, Ando et al., 2019). Specifically, we detect *pdgfrb* single-positive cells at 2 dpf, where they then expand in number and increase expression of *tagln* starting at 3 dpf (Fig. 2). Moreover, we observe a slow and sporadic increase of *pdgfrb* single-positive cells on the PCV between 1 to 5 dpf (Fig. 2). At 5 dpf, there are twice as many *pdgfrb*-positive mural cells associated with the ventral side of the DA than the entire PCV, which highlights the clear difference in the magnitude of mural cell recruitment between arteries and veins early in development.

Our analysis of both the anterior and caudal axial vessels also reveals that accumulation and maturation of these mural cells occurs in an anterior to posterior fashion. By 1 dpf, the anterior and posterior regions of the main axial vessels have formed, somite segmentation has been completed, and the sclerotome is distinct from other somitic tissues (Stickney et al., 2000). Despite this, DA vSMCs—which are mainly derived from the sclerotome (Stratman et al., 2017)—mature one to two days earlier than the vSMCs of the CA (Fig. 2). There may be several reasons for this delay, with one hypothesis being that the DA provides earlier signals than the CA to initiate maturation of vSMCs. Alternatively, since somites develop in an anterior to posterior wave (Stickney et al., 2000), the age of the somites from which mural cells are derived could influence the timing of vSMC maturation, even after vSMCs have accumulated on the vessels. More work will be needed in this area to resolve the cues driving this temporal differentiation pattern.

The Hypochord as a Source of *tagln*-Positive Cells

Through use of the nuclear-GFP *tagln* reporter, we found that the hypochord contributes to *tagln*-positive cells along the dorsal side of the DA up to at least 5 dpf (Fig. 2). Interestingly, the hypochord and medial floor plate—which we also show is *tagln*-positive—are among the earliest structures in the zebrafish embryo to co-express *pdgfrb* and *tagln*. The hypochord, floor plate, and notochord have been shown to be derived from the multipotent midline progenitor cells in the dorsal organizer or the tailbud (Row et al., 2016, Latimer and Appel, 2006)—a common origin which may, in part, explain how two

structures that are dissimilar in function share this unique expression pattern.
Furthermore, in the invertebrate *Ciona*, researchers show that a series of
calponin/transgelin family genes (*tagln-r.b, tagln-r.c, tagln-r.d, tagln-r.e*) are expressed in
the *Ciona* floor plate, notochord, and endodermal strand (the *Ciona* analogue for the
hypochord), indicating that Tagln expression in midline structures may have a conserved
role in embryogenesis (Oonuma et al., 2021). As Tagln is known primarily as an actin-
binding protein in contractile cells, the role of Tagln in these non-contractile, non-
muscular, primitive midline structures remains unknown.

The persistence of the elongated *tagln*-positive cells on the dorsal side of the DA
indicates that the cells of the hypochord are not entirely transient, as previously assumed.
While previous studies have observed hypochord cells undergoing apoptosis (Löfberg
and Collazo, 1997, Cleaver et al., 2000), we propose that a subset of hypochord cells
survive and reprogram their cellular function and morphology after the DA has been
appropriately oriented. Due to the maintained expression of *tagln* and proximity to the DA,
we propose these cells may undertake a mural cell role; however, further research is
needed to determine whether these cells function similarly to other vSMC populations.

As the hypochord is present exclusively in fish and amphibians, we questioned
whether there is an analogous mammalian structure that could contribute to Tagln-
positive cells on the DA of amniotes. Previous reports speculate that the dorsal
endoderm—a midline structure that originates from the midline precursor cell population
and forms ventral to the notochord (Peyrot et al., 2011)—may play a role in angioblast
and DA formation (Poole et al., 2001), similar to the hypochord. However, while the
endoderm gives rise to the intestines, lung, liver, and other organs (Wlizla and Zorn,
2015), there are no known reports of endoderm contributing to mural cell populations in
amniotes (Holm et al., 2018).

In the mouse embryo, the dorsal and ventral sides of the DA are populated by
vSMCs from different sources: the ventral side is first populated by cells from the lateral
plate mesoderm, with dorsal vSMCs originating from the somites. Over time, the vSMCs
from the lateral plate mesoderm disappear, and the ventral side then becomes populated
with somitic vSMCs (Wasteson et al., 2008). While this process is not identical on the
zebrafish DA—given that somitic vSMCs are the initial populators of the ventral side of

the DA in zebrafish—it does indicate that dorsal and ventral vSMCs can arise from
divergent origins and supports that this could be a conserved process.

Mural Cells of the Intersegmental Vessels

In the ISV region, we note perivascular cells associated with the vessels exhibiting
all combinations of high and low expression of *pdgfrb* and *tagln*. To summarize, *pdgfrb*-
high cells were more common on arterial than venous ISVs and increased in number over
time. *pdgfrb/tagln* double-positive cells were also more common on aISVs and were more
likely to be *pdgfrb*-high on aISVs than vISVs. *tagln*-high cells were constant in number
between 2 to 5 dpf, and *tagln*-low cells—the most common expression pattern
represented—were revealed to be sclerotome-derived cells that were migrating to or
associated with the ISVs. Further, expression patterns were related to mural cell location
on the ISVs, with *pdgfrb* single-positive cells more likely to be found on the dorsal ISV
region, double-positive cells nearest to the notochord-neural tube interface, and *tagln*
single-positive cells primarily near the notochord ISV region by 5 dpf (Fig. 5H).

These data indicate that *tagln* may not exclusively label mature mural cells, but
instead could also label a subset of mural cell precursors in the trunk. These precursors
likely overlap with the perivascular fibroblast population described by Rajan et al (Rajan
et al., 2020), as they are both sclerotome-derived and populate the ISVs prior to reported
mural cell maturation. This would suggest that a subset, but not all, of the *tagln*-positive
early precursors on the ISVs will mature into mural cells.

Sclerotome-Derived *tagln*-Positive Cell Populations in the Trunk

Use of our nuclear-GFP *tagln* reporter allowed us to ascertain that sclerotome-
derived cells in the zebrafish trunk are primarily *tagln*-positive as early as 1 dpf. Due to
the main localization of these *tagln*-positive cells at the notochord-neural tube interface
or below, we speculate that the *twist1a/tagln* double-positive cells arise from the ventral
sclerotome, with *twist1a*-positive/*tagln*-negative cells descending from the dorsal
sclerotome. Of note, there are two unique sites where sclerotome-derived *tagln*-positive
cells accumulate during development.

First, we identified a *twist1a*/*tagln* double-positive midline population between the
neural tube and notochord. This population begins to accumulate beneath the medial floor
plate cells at 3 dpf and exhibits high *tagln* expression (Fig. 6, *tagln*+ midline cells). Though
this structure's function is unknown, its location would suggest that it may exist as a
cushion or as structural support to the notochord-neural tube interface.

Second, we identified a mural cell progenitor population that resides adjacent to
the midline at the notochord-neural tube interface. This *tagln*-positive population
upregulates *pdgfrb* expression sporadically in an increasing number of cells between 2 to
5 dpf and migrates to the ISVs (Fig. 6, MC progenitor population; Movies 2,3). Due to the
location of this mural cell progenitor population in relation to the ISVs, we suspect that a
proportion of the *pdgfrb*/*tagln* double-positive cells that we observe on ISVs in the region
closest to the notochord-neural tube interface likely originate from this progenitor source.
Uniquely, the upregulation of *pdgfrb* expression in this mural cell progenitor population
occurs independently of contact with arterial endothelium. We hypothesize that
upregulation of *pdgfrb* confers motility cues to the mural cell progenitors to then migrate
to blood vessels; however, further studies are needed to determine the mechanisms
behind this seemingly stochastic *pdgfrb* upregulation.

This progenitor population also demonstrates that mural cell precursors can
express *tagln* prior to upregulating *pdgfrb*. Since *pdgfrb* is associated with precursor
populations and *tagln* is regarded as a mature vSMC marker, this finding suggests that
the expression patterns related to mural cell maturation are more nuanced than previously
reported. Therefore, we ask: what role could Tagln play prior to Pdgfrb upregulation?
Several studies report that Tagln assists in cell motility: loss of Tagln decreases motility
(Elsafadi et al., 2016), whereas overexpression leads to pathological motility in situations
such as metastatic cancer (Zhou et al., 2016, Chen et al., 2019). Since sclerotome cells
are inherently migratory as they populate and form various tissues, this could help explain
the early expression of Tagln in mural cell precursors during embryogenesis. Future work
in this area will continue to reveal exciting new findings driving site-specific and temporal
regulation of mural cell biology.

Materials and Methods:

Zebrafish

Zebrafish (*Danio rerio*) embryos were raised and maintained as described (Westerfield, 2000, Kimmel et al., 1995). Zebrafish husbandry and research protocols were reviewed and approved by the WUSTL Animal Care and Use Committee. Established transgenic lines that were used in this study are listed in the table below.

Genotype	Allele	Origin
Tg(-2.4shha-ABC:GFP)	*sb15*	(Shkumatava et al., 2004)
Tg(col9a2:mCherry)	*pd1150*	(Garcia et al., 2017)
Tg(kdrl:mCherry-CAAX)	*y171*	(Fujita et al., 2011)
Tg(Ola.Twist:GAL4)	*sq11*	(Lee et al., 2013)
Tg(sox10:mRFP)	*vu234*	(Kucenas et al., 2008)
Tg(tagln:eGFP)	*p151*	(Seiler et al., 2010)
Tg(tagln:NLS-EGFP-2A-CFP-FTASE)	*y450*	(Stratman et al., 2017)
Tg(5xUAS:RFP)	*zf83*	(Asakawa et al., 2008)
TgBAC(pdgfrb:Gal4FF)	*ncv24*	(Ando et al., 2016)

Image Acquisition and Processing

Embryos and larvae were anesthetized in buffered 0.2 mg/mL Tricaine-S (Western Chemical, Inc; NC0872873) and mounted in 0.8% low-melting point agarose (IBI Scientific; #IB70056) on a 35-mm glass-bottom petri dish (MatTek; #P35G-1.5-20-C). Confocal images were obtained using a 4x or 40x objective with a W1 Spinning Disk confocal microscope, a Fusion camera, and the Nikon Eclipse Ti2-E base. Images were then deconvolved using Nikon's NIS-Elements software. Fiji image processing software was used for counting number of cells per vessel length and for generating final images. Embryos were consistently imaged in the afternoon hours, thus: '1 dpf' represents embryos between 28-34 hours post-fertilization (hpf); '2 dpf' represents 52-58 hpf; '3 dpf' represents 76-82 hpf; '4 dpf' represents 100-106 hpf; '5 dpf' represents 124-130 hpf. At least 6 embryos were imaged and analyzed per developmental timepoint.

Acknowledgements:

We thank Tony Tsai's lab for the *Tg(-2.4shha-ABC:GFP)* line, Shawn Burgess's lab for

the *Tg(col9a2:mCherry)* line, and Mayssa Mokalled's lab for the *Tg(sox10:mRFP)* line.

Figure schematics were generated using BioRender.com.

Competing Interests:

The authors declare that they have no conflicts of interest.

Funding:

This work was supported by grants from the NIH/NIGMS R35 GM137976

(A.N.S.); Children's Discovery Institute of Washington University and St. Louis Children's

Hospital (A.N.S.); the Washington University Institute of Clinical and Translational

Sciences which is, in part, supported by the NIH/National Center for Advancing

Translational Sciences (NCATS), CTSA grant #UL1TR002345 (A.N.S.)*;* and American

Heart Association Postdoctoral Fellowship Award #906436 (S.C.).

References:

Ando, K., Fukuhara, S., Izumi, N., Nakajima, H., Fukui, H., Kelsh, R. N. & Mochizuki, N. (2016). Clarification of mural cell coverage of vascular endothelial cells by live imaging of zebrafish. *Development,* 143**,** 1328-1339. 10.1242/dev.132654

Ando, K., Ishii, T. & Fukuhara, S. (2021a). Zebrafish Vascular Mural Cell Biology: Recent Advances, Development, and Functions. *Life (Basel),* 11. 10.3390/life11101041

Ando, K., Shih, Y.-H., Ebarasi, L., Grosse, A., Portman, D., Chiba, A., Mattonet, K., Gerri, C., Stainier, D. Y. R., Mochizuki, N., Fukuhara, S., Betsholtz, C. & Lawson, N. D. (2021b). Conserved and context-dependent roles for pdgfrb signaling during zebrafish vascular mural cell development. *Developmental Biology,* 479**,** 11-22. https://doi.org/10.1016/j.ydbio.2021.06.010

Ando, K., Wang, W., Peng, D., Chiba, A., Lagendijk, A. K., Barske, L., Crump, J. G., Stainier, D. Y. R., Lendahl, U., Koltowska, K., Hogan, B. M., Fukuhara, S., Mochizuki, N. & Betsholtz, C. (2019). Peri-arterial specification of vascular mural cells from naïve mesenchyme requires Notch signaling. *Development,* 146. 10.1242/dev.165589

Armulik, A., Abramsson, A. & Betsholtz, C. (2005). Endothelial/Pericyte Interactions. *Circulation Research,* 97**,** 512-523. doi:10.1161/01.RES.0000182903.16652.d7

Asakawa, K., Suster, M. L., Mizusawa, K., Nagayoshi, S., Kotani, T., Urasaki, A., Kishimoto, Y., Hibi, M. & Kawakami, K. (2008). Genetic dissection of neural circuits by *Tol2* transposon-mediated Gal4 gene and enhancer trapping in zebrafish. *Proceedings of the National Academy of Sciences,* 105**,** 1255-1260. doi:10.1073/pnas.0704963105

Chakraborty, R., Saddouk, F. Z., Carrao, A. C., Krause, D. S., Greif, D. M. & Martin, K. A. (2019). Promoters to Study Vascular Smooth Muscle. *Arteriosclerosis, Thrombosis, and Vascular Biology,* 39**,** 603-612. doi:10.1161/ATVBAHA.119.312449

Chen, Z., He, S., Zhan, Y., He, A., Fang, D., Gong, Y., Li, X. & Zhou, L. (2019). TGF-b-induced transgelin promotes bladder cancer metastasis by regulating epithelial-mesenchymal transition and invadopodia formation. *eBioMedicine,* 47**,** 208-220. 10.1016/j.ebiom.2019.08.012

Cleaver, O., Seufert, D. W. & Krieg, P. A. (2000). Endoderm patterning by the notochord: development of the hypochord in Xenopus. *Development,* 127**,** 869-879. 10.1242/dev.127.4.869

Donadon, M. & Santoro, M. M. (2021). The origin and mechanisms of smooth muscle cell development in vertebrates. *Development,* 148. 10.1242/dev.197384

Elsafadi, M., Manikandan, M., Dawud, R. A., Alajez, N. M., Hamam, R., Alfayez, M., Kassem, M., Aldahmash, A. & Mahmood, A. (2016). Transgelin is a TGFβ-inducible gene that regulates osteoblastic and adipogenic differentiation of human skeletal stem cells through actin cytoskeleston organization. *Cell Death Dis,* 7**,** e2321. 10.1038/cddis.2016.196

Eriksson, J. & Löfberg, J. (2000). Development of the hypochord and dorsal aorta in the zebrafish embryo (Danio rerio). *Journal of Morphology,* 244**,** 167-176. https://doi.org/10.1002/(SICI)1097-4687(200006)244:3<167::AID-JMOR2>3.0.CO;2-J

Ertzer, R., Müller, F., Hadzhiev, Y., Rathnam, S., Fischer, N., Rastegar, S. & Strähle, U. (2007). Cooperation of sonic hedgehog enhancers in midline expression. *Developmental Biology,* 301**,** 578-589. https://doi.org/10.1016/j.ydbio.2006.11.004

Fujita, M., Cha, Y. R., Pham, V. N., Sakurai, A., Roman, B. L., Gutkind, J. S. & Weinstein, B. M. (2011). Assembly and patterning of the vascular network of the vertebrate hindbrain. *Development,* 138**,** 1705-1715. 10.1242/dev.058776

Garcia, J., Bagwell, J., Njaine, B., Norman, J., Levic, D. S., Wopat, S., Miller, S. E., Liu, X., Locasale, J. W., Stainier, D. Y. R. & Bagnat, M. (2017). Sheath Cell Invasion and Trans-differentiation Repair Mechanical Damage Caused by Loss of Caveolae in the Zebrafish Notochord. *Current Biology,* 27**,** 1982-1989.e3. https://doi.org/10.1016/j.cub.2017.05.035

Hellstrom, M., Kal N, M., Lindahl, P., Abramsson, A. & Betsholtz, C. (1999). Role of PDGF-B and PDGFR-beta in recruitment of vascular smooth muscle cells and pericytes during embryonic blood vessel formation in the mouse. *Development,* 126**,** 3047-3055.

Holm, A., Heumann, T. & Augustin, H. G. (2018). Microvascular Mural Cell Organotypic Heterogeneity and Functional Plasticity. *Trends in Cell Biology,* 28**,** 302-316. 10.1016/j.tcb.2017.12.002

Isogai, S., Horiguchi, M. & Weinstein, B. M. (2001). The Vascular Anatomy of the Developing Zebrafish: An Atlas of Embryonic and Early Larval Development. *Developmental Biology,* 230**,** 278-301. https://doi.org/10.1006/dbio.2000.9995

Isogai, S., Lawson, N. D., Torrealday, S., Horiguchi, M. & Weinstein, B. M. (2003). Angiogenic network formation in the developing vertebrate trunk. *Development,* 130**,** 5281-5290. 10.1242/dev.00733

Kimmel, C. B., Ballard, W. W., Kimmel, S. R., Ullmann, B. & Schilling, T. F. (1995). Stages of embryonic development of the zebrafish. *Dev Dyn,* 203**,** 253-310. 10.1002/aja.1002030302

Kucenas, S., Takada, N., Park, H.-C., Woodruff, E., Broadie, K. & Appel, B. (2008). CNS-
derived glia ensheath peripheral nerves and mediate motor root development.
Nature Neuroscience, 11**,** 143-151. 10.1038/nn2025

Latimer, A. J. & Appel, B. (2006). Notch signaling regulates midline cell specification and
proliferation in zebrafish. *Developmental Biology,* 298**,** 392-402.
https://doi.org/10.1016/j.ydbio.2006.05.039

Lee, R. T. H., Knapik, E. W., Thiery, J. P. & Carney, T. J. (2013). An exclusively
mesodermal origin of fin mesenchyme demonstrates that zebrafish trunk neural
crest does not generate ectomesenchyme. *Development,* 140**,** 2923-2932.
10.1242/dev.093534

Leonard, E. V., Figueroa, R. J., Bussmann, J., Lawson, N. D., Amigo, J. D. & Siekmann,
A. F. (2022). Regenerating vascular mural cells in zebrafish fin blood vessels are
not derived from pre-existing mural cells and differentially require Pdgfrb signalling
for their development. *Development,* 149. 10.1242/dev.199640

Li, L., Miano, J. M., Cserjesi, P. & Olson, E. N. (1996). SM22α, a Marker of Adult Smooth
Muscle, Is Expressed in Multiple Myogenic Lineages During Embryogenesis.
Circulation Research, 78**,** 188-195. doi:10.1161/01.RES.78.2.188

Löfberg, J. & Collazo, A. (1997). Hypochord, an enigmatic embryonic structure: Study of
the axolotl embryo. *Journal of Morphology,* 232**,** 57-66.
https://doi.org/10.1002/(SICI)1097-4687(199704)232:1<57::AID-
JMOR3>3.0.CO;2-L

Ma, R. C., Jacobs, C. T., Sharma, P., Kocha, K. M. & Huang, P. (2018). Stereotypic
generation of axial tenocytes from bipartite sclerotome domains in zebrafish. *PLoS
Genet,* 14**,** e1007775. 10.1371/journal.pgen.1007775

Majesky, M. W. (2007). Developmental Basis of Vascular Smooth Muscle Diversity.
Arteriosclerosis, Thrombosis, and Vascular Biology, 27**,** 1248-1258.
doi:10.1161/ATVBAHA.107.141069

Oonuma, K., Yamamoto, M., Moritsugu, N., Okawa, N., Mukai, M., Sotani, M., Tsunemi,
S., Sugimoto, H., Nakagome, E., Hasegawa, Y., Shimai, K., Horie, T. & Kusakabe,
T. G. (2021). Evolution of Developmental Programs for the Midline Structures in
Chordates: Insights From Gene Regulation in the Floor Plate and Hypochord
Homologues of Ciona Embryos. *Front Cell Dev Biol,* 9**,** 704367.
10.3389/fcell.2021.704367

Owens, G. K., Kumar, M. S. & Wamhoff, B. R. (2004). Molecular Regulation of Vascular
Smooth Muscle Cell Differentiation in Development and Disease. *Physiological
Reviews,* 84**,** 767-801. 10.1152/physrev.00041.2003

Peyrot, S. M., Wallingford, J. B. & Harland, R. M. (2011). A revised model of Xenopus
dorsal midline development: Differential and separable requirements for Notch and

709   Shh   signaling.   *Developmental*   *Biology,*   352**,**   254-266.
710   https://doi.org/10.1016/j.ydbio.2011.01.021

711 Poole, T. J., Finkelstein, E. B. & Cox, C. M. (2001). The role of FGF and VEGF in
712   angioblast induction and migration during vascular development. *Developmental*
713   *Dynamics,* 220**,** 1-17. https://doi.org/10.1002/1097-0177(2000)9999:9999<::AID-
714   DVDY1087>3.0.CO;2-2

715 Rajan, A. M., Ma, R. C., Kocha, K. M., Zhang, D. J. & Huang, P. (2020). Dual function of
716   perivascular fibroblasts in vascular stabilization in zebrafish. *PLOS Genetics,* 16**,**
717   e1008800. 10.1371/journal.pgen.1008800

718 Rensen, S. S., Doevendans, P. A. & Van Eys, G. J. (2007). Regulation and characteristics
719   of vascular smooth muscle cell phenotypic diversity. *Neth Heart J,* 15**,** 100-8.
720   10.1007/bf03085963

721 Row, R. H., Tsotras, S. R., Goto, H. & Martin, B. L. (2016). The zebrafish tailbud contains
722   two independent populations of midline progenitor cells that maintain long-term
723   germ layer plasticity and differentiate in response to local signaling cues.
724   *Development,* 143**,** 244-54. 10.1242/dev.129015

725 Santoro, M. M., Pesce, G. & Stainier, D. Y. (2009). Characterization of vascular mural
726   cells   during   zebrafish   development.   *Mech*   *Dev,*   126**,**   638-49.
727   10.1016/j.mod.2009.06.1080

728 Seiler, C., Abrams, J. & Pack, M. (2010). Characterization of zebrafish intestinal smooth
729   muscle development using a novel sm22α-b promoter. *Developmental Dynamics,*
730   239**,** 2806-2812. https://doi.org/10.1002/dvdy.22420

731 Shih, Y.-H., Portman, D., Idrizi, F., Grosse, A. & Lawson, N. D. (2021). Integrated
732   molecular analysis identifies a conserved pericyte gene signature in zebrafish.
733   *Development,* 148. 10.1242/dev.200189

734 Shkumatava, A., Fischer, S., MüLler, F., Strahle, U. & Neumann, C. J. (2004). Sonic
735   hedgehog, secreted by amacrine cells, acts as a short-range signal to direct
736   differentiation and lamination in the zebrafish retina. *Development,* 131**,** 3849-
737   3858. 10.1242/dev.01247

738 Sinclair, J. W., Hoying, D. R., Bresciani, E., Nogare, D. D., Needle, C. D., Berger, A., Wu,
739   W., Bishop, K., Elkahloun, A. G., Chitnis, A., Liu, P. & Burgess, S. M. (2021). The
740   Warburg effect is necessary to promote glycosylation in the blastema during
741   zebrafish tail regeneration. *NPJ Regen Med,* 6**,** 55. 10.1038/s41536-021-00163-x

742 Smith, C. L. & Tallquist, M. D. (2010). PDGF function in diverse neural crest cell
743   populations. *Cell Adh Migr,* 4**,** 561-6. 10.4161/cam.4.4.12829

Stickney, H. L., Barresi, M. J. F. & Devoto, S. H. (2000). Somite development in zebrafish.
Developmental Dynamics, 219**,** 287-303. https://doi.org/10.1002/1097-
0177(2000)9999:9999<::AID-DVDY1065>3.0.CO;2-A

Strähle, U., Lam, C. S., Ertzer, R. & Rastegar, S. (2004). Vertebrate floor-plate
specification: variations on common themes. *Trends in Genetics,* 20**,** 155-162.
https://doi.org/10.1016/j.tig.2004.01.002

Stratman, A. N., Burns, M. C., Farrelly, O. M., Davis, A. E., Li, W., Pham, V. N.,
Castranova, D., Yano, J. J., Goddard, L. M., Nguyen, O., Galanternik, M. V., Bolan,
T. J., Kahn, M. L., Mukouyama, Y. S. & Weinstein, B. M. (2020). Chemokine
mediated signalling within arteries promotes vascular smooth muscle cell
recruitment. *Commun Biol,* 3**,** 734. 10.1038/s42003-020-01462-7

Stratman, A. N., Pezoa, S. A., Farrelly, O. M., Castranova, D., Dye, L. E., Butler, M. G.,
Sidik, H., Talbot, W. S. & Weinstein, B. M. (2017). Interactions between mural cells
and endothelial cells stabilize the developing zebrafish dorsal aorta. *Development,*
144**,** 115-127. 10.1242/dev.143131

Tani, S., Chung, U.-I., Ohba, S. & Hojo, H. (2020). Understanding paraxial mesoderm
development and sclerotome specification for skeletal repair. *Experimental &*
Molecular Medicine, 52**,** 1166-1177. 10.1038/s12276-020-0482-1

Tsuji-Tamura, K., Morino-Koga, S., Suzuki, S. & Ogawa, M. (2021). The canonical smooth
muscle cell marker TAGLN is present in endothelial cells and is involved in
angiogenesis. *Journal of Cell Science,* 134. 10.1242/jcs.254920

Ulrich, F., Ma, L.-H., Baker, R. G. & Torres-Vázquez, J. (2011). Neurovascular
development in the embryonic zebrafish hindbrain. *Developmental Biology,* 357**,**
134-151. https://doi.org/10.1016/j.ydbio.2011.06.037

Wang, Y., Pan, L., Moens, C. B. & Appel, B. (2014). Notch3 establishes brain vascular
integrity by regulating pericyte number. *Development,* 141**,** 307-317.
10.1242/dev.096107

Wasteson, P., Johansson, B. R., Jukkola, T., Breuer, S., AkyüRek, L. M., Partanen, J. &
Lindahl, P. (2008). Developmental origin of smooth muscle cells in the descending
aorta in mice. *Development,* 135**,** 1823-1832. 10.1242/dev.020958

Westerfield, M. (2000). *The zebrafish book. A guide for the laboratory use of zebrafish*
(Danio rerio), Eugene, Univ. of Oregon Press.

Whitesell, T. R., Kennedy, R. M., Carter, A. D., Rollins, E. L., Georgijevic, S., Santoro, M.
M. & Childs, S. J. (2014). An α-smooth muscle actin (acta2/αsma) zebrafish
transgenic line marking vascular mural cells and visceral smooth muscle cells.
PLoS One, 9**,** e90590. 10.1371/journal.pone.0090590

Wiens, K. M., Lee, H. L., Shimada, H., Metcalf, A. E., Chao, M. Y. & Lien, C.-L. (2010).
Platelet-Derived Growth Factor Receptor β Is Critical for Zebrafish Intersegmental
Vessel Formation. *PLOS ONE,* 5**,** e11324. 10.1371/journal.pone.0011324

Wlizla, M. & Zorn, A. M. (2015). Chapter 13 - Vertebrate Endoderm Formation. *In:* Moody,
S. A. (ed.) *Principles of Developmental Genetics (Second Edition).* Oxford:
Academic Press.

Zheng, P., Xie, H., Zhu, P. & Zhao, C. (2022). Distribution and characteristics of neurons
at the nerve base plate of zebrafish. *Heredity,* 44**,** 510-520. 10.16288/j.yczz.22-
074

Zhou, H.-M., Fang, Y.-Y., Weinberger, P. M., Ding, L.-L., Cowell, J. K., Hudson, F. Z.,
Ren, M., Lee, J. R., Chen, Q.-K., Su, H., Dynan, W. S. & Lin, Y. (2016). Transgelin
increases metastatic potential of colorectal cancer cells in vivo and alters
expression of genes involved in cell motility. *BMC Cancer,* 16**,** 55.
10.1186/s12885-016-2105-8
794
795
796

 Figure Legends:

 Figure 1: Coronal views of *Tg(tagln:NLS-EGFP); Tg(pdgfrb:Gal4FF; UAS:RFP)*
 zebrafish brains from 1 to 5 dpf. (A) Schematic diagram illustrating the orientation for
 imaging brain coronal sections: the lower middle region (yellow) and upper middle region
 (purple). (B-J) Maximum intensity projection images of the mid- and hindbrain from 1 to 5
 dpf. (B'-J') Outlines of the blood vessels visualized in the confocal images; pink
 represents arteries and blue represents veins. At 1 dpf (B) *pdgfrb*-positive cells are
 associated with the PHBC. At 2 dpf (C,D) *pdgfrb*-positive cells begin to accumulate
 around the CoW (C, comprised of the BCA and PCS) and on the CtAs (D). *pdgfrb*-positive
 cells continue to accumulate on the CoW and CtAs from 3 to 5 dpf (E-J). A single
 pdgfrb/tagln double-positive cell is evident on the MMCtA at 4 dpf (H, inset). Schematic
 created with BioRender.com. AMCtA, anterior mesencephalic central artery. BA, basilar
 artery. BCA, basal communicating artery. CaDI, caudal division of the internal carotid
 artery. CCtA, cerebellar central artery. CoW, Circle of Willis. MMCtA, middle
 mesencephalic central artery. MtA, metencephalic artery. PCS, posterior communicating
 segment. PHBC, primordial hindbrain channel. PMCtA, posterior mesencephalic central
 artery. Scale bars: 50 μm.

 Figure 2: Summary of mural cell accumulation on the axial vasculature. The
 hypochord—which begins as a *pdgfrb/tagln* double-positive structure—is a novel source
 of *tagln*-positive cells on the dorsal side of the dorsal aorta and caudal artery. At 2 dpf,
 the hypochord cells began to delaminate on the dorsal aorta. At 3 dpf, the *pdgfrb/tagln*
 double-positive cells begin to decrease *pdgfrb* expression. By 4 dpf, these cells were *tagln*
 single-positive. In the caudal region, this hypochord reprogramming event began 1 day
 later than the anterior region. On the ventral side of the dorsal aorta and caudal vein,
 pdgfrb single-positive vSMCs were observed beginning at 2 dpf, where they then
 expanded in number and increased expression of *tagln*. By 5 dpf, *pdgfrb/tagln* double-
 positive cells accounted for approximately 60% of ventral vSMCs on the dorsal aorta, and
 no *tagln* single-positive cells were detected. The vSMCs on the ventral side of the CA
 accumulated more slowly and did not begin co-expressing *tagln* until 5 dpf, which

amounted to a 2-day delay compared to the anterior region. Finally, we observed a slow

accumulation of *pdgfrb* single-positive cells on the posterior cardinal vein and caudal vein

between 1 to 5 dpf. *tagln*-positive cells were particularly sparse on the veins, but visible

at later stages.

Figure 3: *tagln* and *pdgfrb* expression along the dorsal aorta and cardinal vein of 1

to 5 dpf *Tg(tagln:NLS-EGFP); Tg(pdgfrb:Gal4FF; UAS:RFP)* zebrafish. (A-E)

Maximum intensity projection images of the DA and PCV from 1 to 5 dpf. (A'-E') The

images from A-E without the brightfield overlay. (F-G) Bar and line graph representations

of the number of single- and double-positive cells on the dorsal side of the DA (F), the

ventral side of the DA (G), and the PCV (H). (I) Schematic diagram illustrating the imaged

region. Schematic created with BioRender.com. At least 6 embryos were analyzed at

each age. DA, dorsal aorta. PCV, posterior cardinal vein. HC, hypochord. NC, notochord.

Scale bars: 50 µm.

Figure 4: *tagln* and *pdgfrb* expression along the caudal artery and vein of 1 to 5 dpf

***Tg(tagln:NLS-EGFP); Tg(pdgfrb:Gal4FF; UAS:RFP)* zebrafish.** (A-E) Maximum

intensity projection images of the CA and CV from 1 to 5 dpf. (A'-E') The images from A-

E without the brightfield overlay. (F-G) Bar and line graph representations of the number

of single- and double-positive cells on the dorsal side of the CA (F), the ventral side of the

CA (G), and the CV (H). (I) Schematic diagram illustrating the imaged region. Schematic

created with BioRender.com. At least 6 embryos were analyzed at each age. CA, caudal

artery. CV, caudal vein. HC, hypochord. NC, notochord. Scale bars: 50 µm.

Figure 5. Low and high *pdgfrb/tagln* expression is evident along arterial and

venous ISVs from 2 to 5 dpf. (A,B) Maximum intensity projection images of arterial and

venous ISVs at 5 dpf. ISVs are outlined by white dotted lines. *tagln*-positive skeletal

muscle nuclei are particularly evident in the trunk region near the ISVs, thus requiring

frame-by-frame analysis to identify *tagln*-positive perivascular cells. (C,D) Bar graph

representations of the number of low and high single- and double-positive cells on the

arterial (C) and venous (D) ISVs. Arterial ISVs accumulate *pdgfrb*-high mural cells more

robustly than venous ISVs. (E) Representative images of the combinations of *pdgfrb*/*tagln*

double-positive cells. (F,G) Bar graph representations of the number double-positive cells

on the arterial (F) and venous (G) ISVs. *pdgfrb*/*tagln* double-positive cells were more

apparent on arterial ISVs. (H) Schematic diagram summarizing the general localization

of perivascular cells on the ISVs at 2 and 5 dpf. *pdgfrb* single-positive cells were more

likely to be found on the dorsal ISV region, *tagln* single-positive cells primarily near the

notochord ISV region, and double-positive cells nearest to the notochord-neural tube

interface by 5 dpf. Schematic created with BioRender.com. At least 6 embryos were

analyzed at each age. ISVs, intersegmental vessels. Scale bars for A,B: 50 μm; scale

bars for E: 5 μm.

Figure 6: Summary of the midline cell populations at the notochord-neural tube

interface. We identify the medial floor plate cells as *pdgfrb*/*tagln* double-positive in early

zebrafish development (MFP). Additionally, we observe a *tagln* single-positive population

of cells along the midline that accumulates below the medial floor plate cells beginning at

3 dpf (*tagln*+ Midline Cells). Finally, we describe a mural cell progenitor population near

the notochord-neural tube interface that borders either side of the midline. This

population—which begins as *tagln* single-positive cells before acquiring *pdgfrb*

expression—can be observed migrating to the nearby ISVs from 2 to 5 dpf (MC Progenitor

Population). Schematic created with BioRender.com. NC, notochord. NT, neural tube.

ISV, intersegmental vessel. MFP, medial floor plate. MC, mural cell.

Figure 7. A *tagln*-positive sclerotome-derived mural cell progenitor population at

the notochord-neural tube interface upregulates *pdgfrb* and migrates to the ISVs.

(A-J) Maximum intensity projection images of cell populations near the notochord-neural

tube interface. Sagittal (A,C,E,G,I) and orthogonal (B,D,F,H,J) views of the medial floor

plate from 1-5 dpf reveal *pdgfrb*/*tagln* co-expression (cyan arrows). A separate *tagln*

single-positive population of cells accumulates immediately below the MFP starting at 3

dpf (E-J, white outlined arrows). Finally, an independent *tagln* single-positive population—

hereafter the mural cell progenitor population—aligns near the notochord-neural tube

interface (A'-I',B,D,F,H,J, yellow arrows) and begins to elongate and upregulate *pdgfrb*

over time. (K) Schematic diagram illustrating the three described cell populations from a

sagittal and orthogonal view. (L-O) The *tagln*-positive midline cells (white outlined arrows)

and mural cell progenitors (yellow arrows) are *twist1a*-positive, indicating they are derived

from the sclerotome, unlike the medial floor plate (pink arrows). (P-W) Cells from the

mural cell progenitor population can be seen migrating to the ISVs using *pdgfrb* and *tagln*

as markers (P-S) or *twist1a* and *tagln* (T-W). ISVs are outlined. White arrows indicate

migrating cells. Schematic created with BioRender.com. NC, notochord. NT, neural tube.

ISV, intersegmental vessel. MFP, medial floor plate. MC, mural cell. Scale bars: 10 μm.

Figure 1

Figure 2

pdgfrb   tagln

Figure 3

I

dorsal

anterior ←→ posterior

ventral

Figure 4

I

dorsal

anterior ←→ posterior

ventral

Figure 5

Tg(tagln:NLS-EGFP); Tg(pdgfrb:Gal4FF;UAS:RFP)

Arterial ISVs

A | 5 dpf | Merge | *pdgfrb* | *tagln*

Venous ISVs

B | Merge | *pdgfrb* | *tagln*

E | *pdgfrb+/tagln+* Double-Positive Cells | *pdgfrb^high^/tagln^high^* | *pdgfrb^high^/tagln^low^* | *pdgfrb^low^/tagln^high^* | *pdgfrb^low^/tagln^low^*

Total Arterial ISV Cells

- *pdgfrb^low^*
- *pdgfrb^high^*
- *pdgfrb+/tagln+*
- *tagln^low^*
- *tagln^high^*

Total Venous ISV Cells

Double-Positive Arterial ISV Cells

- *pdgfrb^high^/tagln^high^*
- *pdgfrb^high^/tagln^low^*

Double-Positive Venous ISV Cells

- *pdgfrb^low^/tagln^high^*
- *pdgfrb^low^/tagln^low^*

C # of cells per ISV — 15, 0

D # of cells per ISV — 15, 10, 5, 0 — dpf: 2 3 4 5

F # of cells per ISV — 6, 4, 2, 0 — dpf: 2 3 4 5

G # of cells per ISV — 6, 4, 2, 0 — dpf: 2 3 4 5

H

Arterial ISV — Venous ISV — Arterial ISV — Venous ISV

Dorsal — Ventral

tagln-high
tagln-low
pdgfrb-high
pdgfrb-low

Notochord (NC)
Neural Tube (NT)
Intersegmental Vessel (ISV)

2 dpf — **5 dpf**

Figure 6

**Midline Cell Populations at the
Notochord-Neural Tube Interface**

1 dpf

2 dpf

3 dpf

4 dpf

Dorsal

Ventral

Migration to ISVs

NC   NT   ISV   MFP   *tagln+* Midline Cells   MC Progenitor Population

tagln   *pdgfrb*

Figure 7