p53 translational-isoform Δ40p53 regulates cell cycle by modulating the miR-4671-5p/SGSH axis

Apala Pal[1], Pritam Kumar Ghosh[1], Sahana Ghosh[2], Sachin Kumar Tripathi[1], Sohini Guha[3], Pragnya Coca[4], Subrata Patra[2], K.M. Prathima[5], Debjit Khan[1,6], Manjula Das[3,7], Arindam Maitra[2] and Saumitra Das[1, *]

[1]Department of Microbiology and Cell Biology, Indian Institute of Science, Bangalore-560012, India.

[2]National Institute of Biomedical Genomics, Kalyani-741251, India.

[3]Molecular Immunology, Mazumdar Shaw Medical Foundation, Narayana Health City, Bommasandra, Bangalore-560099, India.

[4]Mazumdar Shaw Medical Center, Narayana Health City, Bommasandra, Bangalore-560099, India.

[5]Manipal Hospital, Miller's Road, Bangalore-560052, India.

[6]Department of Life Sciences, GITAM School of Science, GITAM (Deemed to be University), Bangalore-562163, India.

[7]Center for Precision Diagnostic and Therapeutic Research, India (CPDTRI), Electronic city, Bangalore-560100, India

*Address for correspondence:

Dr. Saumitra Das, Professor,

Department of Microbiology and Cell Biology,

Indian Institute of Science, Bangalore-560012, India

Email: sdas@iisc.ac.in

Phone: 91 80 2293 2886

Fax: 91 80 2360 2697

ABSTRACT

Parsing functions of tumour suppressor gene *TP53* is inherently complicated owing to multiple p53 isoforms it encodes. Δ40p53, the only translational isoform, alters p53 target gene expression both positively and negatively, thereby modulating p53 functional activity. Independent of p53, Δ40p53 regulates miR-186-5p–YY1 axis to decrease cell proliferation. To uncover more such miRNAs that are differentially regulated by Δ40p53 and p53, in this study, we performed small RNA sequencing. Overexpression of Δ40p53, but not full-length (FL) p53, substantially downregulated expression of miR-4671-5p. However, upon overexpression of different ratios of Δ40p53 and FLp53, changes in miR-4671-5p levels implied that it is a FLp53 modifiable target. In an *in-silico* approach, we identified N-sulfoglucosamine sulfohydrolase (*SGSH*) as a candidate miR-4671-5p target. The expression levels of SGSH show potential prognostic significance for survival that trends oppositely to miR-4671-5p levels associated with the same cancer types, which supports a direct physiological relevance of their interaction. Furthermore, decreased *SGSH* mRNA and protein levels were observed with siΔ40p53 as well as upon overexpression of miR-4671-5p, confirming the molecular interplay. Notably, overexpression of miR-4671-5p led to intra-S-phase cell cycle arrest. Therefore, the Δ40p53-miR-4671-5p-SGSH axis emerges as a novel regulatory pathway influencing cell cycle progression. We also expanded our study to a cohort of colorectal cancer patient samples. Results indicate that the expression level of Δ40p53 can stratify miR-4671-5p and *SGSH* mRNA levels, validating their inverse correlation observed in the cellular scenario. Collectively, these findings significantly enhance our understanding of the functions of Δ40p53 mediated by miRNAs, contributing to the maintenance of metabolic and cellular homeostasis independent of full-length p53.

Keywords: p53 isoform, Δ40p53 role, miR-4671-5p expression, SGSH role in cell cycle regulation

INTRODUCTION

Cellular stress responses are multifaceted and involve damage recognition and repair, including that of DNA, to minimize the risk of genetic instability [1]. p53 binds DNA and plays a central role in protecting genomic integrity. Under genomic stress, p53 causes cell cycle arrest and hinders the proliferation of cells with damaged DNA [2]. The functional activities of p53 are lost in multiple human cancers due to protein inactivation or gene mutations [3]. The p53 pathway is regulated by several factors, among which the role of p53 isoforms is crucial because these can alter transcriptional activation of p53 targets and thus affect its functions [4]. Twelve p53 isoforms are produced via various mechanisms to bear N-terminal (FL/Δ40/Δ133/Δ160) and C-terminal (α/β/γ) diversity [5], and each is differentially expressed in normal and tumor tissues in various types of cancer [6, 7]. Among these, the only translational isoform of p53 is Δ40p53, and it critically sculpts p53 functions [8]. We previously identified two internal ribosome entry sites (IRESs) within p53 mRNA, where one IRES generates the full-length p53 (FLp53) protein, whereas initiation at the other generates Δ40p53 [9]. We have also shown that several translation initiation factors are critical for the differential regulation of IRES activity under various types of stress [10, 11].

FLp53 has a strong affinity for, and can form homo-tetramers and hetero-tetramers with Δ40p53, because they share oligomerization domains [12]. Furthermore, Δ40p53 can regulate the functional activity of p53, which directly affects cellular processes [13], as well as critical cellular processes, such as the cell cycle, proliferation, and senescence. The IRESs driving p53 isoforms have different cell cycle phase-dependent activities, with IRES for FLp53 being active at the G2–M transition and IRES for Δ40p53 being the most active at the G1–S transition [9]. In addition to regulating p53 activity, Δ40p53 retains the second transcriptional activation domain in FLp53 and induces the expression of various genes independently of FLp53,—[14, 15, 16]. Δ40p53 also induces 14-3-3σ and G2 arrest but does not affect G1 progression [17, 18]. It transactivates Bcl-2 Associated X-protein and Growth Arrest and DNA Damage-inducible 45 in p53-null cells [8]. The Nanog-IGF1 axis is also modulated by Δ40p53/FLp53 hetero-oligomers, thus controlling the switch from pluripotency to differentiation [19]. The proliferation of pancreatic β cells and glucose homeostasis is controlled by Δ40p53 in mice [20]. At basal levels, Δ40p53 and p53 similarly suppress cellular mobility and proliferation in breast cancer [21]. We found that Δ40p53 (and not FLp53) uniquely regulates miR-186-5p, which can downregulate the expression of the known oncogene Ying

Yang 1 (YY1) protein, thereby decreasing cell proliferation [22]. In another study, we have also

found that Δ40p53 regulates lncRNA (LINC00176), which can control cell fate mediated

through multiple miRNAs and mRNAs [23]. Contrary to its tumor suppressive role Δ40p53

independently activates the netrin-1 promoter and promotes cell survival [24].

This study aimed to globally identify those miRNAs that are differentially regulated by

Δ40p53(α) and p53(α). We assessed changes in miRNA expression in lung and colon cancer

cell lines cells expressing either or both FLp53(α) and Δ40p53(α) to identify such differential

miRNA regulation.

MATERIALS AND METHODS

Cell lines and transfections

We have used three cell lines in the current study: H1299 (lung adenocarcinoma cell line

lacking p53 or Δ40p53), HCT116 p53+/+ cells (colon carcinoma cell line harboring wild-type

p53; hereafter called HCT116+/+); and HCT116 p53-/- cells (colon carcinoma cell line

harboring only Δ40p53; hereafter called HCT116-/-). These cells were maintained in DMEM

(Sigma) with 10% Foetal Bovine Serum (GIBCO, Invitrogen). 70%-80% confluent

monolayer of cells was transfected with various plasmid constructs using Lipofectamine 2000

(Invitrogen) and Turbofectamine in Opti-MEM (Invitrogen). The medium was replaced with

DMEM (with antibiotic) and 10% FBS four hours later. The cells were harvested and

processed as required at the desired time point.

Small RNA Sequencing

RNA Extraction: Total RNA was isolated from control (only vector), p53, Δ40p53, 14A (p53

and Δ40p53) transfected H1299 cells using TRIzol and PureLink RNA mini kit (Ambion)

according to the manufacturer's instructions. Quality of isolated total RNA was checked

using Agilent RNA 6000 Nano chips in 2100 Bioanalyzer (Agilent) and NanoDrop

spectrophotometer (Thermo Scientific) and quantitation was performed in Qubit using Quant-

iT RNA assay kit broad range (Thermo Scientific). Total RNA samples with RNA integrity

number (RIN) >7 was selected for Illumina small RNA sequencing library preparation.

Library Preparation: Small RNA sequencing library preparation was performed using

Illumina® TruSeq® Small RNA Library Prep Kit according to the manufacturer's protocol.

One microgram of the total RNA sample was used for library preparation, and adapters were ligated to 3' and 5'ends of each RNA molecule. Subsequent reverse transcription and amplification were performed to generate a cDNA library, which was further gel purified to select bands between 145-160 bp to prepare a final small RNA sequencing library for clustering and sequencing. Single read 1 x 50 bp sequencing of these libraries was performed in HiSeq-2500 (Illumina).

RNA-seq data analysis: FASTQ files generated for each sample were checked for quality, and then adapter trimming was performed using the cutadapt2 tool (v1.8.1). Adapter trimmed FASTQ files were aligned to the hg19 reference sequence using the mapper module of the miRDeep2 software package. The accepted aligned FASTA files were processed using miRDeep2 module where known mature, precursor miRNAs from humans and mature miRNAs from other species were provided (downloaded from miRBase database) as input. It identified known and novel miRNAs expressed. miRDeep2 module quantifies the expression of the known miRNAs and gives normalized read counts (reads per million: RPM) for each known precursor miRNA. Average RPM values were calculated where a single mature miRNA had multiple precursor miRNAs. DESeq2 was used to identify differentially expressed miRNAs.

Plasmids and constructs

pGFP-hp-p53- 5'UTR-cDNA (14A): It expresses both FLp53 and Δ40p53 (a generous gift from Dr. Robin Fahraeus, INSERM, France). pGFP-hp-p53-5'UTR (A135T)-cDNA which expresses only Δ40p53 and pGFP-hp-p53- 5'UTR (A251G/T252 C/G253T)-cDNA which expresses only full-length p53 (p53FL) [22]. miR-4671-5p overexpression construct cloning was done by annealing double-stranded oligos of a mature miR-4671-5p, along with the hairpin sequence in the middle between BglII and XhoI in pSUPER vector.

siRNA transfections

HCT116+/+ and HCT116-/- cells were transfected with 30 nM si p53 RNA (IDT). 5'-AACCUCUUGGUGAACCUUAGUACCU-3' is the sequence for p53 siRNA, directed against the 3'UTR of p53; therefore, it targets both p53 as well as Δ40p53. In the experiments, a non-specific siRNA (Dharmacon) was used as a control.

Western Blot analysis

Protein concentrations of the extracts were assayed by Bradford (Bio-Rad), and equal amounts of cell extracts were separated by SDS 12% PAGE and transferred to nitrocellulose membrane (Bio-Rad). Samples were then analyzed by Western blotting using a rabbit-raised anti-p53 polyclonal antibody (CM1, a kind gift from Dr. Robin Fahraeus, INSERM, France and Prof. J.C. Bourdon, of University of Dundee, UK), followed by secondary antibody (horseradish peroxidase-conjugated anti-mouse or anti-rabbit IgG; Sigma). SGSH protein has been detected using anti-SGSH Antibody (Abclonal, Cat. No. A8148). CDK11B protein has been detected using anti-CDK11B Antibody (Abclonal Cat. No. A12830). CDK5R1 protein has been detected using anti-CDK5R1 Antibody (Abclonal Cat. No. A14497). Densitometric analysis was done on Fiji (https://fiji.sc/) by calculating band intensity and deducting the averaged background value from equal areas at the top and bottom of the band.

RNA isolation and Real-time PCR

According to the manufacturer's protocol, total RNA was isolated from cells with TRI Reagent TM (Sigma). The isolated RNA was treated with 10 units of DNase I (Promega), extracted with acidic phenol-chloroform, and precipitated with 3 M sodium acetate (pH 5.2) and 2.5 volumes of absolute alcohol. RNA amount was quantified with Nano-spectrophotometer, and cDNAs were synthesized using specific reverse primers and MMLV RT (Revertaid ^{TM} Thermo Scientific) at 42°C for 1 h, according to standard protocol. At the same time, 2-5µg of total RNA has been used to check mRNAs' expression. 50ng of total RNA has been used to check the expression of miRNAs.

SYBR green Assay System was used for mRNA, lncRNA and miRNA detection and quantification. We have used actin as an endogenous control for mRNA and 5SrRNA as an endogenous control for miRNAs. The thermocycling conditions for SYBR green Assay system include 1 cycle of 95°C for 10 min, 40 cycles of 95°C for 15 s, 60°C for 30 s and 72°C for 30 s (Applied Biosystems). $2^{-\Delta\Delta Ct}$ method algorithm was used to analyze the relative changes in expressions, where actin/5S served as an endogenous control. The fold change was calculated using $2^{-\Delta\Delta Ct}$. $\Delta Ct = Ct$ (target gene) $-Ct$ (endogenous control) and $\Delta\Delta Ct = \Delta Ct$ target sample) $-\Delta Ct$ (control sample). For patient samples, normalized m gciR-4671 (ΔCt) = Ct (miR-4671) $-Ct$ (U6 snRNA) and normalized mRNA of (ΔCt) = Ct (mRNA) $-Ct$ (actin). Melting curve analysis of every (q)PCR was conducted after each cycle.

Cell cycle Analysis

HCT116-/- cells were transfected with miR-4671-5p or miR-34a-5p overexpression construct 48 h post-transfection; the cells were harvested and fixed with methanol. The fixed cells were treated with RNase A (10 mg/ml) for 1 h at 37°C, following which PI (Propidium iodide-1mg/ml) was added to the cells and taken for Flow cytometric analysis.

Bioinformatic Analysis

dbDEMC2.0 (https://www.biosino.org/dbDEMC/index) has been used to perform a meta-profiling of selected miRNAs [25]. TargetScan 7.2 (https://www.targetscan.org/vert_72/) has been used to predict target mRNAs of miRNAs [26]. miRDB has been used to predict target miRNAs (https://mirdb.org/cgi-bin/search.cgi) [27, 28]. GO-term analysis was done on PANTHER release 17.0 (http://www.pantherdb.org/). Kaplan-Meier survival analysis was done on R2: Genomics Analysis and Visualization Platform (http://r2.amc.nl). Heat map for Figure 1E has been generated using Heatmapper (http://heatmapper.ca/).

Collection, processing, and analysis of colorectal carcinoma samples

Sample collection: Tumor samples, with informed consent [approval no. NHH/AEC-CL-2020-458 of ethics committee of Narayana HealthBangalore, India] from patients were cut into 3-5 small sections weighing ~30 mg and snap frozen in liquid nitrogen for storage The clinical and pathological details of the patients are mentioned in **Figure S1, Table 1**.

RNA isolation: RNA was isolated from fresh frozen tissue samples using TRIzol (Ambion, #15596018) as per manufacturer's instructions. Briefly, ~30 mg of the fresh frozen tissue samples were minced and dissolved in 1 ml TRIzol followed by addition of 200 µl of chloroform (Sigma, #496189) and centrifugation at 13,000 g for 15 minutes at 4°C. The aqueous layer was transferred to a fresh tube containing an equal volume of 100% isopropanol (Qualigen, #Q26897) without disturbing either the middle interface or the lower organic phase. The tube was gently invert-mixed and incubated at room temperature for 10 minutes followed by centrifugation at 13,000 g for 10 minutes at 4°C. The supernatant was discarded, and the pellet was washed with 75% ethanol (HiMedia, #MB228) followed by centrifugation at 7500 g for 5 minutes at 4°C. The pellet was air dried and resuspended in nuclease free water (Ambion, AM9937) for further processing.

Protein isolation and Quantification: Protein was isolated from a fresh frozen tissue piece of

~30 mg using NucleoSpin RNA/Protein purification kit (Macharey-Nagel, #740933.5), as per

manufacturer's instructions. Quantification was done using the Pierce BCA protein assay kit

(ThermoScientific #23227) as per manufacturer's instructions. Absorbance was measured at

562 nm.

Histopathology analysis: Tumor tissues fixed in 10% (v/v) formalin (Sigma, #HT501128)

and embedded in paraffin (Qualigen, #Q19215) were used in the study. Sections of 5 µm

thickness were stained with hematoxylin and eosin. Slides were analysed for the extent of

tumor boundary and regions of necrosis.

Statistical analysis

213 The data for the experimental section are expressed as mean ± SD. Statistical significance for

214 the experiments was determined using two-tailed Student's t-test. The criterion for statistical

215 significance was $p \leq 0.05$ (*) or $p \leq 0.01$ (**) or $p \leq 0.001$(***)

RESULTS

Small RNA sequencing reveals miRNA targets of p53 and Δ40p53

H1299 cells (that do not endogenously express any p53 isoforms) were transfected with

vectors (in triplicate) to ensure that they expressed only FLp53, only Δ40p53, both p53 and

Δ40p53 (vector 14A), or GFP (Control, C) **(Figure 1A-C)**. We then investigated the effect of

differential expression of p53 isoforms on miRNA gene expression using RNA isolated from

these cells and processed for quality control and sequencing of small RNAs. A heatmap was

generated for those miRNAs with unidirectional fold changes for at least one condition of

p53 isoform overexpression, in all triplicates, compared to control **(Figure 1D)**. Among 135

miRNAs, we selected a significantly expressed cluster which is differentially regulated under

at least one condition. ($p \leq 0.05$; **Figure 1E**).

We primarily aimed to identify miRNAs with an expression that was differentially regulated

by p53 and Δ40p53 and thus selected miR-4671-5p, miR-301b-5p, miR-34a-5p, and miR-

548ae-5p, with fold changes ranging between ≥ 0.58 and < -1.0 ($p \leq 0.05$). The expression of

miR-4671-5p was significantly downregulated and was unchanged on the Δ40p53 and p53

backgrounds, respectively. The expression miR-301b-5p was significantly downregulated and

unchanged on the p53 and Δ40p53 backgrounds, respectively. We also analysed the novel miR-548-ae-5p, the expression of which was significantly upregulated by p53 and the other two conditions. We then investigated the mechanism of co-regulation by the isoforms (if any) on miR-548ae-5p, and the positive control was miR-34a-5p that is regulated by p53 [29].

Selection and validation of miRNAs for further studies

Control (vector only), p53, Δ40p53, and 14A (a combination of both isoforms) were transfected in H1299 cells **(Figure 2A)** and the miRNA levels were examined using qRT-PCR. The expression of miR-4671-5p decreased when Δ40p53 and 14A were overexpressed and remained unchanged upon p53 overexpression compared with controls **(Figure 2B)**. The expression of miR-34a-5p and miR-548ae-5p increased when p53, Δ40p53, and 14A were overexpressed **(Figure 2C and D)**. The expression of miR-301b-5p was not altered by overexpressed p53, whereas overexpressed Δ40p53 and 14A increased it **(Figure 2E)**. Except for miR-301b-5p, the expression of miR-4671-5p, miR-548ae-5p, and miR-34a-5p was similar to that obtained using the sequencing platform **(Figure 2F)**. Therefore, we continued our investigation with these three miRNAs.

Regulation of miRNAs by Δ40p53 and p53

Colon cancer cell lines HCT116+/+ (predominantly expressing p53 and some Δ40p53) and HCT116-/- (expressing only 40p53) were used to determine how p53 isoforms regulate the expression of these miRNAs and to render our study more physiologically relevant **(Figure 2G)**. We initially found less endogenous expression of miR-4671-5p and more miR-548ae-5p and miR-34a-5p in HCT116-/-, than HCT116+/+ cells **(Figure 2H)**. Furthermore, p53 or Δ40p53 was partially silenced by siRNA-mediated knockdown in HCT116+/+ **(Figure 2J)** and HCT116-/- **(Figure 2L)** cells and miRNA levels were detected using qRT-PCR. The expression of miR-548ae-5p and miR-34a-5p significantly decreased upon silencing p53 in HCT116+/+ cells, whereas that of miR-4671-5p did not significantly change **(Figure 2I)**. Silencing Δ40p53 increased miR-4671-5p and decreased miR-548ae-5p and miR-34a-5p expression in HCT116-/-cells. These data suggested that Δ40p53 regulates miR-4671-5p negatively and miR-548ae-5p and miR-34a-5p positively **(Figure 2K)**. Importantly, Δ40p53 exclusively altered miR-4671-5p expression, with no effect by FLp53. These results also indicated that changing relative p53 and Δ40p53 levels differentially affects miRNA expression. The abundance of p53 isoforms notably differs among various types of cancer [6]. Therefore, target miRNAs should vary accordingly.

Bioinformatic analysis of miR-4671-5p, miR-548ae-5p, and miR-34a-5p expression in

various types of cancer using the database of differentially expressed miRNAs (dbDEMC)

2.0 revealed that miR-34a-5p is expressed in many types of cancer, whereas miR-4671-5p is

expressed only in colon, lung, and pancreatic cancers. In contrast, miR-548ae-5p was not

detected in any cancer databases **(Figure 3A)**. The results indicated that the varying levels of

miRNAs among types of cancer could be attributed to the varying abundance of p53 and

Δ40p53. To confirm this, we overexpressed p53 or Δ40p53 at different ratios in H1299 cells

and measured the expression of these miRNAs **(Figure 3B)**. We found that miRNA

expression significantly changed according to the abundance of p53 or Δ40p53. The

expression of miR-34a-5p was upregulated by either overexpressed p53 or Δ40p53 alone.

However, varying ratios of both p53 and Δ40p53 isoforms reduced miR-34a-5p expression,

while equal amounts did not result in much change. **(Figure 3C)**. These results agree with

previous findings suggesting that p53 and Δ40p53 co-regulate their activities, thus leading to

the modulation of target gene expression [12].

Furthermore, overexpressed FLp53 did not significantly change miR-4671-5p expression,

which was (similar to our findings in **Figure 2B**). However, increasing Δ40p53 levels in the

absence or presence of FLp53 consistently decreased miRNA expression **(Figure 3D)**. The

combined effects of these isoforms on the expression miR-4671-5p and 14A were similar.

Thus, in the context of miR-4671-5p, p53 might act as a regulator of the Δ40p53-mediated

gene expression. However, p53 alone did not affect miR-4671-5p.

Cellular role of miR-4671-5p

Micro RNAs function by base pairing with their target mRNAs and negatively regulating

their expression. The miR-34a family is regulated by p53 and miR-34a-5p is involved in

multiple cellular processes[29]. However, miR-4671-5p is novel, and its regulation has

remained unknown. Therefore, we investigated miR-4671-5p and elucidated its downstream

functions using the online databases Human TargetScan 7.2 and miRDB to identify potential

mRNA targets of miR-4671-5p. These two databases respectively provided 780 and 46

targets **(Figure S1, Table 2)**. 40/46 targets from miRDB targets were found in the

TargetScan database and 6/46 were absent. Next, we performed GO term analysis by

PANTHER statistical overrepresentation test using the curated 786 unique-target gene. GO

molecular function "protein serine kinase activity" was significantly enriched in the predicted

target list of miR-4671-5p that includes multiple kinases **(Figure S1, Table 3)**.

CDK11B (Cyclin Dependent Kinase 11B) and CDK5R1 (Cyclin Dependent Kinase 5
Regulatory Subunit 1) were selected as potential miR-4671-5p targets **(Figure 3E)**.
Importantly, PANTHER analysis showed an overrepresentation of serine kinases as predicted
targets of miR-4671-5p, highlighting the relevance of selecting these candidates **(Figure S1,**
Table 3). Further, we also shortlisted SGSH (N-Sulfoglucosamine Sulfohydrolase) as it is the
topmost target for miR-4671-5p in Targetscan. Mutations in SGSH cause a lysosomal sulfite
enzyme deficiency that leads to glycosaminoglycan (GAG) and heparan sulfate accumulation
in various organs of the body, ultimately causing mucopolysaccharidosis III A (MPS)
syndrome that is characterized by sever neurological symptoms and skeletal deformities [30].
While an association of SGSH to monkeypox and interstitial lung disease has been predicted
[31, 32] a role for SGSH in cancer is yet unknown. An analysis of existing primary tumor
datasets for SGSH gene expression revealed that low *SGSH* is significantly correlated with
poor survival in pancreatic adenocarcinoma **(Figure 3 F, G and Figure S1-Table 4)** in 3/4
datasets, while high *SGSH* is significantly correlated with poor survival in multiple colon
tumor datasets **(Figure 3 H, I)**. Importantly, in these tumor types, prognostic *SGSH* levels
trend opposite to miR-4671-5p as revealed in dbDEMC2.0 analysis **(Figure 3A)** suggestive
of a link between the miRNA and mRNA. Thus, SGSH, CDK11B and CDK5R1 were
selected for further studies.

We validated miR-4671-5p target levels under the same experimental conditions as in Figure
3B, with different ratios of overexpressed p53/Δ40p53 in H1299 cells. The results showed
increased SGSH, CDK11B, and CDK5R1 levels only under Δ40p53 overexpression **(Figure**
3 J-L). Moreover, we also checked for the protein levels of these targets in HCT116-/- (with
endogenous expression of Δ40p53), we observe that the abundance of SGSH, CDK11B and
CDK5R1 was very high compared to HCT116+/+ **(Figure 3M)**. This observation was in line
with Figure 3A, where miR-4671-5p levels were lower in HCT116-/- than in HCT116+/+.
This contrasting regulation of miR-4671-5p and its targets mediated by the presence of
Δ40p53 confirmed the involvement of these molecules in a single pathway.To confirm the
regulation of target mRNAs by Δ40p53, we partially silenced Δ40p53 in HCT116-/- and
miR-4671-5p target mRNA expression mediated by siRNA. The results showed that SGSH,
CDK11B and CDK5R1 mRNA expression significantly decreased under silenced Δ40p53
(Figure 4 A-B). These findings confirmed that Δ40p53 positively regulates SGSH, CDK11B,
and CDK5R1 mRNA expression. The Δ40p53-mediated positive regulation of mRNAs was
directed via downregulated miR-4671-5p expression **(Figure 2 B)**. We overexpressed miR-

4671-5p in HCT116 -/- cells to determine the direct effects of miRNAs on their targets. The

expression of SGSH, CDK11B and CDK5R1 mRNAs was decreased **(Figure 4 C-D)** and the

effect was the most significant on SGSH mRNA.

As mentioned earlier **(Figure 3 F-I)**, survival-correlated SGSH levels trend opposite to the

levels of miR-4671-5p in certain cancer types, which suggests a direct possible link between

the miRNA and the mRNA. However, the functional effect of mRNA is mainly mediated

through its protein expression, therefore we checked the effect of siΔ40p53 or overexpression

of miR-4671-5p on the respective protein levels. In both conditions, we observed a

substantial and consistent reduction in SGSH protein levels reflecting the mRNA levels

(**Figure 4 E, F**). This result suggested that SGSH mRNA could be a significant target

downstream to Δ40p53-miR-4671-5p pathway. Finally, it was essential to decipher the direct

role of miR-4671-5p on any cellular process. Since most of the selected targets of miR-4671-

5p were directed toward cell cycle regulation, we checked for the changes in cell cycle phases

upon miR-4671-5p overexpression. We observed that the percentage of cells in the S phase

was significantly increased, whereas the number of cells in the G2 phase decreased **(Figure

4G),** indicating the activation of a potential S-phase checkpoint which accumulated the cells

in the S phase and prevented them from proceeding to G2. Thus, miR-4671-5p can regulate

the cell cycle and possibly other associated cellular processes.

Δ40p53 expression in colorectal cancer stratifies miR-4671-5p and *SGSH* levels

To extend our findings in colon cancer cell lines and to ascertain the relevance of Δ40p53-

related changes in miR-4671-5p and SGSH, we analysed a small cohort of colorectal cancer

(CRC) patient samples available to us. Relevant clinical findings in this cohort are tabulated

in **Figure S1, Table 1.** Processed tissue samples from the patients were interrogated for

Δ40p53 protein expression **(Figure 5 A)** and binned into Δ40p53-high (C4, C1, C14, C5) and

Δ40p53-low (C8, C2, C3, C11, C15, C16) groups by densitometric analysis of immunoblots

(Figure 5 B). C11 was excluded on account of near-absence of p53 as well as Δ40p53

expression relative to other CRC samples. Normalized expression of miR-4671-5p and *SGSH*

mRNA levels from tumours revealed low miRNA and high mRNA levels in the Δ40p53-high

compared to Δ40p53-low group **(Figure 5 C-D)**, supporting our earlier findings in

HCT116+/+ and -/- cells. Importantly, the levels of miR-4671-5p and *SGSH* were inversely

correlated (Pearson r = -0.8061, two-tailed p-value =0.0087) in CRC patients as well (**Figure**

5 E).

Further, to analyse association of Δ40p53 in metastatic progression of CRC, we correlated its

levels with TNM clinical classification of tumours that denotes anatomic extent of the disease

as well as prognostic cancer staging. Here we observed that the samples with relatively high

Δ40p53 (C1, C4, C5 and C14) had lymph node metastasis; histopathological analysis

revealed normal appearing colonic mucosal glands in adjacent normal tissues (**Figure S2 A**),

whereas tumor sections obtained from the qualified tumour blocks show glandular pattern of

adenocarcinoma enriched in pleomorphic cells with a vesicular nucleus with nucleoli

surrounded by desmoplastic (fibrous) stroma and inflammatory cells. Relevantly, these

samples (C1, C4, C5 and C14) demonstrate extensive tumour infiltration and deposits,

indicating nodal metastasis and more advanced disease (**Figure S2 B**), implying that Δ40p53

is associated with aggressive disease progression in CRC.

DISCUSSION

MicroRNAs regulate diverse cellular processes, including embryonic development, cell

cycle, proliferation, and survival. Cancer is often associated with changes in miRNA

expression [33]. Several studies have identified p53-regulated miRNAs and their roles in cancer

[34]. Common malignancies express different p53 isoforms [6]. However, miRNA regulation by

p53 isoforms had remained unknown until recently, and we showed that Δ40p53 specifically

upregulates miR-186-5p and downregulates expression of the oncogene YY1, leading to a

decrease in cell proliferation [22]. The function(s) of Δ40p53 has not been studied in detail,

although its importance has been acknowledged given its roles in cellular functions such as

apoptosis, cell cycle regulation, senescence, migration, and invasion [8]. If Δ40p53 is an

important regulatory molecule, the pertinent question is which cellular pool of RNAs,

specifically long non-coding (lnc) and microRNAs, are regulated. This has gained increasing

importance in many areas of research.

Here, we identified several miRNAs whose expression is regulated by Δ40p53, either

differentially or to the same extent as p53. We then assessed the role of miR-4671-5p in

cellular processes, as it was regulated exclusively by Δ40p53. Overexpressed Δ40p53

decreased, whereas p53 did not change the expression of the novel miR-4671-5p. However,

overexpressed 14A (combination of Δ40p53 and p53) decreased in miR-4671-5p expression,

indicating that p53 facilitates the Δ40p53-mediated regulation of miR-4671-5p **(Figure 2B)**.

Likewise, different ratios of p53 and Δ40p53 reduced miR-4671-5p under all conditions

where both isoforms were present **(Figure 3D)**. This indicated that a differential abundance

of p53 and Δ40p53 expression in cancer will accordingly lead to variable expression of

downstream targets. To characterize the role of miR-4671-5p, we identified potential mRNA

targets using bioinformatic analysis **(Figure S1, Table 2)** and selected a unique target,

SGSH, that plays a metabolic role in governing cellular function. We report possible

prognostic value of SGSH gene expression in certain tumor types **(Figure 3 F-I)**; miR-4671-

5p levels trend oppositely in these cancer types **(Figure 3 A)**. High SGSH expression was

also significantly correlated with poor survival in glioblastoma multiforme (GBM), a highly

malignant adult brain tumor **(Figure S1, Table 4)**. In addition, high hsa-miR-4671-5p is

correlated with poor survival in a miRNA expression dataset in neuroblastoma, a common

pediatric tumor, while low SGSH levels are significantly correlated with poor survival in

multiple neuroblastoma gene expression datasets **(Figure S1, Table 4)**, providing further

evidence of a putative link between the two. While the prognostic value of *SGSH* is

significant, the seemingly confounding directionality in survival-correlation (high levels

correlated to better survival in pancreatic adenocarcinoma and neuroblastoma versus low

levels, in colon tumor and GBM) hints that SGSH is not a bona fide proto-oncogene or tumor

suppressor; rather such differences may be explained by unique cancer-specific adaptation

programs modifying Δ40p53-miR-4671-5p-SGSH axis that requires further investigation.

Further, from enrichment analysis, we selected CDK11B and CDK5R1 as potential targets.

We found that overexpressed Δ40p53 increased the abundance of all targets **(Figure 3 J-L)**,

and the proteins also had a higher abundance in the presence of endogenous Δ40p53 **(Figure**

3 M). Although miR-4671-5p expression was decreased by co-expressed Δ40p53 and FLp53

(Figure 3 D), this trend was not apparent at the target mRNA level **(Figure 3 J-L)**. This

might stem from the activated or deactivated expression of other miRNA by combined p53

isoforms, which in turn would influence target mRNA regulation by miR-4671-5p. We then

determined the relevance of the Δ40p53-miR4671-5p-mRNA axis. Target mRNA expression

was directly decreased by overexpressed miRNA and silenced Δ40p53 **(Figure 4 A, D)**,

confirming a link between them. Further, we measured protein levels of target mRNAs to

determine the cellular effects of interactions. Direct miRNA overexpression and silenced

Δ40p53 decreased SGSH protein levels **(Figure 4 E, F)**. We assessed the physiological

relevance of SGSH and miRNA interactions before delving into the cellular effects of this

regulation. The SGSH encodes a sulfamidase enzyme associated with the lysosomal

degradation of heparan sulfate, a glycosaminoglycan. Mutations in SGSH cause impaired

degradation of heparan sulfate in MPS Type IIIA [35, 36]. How GAGs affect the cell cycle, a

complex process that controls cell growth and division, in MPS disorders has been

investigated. Flow cytometry findings have revealed disturbed cell cycle in all MPS types [37].

These findings suggest that SGSH is directly involved in cell cycle regulation. Along with

SGSH, other targets of miR-4671-5p, namely CDK11B and CDK5R1are also associated with

cell cycle regulation. Therefore, we directly examined whether miR-4671-5p mediates cell

cycle effects on different cell cycle phases. Upon miR-4671-5p overexpression, the ratios (%)

of cells respectively increased and decreased in the S and G2 phases **(Figure 4G)**. The

increased abundance of S-phase cells may be attributed to cell cycle arrest in the intra-S

phase. While this is expected from regulation impinging on CDK5R1 and CDK11B, the

effect of SGSH on cell cycle has been relatively obscure. Populations in the S and G2 phases

are respectively increased and decreased in MPS Type IIIA, where SGSH is mutated [37].

Heparan sulfate inhibits topoisomerase 1 activity during DNA replication [38], and blocking

this activity causes cell cycle arrest at the intra-S phase [39]. Thus, the inhibition of SGSH

expression by miR-4671-5p, apart from inhibiting cell-cycle regulatory kinase functions,

might increase the heparan sulfate content in cells, combinatorically causing an increase in S-

phase cell counts (intra-S-phase arrest).

Building on our cellular findings and based on the reported prognostic value of SGSH, we

expanded our research to include cancer tissue samples to explore the physiological relevance

of Δ40p53/ miR-4671-5p and SGSH axis. Analysis of an Indian cohort of colorectal

carcinoma (CRC) patients pointed to the stratification of samples with both low miR-4671-5-

p and high SGSH mRNA levels into the subgroup of patients where Δ40p53 expression was

high **(Figure 5A-D)**. Overall, miR-4671-5p and SGSH levels were inversely correlated in our

cohort **(Figure 5E).** Additionally, TNM staging and histopathological staining from the

patient samples showed high Δ40p53 (C1, C4, C5 and C14) in patients with lymph node

metastasis, implying that high Δ40p53 is associated with more advanced CRC. Demographic

and clinical profile of colorectal cancers in India differs from the western world, with higher

incidence in younger patients, higher proportion of signet ring carcinomas, and advanced

stage presentation and future comparative studies may be of clinical and mechanistic

relevance[40]. For clinical settings, a neoepitope-like antibody may be developed leveraging

potentially unique structural features of the N-terminus of Δ40p53 as a more advanced
detection tool.

Thus, Δ40p53 ideally reduces miR-4671-5p expression, resulting in elevated levels of target
mRNAs such as *SGSH*, which support proper cell cycle regulation. However, in a
physiological context, elevated Δ40p53 levels can inhibit intra-S-phase cell cycle arrest,
leading to unchecked cell cycle progression, uncontrolled division, and potentially metastasis
in advanced stages of cancer **(Figure 5F)**. This role of Δ40p53 needs further investigation to
clarify its connection to cellular mechanisms and the context of cancer tissue samples.

ACKNOWLEDGEMENTS

We are grateful to Prof. J.C. Bourdon, University of Dundee, UK and Dr. Robin Fahraeus,
INSERM, France, for providing us with the anti-p53 antibody. We want to acknowledge
NIBMG (National Institute of Biomedical Genomics, Kalyani, WB, India) for providing an
RNA sequencing facility. We also thank the present and past SD lab members for their
helpful discussion of the work. A research grant supports this work to SD from the
Department of Biotechnology (DBT), Govt. of India. SD also acknowledges the J.C. Bose
fellowship for research support. The DBT-IISc partnership program also supported this study.
DST Fund for Improvement of Science and Technology Infrastructure (FIST) level II
infrastructure and University Grants Commission Centre of Advanced Studies support is
acknowledged.

AUTHOR CONTRIBUTIONS

AP and SD: Conception and design of studies analysis, interpretation, and article writing. AP,
SKT and PKG: performing experiments, interpretation of results and article editing. DK:
bioinformatic analysis, interpretation of results, article writing and editing. SaG, SP and AM:
RNA sequencing and analysis. SoG: Colorectal cancer sample collection and experimental
procedures; PC: Patient recruitment; KMP: Histological analysis; MD: technical review and
coordination for colorectal cancer samples. SD: Funding acquisition, supervision and project
management.

DATA AVAILABILITY STATEMENT

The sequencing data files obtained is available in the online biorepository forum with the SRP BioProject ID **PRJEB47067.**

CONFLICT OF INTEREST

The authors declare no conflict of interest.

REFERENCE

1.   Kops GJ, Weaver BA, Cleveland DW. On the road to cancer: aneuploidy and the mitotic checkpoint. *Nat Rev Cancer* 2005, **5**(10): 773-785.

2.   Powell DJ, Hrstka R, Candeias M, Bourougaa K, Vojtesek B, Fahraeus R. Stress-dependent changes in the properties of p53 complexes by the alternative translation product p53/47. *Cell Cycle* 2008, **7**(7): 950-959.

3.   Rivlin N, Brosh R, Oren M, Rotter V. Mutations in the p53 Tumor Suppressor Gene: Important Milestones at the Various Steps of Tumorigenesis. *Genes Cancer* 2011, **2**(4): 466-474.

4.   Bourdon JC. p53 and its isoforms in cancer. *Br J Cancer* 2007, **97**(3): 277-282.

5.   Khoury MP, Bourdon JC. p53 Isoforms: An Intracellular Microprocessor? *Genes Cancer* 2011, **2**(4): 453-465.

6.   Vieler M, Sanyal S. p53 Isoforms and Their Implications in Cancer. *Cancers (Basel)* 2018, **10**(9).

7.   Bourdon JC, Fernandes K, Murray-Zmijewski F, Liu G, Diot A, Xirodimas DP, *et al.* p53 isoforms can regulate p53 transcriptional activity. *Genes Dev* 2005, **19**(18): 2122-2137.

8.   Steffens Reinhardt L, Zhang X, Wawruszak A, Groen K, De Iuliis GN, Avery-Kiejda KA. Good Cop, Bad Cop: Defining the Roles of Delta40p53 in Cancer and Aging. *Cancers (Basel)* 2020, **12**(6).

9.   Ray PS, Grover R, Das S. Two internal ribosome entry sites mediate the translation of p53 isoforms. *EMBO Rep* 2006, **7**(4): 404-410.

517 10. Khan D, Katoch A, Das A, Sharathchandra A, Lal R, Roy P, *et al.* Reversible induction of
518   translational isoforms of p53 in glucose deprivation. *Cell Death Differ* 2015, **22**(7): 1203-
519   1218.

520
521 11. Sharathchandra A, Katoch A, Das S. IRES mediated translational regulation of p53 isoforms.
522   *Wiley Interdiscip Rev RNA* 2014, **5**(1): 131-139.

523
524 12. Ghosh A, Stewart D, Matlashewski G. Regulation of human p53 activity and cell localization
525   by alternative splicing. *Mol Cell Biol* 2004, **24**(18): 7987-7997.

526
527 13. Hafsi H, Santos-Silva D, Courtois-Cox S, Hainaut P. Effects of Delta40p53, an isoform of p53
528   lacking the N-terminus, on transactivation capacity of the tumor suppressor protein p53.
529   *BMC Cancer* 2013, **13**: 134.

530
531 14. Ohki R, Kawase T, Ohta T, Ichikawa H, Taya Y. Dissecting functional roles of p53 N-terminal
532   transactivation domains by microarray expression analysis. *Cancer Sci* 2007, **98**(2): 189-200.

533
534 15. Ota A, Nakao H, Sawada Y, Karnan S, Wahiduzzaman M, Inoue T, *et al.* Delta40p53alpha
535   suppresses tumor cell proliferation and induces cellular senescence in hepatocellular
536   carcinoma cells. *J Cell Sci* 2017, **130**(3): 614-625.

537
538 16. Maier B, Gluba W, Bernier B, Turner T, Mohammad K, Guise T, *et al.* Modulation of
539   mammalian life span by the short isoform of p53. *Genes Dev* 2004, **18**(3): 306-319.

540
541 17. Bourougaa K, Naski N, Boularan C, Mlynarczyk C, Candeias MM, Marullo S, *et al.*
542   Endoplasmic reticulum stress induces G2 cell-cycle arrest via mRNA translation of the p53
543   isoform p53/47. *Mol Cell* 2010, **38**(1): 78-88.

544
545 18. Thomas SE, Malzer E, Ordonez A, Dalton LE, van TWEFA, Liniker E, *et al.* p53 and translation
546   attenuation regulate distinct cell cycle checkpoints during endoplasmic reticulum (ER) stress.
547   *J Biol Chem* 2013, **288**(11): 7606-7617.

548
549 19. Ungewitter E, Scrable H. Delta40p53 controls the switch from pluripotency to differentiation
550   by regulating IGF signaling in ESCs. *Genes Dev* 2010, **24**(21): 2408-2419.

551
552 20. Hinault C, Kawamori D, Liew CW, Maier B, Hu J, Keller SR, *et al.* Delta40 Isoform of p53
553   controls beta-cell proliferation and glucose homeostasis in mice. *Diabetes* 2011, **60**(4): 1210-
554   1222.

555
556 21. Zhang X, Groen K, Morten BC, Steffens Reinhardt L, Campbell HG, Braithwaite AW, *et al.*
557   Effect of p53 and its N-terminally truncated isoform, Delta40p53, on breast cancer migration
558   and invasion. *Mol Oncol* 2022, **16**(2): 447-465.

560   22.   Katoch A, Tripathi SK, Pal A, Das S. Regulation of miR-186-YY1 axis by the p53 translational
561   isoform Δ40p53: implications in cell proliferation. *Cell Cycle* 2021, **20**(5-6): 561-574.

562
563   23.   Pal A, Ghosh PK, Das S. The "LINC" between Δ40p53-miRNA Axis in the Regulation of Cellular
564   Homeostasis. *Mol Cell Biol* 2023, **43**(7): 335-353.

565
566   24.   Sun Y, Manceau A, Frydman L, Cappuccio L, Neves D, Basso V, *et al.* Delta40p53 isoform up-
567   regulates netrin-1/UNC5B expression and potentiates netrin-1 pro-oncogenic activity. *Proc*
568   *Natl Acad Sci U S A* 2021, **118**(36).

569
570   25.   Xu F, Wang Y, Ling Y, Zhou C, Wang H, Teschendorff AE, *et al.* dbDEMC 3.0: Functional
571   Exploration of Differentially Expressed miRNAs in Cancers of Human and Model Organisms.
572   *Genomics, Proteomics & Bioinformatics* 2022.

573
574   26.   Agarwal V, Bell GW, Nam J-W, Bartel DP. Predicting effective microRNA target sites in
575   mammalian mRNAs. *eLife* 2015, **4**: e05005.

576
577   27.   Chen Y, Wang X. miRDB: an online database for prediction of functional microRNA targets.
578   *Nucleic Acids Res* 2020, **48**(D1): D127-d131.

579
580   28.   Liu W, Wang X. Prediction of functional microRNA targets by integrative modeling of
581   microRNA binding and target expression data. *Genome Biology* 2019, **20**(1): 18.

582
583   29.   Zhang L, Liao Y, Tang L. MicroRNA-34 family: a potential tumor suppressor and therapeutic
584   candidate in cancer. *Journal of Experimental & Clinical Cancer Research* 2019, **38**(1): 53.

585
586   30.   Li X, Xiao R, Chen B, Yang G, Zhang X, Fu Z, *et al.* A novel mutation of SGSH and clinical
587   features analysis of mucopolysaccharidosis type IIIA. *Medicine* 2018, **97**(52): e13758.

588
589   31.   Anuraga G, Lang J, Xuan DTM, Ta HDK, Jiang JZ, Sun Z, *et al.* Integrated bioinformatics
590   approaches to investigate alterations in transcriptomic profiles of monkeypox infected
591   human cell line model. *Journal of infection and public health* 2024, **17**(1): 60-69.

592
593   32.   Ye J, Liu P, Li R, Liu H, Pei W, Ma C, *et al.* Biomarkers of connective tissue disease-associated
594   interstitial lung disease in bronchoalveolar lavage fluid: A label-free mass spectrometry-
595   based relative quantification study. *Journal of clinical laboratory analysis* 2022, **36**(5):
596   e24367.

597
598   33.   Israel A, Sharan R, Ruppin E, Galun E. Increased MicroRNA Activity in Human Cancers. *PLOS*
599   *ONE* 2009, **4**(6): e6045.

600
601   34.   Takwi A, Li Y. The p53 Pathway Encounters the MicroRNA World. *Current genomics* 2009,
602   **10**(3): 194-197.

603
604   35.   Griffin LS, Gloster TM. The Enzymatic Degradation of Heparan Sulfate. *Protein and peptide*
605   *letters* 2017, **24**(8): 710-722.

606
607   36.   Boado RJ, Lu JZ, Hui EK-W, Pardridge WM. Reduction in Brain Heparan Sulfate with Systemic
608   Administration of an IgG Trojan Horse–Sulfamidase Fusion Protein in the
609   Mucopolysaccharidosis Type IIIA Mouse. *Molecular Pharmaceutics* 2018, **15**(2): 602-608.

610
611   37.   Brokowska J, Gaffke L, Pierzynowska K, Cyske Z, Węgrzyn G. Cell cycle disturbances in
612   mucopolysaccharidoses: Transcriptomic and experimental studies on cellular models.
613   *Experimental biology and medicine (Maywood, NJ)* 2022, **247**(18): 1639-1649.

614
615   38.   Kovalszky I, Dudás J, Oláh-Nagy J, Pogány G, Töváry J, Timár J, *et al.* Inhibition of DNA
616   topoisomerase I activity by heparan sulfate and modulation by basic fibroblast growth
617   factor. *Mol Cell Biochem* 1998, **183**(1-2): 11-23.

618
619   39.   Lin R-W, Yang C-N, Ku S, Ho C-J, Huang S-B, Yang M-C, *et al.* CFS-1686 Causes Cell Cycle
620   Arrest at Intra-S Phase by Interference of Interaction of Topoisomerase 1 with DNA. *PLOS*
621   *ONE* 2014, **9**(12): e113832.

622
40.   Patil PS, Saklani A, Gambhire P, Mehta S, Engineer R, De'Souza A, *et al.* Colorectal Cancer in
India: An Audit from a Tertiary Center in a Low Prevalence Area. *Indian journal of surgical*
oncology 2017, **8**(4): 484-490.

FIGURE LEGENDS

Figure 1. Regulation of miRNA expression by p53 and its isoform Δ40p53

(A) Schematic of constructs used in study (B) Experimental setup prior to RNA sequencing.

(C) Western blot analysis of cell extracts from H1299 cells expressing control, p53 only,

Δ40p53 only and 14A construct, probed with CM1 after 48 h. (D) The cluster of differentially

expressed miRNAs obtained from the small RNA sequencing data set. (E) The compressed

cluster of significant differentially expressed miRNAs with p-value ≤ 0.05 (generated using

Heatmapper).

Figure 2. Selection and validation of miRNAs for further studies

(A) Western blot analysis of cell extracts from H1299 cells expressing control, p53 only,

Δ40p53 only and 14A construct, probed with CM1 after 48 h. (B-E) Quantitative PCR for

validation of miR-4671-5p, miR-34a-5p, miR-548ae-5p, miR-301b-5p, respectively, in H1299 cells expressing control, p53 only, Δ40p53 only and 14A construct. (F) Table for comparative analysis of fold changes obtained in sequencing versus validation result. (G) Western blot analysis of cell extracts from HCT116+/+ and HCT116-/-cells probed with CM1. (H) Quantitative PCR for validations of miR-4671-5p, miR-34a-5p and miR-548ae-5p in HCT116+/+ and HCT116-/-cells. (I) Quantitative PCR of miR-4671-5p, miR-34a-5p and miR-548ae-5p in HCT116+/+ cells transfected with si p53 (30nM) and non-specific si (si Nsp). (J) Western blot analysis of cell extracts from HCT116+/+ cells transfected with either si p53 (30nM) and non-specific si (si Nsp), probed with CM1. (K) Quantitative PCR of miR-4671-5p, miR-34a-5p and miR-548ae-5p in HCT116-/- cells transfected with si Δ40p53 (30nM) and non-specific si (si Nsp). (L) Western blot analysis of cell extracts from HCT116-/- cells transfected with siΔ40p53 (30nM) and non-specific si (si Nsp), probed with CM1 The criterion for statistical significance was $p≤0.05$ (*) or $p≤0.01$ (**) or $p≤0.001$(***)

Figure 3. Regulation of miRNAs by Δ40p53 and p53

(A) Differential expression Profile of miRNAs in cancer vs. normal obtained from dbDEMC 3.0 database. (B) Western blot analysis of cell extracts from H1299 transfected cells with different ratios of p53 and Δ40p53 probed with CM1. (C-D) Quantitative PCR of miR-34a-5p and miR-4671-5p respectively in H1299 transfected with different ratios of p53 and Δ40p53. (E) Table for miR-4671-5p targets selected in the study. (F-I) Kaplan-Meier estimates of survival for pancreatic adenocarcinoma (F-G) and colon tumor (H-I) classified by SGSH mRNA expression levels. The number of patients at risk is indicated for time increments of 12 or 24 months. p values were calculated using a log rank test. (J-L) Quantitative PCR of mRNA targets (SGSH, CDK11B and CDK5R1) in H1299 transfected cells with different ratios of p53 and Δ40p53. (M) Western blot analysis of cell extracts from HCT116+/+ and HCT116-/- cells probed with CM1, SGSH, CDK11B, CDK5R1 antibodies. The criterion for statistical significance was $p≤0.05$ (*) or $p≤0.01$ (**) or $p≤0.001$(***).

Figure 4. role of miR-4671-5p

(A) Quantitative PCR of SGSH, CDK11B and CDK5R1 in HCT116-/- cells transfected with si Δ40p53 (30nM) and non-specific si (si Nsp). (B) Western blot analysis of cell extracts from HCT116-/- cells transfected with siΔ40p53 (30nM) and non-specific si (si Nsp), probed with CM1. (C) Quantitative PCR of miR-4671-5p in HCT116-/- cells transfected with miR-4671-5p overexpression construct. (D) Quantitative PCR of SGSH, CDK11B and CDK5R1 in

HCT116-/- cells transfected with miR-4671-5p overexpression construct. (E) Western blot analysis of cell extracts from HCT116-/- cells transfected with siΔ40p53 (30nM) and non-specific si (si Nsp), probed with CM1 and SGSH. (F) Western blot analysis of cell extracts from HCT116-/- cells transfected with miR-4671-5p overexpression construct, probed with SGSH. (G) Analysis of different cell cycle phases of fixed cell extracts from HCT116-/- cells transfected with miR-4671-5p overexpression construct. The criterion for statistical significance was $p \leq 0.05$ (*) or $p \leq 0.01$ (**) or $p \leq 0.001$ (***).

Figure 5. Δ40p53 expression level in colorectal cancer (CRC) patients stratifies miR-4671-5p and SGSH levels

(A) Western blot analysis of p53 and Δ40p53 protein expressions in fresh frozen resected tissue samples from CRC patients (C1-C5, C8, C11, C14-16). Western blots from HCT116+/+ (expresses both isoforms) and HCT116-/- (expresses only Δ40p53) colon cancer cells as markers for relative migration of p53 and Δ40p53 done in same gel and same membrane as C1-C5, at lower exposure than C1-C5. (B) Densitometric analysis of Δ40p53 band in CRC patients (arb. unit: arbitrary units), p-value from unpaired t-test (C) Quantitative PCR Ct value of miR-4671-5p levels normalized to U6 RNA, p-value from unpaired t-test (D) Quantitative PCR Ct value of SGSH levels normalized to ACTB RNA, p-value from unpaired t-test. (E) Correlation of normalized miR-4671-5p Ct values with normalized SGSH Ct values across CRC patients. Simple linear regression line in grey (R^2=0.6498).

Figure 5F. Model depicting the regulation of Δ40p53 mediated regulation of miR-4671-5p and their impact on cell cycle

(A)When Δ40p53 is present, it lowers miR-4671-5p levels, inhibiting the degradation of target mRNAs. This increase in residual mRNAs leads to higher protein production (e.g., SGSH), facilitating normal cell cycle progression. In this context, the mechanism may involve enhanced degradation of HS, contributing to the stabilization of metabolic and cellular homeostasis. (B) Cancer Tissue Samples Comparison of patient samples based on Δ40p53 expression levels. Samples with relatively high Δ40p53 (C1, C4, C5, and C14) exhibited low levels of miR-4671-5p mRNA and high levels of *SGSH* mRNA. In contrast, patients with low Δ40p53 displayed higher miR-4671-5p levels and lower *SGSH* mRNA levels. Moreover, samples with high Δ40p53 revealed lymph node metastasis implying that high Δ40p53 is associated with more advanced CRC.

Figure S1: (Table 1): Clinical and pathological details of the patient samples. *T, size and extent of the main tumor; N, number of nearby lymph nodes that have cancer; M, metastasis. **(Table 2)**: miR-4671-5p targets obtained from TargetScan 7.2 database and miRDB. **(Table 3)**: Result of PANTHER Statistical Overrepresentation Test of "GO molecular function"-terms obtained from 786 predicted miR-4671-5p targets, using all *Homo sapiens* genes as reference list. FDR: False Discovery Rate. Results for FDR $P < 0.05$ reported. **(Table 4)**: Survival analysis of patients as classified by SGSH mRNA expression levels in pancreatic adenocarcinoma, colon tumor, glioblastoma and neuroblastoma gene expression datasets available on R2: Genomics Analysis and Visualization Platform. p-values <0.07 reported, below 0.05 considered significant. Similar analysis for hsa-miR-4671-5p expression, as available in the sole miRNA expression dataset on R2.

Figure S2: (A) H&E images of FFPE samples from tumor and adjacent normal tissues. (Yellow arrow indicates tumor features) (B) H&E images of FFPE samples from lymph nodes with metastasized and normal lymph node tissues (Yellow arrow indicates deposited tumor cells)

a

b

WORK FLOW

p53   Δ40p53   14A (p53/Δ40p53)

H1299 cells
(p53/Δ40p53
null cell line)

Isolation of total RNA

Small RNA Sequencing

c

kDa	Control	p53	Δ40p53	14A (p53 +Δ40p53)	
63-					
48-					p53
					Δ40p53
48-					β-Actin

d

Color Key

-2 0 2
Value

Control   1 2 3   1 2 3   1 2 3
          14A (p53   Δ40p53   p53
          +Δ40p53)

e

Δ40p53   p53   14A (p53 +Δ40p53)

-1 0.5
Value

hsa-miR-34a-5p
hsa-miR-548ae-5p
hsa-miR-182-5p
hsa-miR-34c-5p
hsa-miR-561-5p
hsa-miR-148b-5p
hsa-miR-30e-5p
hsa-miR-7-5p
hsa-miR-708-5p
hsa-miR-629-5p
hsa-miR-378a-5p
hsa-miR-129-5p
hsa-miR-320b
hsa-miR-3909
hsa-miR-128-1-5p
hsa-miR-1270
hsa-miR-3180-5p
hsa-miR-4661-5p
hsa-miR-103a-2-5p
hsa-miR-1284
hsa-miR-4671-5p
hsa-miR-3940-5p
hsa-miR-301b-5p