Rice stripe virus utilizes an *Laodelphax striatellus* salivary carbonic anhydrase to facilitate plant infection by direct molecular interaction

Authors' names

Jing Zhao[1, 2, #], Xiangyi Meng[1, 2, #], Jie Yang[1, 2], Rongxiang Fang[1, 2], Yan Huo[1*], Lili Zhang[1, 3*]

Authors' affiliations

[1] Department of Agri-microbiomics and Biotechnology, State Key Laboratory of Microbial Diversity and Innovative Utilization, Institute of Microbiology, Chinese Academy of Sciences, Beijing 100101, China.

[2] College of Life Sciences, University of the Chinese Academy of Sciences, Beijing 100049, China.

[3] College of Advanced Agricultural Sciences, University of the Chinese Academy of Sciences, Beijing 100049, China.

*Corresponding author

* Address correspondence to: Dr. Lili Zhang, Institute of Microbiology, Chinese Academy of Sciences, No 3, Yard 1, West Beichen Road, Chaoyang, Beijing 100101, China. Phone (+86)10-64806916; Fax (+86)10-64806543; E-mail: Zhangll@im.ac.cn

* Address correspondence to: Dr. Yan Huo, Institute of Microbiology, Chinese Academy of Sciences, No 3, Yard 1, West Beichen Road, Chaoyang, Beijing 100101, China. Phone (+86)10-64806915; Fax (+86)10-64806543; E-mail: huoy@im.ac.cn

Abstract

Plant viruses typically rely on insect vectors for transmission between plants, with insect salivary proteins playing critical roles in this process. In this study, we demonstrate how *Laodelphax striatellus* salivary carbonic anhydrase (LssaCA) promotes rice stripe virus (RSV) infection in plants. We discovered that LssaCA directly binds to RSV nucleocapsid protein (NP) in insect salivary glands. This LssaCA-NP complex interacts with a rice thaumatin-like protein (OsTLP) that possess endo-β-1,3-glucanase activity potentially degrading callose. Upon binding, the LssaCA-NP complex significantly enhances OsTLP enzymatic activity. We further clarify that both *L. striatellus* feeding and RSV infection induce callose deposition. The tripartite LssaCA-NP-OsTLP interaction enhances callose degradation, thereby facilitating RSV infection via its insect vector. This study provides new insights into complex virus-insect-plant tripartite interactions mediated by insect salivary proteins, with broad implications for numerous plant viruses transmitted by insect vectors.

Keywords: rice stripe virus, *Laodelphax striatellus*, viral infection of plant, salivary carbonic anhydrase, rice thaumatin-like protein, callose deposition

Introduction

During plant virus transmission by piercing-sucking insects, most viruses are inoculated into the plant phloem via the insect's secreted saliva (Arca & Ribeiro, 2018; Conway et al., 2016; Wu et al., 2022). Thus, insect saliva acts as an interface for the virus-insect-host tripartite interaction and can directly promote viral transmission to, and infection of, the host plants (Sun et al., 2020; Wu et al., 2022). However, despite the importance of insect salivary proteins in this tripartite interaction, there is still much to learn about how these proteins enable successful viral infection.

Previous studies have revealed that there are two ways in which insect saliva facilitates viral infection. One is an indirect approach whereby the saliva modulates the host microenvironment at the feeding site, and saliva effectors work together to allow the arthropod to go unnoticed while it feeds on the host plant (Acevedo et al., 2019; Arca & Ribeiro, 2018; Sun et al., 2020). For example, our recent work indicated that a *Laodelphax striatellus* mucin protein, LssaMP, enables the formation of the salivary sheath and facilitates the transmission of rice stripe virus (RSV) into the rice phloem (Huo et al., 2022). The study on leafhoppers revealed that the expression of a saliva calcium-binding protein is inhibited by rice gall dwarf virus (RGDV), thus causing an increase of cytosolic Ca^{2+} levels in rice and triggering callose deposition and H_2O_2 production. This increases the frequency of insect probing, thereby enhancing viral horizontal transmission into the rice phloem (Wu et al., 2022). The other mechanism by insect saliva to facilitate virus infection is direct regulation, whereby saliva proteins promote virus transmission through specific molecular interactions (Wen et al., 2019). Direct saliva protein-pathogen interactions have been reported in animal pathogens. For example, during transmission of *Borrelia burgdorferi* by *Ixodes scapularis*, the saliva protein Salp15 of *I. scapularis* binds to the bacterial outer surface protein C, which prevents the bacterium from being recognized by the animal immune system. In this way, the saliva protein enables the pathogen to infect the animal host (Ramamoorthi et al., 2005; Schuijt et al., 2008).

Although most plant viruses are heavily dependent on insect vectors for plant-to-plant

transmission (Gray, 2008), the direct function of insect saliva proteins in mediating

virus transmission remains largely uninvestigated.

During sap-feeding, arthropods produce two distinct types of saliva at different

stages of the feeding process: gel saliva and watery saliva (Bonaventure, 2012; Lou et

al., 2019). The former forms a salivary sheath to provide a smooth path for the stylet

penetration (Lou et al., 2019). The latter is mainly secreted into the phloem sieve

elements to prevent them from plugging up and suppress plant defense responses.

Some salivary components may act as herbivore-associated molecular patterns that

can trigger pattern-triggered immunity, and certain salivary effectors may be

recognized by plant resistance proteins to induce effector-triggered immunity, etc.

(Huang et al., 2017; Ji et al., 2017; Yi et al., 2021).

For phloem-feeding insects, callose deposited on phloem sieve plate and

plasmodesmata of sieve elements functions as a defense mechanism by reducing

insect feeding and preventing viral movement (Hao et al., 2008; Hipper et al., 2013;

Will & Vilcinskas, 2015; Zavaliev et al., 2011; Yue et al., 2022). Callose is a

β-(1,3)-D-glucan polysaccharide that is synthesized by callose synthases and

degraded by β-(1,3)-glucanases. Plants defend themselves by depositing callose at the

sieve plates and plasmodesmata in response to virus infection, whereas viruses

counter this defense by activating β-(1,3)-glucanases to degrade callose (Bucher et al.,

2001; Hao et al., 2008; Wu et al., 2022; Zavaliev et al., 2011).

RSV is the causative agent of rice stripe disease, a serious disease of rice crops

that has occurred repeatedly in China, Japan, and Korea (Xu et al., 2021). RSV is

completely dependent on insect vectors for transmission among its host plants, and *L.*

striatellus is the main vector (Xu et al., 2021; Zhao et al., 2018). *L. striatellus*

transmits RSV in a persistent-propagative manner. The virus initially infects the

midgut, then disperses from the hemolymph into the salivary glands, and is inoculated

into the plant host during *L. striatellus* feeding (Huo et al., 2022). *L. striatellus*

belongs to the order Hemiptera, whose members mainly feed from sieve tubes

through their mouthparts (stylets) that penetrate plant tissues and reach sieve tubes to

ingest the phloem sap (Tjallingii, 2006; Van Bel & Will, 2016). RSV is mainly

secreted into the rice phloem via the watery saliva (Huo et al., 2022; Wang & Blanc,

2021).

In this study, we identified a molecular interaction among RSV, an *L. striatellus*

saliva protein, and a plant β-1,3-glucanase. The insect saliva protein directly binds to

the RSV nucleocapsid protein (NP) and then binds to a rice thaumatin-like protein to

activate its β-1,3-glucanase activity. The activation of β-1,3-glucanase helps RSV

infection by inhibiting callose deposition in response to viral infection.

Results

L. striatellus *salivary carbonic anhydrase directly bind to RSV NP*

Insect-secreted watery saliva is inoculated into plant phloem sieve tubes along with RSV (Huo et al., 2022). We identified the salivary proteins that directly interact with RSV to promote viral plant infection. Among the saliva proteins identified in our previous study from RSV-free *L. striatellus* (Huo et al., 2022), Sap6 (GenBank accession No. RZF48846.1) was found to co-localize with RSV in infected salivary glands, suggesting a potential molecular interaction between them (Fig. 1A and S2).

Sequence homology analysis revealed Sap6 as a carbonic anhydrase-like protein, with a signal peptide located at the N-terminus (amino acids 1-24) (Fig. 1B and S1A). We recombinantly expressed this protein in *E.coli* or in sf9 cells as a secreted protein. The purified proteins were measured enzymatic activity in hydrolyzing *p*-nitroethyl acetate. The observed hydrolyzing activity confirmed Sap6 as a carbonic anhydrase (abbreviated as CA, Fig. 1C). Sequence analysis indicated seven residues constituting the enzymatic active center: H_{111}, N_{139}, H_{141}, H_{143}, E_{153}, H_{166}, and T_{253}. We then constructed a mutant with substations of all seven residues (Fig. S1B. H111D, N139H, H141D, H143D, E153H, H166D, and T253E). The mutant protein lost enzymatic activity, indicating the necessity of the enzymatic active center (Fig. 1C). Sap6 was subsequently named *L. striatellus* salivary carbonic anhydrase (LssaCA).

We performed pull-down and Microscale thermophoresis (MST) assays to confirm the molecular interaction between LssaCA and RSV NP. Pull-down assays revealed a specific interaction between LssaCA and NP (Fig. 1D). MST assays determined that the K_D of the interaction between LssaCA and NP was 2.7 ± 2.2 μM, further confirming the molecular interaction (Fig. 1E and S3).

We then performed quantitative reverse-transcriptional PCR (RT-qPCR) to determine the gene expression and western blotting to assess protein distribution. RT-qPCR revealed specific expression of *LssaCA* in *L. striatellus* salivary glands (Fig. 1F), while western blotting confirmed the protein's presence in this tissue (Fig. 1F).

When *LssaCA* expression was knocked down using *LssaCA*-specific double-stranded RNA (dsLssaCA) microinjection, LssaCA protein levels in salivary glands decreased dramatically (Fig. 1G). To investigate LssaCA secretion, we allowed RSV-free insects to feed on an artificial diet, collecting watery saliva proteins from the diet solution and gel saliva proteins from the feeding parafilm. Western blotting demonstrated that the LssaCA protein was secreted as a component of watery saliva (Fig. 1H). Furthermore, by measuring LssaCA protein levels in plants after feeding by RSV-free insects, we confirmed that the LssaCA proteins can be secreted into plants (Fig. 1I). We also examined the regulation of *LssaCA* expression by RSV. RSV infection did not significantly influence *LssaCA* mRNA levels in the salivary glands; however, the protein levels was significantly higher in RSV-infected salivary glands (Fig. 1J).

In conclusion, these results indicate that when RSV NP is secreted by *L. striatellus*, a salivary-gland-specifically expressed protein, LssaCA, can directly bind to RSV and be secreted via saliva.

LssaCA enhances RSV infection in plants

To determine the role of LssaCA in RSV infection of rice plants, 36.8 nL dsLssaCA at 1 ng/nL was microinjected into the hemocoel of RSV-infected 3^{rd}-instar nymphs to interfere with gene expression. At 2 days post-microinjection (dpi), *LssaCA* mRNA levels were measured by RT-qPCR and compared with the dsGFP control. Results showed that dsLssaCA treatment reduced *LssaCA* mRNA levels by 80% (Fig. 2A). The two groups of insects were then allowed to feed on healthy rice seedlings (5 insects per rice seedling) for 2 days. RSV NP RNA levels (indicative of RSV titers) in the plants were measured at 14 days post-feeding (dpf), when inoculated viruses had infected the plant host, replicated in plant cells, and viral titers were high enough to be detected by RT-qPCR and western blotting. Compared to rice seedlings fed by dsGFP-treated insects, those fed by dsLssaCA-treated insects had significantly reduced RSV titers (Fig. 2B-C), indicating that LssaCA plays an essential role in mediating RSV infection of rice plants.

To determine at which step LssaCA affected RSV infection, we analyzed the influence of LssaCA-deficiency on RSV titers during several steps of the RSV transmission and infection process, including virus load in salivary glands and saliva, and initial inoculation levels *in planta*. First, at 2 days after dsLssaCA treatment of viruliferous insects, when *LssaCA* was significantly downregulated (Fig. 2D), RT-qPCR was used to detect RSV titer in insect salivary glands. Results showed that *LssaCA* deficiency did not reduce RSV titers in this tissue (Fig. 2E). Then, LssaCA-deficient insects were allowed to feed on an artificial diet, and their saliva was collected for RSV level analysis. Results revealed that LssaCA-deficiency did not reduce RSV titer in the secreted saliva (Fig. 2F). Finally, a rice plant was fed with 10 viruliferous insects for 24 h, with insects limited to a specific area, and RSV titers were measured immediately after feeding. We found that LssaCA-deficiency did not affect RSV initial inoculation levels *in planta* (Fig. 2G). We also monitored the influence of LssaCA deficiency on the insect feeding behaviors by using Electrical penetration graph (EPG). The EPG waveforms can be classified into three different periods: nonpenetration (NP), penetration (N1, N2, N3) and phloem feeding (N4: N4-a, N4-b) periods. The NP waveform indicates a lack of feeding, the N waveform indicates penetration of mesophyll cells to seek the phloem and the N4 waveform indicates digestion of the sap from the phloem. Results indicated that LssaCA deficiency did not affect the insect feeding behaviors (Fig. S4). These results collectly suggest that LssaCA promotes RSV infection through a mechanism occurring not in insects or during early stages of viral entry in plants, but *in planta* after viral inoculation.

LssaCA protein levels were measured after insect feeding. Plants were fed on by 50 RSV-free insects for 2 days, and then LssaCA protein levels were measured at 1 and 3 dpf by western blotting. We found that LssaCA was retained for at least 3 days, and protein levels decreased rapidly from 1 dpf to 3 dpf (Fig. 2H). We then measured RSV infection levels at 3 dpf to assess the *in-planta* influence after viral inoculation. *LssaCA* expression was knocked down using dsLssaCA in RSV-infected 3[rd]-instar

nymphs, and 5 insects were allowed to feed on healthy rice for 2 days. We measured

RSV titers in the whole plant shoot at 3 dpf. We found significantly lower RSV titers

in the dsLssaCA-treatment group at 3 dpf (Fig. 2I-J). These results further confirmed

the enhancement of RSV's plant infection process by LssaCA.

LssaCA interacts with rice thaumatin-like protein to increase its

***endo-β-1,3-glucanase activity and facilitate RSV infection* in planta**

To investigate the mechanism by which LssaCA facilitates RSV infection in

plants, we performed yeast two-hybrid screening to identify rice proteins that interact

with LssaCA. A rice thaumatin-like protein (XP_025879846.1), designated as *Oryza*

sativa thaumatin-like protein (OsTLP) (Fig. 3A-B and S5, Table S1), was identified.

The open reading frame of OsTLP encodes a polypeptide of 327 amino acids with a

thaumatin family (THM) domain from residues 27 to 267. A BLAST algorithm-based

analysis revealed sequence identities to TLP 1b. The N-terminal 24 amino acids

constitute a secretion signal peptide, suggesting OsTLP is a secreted protein.

Pull-down and MST assays were performed to confirm the molecular interaction

between OsTLP and LssaCA. Pull-down assays showed that OsTLP and LssaCA have

a specific interaction (Fig. 3C). The MST assay revealed a K_D of 1.3 ± 2.2 μM for the

OsTLP-LssaCA interaction, further confirming the molecular interaction (Fig. 3D).

Previous studies have shown that TLP orthologs in various plants, including

apples, cherries, tomatoes, and barley, possess endo-β-1,3-glucanase activity that

degrades callose (de Jesus-Pires et al., 2020). To verify the endo-β-1,3-glucanase

activity of OsTLP, we conducted enzymatic assays using recombinantly expressed

OsTLP protein. The purified protein exhibited an average activity of 60 units/mg (Fig.

3E). To examine OsTLP activity *in planta*, we constructed transgenic plants

overexpressing OsTLP (OsTLP OE) by using the *Japonica* rice cultivar Nipponbare

(Fig. S6A-B). These transgenic plant showed significantly increased β-1,3-glucanase

activity compared to wild-type controls (Fig. 3F).

To determine whether the interaction between LssaCA and OsTLP affects

enzymatic activity, we pre-incubated OsTLP with either LssaCA or BSA (control) for

2 h before measuring activity. Pre-incubation with LssaCA significantly enhanced

OsTLP activity compared to the BSA control (Fig. 3G). Since CA family proteins

catalyze the conversion of CO_2 and H_2O to HCO_3^- and H^+, potentially lowering pH,

we examined the pH dependence of OsTLP activity. Activity assays conducted across

a pH range of 5.0-7.5 revealed that OsTLP activity increased at lower pH values (Fig.

3H). However, pH monitoring during OsTLP-LssaCA interaction experiments showed

no change from the initial pH of 7.4, indicating that the enhanced enzymatic activity

was not due to pH changes but rather to the direct interaction between OsTLP and

LssaCA.

Moreover, we measured RSV infection levels in plants with increased OsTLP

activity. Viruliferous *L. striatellus* fed on either OsTLP OE plants or wild-type plants

for 2 days, and RSV titers were measured at 14 dpf. RT-qPCR analyses indicated that

compared with wild-type plants, plants overexpressing OsTLP had significantly

higher RSV titers (Fig. 3I), indicating that OsTLP facilitates RSV infection *in planta*.

Collectively, these results suggest that LssaCA may promote RSV infection by

enhancing OsTLP's callose-degrading activity.

RSV NP-LssaCA-OsTLP tripartite interaction further enhances OsTLP enzymatic

activity

Given that LssaCA interacts with both RSV NP and OsTLP, we investigated the

possibility of a tripartite interaction. Through pull-down assays, we discovered that

when LssaCA was pre-incubated with NP to form an NP-LssaCA complex, this

complex retained the ability to bind to OsTLP (Fig. 4A). This result demonstrates a

tripartite interaction among these three proteins.

We then explored how the tripartite interaction affects OsTLP enzymatic activity.

OsTLP was pre-incubated with either the LssaCA-NP complex or LssaCA alone as a

control. After 2 h of incubation, enzyme activity was measured. The results showed

that pre-incubation with the LssaCA-NP complex significantly increased OsTLP

activity compared to the LssaCA control (Fig. 4B). This indicates that stronger
activation of β-1,3-glucanase activity is achieved through the NP-LssaCA-OsTLP
tripartite interaction.

LssaCA inhibits insect-feeding and RSV-infection induced callose deposition in rice
plants

Callose deposition on sieve plates and plasmodesmata serves as an critical
defense mechanism in plants, reducing insect feeding, viral spread, and infection
severity (Li et al., 2012; Zavaliev et al., 2011). Given that the LssaCA or LssaCA-NP
complex enhances OsTLP's β-1,3-glucanase activity and potentially promotes callose
degradation, we examined callose regulation during feeding by both RSV-free and
RSV-infected *L. striatellus*. To assess feeding-induced callose deposition, we exposed
rice seedlings to either RSV-free or RSV-infected *L. striatellus* for 24 h and analyzed
callose levels around the feeding sites. ELISA assays showed that while feeding by
RSV-free insects increased callose concentration, feeding by RSV-infected insects
triggered an even greater increase (Fig. 5A). Consistent with these findings,
microscopic analysis revealed enhanced callose deposition in phloem cells of plants
fed upon by RSV-infected *L. striatellus* compared to those fed by RSV-free insects
(Fig. 5B-C and S7). To determine whether this response was localized, we measured
callose levels in tissues distant from the feeding sites. No significant differences in
callose deposition were observed between control plants and the non-feeding areas
(Fig. S8A), indicating that callose accumulation was confined to feeding sites and
their immediate vicinity.

To evaluate the molecular mechanisms underlying the induced callose deposition,
we analyzed transcript levels of callose synthase genes using RT-qPCR. In agreement
with the callose deposition patterns, feeding by RSV-free insects evaluated the
expression of callose synthase genes, including *gsl3, gsl4, gsl5,* and *gsl10*, while
RSV-infection further enhanced their upregulation (Fig. 5D–G and S8B-E). These
results further support that both insect feeding and RSV infection promote callose

deposition. Notably, the transcriptional levels of these genes remained unchanged in non-feeding areas (Fig. S8B-E). We also measured the *OsTLP* gene expression and found that neither SBPH feeding nor RSV infection significantly induced *OsTLP* expression (Fig. 5H).

To investigate whether LssaCA influences callose deposition induced by insect-transmitted RSV, we downregulated *LssaCA* in RSV-infected *L. striatellus* using dsRNA microinjection. At 2 dpi, when *LssaCA* expression was significantly reduced, we allowed the insects to feed on healthy plants for 24 h before measuring callose deposition. Our results showed that plants fed upon by *LssaCA*-deficient insects accumulated more callose compared to those fed upon by control insects (dsGFP treatment) (Fig. 5I-K). These results indicate that LssaCA inhibits callose deposition.

In conclusion, these results demonstrate that both *L. striatellus* feeding and RSV infection induce callose deposition, which can be suppressed by LssaCA.

LssaCA mediates in planta *callose degradation by enhancing OsTLP enzymatic activity*

To confirm whether OsTLP degrades the deposited callose induced by insect-transmitted RSV *in planta*, we constructed *ostlp* mutant plants (Fig. S6C). Both ELISA assays and microscopic analysis showed that compared to wild-type plants, *ostlp* plants exhibited significantly higher callose levels (Fig. 6A-B).

Given that LssaCA mediates callose degradation by enhancing OsTLP enzymatic activity, we hypothesized that when OsTLP is knocked out in plant, LssaCA would not affect callose levels induced by RSV infection. To test this, we allowed dsLssaCA-treated viruliferous *L. striatellus* or dsGFP-treated control insects to feed on *ostlp* seedlings. ELISA assays indicated that LssaCA deficiency did not significantly change callose accumulation (Fig. 6C). Microscopic analysis confirmed higher callose deposition in phloem cells of *ostlp* seedlings, which remained unaffected by LssaCA deficiency (Fig. 6D). These results indicate that OsTLP is required *in planta* for LssaCA-mediated effects on callose deposition.

Discussion

Insects are known to be vectors of many plant viruses, and some of these plant viruses rely entirely on insect vectors for their plant-to-plant transmission. Insect salivary proteins have been reported to facilitate viral infection of plants through various mechanisms, such as breaking down plant cell walls, inhibiting plant immune defenses and callose deposition, directly binding to viruses to prevent their detection by plant immune systems, and more. In this study, we found that RSV infection of rice plants triggers the deposition of callose. However, this process is reversed by a tripartite interaction involving the central component of an insect salivary protein LssaCA, which subsequently binds to RSV in insect salivary glands and then OsTLP in the plants. The activity of OsTLP's β-1,3-glucanase is significantly increased when this tripartite interaction occurs; leading to the degradation of callose and thus allowing for efficient RSV infection (Fig. 7).

Callose is a complex polysaccharide, and its deposition in plant phloem plays an important role in defending against insect attacks and viral spread (Zavaliev et al., 2011). When attacked by piercing insects or viruses, plants activate callose deposition on the sieve plates to occlude the flow of phloem sap, thus discouraging feeding and reducing viral spread. This accumulation of callose in plant phloem can be switched on or off through the action of callose synthases and hydrolases (Li et al., 2012; Shangguan et al., 2018). Studies have shown that expression levels of these glucanase enzymes differ between resistant and susceptible plant species. This suggests that some plants may be more successful at resisting infection due to high levels of glucanase enzymes in their tissues, which allow them to more quickly and effectively form callose barriers against insects and viruses (Hao et al., 2008). Some insects have evolved β-1,3-glucanases enzymes that can degrade the deposited callose, thus enabling them to bypass the plant's defense mechanisms and continue feeding (Bucher et al., 2001; Hao et al., 2008). In this study, we demonstrate a novel mechanism by which an insect saliva protein binds to a plant β-1,3-glucanases protein,

activating the enzymatic activity and promoting callose degradation.

In this study, we found that plants fed on by viruliferous insects exhibited higher
callose deposition than those fed on by uninfected insects (Fig. 5A–C), suggesting
that viral infection *via* insect saliva may constitute a dual stress to the plant, inducing
enhanced callose deposition that must be overcome for successful insect feeding and
viral spread. To investigate the underlying mechanism, we examined the tripartite
interaction between LssaCA-NP-OsTLP. We found that this interaction induced
significantly higher β-1,3-glucanase activity than the LssaCA-OsTLP interaction
alone, suggesting that SBPH and RSV have evolved mechanisms to enhance
β-1,3-glucanase activity in order to bypass the callose-mediated defense deposition
and facilitate successful viral spread.

Carbonic anhydrase catalyzes the reversible hydration of carbon dioxide
according to the reaction $CO_2 + H_2O \rightleftharpoons HCO_3^- + H^+$. This enzymatic activity is
essential for maintaining pH homeostasis in biological systems by regulating the
balance between CO_2, bicarbonate, and protons. The importance of carbonic
anhydrases in pH regulation is exemplified in mosquito larvae, which utilize an
extremely alkaline digestive strategy (pH 10.5-11) that is relatively rare in nature,
with carbonic anhydrases serving as the central enzymatic system for generating and
maintaining bicarbonate/carbonate buffering in the midgut lumen (Linser et al., 2009).
Similarly, carbonic anhydrases in arthropod saliva can significantly impact
host-pathogen interactions through pH modulation. A recent study elucidated how
aphid salivary carbonic anhydrase II (CA-II) facilitates virus transmission, with the
enzyme showing significantly higher expression levels in winged morphs compared to
wingless counterparts. When secreted into the plant apoplastic space, CA-II catalyzes
H^+ accumulation, resulting in apoplastic acidification from 5.5 to 5.0. This pH
reduction enhances polygalacturonase activity, thereby promoting the degradation of
demethylesterified homogalacturonan components. Consequently, plants respond by
accelerating vesicle trafficking to transport pectin for cell wall reinforcement, a
process that simultaneously facilitates viral translocation from the endomembrane

system to the apoplastic compartment (Guo et al., 2023). In contrast to these pH-dependent mechanisms, our study demonstrated that LssaCA's biological function in mediating RSV infection is, if not completely, at least partially independent of its enzymatic activity. We found that although OsTLP enzymatic activity is highly pH-dependent and exhibits increased enzymatic activity as pH decreases from 7.5 to 5.0 (Fig. 3H), the LssaCA-OsTLP interaction occurring at pH 7.4 significantly enhanced OsTLP enzymatic activity without any change in buffer pH (Fig. 3G). These results demonstrate the crucial importance of LssaCA-OsTLP protein interactions, rather than enzymatic activity alone, in mediating RSV infection. While the exact role of LssaCA enzymatic activity in aiding RSV infection remains unclear, the local low-pH plant environment would enhance OsTLP enzymatic activity, suggesting potential synergistic effects between enzymatic and non-enzymatic functions.

In conclusion, this study elucidated a tripartite interaction among a plant virus, an insect saliva protein, and a host phloem β-1,3-glucanase. This interaction inhibited the plant defense of callose deposition and enabled viral phloem transmission. This study provides new insights into the tripartite virus-insect vector-plant interaction, which is relevant to many agriculturally important plant arboviruses whose transmission is facilitated *via* insect saliva proteins.

Figure 1.   An *L. striatellus* salivary carbonic anhydrase (LssaCA) directly binds to RSV NP in salivary glands

A. Immunofluorescence showing co-localization of RSV (red) and LssaCA (green) in *L. striatellus* salivary glands. Images represent three independent experiments with 30 SBPHs analyzed. Scale bar: 20 μm. psg, principal salivary gland; asg, accessory salivary gland.

B. Schematic of LssaCA protein sequence. SP, signal peptide; Carb_anhydrase domain, conserved eukaryotic-type carbonic anhydrase sequence. Triangles indicate seven predicted catalytically active residues.

C. Esterase activity of recombinant LssaCA protein expressed in sf9 insect cell and *E.*

coli system. In LssaCA mutant protein, all seven predicted catalytically active residues were replaced. Mean and SD were calculated from 2 independent enzymatic assays. ****, P<0.0001.

D. GST pull-down assays demonstrating interaction between LssaCA (GST-LssaCA) and RSV NP (NP-His). GST served as negative control. Anti-GST and anti-His antibodies detected corresponding proteins.

E. MST assay showing binding between LssaCA (LssaCA-His) and RSV NP (GST-NP). GST served as negative control. Error bars represent SD.

F. Measurement of *LssaCA* tissure-specific expression by RT-qPCR and western blotting. Third-instar RSV-free nymphs were used to measure the gene expression in gut, salivary gland, fatbody, and remaining tissues (R-body) after dissection of the aforementioned tissues. Adult females and adult males were used to measure gene expression in ovaries and testes, respectively. Mean and SD were calculated from three independent experiments with 5 tissue samples each experiment. LssaCA protein was detected using anti-LssaCA antibodies.

G. Western blot analysis of LssaCA protein in RSV-free SBPH salivary glands. dsLssaCA treatment significantly reduces target protein levels compared to controls. LssaCA protein was detected using anti-LssaCA antibodies.

H. Western blot analysis showing LssaCA distribution in watery and gel saliva of RSV-free insects. LssaCA protein was detected using anti-LssaCA antibodies. Total saliva proteins were visualized by silver staining.

I. Western blot detection of LssaCA proteins in rice phloem following *L. striatellus* feeding. LssaCA protein was detected using anti-LssaCA antibodies.

J. Analyses of LssaCA levels regulated by RSV infection. Third-instar RSV-free and RSV-infected nymphs were used to measure gene expression (RT-qPCR) and protein accumulation (Western blotting). Mean and SD were calculated from three independent experiments with totally 14 tissue samples each experiment. ns, not significant.

All western blot images are representative of three independent experiments.

Figure 2. LssaCA enhances RSV infection in rice plants

A. Gene silencing efficiency determined by RT-qPCR.

B-C. Analysis of RSV infection levels in rice seedlings. Each plant seedling was fed upon by 5 infected insects for 2 days, followed by 14 days of cultivation. RSV levels were determined by RT-qPCR (B) and western blotting (C). RSV NP detected by anti-NP polyclonal antibodies indicated viral titers. Rubisco large subunit (RBCL) served as protein loading control.

D. Gene silencing efficiency determined by RT-qPCR.

E. RT-qPCR analysis of RSV titers in salivary glands. Each dot represents salivary glands from 5 RSV-infected third-instar nymphs.

F. RSV titers in *L. striatellus* saliva determined by RT-qPCR. Each dot represents one saliva sample from 10 insects fed with artificial diet.

G. RT-qPCR analysis of RSV titers in rice. Plants were fed upon by infected insects for 24 h (Left: schematic of *L. striatellus* feeding on specific rice stem sites), with titers assayed immediately post-feeding (Right). Each dot represents an individual rice seedling..

H. Western blot analysis of LssaCA in plants fed upon by 50 RSV-free insects for 2 days and continuously cultured for an additional 1 and 3 days. LssaCA protein was detected using anti-LssaCA antibodies. dpf, days post-feeding.

I-J. RSV infection in plants at 3 dpf. Five RSV-infected third-instar *L. striatellus* nymphs were fed upon healthy rice plants for 2 days, and viral titers in whole plant shoots were measured at 3 dpf by RT-qPCR (I) and western blotting (J). Each dot represents an individual plant. RSV NP detected by anti-NP polyclonal antibodies indicated viral titers. RBCL protein served as a protein loading control.

Mean and SD were calculated from 3 independent microinjection experiments. ns, not significant; *, P<0.05; **, P<0.01; ****, P<0.0001. All western blot images are

445 representative of three independent experiments.

Figure 3. LssaCA interacts with rice thaumatin-like protein (OsTLP) to enhance its endo-β-1,3-glucanase activity

A. Schematic diagram of OsTLP protein sequence. SP, signal peptide; THM, conserved thaumatin family protein sequence. Glyco_hydro_64 domain, glycoside hydrolases of family 64.

B. Yeast two-hybrid assay showing interaction between OsTLP and LssaCA. SD, synthetically defined medium; Leu, Leucine; Trp, Tryptophan; His, Histidine; 3'AT, 3-amino-1,2,4-triazole; AD, transcription activation domain; BD, DNA-binding domain.

C. Pull-down assays showing interactions between LssaCA (LssaCA-His) and OsTLP (MBP-OsTLP). MBP served as a negative control. Anti-His and anti-MBP polyclonal antibodies were used for protein detection. Western blot images are representative of three independent experiments.

D. MST assay showing molecular interactions between LssaCA (LssaCA-His) and OsTLP (MBP-OsTLP). MBP served as a negative control. Error bars represent SD.

E. Endo-β-1,3-glucanase activity of purified recombinant OsTLP protein expressed as MBP-fusion protein. MBP was used for normalization.

F. Endo-β-1,3-glucanase activity in OsTLP-overexpressing transgenic plants (OsTLP OE) or wild-type plants (WT). Plant total soluble proteins were used for enzymatic activity assays. Units mg^{-1} means units per mg of plant total soluble proteins.

G. Regulation of OsTLP enzymatic activity by LssaCA binding.

H. OsTLP enzymatic activity measured at pH 5.0, 5.5, 6.0, 6.5, 7.0 and 7.5 to assess pH dependence.

I. RT-qPCR analysis of RSV infection in OsTLP OE and WT plants. RSV titers (NP copy number) were measured at 14 dpf.

ns, not significant; *, P<0.05; ***, P<0.001; ****, P<0.0001.

Figure 4. RSV NP-LssaCA-OsTLP tripartite interaction further enhances OsTLP enzymatic activity

A. Pull-down assays showing interactions between OsTLP (MBP-OsTLP) and LssaCA-NP complex. LssaCA was expressed with a His tag, and RSV NP was expressed as a GST-fusion protein. LssaCA and NP were pre-incubated before co-incubation with OsTLP. Anti-His, anti-GST, and anti-MBP antibodies were used to detect corresponding proteins.

B. Endo-β-1,3-glucanase activity of OsTLP after co-incubation with LssaCA (LssaCA + GST) and LssaCA-NP complex (LssaCA + NP). LssaCA was pre-incubated with either GST or NP before addition to OsTLP. ****, P<0.0001.

Figure 5. LssaCA inhibits *L. striatellus*-transmitted RSV-induced callose deposition in rice plants

A. Callose concentration quantified by ELISA in leaves of rice plants fed on by RSV-free or RSV-infected *L. striatellus*. Non-fed plants served as the control (CK).

B. Bright blue fluorescence of cross-sections showing callose deposition at feeding sites. Samples were prepared from the leaf phloem of plants that were not fed or were fed on by RSV-free or RSV-infected *L. striatellus*. Thin sections were stained with 0.1% aniline blue at 24 h post-feeding and examined under a fluorescence microscope. xy, xylem; ph, phloem. Scale bars: 10 μm.

C. Average fluorescence intensity of callose deposition measured using ImageJ. Eight to ten random sites per sample were selected for intensity quantification.

D–H. Transcript levels of callose synthase genes (*gls3, gls4, gls5, gls10*) and *OsTLP* determined by RT-qPCR. Insects were allowed to feed on rice plants for 24 h prior to RNA extraction.

I. Callose concentration quantified by ELISA in leaves of rice plants fed on by

dsGFP- or dsLssaCA-treated RSV-infected *L. striatellus*.

J. Bright blue fluorescence of cross-sections showing callose deposition at feeding sites. Samples prepared from the leaf phloem of plants fed on by dsGFP- or dsLssaCA-treated RSV-infected *L. striatellus*. Scale bars: 20 μm.

K. Average fluorescence intensity of callose deposition measured using ImageJ. Eight to ten random sites per sample were selected for intensity quantification.

Mean and SD were calculated from three independent experiments. ns, not significant; *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; ****, $p < 0.0001$.

Figure 6. LssaCA mediates *in planta* callose degradation by enhancing OsTLP enzymatic activity

A. Callose concentration quantified by ELISA in leaves of wild-type (WT), *OsTLP* overexpression (OsTLP OE), and *OsTLP* mutant (*ostlp*) plants.

B. Bright blue fluorescence of cross-sections showing callose deposition in leaf phloem. Plants were not fed on by insects. Scale bars: 10 μm. Average fluorescence intensity of callose deposition was measured using ImageJ. Eight to ten random sites per sample were selected for intensity quantification. xy, xylem; ph, phloem.

C. Callose concentration quantified by ELISA in leaves of *ostlp* plants fed on by dsGFP- or dsLssaCA-treated RSV-infected *L. striatellus*.

D. Bright blue fluorescence of cross-sections showing callose deposition at feeding sites. Samples were prepared from the leaf phloem of *ostlp* plants fed on by dsGFP- or dsLssaCA-treated RSV-infected *L. striatellus*. Scale bars: 10 μm. Average fluorescence intensity was quantified using ImageJ. Eight to ten random sites per sample were selected for intensity quantification.

Mean and SD were calculated from three independent experiments. ns, not significant; *, $p < 0.05$. ****, $p < 0.0001$.

Figure 7. Proposed model of RSV NP-LssaCA-OsTLP tripartite interaction facilitating callose degradation and promoting RSV infection.

In *L. striatellus* salivary glands (SG), RSV (red star) binds to LssaCA proteins (blue triangle) via the nucleocapsid protein (NP). Upon delivery into the rice phloem, RSV infection triggers callose synthesis, leading to callose deposition that restricts viral spread. However, LssaCA binds to OsTLP (yellow scissors), enhancing its β-1,3-glucanase activity and promoting callose degradation. The RSV NP-LssaCA complex formed in the insect's SG can still interact with OsTLP *in planta*, resulting in significantly higher OsTLP β-1,3-glucanase activity. This enhanced enzymatic activity efficiently degrades callose, thereby facilitating RSV infection.

Figure S1. Characteristics of LssaCA.

A. Amino acid sequence of LssaCA. The signal peptide is underlined in red. Catalytically active residues are shown in green, and zinc binding sites in blue. The alpha-carbonic anhydrase domain spans residues 91-312.

B. Summary of amino acid substations in the LssaCA mutant. Seven catalytically active residues were replaced with the indicated amino acids to generate an enzymatically inactive mutant.

Figure S2. Immunofluorescence localization of RSV and LssaCA in uninfected *L. striatellus* salivary glands.

RSV (red) is absent in uninfected salivary glands, while LssaCA (green) is detected. Scale bar: 20 μm. Abbreviations: psg, principal salivary gland; asg, accessory salivary gland.

Figure S3. SDS-PAGE analysis of purified recombinant proteins used in Microscale thermophoresis (MST) assays.

Protein bands were visualized by Coomassie Brilliant Blue staining.

Figure S4.   LssaCA does not affect *L. striatellus* feeding behavior.

A-B. Electrical penetration graph (EPG) recordings of *L. striatellus* feeding on rice plants. Waveforms were categorized into non-penetration (NP), penetration (N1, N2, N3), and phloem feeding (N4a & N4b) phases. A, dsGFP-treated insects; B, dsLssaCA-treated insects.

C-D. Quantitative analysis of the time to first non-phloem phase (C) and time to first phloem feeding (D). Mean and SD were calculated from six independent EPG recordings. ns, not significant.

Figure S5. Nucleotide and deduced amino acid sequence of OsTLP

The predicted signal peptide is underlined. Conserved thaumatin-like protein sequences are shown in bold.

Figure S6. Generation of OsTLP-overexpressing and knockout rice lines.

A. Schematic representation of OsTLP overexpression vector construction using the pCAMBIA1300 backbone, driven by CaMV 35S promoter.

B. RT-qPCR and western blot analyses of OsTLP expression in transgenic lines (OsTLP OE) compared to wild type (WT). OsTLP protein was detected using an anti-His antibody.

C. Summary of mutations in *ostlp* knockout lines generated via CRISPR/Cas9. Exons are shown as black boxes, with start (ATG) and stop (TGA) codons labeled. The sgRNA target site is marked with a triangle.

Figure S7. Detection of callose deposition following *L. striatellus* feeding.

Bright blue fluorescence indicates callose deposition at feeding sites in longitudinal leaf cross-sections. Samples were collected from plants that were unfed, fed on by RSV-free insects, or fed on by RSV-infected *L. striatellus*. Observation were made using fluorescence microscopy. Abbreviations: xy, xylem; ph, phloem. Scale bars: 10 μm.

Figure S8. Callose deposition analysis in non-feeding areas

A. Callose concentration quantified by ELISA in leaf areas 1 cm away from insect feeding sites. CK: control (non-fed plants).

B-E. RT-qPCR analysis of callose synthase gene expression in non-fed areas. Insects were allowed to feed for 24 h prior to RNA extraction. ns, not significant; *, $p < 0.05$.

Table S1. Yeast two-hybrid screening for rice proteins interacting with LssaCA.

Table S2. List of primers used in this study.

Materials and methods

Viruses, insects, host plants, and antibodies

The RSV-free and RSV-infected *L. striatellus* individuals used in this study were originally captured in Jiangsu Province, China, and were maintained in our laboratory. All plants used to rear *L. striatellus* were grown in a growth incubator (14.5 cm height, 3.2 cm radius) at 25°C under a 16-h light/8-h dark photoperiod. Insects were transferred to fresh seedlings every 15 days to ensure adequate nutrition and insect activity.

RSV ribonucleoproteins (RNPs) were purified from RSV-infected rice plants. These purified RNPs were subsequently used as the antigen to immunize rabbits for the production of an RSV-specific antibody. The LssaCA polyclonal antibody was produced using recombinantly expressed LssaCA protein (with a His-tag at the C-terminus) as the antigen. Anti-His, anti-GST, and anti-MBP antibodies were purchased from ABclonal.

Tissue collection

Insect dissection and tissue collection were performed using previously described procedures (Huo et al., 2018). In brief, insects were anesthetized at 4 °C for 10 min, and the forelegs were severed at the coxa-trochanter joint using forceps. The hemolymph was expelled and drawn to the tip of clean forceps. The insects were then dissected from the abdomen in pre-chilled PBS buffer. The fat body was collected

with forceps, aided by the tension of the liquid. Tissues, including the guts, salivary glands, and ovaries, were washed three times in PBS to remove contaminants from the hemolymph.

Bioinformatic analysis

The signal peptide and transmembrane helix were predicted using tools at the SignalP 5.0 server (https://services.healthtech.dtu.dk/service.php?SignalP-5.0) and the TMHMM-2.0 server (https://services.healthtech.dtu.dk/service.php?TMHMM-2.0). Homology searches of the protein sequence of LssaCA and OsTLP were conducted using the BLAST algorithm at the National Center for Biotechnology Information (http://blast.ncbi.nlm.nih.gov/Blast.cgi). Conserved domains were predicted with SMART (http://smart.embl-heidelberg.de/).

Collection of watery and gel saliva

Saliva was collected as described previously (Huo et al., 2022). Watery saliva was obtained by centrifuging the collected artificial diet at 10,000 ×g for 10 min to remove the solid gel. To collect gel saliva proteins, the membrane was washed twice with PBS. Sheathes adhering to the parafilm were denatured and solubilized in a solution consisting of 8 M urea, 4% v/v CHAPS, 0.1% w/v sodium dodecyl sulfate (SDS), and 2% v/v DTT. The parafilm was incubated on an orbital shaker at room temperature for 45 min. Then, the denatured gel saliva was concentrated to a final volume of 100 μL. Saliva samples were collected from 2,000 insects for silver staining and western blotting analyses.

Measurement of carbonic anhydrase activity

DNA fragments encoding full-length LssaCA and a LssaCA mutant (H111D, N139H, H141D, H143D, E153H, H166D, and T253E) were amplified by PCR. The resulting products were purified and cloned into the vector pET-30a(+) and pFastBac1 vectors to generate His-tagged recombinant proteins. The pET-30a(+)-LssaCA and pET-30a(+)-LssaCA(mutant) plasmids were transformed into *E. coli* for protein

expression. In parallel, the pFastBac1-LssaCA and the pFastBac1-LssaCA(mutant)

plasmids were transformed into *E. coli* DH10Bac for transposition into the bacmid.

Recombinant bacmids were then transfected into *Spodoptera frugiperda* (Sf9) cells

using Cellfectin II Reagent (Invitrogen, USA). Recombinant proteins expressed in

both systems were purified using Ni Sepharose resin (GE Healthcare, Uppsala,

Sweden).

Carbonic anhydrase activity was assessed by measuring esterase activity using

p-nitroethyl acetate (*p*-NPA) as the substrate. LssaCA and its mutant were expressed

in *E. coli* and Sf9 cells with C-terminal His-tag, and the purified proteins were diluted

in PBS buffer (pH 7.0) to a final concentration of 1 mg/mL. Reaction conditions

followed the method described by Verpoorte et al. (1967), with minor modifications.

Briefly, 1 mL of freshly prepared 3 mM *p*-NPA (dissolved in acetone) was added to

1.9 mL of 100 mM PBS (pH 7.0), followed by 0.1 mL of LssaCA enzyme solution

(0.5 mg/mL in 100 mM PBS). Absorbance at 384 nm was measured at the start of the

reaction and after 3 minutes. One unit (U) of enzyme activity was defined as the

amount of enzyme required to hydrolyze 1 μmol of *p*-NPA per minute at pH 7.0 and

25 ℃. A blank control consisted of 2 mL of 100 mM PBS (pH 7.0) mixed with 1 mL

of 3 mM *p*-NPA.

Immunofluorescence assays

Immunofluorescence assays (IFA) were performed to determine the localization

of RSV and LssaCA in *L. striatellus* salivary glands. The dissected salivary gland

samples were fixed in 4% v/v paraformaldehyde at 4 ℃ overnight and then

permeabilized in 2% v/v Triton X-100 for 2 h. The samples were incubated with

rabbit anti-RSV serum and mouse anti-LssaCA serum (1:250 dilution in PBST/FBS:

PBS containing 2% v/v Triton X-100 and 2% fetal bovine serum) for 1 h at room

temperature. After washing twice with PBST, the samples were incubated with 1:200

diluted Alexa Fluor 488-conjugated goat anti-mouse antibody and Alexa Fluor

594-conjugated goat anti-rabbit antibody (Invitrogen, USA) for 1 h. After washing

three times with PBST, the samples were examined under a Leica TCS SP8

microscope (Leica, Wetzlar, Germany) with the following settings: 1) Alexa Fluor 488 channel: excitation at 488 nm using an argon laser; emission detected with a 493-553 nm bandpass filter. Alexa Fluor 594 channel: excitation at 594 nm using a HeNe laser; excitation detected with a 600-650 nm bandpass filter.

Evaluation of callose deposition in rice plants

Five insects were allowed to feed for 24 h on a 2×2 cm leaf area of 10-d-old rice plants. After removal of the insects, leaf tissues were collected from both the feeding site and an adjacent undamaged area located 1 cm away (non-fed site). Samples were immersed in 5 mL alcoholic lactophenol solution (1 volume of phenol:glycerol:lactic acid:water in a 1:1:1:1 ratio, mixed with 2 volumes of ethanol), evacuated with a syringe for 15 min, and then incubated at 65°C for 30 min until the chlorophyll was completely cleared. Cleared leaf samples were rinsed with 50% v/v ethanol, then with water, and then stained in 150 mM K_2HPO_4 (pH 9.5) containing 0.01% w/v aniline blue for 30 min in the dark. Stained leaf samples were sectioned and mounted in 50% v/v glycerol on slides. The slides were examined under a Zeiss observer Z1 immunofluorescence microscope (Carl Zeiss MicroImaging GmbH, Gottingen, Germany) with excitation at 405 nm. Fluorescence intensity was quantified using ImageJ software. For ImageJ analysis, images were converted to 8-bit grayscale and adjusted for brightness. Phloem-specific regions of interest (ROIs) were outlined using the freehand selection tools. A standardized fluorescence threshold (set at $>2\times$ the mean background intensity) was applied to isolate callose signals. Integrated density and area values were used to calculate normalized fluorescence intensity. Identical exposure times were maintained across all independent experiments.

Callose concentrations were quantified using a plant callose ELISA kit (Catalog No. JL46407; Jonln, Shanghai, China), following the manufacturer's instruction. Leaf tissues from feeding and non-fed sites were homogenized in PBS buffer (pH 7.4), and the homogenate was centrifuged at 500–1000 x g for 20 min. The supernatant was collected, and absorbance at 450 nm was measured. Callose concentrations were

684   calculated using a standard curve.

RNA interference

We conducted RNAi analyses to determine the function of LssaCA in mediating

RSV infection and regulating callose deposition. The *LssaCA*-specific gene fragment

was PCR-amplified with the primer pair LssaCA-T7F/LssaCA-T7R (Table S1). Then,

dsRNA was synthesized using a commercial kit (T7 RiboMAX Express RNAi System,

Promega, Madison, WI, USA) and purified by phenol: chloroform extraction and

isopropanol precipitation. Finally, 36.8 nL dsRNA at 1 ng/nL was delivered into the

insect hemocoel for gene silencing. Microinjection was performed using a glass

needle through a Nanoliter 2000 (World Precision Instruments, Sarasota, FL, USA).

The negative control, GFP dsRNA, was synthesized and microinjected following the

same protocol.

RT-qPCR

Gene transcript levels were determined by RT-qPCR. The RNA was extracted

from *L. striatellus* tissues using a RNeasy Mini Kit (Qiagen, Hilden, Germany), and

from plant leaves using TRIzol. Reverse transcriptional PCR (iScript cDNA Synthesis

Kit, Bio-Rad, Hercules, CA, USA) and SYBR-Green-based qPCR (SYBR Green

Supermix, Bio-Rad) was performed according to the manufacturer's protocols. The

primer pair used to amplify *LssaCA* was LssaCA-q-F/LssaCA-q-R (Table S1). Viral

RNA copies were measured using the primers NP-q-F/NP-q-R (Table S1). *L.*

striatellus translation elongation factor 2 (*EF2*) and rice *actin* were amplified as

internal controls to ensure equal loading of cDNA isolated from different samples.

The primer pairs EF2-q-F/EF2-q-R and actin-q-F/actin-q-R (Table S1) were used to

amplify *EF2* and *actin*, respectively. Water was used as a negative control.

Yeast two-hybrid assays

709   Yeast two-hybrid screening was performed to identify rice proteins interacting

710   with LssaCA. A high-complexity rice cDNA library was fused to *Gal4 AD* and

transformed into the yeast strain Y187. The *LssaCA* gene lacking the signal-peptide sequence (*LssaCA*[sp-]) was fused to *Gal4 DNA-BD* and transformed into the yeast strain Y2HGold. The two yeast strains were co-cultured overnight to allow mating to create diploids. After library screening, positive clones were selected on quadruple dropout medium: SD/–Ade/–His/–Leu/–Trp supplemented with X-a-Gal and aureobasidin A (QDO/X/A), and prey plasmids were isolated for sequencing. To further confirm the interaction between LssaCA and OsTLP, *LssaCA*[sp-] was cloned into the prey vector pDB-Leu, and the *OsTLP* gene lacking the signal-peptide sequence (*OsTLP*[sp-]) was cloned into the bait vector pPC86. The two plasmids were co-transformed into the yeast stain Mav203 and the transformants were selected on SD/–His–Leu–Trp + 20 mM 3-AT medium.

Pull-down assays

Pull-down assays were performed to determine the molecular interactions between LssaCA and RSV NP, LssaCA and OsTLP, and OsTLP and the LssaCA-NP complex. The interaction between LssaCA and RSV NP was determined using GST pull-down analyses as described previously (Huo et al., 2014). Briefly, *LssaCA*[sp-] was cloned into the pGEX-3x vector and LssaCA was expressed as a GST-fusion protein (GST-LssaCA). The *NP* gene was cloned into the pET-30a (+) expression vector and the protein was expressed with an N-terminal His-tag (NP-His). The GST-LssaCA construct was expressed in, and purified from, *E. coli* stain BL21. The GST-LssaCA protein or the GST control protein was bound to Glutathione Sepharose beads (Cytiva, Marlborough, MA, USA), then NP-His protein was added to the beads. The bead-bound proteins were separated by SDS-PAGE and detected by western blotting with antibodies against GST or His-tag.

Next, MBP pull-down analyses were performed to determine the interaction between OsTLP and LssaCA. The *OsTLP*[sp-] sequence was cloned into pMAL-p2X to produce the MBP-fusion protein (MBP-OsTLP). The MBP-OsTLP protein or the MBP control protein was bound to amylose resin (New England Biolabs, Beverley,

MA, USA), and then the LssaCA-His protein was added to the beads. To determine

the interaction between OsTLP and the LssaCA-NP complex, the LssaCA-His and

GST-NP proteins were preincubated before being added to the beads. The bead-bound

proteins were separated by SDS-PAGE and detected by western blotting with the

MBP antibody (Cell Signaling Technology, Danvers, MA, USA), His-tag antibody

(Tiangen, Beijing, China) and GST antibody (Tiangen).

Microscale thermophoresis (MST) assays

We conducted MST assays to detect the interactions between LssaCA and NP,

and between LssaCA and OsTLP. First, 10 μM purified LssaCA^{sp-} protein (His-fusion

protein) was labelled with a Monolith NT Protein Labeling Kit RED-NHS (Nano

Temper Technologies GMBH, München, Germany) using red fluorescent dye NT-647

N-hydroxysuccinimide (amine-reactive), according to the manufacturer's instructions.

The binding assays were performed on a Monolith NT.115 Microscale

Thermophoresis instrument (Nano Temper Technologies GMBH) using capillaries

treated using the standard method. The labeled protein LssaCA^{sp-} was added to

serially diluted NP (GST-NP, GST was used as a negative control) or OsTLP^{sp-}

(MBP-fusion protein, MBP was used as a negative control). The initial concentration

of each protein was 40 μM with 0.1% (v/v) Tween 20. The KD Fit function of

NanoTemper Analysis software version 1.2.214.1 was used for curve fitting and to

calculate the dissociation constant (Kd).

β-1,3-glucanase activity assays

β-1,3-glucanase hydrolyzes laminarin by cleaving β-1,3-glucoside bond, thereby

generating sugar termini. Enzyme activity was quantified by measuring the rate of

reducing sugars production. One unit (U) of β-1,3-glucanase activity was defined as

the amount of enzyme that hydrolyses laminarin to generate 1 μg reducing sugars per

minute.

The β-1,3-glucanase activity of OsTLP was measured using a beta-1,3-glucanase

microplate assay kit (Cat. ASK1028, Bioworld, Bloomington, MN, USA) according

to the manufacturer's instructions. To determine the enzymatic activity of

recombinant OsTLP proteins, MBP-fused OsTLP^{sp-} protein and MBP control protein

were adjusted to 0.5 mg/mL. The MBP-OsTLP^{sp-} fusion protein was equilibrated in

buffers of varying pH for the assay. To determine the enzymatic activity of OsTLP

expressed *in planta*, 0.1 g of rice leaf tissue from either OsTLP over-expressing or

wild-type plants was homogenized with 1 mL of assay buffer on ice. The homogenate

was centrifuged at 12,000g at 4 ℃ for 10 min, and the resulting supernatant was

transferred into a new tube and kept on ice until analysis.

For the assay, 50 μL of the protein solution was mixed with laminarin substrate

and incubated at 37 ℃ for 30 min. The reaction was terminated by incubating the

reaction tube in a boiling water bath for 10 min. The reaction supernatant was

transferred into a microplate and incubated at 90 ℃ for 10 min before measuring the

absorbance at 540 nm.

Generation of transgenic plants

The full-length *OsTLP* gene was PCR-amplified with the primer pair

TLP-F/TLP-R (Table S2) and cloned into the plant expression vector pCAMBIA1300

to produce the plasmid pCAMBIA1300-OsTLP. This construct was transformed into

seedlings of the *Japonica* rice cultivar Nipponbare through *Agrobacterium*-mediated

transformation, following the protocol described by Hiei et al. (1994). The *ostlp*

knockout line was generated using CRISPR-Cas9 technology according to established

protocols. Thirty independent T_0 transformants were obtained and screened by

germinating seeds from T_2 lines on medium containing 50 μg/ml hygromycin.

Hygromycin-resistant homozygous lines were selected for further experiments.

Knockout events were confirmed by sequencing with specific primers (Fig S6C).

Seeds from homozygous lines transformed with the vector pCAMBIA1300-eGFP

were used as negative control.

Statistical analyses

Graphs were generated, and statistical analyses were conducted using Prism 9.0

software (GraphPad Software, San Diego, CA, USA). Data are presented as mean ± standard deviation (SD) from two or three independent experiments. Statistical significance between two groups was assessed using an unpaired Student's t-test, while comparisons among three or more groups were performed using one-way analysis of variance (ANOVA).

Acknowledgments

This work is supported by the National Key R&D Program of China（2022YFD1400800）, Major Program of the National Natural Science Foundation of China (32090013) and Youth Innovation Promotion Association CAS (2021084).

Author contributions

Conceptualization, Z.L.L., H.Y., Z.J.; Methodology, H.Y., Z.J. and Y.J.; Investigation, Z.J., Y.J., M.X.Y., H.Y.; Writing-Original Draft, H.Y. and Z.J.; Writing-Review & Editing, Z.L.L. and H.Y.; Funding Acquisition, Z.L.L., H.Y. and F.R.X.; Supervision, Z.L.L., and F.R.X.

Declarations of competing interests
The authors declare no competing interests.

References

Acevedo, F.E., Smith, P., Peiffer, M., Helms, A., Tooker, J., and Felton, G.W. (2019). Phytohormones in fall armyworm saliva modulate defense responses in plants. J. Chem. Ecol. 45, 598-609. 10.1007/s10886-019-01079-z.

Arca, B., and Ribeiro, J.M. (2018). Saliva of hematophagous insects: a multifaceted toolkit. Curr. Opin. Insect Sci. 29, 102-109. 10.1016/j.cois.2018.07.012.

Bonaventure, G. (2012). Perception of insect feeding by plants. Plant Biol. (Stuttg.) 14, 872-880. 10.1111/j.1438-8677.2012.00650.x.

Bucher, G.L., Tarina, C., Heinlein, M., Serio, F.D., Jr, F.M., and Iglesias, V.A. (2001). Local expression of enzymatically active class I β-1, 3-glucanase enhances symptoms of TMV infection in tobacco. Plant J. 28, 361-369. 10.1046/j.1365-313x.2001.01181.x.

Conway, M.J., Londono-Renteria, B., Troupin, A., Watson, A.M., Klimstra, W.B., Fikrig, E., and Colpitts, T.M. (2016). *Aedes aegypti* D7 saliva protein inhibits *Dengue virus* infection. PLoS Negl. Trop. Dis. 10, e0004941. 10.1371/journal.pntd.0004941.

de Jesus-Pires, C., Ferreira-Neto, J.R.C., Pacifico Bezerra-Neto, J., Kido, E.A., de Oliveira Silva, R.L., Pandolfi, V., Wanderley-Nogueira, A.C., Binneck, E., da Costa, A.F., Pio-Ribeiro, G., et al. (2020). Plant thaumatin-like proteins: function, evolution and biotechnological applications. Curr. Protein Pept. Sci. 21, 36-51. 10.2174/1389203720666190318164905.

Guo, H.J., Zhang, Y.J., Li, B.Y., Li, C.W., Shi, Q.Y., Salzman, K.Z., Ge, F., and Sun, Y.C. (2023). Salivary carbonic anhydrase II in winged aphid morph facilitates plant infection by viruses. Proc Nat Acad Sci USA 120(14), e2222040120. doi:10.1073/pnas.2222040120.

Hao, P.Y., Liu, C.X., Wang, Y.Y., Chen, R.Z., Tang, M., Du, B., Zhu, L.L., and He, G.C. (2008). Herbivore-induced callose deposition on the sieve plates of rice: an important mechanism for host resistance. Plant Physiol. 146, 1810-1820. 10.1104/pp.107.111484.

Hermance, M.E., and Thangamani, S. (2015). Tick saliva enhances *Powassan virus* transmission to the host, influencing its dissemination and the course of disease. J. Virol. 89, 7852-7860. 10.1128/JVI.01056-15.

Hiei, Y., Ohta, S., Komari, T., and Kumashiro, T. (1994). Efficient transformation of rice (*Oryza sativa L.*) mediated by *Agrobacterium* and sequence analysis of the boundaries of the T-DNA. Plant J. 6, 271-282. 10.1046/j.1365-313x.1994.6020271.x.

Hipper, C., Brault, V., Ziegler-Graff, V., and Revers, F. (2013). Viral and cellular

factors involved in Phloem transport of plant viruses. Front. Plant Sci. 4, 154. 10.3389/fpls.2013.00154.

Huang, H.J., Liu, C.W., Xu, H.J., Bao, Y.Y., and Zhang, C.X. (2017). Mucin-like protein, a saliva component involved in brown planthopper virulence and host adaptation. J. Insect Physiol. 98, 223-230. 10.1016/j.jinsphys.2017.01.012.

Huo, Y., Zhao, J., Meng, X.Y., Yang, J., Zhang, Z.Y., Liu, Z.W., Fang, R.X., and Zhang, L.L. (2022). *Laodelphax striatellus* saliva mucin enables the formation of stylet sheaths to facilitate its feeding and *rice stripe virus* transmission. Pest Manag. Sci. 78(8), 3498-3507. 10.1002/ps.6990.

Ji, R., Ye, W.F., Chen, H.D., Zeng, J.M., Li, H., Yu, H.X., Li, J.C., and Lou, Y.G. (2017). A salivary endo-beta-1,4-glucanase acts as an effector that enables the Brown Planthopper to feed on rice. Plant Physiol. 173, 1920-1932. 10.1104/pp.16.01493.

Linser, P.J., Smith, K.E., Seron, T.J., and Oviedo, N.M. (2009). Carbonic anhydrases and anion transport in mosquito midgut pH regulation. J Exp Biol. 212(Pt 11):1662-1671. doi:10.1242/jeb.028084.

Lou, Y.H., Shen, Y., Li, D.T., Huang, H.J., Lu, J.B., and Zhang, C.X. (2019). A mucin-like protein is essential for oviposition in *Nilaparvata lugens*. Front. Physiol. 10, 551. 10.3389/fphys.2019.00551.

Ramamoorthi, N., Narasimhan, S., Pal, U., Bao, F., Yang, X.F., Fish, D., Anguita, J., Norgard, M.V., Kantor, F.S., Anderson, J.F., et al. (2005). The Lyme disease agent exploits a tick protein to infect the mammalian host. Nature 436, 573-577. 10.1038/nature03812.

Schuijt, T.J., Hovius, J.W., van Burgel, N.D., Ramamoorthi, N., Fikrig, E., and van Dam, A.P. (2008). The tick salivary protein Salp15 inhibits the killing of serum-sensitive *Borrelia burgdorferi* sensu lato isolates. Infect. Immun. 76, 2888-2894. 10.1128/IAI.00232-08.

Shangguan, X.X, Zhang, J., Liu, B.F., Zhao, Y., Wang, H.Y., Wang, Z.Z., Guo, J.P., Rao, W.W., Jing, S.L., Guan, W., et al. (2018). A mucin-Like protein of planthopper is required for feeding and induces immunity response in plants. Plant

Physiol. 176, 552-565. 10.1104/pp.17.00755.

Sun, P., Nie, K.X., Zhu, Y.B., Liu, Y., Wu, P., Liu, Z.W., Du, S.Y., Fan, H.H., Chen, C.H., Zhang, R.L., et al. (2020). A mosquito salivary protein promotes flavivirus transmission by activation of autophagy. Nat. Commun. 11, 260. 10.1038/s41467-019-14115-z.

Tjallingii, W.F. (2006). Salivary secretions by aphids interacting with proteins of phloem wound responses. J. Exp. Bot. 57, 739-745. 10.1093/jxb/erj088.

van Bel, A.J., and Will, T. (2016). Functional evaluation of proteins in watery and gel saliva of aphids. Front. Plant Sci. 7, 1840. 10.3389/fpls.2016.01840.

Verpoorte, J.A., Mehta, S., and Edsall, J.T. (1967). Esterase activities of human carbonic anhydrases B and C. J. Biol. Chem. 242, 4221-4229. 10.1016/s0021-9258(18)95800-x.

Wang, X.W., and Blanc, S. (2021). Insect transmission of plant single-stranded DNA viruses. Annu. Rev. Entomol. 66, 389-405. 10.1146/annurev-ento-060920-094531.

Wen, S.Y., Wang, F., Ji, Z.H., Pan, Y.Y., Jian, M.M, Bi, Y.F., Zhou, G.Z., Luo, L.S., Chen, T.G., Li, L.B., et al. (2019). Salp15, a multifunctional protein from tick saliva with potential pharmaceutical effects. Front. Immunol. 10, 3067. 10.3389/fimmu.2019.03067.

Will, T., and Vilcinskas, A. (2015). The structural sheath protein of aphids is required for phloem feeding. Insect. Biochem. Mol. Biol. 57, 34-40. 10.1016/j.ibmb.2014.12.005.

Wu, W., Zheng, L., Chen, H., Jia, D., Li, F., and Wei, T. (2014). Nonstructural protein NS4 of *rice stripe virus* plays a critical role in viral spread in the body of vector insects. PLoS One 9(2), e88636. 10.1371/journal.pone.0088636.

Wu, W., Yi, G., Lv, X.W., Mao, Q.Z., and Wei, T.Y. (2022). A leafhopper saliva protein mediates horizontal transmission of viral pathogens from insect vectors into rice phloem. Commun. Biol. 5, 204. 10.1038/s42003-022-03160-y.

Xu, Y., Fu, S., Tao, X.R, and Zhou, X. P. (2021). *Rice stripe virus*: exploring

molecular weapons in the arsenal of a negative-sense RNA virus. Annu. Rev.

Phytopathol. 59, 351-371. 10.1146/annurev-phyto-020620-113020.

Yi, G., Wu, W., and Wei, T.Y. (2021). Delivery of *rice gall dwarf virus* into plant

phloem by its leafhopper vectors activates callose deposition to enhance viral

transmission. Front. Microbiol. 12, 662577. 10.3389/fmicb.2021.662577.

Yue, N., Jiang, Z., Zhang, X., Li, Z., Wang, X., Wen, Z., Gao, Z., Pi, Q., Zhang,

Y., Wang, X. B., Han, C., Yu, J., and Li, D. (2022). Palmitoylation of γb protein

directs a dynamic switch between Barley stripe mosaic virus replication and

movement. EMBO J. 41(13), e110060. 10.15252/embj.2021110060

Zavaliev, R., Ueki, S., Epel, B.L., and Citovsky, V. (2011). Biology of callose

(beta-1,3-glucan) turnover at plasmodesmata. Protoplasma 248, 117-130.

10.1007/s00709-010-0247-0.

Zhao, W., Kang, L., and Cui, F. (2017). Isolation of *rice stripe virus* preparation

from viruliferous Small Brown Planthoppers and mechanic inoculation on rice. Bio.

Protoc. 7, e2597. 10.21769/BioProtoc.2597.

Figure 1

Figure 1. An *L. striatellus* salivary carbonic anhydrase (LssaCA) directly binds to RSV NP in salivary glands

A.Immunofluorescence showing co-localization of RSV (red) and LssaCA (green) in *L. striatellus* salivary glands. Images represent three independent experiments with 30 SBPHs analyzed. Scale bar: 20 μm. psg, principal salivary gland; asg, accessory salivary gland.

B.Schematic of LssaCA protein sequence. SP, signal peptide; Carb_anhydrase domain, conserved eukaryotic-type carbonic anhydrase sequence. Triangles indicate seven predicted catalytically active residues.

C.Esterase activity of recombinant LssaCA protein expressed in sf9 insect cell and *E. coli* system. In LssaCA mutant protein, all seven predicted catalytically active residues were replaced. Mean and SD were calculated from 2 independent enzymatic assays. ****, P<0.0001.

D.GST pull-down assays demonstrating interaction between LssaCA (GST-LssaCA) and RSV NP (NP-His). GST served as negative control. Anti-GST and anti-His antibodies detected corresponding proteins.

E.MST assay showing binding between LssaCA (LssaCA-His) and RSV NP (GST-NP). GST served as negative control. Error bars represent SD.

F.Measurement of *LssaCA* tissure-specific expression by RT-qPCR and western blotting. Third-instar RSV-free nymphs were used to measure the gene expression in gut, salivary gland, fatbody, and remaining tissues (R-body) after dissection of the aforementioned tissues. Adult females and adult males were used to measure gene expression in ovaries and testes, respectively. Mean and SD were calculated from three independent experiments with 5 tissue samples each experiment. LssaCA protein was detected using anti-LssaCA antibodies.

G.Western blot analysis of LssaCA protein in RSV-free SBPH salivary glands. dsLssaCA treatment significantly reduces target protein levels compared to controls. LssaCA protein was detected using anti-LssaCA antibodies.

H.Western blot analysis showing LssaCA distribution in watery and gel saliva of RSV-free insects. LssaCA protein was detected using anti-LssaCA antibodies. Total saliva proteins were visualized by silver staining.

I.Western blot detection of LssaCA proteins in rice phloem following *L. striatellus* feeding. LssaCA protein was detected using anti-LssaCA antibodies.

J.Analyses of LssaCA levels regulated by RSV infection. Third-instar RSV-free and RSV-infected nymphs were used to measure gene expression (RT-qPCR) and protein accumulation (Western blotting). Mean and SD were calculated from three independent experiments with totally 14 tissue samples each experiment. ns, not significant.

All western blot images are representative of three independent experiments.

Figure 2

Figure 2. LssaCA enhances RSV infection in rice plants

A.Gene silencing efficiency determined by RT-qPCR.

B-C.Analysis of RSV infection levels in rice seedlings. Each plant seedling was fed upon by 5 infected insects for 2 days, followed by 14 days of cultivation. RSV levels were determined by RT-qPCR (B) and western blotting (C). RSV NP detected by anti-NP polyclonal antibodies indicated viral titers. Rubisco large subunit (RBCL) served as protein loading control.

D.Gene silencing efficiency determined by RT-qPCR.

E.RT-qPCR analysis of RSV titers in salivary glands. Each dot represents salivary glands from 5 RSV-infected third-instar nymphs.

F.RSV titers in *L. striatellus* saliva determined by RT-qPCR. Each dot represents one saliva sample from 10 insects fed with artificial diet.

G.RT-qPCR analysis of RSV titers in rice. Plants were fed upon by infected insects for 24 h (Left: schematic of *L. striatellus* feeding on specific rice stem sites), with titers assayed immediately post-feeding (Right). Each dot represents an individual rice seedling..

H.Western blot analysis of LssaCA in plants fed upon by 50 RSV-free insects for 2 days and continuously cultured for an additional 1 and 3 days. LssaCA protein was detected using anti-LssaCA antibodies. dpf, days post-feeding.

I-J.RSV infection in plants at 3 dpf. Five RSV-infected third-instar *L. striatellus* nymphs were fed upon healthy rice plants for 2 days, and viral titers in whole plant shoots were measured at 3 dpf by RT-qPCR (I) and western blotting (J). Each dot represents an individual plant. RSV NP detected by anti-NP polyclonal antibodies indicated viral titers. RBCL protein served as a protein loading control.

Mean and SD were calculated from 3 independent microinjection experiments. ns, not significant; *, P<0.05; **, P<0.01; ****, P<0.0001. All western blot images are representative of three independent experiments.

Figure 3

Figure 3. LssaCA interacts with rice thaumatin-like protein (OsTLP) to enhance its endo-β-1,3-glucanase activity

A.Schematic diagram of OsTLP protein sequence. SP, signal peptide; THM, conserved thaumatin family protein sequence. Glyco_hydro_64 domain, glycoside hydrolases of family 64.

B.Yeast two-hybrid assay showing interaction between OsTLP and LssaCA. SD, synthetically defined medium; Leu, Leucine; Trp, Tryptophan; His, Histidine; 3'AT, 3-amino-1,2,4-triazole; AD, transcription activation domain; BD, DNA-binding domain.

C.Pull-down assays showing interactions between LssaCA (LssaCA-His) and OsTLP (MBP-OsTLP). MBP served as a negative control. Anti-His and anti-MBP polyclonal antibodies were used for protein detection. Western blot images are representative of three independent experiments.

D.MST assay showing molecular interactions between LssaCA (LssaCA-His) and OsTLP (MBP-OsTLP). MBP served as a negative control. Error bars represent SD.

E.Endo-β-1,3-glucanase activity of purified recombinant OsTLP protein expressed as MBP-fusion protein. MBP was used for normalization.

F.Endo-β-1,3-glucanase activity in OsTLP-overexpressing transgenic plants (OsTLP OE) or wild-type plants (WT). Plant total soluble proteins were used for enzymatic activity assays. Units mg^{-1} means units per mg of plant total soluble proteins.

G.Regulation of OsTLP enzymatic activity by LssaCA binding.

H.OsTLP enzymatic activity measured at pH 5.0, 5.5, 6.0, 6.5, 7.0 and 7.5 to assess pH dependence.

I.RT-qPCR analysis of RSV infection in OsTLP OE and WT plants. RSV titers (NP copy number) were measured at 14 dpf.

ns, not significant; *, P<0.05; ***, P<0.001; ****, P<0.0001.

Figure 4

A

B

Figure 4. RSV NP-LssaCA-OsTLP tripartite interaction further enhances OsTLP enzymatic activity

A.Pull-down assays showing interactions between OsTLP (MBP-OsTLP) and LssaCA-NP complex. LssaCA was expressed with a His tag, and RSV NP was expressed as a GST-fusion protein. LssaCA and NP were pre-incubated before co-incubation with OsTLP. Anti-His, anti-GST, and anti-MBP antibodies were used to detect corresponding proteins.

B.Endo-β-1,3-glucanase activity of OsTLP after co-incubation with LssaCA (LssaCA + GST) and LssaCA-NP complex (LssaCA + NP). LssaCA was pre-incubated with either GST or NP before addition to OsTLP. ****, P<0.0001.

Figure 5

Figure 5. LssaCA inhibits *L. striatellus*-transmitted RSV-induced callose deposition in rice plants

A.Callose concentration quantified by ELISA in leaves of rice plants fed on by RSV-free or RSV-infected *L. striatellus*. Non-fed plants served as the control (CK).

B.Bright blue fluorescence of cross-sections showing callose deposition at feeding sites. Samples were prepared from the leaf phloem of plants that were not fed or were fed on by RSV-free or RSV-infected *L. striatellus*. Thin sections were stained with 0.1% aniline blue at 24 h post-feeding and examined under a fluorescence microscope. xy, xylem; ph, phloem. Scale bars: 10 μm.

C.Average fluorescence intensity of callose deposition measured using ImageJ. Eight to ten random sites per sample were selected for intensity quantification.

D–H. Transcript levels of callose synthase genes (*gls3, gls4, gls5, gls10*) and *OsTLP* determined by RT-qPCR. Insects were allowed to feed on rice plants for 24 h prior to RNA extraction.

I. Callose concentration quantified by ELISA in leaves of rice plants fed on by dsGFP- or dsLssaCA-treated RSV-infected *L. striatellus*.

J. Bright blue fluorescence of cross-sections showing callose deposition at feeding sites. Samples prepared from the leaf phloem of plants fed on by dsGFP- or dsLssaCA-treated RSV-infected *L. striatellus*. Scale bars: 20 μm.

K. Average fluorescence intensity of callose deposition measured using ImageJ. Eight to ten random sites per sample were selected for intensity quantification.

Mean and SD were calculated from three independent experiments. ns, not significant; *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; ****, $p < 0.0001$.

Figure 6

Figure 6. LssaCA mediates *in planta* callose degradation by enhancing OsTLP enzymatic activity

A. Callose concentration quantified by ELISA in leaves of wild-type (WT), *OsTLP* overexpression (OsTLP OE), and *OsTLP* mutant (*ostlp*) plants.

B. Bright blue fluorescence of cross-sections showing callose deposition in leaf phloem. Plants were not fed on by insects. Scale bars: 10 μm. Average fluorescence intensity of callose deposition was measured using ImageJ. Eight to ten random sites per sample were selected for intensity quantification. xy, xylem; ph, phloem.

C. Callose concentration quantified by ELISA in leaves of *ostlp* plants fed on by dsGFP- or dsLssaCA-treated RSV-infected *L. striatellus*.

D. Bright blue fluorescence of cross-sections showing callose deposition at feeding sites. Samples were prepared from the leaf phloem of *ostlp* plants fed on by dsGFP- or dsLssaCA-treated RSV-infected *L. striatellus*. Scale bars: 10 μm. Average fluorescence intensity was quantified using ImageJ. Eight to ten random sites per sample were selected for intensity quantification.

Mean and SD were calculated from three independent experiments. ns, not significant; *, $p < 0.05$. ****, $p < 0.0001$.

Figure 7

Figure 7. Proposed model of RSV NP-LssaCA-OsTLP tripartite interaction facilitating callose degradation and promoting RSV infection.

In *L. striatellus* salivary glands (SG), RSV (red star) binds to LssaCA proteins (blue triangle) via the nucleocapsid protein (NP). Upon delivery into the rice phloem, RSV infection triggers callose synthesis, leading to callose deposition that restricts viral spread. However, LssaCA binds to OsTLP (yellow scissors), enhancing its β-1,3-glucanase activity and promoting callose degradation. The RSV NP-LssaCA complex formed in the insect's SG can still interact with OsTLP *in planta*, resulting in significantly higher OsTLP β-1,3-glucanase activity. This enhanced enzymatic activity efficiently degrades callose, thereby facilitating RSV infection.