Sequential effect in pre-stimulus oculomotor inhibition

Noam Tal-Perry[1,2] & Shlomit Yuval-Greenberg[1, 3]

[1]School of Psychological Sciences, Tel Aviv University, Tel Aviv, Israel

[2]Department of Experimental Psychology, University of Oxford, Oxford, United Kingdom

[3]Sagol School of Neuroscience, Tel Aviv University, Tel Aviv, Israel

Noam Tal-Perry https://orcid.org/0000-0003-2521-9546

Shlomit Yuval-Greenberg https://orcid.org/0000-0001-6455-7578

Correspondence concerning this article should be addressed to Shlomit Yuval-Greenberg, School of Psychological Sciences and Sagol School of Neuroscience, Tel-Aviv University, P.O. box 39040, Tel Aviv 6997801. Email: shlomitgr@tau.ac.il. This study was funded by the Israel Science Foundation grant 1960/19 to S.Y-G.

Abstract

When faced with unfamiliar circumstances, we often turn to our past experiences with similar situations to shape our expectations. This results in the well-established sequential effect, in which previous trials influence the expectations of the current trial. Studies have revealed that, in addition to the classical behavioral metrics, the inhibition of eye movement could be used as a biomarker to study temporal expectations. This pre-stimulus oculomotor inhibition is found a few hundred milliseconds prior to predictable events, with a stronger inhibition for predictable than unpredictable events. The phenomenon has been found to occur in various temporal structures, such as rhythms, cue-association and conditional probability, yet it is still unknown whether it reflects local sequential information of the previous trial. To explore this, we examined the relationship between the sequential effect and the pre-stimulus oculomotor inhibition. Our results (N=40) revealed that inhibition was weaker when the previous trial was longer than the current trial, in line with findings of behavioral metrics. These findings indicate that the pre-stimulus oculomotor inhibition covaries with expectation based on local sequential information, demonstrating the tight connection between this phenomenon and expectation and providing a novel measurement for studying sequential effects in temporal expectation.

Keywords: Temporal expectation; Temporal attention; Temporal orientation; Microsaccades; Fixational saccades

Introduction

Our day-to-day life is full of temporal uncertainty. We find ourselves asking, when will the next bus arrive? How long do I need to stand in the queue? When is my manuscript expected to be back from review? To guide our behavior through this uncertainty, e.g., to decide whether we should take a cab, switch to another queue, or contact our editor, we form temporal predictions regarding the likely onset time of expected events (Nobre & van Ede, 2018). The formation of temporal expectations can be based on different sources of information, i.e., temporal priors, such as associations with an informative cue (Amit et al., 2019; Coull et al., 2000; Miniussi et al., 1999) or prior statistical knowledge (Tal-Perry & Yuval-Greenberg, 2022). One such source of information is our experience, acquired through previous encounters with similar events. This form

of temporal expectation is often called *sequential effects* – the expectation that the timing of the

present event will resemble that of the previous one.

Sequential effects in the temporal domain are mostly studied using reaction time (RT)

measurements (Capizzi et al., 2015; Possamai et al., 1973; Steinborn & Langner, 2012; Tal-Perry

& Yuval-Greenberg, 2022). The typical trial design of these studies includes a warning signal

followed, after a varying interval called *a foreperiod*, with a target. A common finding in this field

is that when the foreperiod of one trial is shorter than its precedent, perceptual decisions regarding

the present target will be slower. This was interpreted as reflecting the expectation that the duration

of the present foreperiod will resemble that of the previous one. However, the opposite effect does

not always occur, i.e., when the foreperiod of the present trial is longer than the precedent trial is,

RTs remain unchanged relative to when the two trials have the same foreperiod (Niemi &

Näätänen, 1981; Tal-Perry & Yuval-Greenberg, 2022).

While RT provides a useful behavioral measurement of temporal expectation, it measures

temporal expectations, by definition, retrospectively, after the event has already occurred and

expectations have already been formed. In a series of studies, we have shown that saccade rate can

be used as a marker of temporal expectation, termed *the pre-target oculomotor inhibition effect*,

which can be measured while expectations are formed rather than retrospectively. The pre-target

oculomotor inhibition effect is a reduction in the rate of eye movements which occurs a few

hundred milliseconds prior to the appearance of a predictable, relative to an unpredictable, target

(Abeles et al., 2020; Amit et al., 2019; Badde et al., 2020; Dankner et al., 2017; Tal-Perry & Yuval-

Greenberg, 2020, 2021).

The oculomotor inhibition effect was shown to be modulated by various types of temporal

expectations, including those driven by rhythm (Dankner et al., 2017), by associations between a

cue and a target, and by hazard rate – the change in the conditional probability for target occurrence

that changes when times passes by and the target has not yet appeared (Amit et al., 2019; Tal-Perry

& Yuval-Greenberg, 2020). However, to date, it remains unknown whether the pre-stimulus

oculomotor inhibition marker is also modulated by local sequential information, i.e., temporal

predictions induced by the recent previous trials. Finding a link between sequential effects and the

oculomotor inhibition effect would indicate that this marker is associated with the formation of

temporal expectations based on recent previous experiences. Such a finding would lend further

support to the validity of this index as a measurement of temporal expectations and provide a novel

metric for studing the effect of local sequential information.

In this study, we examined whether oculomotor inhibition covaries with sequential effects

in two experiments. In Experiment 1 (N=40), we examined saccade rate as a function of the

difference between the foreperiod of one trial and that of its precedent trial. We hypothesized that

local sequential information is reflected in eye movements dynamics – that when the foreperiod of

a given trial is shorter than that of its precedent, there would be less pre-target oculomotor

inhibition (i.e., more pre-target saccades) relative to when it was longer or equal. Our findings

confirmed this hypothesis, indicating the oculomotor measurements are modulated by sequential

effects as RTs. In Experiment 2 (*N*=40) we examined whether this first-order oculomotor

sequential effect could provide an alternative interpretation to our previous findings of the pre-

target oculomotor inhibition effect. This study joins previous studies by showing that the pre-

stimulus oculomotor inhibition covaries with temporal expectation of various temporal priors and

modalities, supporting its validity as an index for the formation of temporal expectations.

Methods

Participants

A total of 80 participants were included in the study: 40 participants were included in Experiment

1 (25 females, 3 left-handed, Mean age 24.95±3.82 standard deviations [SD]) and 40 participants

in Experiment 2 (24 females, one left-handed, Mean age 22.55±3.12 SD). All participants were

healthy, reported normal or corrected-to-normal vision, and reported no history of neurological or

psychiatric disorders. Participants received payment or course credit for their participation. The

experimental protocols were approved by the ethics committees of Tel-Aviv University and the

School of Psychological Sciences. Prior to participation, participants signed informed consent

forms. The present study includes two experiments that are based on reanalyses of three published

data sets as follows: Exp. 1 consisted of a novel analysis of eye tracking data of participants who

were originally included in Tal-Perry & Yuval-Greenberg (2022), but no eye tracking analysis was

performed in the original study; Exp. 2 consisted of a reanalysis of the eye tracking data of datasets

published originally in Amit et al. (2019) and Tal-Perry & Yuval-Greenberg (2020).

Stimuli

Experiment 1. The fixation object consisted of a dot (0.075° radius) within a ring (0.15°

radius), embedded within a diamond shape (0.4x0.4°). The edges of the diamond changed color

from black to white, cueing attention to the left (two left edges became white) or right (two right

edges became white) side of the fixation object, or remaining neutral with respect to the target

location (all four edges became white). The target was a black asterisk (0.4x0.4°) presented at 4°

eccentricity to the right or left of the fixation object. A 1000 Hz pure tone was played for 60 ms as

negative feedback following errors. Fixation objects and targets were presented on a mid-gray

background.

Experiment 2. The fixation object in this experiment was a cross (black or blue, $0.4° \times$

$0.4°$), and the target was a Gabor grating patch (2° diameter, 30% contrast, spatial frequency of 5

cycles/degree) slightly tilted clockwise (CW) or counter-clockwise (CCW) from vertical, with tilt

degree determined individually via 1-up 3-down staircase procedure. All stimuli were displayed

at screen center on a mid-gray background.

Procedure

The datasets that we have used originated from three previously published studies. Exp. 1 was

based on the dataset reported in Tal-Perry & Yuval-Greenberg (2022). Exp. 2 was based on the

combined datasets of Amit et al., (2019) and part of the dataset of Tel-Perry & Yuval-Greenberg

(2020), both based on identical procedures.

General procedure. In both experiments, participants were seated in a dimly lit room, with

a computer monitor placed 100 cm in front of them (24" LCD ASUS VG248QE, $1,920 \times 1,080$

pixels resolution, 120 Hz refresh rate, mid-gray luminance was measured to be 110 cd/m^2). During

the session, participants rested their heads on a chinrest. MATLAB R2015a (Mathworks, USA)

was used to code and control the experiment, with stimuli displayed using Psychophysics Toolbox

v3 (Brainard, 1997).

Experiment 1. Each trial started with a central black fixation object, presented until an

online gaze-contingent procedure verified 1000 ms of stable fixation, defined by the placement of

gaze within a radius of 1.5° of screen center. Following this, the edges of the fixation object

changed color for 200 ms to represent a spatial cue (right or left in 75% of trials or neutral in the

remaining 25% of trials). After a varying foreperiod (500 / 900 / 1300 / 1700 / 2100 ms) the target

was briefly (33 ms) presented at 4° eccentricity to the left or right of the screen center, with the

cue being valid with the target location in 75% of informative trials. Participants were instructed

Figure 1 *Trial progression*. (A) Trial progression for Exp. 1, adapted from Tal-Perry & Yuval-Greenberg (2022); (B) Trial progression for Exp. 2, adapted from Tal-Perry & Yuval-Greenberg (2020), which was identical to the trial progression used in Amit et al. (2019).

to respond as fast as possible upon detecting the target via a single button press. Between groups,

participants were presented with the five foreperiods in either a uniform distribution (20%

probability for each foreperiod) or an inverse-U-shaped distribution (a ratio of 1:2:3:2:1 between

the five foreperiods, leading to trial percentages of approximately 11%, 22%, 33%, 22%, and 11%,

respectively). Fixation was monitored throughout the foreperiod, using an online gaze-contingent

procedure, and trials that included ≥ 1.5° gaze-shift for more than 10 ms during this period were

aborted and repeated at a later stage of the session. An error feedback tone was played when

participants responded before the target onset or did not respond within 1000 ms following the

target onset. These trials were not included in the analysis. Participants of the uniform distribution

group (N=20) performed 10 blocks of 160 trials each, divided into two sessions. Participants of

the inverse-U-shaped distribution group (N=20) performed 18 blocks of 144 trials each, divided

into three sessions done on separate days. A short break was given after each block. A practice

block of 10 trials with random conditions was administered at the beginning of each session.

Figure 1A summarizes the trial procedure used in this experiment.

Experiment 2. A central black fixation cross was presented between trials for a jittered

inter-trial interval of 700-1200 ms. At trial onset, the fixation cross changed color from black to

blue marking the onset of the foreperiod interval. After the foreperiod has elapsed, the target (tilted

Gabor patch) was briefly (33 ms) presented and followed by a blank screen, and participants were

requested to perform a 2AFC discrimination on the Gabor tilt by pressing one of two keyboard

keys. Foreperiod was set to be between 1000-3000 ms in 500 ms increments. In fixed blocks,

foreperiod was constant throughout the block. In random blocks, the foreperiod randomly varied

from trial to trial. In part of the dataset (originally published in Amit et al., 2019), participants

completed a total of five fixed blocks (one of each foreperiod) and five random blocks, with 100

trials per block. In the rest of the dataset (originally published in Tal-Perry & Yuval-Greenberg,

2020), participants completed two fixed blocks (one of 1000 ms and one of 2000 ms), and five

random blocks, with 80 trials per block. Ten additional blocks performed by the participants in

Tal-Perry & Yuval-Greenberg (five blocks with a distribution of 80% 1000 ms foreperiods and

20% 2000 ms foreperiods, and five blocks with the opposite ratio) were not included in the present

study. The collapsed datasets of the two original studies were imbalanced in their number of trials

per condition: there were 500 random trials and 500 fixed trials of five different foreperiods per

participant in Amit et al. and there were 400 random trials and 160 fixed trials of two different

foreperiods per participant in Tal-Perry & Yuval-Greenberg. This imbalance was accounted for by

the generalized linear mixed model analysis which takes into account trial-wise variance (see

Statistical analysis). **Figure 1B** summarizes the trial procedure used in this experiment.

Eye-tracking

Eye movements were monitored using EyeLink 1000 Plus infrared video-oculographic desktop

mounted system (SR Research Ltd., Oakville, ON, Canada), with a 1000 Hz sampling rate and the

standard online and offline analog filters provided by Eyelink (Stampe, 1993). A nine-point

calibration was performed at the beginning of the experiment and when necessary. Raw gaze data

was low-pass filtered at 60 Hz and segmented between -500 ms relative to cue onset and 500 ms

relative to target onset. Blinks were detected based on the built-in algorithm provided by EyeLink,

plus an additional criterion requiring a binocular change in pupil size that exceeded 2.5 standard

deviations from the segment's mean pupil size for 3 or more consecutive samples (Hershman et

al., 2018). Saccades were detected using a published algorithm (Engbert & Kliegl, 2003; Engbert

& Mergenthaler, 2006), with saccade onset defined as the point in which the absolute standardized

eye velocity exceeded the segment's median eye velocity by six or more standard deviations, for a

minimum of six consecutive samples. Only binocular saccades were included in the analysis. A 50

ms interval between saccade offset and the next saccade onset was imposed to prevent detection

of overshoots. Intervals with blinks and 200 ms before the onset of blinks and after their offsets,

were excluded from the saccades analysis. The analysis included saccades of all sizes, although

most saccades were minuscule (<1°) due to the instruction to maintain fixation. Correlation

between saccade amplitude and peak velocity (main sequence, Zuber et al., 1965) was high ($r >$

0.9) for all participants, verifying the validity of the saccade detection procedure.

For each trial, we determined whether a saccade was detected in the period of -300 to 0 ms

relative to the target onset. This range was based on the time period known to include the

oculomotor inhibition effect as reported in Tal-Perry & Yuval-Greenberg (2021). Trials that

included a blink or missing data during this period were discarded from the analysis.

Statistical analysis

Experiment 1. For each trial, we calculated the difference between its foreperiod and the

foreperiod of the previous trial (FP-difference, $FP_n - FP_{n-1}$). For this purpose, the first trial of

each session was discarded from the analysis. The resulting continuous factor ranged between -

1600 and 1600 ms in 400 ms increments and was Z-scaled to reduce computational complexity

(standardized foreperiod difference, SFD). The probability of a saccade onset in the pre-target time

interval was then analyzed using a generalized linear mixed model (GLMM), assuming a binomial

family of responses (saccade present / absent) with a logit link, i.e., a logistic mixed model. We

based our choice of the model on the assumption that due to pre-stimulus oculomotor inhibition,

in the vast majority of trials, we are not expecting more than a single saccade to occur in the

analyzed duration, with trials deviating from this assumption being rare and of little impact on the

overall results. The following fixed factors were included in the model: (1) a scaled linear and

quadratic relation between the current and previous foreperiod (SFD), to model the sequential

effect; (2) the foreperiod distribution (uniform / inverse-U-shaped), a between-subject factor, using

sum contrasts; (3) The interaction between the two factors, to model the effect of foreperiod

distribution on the sequential effect. As oculomotor behavior was examined prior to target onset,

the trial's spatial cueing condition used in the original experiment (valid / invalid / neutral) was

not included as a factor.

Experiment 2. Trials of the random block were screened according to the foreperiod of the

previous trial, such that only trials in which the previous foreperiod was equal to the current trial's

foreperiod were included in the analysis. The probability of a saccade onset was then analyzed

using a logistic mixed model, with the following factors: (1) the current trial's foreperiod; (2)

condition (fixed / random); and (3) the interaction between the two factors. Difference contrast

was used for Foreperiod, and sum contrast was used for Condition.

General statistical analysis. For all models, statistical significance for main effects and

interactions was determined via a likelihood-ratio (LR) test against a reduced nested model

excluding the fixed term (i.e., type-II sum of squares, SS), and statistical significance for parameter

coefficients was determined according to Wald z-test (Fox, 2016). To provide support for null

results ($p > 0.05$), we additionally calculated the Bayes Factor (BF) between the full and reduced

model, using BIC approximation (Wagenmakers, 2007). BF is reported with the null result in the

denominator (BF_{01}), representing how much the data is supported by the null model relative to the

full model. The model's random effect structure was selected according to the model that was

found to be most parsimonious with the data, i.e., the fullest model that the data permits while still

converging with no singular estimates (Bates, Kliegl, et al., 2015), in order to balance between

type-I error and statistical power (Matuschek et al., 2017). This was achieved by starting with a

random intercept by subject only model, and continuing to a model with random slopes for fixed

terms by subject and their correlation parameters, and from there to a random-interaction-slopes

by-subject model, testing for model convergences in each step. Models that failed to converge

were trimmed by the random slope with the least explained variance and were retested. Analyses

were performed in R v4.0.3 using R-studio v1.3.959 (R Core Team, 2018). Modelling was

performed using the lme4 (Bates, Mächler, et al., 2015) package, BF was calculated using the

BayesFactor package (Morey & Rouder, 2018), and model diagnostics were performed using the

performance package (Lüdecke et al., 2020). An R-markdown file describing all the model fitting

steps and diagnostic checks on the final model is available at the project's OSF repository (see

Data Availability Statement)

222   **Results**

Experiment 1: the oculomotor sequential effect

In the first experiment, we tested whether the pre-stimulus oculomotor inhibition is affected by

contextual information about the previous trial in a speeded detection task. For this goal, we

analyzed the eye movement data from Tal-Perry & Yuval-Greenberg (2022). We first start by

examining the behavioral sequential effect in the same dataset.

Behavioral results. The behavioral results were reported in a previous publication (Tal-

Perry & Yuval-Greenberg, 2022) and are summarized here. RTs were modeled as a factor of the

Standardized Foreperiod Difference between the current and the preceding trial (SFD, continuous;

linear and quadratic terms), Foreperiod Distribution (uniform / inverse-U-shaped), Condition

(valid / invalid / neutral), and their interaction terms. Findings revealed a strong sequential effect,

such that RT was slower in trials that were preceded by a longer trial, while the opposite was not

found. This effect was modulated by the Foreperiod Distribution, but not by Condition. These

results confirm that a behavioral sequential effect was present in the analyzed dataset.

Eye tracking results. The probability of a saccade occurring prior to target onset was

examined using a logistic mixed model, with SFD (continuous; linear and quadratic terms),

Foreperiod Distribution (uniform / inverse-U-shaped), and the interaction between them set as

fixed factors, allowing for a random intercept per subject and a random slope to SFD per subject.

The saccade probability for each distribution as a factor of the standardized difference

between the current and previous foreperiod is displayed in **Figure 2**. Results showed that SFD

had a strong influence on saccade likelihood ($\chi^2(2) = 30.225$, $p < .001$), with a strong negative

linear slope (log estimate -0.238, $z = -6.061$, $p < .001$), indicating that saccade probability is

reduced as the difference between the current and previous trial gradually turns from being

negative (previous is longer) to positive (previous is shorter). This indicates that oculomotor

inhibition is enhanced when the previous trial becomes shorter relative to the current trial. This

negative slope was accompanied by a weaker positive quadratic slope (log estimate 0.081, $z =$

4.483, $p < .001$), reflecting that the degree of saccade inhibition is gradually decreased until it

changes direction. The combination of the negative linear and the positive quadratic trends resulted

in the asymmetry between negative and positive SFD observed.

Figure 2 *Experiment 1 results*. The probability of performing a saccade during the -300 to 0 ms period relative to target onset, as a function of the difference between the current and previous foreperiod, and the Foreperiod Distribution. Negative values indicate that previous foreperiod was longer than current foreperiod, and vice-versa for positive values. Error bars depict ± 1 standard error from the mean, correcting for within-subject variability (Cousineau & O'Brien, 2014). Lines depict 2^{nd} polynomial fit to the observed data. $N = 20$ in each distribution.

To examine this gradual change, we modeled SFD as a categorical factor and contrasted
each level with an SFD of zero (i.e., previous foreperiod equals current; see **Table 1**). As can be
observed, the negative SFDs (longer foreperiod at previous trial) resulted in higher saccade
probability compared to the zero SFD (no difference in foreperiod between this and the previous
trial), with a gradual decrease in saccade probability difference as the SFD decreases. The positive
SFDs (shorter foreperiod at previous trial) resulted in a significantly smaller saccade probability
for the 400 and 800 ms difference relative to the zero SFD, with the effect gradually decreasing as
the SFD increases, such that no significant difference was observed for the larger positive SFD
values. These findings are consistent with the hypothesis that oculomotor inhibition reflects
anticipation – in negative SFD trials, participants are anticipating the target to occur at a later stage
of the trial, thus there is a weaker inhibition of saccades prior to the actual target onset as compared
to trials where the target appeared at the expected time. The reduced saccade rate observed in the
first two positive SFD levels may reflect the effect of conditional probability (hazard rate) – as
target was expected to appear but has yet to appear, anticipation continues to be built up, and with
it, the inhibition of saccades continues to increase. Lastly, the lack of significant difference in the
last two positive SFD levels may reflect the added time uncertainty in expectation – at these trials,

SFD	Log difference [95% CI]	z-ratio	FDR-corrected p
-1600	0.796 [0.634 0.958]	13.107	< .001
-1200	0.562 [0.455 0.669]	14.075	< .001
-800	0.352 [0.263 0.441]	10.520	< .001
-400	0.165 [-0.082 0.248]	5.285	< .001
400	-0.169 [-0.258 -0.080]	-5.073	< .001
800	-0.213 [-0.315 -0.111]	-5.587	< .001
1200	-0.091 [-0.219 0.037]	-1.888	.067
1600	0.091 [-0.157 0.260]	0.654	.513

Table 1 *SFD from zero level.* The model estimates represent the difference in log-odds saccade probability of [SFD level minus zero level] along with a 95% confidence interval (CI). Reported statistics based on log odds-ratio with the zero SFD level in the denominator. Reported *p* values were false discovery-rate (FDR) corrected. SFD = standardized foreperiod difference

target appeared long after the expected time, such that anticipating when would it occur became increasingly more difficult.

Importantly, the SFD significantly interacted with the Foreperiod Distribution ($\chi^2(2) = 17.635$, $p < .001$), stemming from both a difference in the linear (log estimate 0.038, $z = 3.529$, $p < .001$) and quadratic (log estimate 0.022, $z = 2.554$, $p = .011$) components between the foreperiod distributions. To explore this interaction, we contrasted each SFD between distributions (see **Table 2**). As can be observed, the two foreperiod distributions significantly differed only for the positive SFD values starting with 800 ms. Lastly, we observed a significant effect for Foreperiod Distribution ($\chi^2(1) = 23.588$, $p < .001$), stemming from the higher probability of saccade occurrence for the inverse-U-shaped distribution. These results suggest that the pre-stimulus oculomotor inhibition reflects the combination of short-term expectations from the previous trials, and long-term expectations from the distribution of intervals and higher-order sequential effects.

SFD	Log difference [95% CI]	z-ratio	FDR-corrected p
-1600	0.119 [-0.342 0.103]	1.054	.292
-1200	-0.077 [-.055 0.201]	-1.140	.254
-800	-0.093 [-0.009 0.195]	-1.792	.073
-400	-0.087 [-0.003 0.178]	-1.894	.058
0	-0.030 [-0.057 0.116]	-0.670	0.503
400	-0.078 [-0.023 0.179]	-1.506	0.132
800	-0.334 [-0.209 -0.459]	-5.244	<.001
1200	-0.243 [-0.074 -0.411]	-2.821	.005
1600	-0.524 [-0.232 -0.815]	-3.524	<.001

Table 2 *SFD contrasts by Foreperiod Distribution.* The model estimates represent the difference in log-odds saccade probability [uniform minus inverse-U-shaped] conditions along with a 95% confidence interval (CI). Reported statistics based on log odds-ratio with uniform distribution in the nominator. Reported *p* values were false discovery-rate (FDR) corrected. SFD = standardized foreperiod difference.

Experiment 2: Does the fixed effect stem from the sequential effect?

280
Previous studies have demonstrated that the pre-stimulus oculomotor inhibition is stronger when
the foreperiod is fixed throughout the block compared to when it varies randomly (Abeles et al.,
2020; Amit et al., 2019; Badde et al., 2020; Dankner et al., 2017; Tal-Perry & Yuval-Greenberg,
2020, 2021). In Exp. 1, we observed that saccades were inhibited to a larger degree when the
previous foreperiod matched the current foreperiod (SFD of zero) compared to when the previous
foreperiod was longer in duration (negative SFD), but less so when it was shorter in duration
(positive SFD). This raises the question of whether the fixed vs. random effect observed in
previous studies stemmed from local sequential information rather than from target probability. In
these previous studies, the fixed and random trials were included in different blocks: in the fixed
blocks the previous trial always matched the present trial, i.e., had SFD zero; In contrast, in the
random block, the previous trial could have been longer, equal, or shorter than the present trial.
Since the sequential effect is asymmetrical (see **Figure 2**), averaging trials with positive and

negative SFDs (as in the random condition) is expected to lead to a higher saccade rate than trials

of zero SFD (as in the fixed condition). It could therefore be hypothesized that the sequential effect

is at the basis of the difference in saccade rate between the fixed and random conditions (higher

pre-stimulus saccade rate for random relative to fixed) rather than temporal expectation, as was

previously suggested.

To examine this hypothesis, we reexamined the fixed vs. random effect reported in two

previous studies (Amit et al., 2019; Tal-Perry & Yuval-Greenberg, 2020), while controlling for

the previous trial. We compared the probability of performing a saccade prior to target onset

between the fixed and the random trials while including only trials in which the previous foreperiod

was equal to the current foreperiod. Thus, the n-1 identity was matched between the two

conditions. Results were analyzed using a GLMM assuming a binomial family of response, with

Condition (fixed / random), Foreperiod (continuous), and the interaction between them as within-

subject fixed factors, allowing for a random intercept by subject and random slope for each of the

main effects by subject.

Figure 3 depicts the descriptive results of this analysis. As can be observed, we found

saccade probability to be significantly lower in the fixed compared to the random condition ($\chi^2(1)$

$= 6.700$, $p = .01$). There was no significant effect for foreperiod ($\chi^2(1) = 0.916$, $p = .338$, $BF_{01} =$

77.675), but foreperiod significantly interacted with condition ($\chi^2(1) = 20.500$, $p < .001$), owing

to the positive slope in the fixed condition compared to the negative slope in the random condition.

This reversal in slopes matches what was observed in Amit et al. (2019) – the decrease in

oculomotor inhibition in the fixed condition as foreperiod increases could be explained by the

increase in temporal uncertainty, while the increase in oculomotor inhibition in the random

condition might be the result increasing hazard rate, i.e., the likelihood of an event to occur given

that it has yet to occur. Overall, these results are consistent with those found when not controlling

for the previous trial foreperiod and therefore suggest that the fixed effect observed in previous

studies cannot be explained solely as the result of a sequential effect, but likely reflects target

predictability.

Figure 3 *Experiment 2 results.* The probability of performing a saccade during the -300 to 0 ms period relative to target onset, as a function of Condition and Foreperiod, with the n-1 trial's foreperiod matching the current trial's foreperiod for both conditions. Error bars depict ± 1 standard error from the mean, correcting for within-subject variability (Cousineau & O'Brien, 2014). Lines depict linear fit to the observed data. *N* = 40

Discussion

In this study, we examined whether the pre-stimulus oculomotor inhibition is affected by local
sequential information, i.e., the sequential effect – the previous trial history with respect to the
current trial. In Exp. 1, we demonstrated that the likelihood of performing a saccade prior to target
onset changed as a factor of the relation between the current trial's foreperiod and the previous
trial's foreperiod, consistent with the typical pattern observed for RT with respect to the sequential
effect and demonstrating once more that the pre-stimulus oculomotor inhibition reflects
anticipation similarly to RTs.
327
As a follow-up question, in Exp. 2 we examined whether the sequential effect could
provide an alternative explanation to findings observed in previous cue-based temporal expectation
studies. In these studies, we found that when the foreperiod was fixed throughout the block, pre-
target saccades were inhibited to a larger degree than when the foreperiod varied within the block,
for short foreperiods (Abeles et al., 2020; Amit et al., 2019; Badde et al., 2020; Dankner et al.,
2017; Tal-Perry & Yuval-Greenberg, 2020, 2021). These findings were interpreted as the result of
a higher expectation as target predictability increased. However, in the fixed block of all these
cases, the previous trial's foreperiod always matched the current trial's foreperiod. Thus, the lower

saccade rate found in the fixed condition could be interpreted as resulting from a sequential effect

of the previous trial. In Exp. 2, we demonstrated that this interpretation is unlikely – trials from

the random condition in which the previous foreperiod matched the current trial's, nevertheless

exhibited a higher saccade rate compared to the fixed condition. Together, these two experiments

indicate that oculomotor inhibition is modulated both by local sequential information, and by more

global temporal priors, such as target probability.

Models of the sequential effect

In Exp. 1, we found an asymmetrical sequential effect of pre-target saccade rate: pre-target saccade

rate was higher in trials succeeding trials with longer foreperiod but it was either similar or only

slightly lower in trials succeeding a shorter foreperiod. This asymmetry is consistent with the

asymmetric sequential effect reported previously with RTs: RTs in trials that follow a trial with

longer foreperiod tend to be slower than the RT in trials that follow trials of identical foreperiods,

but the opposite is typically not found (e.g., Los et al., 2001; Steinborn & Langner, 2012; Tal-

Perry & Yuval-Greenberg, 2022; Vallesi & Shallice, 2007). Two competing theories have tried to

explain this asymmetrical pattern in RT data – the *dual-process model* (Vallesi et al., 2007; Vallesi

& Shallice, 2007) and the *trace-conditioning model* (Los et al., 2001) which was later updated to

form the *multiple-trace theory* (Los et al., 2014, 2017, 2021; Salet et al., 2022). While these models

were developed to explain results in behavioural data, the two explanations may be adapted to

interpret the sequential effect observed for the pre-stimulus oculomotor inhibition in the current

study.

According to the *dual-process model*, sequential effects stem from two factors: the

monitoring of conditional probability that varies during the given trial, along with an increase in

arousal from the previous trial. The effect of conditional probability could explain the finding of

an asymmetrical sequential effect, demonstrated in Exp. 1. When the previous foreperiod is shorter

than the current foreperiod, participants orient their expectations toward the short period in the

current trial, but given that the event has not occured, they reorient their expectations to the next

probable target onset, whose conditional probability is typically higher (e.g., under uniform

foreperiod distribution), thus leading to higher expectations and inhibition of eye movements prior

to the next target. This explanation is also in line with the observed difference in asymmetry

between the two foreperiod distributions in Exp. 1, as different foreperiod distributions lead to

different conditional probabilities. This view is consistent with our previous studies in which we

have demonstrated other effects of conditional probabilities on patterns of pre-stimulus oculomotor

inhibition (Abeles et al., 2020; Amit et al., 2019; Tal-Perry & Yuval-Greenberg, 2020).

369   The second process postulated by the dual-process model is arousal or alertness, defined

by the model as the readiness to respond. According to the model, arousal from the previous trial

affects expectations in the current trial. When preceded by a short trial, arousal tends to be high,

and RT is accordingly fast. When preceded by a long trial, the prolonged preparation causes

exhaustion of alertness, leading to poorer RT in the following trial (Steinborn & Langner, 2012;

Vallesi & Shallice, 2007). To provide a similar explanation in the case of pre-stimulus oculomotor

inhibition, one has to assume that arousal or readiness to respond is positively correlated with

oculomotor rate. In a previous study we showed that the pre-stimulus oculomotor inhibition is

independent of motor readiness (Tal-Perry & Yuval-Greenberg, 2021) and thus response readiness

is unlikely to explain the higher saccade probabilities observed for negative SFD trials (longer

previous trial) observed in Exp. 1 of the present study (see **Figure 2**). Thus, the dual-process model

falls short of explaining the full pattern of results in this study.

381   As an alternative, the *trace-conditioning model* (Los et al., 2001) and its recent expansion,

382   the *multiple-trace theory* (Los et al., 2014, 2017, 2021; Salet et al., 2022), suggest that the

383   asymmetrical sequential effect is the result of a single process – the activation of the weighted

384   memory traces of previous trials starting at cue onset, and decaying as time progresses. We begin

385   by explaining how the model interprets RT data in temporal expectation tasks in general, and then

386   move on to focus on the model's explanation of the sequential effect and its relation to our data.

387   According to the *multiple-trace theory*, the onset of the cue in each trial triggers a motor

inhibition which prevents the execution of a premature response during the foreperiod interval.

Simultaneously, the trial onset also activates a second neuronal population, whose activity peaks

at the time of the expected target onset. This pattern of inhibition followed by activation constitutes

the preparatory temporal profile, which is saved as a trace in memory after each trial. At each cue

onset, the existing memory traces, accumulated over previous trials, are reactivated and are

aggregated to a preparatory pattern which determines when will inhibition wane and activation

peak within the current trial. Due to the dissipation of memory over time, recent trials contribute

more strongly to the activation than older trials. As the preparatory pattern is an aggregation of

previous trials, different foreperiod distributions are expected to lead to different preparatory

patterns, as the mixture of previous foreperiod (i.e., the higher-order sequential effects) should

vary according to distribution. This, in turn, explains the different RT patterns that are induced by

different foreperiod distributions.

Given this model, it is clear to see how sequential effects come about. Trials in which the

previous foreperiod matches the current foreperiod would lead to a better preparation (more

activation and less inhibition) at target onset, compared to trials where there is a mismatch. To

explain the asymmetry in sequential effects of RT data, the multiple-trace theory postulates that

the activation pattern builds up and dissipates slowly over the trial. Thus, trials whose previous

foreperiod was shorter than the current foreperiod would still benefit from a relatively high

activation at target onset, leading to a relatively fast RT.

Can the multiple trace model explain the results observed in our study? First, it is possible

that the inhibitory process postulated by the model could manifest as inhibition of eye movements,

leading to a lower saccade probability prior to target onset when the previous and current trials

match, as observed in Exp. 1 (see **Figure 2**, negative vs. zero value). Thus, eye movements may

form a window into one of the latent variables postulated by the model. The observed asymmetry

in pre-stimulus oculomotor inhibition in Exp. 1 (**Figure 2**, positive values) can also be explained

by the model – as activation dissipates slowly over the trial, current trials whose foreperiod is

longer than the previous trial still feature relatively higher activation prior to the target onset, which

would translate into relatively lower saccade rate. The increase in saccade rate observed for

increasing positive SFDs (shorter previous trial) could likewise be accounted for by the dissipation

of activation postulated by the model. Lastly, the observed difference in oculomotor inhibition for

the uniform and inverse-U-shaped distributions can be explained by differences in the aggregated

traces between the two distributions – there are fewer trials with an extreme foreperiod difference

in the inverse-U-shaped distribution compared to the uniform distribution, meaning that

aggregated activity is predicted to be low at late time points during the trial in the inverse-U-shaped

condition, and this translates into a higher saccade probability at extreme positive values as

depicted in **Figure 2**.

Thus, of the two alternatives, our results are best explained by the multiple trace theory.

Furthermore, the pre-stimulus oculomotor inhibition may provide means to study the latent

inhibitory process postulated by the theory, thus providing a mean by which the theory could be

tested in future studies.

Conclusions

Our results demonstrate that the pre-stimulus oculomotor inhibition is modulated by temporal

information that stems from a recent experience. This study joins a growing body of studies that

demonstrate that pre-stimulus oculomotor inhibition reflects different types of temporal

expectation processes, based on rhythms (Dankner et al., 2017), cue associations (Abeles et al.,

2020; Amit et al., 2019; Tal-Perry & Yuval-Greenberg, 2020, 2021) or hazard rate function (Tal-

Perry & Yuval-Greenberg, 2020); with the degree of anticipation correlated with the degree of

inhibition (Tal-Perry & Yuval-Greenberg, 2020). The present study expands this list by showing

that anticipation based on local sequential information is similarly correlated with pre-stimulus

oculomotor inhibition. Together, this series of studies demonstrate that whenever there is temporal

anticipation, there is also inhibition of eye movements. This study additionally provides a metric

to study sequential effects without requiring a response from the participant. This metric may allow

studying the sequential effect in uncooperative populations, such as infants and toddlers. An open

question that remains following this study is whether the oculomotor sequential effect depends on

attention. Future studies could manipulate attention to the target and examine the effect of this

manipulation on the oculomotor inhibition effect. Lastly, the inhibitory nature of this phenomenon

is in line with the inhibitory process postulated by the multiple trace theory, therefore, could be

used as a direct way to test the multiple trace theory.

Data Availability statement. The datasets used in Exp. 1 and 2 and an R-markdown file that

reproduces all the reported modeling, statistical analyses, and graphs within the paper are uploaded

to the Open Science Foundation repository and are available at doi.org/10.17605/OSF.IO/PV3N2

449   **References**

450   Abeles, D., Amit, R., Tal-Perry, N., Carrasco, M., & Yuval-Greenberg, S. (2020). Oculomotor

451   inhibition precedes temporally expected auditory targets. *Nature Communications*, *11*(1),

452   3524. https://doi.org/10.1038/s41467-020-17158-9

453   Amit, R., Abeles, D., Bar-Gad, I., & Yuval-Greenberg, S. (2017). Temporal dynamics of

454   saccades explained by a self-paced process. *Scientific Reports*, *7*(1), 1–15.

455   https://doi.org/10.1038/s41598-017-00881-7

Amit, R., Abeles, D., Carrasco, M., & Yuval-Greenberg, S. (2019). Oculomotor inhibition

reflects temporal expectations. *NeuroImage*, *184*(March 2018), 279–292.

https://doi.org/10.1016/j.neuroimage.2018.09.026

Badde, S., Myers, C. F., Yuval-Greenberg, S., & Carrasco, M. (2020). Oculomotor freezing

reflects tactile temporal expectation and aids tactile perception. *Nature Communications*,

11(1), 1–9. https://doi.org/10.1038/s41467-020-17160-1

Bates, D., Kliegl, R., Vasishth, S., & Baayen, H. R. (2015). Parsimonious Mixed Models. *ArXiv*.

http://arxiv.org/abs/1506.04967

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models

Using {lme4}. *Journal of Statistical Software*, *67*(1), 1–48.

https://doi.org/10.18637/jss.v067.i01

Brainard, D. H. (1997). The Psychophysics Toolbox. *Spatial Vision*, *10*, 433–436.

https://doi.org/10.1163/156856897x00357

Capizzi, M., Correa, Á., Wojtowicz, A., & Rafal, R. D. (2015). Foreperiod priming in temporal

preparation: Testing current models of sequential effects. *Cognition*, *134*, 39–49.

https://doi.org/10.1016/j.cognition.2014.09.002

Coull, J. T., Frith, C. D., Büchel, C., & Nobre, A. C. (2000). Orienting attention in time:

Behavioural and neuroanatomical distinction between exogenous and endogenous shifts.

Neuropsychologia, *38*(6), 808–819. https://doi.org/10.1016/S0028-3932(99)00132-3

Dankner, Y., Shalev, L., Carrasco, M., & Yuval-Greenberg, S. (2017). Prestimulus Inhibition of

Saccades in Adults With and Without Attention-Deficit / Hyperactivity Disorder as an

Index of Temporal Expectations. *Psychological Science*, *28*(7), 835–850.

https://doi.org/10.1177/0956797617694863

Engbert, R., & Kliegl, R. (2003). Microsaccades uncover the orientation of covert attention.

Vision Research, *43*(9), 1035–1045. https://doi.org/10.1016/s0042-6989(03)00084-1

Engbert, R., & Mergenthaler, K. (2006). Microsaccades are triggered by low retinal image slip.

Proceedings of the National Academy of Sciences of the USA, *103*(18), 3–8.

https://doi.org/10.1073/pnas.0509557103

Fox, J. (2016). Applied regression analysis & genealized linear models. In *Sage*.

https://doi.org/10.1017/CBO9781107415324.004

Hershman, R., Henik, A., & Cohen, N. (2018). A novel blink detection method based on

pupillometry noise. *Behavior Research Methods*, *50*(1), 107–114.

https://doi.org/10.3758/s13428-017-1008-1

Los, S. A., Heuvel, C. E. V. D., & Van Den Heuvel, C. E. (2001). Intentional and unintentional

contributions to nonspecific preparation during reaction time foreperiods. *Journal of*

Experimental Psychology: Human Perception and Performance, *27*(2), 370–386.

https://doi.org/10.1037/0096-1523.27.2.370

Los, S. A., Kruijne, W., & Meeter, M. (2014). Outlines of a multiple trace theory of temporal

preparation. *Frontiers in Psychology*, *5*(September), 1–13.

https://doi.org/10.3389/fpsyg.2014.01058

Los, S. A., Kruijne, W., & Meeter, M. (2017). Hazard versus history: Temporal preparation is

driven by past experience. *Journal of Experimental Psychology: Human Perception and*

Performance, *43*(1), 78–88. https://doi.org/10.1037/xhp0000279

Los, S. A., Nieuwenstein, J., Bouharab, A., Stephens, D. J., Meeter, M., & Kruijne, W. (2021).

The warning stimulus as retrieval cue: The role of associative memory in temporal

preparation. *Cognitive Psychology*, *125*(April 2019), 101378.

https://doi.org/10.1016/j.cogpsych.2021.101378

Lüdecke, D., Makowski, D., & Waggoner, P. (2020). *performance: Assessment of Regression*

Models Performance. https://cran.r-project.org/package=performance

Matuschek, H., Kliegl, R., Vasishth, S., Baayen, H. R., & Bates, D. (2017). Balancing Type I

error and power in linear mixed models. *Journal of Memory and Language*, *94*, 305–315.

https://doi.org/10.1016/j.jml.2017.01.001

Miniussi, C., Wilding, E. L., Coull, J. T., & Nobre, A. C. (1999). Orienting attention in time.

Modulation of brain potentials. *Brain*, *122*, 1507–1518.

https://doi.org/10.1093/brain/122.8.1507

Morey, R. D., & Rouder, J. N. (2018). *BayesFactor: Computation of Bayes Factors for Common*

Designs. https://cran.r-project.org/package=BayesFactor

Niemi, P., & Näätänen, R. (1981). Foreperiod and Simple Reaction Time. *Psychological*

Bullettin, *89*(1), 133–162. https://doi.org/10.1037/0033-2909.89.1.133

Nobre, A. C., & van Ede, F. (2018). Anticipated moments: Temporal structure in attention.

Nature Reviews Neuroscience, *19*(1), 34–48. https://doi.org/10.1038/nrn.2017.141

Possamai, C. A., Granjon, M., Requin, J., & Reynard, G. (1973). Sequential effects related to

foreperiod duration in simple reaction time. *Perceptual and Motor Skills*, *36*(3 (II)),

1185–1186. https://doi.org/10.2466/pms.1973.36.3c.1185

R Core Team. (2018). *R: A Language and Environment for Statistical Computing*. R Foundation

for Statistical Computing. https://www.r-project.org/

Salet, J. M., Kruijne, W., van Rijn, H., Los, S. A., & Meeter, M. (2022). FMTP: A unifying

computational framework of temporal preparation across time scales. *Psychological*

Review, *129*(5), 911–948. https://doi.org/10.1037/rev0000356

Stampe, D. M. (1993). Heuristic filtering and reliable calibration methods for video-based pupil-

tracking systems. *Behavior Research Methods, Instruments, & Computers*, *25*(2), 137–

142. https://doi.org/10.3758/BF03204486

Steinborn, M. B., & Langner, R. (2012). Arousal modulates temporal preparation under

increased time uncertainty: Evidence from higher-order sequential foreperiod effects.

Acta Psychologica, *139*(1), 65–76. https://doi.org/10.1016/j.actpsy.2011.10.010

Tal-Perry, N., & Yuval-Greenberg, S. (2020). Pre-target oculomotor inhibition reflects temporal

orienting rather than certainty. *Scientific Reports*, *10*(1), 1–9.

https://doi.org/10.1038/s41598-020-78189-2

Tal-Perry, N., & Yuval-Greenberg, S. (2021). Prestimulus inhibition of eye movements reflects

temporal expectation rather than time estimation. *Attention, Perception, &*

Psychophysics, 1–13. https://doi.org/10.3758/s13414-021-02319-9

Tal-Perry, N., & Yuval-Greenberg, S. (2022). The Spatiotemporal Link of Temporal

Expectations: Contextual Temporal Expectation Is Independent of Spatial Attention. *The*

Journal of Neuroscience, *42*(12), 2516–2523. https://doi.org/10.1523/jneurosci.1555-

21.2022

Vallesi, A., & Shallice, T. (2007). Developmental Dissociations of Preparation Over Time:

Deconstructing the Variable Foreperiod Phenomena. *Journal of Experimental*

Psychology: Human Perception and Performance, *33*(6), 1377–1388.

https://doi.org/10.1037/0096-1523.33.6.1377

Vallesi, A., Shallice, T., & Walsh, V. (2007). Role of the prefrontal cortex in the foreperiod

effect: TMS evidence for dual mechanisms in temporal preparation. *Cerebral Cortex*,

17(2), 466–474. https://doi.org/10.1093/cercor/bhj163

Wagenmakers, E. J. (2007). A practical solution to the pervasive problems of p values.

Psychonomic Bulletin and Review, *14*(5), 779–804. https://doi.org/10.3758/BF03194105

Zuber, B. L., Stark, L., & Cook, G. (1965). Microsaccades and the velocity-amplitude

relationship for saccadic eye movements. *Science*, *150*(702), 1459–1460.

https://doi.org/10.1126/science.150.3702.1459

553