Dolphin social phenotypes typically show individual variation in response to environmental change alongside population-level stability

David N. Fisher*

School of Biological Sciences, University of Aberdeen, King's College, Aberdeen, AB243FX

* corresponding author: david.fisher@abdn.ac.uk

Barbara J. Cheney

School of Biological Sciences, University of Aberdeen, Lighthouse Field Station, George Street, Cromarty, IV11 8YL

Abstract

Social behaviours can allow individuals to flexibly respond to environmental change, potentially buffering adverse effects. However, individuals may respond differently to the same environmental stimulus, complicating our ability to predict population-level response to environmental change. Here we show that bottlenose dolphins (*Tursiops truncatus*) show a high degree of individual variability in responses to variation in proxies of food availability and climate, against a background of population-level stability. Individual dolphins did not consistently respond to yearly variation in environmental factors. However, at the monthly scale individuals became better connected to their local social environment in months of high salmon abundance, but their overall gregariousness and connectedness to the wider network did not change. This stability in social network position suggests there are important drivers maintaining a consistent structure. These results indicate that forecasting population responses to environmental change should account for among-individual variation in mean and plasticity of social behaviour.

Keywords: dolphin, environmental change, individual variation, plasticity, social network, Tursiops

Introduction

Animals engage in social interactions with conspecifics which are fundamental for determining health, access to resources, and reproductive success (Snyder-Mackler *et al.* 2020). Consequently, social interactions have a strong influence on ecological processes such as population dynamics and evolutionary processes such as the response to selection (Moore *et al.* 1997; Wolf *et al.* 1999; McGlothlin *et al.* 2010). To maintain the best fit with their environment, animals may adjust their social behaviour as conditions change (Komdeur & Ma 2021), for instance being more gregarious when resources are plentiful but less tolerant of conspecifics when resources are scarce (Fisher *et al.* 2021). Animals may also change their behaviour through development and during senescence (White *et al.* 2012). Such plasticity is a hallmark of behavioural traits and gives behaviour an important role in how animals interact with their environment.

When trying to understand how animals may respond through plasticity to changing environments, most examine responses at the population level (e.g., Abrahms *et al.* 2022), implying that all individuals respond similarly. However, individuals typically show variation in plasticity, and so each will respond differently to change (Nussey *et al.* 2007; Dingemanse *et al.* 2010, sometimes referred to as "I x E" i.e., individual by environment interactions; Stamps 2016; Stamps & Biro 2016). For example, European field crickets (*Gryllus campestris*) become bolder and more active as they age, but individuals vary in the extent of this, with some not increasing or even decreasing (Fisher *et al.* 2015). This individuality can influence biological processes such as population growth and adaptive change at a range of scales (Vindenes & Langangen 2015; see: Hamel *et al.* 2018 and accompanying papers), giving fundamentally different results to when population-level only effects are assumed (Brommer 2013b; Dingemanse & Wolf 2013). Additionally, the extent of among-individual variation in plasticity gives an upper-limit to the heritability of plasticity, which indicates how rapidly plasticity itself can evolve (Nussey *et al.* 2005; Chevin 2013; Araya-Ajoy & Dingemanse 2017). Understanding how plasticity as a trait in its own right can evolve is key in understanding how animals will adapt to more variable climates (Nussey *et al.* 2005; Brommer 2013a).

Marine mammals are a key group to study individual variation in response to environmental change.

They are typically long-lived, increasing the relative importance of plasticity versus adaptive

evolution for coping with contemporary environmental change (Hoffmann & Sgrò 2011). They are

also exposed to a wide range of changing conditions including climate, food availability, and

pollution, and their populations are often of conservation concern. All of these factors increase the

need for us to understand how they respond to changes in their environment (Simmonds & Isaac

2007; Tyack 2008; MacLeod 2009; Evans & Bjørge 2013; Avila *et al.* 2018; van Weelden *et al.* 2021;

Kebke *et al.* 2022).

Here, we studied a population of bottlenose dolphins (*Tursiops truncatus*) in the North Sea for 30

years, and regularly recorded their social associations. We use social network analysis to quantify

three different dimensions of individual social behaviour: an individual's gregariousness (strength),

how tightly its immediate social group interact together (clustering coefficient), and how well

connected an individual is to the entire population (closeness). We then used random regression

models to quantify individual social phenotypes and determine how these social phenotypes depend

on yearly and monthly variation in available proxies for climate and food availability. These models

also indicated whether bottlenose dolphins show individual changes in response to the

environment, or if population-level change was more prominent.

Material and methods

Study site & group data collection

This study used data from a bottlenose dolphin population on the east coast of Scotland (Fig. 1). The

population of over 200 individuals (Arso Civil *et al.* 2021) has been studied intensively as part of a

long term individual based study (Wilson *et al.* 1999; Cheney *et al.* 2013; Arso Civil *et al.* 2019b). We

use data from boat-based photo-identification surveys carried out annually between 1990 and 2020

which regularly recorded dolphin groups within the Moray Firth Special Area of Conservation (SAC)

(92/43/EEC) (Cheney *et al.* 2014). The SAC is a core part of the population's range with over 50% of

the population using this area each year, with regular annual summer survey effort aimed at

maximising sighting probability, and consistent and adaptive protocols (Cheney *et al.* 2014).

Additional data were also available from Whale and Dolphin Conservation land based photo-

identification surveys (Cheney *et al.* 2013). Data were available from a total of 913 surveys (between

10 and 90 surveys each year; average of 30) with the majority carried out between May to

September.

Figure 1. Images depicting a photo-identification survey (top let), a newborn calf with mother

(middle left), unique markings used to identify individuals (bottom left). and location of encounters

with bottlenose dolphins within the Moray Firth Special Area of Conservation from May to

September 1990 to 2020 (right).

During surveys, when we located a bottlenose dolphin group (one or more individuals in close

proximity and engaging in similar behaviours such as travelling and foraging) we collected photo-

identification data following a standardised protocol (Cheney *et al.* 2014). We identified individuals

from high-quality photographs based on unique markings matched against a photo catalogue of

previously identified individuals from the area (Würsig & Jefferson 1990; Wilson *et al.* 1999; Cheney

et al. 2013). On average, in a group 84% are successfully photographed, and all individual

identifications are confirmed by at least two experienced researchers. For individuals first sighted as

calves we could determine their year of birth and so their age (Cheney *et al.* 2022), but for

individuals first sighted as juveniles or adults their exact age is unknown. Sex was determined using

genital photographs or if an adult was seen in repeat associations with a known calf (Cheney *et al.*

2022).

Social network construction

Individuals sighted during the same encounter were assumed to be in the same group and therefore

associating (known as the "Gambit of the Group"; Whitehead & Dufault 1999). Aggregating many of

these records of groups allows one to infer which individuals are frequently associated and which

individuals infrequently or never associate. We removed observations of individuals younger than

three years old (n = 2634 observations of 233 individuals), as these individuals are not likely to be

independent of the mother and so their social associations most likely represent her preferences.

We then converted the records of encounters into group by individual matrices (indicating which

individuals were seen together in each encounter) and then into weighted, undirected social

networks using the R package *asnipe* (Farine 2013). Edge weights were set as the simple ratio index,

where the number of times two individuals are seen together is divided by the total number of times

they are seen, both together and apart (Cairns & Schwager 1987). This measure ranges from 0

(individuals never seen together) to 1 (individuals always seen together). We did this separately for

each year, creating a social network per year to assess how social phenotypes vary at the yearly

scale in response to environmental conditions. To assess how social phenotypes vary at the monthly

scale in response to environmental conditions we then reconstructed social networks per month and

removed any months with fewer than 10 encounters (excluding 98 out of 217 months; as networks

constructed using fewer than 10 observations can be biased; Franks *et al.* 2010). Histograms of the

frequency of the number of encounters per year and per month are shown in Fig. S1.

For each individual present in the network for each year, we calculated the following: strength (the

sum of all an individual's associations, representing its gregariousness), weighted clustering

coefficient (the rate at which an individual associates with other individuals who also associate with

each other, representing how tightly individuals in its immediate social environment interact), and

closeness (the mean of the path lengths between that individual and each other individual in the

network, corrected for network size to allow comparisons among networks which vary in the

number of individuals, representing its connectedness to the wider population). We removed an

individual's scores for a given year if they had fewer than three observations in that year (removing

498 and leaving 1133), as the social network position of those individuals would be highly uncertain.

They would however still contribute to the social environments and therefore social network

measures of individuals in that year who had three or more observations. We repeated this at the

month level, again removing individuals' monthly scores when they had fewer than three

observations that month (removing 2215 and leaving 1704). Histograms of the frequency of the

number of encounters per individual per year and per month are shown in Fig. S2. We also removed

instances where an individual's closeness was over 1, which occurs when an individual is only

connected to a single other (n = 2 in the yearly networks, n = 18 in the monthly networks). The social

network measures were not well correlated; Pearson correlations between individuals' strength and

clustering coefficient were 0.136 (yearly) and 0.101 (monthly), for strength and closeness they were

-0.042 (yearly) and -0.314 (monthly), and for clustering coefficient and closeness they were 0.157

(yearly) and 0.032 (monthly).

Environmental data

Following Lusseau *et al.* (2004), we used the North Atlantic Oscillation (NAO) index as a measure of

climate. We used monthly and yearly measures of the NAO index between 1990 and 2020

downloaded from https://www.cpc.ncep.noaa.gov/products/precip/CWlink/pna/nao.shtml (Fig. S3).

This index indicates the atmospheric pressure difference between the low pressure zone over

Iceland and the high pressure zone over the Azores (Hurrell 1995; Stenseth *et al.* 2003). This index

has frequently been linked to the ecology of animal populations (Gordo *et al.* 2011; Wan *et al.* 2022),

for example influencing the foraging behaviour of Cory's shearwaters (Calonectris borealis; Pereira

et al. 2020). While climatic effects on cetaceans are typically thought to occur via changes in prey

species (Víkingsson *et al.* 2015; Kebke *et al.* 2022), a phenomenon also observed in our study

population (Lusseau *et al.* 2004), it is also possible that cetaceans respond directly to climate,

sometimes at even faster rates than their prey species (Sprogis *et al.* 2018; Thorne & Nye 2021;

Methion *et al.* 2022).

For our index of food availability we again followed Lusseau *et al.* (2004), using data on monthly

catches by fishing rods (as opposed to nets) from the wild of both one season and multiple season

adult Atlantic salmon (*Salmo salar*) from the Alness, Beauly, Canon, Ness, and Nairn rivers. These

feed into the sea where observations take place and hence are expected to be a good proxy for

salmon availability in that area. Further, catches on rods are positively correlated among rivers and

among months and with automatic counter data, indicating they are a good proxy for actual

abundance (Youngson *et al.* 2002; Thorley *et al.* 2005). Atlantic salmon are an important food source

for this dolphin population (Janik 2000; Santos *et al.* 2001), with dolphins forming larger groups

when salmon are more abundant (which is correlated with the two-year lagged winter NAO index;

Lusseau *et al.* 2004). We downloaded monthly data from https://scotland.shinyapps.io/sg-salmon-

sea-trout-catch/ (summing retained and released fish) and summed monthly catches within a

calendar year for yearly measures of fish abundance (Fig. S4).

All analyses were performed in R (ver 4.1.0; R Development Core Team & Team 2016) using linear

mixed effect models in *glmmTMB* (Brooks *et al.* 2017). Using regression-based models as opposed to

randomisation-based tests has been recommended for analysing questions about node-level social

network traits as it improves the ability to make inference while accounting as well as node

permutations for common types of data non-independence (Hart *et al.* 2021). We fitted 12 models,

with all combinations across the three social traits, the two environmental variables, and the yearly

and monthly timescales. Clustering coefficient cannot be calculated when an individual only

associates with one other individual, and closeness cannot be calculated for individuals without any

connections, and so the datasets for these models were slightly smaller than the dataset for strength

(see below). We included individuals with unknown birth and death dates to maximise our sample

size, and so neither age nor lifespan could be included as predictor variables. In all models we

included the fixed effect of sex (as males and females can differ in social behaviour; Shane *et al.*

1986; Mann *et al.* 2012; ranging behaviour, and survival; Arso Civil *et al.* 2019a), and either the NAO

index or the count of caught salmon for that month or year. We did not include both the NAO index

and salmon abundance in the same model as we encountered estimation problems with two

random slopes. Including sex meant we excluded individuals of unknown sex (258 observations of

103 individuals for the yearly networks, 281 observations of 91 individuals for the monthly

networks), but the interactions between individuals of known and unknown sex were still used to

build the networks and so associations with individuals of unknown sex still influenced the social

network traits of males and females. We mean centred and scaled to unit variance the

environmental variable (Schielzeth 2010), and included the interaction between it and sex, to see if

male and female social behaviour responded differently to environmental variation. Random effects

were the random intercept for individual ID, the random slope of individual ID with the

environmental variable fitted as a fixed effect in the model (again mean centred and with a standard

deviation of 1), and the correlation between these two terms. We also included a temporal

autocorrelation term (ar1) among years to account for variation among years and that adjacent

years may be more similar than non-adjacent years. Similarly, in the models for monthly variation

we included a random effect of month. We used a Gaussian error structure for all models. For

strength and closeness, we used log link functions as the distributions were right skewed, adding

0.001 to all strength measures before transformation to allow us to include zero values (although

zeroes were rare; n = 1 for the yearly analysis and n = 6 for the monthly analysis). For clustering

coefficient, which is bounded between 0 and 1, we used a logit link function, which is preferable to

an arcsine transformation when handling response variables bounded in this way (Warton & Hui

2011). We used the default optimising algorithm for all models except the model for strength in

response to monthly variation in salmon abundance, and the models for clustering coefficient in

response to monthly variation in both NAO index and salmon abundance, where we used the "BFGS"

optimiser, as otherwise the model did not converge (Brooks *et al.* 2017).

We used the p-values from the Anova function of the *car* package (Fox & Weisberg 2019), using a

Chi-squared test with type III sum of squares. We describe these p-values in terms of "clarity" rather

than "significance", see Dushoff et al. (2019) for a discussion on this. To test whether including

random slopes improved our models, and so whether individuals clearly differed in their response to

the environmental variable, for each model (12 in total) we re-fitted it without the random slope

(and the covariance between random intercepts and slopes) and conducted a likelihood ratio test

between the two models. If random slopes did not improve the model we concluded the intercept-

slope correlation was zero.

Results

Change with environmental variables at a yearly scale

For the analysis of how dolphin social phenotypes change at the yearly scale in response to

environmental variables, our dataset included 155 unique individuals. For strength there were 1133

measures, for clustering coefficient there were 1129 and for closeness there were 1132. Traits had a

mean of 7.31 (sd = 5.99), 7.28 (sd = 5.92), and 7.30 (sd = 5.99) measures per individual for strength,

clustering coefficient and closeness respectively

Dolphins' strength and clustering coefficient were not affected by the NAO index (Fig. 2a and b) or

salmon abundance (Fig. 3a and b) in both sexes, and the sexes did not differ in mean strength or

clustering coefficient (Table 1, full model results in Tables S1-6, supplementary materials). In

addition, the random slopes were not statistically clear, suggesting that individual dolphins did not

differ in their (lack of) response to the NAO index. A lack of variation in individual slopes leads us to

assume that the intercept slope correlations for the NAO index were not different from zero (Table

2), indicating an individual's mean behaviour is not related to how it responds to the NAO index. But

individuals did differ in their responses to salmon abundance, with a negative intercept slope

correlation for strength (individuals with lower mean strength increased their strength at higher

values of the NAO index) and a positive one for clustering coefficient (individuals with lower mean

clustering coefficient decreased it with increasing NAOI index; Table 2).

Similarly, closeness did not vary with NAO index (Fig. 2c) or salmon numbers (Fig. 3c) in either sex

(Table 1). Although females may have had higher mean closeness than males, the effect was not

statistically clear (Supplementary Material Tables S5 & S6). Again, individuals did not differ in their

response to the NAO index, but they did show different responses to salmon abundance (Table 2).

Like strength, individuals with lower mean closeness increased their closeness more with increasing

salmon abundance, and there was no relationship between an individual's mean closeness and its

response to the NAO index model (Table 2).

Figure 2. Plots of each of the three social network traits and yearly variation in the North Atlantic

Oscillation (NAO) index (a. strength, b. clustering coefficient, c. closeness). For each individual

dolphin we have predicted its network trait based on the model results, using the "predict" function

in R with the individual's sex, the range of NAO values that individual was exposed to, and setting

the year to 2005.

Figure 3. Plots of each of the three social network traits and yearly variation in salmon abundance (a.

strength, b. clustering coefficient, c. closeness). For each individual dolphin we have predicted its

network trait based on the model results, using the "predict" function in R with the individual's sex,

the range of salmon counts that individual was exposed to, and setting the year to 2005.

		Strength	Clustering coefficient	Closeness
NAO index	Yearly	M: β = 0.013 ± 0.075, χ^2 = 0.032, p = 0.858	M: β = 0.012 ± 0.086, χ^2 = 0.019, p = 0.891	M: β = -0.053 ± 0.093, χ^2 = 0.324, p = 0.569
		I: β = 0.009 ± 0.023, χ^2 = 0.163, p = 0.687	I: β = 0.005 ± 0.034, χ^2 = 0.019, p = 0.889	I: β = -0.011 ± 0.013, χ^2 = 0.733, p = 0.392
	Monthly	M: β = 0.029 ± 0.021, χ^2 = 1.902, p = 0.168	M: β = 0.038 ± 0.038, χ^2 = 0.968, p = 0.325	M: *β = -0.077 ± 0.044, χ^2 = 3.004, p = 0.083*
		I: β = -0.002 ± 0.024, χ^2 = 0.009, p = 0.924	I: β = 0.009 ± 0.042, χ^2 = 0.045, p = 0.832	I: *β = 0.103 ± 0.053, χ^2 = 3.724, p = 0.054*
Salmon	Yearly	M: β = 0.123 ± 0.077, χ^2 = 2.523, p = 0.112	M: β = 0.040 ± 0.088, χ^2 = 0.019, p = 0.891	M: β = -0.142 ± 0.088, χ^2 = 2.609, p = 0.106
		I: β = -0.009 ± 0.031, χ^2 = 0.091, p = 0.762	I: β = 0.012 ± 0.043, χ^2 = 0.019, p = 0.890	I: β = -0.011 ± 0.013, χ^2 = 0.733, p = 0.392
	Monthly	M: β = 0.015 ± 0.034, χ^2 = 0.199, p = 0.655	M: **β = 0.170 ± 0.051, χ^2 = 11.290, p = 0.001**	M: β = 0.019 ± 0.064, χ^2 = 0.086, p = 0.769
		I: β = 0.050 ± 0.035, χ^2 = 2.131, p = 0.144	I: β = -0.060 ± 0.049, χ^2 = 1.495, p = 0.221	I: β = 0.006 ± 0.069, χ^2 = 0.006, p = 0.936

Table 1. Main effects (prefixed with "M:") and interactions with sex (prefixed with "I:" and shaded in

grey) for the two environmental variables effects on the three social network traits, at each of the

monthly and yearly scales. Effects clearly different from zero (p < 0.05) are highlighted in bold, while

effects with p values between 0.1 and 0.05 are highlight in italics.

For all social traits there was substantial among-year variation and consecutive years were positively

correlated (year to year correlations: strength; r_{NAO} = 0.314, r_{Salmon} = 0.429; clustering coefficient; r_{NAO}

= 0.559, r_{Salmon} = 0.560; and closeness; r_{NAO} =0.455, r_{Salmon} = 0.543), showing that, as expected,

adjacent years were more similar than non-adjacent years.

In summary, there were no trait-environment association at the yearly scale. However, individuals

always differed in their response to salmon abundance but never in their (lack of) response to the

NAO index.

		Strength	Clustering coefficient	Closeness
NAO index	Yearly	- (χ^2 = 1.5345, df = 2, p = 0.464)	- (χ^2 = 0.896, df = 2, p = 0.639)	- (χ^2 = 1.490, df = 2, p = 0.475)
	Monthly	-0.963 (χ^2 = 6.904, df = 2, p = 0.032)	- (χ^2 = 1.632, df = 2, p = 0.442)	-0.792 (χ^2 = 49.643, df = 2, p < 0.001)
Salmon	Yearly	-0.321 (χ^2 = 18.927, df = 2, p < 0.001)	0.268 (χ^2 = 8.568, df = 2, p = 0.014)	-0.510 (χ^2 = 71.013, df = 2, p < 0.001)
	Monthly	-0.299 (χ^2 = 31.096, df = 2, p < 0.001)	- (χ^2 = 1.946, df = 2, p = 0.378)	-0.350 (χ^2 = 194.29, df = 2, p < 0.001)

Table 2. Intercept slope correlations and statistical tests of clarity of random slopes. Positive

correlations in blue, negative in red, those where slopes were not clearly different from zero and so

intercept slope correlation assumed to be not different from zero are indicated with a dash and not

coloured in.

Change with environmental variables at a monthly scale

We analysed how dolphin social phenotypes change in response to environmental variables at the

monthly scale with a dataset of 151 unique individuals for strength and closeness and 149 for

clustering coefficient. For strength there were 1704 measures, for clustering coefficient 1688, and

for closeness 1696. Traits had a mean of 11.3, 11.3, and 11.2 measures per individual for strength,

clustering coefficient, and closeness respectively (sd = 10.7 for all).

Strength was not affected by the monthly NOA index (Fig. 4a) or monthly salmon abundance (Fig. 5a)

in either sex and the sexes did not differ in their mean strength (Table 1, full model results in Tables

S7-12 in the supplementary materials). Individuals differed in their responses to the monthly NAO

index and salmon abundances, with individuals with high values decreasing their strength with

higher values of the environmental factor and individuals with low values increasing their strength

(Table 2).

Months of higher salmon abundance led to higher clustering coefficients in both sexes (Figs. 5b & 6

and Table 1). Clustering coefficient did not however vary in response to monthly NOA index in either

sex (Fig. 4b and Table 1). The sexes did not differ in their mean clustering coefficient. Individuals did

not vary in their response to salmon abundance, or in their lack of response to the NAO index, and

so in both cases the intercept-slope correlation is not different from zero (Table 2).

Figure 4. Plots of each of the three social network traits and monthly variation in the North Atlantic

Oscillation (NAO) index (a. strength, b. clustering coefficient, c. closeness). In c., males are shown in

purple, females in blue, to visualise the possible difference between the sexes (see Results). For each

individual dolphin we have predicted its network trait based on the model results, using the

"predict" function in R with the individual's sex, the range of NAO values that individual was exposed

to, setting the year to 2005, and the month to June.

Females had a lower average closeness than males, although this effect was only clear in the model

for the NAO index (Table S11). Further, the sexes may have differed in their response to the NAO

index; while males showed a marginally unclear decrease with higher NAO index, females may have

shown no change or even a slight increase (Fig. 4c and Table 1). Closeness did not respond to

monthly salmon numbers in either sex (Fig. 5c and Table 1). Individuals differed in their responses to

the NAO index and to salmon abundance (Table 2), with individuals with high values decreasing their

closeness with higher values of the environmental factor and individuals with low values increasing

their closeness.

Figure 5. Plots of each of the three social network traits and monthly variation in salmon abundance

(a. strength, b. clustering coefficient, c. closeness). For each individual dolphin we have predicted its

network trait based on the model results, using the "predict" function in R with the individual's sex,

the range of salmon counts that individual was exposed to, setting the year to 2005, and the month

to June.

There was among-month and among-year variation, and consecutive years were always positive

correlated (year to year correlations: strength: r_{NAO} = 0.553, r_{Salmon} = 0.529; clustering coefficient: r_{NAO}

= 0.228, r_{Salmon} =0.074; closeness: r_{NAO} = 0.467, r_{Salmon} = 0.475).

In summary, months with higher salmon numbers led to higher clustering coefficient in both sexes

but did not impact strength or closeness. Meanwhile, months with a higher NAO index were

associated with a higher closeness in females and a lower closeness in males, although both these

effects were not clearly different from zero. The NAO index did not affect strength or clustering

coefficient. Individuals differed in how their strength and closeness changed in response to both the

NAO index and salmon abundance, but they did not differ in how their clustering coefficient changed

for either variable.

Figure 6. Plots of dolphin social networks in a month of low salmon abundance (May 1990) and a

month of high salmon abundance (August 2007). Circles are individual dolphins and grey lines

indicate associations i.e., those seen in the same group at least once.

Discussion

We explored whether bottlenose dolphin social behaviour responded to environmental variation.

Typically, social behaviour was stable, with no variation at the yearly scale in response to the NAO

index or salmon abundance. Similarly, two social behaviours showed no variation to environmental

change at the monthly scale, although, how tightly individuals in the immediate social environment

interacted (clustering coefficient) did increase in response to increases in monthly salmon

abundance. We also showed that, alongside differences in mean behaviour, individuals differed in

their behavioural responses to changes in salmon abundance at both yearly and monthly scales, and

to the NAO index at the monthly scale, indicating variation in plasticity. This plasticity was often, but

not always, negatively associated with mean behaviour, causing among individual differences to

typically be greater for low values of salmon abundance and the NAO index. Interestingly, when

males and females increased their clustering coefficient in response to higher monthly salmon

abundances, there was limited among-individual variation in this change. This conformity of

response perhaps suggests a particularly important or unavoidable change. Otherwise, the stability

of social behaviour implies that maintaining a consistent social structure is important (see also Evans

et al. 2021 who found consistency over lifetime in the social behaviours of Indo-Pacific bottlenose

dolphins, T. aduncus), and so testing the importance of social network stability on fitness in dolphins

is key.

Despite the population-level stability, in most cases individual dolphins differed in how their social

behaviour changed in response to environmental variation (i.e., they showed among-individual

variation in plasticity). Therefore, while the population as a whole showed no overall change, some

individuals would show an increase in the social behaviour, while others a decrease. Variation in

individual plasticity leads to environment-dependent repeatability (and possibly heritability), can

dampen population responses to environmental variability, and enhance population persistence

(Dingemanse & Wolf 2013). For example, due to the negative intercept-slope correlation, individual

strength shows the most among-individual variation at low monthly salmon abundances. This wider

range of social phenotypes could increase the possibility that at least some individuals are successful

despite low food availability. Determining the genetic basis and importance of early life conditions

for the development of these different responses (Stamps 2016) and how this variation impacts

population dynamics (Vindenes *et al.* 2008) is key. Models projecting how dolphin populations will

change in the future should account for both among-individual variation in mean behaviour and in

behavioural plasticity.

In general, the lack of plasticity in social network traits suggests dolphin social behaviour will show

limited changes at the population level if salmon abundances or climate change over time (based on

similar changes to environmental variables studied here). Populations of animals with relatively long

generation times are more reliant on phenotypic plasticity than adaptive evolution to track rapid

environmental change (Hoffmann & Sgrò 2011). Therefore, if dolphins do not change plastically

there is limited scope for evolution to compensate to enhance phenotype-environment fit. If social

behaviours are important for coping with detrimental environments, this could potentially be

problematic, leading to reduced reproduction and possibly population decline.

The one exception to stability was that individual clustering coefficient increased in months with

higher salmon abundance (see Fig. 6 for a comparison of months with low vs. high salmon

abundance). This indicates that dolphins were "closing triads" and interacting more often with those

who they shared a mutual third association. The increase in clustering coefficient occurred without a

change in strength or closeness, suggesting a nuanced change in behaviour. Interestingly, individuals

did not show variability in this increase, suggesting it is important or unavoidable for all. Similar

results have been found in spotted hyenas (*Crocuta crocuta*) where seasons with high prey

availability have denser social networks (Holekamp *et al.* 2012). However, other systems have found

lower food availability leads to decreased sociability (Randall *et al.* 2005; Williams *et al.* 2017), and it

is presumed this response reduces the intensity of resource competition (Lin *et al.* 2019; reviewed

in: Fisher *et al.* 2021). Therefore, higher clustering may be beneficial for non-foraging reasons, such

as mating. Both mating and calving seasons approximately align with the months of highest salmon

abundance in the summer (Fig. S4), supporting this suggestion. Dolphins could also reduce travel

distances when food availability is high (Shane *et al.* 1986), leading to them stay in the same place

and so interact more with a tight circle of others; increasing their clustering coefficient.

Alternatively, it is possible that months with fewer salmon also differ in an unidentified variable

which causes dolphins to cluster less. However, it is unlikely that this unidentified factor is predation

threat that changes month to month, as predators are absent in this area (Cheney *et al.* 2019).

Therefore, a change in social behaviour and/or movement related to the seasonal availability of

salmon, perhaps influenced by reproductive behaviour, seems the most likely.

Notwithstanding the response of dolphins to changes in environmental factors on a monthly scale,

we did not see any responses to environmental variation on a yearly scale. Both factors varied

considerably year-to-year; the NAO index ranged from -2.35 to 1.23 (Fig. S3) and salmon abundance

varied by more than 2.8 times between years (Fig. S4), hence a lack of relevant variability seems

unlikely. Lusseau *et al.* (2004) observed larger groups in years of higher salmon abundance, but this

was over fewer years and was not an individual-level analysis. The robustness or inflexibility of social

behaviour in response to year-to-year variation in climatic conditions and food availability might

indicate that the yearly environmental variation is inconsequential and so they have no need to

respond to it. However, it seems unlikely that variation in food availability of 2.8 times does not

matter to this population, as their distribution in space reflects foraging opportunities (Hastie *et al.*

2004). Alternatively, dolphins may change other phenotypes, such as foraging behaviour or

metabolism, to cope with these stressors (Kebke *et al.* 2022), leaving social behaviour unchanged.

For example, salmon are not this dolphin population's only prey (Santos *et al.* 2001), and so they

may switch to other prey if salmon are scarce, obviating the need to modify their social networks.

We estimated correlations between an individual's mean behaviour (its intercept) and its plasticity in

behaviour (its slope). For strength and closeness, intercept-slope correlations were negative if they

were different from zero. Therefore, individuals typically converged to similar behaviour at higher

NAO index values and salmon abundances. Negative intercept-slope correlations can constrain

evolution if higher mean values *and* higher plasticities are favoured, as selection is favouring a

combination of phenotypes that is rare (Dingemanse & Wolf 2013). In contrast, intercept-slope

correlations for clustering coefficient were either zero or, in the case of the response to salmon

abundance at the yearly scale, positive. The positive correlation shows behaviours diverging among

individuals at higher yearly salmon abundances. Additionally, while a positive intercept slope

correlation means that selection on either intercept or slope causes the other to increase too, no

correlation means intercepts and slopes can be independent targets of selection. Kebke *et al.* (2022)

suggest that cetacean ranging and foraging behaviour may well be under selection for increased

plasticity as environments change, and so testing this directly is a logical and important next step.

In conclusion, we found that at the population-level dolphin social behaviour is relatively stable

across years despite year-to-year variation in environmental conditions. At the monthly scale, we

observed all individuals increased clustering of local social associations due to higher salmon

abundance. However, at both monthly and yearly scales there was lots of variability at the individual

level in responses to environmental factors. As such, individual heterogeneity in both mean and

plasticity in behaviour both should be more widely accounted for when predicting species responses

to environmental change, and individual plasticities as well as means may be targets of selection and

hence evolvable.

Acknowledgements

We are indebted to Paul Thompson for conception and development of this long-term individual

based research study, and for advice during this work. We are especially grateful to our colleagues at

the Lighthouse Field Station, past and present, who have contributed to and advanced this long-term

study, and to Charlie Phillips who contributed land-based photographs for photo-identification.

Matthew Silk made numerous useful comments and suggestions on an earlier draft. This long-term

study has depended upon a number of funders including University of Aberdeen, NatureScot,

Beatrice Offshore Windfarm Ltd., Moray Offshore Renewables Ltd., Marine Scotland, the Crown

Estate, Highlands and Islands Enterprise, the BES, ASAB, Greenpeace Environmental Trust, Whale

and Dolphin Conservation, Talisman Energy (UK) Ltd., Department of Energy and Climate Change,

Chevron and Natural Environment Research. Survey work was conducted under NatureScot Animal

Scientific Licences. We have no conflicts of interest.

Statement of authorship

441   DF conceived the idea for the manuscript, BC provided data, DF ran analyses with guidance from BC,

442   DF wrote first draft of the manuscript and both authors contributed to revisions.

Data accessibility statement

Data and R code used for the analysis will be made publicly available upon acceptance.

References

Abrahms, B., Rafiq, K., Jordan, N.R. & McNutt, J.W. (2022). Long-term, climate-driven phenological shift in a tropical large carnivore. *Proceedings of the National Academy of Sciences*, 119.

Araya-Ajoy, Y.G. & Dingemanse, N.J. (2017). Repeatability, heritability, and age-dependence of seasonal plasticity in aggressiveness in a wild passerine bird. *Journal of Animal Ecology*, 86, 227–238.

Arso Civil, M., Cheney, B., Quick, N.J., Islas-Villanueva, V., Graves, J.A., Janik, V.M., *et al.* (2019a). Variations in age- and sex-specific survival rates help explain population trend in a discrete marine mammal population. *Ecology and Evolution*, 9, 533–544.

Arso Civil, M., Quick, N.J., Cheney, B., Pirotta, E., Thompson, P.M. & Hammond, P.S. (2019b). Changing distribution of the east coast of Scotland bottlenose dolphin population and the challenges of area-based management. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 29, 178–196.

Arso Civil, M., Quick, N.J., Mews, S., Hague, E., Cheney, B.J., Thompson, P.M., *et al.* (2021). *Improving understanding of bottlenose dolhin movements along the east coast of Scotland.* (Final Report No. SMRUC-VAT-2020-10). European Offshore Wind Deployment Centre (EOWDV).

Avila, I.C., Kaschner, K. & Dormann, C.F. (2018). Current global risks to marine mammals: Taking stock of the threats. *Biological Conservation*, 221, 44–58.

Brommer, J.E. (2013a). Phenotypic plasticity of labile traits in the wild. *Current Zoology*, 59, 485–505.

Brommer, J.E. (2013b). Variation in plasticity of personality traits implies that the ranking of personality measures changes between environmental contexts: calculating the cross-environmental correlation. *Behavioral Ecology and Sociobiology*, 67, 1709–1718.

Brooks, M.E., Kristensen, K., van Benthem, K.J., Magnusson, A., Berg, C.W., Nielsen, A., *et al.* (2017). glmmTMB Balances speed and flexibility among packages for modeling. *The R Journal*, 9, 378–400.

Cairns, S.J. & Schwager, S.J. (1987). A comparison of association indices. *Animal Behaviour*, 35, 1454–1469.

Cheney, B., Corkrey, R., Durban, J.W., Grellier, K., Hammond, P.S., Islas-Villanueva, V., *et al.* (2014). Long-term trends in the use of a protected area by small cetaceans in relation to changes in population status. *Global Ecology and Conservation*, 2, 118–128.

Cheney, B., Thompson, P.M., Ingram, S.N., Hammond, P.S., Stevick, P.T., Durban, J.W., *et al.* (2013). Integrating multiple data sources to assess the distribution and abundance of bottlenose dolphins Tursiops truncatus in Scottish waters. *Mammal Review*, 43, 71–88.

Cheney, B.J., Dale, J., Thompson, P.M., Quick, N.J. & Barbara Cheney, C.J. (2022). Spy in the sky: a method to identify pregnant small cetaceans. *Remote Sensing in Ecology and Conservation*.

Cheney, B.J., Thompson, P.M. & Cordes, L.S. (2019). Increasing trends in fecundity and calf survival of bottlenose dolphins in a marine protected area. *Scientific Reports*, 9, 1767.

Chevin, L.-M. (2013). Genetic constraints on adaptation to a changing environment. *Evolution*, 67, 708–721.

Dingemanse, N.J., Kazem, A.J.N., Réale, D. & Wright, J. (2010). Behavioural reaction norms: animal personality meets individual plasticity. *Trends in Ecology & Evolution*, 25, 81–89.

Dingemanse, N.J. & Wolf, M. (2013). Between-individual differences in behavioural plasticity within
populations: causes and consequences. *Animal Behaviour*, 85, 1031–1039.
Dushoff, J., Kain, M.P. & Bolker, B.M. (2019). I can see clearly now: Reinterpreting statistical
significance. *Methods in Ecology and Evolution*, 10, 756–759.
Evans, P.G. & Bjørge, A. (2013). Impacts of climate change on marine mammals. *Marine Climate*
Change Impacts Partnership (MCCIP) Science Review, 2013, 134–148.
Evans, T., Krzyszczyk, E., Frère, C. & Mann, J. (2021). Lifetime stability of social traits in bottlenose
dolphins. *Commun Biol*, 4, 1–8.
Farine, D.R. (2013). Animal social network inference and permutations for ecologists in R using
asnipe. *Methods in Ecology and Evolution*, 4, 1187–1194.
Fisher, D.N., David, M., Tregenza, T. & Rodriguez-Munoz, R. (2015). Dynamics of among-individual
behavioral variation over adult lifespan in a wild insect. *Behavioral Ecology*, 26, 975–985.
Fisher, D.N., Kilgour, R.J., Siracusa, E.R., Foote, J.R., Hobson, E.A., Montiglio, P.O., *et al.* (2021).
Anticipated effects of abiotic environmental change on intraspecific social interactions.
Biological Reviews, 96, 2661–2693.
Fox, J. & Weisberg, S. (2019). *An R Companion to Applied Regression*. Sage, Thousand Oaks (CA).
Franks, D.W., Ruxton, G.D. & James, R. (2010). Sampling animal association networks with the
gambit of the group. *Behavioral Ecology and Sociobiology*, 64, 493–503.
Gordo, O., Barriocanal, C. & Robson, D. (2011). Ecological Impacts of the North Atlantic Oscillation
(NAO) in Mediterranean Ecosystems. In: *Hydrological, Socioeconomic and Ecological Impacts*
of the North Atlantic Oscillation in the Mediterranean Region, Advances in Global Change
Research (eds. Vicente-Serrano, S.M. & Trigo, R.M.). Springer Netherlands, Dordrecht, pp.
153–170.
Hamel, S., Gaillard, J.-M. & Yoccoz, N.G. (2018). Introduction to: Individual heterogeneity - the
causes and consequences of a fundamental biological process. *Oikos*, 127, 643–647.
Hart, J.D.A., Weiss, M.N., Brent, L.J.N. & Franks, D.W. (2021). Common Permutation Methods in
Animal Social Network Analysis Do Not Control for Non-independence. *bioRxiv*.
Hastie, G.D., Wilson, B., Wilson, L.J., Parsons, K.M. & Thompson, P.M. (2004). Functional mechanisms
underlying cetacean distribution patterns: hotspots for bottlenose dolphins are linked to
foraging. *Marine Biology*, 144, 397–403.
Hoffmann, A.A. & Sgrò, C.M. (2011). Climate change and evolutionary adaptation. *Nature*, 470, 479–
485.
Holekamp, K.E., Smith, J.E., Strelioff, C.C., Van Horn, R.C. & Watts, H.E. (2012). Society, demography
and genetic structure in the spotted hyena. *Molecular Ecology*, 21, 613–632.
Hurrell, J.W. (1995). Decadal Trends in the North Atlantic Oscillation: Regional Temperatures and
Precipitation. *Science*, 269, 676–679.
Janik, V.M. (2000). Food–related bray calls in wild bottlenose dolphins (Tursiops truncatus).
Proceedings of the Royal Society of London. Series B: Biological Sciences, 267, 923–927.
Kebke, A., Samarra, F. & Derous, D. (2022). Climate change and cetacean health: impacts and future
directions. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 377.
Komdeur, J. & Ma, L. (2021). Keeping up with environmental change: The importance of sociality.
Ethology, 127, 790–807.
Lin, Y.H., Chan, S.F., Rubenstein, D.R., Liu, M. & Shen, S.F. (2019). Resolving the paradox of
environmental quality and sociality: The ecological causes and consequences of cooperative
breeding in two lineages of birds. *American Naturalist*, 194, 207–216.
Lusseau, D., Williams, R., Wilson, B., Grellier, K., Barton, T.R., Hammond, P.S., *et al.* (2004). Parallel
influence of climate on the behaviour of Pacific killer whales and Atlantic bottlenose
dolphins. *Ecology Letters*, 7, 1068–1076.
MacLeod, C.D. (2009). Global climate change, range changes and potential implications for the
conservation of marine cetaceans: a review and synthesis. *Endangered Species Research*, 7,
125–136.

Mann, J., Stanton, M.A., Patterson, E.M., Bienenstock, E.J. & Singh, L.O. (2012). Social networks
reveal cultural behaviour in tool-using dolphins. *Nat Commun*, 3, 980.
McGlothlin, J.W., Moore, A.J., Wolf, J.B. & Brodie, E.D. (2010). Interacting phenotypes and the
evolutionary process. III. Social evolution. *Evolution; international journal of organic*
evolution, 64, 2558–74.
Methion, S., Paradell, O.G., Padín, X.A., Corrège, T. & López, B.D. (2022). *Group size varies with*
climate and oceanographic conditions in bottlenose dolphins (preprint). In Review.
Moore, A.J., Brodie, E.D.I. & Wolf, J.B. (1997). Interacting phenotypes and the evolutionary process:
I. Direct and indirect genetic effects of social interactions. *Evolution*, 51, 1352–1362.
Nussey, D.H., Postma, E., Gienapp, P. & Visser, M.E. (2005). Selection on Heritable Phenotypic
Plasticity in a Wild Bird Population. *Science*, 310, 304–306.
Nussey, D.H., Wilson, A.J. & Brommer, J.E. (2007). The evolutionary ecology of individual phenotypic
plasticity in wild populations. *Journal of Evolutionary Biology*, 20, 831–844.
Pereira, J.M., Paiva, V.H., Ceia, F.R. & Ramos, J.A. (2020). Facing extremes: Cory's shearwaters adjust
their foraging behaviour differently in response to contrasting phases of North Atlantic
Oscillation. *Reg Environ Change*, 20, 77.
R Development Core Team & Team, R.D.C. (2016). R: A language and environment for statistical
computing. *R Foundation for Statistical Computing*, R Foundation for Statistical Computing.
Randall, J.A., Rogovin, K., Parker, P.G. & Eimes, J.A. (2005). Flexible social structure of a desert
rodent, Rhombomys opimus: philopatry, kinship, and ecological constraints. *Behavioral*
Ecology, 16, 961–973.
Santos, M.B., Pierce, G.J., Reid, R.J., Patterson, I. a. P., Ross, H.M. & Mente, E. (2001). Stomach
contents of bottlenose dolphins (Tursiops truncatus) in Scottish waters. *Journal of the*
Marine Biological Association of the United Kingdom, 81, 873–878.
Schielzeth, H. (2010). Simple means to improve the interpretability of regression coefficients.
Methods in Ecology and Evolution, 1, 103–113.
Shane, S.H., Wells, R.S. & Würsig, B. (1986). Ecology, Behavior and Social Organization of the
Bottlenose Dolphin: A Review. *Marine Mammal Science*, 2, 34–63.
Simmonds, M.P. & Isaac, S.J. (2007). The impacts of climate change on marine mammals: early signs
of significant problems. *Oryx*, 41, 19–26.
Snyder-Mackler, N., Burger, J.R., Gaydosh, L., Belsky, D.W., Noppert, G.A., Campos, F.A., *et al.* (2020).
Social determinants of health and survival in humans and other animals. *Science*, 368.
Sprogis, K.R., Christiansen, F., Wandres, M. & Bejder, L. (2018). El Niño Southern Oscillation
influences the abundance and movements of a marine top predator in coastal waters. *Global*
Change Biology, 24, 1085–1096.
Stamps, J. & Biro, P. (2016). Personality and individual differences in plasticity. *Current Opinion in*
Behavioral Sciences.
Stamps, J.A. (2016). Individual differences in behavioural plasticities. *Biological Reviews*, 91, 534–
567.
Stenseth, N.Chr., Ottersen, G., Hurrell, J.W., Mysterud, A., Lima, M., Chan, K., *et al.* (2003). Review
article. Studying climate effects on ecology through the use of climate indices: the North
Atlantic Oscillation, El Niño Southern Oscillation and beyond. *Proceedings of the Royal*
Society of London. Series B: Biological Sciences, 270, 2087–2096.
Thorley, J.L., Eatherley, D.M.R., Stephen, A.B., Simpson, I., MacLean, J.C. & Youngson, A.F. (2005).
Congruence between automatic fish counter data and rod catches of Atlantic salmon (Salmo
salar) in Scottish rivers. *ICES Journal of Marine Science*, 62, 808–817.
Thorne, L.H. & Nye, J.A. (2021). Trait-mediated shifts and climate velocity decouple an endothermic
marine predator and its ectothermic prey. *Sci Rep*, 11, 18507.
Tyack, P.L. (2008). Implications for marine mammals of large-scale changes in the marine acoustic
environment. *Journal of Mammalogy*, 89, 549–558.

Víkingsson, G.A., Pike, D.G., Valdimarsson, H., Schleimer, A., Gunnlaugsson, T., Silva, T., *et al.* (2015).
Distribution, abundance, and feeding ecology of baleen whales in Icelandic waters: have
recent environmental changes had an effect? *Frontiers in Ecology and Evolution*, 3.
Vindenes, Y., Engen, S. & Sæther, B. (2008). Individual Heterogeneity in Vital Parameters and
Demographic Stochasticity. *The American Naturalist*, 171, 455–467.
Vindenes, Y. & Langangen, Ø. (2015). Individual heterogeneity in life histories and eco-evolutionary
dynamics. *Ecology Letters*, 18, 417–432.
Wan, X., Holyoak, M., Yan, C., Le Maho, Y., Dirzo, R., Krebs, C.J., *et al.* (2022). Broad-scale climate
variation drives the dynamics of animal populations: a global multi-taxa analysis. *Biological*
Reviews, n/a.
Warton, D. & Hui, F. (2011). The arcsine is asinine: the analysis of proportions in ecology. *Ecology*.
van Weelden, C., Towers, J.R. & Bosker, T. (2021). Impacts of climate change on cetacean
distribution, habitat and migration. *Climate Change Ecology*, 1, 100009.
White, D.J., Gersick, A.S. & Snyder-Mackler, N. (2012). Social networks and the development of social
skills in cowbirds. *Philosophical Transactions of the Royal Society of London B: Biological*
Sciences, 367.
Whitehead, H. & Dufault, S. (1999). Techniques for Analyzing Vertebrate Social Structure Using
Identified Individuals: Review and Recommendations. *Advances in the Study of Behavior*, 28.
Williams, A.E., Worsley-Tonks, K.E.L. & Ezenwa, V.O. (2017). Drivers and consequences of variation in
individual social connectivity. *Animal Behaviour*, 133, 1–9.
Wilson, B., Hammond, P.S. & Thompson, P.M. (1999). Estimating Size and Assessing Trends in a
Coastal Bottlenose Dolphin Population. *Ecological Applications*, 9, 288–300.
Wolf, J.B., Brodie III, E.D. & Moore, A.J. (1999). Interacting Phenotypes and the Evolutionary Process.
II. Selection Resulting from Social Interactions. *The American Naturalist*, 153, 254–266.
Würsig, B. & Jefferson, T.A. (1990). *Methods of photo-identification for small cetaceans* (No.
SC/A88/ID13). Reports of the International Whaling Commission.
Youngson, A.F., MacLean, J.C. & Fryer, R.J. (2002). Rod catch trends for early-running MSW salmon in
Scottish rivers (1952–1997): divergence among stock components. *ICES Journal of Marine*
Science, 59, 836–849.
619