# Dolphin social phenotypes typically show individual variation in

# response to environmental change alongside population-level stability

David N. Fisher*

School of Biological Sciences, University of Aberdeen, King's College, Aberdeen, AB243FX

* corresponding author: david.fisher@abdn.ac.uk

Barbara J. Cheney

School of Biological Sciences, University of Aberdeen, Lighthouse Field Station, George Street,
Cromarty, IV11 8YL

## Abstract

Social behaviours can allow individuals to flexibly respond to environmental change, potentially

buffering adverse effects. However, individuals may respond differently to the same environmental

stimulus, complicating our ability to predict population-level response to environmental change.

Here we show that bottlenose dolphins (*Tursiops truncatus*) show a high degree of individual

variability in responses of social network traits to variation in proxies of food availability (salmon

abundance) and climate (the North Atlantic Oscillation index), against a background of population-

level stability. Individual dolphins did not reliably respond to yearly variation in environmental

factors and showed limited consistency in their overall gregariousness or connectedness to distant

parts of the network. However, individuals showed consistent differences in mean connectedness to

their immediate social environment and at the monthly scale these connections became denser in

months of high salmon abundance. Negative relationships between mean behaviour and plasticity in

behaviour suggest greater levels of variation during times of low North Atlantic Oscillation index or

salmon abundance. These results indicate that forecasting population responses to environmental

change should account for among-individual variation in both mean and plasticity of social

behaviour.

29    **Keywords**: dolphin, environmental change, individual variation, plasticity, social network, Tursiops

## Introduction

Animals engage in social interactions with conspecifics which are fundamental for determining

health, access to resources, and reproductive success (1). Consequently, social interactions have a

strong influence on ecological processes such as population dynamics and evolutionary processes

such as the response to selection (2–4). To maintain the best fit with their environment, animals may

adjust their social behaviour as conditions change (5), for instance being more gregarious when

resources are plentiful but less tolerant of conspecifics when resources are scarce (6). Animals may

also change their behaviour through development and during senescence (7). Such plasticity is a

hallmark of behavioural traits and gives behaviour an important role in how animals interact with

their environment.

When trying to understand how animals may respond through plasticity to changing environments,

most examine responses at the population level (e.g., 8), implying that all individuals respond

similarly. However, individuals typically show variation in plasticity, and so each will respond

differently to change (9,10, sometimes referred to as 'I x E' i.e., individual by environment

interactions,11,12). For example, European field crickets (*Gryllus campestris*) become bolder and

more active as they age, but individuals vary in the extent of this, with some not increasing or even

decreasing (13). The degree of plasticity animals show can be correlated with their mean behaviour

(an "intercept-slope correlation") which determines how the magnitude of among-individual

differences varies across environments and indicates the extent to which plasticity is a separate trait

in its own right (10,14). Individuality in plasticity can influence biological processes such as

population growth and adaptive change at a range of scales (15,see: 16 and accompanying papers),

giving fundamentally different results to when population-level only effects are assumed (14,17).

Additionally, the extent of among-individual variation in plasticity gives an upper-limit to the

heritability of plasticity, which indicates how rapidly plasticity itself can evolve (18–20).

Understanding how plasticity as a trait in its own right can evolve is key in understanding how

animals will adapt to more variable climates (20,21).

Marine mammals are a key group to study individual variation in response to environmental change.

They are typically long-lived, increasing the relative importance of plasticity versus adaptive

evolution for coping with contemporary environmental change (22). They are also exposed to a wide

range of changing conditions including climate, food availability, and pollution, and their populations

are often of conservation concern. All of these factors increase the need for us to understand how

they respond to changes in their environment (23–30).

Here, we studied a population of bottlenose dolphins (*Tursiops truncatus*) in the North Sea for 30

years, and regularly recorded their social associations. We use social network analysis to quantify

three different dimensions of individual social behaviour: an individual's gregariousness (strength),

how tightly its immediate social group interact together (clustering coefficient), and how well

connected an individual is to the entire population (closeness). We then used random regression

models to quantify individual social phenotypes and determine how these social phenotypes depend

on yearly and monthly variation in available proxies for climate and food availability. These models

also indicated whether bottlenose dolphins show individual changes in response to the

environment, or if population-level change was more prominent.

## Material and methods

### Study site & group data collection

This study used data from a bottlenose dolphin population on the east coast of Scotland (Fig. 1). The

population of over 200 individuals (31) has been studied intensively as part of a long term individual

based study (32–34). We use data from boat-based photo-identification surveys carried out annually

between 1990 and 2020 which regularly recorded dolphin groups within the Moray Firth Special

Area of Conservation (SAC) (92/43/EEC) (35). The SAC is a core part of the population's range with

over 50% of the population using this area each year, with regular annual summer survey effort

aimed at maximising sighting probability, and consistent and adaptive protocols (35). Additional data

were also available from Whale and Dolphin Conservation land based photo-identification surveys

(33). Data were available from a total of 913 surveys (between 10 and 90 surveys each year; average

of 30) with the majority carried out between May to September.

**Figure 1**. Images depicting a photo-identification survey (top let), a newborn calf with mother

(middle left), unique markings used to identify individuals (bottom left). and location of encounters

with bottlenose dolphins within the Moray Firth Special Area of Conservation from May to

September 1990 to 2020 (right).

During surveys, when we located a bottlenose dolphin group (one or more individuals in close

proximity and engaging in similar behaviours such as travelling and foraging) we collected photo-

identification data following a standardised protocol (35). We identified individuals from high-quality

photographs based on unique markings matched against a photo catalogue of previously identified

individuals from the area (32,33,36). On average, in a group 84% are successfully photographed, a

rate of identification well above the level at which social network metrics in incomplete networks

are reliable (37). All individual identifications from photographs are confirmed by at least two

experienced researchers. For individuals first sighted as calves we could determine their year of birth

and so their age (38), but for individuals first sighted as juveniles or adults their exact age is

unknown. Sex was determined using genital photographs or if an adult was seen in repeat

associations with a known calf (38).

Social network construction

Individuals sighted during the same encounter were assumed to be in the same group and therefore

associating (known as the 'Gambit of the Group'; ,39). Aggregating many of these records of groups

allows one to infer which individuals are frequently associated and which individuals infrequently or

never associate. We removed observations of individuals younger than three years old (n = 2634

observations of 233 individuals), as these individuals are not likely to be independent of the mother

and so their social associations most likely represent her preferences. We then converted the

records of encounters into group by individual matrices (indicating which individuals were seen

together in each encounter) and then into weighted, undirected social networks using the R package

*asnipe* (40). Edge weights were set as the simple ratio index, where the number of times two

individuals are seen together is divided by the total number of times they are seen, both together

and apart (41). This measure ranges from 0 (individuals never seen together) to 1 (individuals always

seen together). We did this separately for each year, creating a social network per year to assess

how social phenotypes vary at the yearly scale in response to environmental conditions. To assess

how social phenotypes vary at the monthly scale in response to environmental conditions we then

reconstructed social networks per month and removed any months with fewer than 10 encounters

(excluding 98 out of 217 months; as networks constructed using fewer than 10 observations can be

biased; Franks *et al.* 2010). Histograms of the frequency of the number of encounters per year and

per month are shown in Fig. S1.

For each individual present in the network for each year, we calculated the following: strength (the

sum of all an individual's associations, representing its gregariousness), weighted clustering

coefficient (the rate at which an individual associates with other individuals who also associate with

each other, representing how tightly individuals in its immediate social environment interact), and

closeness (the mean of the path lengths between that individual and each other individual in the

network, corrected for network size to allow comparisons among networks which vary in the

number of individuals, representing its connectedness to the wider population). We removed an

individual's scores for a given year if they had fewer than three observations in that year (removing

498 and leaving 1133), as the social network position of those individuals would be highly uncertain.

They would however still contribute to the social environments and therefore social network

measures of individuals in that year who had three or more observations. We repeated this at the

month level, again removing individuals' monthly scores when they had fewer than three

observations that month (removing 2215 and leaving 1704). Histograms of the frequency of the

number of encounters per individual per year and per month are shown in Fig. S2. We also removed

instances where an individual's closeness was over 1, which occurs when an individual is only

connected to a single other (n = 2 in the yearly networks, n = 18 in the monthly networks). The social

network measures were not well correlated; Pearson correlations between individuals' strength and

clustering coefficient were 0.136 (yearly) and 0.101 (monthly), for strength and closeness they were

-0.042 (yearly) and -0.314 (monthly), and for clustering coefficient and closeness they were 0.157

(yearly) and 0.032 (monthly).

Environmental data

Following Lusseau *et al*. (43), we used the North Atlantic Oscillation (NAO) index as a measure of

climate. We used monthly and yearly measures of the NAO index between 1990 and 2020

downloaded from https://www.cpc.ncep.noaa.gov/products/precip/CWlink/pna/nao.shtml (Fig. S3).

This index indicates the atmospheric pressure difference between the low pressure zone over

Iceland and the high pressure zone over the Azores (44,45). This index has frequently been linked to

the ecology of animal populations (46,47), for example influencing the foraging behaviour of Cory's

shearwaters (Calonectris borealis; ,48).  While climatic effects on cetaceans are typically thought to

occur via changes in prey species (29,49), a phenomenon also observed in our study population (43),

it is also possible that cetaceans respond directly to climate, sometimes at even faster rates than

their prey species (50–52).

For our index of food availability we again followed Lusseau *et al*. (43), using data on monthly

catches by fishing rods (as opposed to nets) from the wild of both one season and multiple season

adult Atlantic salmon (*Salmo salar*) from the Alness, Beauly, Canon, Ness, and Nairn rivers. These

feed into the sea where observations take place and hence are expected to be a good proxy for

salmon availability in that area. Further, catches on rods are positively correlated among rivers and

among months and with automatic counter data, indicating they are a good proxy for actual

abundance (53,54). Atlantic salmon are an important food source for this dolphin population (55,56),

with dolphins forming larger groups when salmon are more abundant (which is correlated with the

two-year lagged winter NAO index; ,43). We downloaded monthly data from

https://scotland.shinyapps.io/sg-salmon-sea-trout-catch/ (summing retained and released fish) and

summed monthly catches within a calendar year for yearly measures of fish abundance (Fig. S4).

Data analysis

All analyses were performed in R (ver 4.1.0; ,57) using linear mixed effect models in *glmmTMB* (58).

Using regression-based models as opposed to randomisation-based tests has been recommended

for analysing questions about node-level social network traits as it improves the ability to make

inference while accounting as well as node permutations for common types of data non-

independence (59). We fitted 12 models, with all combinations across the three social traits, the two

environmental variables, and the yearly and monthly timescales. Clustering coefficient cannot be

calculated when an individual only associates with one other individual, and closeness cannot be

calculated for individuals without any connections, and so the datasets for these models were

slightly smaller than the dataset for strength (see below). We included individuals with unknown

birth and death dates to maximise our sample size, and so neither age nor lifespan could be included

as predictor variables. In all models we included the fixed effect of sex (as males and females can

differ in social behaviour; ,60,61; ranging behaviour, and survival; ,62), and either the NAO index or

the count of caught salmon for that month or year. We did not include both the NAO index and

salmon abundance in the same model as we encountered estimation problems with two random

slopes. Including sex meant we excluded individuals of unknown sex (258 observations of 103

individuals for the yearly networks, 281 observations of 91 individuals for the monthly networks),

but the interactions between individuals of known and unknown sex were still used to build the

networks and so associations with individuals of unknown sex still influenced the social network

traits of males and females. We mean centred and scaled to unit variance the environmental

variable (63), and included the interaction between it and sex, to see if male and female social

behaviour responded differently to environmental variation. Random effects were the random

intercept for individual ID, the random slope of individual ID with the environmental variable fitted

as a fixed effect in the model (again mean centred and with a standard deviation of 1), and the

correlation between these two terms. We also included a temporal autocorrelation term (ar1)

among years to account for variation among years and that adjacent years may be more similar than

non-adjacent years. Similarly, in the models for monthly variation we included a random effect of

month. We used a Gaussian error structure for all models. For strength and closeness, we used log

link functions as the distributions were right skewed, adding 0.001 to all strength measures before

transformation to allow us to include zero values (although zeroes were rare; n = 1 for the yearly

analysis and n = 6 for the monthly analysis). For clustering coefficient, which is bounded between 0

and 1, we used a logit link function, which is preferable to an arcsine transformation when handling

response variables bounded in this way (64). We used the default optimising algorithm for all models

except the model for strength in response to monthly variation in salmon abundance, and the

models for clustering coefficient in response to monthly variation in both NAO index and salmon

abundance, where we used the "BFGS" optimiser, as otherwise the model did not converge (58).

We report the coefficients and standard errors for each fixed effect, along with p-values, to give an

idea of the magnitude and uncertainty of each effect. We used the p-values from the Anova function

of the *car* package (65), using a Chi-squared test with type III sum of squares. We describe these p-

values in terms of "clarity" rather than "significance", see Dushoff et al. (66) for a discussion on this.

To test whether individuals clearly differed in their response to the environmental variable, we first

tested whether there was a correlation between an individual's plasticity and its mean behaviour by

re-fitting each model (12 in total) with the correlation between random intercepts and slopes

supressed to zero and conducted a likelihood ratio test between the full model and this reduced

model with a single degree of freedom. If there was no clear difference between the models, we

then tested the importance of the random slopes by comparing the model with an intercept-slope

correlation of zero to a model without the random slopes (but still with the random intercepts) using

a likelihood ratio test with a mix of zero and one degrees of freedom (as is appropriate for testing

the clarity of single variance components, 67). When there is variation in plasticity, the magnitude of

among-individual differences varies across environments. We therefore calculate the marginal

repeatability for each trait following Schielzeth and Nakagawa (68). We did this for all trait-

environmental variable-time scale combinations (12 in total), even when there was no evidence for

random slopes, to aid comparison among traits.

## Results

### Change with environmental variables at a yearly scale

For the analysis of how dolphin social phenotypes change at the yearly scale in response to environmental variables, our dataset included 155 unique individuals. For strength there were 1133 measures, for clustering coefficient there were 1129 and for closeness there were 1132. Traits had a mean of 7.31 (sd = 5.99), 7.28 (sd = 5.92), and 7.30 (sd = 5.99) measures per individual for strength, clustering coefficient and closeness respectively

Dolphins' strength and clustering coefficient were not affected by the NAO index (Fig. 2a and b) or salmon abundance (Fig. 2d and e) in both sexes, and the sexes did not differ in mean strength or clustering coefficient (Table 1, full model results in Tables S1-6, supplementary materials). There were no intercept-slope correlations, and the random slopes were not statistically clear, suggesting that individual dolphins did not differ in their (lack of) response to the NAO index (Table 2). However, individuals did differ in their responses to salmon abundance, with clear random slopes for both strength and clustering coefficient. For both traits the intercept-slope correlation was not clearly different from zero, although it was negative and close to clarity for strength (suggesting individuals with lower mean strength increased their strength at higher values of the NAO index; Table 2). The marginal repeatabilities for strength were low: 0.017 in the NAO index model and 0.019 in the salmon abundance model, suggesting the trait has limited repeatability. The marginal repeatabilities for clustering coefficient were higher: 0.173 and 0.204 for the NAO index and salmon abundance models respectively.

Similarly, closeness did not vary with NAO index (Fig. 2c) or salmon numbers (Fig. 2f) in either sex (Table 1). Although females may have had higher mean closeness than males, the effect was not statistically clear (Supplementary Material Tables S5 & S6). Again, individuals' mean behaviour and plasticities in response to the NAO were not related, and dolphins did not show differences in their

plasticities (Table 2). However, dolphins did show a mean-plasticity relationship for salmon

abundance, with individuals with lower mean closeness increasing their closeness more with

increasing salmon abundance. This also indicates there was individual variation in plasticity (Table 2).

The marginal repeatabilities were low for closeness: 0.030 in the NAO index model and 0.023 in the

salmon abundance model.

**Figure 2.** Plots of each of the three social network traits and yearly variation in the North Atlantic

Oscillation (NAO) index (a. strength, b. clustering coefficient, c. closeness) and salmon abundances

(d. strength, e. clustering coefficient, f. closeness). For each individual dolphin we have predicted its

network trait on the observed scale based on the model results, using the "predict" function in R

with the individual's sex, the range of NAO values or salmon counts that individual was exposed to,

and setting the year to 2005.

For all social traits there was substantial among-year variation and consecutive years were positively

correlated (year to year correlations: strength; $r_{NAO}$ = 0.314, $r_{Salmon}$ = 0.429; clustering coefficient; $r_{NAO}$

= 0.559, $r_{Salmon}$ = 0.560; and closeness; $r_{NAO}$ =0.455, $r_{Salmon}$ = 0.543), showing that, as expected,

adjacent years were more similar than non-adjacent years.

In summary, there were no trait-environment association at the yearly scale. However, individuals

always differed in their response to salmon abundance but never in their (lack of) response to the

NAO index. Clustering coefficient showed some repeatability across environments, but strength and

closeness did not.

**Table 1**. Main effects ("M") and interactions with sex ("I", also shaded in grey) for the two

environmental variables effects on the three social network traits, at each of the monthly and yearly

scales. Effects clearly different from zero (p < 0.05) are highlighted in bold, while

effects with p values between 0.1 and 0.05 are highlight in italics.

|  |  |  | Strength | Clustering coefficient | Closeness |
|---|---|---|---|---|---|
| NAO index | Yearly | M | $\beta$ = 0.013 ± 0.075, $\chi^2$ = 0.032, p = 0.858 | $\beta$ = 0.012 ± 0.086, $\chi^2$ = 0.019, p = 0.891 | $\beta$ = -0.053 ± 0.093, $\chi^2$ = 0.324, p = 0.569 |
|  |  | I | $\beta$ = 0.009 ± 0.023, $\chi^2$ = 0.163, p = 0.687 | $\beta$ = 0.005 ± 0.034, $\chi^2$ = 0.019, p = 0.889 | $\beta$ = -0.011 ± 0.013, $\chi^2$ = 0.733, p = 0.392 |
|  | Monthly | M | $\beta$ = 0.029 ± 0.021, $\chi^2$ = 1.902, p = 0.168 | $\beta$ = 0.038 ± 0.038, $\chi^2$ = 0.968, p = 0.325 | *$\beta$ = -0.077 ± 0.044, $\chi^2$ = 3.004, p = 0.083* |
|  |  | I | $\beta$ = -0.002 ± 0.024, $\chi^2$ = 0.009, p = 0.924 | $\beta$ = 0.009 ± 0.042, $\chi^2$ = 0.045, p = 0.832 | *$\beta$ = 0.103 ± 0.053, $\chi^2$ = 3.724, p = 0.054* |
| Salmon | Yearly | M | $\beta$ = 0.123 ± 0.077, $\chi^2$ = 2.523, p = 0.112 | $\beta$ = 0.040 ± 0.088, $\chi^2$ = 0.019, p = 0.891 | $\beta$ = -0.142 ± 0.088, $\chi^2$ = 2.609, p = 0.106 |
|  |  | I | $\beta$ = -0.009 ± 0.031, $\chi^2$ = 0.091, p = 0.762 | $\beta$ = 0.012 ± 0.043, $\chi^2$ = 0.019, p = 0.890 | $\beta$ = -0.011 ± 0.013, $\chi^2$ = 0.733, p = 0.392 |
|  | Monthly | M | $\beta$ = 0.015 ± 0.034, $\chi^2$ = 0.199, p = 0.655 | **$\beta$ = 0.170 ± 0.051, $\chi^2$ = 11.290, p = 0.001** | $\beta$ = 0.019 ± 0.064, $\chi^2$ = 0.086, p = 0.769 |
|  |  | I | $\beta$ = 0.050 ± 0.035, $\chi^2$ = 2.131, p = 0.144 | $\beta$ = -0.060 ± 0.049, $\chi^2$ = 1.495, p = 0.221 | $\beta$ = 0.006 ± 0.069, $\chi^2$ = 0.006, p = 0.936 |

272   Change with environmental variables at a monthly scale

273   We analysed how dolphin social phenotypes change in response to environmental variables at the

274   monthly scale with a dataset of 151 unique individuals for strength and closeness and 149 for

clustering coefficient. For strength there were 1704 measures, for clustering coefficient 1688, and

for closeness 1696. Traits had a mean of 11.3, 11.3, and 11.2 measures per individual for strength,

clustering coefficient, and closeness respectively (sd = 10.7 for all).

**Table 2**. Intercept slope correlations ("ISC") and their statistical tests, and whether random slopes

("RS") were present or absent and their statistical tests (likelihood ratio tests in all cases, an asterix is

given for the test of the random slopes if it was not performed as the intercept-slope correlation was

first found to be clear). Clear positive correlations are noted in bold text and highlighted with blue

fill, and clear negative correlations with bold text and orange fill, with the two intercept-slope

correlations with p values between 0.1 and 0.05 indicted in italics.

| | | | Strength | Clustering coefficient | Closeness 284 |
|---|---|---|---|---|---|
| NAO index | Yearly | ISC | -0.539 ($\chi_1^2$ = 1.299, p = 0.254) | -0.999 ($\chi_1^2$ = 0.896, p = 0.344) | -0.784 ($\chi_1^2$ = 1.477, p = 0.224) |
| | | RS | Absent ($\chi_{0,1}^2$ = 0.236, p = 0.314) | Absent ($\chi_{0,1}^2$ = 0.000, p = 0.500) | Absent ($\chi_{0,1}^2$ = 0.013, p = 0.455) |
| | Monthly | ISC | **-0.963 ($\chi_1^2$ = 6.864, p = 0.009)** | 0.975 ($\chi_1^2$ = 1.660, p = 0.198) | **-0.792 ($\chi_1^2$ = 22.911, p < 0.001)** |
| | | RS | **Present \*** | Absent ($\chi_{0,1}^2$ = -0.027, p = 0.500) | **Present \*** |
| Salmon | Yearly | ISC | *-0.321 ($\chi_1^2$ = 3.002, p = 0.083)* | 0.268 ($\chi_1^2$ = 1.182, p = 0.277) | **-0.510 ($\chi_1^2$ = 9.389, p = 0.002)** |
| | | RS | **Present ($\chi_{0,1}^2$ = 15.925, p < 0.001)** | **Present ($\chi_{0,1}^2$ = 7.385, p = 0.003)** | **Present \*** |
| | Monthly | ISC | *-0.299 ($\chi_1^2$ = 2.838, p = 0.092)* | 0.994 ($\chi_1^2$ = 1.946, p = 0.163) | **-0.350 ($\chi_1^2$ = 5.891, p = 0.015)** |
| | | RS | **Present ($\chi_{0,1}^2$ = 28.258, p < 0.001)** | Absent ($\chi_{0,1}^2$ = 0.000, p = 0.500) | **Present \*** |

Strength was not affected by the monthly NOA index (Fig. 3a) or monthly salmon abundance (Fig.

3d) in either sex and the sexes did not differ in their mean strength (Table 1, full model results in

Tables S7-12 in the supplementary materials). Individuals differed in their responses to the monthly

NAO index and salmon abundances (Table 2). For the NAO index, individuals with high values

decreased their strength as the NAO index increased while individuals with low values increased

their strength (Table 2). There was no clear intercept-slope correlation for salmon abundance

although it was also negative. As for the yearly models, the marginal repeatability of strength was

low (0.007 in the NAO index model, 0.009 in the salmon abundance model).

Months of higher salmon abundance led to higher clustering coefficients in both sexes (Figs. 3e and

4 and Table 1). Clustering coefficient did not however vary in response to monthly NOA index in

either sex (Fig. 3b and Table 1). The sexes did not differ in their mean clustering coefficient.

Individuals showed no mean-plasticity relationship and did not vary in their responses to salmon

abundance, or in their lack of response to the NAO index (Table 2). Clustering coefficient was

moderately repeatable, with marginal repeatabilities of 0.337 and 0.284 in the NAO index and

salmon abundance models respectively.

**Figure 3**. Plots of each of the three social network traits and monthly variation in the North Atlantic

Oscillation (NAO) index (a. strength, b. clustering coefficient, c. closeness) and salmon abundances

(d. strength, e. clustering coefficient, f. closeness). In c., males are shown in purple, females in blue,

to visualise the possible difference between the sexes (see Results). For each individual dolphin we

have predicted its network trait on the observed scale based on the model results, using the

"predict" function in R with the individual's sex, the range of NAO values or salmon counts that

individual was exposed to, setting the year to 2005, and the month to June.

Females had a lower average closeness than males, although this effect was only clear in the model

for the NAO index (Table S11). Further, the sexes may have differed in their response to the NAO

index; while males showed a marginally unclear decrease with higher NAO index, females may have

shown no change or even a slight increase (Fig. 3c and Table 1). Closeness did not respond to

monthly salmon numbers in either sex (Fig. 3f and Table 1). Individuals differed in their responses to

the NAO index and to salmon abundance (Table 2), with individuals with high values decreasing their

closeness with higher values of the environmental factor and individuals with low values increasing

their closeness. Unlike for the yearly scale there was some repeatability of closeness at the monthly

scale, with a marginal repeatability of 0.216 in the NAO index model and 0.168 in the salmon

abundance model.

**Figure 4**. Plots of dolphin social networks in a month of low salmon abundance (May 1990) and a

month of high salmon abundance (August 2007). Circles are individual dolphins and grey lines

indicate associations i.e., those seen in the same group at least once in that month.

There was among-month and among-year variation, and consecutive years were always positive

correlated (year to year correlations: strength: $r_{NAO}$ = 0.553, $r_{Salmon}$ = 0.529; clustering coefficient: $r_{NAO}$

= 0.228, $r_{Salmon}$ =0.074; closeness: $r_{NAO}$ = 0.467, $r_{Salmon}$ = 0.475).

In summary, months with higher salmon numbers led to higher clustering coefficient in both sexes

but did not impact strength or closeness. Meanwhile, months with a higher NAO index were

associated with a higher closeness in females and a lower closeness in males, although both these

effects were not clearly different from zero. The NAO index did not affect strength or clustering

coefficient. Individuals differed in how their strength and closeness changed in response to both the

NAO index and salmon abundance, but they did not differ in how their clustering coefficient changed

for either variable. Clustering coefficient was somewhat repeatable across environments, closeness

marginally so, and strength not at all.

Discussion

We explored whether bottlenose dolphin social behaviour responded to environmental variation.

Typically, social behaviour showed no variation at the yearly scale in response to the NAO index or

salmon abundance. Similarly, two social behaviours showed no variation to environmental change at

the monthly scale, although, how tightly individuals in the immediate social environment interacted

(clustering coefficient) did increase in response to increases in monthly salmon abundance. We also

showed that individuals differed in their behavioural responses to changes in salmon abundance at

both yearly and monthly scales, and to the NAO index at the monthly scale, indicating variation in

plasticity. This plasticity was often, but not always, negatively associated with mean behaviour,

causing among individual differences to typically be greater for low values of salmon abundance and

the NAO index. Interestingly, when males and females increased their clustering coefficient in

response to higher monthly salmon abundances, there was limited among-individual variation in this

change. This conformity of response perhaps suggests a particularly important or unavoidable

change. Among-individual variation in mean behaviour was limited for strength at both timescales

and for closeness at the yearly timescale, marginal for closeness at the monthly scale and clustering

coefficient at the yearly scale, and most apparent for clustering coefficient at the monthly scale (see

also 69 who found consistency over lifetime in the social behaviours of Indo-Pacific bottlenose

dolphins, T. aduncus). Social traits differed therefore in either showing limited population-level

plasticity and little repeatability, but some variation in individual slopes (strength and closeness),

and some population-level plasticity and repeatability, and less variation among individual slopes

(clustering coefficient).

Despite the population-level stability in strength and closeness, in most cases individual dolphins

differed in how their social behaviour changed in response to environmental variation (i.e., they

showed among-individual variation in plasticity). Therefore, while the population as a whole showed

no overall change, some individuals would show an increase in their overall gregariousness and/or

connectedness to wider parts of the network, while others a decrease. Variation in individual

plasticity leads to environment-dependent repeatability (and possibly heritability), can dampen

population responses to environmental variability, and enhance population persistence (17). For

example, due to the negative intercept-slope correlation, individual closeness shows the most

among-individual variation at low monthly salmon abundances (the approximate marginal

repeatability two standard deviations below the mean salmon abundance was 0.54, compared to

0.07 at the mean). This wider range of social phenotypes could increase the possibility that at least

some individuals are successful despite low food availability. Determining the genetic basis and

importance of early life conditions for the development of these different responses (11) and how

this variation impacts population dynamics (70) is key. Models projecting how dolphin populations

will change in the future should account for both among-individual variation in mean behaviour and

in behavioural plasticity.

In general, the lack of plasticity in social network traits suggests dolphin social behaviour will show

limited changes at the population level if salmon abundances or climate change over time (based on

similar changes to environmental variables studied here). Populations of animals with relatively long

generation times are more reliant on phenotypic plasticity than adaptive evolution to track rapid

environmental change (22). Therefore, if dolphins do not change plastically there is limited scope for

evolution to compensate to enhance phenotype-environment fit. If social behaviours are important

for coping with detrimental environments, this could potentially be problematic, leading to reduced

reproduction and possibly population decline.

The one exception to stability was that individual clustering coefficient increased in months with

higher salmon abundance (see Fig. 4 for a comparison of months with low vs. high salmon

abundance). This indicates that dolphins were "closing triads" and interacting more often with those

who they shared a mutual third association. The increase in clustering coefficient occurred without a

change in strength or closeness, suggesting a nuanced change in behaviour. Interestingly, individuals

did not show variability in this increase, suggesting it is important or unavoidable for all. Similar

results have been found in spotted hyenas (*Crocuta crocuta*) where seasons with high prey

availability have denser social networks (71). However, other systems have found lower food

availability leads to decreased sociability (72,73), and it is presumed this response reduces the

intensity of resource competition (reviewed in: ,6,see also: ,74). Therefore, higher clustering may be

beneficial for non-foraging reasons, such as mating. Both mating and calving seasons approximately

align with the months of highest salmon abundance in the summer (Fig. S4), supporting this

suggestion. Dolphins could also reduce travel distances when food availability is high (60), leading to

them stay in the same place and so interact more with a tight circle of others; increasing their

clustering coefficient. Alternatively, it is possible that months with fewer salmon also differ in an

unidentified variable which causes dolphins to cluster less. However, it is unlikely that this

unidentified factor is predation threat that changes month to month, as predators are absent in this

area (75). Therefore, a change in social behaviour and/or movement related to the seasonal

availability of salmon, perhaps influenced by reproductive behaviour, seems the most likely.

Notwithstanding the response of dolphins to changes in environmental factors on a monthly scale,

we did not see any responses to environmental variation on a yearly scale. Both factors varied

considerably year-to-year; the NAO index ranged from -2.35 to 1.23 (Fig. S3) and salmon abundance

varied by more than 2.8 times between years (Fig. S4), hence a lack of relevant variability seems

unlikely. Lusseau *et al*. (43) observed larger groups in years of higher salmon abundance, but this

was over fewer years and was not an individual-level analysis. The robustness or inflexibility of social

behaviour in response to year-to-year variation in climatic conditions and food availability might

indicate that the yearly environmental variation is inconsequential and so they have no need to

respond to it. However, it seems unlikely that variation in food availability of 2.8 times does not

matter to this population, as their distribution in space reflects foraging opportunities (76).

Alternatively, dolphins may change other phenotypes, such as foraging behaviour or metabolism, to

cope with these stressors (29), leaving social behaviour unchanged. For example, salmon are not this

dolphin population's only prey (56), and so they may switch to other prey if salmon are scarce,

obviating the need to modify their social networks.

We estimated correlations between an individual's mean behaviour (its intercept) and its plasticity in

behaviour (its slope). For strength and closeness, intercept-slope correlations were negative if they

were different from zero. Therefore, individuals typically converged to similar behaviour at higher

NAO index values and salmon abundances (among-individual differences in mean behaviours were

also low for these traits). Negative intercept-slope correlations can constrain evolution if higher

mean values *and* higher plasticities are favoured, as selection is favouring a combination of

phenotypes that is rare (17). In contrast, intercept-slope correlations for clustering coefficient were

either zero or, in the case of the response to salmon abundance at the yearly scale, positive. The

positive correlation shows behaviours diverging among individuals at higher yearly salmon

abundances. Additionally, while a positive intercept slope correlation means that selection on either

intercept or slope causes the other to increase too, no correlation means intercepts and slopes can

be independent targets of selection. Kebke *et al*. (29) suggest that cetacean ranging and foraging

behaviour may well be under selection for increased plasticity as environments change, and so

testing this directly is a logical and important next step.

In conclusion, we found that at the population-level dolphin social behaviour is relatively stable

across years despite year-to-year variation in environmental conditions. At the monthly scale, we

observed all individuals increased clustering of local social associations due to higher salmon

abundance. However, at both monthly and yearly scales there was lots of variability at the individual

level in responses to environmental factors, and variation among both traits and timescales in how

repeatable traits were. As such, individual heterogeneity in both mean and plasticity in behaviour

both should be more widely accounted for when predicting species responses to environmental

change, and individual plasticities as well as means may be targets of selection and hence evolvable.

## Acknowledgements

We are indebted to Paul Thompson for conception and development of this long-term individual

based research study, and for advice during this work. We are especially grateful to our colleagues at

the Lighthouse Field Station, past and present, who have contributed to and advanced this long-term

study, and to Charlie Phillips who contributed land-based photographs for photo-identification.

Matthew Silk, Dan Blumstein, and two anonymous reviewers made numerous useful comments and

suggestions on an earlier draft. This long-term study has depended upon a number of funders

including University of Aberdeen, NatureScot, Beatrice Offshore Windfarm Ltd., Moray Offshore

Renewables Ltd., Marine Scotland, the Crown Estate, Highlands and Islands Enterprise, the BES,

ASAB, Greenpeace Environmental Trust, Whale and Dolphin Conservation, Talisman Energy (UK) Ltd.,

Department of Energy and Climate Change, Chevron and Natural Environment Research. Survey

work was conducted under NatureScot Animal Scientific Licences. We have no conflicts of interest.

## Statement of authorship

DF conceived the idea for the manuscript, BC provided data, DF ran analyses with guidance from BC,

DF wrote first draft of the manuscript and both authors contributed to revisions.

## Data accessibility statement

Data and R code used for the analysis are available to reviewers and will be made publicly available

upon acceptance.

## References

1.  Snyder-Mackler N, Burger JR, Gaydosh L, Belsky DW, Noppert GA, Campos FA, et al. Social
    determinants of health and survival in humans and other animals. Science [Internet]. 2020 May
    22 [cited 2020 Oct 8];368(6493). Available from:
    https://science.sciencemag.org/content/368/6493/eaax9553

2.  Moore AJ, Brodie EDI, Wolf JB. Interacting phenotypes and the evolutionary process: I. Direct
    and indirect genetic effects of social interactions. Evolution. 1997;51(5):1352–62.

3.  Wolf JB, Brodie III ED, Moore AJ. Interacting Phenotypes and the Evolutionary Process. II.
    Selection Resulting from Social Interactions. The American Naturalist. 1999 Mar;153(3):254–66.

4.  McGlothlin JW, Moore AJ, Wolf JB, Brodie ED. Interacting phenotypes and the evolutionary
    process. III. Social evolution. Evolution; international journal of organic evolution. 2010
    Sep;64(9):2558–74.

5.  Komdeur J, Ma L. Keeping up with environmental change: The importance of sociality. Ethology.
    2021 Oct 1;127(10):790–807.

6.  Fisher DN, Kilgour RJ, Siracusa ER, Foote JR, Hobson EA, Montiglio PO, et al. Anticipated effects
    of abiotic environmental change on intraspecific social interactions. Biological Reviews.
    2021;96(6):2661–93.

7.  White DJ, Gersick AS, Snyder-Mackler N. Social networks and the development of social skills in
    cowbirds. Philosophical Transactions of the Royal Society of London B: Biological Sciences.
    2012;367(1597).

8.  Abrahms B, Rafiq K, Jordan NR, McNutt JW. Long-term, climate-driven phenological shift in a
    tropical large carnivore. Proceedings of the National Academy of Sciences [Internet]. 2022 Jul 5
    [cited 2022 Jun 29];119(27). Available from:
    https://pnas.org/doi/full/10.1073/pnas.2121667119

9.    Nussey DH, Wilson AJ, Brommer JE. The evolutionary ecology of individual phenotypic plasticity
in wild populations. Journal of Evolutionary Biology. 2007 May;20(3):831–44.

10.   Dingemanse NJ, Kazem AJN, Réale D, Wright J. Behavioural reaction norms: animal personality
meets individual plasticity. Trends in Ecology & Evolution. 2010 Feb;25(September):81–9.

11.   Stamps JA. Individual differences in behavioural plasticities. Biological Reviews. 2016
May;91(2):534–67.

12.   Stamps J, Biro P. Personality and individual differences in plasticity. Current Opinion in
Behavioral Sciences [Internet]. 2016 [cited 2016 Nov 22]; Available from:
http://www.sciencedirect.com/science/article/pii/S2352154616301589

13.   Fisher DN, David M, Tregenza T, Rodriguez-Munoz R. Dynamics of among-individual behavioral
variation over adult lifespan in a wild insect. Behavioral Ecology. 2015;26(4):975–85.

14.   Brommer JE. Variation in plasticity of personality traits implies that the ranking of personality
measures changes between environmental contexts: calculating the cross-environmental
correlation. Behavioral Ecology and Sociobiology. 2013 Aug 7;67(10):1709–18.

15.   Vindenes Y, Langangen Ø. Individual heterogeneity in life histories and eco-evolutionary
dynamics. Ecology Letters. 2015;18(5):417–32.

16.   Hamel S, Gaillard JM, Yoccoz NG. Introduction to: Individual heterogeneity - the causes and
consequences of a fundamental biological process. Oikos. 2018 May;127(5):643–7.

17.   Dingemanse NJ, Wolf M. Between-individual differences in behavioural plasticity within
populations: causes and consequences. Animal Behaviour. 2013 May;85(5):1031–9.

18.   Chevin LM. Genetic constraints on adaptation to a changing environment. Evolution. 2013
Mar;67(3):708–21.

19.   Araya-Ajoy YG, Dingemanse NJ. Repeatability, heritability, and age-dependence of seasonal
plasticity in aggressiveness in a wild passerine bird. van de Pol M, editor. Journal of Animal
Ecology. 2017 Mar;86(2):227–38.

20.   Nussey DH, Postma E, Gienapp P, Visser ME. Selection on Heritable Phenotypic Plasticity in a
Wild Bird Population. Science. 2005;310:304–6.

21.   Brommer JE. Phenotypic plasticity of labile traits in the wild. Current Zoology. 2013;59(4):485–
505.

22.   Hoffmann AA, Sgrò CM. Climate change and evolutionary adaptation. Nature. 2011 Feb
24;470(7335):479–85.

23.   Simmonds MP, Isaac SJ. The impacts of climate change on marine mammals: early signs of
significant problems. Oryx. 2007 Jan;41(1):19–26.

24.   Tyack PL. Implications for marine mammals of large-scale changes in the marine acoustic
environment. Journal of Mammalogy. 2008 Jun;89(3):549–58.

25. MacLeod CD. Global climate change, range changes and potential implications for the
conservation of marine cetaceans: a review and synthesis. Endangered Species Research. 2009
May 14;7(2):125–36.

26. Evans PG, Bjørge A. Impacts of climate change on marine mammals. Marine Climate Change
Impacts Partnership (MCCIP) Science Review. 2013;2013:134–48.

27. Avila IC, Kaschner K, Dormann CF. Current global risks to marine mammals: Taking stock of the
threats. Biological Conservation. 2018 May 1;221:44–58.

28. van Weelden C, Towers JR, Bosker T. Impacts of climate change on cetacean distribution, habitat
and migration. Climate Change Ecology. 2021 Jul 1;1:100009.

29. Kebke A, Samarra F, Derous D. Climate change and cetacean health: impacts and future
directions. Philosophical Transactions of the Royal Society B: Biological Sciences [Internet]. 2022
[cited 2022 Nov 10];377. Available from:
https://royalsocietypublishing.org/doi/abs/10.1098/rstb.2021.0249?casa_token=mPcr4-
VPLiAAAAAA:NkbeNvtFMfHsOUzCtFl45ae_Cvvkd5WaxEEDNRIlVgvmhX1f-fjUhSHayK2aBuP-
qrmRlTa3pJow

30. Kershaw JL, Hall AJ. Mercury in cetaceans: Exposure, bioaccumulation and toxicity. Science of
The Total Environment. 2019 Dec 1;694:133683.

31. Arso Civil M, Quick NJ, Mews S, Hague E, Cheney BJ, Thompson PM, et al. Improving
understanding of bottlenose dolhin movements along the east coast of Scotland. 2021 Mar.
(European Offshore Wind Deployment Centre (EOWDV)). Report No.: SMRUC-VAT-2020-10.

32. Wilson B, Hammond PS, Thompson PM. Estimating Size and Assessing Trends in a Coastal
Bottlenose Dolphin Population. Ecological Applications. 1999;9(1):288–300.

33. Cheney B, Thompson PM, Ingram SN, Hammond PS, Stevick PT, Durban JW, et al. Integrating
multiple data sources to assess the distribution and abundance of bottlenose dolphins Tursiops
truncatus in Scottish waters. Mammal Review. 2013;43(1):71–88.

34. Arso Civil M, Quick NJ, Cheney B, Pirotta E, Thompson PM, Hammond PS. Changing distribution
of the east coast of Scotland bottlenose dolphin population and the challenges of area-based
management. Aquatic Conservation: Marine and Freshwater Ecosystems. 2019;29(S1):178–96.

35. Cheney B, Corkrey R, Durban JW, Grellier K, Hammond PS, Islas-Villanueva V, et al. Long-term
trends in the use of a protected area by small cetaceans in relation to changes in population
status. Global Ecology and Conservation. 2014 Dec 1;2:118–28.

36. Würsig B, Jefferson TA. Methods of photo-identification for small cetaceans [Internet]. 1990
[cited 2022 Aug 22] p. 43–52. (Reports of the International Whaling Commission). Report No.:
SC/A88/ID13. Available from:
https://www.researchgate.net/publication/285799303_Methods_of_photo-
identification_for_small_cetaceans

37. Silk MJ, Jackson AL, Croft DP, Colhoun K, Bearhop S. The consequences of unidentifiable
individuals for the analysis of an animal social network. Animal Behaviour. 2015 Jun;104:1–11.

38. Cheney BJ, Dale J, Thompson PM, Quick NJ, Barbara Cheney CJ. Spy in the sky: a method to
identify pregnant small cetaceans. Scales K, Bouchet P, editors. Remote Sensing in Ecology and

Conservation [Internet]. 2022 Feb 8 [cited 2022 Feb 14]; Available from:
https://onlinelibrary.wiley.com/doi/full/10.1002/rse2.258

39. Whitehead H, Dufault S. Techniques for Analyzing Vertebrate Social Structure Using Identified
Individuals: Review and Recommendations. Slater P, Rosenblatt J, Roper, editors. Advances in
the Study of Behavior [Internet]. 1999 [cited 2013 Aug 20];28. Available from:
http://www.sciencedirect.com/science/article/pii/S0065345408602156

40. Farine DR. Animal social network inference and permutations for ecologists in R using asnipe.
O'Hara RB, editor. Methods in Ecology and Evolution. 2013 Dec 22;4(12):1187–94.

41. Cairns SJ, Schwager SJ. A comparison of association indices. Animal Behaviour. 1987;35(5):1454–
69.

42. Franks DW, Ruxton GD, James R. Sampling animal association networks with the gambit of the
group. Behavioral Ecology and Sociobiology. 2010 Oct 17;64(3):493–503.

43. Lusseau D, Williams R, Wilson B, Grellier K, Barton TR, Hammond PS, et al. Parallel influence of
climate on the behaviour of Pacific killer whales and Atlantic bottlenose dolphins. Ecology
Letters. 2004 Sep 22;7(11):1068–76.

44. Hurrell JW. Decadal Trends in the North Atlantic Oscillation: Regional Temperatures and
Precipitation. Science. 1995 Aug 4;269(5224):676–9.

45. Stenseth NChr, Ottersen G, Hurrell JW, Mysterud A, Lima M, Chan K, et al. Review article.
Studying climate effects on ecology through the use of climate indices: the North Atlantic
Oscillation, El Niño Southern Oscillation and beyond. Proceedings of the Royal Society of London
Series B: Biological Sciences. 2003 Oct 22;270(1529):2087–96.

46. Gordo O, Barriocanal C, Robson D. Ecological Impacts of the North Atlantic Oscillation (NAO) in
Mediterranean Ecosystems. In: Vicente-Serrano SM, Trigo RM, editors. Hydrological,
Socioeconomic and Ecological Impacts of the North Atlantic Oscillation in the Mediterranean
Region [Internet]. Dordrecht: Springer Netherlands; 2011 [cited 2022 Aug 22]. p. 153–70.
(Advances in Global Change Research). Available from: https://doi.org/10.1007/978-94-007-
1372-7_11

47. Wan X, Holyoak M, Yan C, Le Maho Y, Dirzo R, Krebs CJ, et al. Broad-scale climate variation
drives the dynamics of animal populations: a global multi-taxa analysis. Biological Reviews
[Internet]. 2022 [cited 2022 Aug 22];n/a(n/a). Available from:
https://onlinelibrary.wiley.com/doi/abs/10.1111/brv.12888

48. Pereira JM, Paiva VH, Ceia FR, Ramos JA. Facing extremes: Cory's shearwaters adjust their
foraging behaviour differently in response to contrasting phases of North Atlantic Oscillation.
Reg Environ Change. 2020 Jun 24;20(3):77.

49. Víkingsson GA, Pike DG, Valdimarsson H, Schleimer A, Gunnlaugsson T, Silva T, et al. Distribution,
abundance, and feeding ecology of baleen whales in Icelandic waters: have recent
environmental changes had an effect? Frontiers in Ecology and Evolution [Internet]. 2015 [cited
2022 Nov 10];3. Available from: https://www.frontiersin.org/articles/10.3389/fevo.2015.00006

50. Sprogis KR, Christiansen F, Wandres M, Bejder L. El Niño Southern Oscillation influences the
abundance and movements of a marine top predator in coastal waters. Global Change Biology.
2018;24(3):1085–96.

51. Thorne LH, Nye JA. Trait-mediated shifts and climate velocity decouple an endothermic marine
predator and its ectothermic prey. Sci Rep. 2021 Sep 16;11(1):18507.

52. Methion S, Paradell OG, Padín XA, Corrège T, López BD. Group size varies with climate and
oceanographic conditions in bottlenose dolphins [Internet]. In Review; 2022 Apr [cited 2022 Nov
10]. Available from: https://www.researchsquare.com/article/rs-1547502/v1

53. Youngson AF, MacLean JC, Fryer RJ. Rod catch trends for early-running MSW salmon in Scottish
rivers (1952–1997): divergence among stock components. ICES Journal of Marine Science. 2002
Jan 1;59(4):836–49.

54. Thorley JL, Eatherley DMR, Stephen AB, Simpson I, MacLean JC, Youngson AF. Congruence
between automatic fish counter data and rod catches of Atlantic salmon (Salmo salar) in Scottish
rivers. ICES Journal of Marine Science. 2005 Jan 1;62(4):808–17.

55. Janik VM. Food–related bray calls in wild bottlenose dolphins (Tursiops truncatus). Proceedings
of the Royal Society of London Series B: Biological Sciences. 2000 May 7;267(1446):923–7.

56. Santos MB, Pierce GJ, Reid RJ, Patterson I a. P, Ross HM, Mente E. Stomach contents of
bottlenose dolphins (Tursiops truncatus) in Scottish waters. Journal of the Marine Biological
Association of the United Kingdom. 2001 Oct;81(5):873–8.

57. R Development Core Team, Team RDC. R: A language and environment for statistical computing
[Internet]. R Foundation for Statistical Computing. Vienna, Austria: R Foundation for Statistical
Computing; 2016 [cited 2013 Sep 17]. p. 409. (R Foundation for Statistical Computing; vol. 1).
Available from: http://www.r-project.org

58. Brooks ME, Kristensen K, van Benthem KJ, Magnusson A, Berg CW, Nielsen A, et al. glmmTMB
Balances speed and flexibility among packages for modeling. The R Journal. 2017;9(2):378–400.

59. Hart JDA, Weiss MN, Brent LJN, Franks DW. Common Permutation Methods in Animal Social
Network Analysis Do Not Control for Non-independence. bioRxiv [Internet]. 2021; Available
from: https://www.biorxiv.org/content/10.1101/2021.06.04.447124v1.abstract

60. Shane SH, Wells RS, Würsig B. Ecology, Behavior and Social Organization of the Bottlenose
Dolphin: A Review. Marine Mammal Science. 1986;2(1):34–63.

61. Mann J, Stanton MA, Patterson EM, Bienenstock EJ, Singh LO. Social networks reveal cultural
behaviour in tool-using dolphins. Nat Commun. 2012 Jul 31;3(1):980.

62. Arso Civil M, Cheney B, Quick NJ, Islas-Villanueva V, Graves JA, Janik VM, et al. Variations in age-
and sex-specific survival rates help explain population trend in a discrete marine mammal
population. Ecology and Evolution. 2019;9(1):533–44.

63. Schielzeth H. Simple means to improve the interpretability of regression coefficients. Methods in
Ecology and Evolution. 2010 Feb 10;1(2):103–13.

64. Warton D, Hui F. The arcsine is asinine: the analysis of proportions in ecology. Ecology [Internet].
2011 Apr 7 [cited 2014 Jul 16]; Available from: http://www.esajournals.org/doi/abs/10.1890/10-
0340.1

65. Fox J, Weisberg S. An R Companion to Applied Regression. Thousand Oaks (CA): Sage; 2019.

66. Dushoff J, Kain MP, Bolker BM. I can see clearly now: Reinterpreting statistical significance.
O'Hara RB, editor. Methods in Ecology and Evolution. 2019 Jun 27;10(6):756–9.

67. Self SG, Liang KY. Asymptotic properties of maximum likelihood estimators and likelihood ratio
tests under nonstandard conditions. Journal of the American Statistical Association. 1987
Jun;82(398):605–10.

68. Schielzeth H, Nakagawa S. Conditional repeatability and the variance explained by reaction norm
variation in random slope models. Methods in Ecology and Evolution. 2022;13(6):1214–23.

69. Evans T, Krzyszczyk E, Frère C, Mann J. Lifetime stability of social traits in bottlenose dolphins.
Commun Biol. 2021 Jun 18;4(1):1–8.

70. Vindenes Y, Engen S, Sæther B. Individual Heterogeneity in Vital Parameters and Demographic
Stochasticity. The American Naturalist. 2008 Apr;171(4):455–67.

71. Holekamp KE, Smith JE, Strelioff CC, Van Horn RC, Watts HE. Society, demography and genetic
structure in the spotted hyena. Molecular Ecology. 2012 Feb;21(3):613–32.

72. Randall JA, Rogovin K, Parker PG, Eimes JA. Flexible social structure of a desert rodent,
Rhombomys opimus: philopatry, kinship, and ecological constraints. Behavioral Ecology. 2005
Nov 1;16(6):961–73.

73. Williams AE, Worsley-Tonks KEL, Ezenwa VO. Drivers and consequences of variation in individual
social connectivity. Animal Behaviour. 2017 Nov 1;133:1–9.

74. Lin YH, Chan SF, Rubenstein DR, Liu M, Shen SF. Resolving the paradox of environmental quality
and sociality: The ecological causes and consequences of cooperative breeding in two lineages
of birds. American Naturalist. 2019 Aug 1;194(2):207–16.

75. Cheney BJ, Thompson PM, Cordes LS. Increasing trends in fecundity and calf survival of
bottlenose dolphins in a marine protected area. Scientific Reports. 2019 Dec 11;9(1):1767.

76. Hastie GD, Wilson B, Wilson LJ, Parsons KM, Thompson PM. Functional mechanisms underlying
cetacean distribution patterns: hotspots for bottlenose dolphins are linked to foraging. Marine
Biology. 2004 Feb 1;144(2):397–403.