Article Type: Original Article

Running Title: Chronosequence of invasion

Chronosequence of invasion reveals minimal losses of population genomic diversity,

niche expansion, and trait divergence in the polyploid, leafy spurge

Thomas A. Lake[1], Ryan D. Briscoe Runquist[1], Lex E. Flagel[12], and David A. Moeller[1]

Affiliations

[1] Department of Plant and Microbial Biology, University of Minnesota, St. Paul, Minnesota, USA

[2] Gencove, Long Island City, New York, USA

Emails and ORCID

Thomas Lake: lakex055@umn.edu, 0000-0002-8836-5164

Ryan Briscoe Runquist: rbriscoe@umn.edu, 0000-0001-7160-9110

Lex Flagel: flag0010@umn.edu, 0000-0003-2106-4196

David Moeller: moeller@umn.edu, 0000-0002-6202-9912

Correspondence

Thomas A. Lake and David A. Moeller

Department of Plant and Microbial Biology, University of Minnesota, St. Paul, Minnesota, USA

Email: lakex055@umn.edu, moeller@umn.edu

Acknowledgments

Funding was provided by the Minnesota Invasive Terrestrial Plants and Pests Center through

the Environment and Natural Resources Trust Fund as recommended by the Legislative-Citizen

Commission on Minnesota Resources. T.A.L. was supported by the University of Minnesota

Doctoral Dissertation Fellowship.The Minnesota Supercomputing Institute provided computing

and data storage resources. We thank Yaniv Brandvain and Ken Kozak for discussions and

advice on analyses. We also thank Erik Runquist, Brooke Kern, Danielle Schoenecker, Ben

Greene, and Isaac Olson for their assistance with data collection.

ABSTRACT

Rapid evolution may play an important role in the range expansion of invasive species and modify forecasts of invasion, which are the backbone of land management strategies. However, losses of genetic variation associated with colonization bottlenecks may constrain trait and niche divergence at leading range edges, thereby impacting management decisions that anticipate future range expansion. The spatial and temporal scales over which adaptation contributes to invasion dynamics remains unresolved. We leveraged detailed records of the ~130 year invasion history of the invasive polyploid plant, leafy spurge (*Euphorbia virgata*), across ~500km in Minnesota, U.S.A. We examined the consequences of range expansion for population genomic diversity, niche breadth, and the evolution of germination behavior. Using genotyping-by-sequencing, we found some population structure in the range core, where introduction occurred, but panmixia among all other populations. Range expansion was accompanied by only modest losses in sequence diversity, with small, isolated populations at the leading edge harboring similar levels of diversity to those in the range core. The climatic niche expanded during most of range expansion, and the niche of the range core was largely non-overlapping with the invasion front. Ecological niche models indicated that mean temperature of the warmest quarter was the strongest determinant of habitat suitability and that populations at the leading edge had the lowest habitat suitability. Guided by these findings, we tested for rapid evolution in germination behavior over the time course of range expansion using a common garden experiment and temperature manipulations. Germination behavior diverged from early to late phases of the invasion, with populations from later phases having higher dormancy at lower temperatures. Our results suggest that trait evolution may have contributed to niche expansion during invasion and that distribution models, which inform future management planning, may underestimate invasion potential without accounting for evolution.

Introduction

63
Invasive species experience considerable changes to genetic variation during the

process of introduction and subsequent invasion (Lee 2002; Suarez and Tsutsui 2008). In

particular, it has been well-documented that founder effects and genetic drift can cause

substantial losses of genetic diversity during the colonization process (Dlugosch and Parker

2008; Uller and Leimu 2011). Following initial establishment, further losses of variation may

occur during range expansion. However, the magnitude of changes in genetic variation depends

upon the number of introductions and the severity of population bottlenecks (Nei et al. 1975;

Uller and Leimu 2011; Welles and Dlugosch 2019). Such changes in genetic variation early in

the invasion process may influence the capacity for adaptation, forecasts of range expansion,

and subsequent management decisions.

Following colonization, some non-native species exhibit rapid population growth and

dispersal into new environments (Sakai et al. 2001). The process of invasion is often highly

variable, involving repeated founder events and density-dependent population growth,

especially Allee effects (Melbourne and Hastings 2009; Sullivan et al. 2017). In addition to

affecting the speed of invasion, these population fluctuations can influence levels of genetic

diversity and structure across an invaded range (Austerlitz et al. 1997; Excoffier 2004). For

example, range expansion is expected to cause a reduction in allelic richness and

heterozygosity with increasing distance from the origin of expansion (Slatkin and Excoffier 2012;

Peter and Slatkin 2013). The prevalence of drift during invasion may also cause populations to

depart from migration-drift equilibrium, resulting in a lack of isolation-by-distance (Wright 1943;

Slatkin 1987, 1993; Hutchison and Templeton 1999). Last, mutations arising at the range edge

may rise in frequency due to genetic drift, "surf" along the expanding front, and travel long

distances (Klopfstein et al. 2006; Excoffier and Ray 2008). Models of allele surfing indicate that

rapid range expansion can produce clinal variation in allele frequencies and increase the

frequency of loci with private alleles (Klopfstein et al. 2006; Excoffier and Ray 2008; Goodsman

et al. 2014). Such clines can emerge for any type of mutation (beneficial, neutral, deleterious),

and therefore could reflect drift and/or selection (Lehe et al. 2012; Peischl et al. 2013; Koski et

al. 2019). Overall, the prevalence of drift during range expansion has the potential to influence

the capacity for adaptation as organisms encounter novel environments, particularly when

functionally-important allelic variation is lost.

An increasing body of evidence suggests that rapid phenotypic evolution can be

important to the process of range expansion (Colautti and Barrett 2013; Hodgins et al. 2018;

Selechnik et al. 2019; Ma et al. 2020). Forecasts of range expansion in invasive species can

underpredict the potential extent of invasion if they assume a species does not evolve or adapt

over short time scales (Chardon et al. 2020; Collart et al. 2020). A recent meta-analysis

indicated that the signature of local adaptation in invasive species was at least as strong as in

native species, even when accounting for variation in life history (Oduor et al. 2016). Invasive

species frequently expand across strong environmental gradients and into novel niche space

(Atwater et al. 2018; Bates and Bertelsmeier 2021). Such niche expansion may require adaptive

evolution at the invasion front (Chown et al. 2015; Moran et al. 2017; Hodgins et al. 2018). For

example, purple loosestrife rapidly diverged in flowering time and plant size during invasion

across latitudinal gradients in growing season length (Colautti and Barrett 2013). Despite

evidence of rapid evolution in some systems, range expansion may not involve any changes in

the organism's niche if the only limit to spatial expansion is dispersal and time. As such, the

apparent expansion of the climate niche with invasion may not actually involve the evolution of

ecologically-important traits. While tests of local adaptation within an invaded range remain few,

there is growing appreciation that rapid evolution is likely to shape the trajectory of range

expansion in non-native species (Hodgins et al. 2018; Woods and Sultan 2022).

Phenotypic evolution during range expansion may be caused by spatially-variable

selection and/or neutral processes (e.g., spatial sorting) (Keller and Taylor 2008; Colautti and

Lau 2015; Hodgins et al. 2018). Adaptive evolution, in particular, may be paramount to the

invasion process if selection in response to novel environments results in trait changes that

enhance a species' capacity to establish in new habitats (Prentis et al. 2008; Williams et al.

2016; Hodgins et al. 2018; Woods and Sultan 2022). While reciprocal transplant experiments

are the gold standard for testing for adaptation, the translocation of invasive species for these

experiments is subject to ethical concerns and legal restrictions in many areas. Alternatively,

researchers have started to use ecological niche models (ENMs) to identify important

environmental gradients that span from optimal to marginal habitat, such as from a range core

to edge. Predictions are then made about traits that may promote adaptation to the novel

environments found in marginal habitats (Searcy and Shaffer 2016; Dixon and Busch 2017;

Capblancq et al. 2020; Morente-López et al. 2022). Finally, common garden experiments can

determine whether the putative traits under selection have differentiated across the key

environmental gradients identified by ENMs. Taken together, this series of approaches can

provide insight into the role of adaptation in the process of niche expansion at leading range

edges.

129   Among invasive plant species, polyploidy is prevalent (Pandit et al. 2011) and can

influence the process of range expansion (Van de Peer et al. 2021). The frequency of polyploid

species increases with higher latitudes, lower temperatures, and seasonally drier environments

(Brochmann et al. 2004; Rice et al. 2019). Direct effects of polyploidy on physiological,

morphological, and phenological traits may facilitate niche shifts (Glennon et al. 2014; Blaine

Marchant et al. 2016; Brittingham et al. 2018; Wang et al. 2022) and preadapt polyploids to new

environments (Treier et al. 2009; Lachmuth et al. 2010). Polyploidy may also influence the

capacity for adaptation during range expansion as new environments are encountered (te Beest

et al. 2012; Baniaga et al. 2020). Although genetic drift during range expansion can cause

losses of genetic diversity, drift may have less severe effects (e.g., inbreeding depression) in

polyploids relative to diploids when there is polysomic inheritance (Moody et al. 1993; Soltis and

Soltis 2000). Despite numerous polyploid invaders, they have been the subject of few studies

because of substantial challenges with the application of evolutionary genetic analyses that were developed for diploids (Rutland et al. 2021).

In this study, we used a well-documented chronosequence of invasion to examine the consequences of range expansion for population genomic diversity, climatic niche breadth, and the evolution of germination behavior in the polyploid, leafy spurge (*Euphorbia virgata*). We focused on one area of introduction to southwestern Minnesota, U.S.A and subsequent range expansion to the north and east. We were interested in examining the severity of losses of genetic diversity following introduction and its potential consequences for invasion, particularly since existing species distribution models predict a low probability of range expansion at the current leading range edge (Lake et al. 2020). Introduction to this region occurred in the 1890s in southwestern Minnesota with subsequent range expansion to northeastern Minnesota (ca., 500 km), where populations are currently rare, isolated, and small. Based on a detailed historical occurrence dataset, we defined a range core, area of early expansion, area of late expansion, and invasion front.

First, we sampled populations along the gradient from core to invasion front using two sampling schemes to quantify population genomic diversity and structure (using reduced representation sequencing). Samples of multiple individuals from 14 populations (population samples) allowed us to quantify changes in sequence diversity among populations over the course of range expansion. Samples of single individuals from 157 populations (landscape samples) allowed us to test for fine-scale population structure (e.g., isolation-by-distance) over the time series of range expansion. Second, we tested whether range expansion involved niche expansion - i.e., occurred into novel climatic environments. Third, we developed an ecological niche model (ENM) to test whether habitat suitability declines from range core to invasion front and to determine which environmental gradients are most strongly associated with high versus low habitat suitability. Warm season temperature had the greatest positive contribution to habitat suitability. Because temperature is known to modulate germination behavior in leafy

spurge, and because establishment in new habitats is dependent upon successful germination

timing, we focused on this trait for common garden experiments. Past work has also suggested

that shifts in seed dormancy might facilitate invasion at leading range edges (Mathias and Kisdi

2002; Travis et al. 2021). We examined the responses of seeds from early versus late in

invasion to five temperature regimes in a common garden experiment. We specifically tested

whether there was an interaction between geographic region (early vs. late invasion stages) and

temperature regime, which would indicate divergence in germination behavior over the course

of invasion.

Methods

<u>Invasion and natural history of leafy spurge</u>

Natural history

Leafy spurge has invaded approximately two million hectares across the northern tier of

the United States and southern Canada (Duncan et al. 2004). While it is most commonly found

in dry, open sites with well-drained soils (e.g., prairies), it can occasionally occur in seasonally

wet meadows and riparian areas (Selleck et al. 1962). Leafy spurge impacts rangelands and

natural habitats by competitively displacing native species (Hein and Miller 1992). When

damaged, plants exude a toxic white latex that deters grazing (Lym and Kirby 1987; Lym 1998).

Leafy spurge spreads locally via rhizomes and ballistic seed dispersal (Morrow 1979).

Longer distance dispersal has been proposed to occur via animals or agricultural machinery

(Pemberton 1988; Lacey et al. 1992). Seeds germinate in spring or may remain dormant in soil

for at least two years (Hanson and Rudd 1933; Selleck et al. 1962). Flowers are insect

pollinated and the mating system is primarily outcrossing (Selleck et al. 1962).

Leafy spurge is an auto-allohexaploid that likely originated from hybridization between

closely-related *Euphorbia* species, although the progenitor species are not yet known (Schulz-

Schaeffer and Gerhardt 1989; Riina et al. 2013). It has been the subject of several genomic

investigations but lacks a full genome assembly and annotation (Chao et al. 2005; Horvath et al.

2015, 2018; West et al. 2023).

Invasion history

One introduction of leafy spurge occurred into southwestern Minnesota ca., 1890

purportedly via contaminated grains imported from southern Russia (Batho 1932; Hanson and

Rudd 1933; Dunn 1985). Following introduction, the range expanded to eastern South Dakota

by ca., 1902 (Bakke 1936), eastern North Dakota by ca. 1909 (Hanson and Rudd 1933), and

southern Manitoba and Saskatchewan by the 1920s (Batho 1932; Selleck et al., 1962). Hanson

and Rudd (1933) documented in detail the distribution of leafy spurge across Minnesota and

neighboring regions, providing a baseline for understanding the timeline of subsequent range

expansion (Figure S1). By the late 1970s, leafy spurge had become common throughout

grasslands of the north-central plains (Dunn 1979, 1985). Invasion of the boreal forest region of

northeastern Minnesota began in the 1940s and 1950s with isolated occurrences (Lakela 1965)

and populations were not common until the 1990s. This invasion front has persisted with limited

expansion further northeast.

Delineating the timeline of range expansion

We digitized the earliest known point record map (Hanson and Rudd 1933) using ArcGIS

Pro (ESRI, 2022). We then applied empirical Bayesian kriging to produce a continuous density

surface that represented the density of populations in the north-central plains. From this density

surface, we applied an equal-interval threshold to demarcate a range core, area of early

expansion, area of late expansion, and invasion front that corresponded to four density

categories across Minnesota and surrounding states (Figure 1A). We verified these

demarcations with published accounts of the invasion history (described above).

Population genomic diversity and structure

Sampling and sequencing

In 2019, we collected leaf tissue from six individuals in each of 14 populations distributed evenly across Minnesota (*hereafter:* population samples; Figure 1B). In addition, we collected tissue from one individual in each of 157 populations distributed relatively evenly across Minnesota, eastern South Dakota, eastern North Dakota, and western Wisconsin (*hereafter:* landscape samples; Figure 1C). We sampled tissue from individuals that were at least five meters apart to minimize collecting from the same genet and placed tissues immediately in silica for preservation until DNA extraction (Table S1).

We extracted DNA using QIAGEN DNeasy Plant Mini Kits (QIAGEN Inc.). Dual-indexed GBS (genotyping-by-sequencing) libraries were created using the BamHI + NsiI enzyme combination. All libraries were pooled and sequenced on an Illumina NovaSeq System (Illumina Inc., San Diego, CA, USA) with 1x100-bp sequencing. Once sequenced, the reads were demultiplexed and balanced with a mean quality score ≥ Q30 for all libraries. We filtered low-quality bases using Trimmomatic (Bolger et al. 2014) and used Stacks v.2.5.9 (Rochette et al. 2019) to build loci *de novo* (i.e., without aligning reads to a reference genome). Overall, we obtained 510 million reads across the 241 samples (599,386 – 3,376,078 of raw reads per individual). Mean read depth per locus ranged from 14x to 26x (Supplementary Methods).

We called SNPs using polyRAD v.2.0.0 (Clark et al. 2019), a Bayesian algorithm designed for polyploid GBS data. PolyRAD estimates the genotype probabilities for each individual from read depth distributions with ploidy level as a prior. First, we filtered our dataset using the H_{ind}/H_e statistic to cull likely paralogous loci (Clark et al. 2022). Next, we estimated posterior probabilities for each genotype using the 'IterateHWE' function with Hardy-Weinberg equilibrium as the prior (Gerard and Ferrão 2020; Clark et al. 2022). For each individual at each locus, we exported the most probable genotype for subsequent analyses (Supplementary Methods).

We implemented a second filtering step to account for potential biases caused by

homoeologous loci present in our dataset. Because leafy spurge is an auto-allohexaploid

(Schulz-Schaeffer and Gerhardt 1989), we expect homoeologous loci to have a 2:1 allelic ratio

(e.g. AAAABB genotype; Horvath et al. 2018). Indeed, we identified a peak in the minor allele

frequency spectrum around 0.33 (Figure S2). We removed loci with a minor allele frequency

above 0.26 from our dataset because they are likely to have an excess of homoeologous

genotypes (Figure S2). While essential, this second filtering step limits our capacity to

understand absolute levels of sequence diversity. However, our primary goal was to examine

changes in sequence diversity over the course of invasion rather than absolute quantities. After

filtering, 3,176 loci remained for downstream analyses.

Population structure

We performed an analysis of molecular variance (AMOVA) to quantify the proportion of

genetic variation partitioned among populations, among individuals within populations, and

within individuals (Excoffier et al. 1992; Meirmans 2012, 2020). We estimated genetic variance

components using the rho statistic (Ronfort et al. 1998) and determined significance using

permutation tests (n= 999) using the R package poppr v.2.8.6 (Kamvar et al. 2015). Further, we

checked for clonality among samples using the 'clonecorrect' function in (Kamvar et al. 2015).

We assessed population structure using the Bayesian clustering algorithm STRUCTURE

v.2.3.4 (Pritchard et al. 2000). We ran the analysis 10x for each of K = 1-10 with 500,000

Markov Chain Monte Carlo iterations with a 50,000-run burn-in period, specifying an admixture

model with the assumption of uncorrelated allele frequencies. We used 'structure_threader' in

Python 3 (Pina-Martins et al. 2017) to parallelize runs across clusters (K = 1 – 10). We

determined the most plausible number of clusters using the ∆K method (Evanno et al. 2005)

and STRUCTURE HARVESTER web v.0.6.94 (Earl and vonHoldt 2012).

We performed principal component analyses (PCA) to examine population structure

using GENODIVE v.3.0.4 (Meirmans 2020). We performed PCA separately for the population

samples and the landscape samples.

We tested for isolation by distance (IBD; Wright 1943) in the population samples by

estimating genetic differentiation as G_{ST} (Dufresne et al. 2014) using GENODIVE v.3.0.4

(Meirmans 2020). We also tested for IBD in the landscape samples by calculating Nei's genetic

distance (D; Nei 1972) (Meirmans 2020). We subset the landscape samples according to

successive stages of range expansion and tested for IBD within each subset (i.e., within range

core, then successively including areas of early expansion, late expansion, and invasion front).

We tested for a relationship between the pairwise genetic and geographic distance (kilometers)

matrices using Mantel tests with 9,999 permutations in the R package 'adegenet' v.2.1.8

(Jombart 2008).

Tests for changes in population genetic diversity during range expansion

For each of the 14 population samples, we estimated the inbreeding coefficient (G_{IS}) and

observed heterozygosity (H_o; gametic heterozygosity; Moody et al. 1993), which corrects for

potential overestimates of heterozygosity in polyploids by calculating the fraction of

heterozygotic gametes for each genotype, using GENODIVE v.3.0.4 (Meirmans 2020). We also

calculated the number of private alleles (P) using poppr v.2.8.6 (Kamvar et al., 2014). We

calculated Tajima's *D* (Tajima 1989) using DNASp v.6.0 (Rozas et al. 2017) to gauge if

populations have an excess or deficit of rare alleles, which can be indicative of population

expansion following a bottleneck (negative *D*) or sudden population contraction (positive *D*),

respectively. Because we filtered loci with higher minor allele frequencies, Tajima's *D* should be

biased to lower values. Finally, we tested for evidence of genetic bottlenecks based on linkage

disequilibrium using NeEstimator v.2.0.1 (Do et al. 2014).

We tested for changes in population genetic parameters (H_o, G_{IS}, P, and Tajima's D) as

populations dispersed beyond the range core (Table 1). For each statistic separately, we used a

multiple linear regression that included latitude, longitude, and their interaction as independent

variables with the R package 'car' v.3.1-2 (Weisberg, 2019). As all late expansion and invasion

front populations are located either north or east of the range core, latitude and longitude

describe the northern and eastern invasion spread, respectively. In the model of private alleles,

we identified the 'Winona' population as an outlier using diagnostic plots of residuals (Figure

S3), so we removed it from the analysis. We were unable to conduct such a geographic test of

N_e from the NeEstimator analyses because all but three populations had an N_e of infinity (Table

S2).

Niche breadth and habitat suitability

Environmental data

We downloaded 19 bioclimatic variables at a 30 arcsecond resolution (~1 km) from

Worldclim (http://worldclim.org/version2). We used a principal component analysis based on the

157 landscape samples to examine patterns of correlation among bioclimatic variables. The

biplot of the first two principal components (74.7% variance explained) showed that there were

three sets of correlated variables: one set includes mostly precipitation variables, a second set

includes mostly temperature variables, and a third set includes variables that describe

seasonality and temperature ranges (Figure S4). Because the selection of one variable from a

set of correlated ones is challenging (e.g., choosing one precipitation variable from a set of

correlated precipitation variables), studies have advocated for choosing variables that are

biologically most relevant to the organism and largely uncorrelated (Chapman et al. 2017;

Petitpierre et al. 2017).

We selected minimum temperature of the coldest month (Bio 6), mean temperature of

the warmest quarter (Bio 10), and precipitation of the warmest quarter (Bio 16). These three

variables provide biologically meaningful axes of climate variation that are relevant to key life

history stages for an herbaceous perennial (Petitpierre et al. 2017; Chapman et al. 2017).

Specifically, minimum cold temperatures are important for overwintering and cold tolerance

(Chapman et al. 2017), whereas the temperature of the warmest quarter likely influences seed

germination, plant growth, and phenological transitions (Wolkovich et al. 2013). Precipitation of

the warmest quarter likely also affects plant growth and reproduction, with lower precipitation

associated with reduced growth and increased drought stress in northern temperate

ecosystems (Petitpierre et al. 2017; Gorton et al. 2019).

Tests for niche differentiation during range expansion

We tested for climatic niche differentiation between the range core, early expansion, late

expansion, and invasion front using the 'ecospat' R package v.3.5.1 (Di Cola et al. 2017). We

sampled the total extent of the background environmental space with a principal component

analysis using 1500 random points drawn from a bounding box centered on Minnesota

(Latitude: min: 43°N, max: 50°N; Longitude: min: -98°W, max: -89°W). We then used the

landscape sample localities to calculate the niche boundaries and density of occurrence for

each portion of the range within the environmental PCA space.

For all pairwise comparisons of the range core, early expansion, late expansion, and

invasion front, we quantified four measures of niche differentiation. First, we used Schoener's *D*

to quantify the similarity in niche by incorporating both niche breadth and density (Warren et al.

2008). Values of *D* can vary from 0 (no overlap) to 1 (complete overlap). For each pair of

regions, we then calculated what proportion of the combined niche space represented niche

stability, expansion, and unfilling. In this framework, if 'A' represents the older potion of the

range (range core) and 'B' represents the more recent portion of the range (invasion front),

niche stability is the proportion of niche B that overlaps A, niche expansion is the proportion of

niche B that does not overlap A, and niche unfilling is the proportion niche A that does not

overlap B.

We used permutation tests to determine if values of Schoener's D, expansion, stability,

and unfilling were equivalent between the range core, early expansion, late expansion, and

invasion front. In the niche equivalency tests, the data were pooled and then randomly assigned

to one group of the pairwise range comparisons for 999 permutations. For each permutation, we

computed all four statistics. We rejected the null hypothesis of niche equivalency if the observed

Schoener's D value was less than 95% of permuted D values. Similarly, we rejected niche

equivalency based on the combined niche space if observed stability was less than 95% of

permuted values and observed niche expansion or unfilling was greater than 95% of permuted

values.

Ecological niche model

We developed an ecological niche model (ENM) using MaxEnt v.3.4.3 (Merow et al.

2013; Phillips et al. 2017) with the 'dismo' package v.1.3-14 in R (Hijmans et al. 2022). Our goal

was to identify environmental gradients that could potentially drive phenotypic evolution during

range expansion (Elith and Leathwick 2009; Araújo et al. 2019; Morente-López et al. 2022). We

built ENMs with the same bioclimatic variables and in the same bounding box as the analyses of

niche differentiation. We found that temperature of the warmest quarter (Bio10) had the

strongest influence on the probability of occurrence. To examine whether this result was robust,

we built three other ENMs using alternative sets of environmental variables (Table S3). In every

case, temperature of the warmest quarter was by far the strongest predictor (Table S3).

We built all ENMs using occurrence records from our tissue collection sites and 10,000

background points. We excluded threshold and hinge features during the model building

process as the preliminary models that included these features tended to be overspecified. We

used five-fold cross-validation to assess model performance; data were randomly partitioned

into five equal groupings and 80% of data were used for training and 20% were used for

evaluation. Model predictions are a mean of the five cross-validated models.

We evaluated models using AUC and sensitivity, which were calculated using the

withheld dataset. AUC characterizes model discrimination ability and ranges between 0 and 1,

with higher values indicating greater model performance and a value of 0.5 indicating that model

discrimination is no better than random (Phillips and Dudík 2008). Sensitivity quantifies the

proportion of correctly identified positives. We calculated sensitivity where the sum of the true

positive rate and true negative rate was maximized (threshold = 0.53). We then used the

variable permutation importance and percent contribution to identify which environmental

variable had the greatest contribution to habitat suitability. We also visualized response curves

of each environmental variable to ensure models were not overspecified. Hereafter, "habitat

suitability" refers to the probability of occurrence from our ENMs based upon climatic factors.

To test for divergence in germination behavior during range expansion, we focused on

mean temperature of the warmest quarter since it most strongly affected predicted habitat

suitability in all ENMs. Populations from early in the invasion had high predicted habitat

suitability and higher warm season temperatures (all above 20°C) compared to those from later

in the invasion, which had lower predicted habitat suitability and lower warm season

temperatures (all below 20°C; Fig. 6A,B). Guided by these results, we divided the 14

populations for which we had seed collections into two groups: early (n=8) versus late (n=6) in

invasion and exposed them to five temperature regimes (see below).

Tests for differentiation in germination behavior during range expansion

We collected seeds in 2019 from 14 populations (8 – 24 maternal families/pop; Table

S4). Seeds were collected from individuals that were at least 5 meters apart. We stored and

after-ripened seeds in an indoor environment for two years prior to the germination experiment

(Wicks and Derscheid 1958). Seeds were pooled within populations prior to applying

treatments. We were unable to collect seeds from every population used in the population

genetic survey because some had already dispersed seeds prior to our collection effort.

400   We examined the effects of temperature regime and source geographic region (early vs.

late in invasion) on germination. Five temperature regimes were designed to mimic the full

range of variation in daytime and nighttime temperatures during the spring and summer in this

region (14 hour day /10 hour night periods: 15/5 °C, 20/10 °C, 25/15 °C, 30/20 °C, and 35/25

°C). The experiment was conducted in five successive rounds in two growth chambers

(Conviron Inc.). We conducted each temperature regime twice – i.e., once in each growth

chamber – to control for growth chamber effects.

407   For each round, we placed ten seeds per population in 60 x 15mm polystyrene Petri

dishes containing 2 mL sterile distilled water, lined with one Whatman #1 filter paper, and

sealed with parafilm. Each population was replicated three times per chamber for a total of 42

dishes per treatment per chamber or 84 dishes per round. Dishes were wrapped in aluminum

foil to block light, which can inhibit germination (Selleck et al., 1962). Every 24 hours we

recorded the number of germinated seeds (emergence of radicle) per dish. After each treatment

period ended, we tested whether ungerminated seeds were viable by soaking bisected seeds in

a 1% Tetrazolium solution for 24 hours (Verma and Majee 2013). Red staining of tissues

indicates that seeds are viable. For analyses of germination, we included the number of

germinated seeds out of the total number of germinated plus viable (but ungerminated) seeds.

417   We tested for the effects of temperature regime (categorical), source geographic region,

and their interaction on germination probability using a mixed-effects model with a binomial

family. Experimental round and popullation were included as random effects. We evaluated

significance with Type III tests. All models were run using the 'mixed' function in the 'afex'

package v.1.3-0 (Singmann et al. 2016) in R v.4.0.2 (R Development Core Team, 2015). We

used linear contrasts to test for differences in germination between geographic regions for each

temperature regime category individually.

Results

<u>Population genomic consequences of range expansion</u>

Analysis of molecular variance (AMOVA) revealed significant partitioning of genetic

variance among populations (13.7%; $P < 0.001$), among individuals within populations (8.4%; P

< 0.001), and within individuals (77.8%; $P < 0.001$) (Table S5). No clonal genotypes were

detected in the dataset. STRUCTURE indicated that the optimal number of clusters was three

($K = 3$) (Table S6). All individuals were assigned primarily to one cluster regardless of where the

population was found in the invasive range (i.e., core, early expansion, late expansion, invasion

front). There was some evidence of population structure in the other two clusters; however,

there was no clear geographic pattern (Figure 2). We also examined K=4, which similarly

revealed one major cluster for each individual and a minor one where there was some evidence

of population structure. For example, four northern populations near the invasion front share the

same minority cluster (Koochiching, St. Louis, Duluth, Aitkin); this cluster is also shared with two

populations in the range core (Figure 2).

The PCA did not indicate substantial population structure across the invaded range

based on either population or landscape samples. For the population samples, there was some

evidence of differentiation among three populations in or near the range core (Figure 3; PC1

and PC2 explained: 6.8% and 5.9% of variance, respectively). The PCA of landscape samples

did not reveal a relationship between genetic similarity and geography over the time course of

range expansion (Figure 3; PC1 and PC2 explained 1.7% and 1.3% of variance, respectively).

There was no evidence of isolation by distance (IBD) for either population ($R^2 = 0.02$; P

$= 0.418$; Table 2; Table S7) or landscape samples ($R^2 = 0.05$; $P = 0.676$; Table S7). We also did

not detect IBD when landscape samples were subset according to the four invasion phases that

we defined (Figure 4).

Range expansion from the core area of invasion was accompanied by modest changes

in genetic diversity. Heterozygosity declined from the core to the invasion front, as indicated by

a significant interaction of latitude and longitude ($P = 0.013$; Table 3). However, the number of

private alleles, Tajima's D, and the inbreeding coefficient did not change over the course of

range expansion (Table 3). Estimates of N_e based on linkage disequilibrium using NeEstimator

suggested that N_e was large (estimated at infinity) in all but three populations: Blue Earth (N_e =

134), Crow Wing (N_e = 210), and Koochiching (N_e = 17,836). Those three populations occur in

the range core, late invasion, and invasion front portions of the range, respectively.

<u>Niche differentiation during range expansion</u>

The climatic niche expanded during invasion. Relative to the range core, the early

expansion niche represented a sizable increase in niche breadth (Figure 5A; Table S8; overlap

= 0.43; $P < 0.01$; expansion = 0.37; $P < 0.01$; stability = 0.63; $P < 0.01$). When comparing the

early expansion and late expansion niches, there was similar evidence for a niche shift (Figure

5B; Table S8; overlap = 0.36, $P < 0.01$; expansion = 0.39, $P < 0.01$; stability = 0.61, $P < 0.01$).

Between the late expansion niche and the invasion front, the null hypothesis of niche

equivalency was not rejected (Table S8); the invasion front niche was contained within the late

expansion niche (Figure 5C). When comparing range core to invasion front, there was near-zero

niche overlap (Figure 5D; Table S8; overlap = 0.03, $P < 0.01$; stability = 0.06, $P < 0.01$) and high

expansion (Table S8; expansion = 0.94, $P < 0.01$). Niche differences between the core and

invasion front were most apparent along environmental axes related to temperature of the

warmest quarter and minimum temperature of the coldest month, rather than precipitation

(Figure S6).

Ecological niche model

The model AUC (0.79) and sensitivity (0.75) metrics indicated moderately high

discrimination and accuracy (Figure 6A). The variable response curves indicated the ENM was

not overspecified (Figure S7). The mean temperature of the warmest quarter constituted the

most important variable for the predicted probability of occurrence, hereafter "habitat suitability"

(percent contribution = 81.5%; permutation importance = 67.8%). Warmer temperatures were

associated with an increase in habitat suitability (R = 0.67; Figure S8). The minimum

temperature of the coldest month had the second highest importance (percent contribution =

11.2%; permutation importance = 17.9%) and was modestly associated with increased habitat

suitability (R = 0.31; Figure S8). Precipitation of the warmest quarter had the lowest importance

(percent contribution = 7.3%; permutation importance = 14.3%) and was weakly correlated with

habitat suitability (R = 0.02; Figure S8).

Populations from early versus late in invasion were distinct along the mean temperature

of the warmest quarter axis with a disjunction at 20°C. This temperature also distinguished

highly suitable from less suitable habitat in the ENM (Figure 6A & 6B).

Evolution of germination behavior during range expansion

Populations from early versus late in invasion responded differently to temperature

regimes (source geographic region x temperature regime interaction: P = 0.003; Table S9;

Figure 6C). The linear contrast between geographic regions for the 20/10 °C treatment, the

second coldest treatment, was marginally significant (p = 0.09); contrasts between regions for

all other temperature treatments were not significant (p > 0.77; Table S10). Germination

probability increased with temperature for both geographic regions and plateaued (P < 0.001;

Table S9; Figure 6C).

Discussion

Rapid evolution is increasingly recognized as an important process contributing to range

expansion and influencing forecasts of future invasion (Prentis et al. 2008; van Boheemen et al.

2019; Clements and Jones 2021). However, our understanding of the temporal and spatial scale

over which niche and trait divergence contribute to invasion at leading range edges remains

unresolved. Our study took advantage of a well-documented invasion history to synthesize the

consequences of recent range expansion for population genomic diversity, niche breadth, and

germination behavior, a trait important in the colonization of new habitats. We found that leafy

spurge populations experienced only modest losses in sequence diversity over the

chronosequence of invasion and that recruitment within invaded populations occurred via seeds

rather than strictly from clonal expansion. Range expansion involved climatic niche expansion

and ecological niche models suggested that mean temperature of the warmest quarter had the

strongest influence on habitat suitability. Populations differentiated in germination behavior in

response to temperature, where leading edge populations exhibited a trend of increased

dormancy at low temperatures. Our results suggest that evolution during range expansion may

be important to consider in the development of models forecasting range shifts under current

and future climates.

Loss of genetic diversity during range expansion may have fitness consequences and

limit adaptive capacity (Lee 2002; Dlugosch and Parker 2008). The magnitude of within-

population diversity is also potentially important for the success of eradication measures, which

may not be effective against all genotypes. We found that range expansion was only

accompanied by small losses in heterozygosity but no changes in inbreeding coefficient or the

number of private alleles. We also did not observe evidence of genetic bottlenecks based on

linkage disequilibrium. We identified some divergence among populations in or near the range

core, while others occupied intermediate positions in PCA space. Prevalent long-distance

dispersal from the expanding core to the invasion front could reintroduce allelic variation lost

due to bottlenecks in the colonization process. Additionally, admixture among independently

introduced populations (i.e., invasion fronts not included in our analyses) could contribute to

within-population genetic diversity, especially when there is genetic divergence between

sources of introductions (Gaskin and Schaal 2002; Keller and Taylor 2010; Gibson et al. 2021).

Although we do not observe novel genotypes or elevated numbers of private alleles at the

periphery of our focal region, which would be indicative of admixture between invasion fronts,

more work is needed to evaluate this potential influence.

Regardless of the mechanism, the extent of within-population variation may have

important implications for eradication and management. Although leafy spurge can spread

clonally, it does not appear that populations constitute one or few clones that spread widely

upon colonization. Instead, recruitment from seeds contributes substantially to population

growth. This finding has implications for integrated pest management (IPM) strategies.

Management techniques that reduce above-ground productivity and seed production may be

important in addition to those that reduce clonal spread by disturbing root systems (e.g.

biocontrol agents that feed primarily on roots; Hanson et al. 1997). In addition to natural

disturbance, improperly timed grazing or herbicide applications could create open spaces for

recruitment from seeds and inadvertently increase density and abundance of the invader

(Gaskin et al. 2022). As seeds may remain dormant for as long as five years (Selleck 1962),

seed banks may further complicate control efforts. The combination of recruitment from seeds

and clonal growth contributes to the population dynamics of leafy spurge, and therefore, multiple

eradication approaches may be necessary to cause local population extinction.

In addition to gene flow during invasion, polyploidy may be an important factor

influencing losses of genetic variation during colonization bottlenecks and the capacity for range

expansion. Polyploids often maintain higher levels of genetic variation (Otto and Whitton 2000)

and there is some evidence that phenotypic plasticity is greater in synthetically produced

autopolyploids (Mattingly and Hovick 2023). These factors have been used to explain why

polyploids may be better invaders than diploids (Pandit et al. 2011). Leafy spurge is an auto-

allohexaploid, suggesting that its higher N_e should facilitate the maintenance of genetic diversity

within populations and minimize divergence among them. Polyploidy has also been suggested

to increase the capacity for adaptive evolution (Otto and Whitton 2000), which could also

contribute to success as an invader. It is important to recognize that losses of diversity in DNA

sequences do not necessarily translate to losses of variation in quantitative traits (Reed and

Frankham 2001). Nevertheless, it is possible that polyploidy contributed to rapid invasion in

leafy spurge, but more work is needed to distinguish the contribution of polyploidy from other

factors.

Accumulating evidence suggests that invasive plant species frequently undergo climatic

niche shifts during range expansion (Medley 2010; Atwater et al. 2018; van Boheemen et al.

2019; Bates and Bertelsmeier 2021; but see Petitpierre et al. 2012; Liu et al. 2020). Such niche

shifts can complicate the reliability of models that seek to predict future invasions, as these

models generally assume niche conservation. Consistent with past findings, we observed

climatic niche expansion throughout most of range expansion, except from the late expansion

region to the invasion front. From an ENM, we found that warm season temperature had the

strongest influence on habitat suitability and therefore may be one source of divergent selection

from range core to invasion front. Of course, climate change has already caused poleward shifts

in plant species distributions through climatic niche matching (Parmesan 2006; Clements and

Ditommaso 2011; Parmesan and Hanley 2015). Therefore, some range expansion may simply

involve dispersal to already climatically suitable habitats. However, adaptation may be

necessary for continued range expansion, especially where the species is already at its climatic

niche limit (Clements and Ditommaso 2011). In leafy spurge, minimal range expansion

northward has been observed in the last 30+ years and populations remain small/low density at

the invasion front. In addition, ENMs suggest that invasion front populations have very low

habitat suitability in terms of climate. Overall, our results are inconsistent with the hypothesis

that the leading edge is suitable but expansion is limited by dispersal. Instead, responses to

divergent selection may be important for persistence at the leading range edge and for further

range expansion. These findings also hold implications for management approaches:

recognizing and accounting for niche shifts could improve the accuracy of distribution models for

forecasting future invasions, and by extension, help optimize resource allocation for eradication

efforts. Integrating knowledge of which environmental factors underly niche expansion will be

key to designing more effective management strategies that not only suppress existing

populations but also anticipate future invasions.

Divergence in trait expression at a leading range edge can be driven by local adaptation,

phenotypic plasticity, and/or maternal environmental effects (Des Roches et al. 2017;

Westerband et al. 2021). Phenotypic plasticity is considered important during early stages of

invasion because it allows introduced populations to establish in a broader range of

environmental conditions (Sexton et al. 2002; Richards et al. 2006; Funk 2008; Lande 2015). In

germination traits, plasticity could represent a means of habitat selection and niche construction

(Donohue 2003, 2005) by which leafy spurge in the leading edge germinates optimally at the

onset of spring conditions (warmer temperatures in northern latitudes). Likewise, rapid evolution

during range expansion can result from selection on loci that influence dormancy and/or

germination timing (Clements and Ditommaso 2011; Hodgins et al. 2018; Clements and Jones

2021). We found a trend of increased dormancy at lower temperatures in leading edge

populations (p = 0.09 for 20/10 °C treatment). One possibility is that germination at colder

temperatures exposes seedlings to more unpredictable environments (e.g., late season frost)

and thus that selection favored germination later in the season for leafy spurge at its northern

range limit. Interestingly, other work has suggested that reduced dormancy evolves at leading

range edges (Tabassum and Leishman 2018), contrary to our findings. We cannot exclude the

possibility that maternal environmental effects influenced our estimates of germination

probability because we used field-collected seeds. However, our findings are contrary to what

we would have expected if maternal environmental effects had been important: we would have

expected higher germination for leading edge populations in cold treatments because those

populations come from colder environments. Although warm season temperature is most

strongly associated with habitat suitability and relevant to germination in leafy spurge, it is also

possible that other (correlated) variables influenced the evolution of germination behavior.

Overall, our results suggest that range expansion involved niche expansion and potentially the

evolution of germination timing, which may have been important in establishment at the leading

range edge.

Early life history transitions are thought to be under strong selection because of their

cascading effects on later life stages (Baskin and Baskin 1971; Marks and Prince 1981;

Donohue 2002, 2005). In plant populations, environmental conditions at the time of germination

can alter the strength and direction of natural selection on postgermination traits (Donohue et al.

2010; D'Aguillo and Donohue 2023). In turn, this can affect the competitive environment,

resource availability, and density-dependent selective agents experienced by populations

(Donohue et al. 2010). While we did not investigate postgermination traits, it would be valuable

to determine whether germination timing influences performance at later life history stages (e.g.,

fecundity), especially in leading edge populations. Knowledge of germination timing could help

identify optimal windows for applying herbicides or biocontrol agents. If germination timing or

success is found to affect performance at later life stages, management strategies could aim to

destabilize these windows of opportunity, potentially preventing the establishment of new

invasive populations and limiting the growth and persistence of existing ones.

Our results suggest that even over the course of a fairly rapid invasion losses of

genomic variation may be minimal. In leafy spurge, this may have occurred because of

substantial gene flow during invasion and/or polyploidy. Higher levels of genetic variation can

challenge management when genotypes vary in their responses to eradication measures

(Gaskin et al. 2020). We also found that trait divergence may have contributed to climatic niche

expansion and thus to the spatial extent of invasion. Forecasts of continued invasion typically

rely on species distribution models (SDMs), which rarely take into account evolution. As such,

models may fail to predict the complete extent of range expansion, or the severity of range

infilling. Evolution-free SDMs are likely still valuable for management planning over meaningful

spatial and temporal scales in many systems. However, in systems where local adaptation is

extensive, forecasts of range shifts with climate change may require the construction of regional

SDMs that account for evolution.

Tables

Table 1. Locality information and genetic diversity metrics (observed heterozygosity, Ho; inbreeding coefficient, G_{is}; number of private alleles, and Tajima's D) for 14 leafy spurge population samples from four phases of range expansion (see Figure 1).

Range Position	Population Genetic Samples	Latitude	Longitude	Observed Heterozygosity (Ho)	Inbreeding Coefficient (Gis)	Private Alleles	Tajima's D
Core	Blue Earth	44.16	-94.09	0.068	0.123	27	-0.037
	Lyon	44.33	-95.82	0.067	-0.017	9	-0.020
	Meeker	44.94	-94.64	0.07	-0.017	4	0.205
Early Expansion	Becker	46.88	-96.05	0.068	0.116	10	-0.052
	Big Stone	45.52	-96.55	0.067	0.127	9	-0.028
	Polk	47.75	-96.25	0.07	0.097	11	-0.058
Late Expansion	Aitkin	46.98	-93.72	0.068	0.138	27	0.122
	Anoka	45.29	-93.13	0.071	-0.026	7	0.101
	Crow Wing	46.38	-94.22	0.065	0.081	8	0.001
	Winona	44.04	-91.62	0.071	0.156	42	-0.167
Invasion Front	Duluth	46.76	-92.11	0.066	0.12	23	-0.034
	Koochiching	48.60	-93.40	0.066	0.089	10	-0.008
	Pine	46.04	-92.36	0.069	0.138	21	0.000
	St. Louis	47.72	-91.97	0.065	0.046	6	0.068

656   Table 2. Pairwise estimates of genetic differentiation measured as G_{st} for 14 leafy spurge

657   populations from four phases of range expansion (see Figure 1). Darker versus lighter shading

658   of cells indicates higher versus lower values of pairwise G_{st}.

Phase	Population	Blue Earth	Lyon	Meeker	Becker	Big Stone	Polk	Aitkin	Anoka	Crow Wing	Winona	Duluth	Koochiching	Pine	St. Louis
Core	Blue Earth														
Core	Lyon	0.176													
Core	Meeker	0.215	0.266												
Early Expansion	Becker	0.126	0.173	0.208											
Early Expansion	Big Stone	0.125	0.171	0.209	0.112										
Early Expansion	Polk	0.118	0.171	0.205	0.115	0.119									
Late Expansion	Aitkin	0.124	0.187	0.216	0.126	0.127	0.127								
Late Expansion	Anoka	0.187	0.243	0.277	0.177	0.198	0.185	0.197							
Late Expansion	Crow Wing	0.134	0.188	0.219	0.116	0.128	0.126	0.136	0.201						
Late Expansion	Winona	0.084	0.134	0.181	0.081	0.084	0.081	0.095	0.153	0.096					
Invasion Front	Duluth	0.126	0.163	0.203	0.110	0.116	0.118	0.128	0.179	0.124	0.080				
Invasion Front	Koochiching	0.136	0.197	0.225	0.139	0.138	0.139	0.152	0.213	0.153	0.099	0.137			
Invasion Front	Pine	0.132	0.188	0.204	0.122	0.128	0.130	0.138	0.201	0.135	0.094	0.121	0.153		
Invasion Front	St. Louis	0.178	0.240	0.249	0.169	0.177	0.178	0.177	0.252	0.124	0.143	0.171	0.199	0.182	

665   Table 3. ANOVA testing for a relationship between four metrics of sequence diversity and

666   geography (latitude, longitude, and their interaction) for 14 population samples. Bold indicates p-

667   value less than 0.05.

Model	Heterozygosity (Ho)			Private Alleles			Tajima's D			Inbreeding (Gis)		
Term	Df	F	P	Df	F	P	Df	F	P	Df	F	P
Latitude	1	2.37	0.158	1	0.07	0.803	1	0.24	0.636	1	1.41	0.266
Longitude	1	0.48	0.506	1	1.29	0.285	1	1.22	0.298	1	0.04	0.846
Latitude x Longitude	1	9.49	**0.013**	1	0.84	0.382	1	0.02	0.887	1	0.48	0.508

Figures

Figure 1. A) Invasion history reproduced from Hanson & Rudd (1933). The map was used to delineate four phases of invasion: core, early expansion, late expansion, and invasion front. Tissue collection sites for B) 14 population samples and C) 157 landscape samples of leafy spurge.

Figure 2. Cluster assignment probability from STRUCTURE analyses (K = 3 and K = 4) for population and landscape samples. Each bar represents one individual, and populations are separated by black lines. A graph of the delta K values supports the optimal number of clusters at K = 3.

Figure 3. Principal components analysis (PCA) bi-plots for population genomic data from 14 population samples (n = 6 per population) and 157 landscape samples (n = 1 per population).

Figure 4. Isolation by distance (IBD) displayed as scatterplots of genetic distance versus geographic distance for A) population samples (A) and B-E) landscape samples (B-E). For landscape samples, we subset individuals for analyses by successive stages of invasion: B) range core (B), C) range core plus early expansion (C), D) range core, early, plus late expansion ranges (D), and E) all samples from across the four phases (E).

Figure 5. Niche overlap during range expansion in climate niche space. The extent of the background environment is outlined in black (solid = total niche space; dashed = 90% of extent). A) Core (dark pink) versus early expansion (light pink), B) Early expansion (light pink) versus late expansion (beige), C) Late expansion (beige) versus invasion front (grey), D) Core (dark pink) versus invasion front (grey). In all panels, the arrow represents the direction of shift in the centroid of niche space.

Figure 6. Ecological Niche Model (ENM) habitat suitability projections for seed source

populations and germination probability for source regions. A) Habitat suitability projection from

leafy spurge ENM. Predicted habitat suitability ranges from 0 (purple) to 1 (yellow). Dashed

lines demarcate the chronosequence of range expansion. Seed source populations for the

germination experiment are marked and colored yellow (>20°C) or purple (<20°C) depending

on their mean temperature of the warmest quarter. B) Variable response curve for the mean

temperature of the warmest quarter. Vertical yellow bars correspond to seed source populations

from the warmer south and vertical purple bars correspond to source populations from the

cooler north. C) Mean germination probability (± SE) by temperature treatment (X axis) for seed

source regions. The warmer southern region is shown in yellow and the cooler northern region

is shown in purple.

732   Figure 1.

746   Figure 2.

759   Figure 3.

775   Figure 4.

789   Figure 5.

802   Figure 6.

Seed Sources

Statements and Declarations

Funding: Funding for this project was provided by the Minnesota Invasive Terrestrial Plants and Pests Center through the Environment and Natural Resources Trust Fund as recommended by the Legislative-Citizen Commission on Minnesota Resources (LCCMR). TAL was supported by the University of Minnesota Doctoral Dissertation Fellowship.

Competing interests: The authors have no relevant financial or non-financial interests to disclose.

Author contributions: All authors conceived of the study design, contributed to decisions throughout the analysis process, wrote, and edited the manuscript. T.A.L and R.B.R collected and analyzed data.

Data Accessibility and Benefit-Sharing: All data will be made publicly available upon publication via the Dryad Digital Repository [DOI].

References

834 Araújo MB, Anderson RP, Márcia Barbosa A, et al (2019) Standards for distribution models in
835   biodiversity assessments. Sci Adv 5:eaat4858
836 Atwater DZ, Ervine C, Barney JN (2018) Climatic niche shifts are common in introduced plants.
837   Nat Ecol Evol 2:34–43
838 Austerlitz F, Jung-Muller B, Godelle B, Gouyon P-H (1997) Evolution of Coalescence Times,
839   Genetic Diversity and Structure during Colonization. Theor Popul Biol 51:148–164
840 Bakke AL (1936) Leafy spurge, Euphorbia esula L. Iowa Agricultural Experiment Station
841   189:209–245
842 Baniaga AE, Marx HE, Arrigo N, Barker MS (2020) Polyploid plants have faster rates of
843   multivariate niche differentiation than their diploid relatives. Ecol Lett 23:68–78
844 Baskin JM, Baskin CC (1971) Germination of Winter Annuals in July and Survival of the
845   Seedlings. Bull Torrey Bot Club 98:272–276
846 Bates OK, Bertelsmeier C (2021) Climatic niche shifts in introduced species. Curr Biol
847   31:R1252–R1266
848 Batho G (1932) Leafy spurge. Department of Agriculture and Immigration Winnipeg, Canada
849   106
850 Blaine Marchant D, Soltis DE, Soltis PS (2016) Patterns of abiotic niche shifts in allopolyploids
851   relative to their progenitors. New Phytol 212:708–718
852 Bolger AM, Lohse M, Usadel B (2014) Trimmomatic: a flexible trimmer for Illumina sequence
853   data. Bioinformatics 30:2114–2120
854 Brittingham HA, Koski MH, Ashman T-L (2018) Higher ploidy is associated with reduced range
855   breadth in the Potentilleae tribe. Am J Bot 105:700–710
856 Brochmann C, Brysting AK, Alsos IG, et al (2004) Polyploidy in arctic plants. Biol J Linn Soc
857   Lond 82:521–536
858 Capblancq T, Fitzpatrick MC, Bay RA, et al (2020) Genomic Prediction of (Mal)Adaptation
859   Across Current and Future Climatic Landscapes. Annu Rev Ecol Evol Syst 51:245–269
860 Chao WS, Horvath DP, Anderson JV, Foley ME (2005) Potential model weeds to study
861   genomics, ecology, and physiology in the 21st century. Weed Sci 53:929–937
862 Chapman DS, Scalone R, Štefanić E, Bullock JM (2017) Mechanistic species distribution
863   modeling reveals a niche shift during invasion. Ecology 98:1671–1680
864 Chardon NI, Pironon S, Peterson ML, Doak DF (2020) Incorporating intraspecific variation into
865   species distribution models improves distribution predictions, but cannot predict species
866   traits for a wide-spread plant species. Ecography 43:60–74
867 Chown SL, Hodgins KA, Griffin PC, et al (2015) Biological invasions, climate change and
868   genomics. Evol Appl 8:23–46
869 Clark LV, Lipka AE, Sacks EJ (2019) polyRAD: Genotype Calling with Uncertainty from
870   Sequencing Data in Polyploids and Diploids. G3 9:663–673
871 Clark LV, Mays W, Lipka AE, Sacks EJ (2022) A population-level statistic for assessing
872   Mendelian behavior of genotyping-by-sequencing data from highly duplicated genomes.
873   BMC Bioinformatics 23:101
874 Clements DR, Ditommaso A (2011) Climate change and weed adaptation: can evolution of
875   invasive plants lead to greater range expansion than forecasted? Weed Res 51:227–240
876 Clements DR, Jones VL (2021) Rapid evolution of invasive weeds under climate change:
877   Present evidence and future research needs. Front Agron 3:664034
878 Colautti RI, Barrett SCH (2013) Rapid Adaptation to Climate Facilitates Range Expansion of an
879   Invasive Plant. Science 342:364–366
880 Colautti RI, Lau JA (2015) Contemporary evolution during invasion: evidence for differentiation,
881   natural selection, and local adaptation. Mol Ecol 24:1999–2017
882 Collart F, Hedenäs L, Broennimann O, et al (2020) Intraspecific differentiation: Implications for
883   niche and distribution modelling. J Biogeogr 48:415–426

D'Aguillo M, Donohue K (2023) Changes in phenology can alter patterns of natural selection:
the joint evolution of germination time and postgermination traits. New Phytol.
https://doi.org/10.1111/nph.18711
Des Roches S, Turley NE, Bailey JK, et al (2017) The ecological importance of intraspecific
variation. Nature Ecology & Evolution 2:57–64
Di Cola V, Broennimann O, Petitpierre B, et al (2017) ecospat: an R package to support spatial
analyses and modeling of species niches and distributions. Ecography 40:774–787
Dixon AL, Busch JW (2017) Common garden test of range limits as predicted by a species
distribution model in the annual plant Mimulus bicolor. Am J Bot 104:817–827
Dlugosch KM, Parker IM (2008) Founding events in species invasions: genetic variation,
adaptive evolution, and the role of multiple introductions. Mol Ecol 17:431–449
Do C, Waples RS, Peel D, et al (2014) NeEstimator v2: re-implementation of software for the
estimation of contemporary effective population size (Ne) from genetic data. Mol Ecol
Resour 14:209–214
Donohue K (2003) Setting the Stage: Phenotypic Plasticity as Habitat Selection. Int J Plant Sci
164:S79–S92
Donohue K (2005) Niche construction through phenological plasticity: life history dynamics and
ecological consequences. New Phytol 166:83–92
Donohue K (2002) Germination timing influences natural selection on life-history characters in
Arabidopsis Thaliana. Ecology 83:1006–1016
Donohue K, Rubio de Casas R, Burghardt L, et al (2010) Germination, Postgermination
Adaptation, and Species Ecological Ranges. Annu Rev Ecol Evol Syst 41:293–319
Dufresne F, Stift M, Vergilino R, Mable BK (2014) Recent progress and challenges in population
genetics of polyploid organisms: an overview of current state-of-the-art molecular and
statistical tools. Mol Ecol 23:40–69
Duncan CA, Jachetta JJ, Brown ML, et al (2004) Assessing the Economic, Environmental, and
Societal Losses from Invasive Plants on Rangeland and Wildlands. Weed Technol
18:1411–1416
Dunn PH (1985) Origins of leafy spurge in North America. In: Watson AK (ed) Number 3 of the
monograph series of the weed science society of America. Champaign, IL, pp 7–13
Dunn PH (1979) The Distribution of Leafy Spurge (Euphorbia esula) and Other Weedy
Euphorbia spp. in the United States. Weed Sci 27:509–516
Earl DA, vonHoldt BM (2012) STRUCTURE HARVESTER: a website and program for
visualizing STRUCTURE output and implementing the Evanno method. Conserv Genet
Resour 4:359–361
Elith J, Leathwick JR (2009) Species Distribution Models: Ecological Explanation and Prediction
Across Space and Time. Annu Rev Ecol Evol Syst 40:677–697
Evanno G, Regnaut S, Goudet J (2005) Detecting the number of clusters of individuals using
the software STRUCTURE: a simulation study. Mol Ecol 14:2611–2620
Excoffier L (2004) Patterns of DNA sequence diversity and genetic structure after a range
expansion: lessons from the infinite-island model. Mol Ecol 13:853–864
Excoffier L, Ray N (2008) Surfing during population expansions promotes genetic revolutions
and structuration. Trends Ecol Evol 23:347–351
Excoffier L, Smouse PE, Quattro JM (1992) Analysis of molecular variance inferred from metric
distances among DNA haplotypes: application to human mitochondrial DNA restriction
data. Genetics 131:479–491
Fourcade Y, Besnard AG, Secondi J (2018) Paintings predict the distribution of species, or the
challenge of selecting environmental predictors and evaluation statistics. Glob Ecol
Biogeogr 27:245–256
Funk JL (2008) Differences in Plasticity between Invasive and Native Plants from a Low
Resource Environment. J Ecol 96:1162–1173

Gaskin JF, Espeland E, Johnson CD, et al (2020) Managing invasive plants on Great Plains
grasslands: A discussion of current challenges. Rangeland Ecol Manage.
https://doi.org/10.1016/j.rama.2020.04.003
Gaskin JF, Littlefield JL, Rand TA, West NM (2022) Variation in reproductive mode across the
latitudinal range of invasive Russian knapweed. AoB Plants 14:lac032
Gaskin JF, Schaal BA (2002) Hybrid *Tamarix* widespread in U.S. invasion and undetected in
native Asian range. Proc Natl Acad Sci U S A 99:11256–11259
Gerard D, Ferrão LFV (2020) Priors for genotyping polyploids. Bioinformatics 36:1795–1800
Gibson MJ, Torres M de L, Brandvain Y, Moyle LC (2021) Introgression shapes fruit color
convergence in invasive Galápagos tomato. Elife 10:.
https://doi.org/10.7554/eLife.64165
Glennon KL, Ritchie ME, Segraves KA (2014) Evidence for shared broad-scale climatic niches
of diploid and polyploid plants. Ecol Lett 17:574–582
Goodsman DW, Cooke B, Coltman DW, Lewis MA (2014) The genetic signature of rapid range
expansions: How dispersal, growth and invasion speed impact heterozygosity and allele
surfing. Theor Popul Biol 98:1–10
Gorton AJ, Tiffin P, Moeller DA (2019) Does adaptation to historical climate shape plant
responses to future rainfall patterns? A rainfall manipulation experiment with common
ragweed. Oecologia 190:941–953
Hanson HC, Rudd VE (1933) Leafy spurge: Life history and habits. North Dakota Agricultural
College Experiment Station Bulletin 1–24
Hein DG, Miller SD (1992) Influence of leafy spurge on forage utilization by cattle. J Range
Manage 45:405
Hijmans R, Phillips S, Leathwick J, Elith J (2022) dismo: Species Distribution Modeling. Version
1.3-9URL https://cran.r-project.org/web/packages/dismo/index.html
Hodgins KA, Bock DG, Rieseberg LH (2018) Trait Evolution in Invasive Species. Annual Plant
Reviews online 459–496
Horvath D, Anderson JV, Chao WS, et al (2015) Leafy Spurge Genomics: A Model Perennial
Weed to Investigate Development, Stress Responses, and Invasiveness. In: Sablok G,
Kumar S, Ueno S, et al. (eds) Advances in the Understanding of Biological Sciences
Using Next Generation Sequencing (NGS) Approaches. Springer International
Publishing, Cham, pp 63–78
Horvath DP, Patel S, Doğramaci M, et al (2018) Gene Space and Transcriptome Assemblies of
Leafy Spurge (Euphorbia esula) Identify Promoter Sequences, Repetitive Elements,
High-Quality Markers, and a Full-Length Chloroplast Genome. Weed Sci 66:355–367
Hutchison DW, Templeton AR (1999) Correlation of pairwise genetic and geographic distance
measures: inferring the relative influences of gene flow and drift on the distribution of
genetic variability. Evolution 53:1898–1914
Jombart T (2008) adegenet: a R package for the multivariate analysis of genetic markers.
Bioinformatics 24:1403–1405
Kamvar ZN, Brooks JC, Grünwald NJ (2015) Novel R tools for analysis of genome-wide
population genetic data with emphasis on clonality. Front Genet 6:208
Keller SR, Taylor DR (2010) Genomic admixture increases fitness during a biological invasion. J
Evol Biol 23:1720–1731
Keller SR, Taylor DR (2008) History, chance and adaptation during biological invasion:
separating stochastic phenotypic evolution from response to selection. Ecol Lett 11:852–
866
Klopfstein S, Currat M, Excoffier L (2006) The fate of mutations surfing on the wave of a range
expansion. Mol Biol Evol 23:482–490
Koski MH, Layman NC, Prior CJ, et al (2019) Selfing ability and drift load evolve with range
expansion. Evol Lett 3:500–512

Lacey JR, Wallander R, Olson-Rutz K (1992) Recovery, Germinability, and Viability of Leafy
Spurge (Euphorbia esula) Seeds Ingested by Sheep and Goats. Weed Technol 6:599–
602
Lachmuth S, Durka W, Schurr FM (2010) The making of a rapid plant invader: genetic diversity
and differentiation in the native and invaded range of Senecio inaequidens. Mol Ecol
19:3952–3967
Lake TA, Briscoe Runquist RD, Moeller DA (2020) Predicting range expansion of invasive
species: Pitfalls and best practices for obtaining biologically realistic projections. Divers
Distrib 24:348
Lakela O (1965) A Flora of Northeastern Minnesota, NED-New edition. University of Minnesota
Press
Lande R (2015) Evolution of phenotypic plasticity in colonizing species. Mol Ecol 24:2038–2045
Lee CE (2002) Evolutionary genetics of invasive species. Trends Ecol Evol 17:386–391
Lehe R, Hallatschek O, Peliti L (2012) The rate of beneficial mutations surfing on the wave of a
range expansion. PLoS Comput Biol 8:e1002447
Liu C, Wolter C, Xian W, Jeschke JM (2020) Most invasive species largely conserve their
climatic niche. Proc Natl Acad Sci USA 117:23643–23651
Lym RG (1998) The Biology and Integrated Management of Leafy Spurge (Euphorbia esula) on
North Dakota Rangeland. Weed Technol 12:367–373
Lym RG, Carlson RB (2002) Effect of Leafy Spurge (Euphorbia esula) Genotype on Feeding
Damage and Reproduction of Aphthona spp.: Implications for Biological Weed Control.
Biol Control 23:127–133
Lym RG, Kirby DR (1987) Cattle Foraging Behavior in Leafy Spurge (Euphorbia esula)-Infested
Rangeland. Weed Technol 1:314–318
Ma L, Cao L-J, Hoffmann AA, et al (2020) Rapid and strong population genetic differentiation
and genomic signatures of climatic adaptation in an invasive mealybug. Divers Distrib
26:610–622
Marks M, Prince S (1981) Influence of Germination Date on Survival and Fecundity in Wild
Lettuce Lactuca Serriola. Oikos 36:326–330
Mathias A, Kisdi E (2002) Adaptive diversification of germination strategies. Proc Biol Sci
269:151–155
Mattingly KZ, Hovick SM (2023) Autopolyploids of Arabidopsis thaliana are more phenotypically
plastic than their diploid progenitors. Ann Bot 131:45–58
Medley KA (2010) Niche shifts during the global invasion of the Asian tiger mosquito, Aedes
albopictus Skuse (Culicidae), revealed by reciprocal distribution models. Glob Ecol
Biogeogr 19:122–133
Meirmans PG (2020) genodive version 3.0: Easy-to-use software for the analysis of genetic
data of diploids and polyploids. Mol Ecol Resour 20:1126–1131
Meirmans PG (2012) AMOVA-based clustering of population genetic data. J Hered 103:744–
750
Meirmans PG, Liu S, van Tienderen PH (2018) The Analysis of Polyploid Genetic Data. J Hered
109:283–296
Melbourne BA, Hastings A (2009) Highly variable spread rates in replicated biological invasions:
fundamental limits to predictability. Science 325:1536–1539
Merow C, Smith MJ, Silander JA Jr (2013) A practical guide to MaxEnt for modeling species'
distributions: what it does, and why inputs and settings matter. Ecography 36:1058–1069
Moody ME, Mueller LD, Soltis DE (1993) Genetic variation and random drift in autotetraploid
populations. Genetics 134:649–657
Moran EV, Reid A, Levine JM (2017) Population genetics and adaptation to climate along
elevation gradients in invasive Solidago canadensis. PLoS One 12:e0185539

Morente-López J, Kass JM, Lara-Romero C, et al (2022) Linking ecological niche models and common garden experiments to predict phenotypic differentiation in stressful environments: Assessing the adaptive value of marginal populations in an alpine plant. Glob Chang Biol 28:4143–4162

Morrow AL (1979) Studies on the Reproductive Biology of Leafy Spurge (Euphoriba esula). Weed Sci 27:106–109

Nei M (1972) Genetic Distance between Populations. Am Nat 106:283–292

Nei M, Maruyama T, Chakraborty R (1975) THE BOTTLENECK EFFECT AND GENETIC VARIABILITY IN POPULATIONS. Evolution 29:1–10

Oduor AMO, Leimu R, van Kleunen M (2016) Invasive plant species are locally adapted just as frequently and at least as strongly as native plant species. J Ecol 104:957–968

Otto SP, Whitton J (2000) Polyploid incidence and evolution. Annu Rev Genet 34:401–437

Owen MDK, Zelaya IA (2005) Herbicide-resistant crops and weed resistance to herbicides. Pest Manag Sci 61:301–311

Pandit MK, Pocock MJO, Kunin WE (2011) Ploidy influences rarity and invasiveness in plants. J Ecol 99:1108–1115

Parmesan C (2006) Ecological and Evolutionary Responses to Recent Climate Change. Annu Rev Ecol Evol Syst 37:637–669

Parmesan C, Hanley ME (2015) Plants and climate change: complexities and surprises. Ann Bot 116:849–864

Peischl S, Dupanloup I, Kirkpatrick M, Excoffier L (2013) On the accumulation of deleterious mutations during range expansions. Mol Ecol 22:5972–5982

Pemberton RW (1988) Myrmecochory in the Introduced Range Weed, Leafy Spurge (Euphorbia esula L.). Am Midl Nat 119:431–435

Peter BM, Slatkin M (2013) Detecting range expansions from genetic data. Evolution 67:3274–3289

Petitpierre B, Broennimann O, Kueffer C, et al (2017) Selecting predictors to maximize the transferability of species distribution models: lessons from cross-continental plant invasions: Which predictors increase the transferability of SDMs? Glob Ecol Biogeogr 26:275–287

Petitpierre B, Kueffer C, Broennimann O, et al (2012) Climatic niche shifts are rare among terrestrial plant invaders. Science 335:1344–1348

Phillips SJ, Anderson RP, Dudík M, et al (2017) Opening the black box: an open-source release of Maxent. Ecography 40:887–893

Phillips SJ, Dudík M (2008) Modeling of species distributions with Maxent: new extensions and a comprehensive evaluation. Ecography 31:161–175

Pina-Martins F, Silva DN, Fino J, Paulo OS (2017) Structure_threader: An improved method for automation and parallelization of programs structure, fastStructure and MavericK on multicore CPU systems. Mol Ecol Resour 17:e268–e274

Prentis PJ, Wilson JRU, Dormontt EE, et al (2008) Adaptive evolution in invasive species. Trends Plant Sci 13:288–294

Pritchard JK, Stephens M, Donnelly P (2000) Inference of population structure using multilocus genotype data. Genetics 155:945–959

Reed DH, Frankham R (2001) How closely correlated are molecular and quantitative measures of genetic variation? A meta-analysis. Evolution 55:1095–1103

Rice A, Šmarda P, Novosolov M, et al (2019) The global biogeography of polyploid plants. Nature Ecology & Evolution 3:265–273

Richards CL, Bossdorf O, Muth NZ, et al (2006) Jack of all trades, master of some? On the role of phenotypic plasticity in plant invasions. Ecol Lett 9:981–993

Riina R, Peirson JA, Geltman DV, et al (2013) A worldwide molecular phylogeny and classification of the leafy spurges, Euphorbia subgenus Esula (Euphorbiaceae). Taxon 62:316–342

Rochette NC, Rivera-Colón AG, Catchen JM (2019) Stacks 2: Analytical methods for paired-end sequencing improve RADseq-based population genomics. Mol Ecol 28:4737–4754

Ronfort J, Jenczewski E, Bataillon T, Rousset F (1998) Analysis of population structure in autotetraploid species. Genetics 150:921–930

Rozas J, Ferrer-Mata A, Sánchez-DelBarrio JC, et al (2017) DnaSP 6: DNA Sequence Polymorphism Analysis of Large Data Sets. Mol Biol Evol 34:3299–3302

Rutland CA, Hall ND, McElroy JS (2021) The Impact of Polyploidization on the Evolution of Weed Species: Historical Understanding and Current Limitations. Frontiers in Agronomy 3:626454

Sakai AK, Allendorf FW, Holt JS, et al (2001) The Population Biology of Invasive Specie. Annu Rev Ecol Syst 32:305–332

Schulz-Schaeffer, Gerhardt (1989) Cytotaxonomic analysis of the euphorbia spp leafy spurge complex ii. comparative study of the chromosome morphology. Biol Zent Bl 108:69–76

Searcy CA, Shaffer HB (2016) Do Ecological Niche Models Accurately Identify Climatic Determinants of Species Ranges? Am Nat 187:423–435

Selechnik D, Richardson MF, Shine R, et al (2019) Increased Adaptive Variation Despite Reduced Overall Genetic Diversity in a Rapidly Adapting Invader. Front Genet 10:1221

Selleck GW, Coupland RT, Frankton C (1962) Leafy spurge in Saskatchewan. Ecol Monogr 32:1–29

Sexton JP, McKay JK, Sala A (2002) Plasticity and Genetic Diversity May Allow Saltcedar to Invade Cold Climates in North America. Ecol Appl 12:1652–1660

Singmann H, Bolker B, Westfall J, F. A (2016) afex: Analysis of Factorial Experiments. Version 0.16-1URL https://CRAN.R-project.org/package=afex

Slatkin M (1987) Gene flow and the geographic structure of natural populations. Science 236:787–792

Slatkin M (1993) Isolation by Distance in Equilibrium and Non-Equilibrium Populations. Evolution 47:264–279

Slatkin M, Excoffier L (2012) Serial founder effects during range expansion: a spatial analog of genetic drift. Genetics 191:171–181

Soltis PS, Soltis DE (2000) The role of genetic and genomic attributes in the success of polyploids. Proc Natl Acad Sci U S A 97:7051–7057

Suarez AV, Tsutsui ND (2008) The evolutionary consequences of biological invasions. Mol Ecol 17:351–360

Sullivan LL, Li B, Miller TEX, et al (2017) Density dependence in demography and dispersal generates fluctuating invasion speeds. Proc Natl Acad Sci USA 114:5053–5058

Tabassum S, Leishman MR (2018) Have your cake and eat it too: greater dispersal ability and faster germination towards range edges of an invasive plant species in eastern Australia. Biol Invasions 20:1199–1210

Tajima F (1989) Statistical method for testing the neutral mutation hypothesis by DNA polymorphism. Genetics 123:585–595

te Beest M, Le Roux JJ, Richardson DM, et al (2012) The more the better? The role of polyploidy in facilitating plant invasions. Ann Bot 109:19–45

Travis JMJ, Mynard P, Bocedi G (2021) Evolution of reduced dormancy during range expansions. bioRxiv 2021.10.11.463894

Treier UA, Broennimann O, Normand S, et al (2009) Shift in cytotype frequency and niche space in the invasive plant Centaurea maculosa. Ecology 90:1366–1377

Uller T, Leimu R (2011) Founder events predict changes in genetic diversity during human-mediated range expansions. Glob Chang Biol 17:3478–3485

van Boheemen LA, Atwater DZ, Hodgins KA (2019) Rapid and repeated local adaptation to climate in an invasive plant. New Phytol 222:614–627

Van de Peer Y, Ashman T-L, Soltis PS, Soltis DE (2021) Polyploidy: an evolutionary and ecological force in stressful times. Plant Cell 33:11–26

Verma P, Majee M (2013) Seed germination and viability test in tetrazolium (TZ) assay. Bio Protoc 3:1–4

Wang D, Xu X, Zhang H, et al (2022) Abiotic Niche Divergence of Hybrid Species from Their Progenitors. Am Nat 200:634–645

Warren DL, Glor RE, Turelli M (2008) Environmental niche equivalency versus conservatism: quantitative approaches to niche evolution. Evolution 62:2868–2883

Welles SR, Dlugosch KM (2019) Population Genomics of Colonization and Invasion. In: Rajora OP (ed) Population Genomics: Concepts, Approaches and Applications. Springer International Publishing, Cham, pp 655–683

West NM, Gaskin JF, Milan J, Rand TA (2023) High genetic diversity in the landscape suggests frequent seedling recruitment by Euphorbia virgata Waldst. & Kit. (leafy spurge) in the northern U.S.A. Biol Invasions 25:645–652

Westerband AC, Funk JL, Barton KE (2021) Intraspecific trait variation in plants: a renewed focus on its role in ecological processes. Ann Bot 127:397–410

Wicks GA, Derscheid LA (1958) Leafy spurge seed maturation studies. Proc North Cent Weed Control Conf

Williams JL, Kendall BE, Levine JM (2016) Rapid evolution accelerates plant population spread in fragmented experimental landscapes. Science 353:482–485

Wolkovich EM, Davies TJ, Schaefer H, et al (2013) Temperature-dependent shifts in phenology contribute to the success of exotic species with climate change. Am J Bot 100:1407–1421

Woods EC, Sultan SE (2022) Post-introduction evolution of a rapid life-history strategy in a newly invasive plant. Ecology 103:e3803

Wright S (1943) Isolation by Distance. Genetics 28:114–138