Incorporating environmental heterogeneity and observation effort to predict host

distribution and viral spillover from a bat reservoir

Authors: Rita Ribeiro[1], Jason Matthiopoulos[1*], Finn Lindgren[2], Carlos Tello[3,4], Carlos M.

Zariquiey[3], William Valderrama[3,5], Tonie E. Rocke[6], Daniel G. Streicker[1,7*]

*These authors share senior authorship

Affiliations:

[1]School of Biodiversity, One Health and Veterinary Medicine, College of Medical,

Veterinary and Life Sciences, University of Glasgow, Glasgow, UK

[2]School of Mathematics, University of Edinburgh, Edinburgh, UK

[3]ILLARIY (Asociación para el Desarrollo y Conservación de los Recursos Naturales), Lima,

Perú

[4]Yunkawasi, Lima, Perú

[5]Facultad de Medicina Veterinaria y Zootecnia, Universidad Peruana Cayetano Heredia,

Lima, Perú

[6]U.S. Geological Survey, National Wildlife Health Center, Madison, Wisconsin, USA

[7]Medical Research Council - University of Glasgow Centre for Virus Research, Glasgow, UK

R.R., 0000-0002-9976-3731; J.M., 0000-0003-3639-8172; F.L., 0000-0002-5833-2011;

T.E.R., 0000-0003-3933-1563; D.G.S., 0000-0001-7475-2705

Corresponding author: Rita Ribeiro, School of Biodiversity One Health and Veterinary

Medicine, University of Glasgow, University Avenue, Graham Kerr Building, Glasgow G12

8QQ, UK. Email: ritaclaudia.cardosoribeiro@glasgow.ac.uk

Abstract

Predicting the spatial occurrence of wildlife is a major challenge for ecology and management. In Latin America, knowledge of the true number and locations of vampire bat colonies precludes informed allocation of measures intended to limit lethal rabies spillover to humans and livestock. We inferred the complete spatial distribution of vampire bat roosts while accounting for observation effort and environmental covariates by fitting a Log Gaussian Cox Process model to the locations of 563 roosts in three regions of Peru. Our model explained 47% of the variance in the observed roost distribution and identified landscape correlates of roost establishment. Our model estimated that 1,795 roosts (76%) remain undiscovered and identified hotspots of undetected roosts in currently rabies-free areas, implying high risk for viral incursion. Incorporating the locations of undetected roosts improved spatial predictions of rabies spillover to livestock, revealed areas with disproportionate underreporting to surveillance systems, and indicated a higher rabies burden than previously estimated. We provide a robust approach to infer the distribution of a mostly unobserved bat reservoir that can inform strategies to prevent the re-emergence of an important zoonosis.

Keywords: detection probability, Gaussian random field, partially observed species, species distribution models, vampire bat rabies

Introduction

Mapping the geographic distribution of animal reservoirs of infection is a pre-requisite to anticipate and prevent spillover to other species [1]. For example,

vaccination campaigns for dog rabies use detailed surveys of dog density to allocate vaccine

distributions, and control programmes of bovine, sheep, and goat brucellosis require livestock

census for adequate implementation of testing, vaccination, and slaughter strategies [2–4].

However, for wild or free-ranging species, distribution data are frequently sparse, presence-

only records that do not cover the full geographic range [5,6]. The problem is exacerbated for

species such as bats and rodents that are key reservoirs for zoonoses, but are difficult to

observe given their small body size, reclusive nature, nocturnal activity, or high mobility

[5,7,8]. Species distribution models (SDMs) that use relationships between field observations

and environmental data to predict the occurrence or abundance of disease vectors or

reservoirs are widely used to anticipate disease spread and help policymakers target

prevention and control strategies [9]. More recent extensions of SDMs can deal with

observation bias in input data and explicitly account for autocorrelation arising from

geographic proximity between observations, hence capturing residual autocorrelation in the

data, but have only rarely been applied to wildlife disease reservoirs. Accounting for such

inferential challenges is vital to avoid misguided spatial allocations of interventions

[5,6,10,11].

One of the most important zoonotic viruses transmitted by bats is rabies

(*Rhabdoviridae lyssavirus*), which is capable to cause acute lethal encephalitis in all

mammals [12]. In Latin America, the common vampire bat (*Desmodus rotundus*) is the main

reservoir of rabies due to its population abundance, wide geographic distribution and obligate

blood feeding, which provides a direct route for human and livestock infection via

contaminated saliva [13,14]. Efforts to mitigate the burden of vampire bat rabies (VBR)

include human and livestock vaccination; however, financial and logistical challenges result

in campaigns that are largely reactive to rabies mortality events, and costs in human lives and

livestock persist [15–17]. VBR management also involves controlling vampire bat

populations using topical anticoagulant poisons that spread between individual bats

during social grooming or that are applied to cattle for later consumption by bats

[18,19]. The ability of bat culling to reduce rabies incidence is controversial, and

efficacy has been questioned based on field, phylogenetic, and modelling studies

[15,20–22]. Further, countries with active culling campaigns have seen viral spatial

expansions and growing disease burdens [23,24]. It is hypothesized that culls may be

ineffective because rabies virus is maintained by spatial processes, including wave-

like invasions into historically rabies-free areas and metapopulation maintenance

among patchworks of bat colonies, but culls are reactive to rabies outbreaks and rarely

synchronized across enzootic regions [21,23]. Improving the implementation of culls

or the spatial distribution of human and livestock vaccines requires knowledge of the

spatial distribution of vampire bat populations and how this distribution determines

spillover risk; however, with rare exceptions, neither the location nor the number of

vampire bat roosts are known completely [25].

Previous models have predicted the distribution of *D. rotundus* over large

geographic scales (e.g., across countries) and have explored how climate change

might affect the future distribution of this species [26–28]. Predictions from these

models are necessarily coarse given the wide variety of habitat types considered. In

contrast, informing on-the-ground management requires high resolution predictions

that are robust to spatial autocorrelation and that explicitly consider heterogeneities in

the search effort applied to detect individuals. Here, we used data on the locations of

D. rotundus roosts (any place a wild bat uses for shelter or protection) to predict the

distribution and expected density of detected and undetected roosts while accounting

for spatial correlation structures and heterogeneities in observation effort that

generated the underlying roost data. Roost data were collected in three regions in

southern Peru where vampire bat rabies virus (VBRV) is either endemic or invading

historically uninfected zones. Our specific goals were to (1) identify the

environmental and anthropogenic variables that drive roost distribution; (2) reconstruct the

map of expected *D. rotundus* roost density, controlling for heterogeneous effort; (3) estimate

the total number and spatial variation in undetected roosts; (4) test whether our predictions of

bat roost density improved explanations of the locations and intensity of VBR outbreaks in

livestock; and (5) use the inferred projection of VBR outbreaks to re-assess the burden of

rabies spillover.

Materials and methods

1.   Study area

The study area comprised three regions in southern Peru: Ayacucho, Apurimac, and

Cusco (hereafter 'AAC', joint area = 136,697 km^2, Fig. 1). AAC is largely composed of

inter-Andean valleys, with elevations ranging from ~280 to ~6,000 m above sea level. To

avoid modelling high-elevation areas that are known to exceed the physiological tolerance of

vampire bats, we excluded areas above 4,000 m from the study area (Fig. 1B) [23]. The

smoothr package in R was used to smooth the resulting landscape, which contained complex

boundaries with small holes and sharp corners [29].

2.   Data on vampire bat roosts

We used the geolocations of 563 *D. rotundus* roosts in the AAC, collected between

2007 and 2021 (Fig. 1C). Roosts included natural (68%) and human-made structures (32%)

but excluded foraging locations. Most natural structures inhabited by bats were caves (90%)

with remaining classified as trees (5%) or 'other' (5%). Roosts in human-made structures

were abandoned houses (47%), mines (24%), tunnels (15%), and 'other' (14%). Data

originated from three sources: internally from the research group (2.4% of

observations), from the Regional Government of Apurimac (22%, derived from a 2-

year rabies control campaign [2014-2016], which included active searching for

roosts), and from the National Service of Agrarian Health (SENASA, 75.6%). We

analysed unique roost locations from these three partially overlapping datasets.

Apurimac had the highest roost detection effort because searches were carried out by

both national and regional governments; moreover, our field studies in Apurimac

initiated in 2007 (versus 2014 in Cusco and Ayacucho). Although national data from

SENASA are more spatially extensive and represent information accumulated over

the years, regional data were compiled using similar methods and may have been

collected by the same individuals in some cases. Coordinates of roost locations were

projected in Universal Transverse Mercator (UTM) and measured as northings and

eastings in kilometers.

3.   Environmental and anthropogenic variables

We selected environmental variables hypothesized to affect *D. rotundus* roost

establishment, including (1) climate (annual mean temperature and annual mean

precipitation); (2) topography (elevation, slope, and terrain rugosity index); (3)

landcover (percentages of crop cover, tree cover, and permanent water); and (4)

characteristics related to bat foraging (distance to the nearest river, prey density [a

layer of cattle density and a layer of all livestock species density], number of rural

population settlements per 2 km^2 and human footprint). Table S1 (SI Appendix)

provides the source and initial resolution of all variables. Data were collated as raster

maps, set with the same extent (AAC area) and projection (UTM) (SI Appendix Fig.

S1). Variables expected to act at a local scale such as climate or land cover, were

rescaled to the finest resolution of all the variables (~100 m). Variables expected to influence

roost density at a broader scale (i.e., bat foraging) were used at their original resolution (1 or

2 km) or, in case of prey density, rescaled at 5 km resolution (the original resolution was 10

km). Prior to model fitting, variables were standardised (shifted to zero mean and scaled to

unit variance) and the correlation between variables was assessed. To decrease the correlation

between elevation (0-4,000 m) and other environmental variables, the layer of continuous

elevation was reclassified as a binary variable defining suitable and less suitable altitudes for

D. rotundus considering the threshold of 3,600 m indicated by previous research on the

distribution of VBR in the AAC [23,30].

4.   **Modelling approach**

To explain the heterogeneous and partially observed process of *D. rotundus*

distribution, roost locations were treated as an inhomogeneous point pattern that follows a log

Gaussian Cox process (LGCP) with intensity $\lambda(s)$ at coordinates s. An LGCP is double-

stochastic, because it is a hierarchical combination of a Poisson process at the first level

(where the locations of the points are conditionally independent) and a Gaussian random field

(GRF) at the second level (to account for spatial correlation in the data) [31,32]. We selected

an LGCP to take advantage of point locations to explain the intensity process that creates the

observed roost distribution.

Gaussian random fields are spatially continuous structured random processes that

typically have dense covariance matrices that are computationally demanding, requiring long

run times. An efficient way to introduce these structures in the model is to approximate the

continuously indexed GRF by a spatially tessellated approximation to a Stochastic Partial

Differential Equation (SPDE) [32,33]. Our model framework was developed using the

Integrated nested Laplace approximation (INLA) joint with the SPDE approach, implemented

via the inlabru R package [34]. Model fitting and inference were carried out in R

version 4.1.2, using the packages R-INLA version 22.04.09 and inlabru development

version 2.6.0.9003 [34–36]. We developed two main models that we present below: a

detectability model to account for uneven observation effort in the study area, and the

model of expected roost density, which was informed by the detectability model.

- **Detectability model**

To ensure the patterns identified by our model were not observation artefacts

and to robustly estimate the total number of roosts, it was necessary to correct for the

uneven observation effort. Because data on the number of person-hours spent

searching for roosts was unknown, we used the spatial accessibility of each location to

approximate search effort. We approached this as a pre-analysis step, akin to some

distance sampling analyses that fit a detection function separately from the ensuing

habitat modelling [37]. Specifically, the detectability model described roost detections

as a flexible function of accessibility to the main source of roost detection, measured

as the mean travel time separating each grid cell from any of the 7 SENASA offices

(SI Appendix, Table S1) [38]. To distinguish between regions that had different

overall effort and, to include the effort from the Regional Government of Apurimac,

we defined region as a two-level factor covariate separating Apurimac (highest effort)

from Ayacucho and Cusco (combined). The observation model is defined by the

expression:

$$\lambda_o(s) = \exp\left(\beta_0 + f(z) + \beta_1 x_{Apurimac}\right)$$

where $\lambda_o(s)$ is aimed to capture detection probability at location s, β_0 is the intercept term,

$f(z)$ is the non-linear effect of the accessibility layer modelled using a random walk model of

order 1 latent effect, and $\beta_1 x_{Apurimac}$ represents the effect of region x (Apurimac is the

reference). The predicted posterior mean derived from the observation model was normalised

by assigning the probability 1 at the origin, which implies that a roost at the location of the

SENASA offices is guaranteed to be known, resulting in the spatial probability of roost

detection.

-    **Matérn correlation model**

The Matérn correlation model explains the scale of spatial dependency between

neighbouring locations. This correlation structure is expressed over a discretisation of space

known as the mesh (SI Appendix Fig. S2), a finite grid of triangulations of the spatial domain

that approximates smooth random effects within the model [39]. Here, the mesh was defined

within the smoothed boundaries of the inter-Andean valleys of the AAC area (the domain),

and the resolution of the mesh was driven by the observed roost locations (hence directing

more modelling detail in areas with detections). The correlation structure was defined

according to the following equation:

$$cov(i,j) = \sigma^2 Matérn(d_{ij}, k)$$

where the covariance between any two locations depends on their distance (*d*), on the range

(*k*) of the Matérn function and on the spatial variance (σ^2). This model included Penalised

Complexity-priors (PC-priors) for range (*k*, practical range) and sigma (σ, the marginal

standard deviation) [40].

-    **Roost model**

The expected density of vampire bat roosts ($\lambda(s)$) was fitted as a function of an

intercept (β_0), variables for roost distribution ($X(s)$), a spatial random field ($f(s)$), and a

spatial offset (ε). The offset is the probability of roost detection from the detectability model,

to adjust for the uneven observation effort. The roost model is defined according to

the following expression:

$$\lambda(s) = exp(\beta_0 + \beta X(s) + f(s) + \varepsilon).$$

We used a forward addition procedure of variable selection, based on model

conditional predictive ordinate (CPO), deviance information criteria (DIC), and

Watanabe-Akaike information criteria (WAIC) to identify the most parsimonious

model. If two variables were correlated (Pearson's correlation, $r > 0.6$), the variable

that better explained the response was retained, while the other was excluded from the

model. To explore competition between bat colonies, as this behaviour is sometimes

observed in other colonial species (e.g., seabirds), we considered adding the

possibility of repulsion between roosts; however, the pairwise Euclidean distances

between roosts demonstrated no evidence of such an effect (SI Appendix Fig. S3)

[41].

-     **Spatial predictions and estimates of abundance**

To predict counts of roosts from the LGCP (i.e., an LGCP originates

predictions in continuous space), the mesh was converted into 8-km^2 grid cells, a

dimension based on mesh's resolution. We then used the fitted LGCP model to

predict the number of roosts in each grid cell, generating samples from the posterior

of all model parameters. To integrate the SPDE effect into roost count prediction, we

projected integration weights to the mesh nodes and replaced the integral with a

weighted sum in each grid cell. We predicted both the detected roosts (not correcting

for the uneven effort) and the total expected roosts, in the latter case setting

observation effort in all cells to the maximum such that observation probability was

uniform. The posterior distribution of undetected roosts was calculated by subtracting

the number of observed roosts (i.e., 563) from the density of the expected total count. We

then estimated the posterior distribution for the total count that encompassed systematic

stochasticity and uncertainty in parameter estimates, with and without adjusting for uneven

effort.

-    **Model validation**

We validated predictions of roost expected density within our study area by refitting

the model to a randomly selected 70% of the grid cells and reserving points falling inside the

remaining 30% of the cells as a validation set (SI Appendix Fig. S4). We computed the

distance between predicted and observed number of roosts in the validation set as a pseudo-

R^2 for count data [42]. Although discretising space into a regular grid required some

subjectivity in choosing the size of the discretised units, the grid size was defined based on

the mesh.

5.   Explaining the distribution of past VBRV outbreaks

We hypothesized that our predicted roost distribution would improve understanding

of the distribution of past livestock rabies outbreaks (i.e., defined as a single spillover from

bats to livestock), identify high risk areas for future outbreaks, and identify locations with

excess underreporting of outbreaks. For this purpose, we fitted a generalised linear model to

the number of VBRV outbreaks that occurred between 2003 and 2021 (1,212 outbreaks, Fig.

1C and Fig. S5). We used a negative binomial likelihood to account for the overdispersion in

the counts of rabies outbreaks in each 8-km^2 grid cell. The model included an offset

describing the probability of reporting rabies outbreaks in each grid cell due to accessibility

to the SENASA offices [17]. We compared the performance of four alternative measures of

bat distribution: (1) the observed number of roosts in each grid cell (raw data); (2) a kernel

smooth of the raw data to represent a naive interpolation of bat density based on

observed data without covariates or bias correction; (3) the predictions of our LGCP

model corrected for the uneven effort; and (4) a smooth of these predictions

(hereafter, Bat Utilization Distribution [BUD]), using the same parameters as the

smooth for raw roost data. In addition to one of the four measures of bat distribution,

we included the number of months since the first rabies outbreak occurred in each

district to account for the geographic expansion of VBRV across AAC. We also

included cattle density, reasoning that rabies would be more detectable in areas with

more cattle [13,23,43]. We assessed the goodness of fit of eight competing models

using Akaike information criterion (AIC) and model deviance pseudo-R^2. We used

the best model to predict the spatial distribution of past rabies outbreaks with and

without correcting for accessibility to reporting centres. To assess the burden of rabies

spillover, we estimated the underreporting factor as the ratio between predicted rabies

outbreaks (corrected by uneven accessibility) and the reported rabies outbreaks in

each grid cell. This analysis was restricted to grid cells that had at least one reported

rabies outbreak to exclude predicted outbreaks in areas that as of 2021, remained

rabies free (i.e., areas with high future risk).

Results

–   **Detectability and landscape correlates of vampire bat roost**

establishment

The detectability model showed that roost observation decayed in a sigmoid

shape with increasing travel time to SENASA offices. The inflection point of the

sigmoid indicated high roost detectability up to 55 minutes of travel, with detection

probability approaching zero for roosts that were more than 7 h from any SENASA

office (Fig. 2A). The probability of detecting a roost increased with landscape accessibility

and in areas that had more effort, particularly in Apurimac (Fig. 3A). Consistent with

expectations for bat home range, the correlation structure of the Matérn model showed strong

clustering of roosts up to distances of 10 km (half scale dependency) and positive correlations

up to 30 km (Fig. 2B).

D. rotundus roost density was strongly influenced by climate, increasing with mean

temperature until an optimum of 16.4 °C (Fig. S6) and decreasing in areas with higher

average precipitation (Fig. 2C). Roost density also increased in proximity to rural human

settlements and in areas with high crop cover, possibly both reflecting the availability of

livestock prey. Terrain topography had marginal but non-significant effects on roost density

with more roosts in areas with optimal elevation (<3,600m) and lower slopes (Fig. 2C).

-        **Distribution and density of *D. rotundus* roosts**

The spatial projection of expected roost density resembled the observed roost

distribution (pseudo-R^2 = 0.60; Fig. 3B) and predicted a similar number of roosts as those

used in model fitting (observed: 563 roosts; predicted: mean = 579, credible interval [CI]:

529-628; Fig. S7A). Pseudo-R^2 decreased to 0.36 when removing the spatial random field,

indicating that the GRF captured variability in roost density not explained by the

environmental covariates. In our 30% hold out validation data, the full model with

environmental covariates and the spatial random field explained 47% of the variance in roost

density. Correcting for uneven observation effort revealed a total of 2,358 roosts (CI: 2,136-

2,684), implying that 1,795 roosts (76%) remain undiscovered (Fig. S7A). Importantly, roost

detectability varied considerably over space, with as few as 4% of roosts known in some

areas. Further, undiscovered roosts were spatially clustered, with hotspots identified in

western and northern parts of our study area (Fig. 3C-D). The predictions of the posterior

distribution for the total count of roosts had higher uncertainty when adjusting for

effort, as it predicts the total abundance expected in the AAC area, accounting for

observed and missing roosts (Fig. S7B-C).

-       **Explaining the distribution of past VBRV outbreaks**

In univariate models of the alternative descriptions of the vampire bat

distribution, the smoothed predicted roost distribution (BUD) was the strongest

predictor of rabies outbreaks, explaining 22% of the distribution of past rabies

outbreaks (Table 1). When adding the additional covariates describing the geographic

expansion of VBRV across AAC and cattle density to the BUD model, explanatory

power increased by a further 15% (Full model pseudo-$R^2 = 0.37$). Alternate models

using other versions of the bat distribution and the additional covariates had similar

pseudo-R^2 (0.31–0.36), but consistently higher AIC (ΔAIC=128.4–29.6, Table 1).

Spatial predictions of rabies outbreaks from the best model (model 1), generated with

and without correcting for accessibility to reporting offices (Fig. 4A-B), revealed

strong spatial heterogeneity, signaling probable hotspots of unreported rabies spillover

across our study area or high-risk areas where future outbreaks might go unreported.

Comparison of predicted to observed rabies outbreaks showed that, on average, the

number of rabies cases in livestock may have been 7.1 times (95% CI: 3.8–10.5)

higher than officially reported (Fig. 4C).

Discussion

Understanding the spatial distributions of wildlife reservoirs of zoonoses is

vital to anticipate and prevent spillover but undermined by incomplete, biased, and

spatially autocorrelated observations of animal locations. Focusing on an important

bat reservoir in Latin America, our study provides a transferrable statistical approach to infer

the spatial distribution of partially observed wildlife [7]. In our study, spatial heterogeneity in

vampire bat colonies was associated with topographic, ecological, and anthropogenic factors

but also strongly affected by previously unrecognised observation biases. Indeed, our results

indicate that only 24% of vampire bat roosts are known to authorities and models correcting

these biases revealed unrecognised hotspots in vampire bat distribution. Finally, predictions

of bat density improved models of the locations and intensity of historical rabies outbreaks,

revealed spatial heterogeneity in the reporting of rabies outbreaks, and increased the

projected burden of rabies mortality. By identifying spatial hotspots of bat roosts and

potential future rabies outbreaks, our results may improve the allocation of limited resources

for vampire bat management and rabies prevention.

Our model indicated that temperature and precipitation limit the formation of vampire

bat roosts, corroborating results from earlier analyses conducted across the geographic range

of this species [26–28]. Although roosts of *D. rotundus* require more than 45% humidity, the

mechanism by which precipitation acts on vampire bat distribution is poorly understood [44].

The negative effect of precipitation in our model is most likely explained by the increase in

observed bat roosts from the rainy north to the dry climate in the centre and south of the

valleys [45]. Temperature effects were expected as *D. rotundus* is a poor homeotherm and

environmental temperatures below 10°C or above 37°C are not tolerable [46,47]. Of note,

however, the optimal environmental mean temperature found in our study (i.e., 16.4°C) is

relatively low compared with the recommended temperature for maintaining vampire bats in

captivity, between 21°C and 27°C [48]. It is possible that this difference reflects our use of

ambient temperature in our model, rather than within roost temperature measurements, which

were not measured in our study. Vampire bat roosts would generally be expected to have

more stable and lower temperatures than ambient, particularly in daytime hours when bats

occupy roosts, but this relationship may vary depending on local conditions and roosts

characteristics. Further, AAC valleys represent a range limit for this species in southern Peru

with higher elevation areas farther south experiencing intolerable conditions.

Assuming the recommended captive maintenance temperatures are valid, our results

indicate that in the AAC area, vampire bats exist in sub-optimal temperatures. This

implies the possibility of physiological trade-offs that influence the susceptibility to or

tolerance of vampire bats to infection. For example, female tree swallows

(*Tachycineta bicolor*) persisting in suboptimal sites in Alaska had evidence of

impaired immunity and populations of whistling tree frogs (*Litoria verreauxii*

verreauxii) in low quality habitats were less likely to persist through epizootics of

chytridiomycosis [49,50]. For vampire bats, heightened susceptibility to lethal rabies

infection is predicted to have large dynamical consequences on viral prevalence and

spillover [21]. If impaired immunity in these populations is verified, we speculate this

might contribute to the disproportionate burden of rabies in the AAC, which despite

its small size, accounts for most rabies outbreaks at the national level [23]. It is also

conceivable that the high genetic divergence of AAC vampire bats from those in

tropical areas included physiological adaptations to low temperatures which may have

unpredictable trade-offs with immunity [30,51,52].

We also found effects that are consistent with the influence of human activities

on vampire bat roost establishment. For example, the negative effect of slope may

reflect use of artificial (human-made) roosts, which are generally constructed in flatter

areas. The positive effects of the number of rural human settlements and crop cover

on vampire bat roost establishment likely reflect the availability of livestock, which

are the main food source for *D. rotundus* [43,53,54]. Although it is unexpected that

livestock itself was not retained by the model, we suggest that the global layers used,

while suitable to access broad scale effects of livestock, are less representative of variation in

livestock density at fine spatial scales than the more finely resolved measures of human

presence [43]. As such, our findings support the hypothesis that anthropogenic activities

favour vampire bat roost establishment and may facilitate spatial expansions of this species to

new areas within climatically tolerable regions [30].

Despite the influence of environmental covariates in explaining vampire bat roost

density, the higher pseudo-R^2 after adding the GRF (pseudo-R^2 = 0.36 versus 0.60) indicated

that the spatial dependency among roosts explained considerable variation in roost density.

For example, in cranes and seabirds, such dependencies may arise from dispersal limitations,

inter-specific competition, disturbance, or social interactions [10,11,31]. Here, social

interactions may be particularly important to explain spatial dependency at the roost level, as

vampire bats are social species and often use multiple roosts, mostly within small areas (2 to

3 km radius) [55,56]. In the AAC, correlation between roosts remained positive up to 30 km

(Fig. 2B). The spatial extent of this clustering is somewhat unexpected given vampire bat

home ranges are believed to extend only 5–10 km. It is conceivable that longer distance

connectivity – occasionally reported – may be more common than previously recognised,

which would have important implications for rabies spread [44,47,53,57]. Our results also

corroborate the absence of competition between vampire bat roosts, which would have been

expected to generate a negative correlation at small distances [31]. More generally, our

results illustrate the value of accessing the shape of the Matérn correlation range to

understand and generate hypotheses about animal ecology and social behaviour.

Besides spatial dependency, observation biases are expected in most natural habitats

but have not historically been incorporated into distribution modelling for wildlife. Our

observation model captured a decay in vampire bat roost detection as accessibility to

reporting offices decreased (Fig. 2A) and a sharp decrease in observed roosts in Ayacucho

and Cusco (Fig. 3A). When correcting predictions for the uneven effort, our model

estimated that the number of undetected roosts was more than triple the number of

known roosts. Moreover, missed roosts were spatially clustered, rather than uniformly

spread across the study area. Of note, some hotspots of missing roosts are currently in

rabies free areas that have neighbour areas with current viral circulation, indicating

high risk of rabies spread into areas where outbreaks in bats and thus livestock might

rapidly ensue [23]. This highlights an important outcome of incorporating observation

processes into model projections, enabling interventions such as preventive bat culls

or livestock vaccination that might prevent spillover or reduce its burden in high-risk

areas. More generally, improved understanding of the true spatial distribution of

vampire bats facilitates spatially synchronized control, which is predicted to be central

to successfully managing rabies in endemic areas, but until now, it was precluded by

gaps in our knowledge of bat distribution [21,43,58].

In addition to identifying areas at high risk of rabies outbreaks, our predictions

of detected and undetected roosts improved understanding of the spatial distribution

of VBRV outbreaks in livestock and revealed hotspots of disease underreporting.

Previous work carried out in Brazil, which assessed the implications of the geographic

distribution of vampire bat roosts in the occurrence of rabies in livestock,

demonstrated a positive correlation between detected bat roosts and density of rabies

outbreaks in livestock [53,58]. Here, compared with raw roost data and its smooth

surface, our roost distribution model had a better fit and explained more of the spatial

pattern of past rabies outbreaks in Peru (Table 1). However, differences between the

performance (i.e., pseudo-R^2) of models with the smoothed raw data and roost

predictions were relatively small. This may reflect the fact that roost detection and

notification of rabies outbreaks were generated by similar observation processes, both

being affected by distance from the nearest reporting centre. Nevertheless, the modelling

approach presented here is novel and different from previous studies as we accounted for

undetected bat roosts. This is particularly advantageous for identifying high-risk areas for

disease spillover which can be targeted to strengthen rabies awareness (i.e., increasing

chances of reporting), surveillance, and control (i.e., vaccination and synchronised culling,

bat vaccination) [59].

In Latin America, rabies is a neglected zoonotic disease, and suspected outbreaks in

livestock are notified via passive surveillance. Hence, underreporting of rabies outbreaks

leads to underestimation of the true burden of the disease [12,17,54]. In previous research,

Benavides et al. [17] used questionnaires to estimate underreporting in AAC, predicting that

mortality from VBRV was 4.6 times (95% CI: 4.4– 8.2) higher than officially reported, and

inferring spatial heterogeneity in underreporting at district level. Here, without use of

extensive questionnaires, our model estimated that between 2003 and 2021, mortality from

VBRV in AAC was 7.1 times (95% CI: 3.8–10.5) higher than officially reported. In addition,

our predictions of rabies underreporting demonstrated spatial heterogeneity at a finer

resolution than previously possible (Fig. 4C). These results have two distinct implications.

First, our elevated estimate of underreporting implies an underestimation of the disease's true

burden which can be used to inform VBR management. Secondly, our ability to estimate

underreporting at fine spatial scales empowers more precise geographic allocation of

educational campaigns to encourage reporting where they are most needed.

Although the statistical approach presented here provides advances in modelling

partially observed wildlife disease reservoirs, our projections may be improved in several

ways. Future work could take advantage of more sophisticated extensions of SDMs that

integrate distinct but complementary data on vampire bat occurrence (e.g., roosts and reports

of bat bites) [31]. Additionally, integrating the observation and the roost models, rather than

modelling them separately as done here, would also be advantageous to fully

propagate uncertainty across models. A joint likelihood approach with a parametric

form for the detection function would be a potential approach to extend our

framework [60,61].

In conclusion, we present a transferrable statistical approach to model the

spatial distribution of difficult to observe species, demonstrating the benefits of

consider spatial autocorrelation and observation effort. To our knowledge, this is the

first-time observation effort and spatial autocorrelation have been used to reconstruct

the map of likely roosts for any bat species. Our outputs identified putative hot and

cold spots of vampire bat roosts and risk areas for VBRV spread, and areas with high

number of unreported rabies outbreaks. These results can be valuable in spatial

models that explore determinants of viral maintenance and spillover risk and can

guide effective monitoring of *D. rotundus* and prevention and control strategies for

VBR, such as campaigns to increase awareness of rabies, preventive vaccination of

livestock and in future, vaccination of vampire bat colonies. More generally, our

results show how incorporating existing and routinely collected data into powerful

statistical models can improve the management of zoonoses.

Author contributions

Conceptualisation: R.R., J.M. and D.G.S.; Formal analysis: R.R.; Funding acquisition: T.E.R.

and D.G.S.; Methodology: R.R., J.M., F.L., and D.G.S.; Resources: C.T., C.M.Z., W.V. and

D.G.S.; Supervision: J.M. and D.G.S.; Visualization: R.R.; Writing-original draft: R.R., J.M.

and D.G.S. R.R.; Writing-review and editing: All authors reviewed the and gave final

approval for publication.

Funding sources

The project was funded by the NSF/BBSRC Ecology and Evolution of Infectious Diseases Program (DEB 2011069, BB/V003798/1). D.G.S. was funded by a Wellcome Trust Senior Research Fellowship (217221/Z/19/Z).

Acknowledgments

The authors would like to thank Andrew Seaton (University of Glasgow) for assistance with the process of predicting counts from an LGCP; Luca Nelli (University of Glasgow) for the suggestion of relevant variables; all colleagues and students who provided valuable comments on the model, results, and draft: Iosu Paradinas (Asociación Ipar Perspective), all members of the Streicker lab (Nardus Mollentze, Laura Bergner, Jocelyn Pérez Lazo, Max Farrell, Megan Griffiths, Hollie French, Haris Malik, Alice Broos, and Matthew Arnold), Jorge Osorio (University of Wisconsin-Madison), Zulma E. Rojas-Sereno (Universidad Andres Bello), from the University of Glasgow: Yacob Haddou, Jana Jeglinski, and Landry Green, and Fergus Chadwick (BioSS) and Crinan Jarret (vogelwarte). Paulo Colchao and Manuel Sime for their support in roost detection. The dataset of VBRV outbreaks in livestock between 2003 and 2021 was kindly provided by SENASA.

Conflict of interest

The authors have no conflict of interest to declare.

Data availability statement

The data and R scripts that support the findings of this study are available in Zenodo [62].

References

1.   Plowright RK, Parrish CR, McCallum H, Hudson PJ, Ko AI, Graham AL, Lloyd-

Smith JO. 2017 Pathways to zoonotic spillover. *Nat. Rev. Microbiol.* **15**, 502–510.

(doi:10.1038/nrmicro.2017.45)

2.   European Commission. 2009 Eradication of bovine, sheep and goats brucellosis in the

EU. *Eur. Comm. Heal. Consum. Dir.* **1**.

3.   Thanapongtharm W, Kasemsuwan S, Wongphruksasoong V, Boonyo K,

Pinyopummintr T, Wiratsudakul A, Gilbert M, Leelahapongsathon K. 2021 Spatial

distribution and population estimation of dogs in Thailand: Implications for rabies

prevention and control. *Front. Vet. Sci.* **8**, 1–12. (doi:10.3389/fvets.2021.790701)

4.   Lugelo A, Hampson K, Ferguson EA, Czupryna A, Bigambo M, Duamor CT, Kazwala

R, Johnson PCD, Lankester F. 2022 Development of dog vaccination strategies to

maintain herd immunity against rabies. *Viruses* **14**, 1–17. (doi:10.3390/v14040830)

5.   Razgour O, Rebelo H, Di Febbraro M, Russo D. 2016 Paiting maps with bats: Species

distribution modelling in bat research and conservation. *Hystrix, Ital. J. Mammal.* **27**.

(doi:doi:10.4404/hystrix-27.1-11753)

6.   Mengersen K *et al.* 2017 Modelling imperfect presence data obtained by citizen

science. *Environmetrics* **28**, 1–29. (doi:10.1002/env.2446)

7.   Mollentze N, Streicker DG. 2020 Viral zoonotic risk is homogenous among taxonomic

orders of mammalian and avian reservoir hosts. *Proc. Natl. Acad. Sci. U. S. A.* **117**,

9423–9430. (doi:10.1073/pnas.1919176117)

8.   Roberts M, Dobson A, Restif O, Wells K. 2021 Challenges in modelling the dynamics

of infectious diseases at the wildlife–human interface. *Epidemics* **37**, 100523.

(doi:10.1016/j.epidem.2021.100523)

9.   Purse B V., Golding N. 2015 Tracking the distribution and impacts of diseases with

biological records and distribution modelling. *Biol. J. Linn. Soc.* **115**, 664–677.

(doi:10.1111/bij.12567)

10.   Laxton MR, Rodríguez de Rivera Ó, Soriano-Redondo A, Illian JB. 2022 Balancing

structural complexity with ecological insight in Spatio-temporal species distribution

models. *Methods Ecol. Evol.* **2022**, 1–11. (doi:10.1111/2041-210X.13957)

11.   Miller JA. 2012 Species distribution models: Spatial autocorrelation and non-

stationarity. *Prog. Phys. Geogr.* **36**, 681–692. (doi:10.1177/0309133312442522)

12.   World Health Organization. 2018 *WHO expert consultation on rabies: third report.*

World Health Organization. See https://apps.who.int/iris/handle/10665/272364.

13.   Johnson N, Aréchiga-Ceballos N, Aguilar-Setien A. 2014 Vampire bat rabies:

Ecology, epidemiology and control. *Viruses* **6**, 1911–1928. (doi:10.3390/v6051911)

14.   Johnson N, Montano Hirose JA. 2018 The impact of paralytic bovine rabies

transmitted by vampire bats in Latin America and the Caribbean. *Rev. Sci. Tech.* **37**,

451–459. (doi:10.20506/rst.37.2.2814)

15.   Streicker DG *et al.* 2012 Ecological and anthropogenic drivers of rabies exposure in

vampire bats: Implications for transmission and control. *Proc. R. Soc. B Biol. Sci.* **279**,

3384–3392. (doi:10.1098/rspb.2012.0538)

16.   Schneider MC, Romijn PC, Uieda W, Tamayo H, da Silva DF, Belotto A, da Silva JB,

Leanes LF. 2009 Rabies transmitted by vampire bats to humans: an emerging zoonotic

disease in Latin America? *Rev. Panam. Salud Publica* **25**, 260–269.

(doi:10.1590/s1020-49892009000300010)

17.   Benavides JA, Rojas Paniagua E, Hampson K, Valderrama W, Streicker DG. 2017

Quantifying the burden of vampire bat rabies in Peruvian livestock. *PLoS Negl. Trop.*

Dis. **11**, 1–17. (doi:10.1371/journal.pntd.0006105)

18.   Arellano-Sota C. 1988 Vampire bat-transmitted rabies in cattle. *Rev. Infect. Dis.* **10**

Suppl 4, S707-9. (doi:10.1093/clinids/10.supplement_4.s707)

19.   Linhart SB, Crespo RF, Mitchell GC. 1972 Control of vampire bats by topical

application of an anticoagulant, Chlorophacinone. *Bol. la OSP* **VI**.

20.   Fornes A, Lord RD, Kuns ML, Larghi OP, Fuenzalida E, Lazara L. 1974 Control of

bovine rabies through vampire bat control. *J. Wildl. Dis.* **10**, 310–316.

(doi:10.7589/0090-3558-10.4.310)

21.   Blackwood JC, Streicker DG, Altizer S, Rohani P. 2013 Resolving the roles of

immunity, pathogenesis, and immigration for rabies persistence in vampire bats. *Proc.*

Natl. Acad. Sci. U. S. A. **110**, 20837–20842. (doi:10.1073/pnas.1308817110)

22.   Viana M *et al.* 2023 Effects of culling vampire bats on the spatial spread and spillover

of rabies virus. *Sci. Adv.* , 20. (doi:10.1126/sciadv.add7437)

23.   Benavides JA, Valderrama W, Streicker DG. 2016 Spatial expansions and travelling

waves of rabies in vampire bats. *Proc. R. Soc. B Biol. Sci.* **283**.

(doi:10.1098/rspb.2016.0328)

24.   Rojas-Sereno ZE, Streicker DG, Medina-Rodríguez AT, Benavides JA. 2022 Drivers

of Spatial Expansions of Vampire Bat Rabies in Colombia. *Viruses* **14**, 2318.

(doi:10.3390/v14112318)

25.   Janoušková E, Rokhsar J, Jara M, Entezami M, Horton DL, Dias RA, Machado G,

Prada J. 2022 Quantifying spillover risk with an integrated bat-rabies dynamic

modeling framework. (doi:10.22541/au.165402909.96757900/v1)

26.   Zarza H, Martínez-Meyer E, Suzán G, Ceballos G. 2017 Geographic distribution of

Desmodus rotundus in Mexico under current and future climate change scenarios:

Implications for bovine paralytic rabies infection. *Vet. Mex.* **4**.

(doi:10.21753/vmoa.4.3.390)

27.   Hayes MA, Piaggio AJ. 2018 Assessing the potential impacts of a changing climate on

the distribution of a rabies virus vector. *PLoS One* **13**, 1–17.

600   (doi:10.1371/journal.pone.0192887)

601 28. Lee DN, Papeş M, van Den Bussche RA. 2012 Present and potential future distribution

602   of common Vampire bats in the Americas and the associated risk to cattle. *PLoS One*

603   **7**, 1–9. (doi:10.1371/journal.pone.0042466)

604 29. Strimas-Mackey M. 2021 smoothr: Smooth and Tidy Spatial Features.

605 30. Streicker DG *et al.* 2016 Host-pathogen evolutionary signatures reveal dynamics and

606   future invasions of vampire bat rabies. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 10926–

607   10931. (doi:10.1073/pnas.1606587113)

608 31. Illian JB, Sørbye SH, Rue H. 2012 A toolbox for fitting complex spatial point process

609   models using integrated nested Laplace approximation (INLA). *Ann. Appl. Stat.* **6**,

610   1499–1530. (doi:10.1214/11-AOAS530)

611 32. Lindgren F, Rue H, Lindström J. 2011 An explicit link between Gaussian fields and

612   Gaussian Markov random fields: The stochastic partial differential equation approach.

613   *J. R. Stat. Soc. Ser. B Stat. Methodol.* **73**, 423–498. (doi:10.1111/j.1467-

614   9868.2011.00777.x)

615 33. Rue H, Held L. 2005 *Gaussian Markov Random Fields: Theory and Applications*. 1st

616   edn. New York: Chapman & Hall/CRC. (doi:10.1201/9780203492024)

617 34. Bachl FE, Lindgren F, Borchers DL, Illian JB. 2019 inlabru: an R package for

618   Bayesian spatial modelling from ecological survey data. *Methods Ecol. Evol.* **10**, 760–

619   766. (doi:10.1111/2041-210X.13168)

620 35. R Core Team. 2021 R: A language and environment for statistical computing.

621 36. Rue H, Martino S, Chopin N. 2009 Approximate Bayesian inference for latent

622   Gaussian models using integrated nested Laplace approximations. *J. R. Stat. Soc. Ser.*

623   *B* **71**, 319–392.

624 37. Williams R, Hedley SL, Branch TA, Bravington M V., Zerbini AN, Findlay KP. 2011

Chilean blue whales as a case study to illustrate methods to estimate abundance and

evaluate conservation status of rare species. *Conserv. Biol.* **25**, 526–535.

(doi:10.1111/j.1523-1739.2011.01656.x)

38. Weiss DJ *et al.* 2018 A global map of travel time to cities to assess inequalities in

accessibility in 2015. *Nature* **553**, 333–336. (doi:10.1038/nature25181)

39. Lindgren F, Rue H. 2015 Bayesian spatial modelling with R-INLA. *J. Stat. Softw.* **63**,

1–25. (doi:10.18637/jss.v063.i19)

40. Fuglstad GA, Simpson D, Lindgren F, Rue H. 2019 Constructing priors that penalize

the complexity of gaussian random fields. *J. Am. Stat. Assoc.* **114**, 445–452.

(doi:10.1080/01621459.2017.1415907)

41. Lewis S, Sherratt TN, Hamer KC, Wanless S. 2001 Evidence of intra-specific

competition for food in a pelagic seabird. *Nature* **412**, 816–819.

(doi:10.1038/35090566)

42. Cameron AC, Windmeijer FAG. 1996 R-squared measures for count data regression

models with applications to health-care utilization. *J. Bus. Econ. Stat.* **14**, 209–220.

(doi:10.1080/07350015.1996.10524648)

43. Ulloa-Stanojlovic FM, Dias RA. 2020 Spatio-temporal description of bovine rabies

cases in Peru, 2003–2017. *Transbound. Emerg. Dis.* **67**, 1688–1696.

(doi:10.1111/tbed.13512)

44. Arellano-Sota C. 1988 Biology and control of the vampire bat. *Rev. Infect. Dis.* **10**,

615–619. (doi:10.1201/9780203736371)

45. Castro A *et al.* 2021 *Climas del Perú - Mapa de Clasificación Climática Nacional.*

Red Activa. Lima.

46. McNab BK. 1973 Energetics and the distribution of vampires. *J. Mammal.* **54**, 131–

144. (doi:10.2307/1378876)

47. Greenhall AM, Joermann G, Schmidt U. 1983 *Desmodus rotundus*. *Mamm. Species*

202, 1–6.

48. Wimsatt and Guerriere. 1961 Care and maintenance of the common vampire in

captivity. *J. Mammal.* **42**, 449–455.

49. Ardia DR. 2005 Tree swallows trade off immune function and reproductive effort

differently across their range. *Ecology* **86**, 2040–2046. (doi:10.1890/04-1619)

50. Scheele BC, Guarino F, Osborne W, Hunter DA, Skerratt LF, Driscoll DA. 2014

Decline and re-expansion of an amphibian with high prevalence of chytrid fungus.

Biol. Conserv. **170**, 86–91. (doi:10.1016/j.biocon.2013.12.034)

51. Glanville EJ, Murray SA, Seebacher F. 2012 Thermal adaptation in endotherms:

Climate and phylogeny interact to determine population-level responses in a wild rat.

Funct. Ecol. **26**, 390–398. (doi:10.1111/j.1365-2435.2011.01933.x)

52. Becker DJ *et al.* 2019 Leukocyte profiles reflect geographic range limits in a

widespread neotropical bat. *Integr. Comp. Biol.* **59**, 1176–1189.

(doi:10.1093/icb/icz007)

53. Mantovan KB, Menozzi BD, Paiz LM, Sevá AP, Brandão PE, Langoni H. 2022

Geographic distribution of common vampire bat *Desmodus rotundus* (Chiroptera:

Phyllostomidae) shelters: Implications for the spread of rabies virus to cattle in

southeastern Brazil. *Pathogens* **11**. (doi:10.3390/pathogens11080942)

54. Meske M, Fanelli A, Rocha F, Awada L, Soto PC, Mapitse N, Tizzani P. 2021

Evolution of rabies in south america and inter-species dynamics (2009–2018). *Trop.*

Med. Infect. Dis. **6**. (doi:10.3390/tropicalmed6020098)

55. Ripperger SP, Carter GG. 2021 Social foraging in vampire bats is predicted by long-

term cooperative relationships. *PLoS Biol.* **19**, 1–17.

(doi:10.1371/journal.pbio.3001366)

56.   Trajano E. 1996 Movements of cave bats in southeastern Brazil, with emphasis on the

676   population ecology of the common vampire bat, *Desmodus rotundus* (Chiroptera).

677   *Biotropica* **28**, 121. (doi:10.2307/2388777)

678   57.   Delpietro HA, Russo RG, Carter GG, Lord RD, Delpietro GL. 2017 Reproductive

679   seasonality, sex ratio and philopatry in Argentina's common vampire bats. *R. Soc.*

680   *Open Sci.* **4**. (doi:10.1098/rsos.160959)

681   58.   Ribeiro J, Vieira RGV, Martins CM, Ferreira F, Araujo JP, Ullmann LS, Dos Santos

682   AP, Biondo AW. 2021 Spatial distribution of bat shelters and livestock rabies in

683   southern Brazil. *Vector-Borne Zoonotic Dis.* **21**, 785–795.

684   (doi:10.1089/vbz.2020.2730)

685   59.   Cárdenas-Canales EM, Velasco-Villa A, Ellison JA, Satheshkumar PS, Osorio JE,

686   Rocke TE. 2022 A recombinant rabies vaccine that prevents viral shedding in rabid

687   common vampire bats (Desmodus rotundus). *PLoS Negl. Trop. Dis.* **16**.

688   (doi:10.1371/JOURNAL.PNTD.0010699)

689   60.   Yuan Y, Bachl FE, Lindgren F, Borchers DL, Illian JB, Buckland ST, Rue H,

690   Gerrodette T. 2017 Point process models for spatio-temporal distance sampling data

691   from a large-scale survey of blue whales. *Ann. Appl. Stat.* **11**, 2270–2297.

692   (doi:10.1214/17-AOAS1078)

693   61.   Royle JA, Dawson DK, Bates S. 2004 Modeling abundance effects in distance

694   sampling. *Ecology* **85**, 1591–1597. (doi:10.1890/03-3127)

695   62.   Ribeiro R, Mattiopoulos J, Lindgren F, Tello C, Zariquiey CM, Valderrama W, Rocke

696   TE, Streicker DG. 2023 Code and data for manuscript: Incorporating environmental

697   heterogeneity and observation effort to predict host distribution and viral spillover

698   from a bat reservoir. (Version 1). (doi:10.5281/zenodo.7704510)

Table 1: Estimate and 95% confidence intervals of the covariates included in the eight models used to compare the performance of four measures of bat distribution. The estimates of the fixed effects are presented in the logarithm and standardised forms. Models are displayed in descending order, from higher to lower AIC and pseudo-R^2.

Model	Covariates							AIC	pseudo-R^2
	Bat distribution				Cattle density	Interaction cattle and bat distribution	Months since first rabies outbreak		
	Roost data	Smooth roost data	Roost model	BUD					
1				0.52 [0.45; 0.59]	0.12 [0.03; 0.22]	0.19 [0.09; 0.29]	1.08 [0.96; 1.19]	3800.8	0.37
2			0.52 [0.45; 0.59]		0.13 [0.04; 0.23]	0.14 [0.05; 0.23]	1.13 [1.02; 1.25]	3830.4	0.36
3		0.54 [0.46; 0.62]			0.07 [-0.04; 0.17]	-0.07 [-0.17; 0.04]	1.12 [1.0; 1.24]	3842.6	0.35
4	0.31 [0.23; 0.38]				0.18 [0.08; 0.28]	-0.004 [-0.10; 0.09]	1.26 [1.15; 1.38]	3929.2	0.31
5				0.88 [0.81; 0.96]				4109.2	0.22
6			0.90 [0.83; 0.97]					4166.1	0.18
7		0.80 [0.73; 0.88]						4175.3	0.18
8	0.41 [0.33; 0.49]							4368.0	0.06

Note: The pseudo-R^2 of a multivariable model only with cattle density and the number of months since first rabies outbreaks, is 0.28.

Fig. 1: The study area with detailed spatial locations of vampire bat roosts and rabies outbreaks. A - Elevation in Peru (source https://www.worldclim.org) and the boundaries (white) of the departments (from left to right) of Ayacucho, Apurimac, and Cusco. B – Zoom to the three departments and the study area, the inter-Andean valleys ('amebae shape'). These valleys exclude landscape with elevation above 4,000 m. C - The inter-Andean valleys of the AAC area (Ayacucho in light blue, Apurimac in light green and Cusco in light yellow), with the number of observed vampire bat roosts (n=563) and the number of reported rabies outbreaks between 2003 and 2021 (n=1,212).

Fig. 2: Factors affecting roost detectability and expected density: A – Non-linear effect of accessibility. X-axis is travelling time (log minutes, i.e., exp (4) = 55 min, and exp (6) = 7 h) and Y-axis is the mode of roost density (in the linear predictor scale). B - Range of the Matérn correlation model showing the mean correlation (Y-axis) in function of the distance between points. As the correlation range is on the log-intensity scale, on the intensity scale the range of correlation is shorter. Both in A and B, the grey shaded areas show the 95% credible interval. C - Posterior mean and respective 95% credible intervals of fixed effects of roost density. Red for positive significant coefficients (i.e., positive credible intervals), black for non-significant coefficients (i.e., include 0), and blue for negative significant coefficients (i.e., negative credible intervals).

Fig. 3: Spatial predictions of the detectability and roost models. A - Probability of detecting one roost (output from the detectability model). The detection probability is a combination of accessibility and surveillance effort in each region. Yellow for high probability of roost observation and dark purple for low probability of roost detection. B - Predictions of detected roosts. The black points are the 563 observed roosts. C – Predictions of the expected roosts when adjusting for the uneven effort. In both B and C, the predicted posterior mean of roosts was mapped; also, maps have the same scale, and the same continuous colour gradient was applied, with red indicating a higher mean of predicted roosts. D – Missing roosts (C-B). Light colour indicates higher number of missing roosts.

Fig. 4: Spatial predictions of rabies outbreaks and heterogeneity in underreporting. A – Predictions of reported rabies outbreaks. The black points are the 1,212 reported rabies outbreaks in livestock between 2003 and 2021. B – Predictions of rabies outbreaks after correcting for accessibility to reporting offices. A and B have the same scale. A continuous colour gradient was applied, with red indicating a higher number of outbreaks. C – Underreporting in cells where outbreaks were reported. The cells were grouped according to the ratio between predictions corrected for accessibility and reported outbreaks. The two thresholds used were the average (i.e., 7.1) and the median (i.e., 2.4) of underreporting in the cells where outbreaks were reported (493 cells out of 4,797).