Three-dimensional chromatin organization promotes genome evolution in a fungal plant pathogen

David E. Torres[1,2,#], H. Martin Kramer[1,#], Vittorio Tracanna[3#], Gabriel L. Fiorin[1], David E. Cook[1,4], Michael F. Seidl[1,2,§,*], Bart P.H.J. Thomma[1,3,§,*]

[1]Laboratory of Phytopathology, Wageningen University and Research, Droevendaalsesteeg 1, 6708 PB Wageningen, the Netherlands

[2]Theoretical Biology & Bioinformatics Group, Department of Biology, Utrecht University, Utrecht, The Netherlands

[3]University of Cologne, Institute for Plant Sciences, Cluster of Excellence on Plant Sciences (CEPLAS), 50674 Cologne, Germany

[4]Department of Plant Pathology, Kansas State University, 1712 Claflin Road, Manhattan, Kansas 66506, USA

[#]These authors contributed equally

[§]These authors contributed equally

[*]For correspondence: bthomma@uni-koeln.de and m.f.seidl@uu.nl

ABSTRACT

The spatial organization of eukaryotic genomes is linked to their biological functions,

although it is not clear how this impacts the overall evolution of a genome. Here, we uncover

the three-dimensional (3D) genome organization of the phytopathogen *Verticillium dahliae*,

known to possess distinct genomic regions, designated adaptive genomic regions (AGRs),

enriched in transposable elements and genes that mediate host infection. Short-range DNA

interactions form clear topologically associating domains (TADs) with gene-rich boundaries

that show reduced levels of gene expression and reduced genomic variation. Intriguingly,

TADs are less clearly structured in in AGRs than in the core genome. At a global scale, the

genome contains bipartite long-range interactions, particularly enriched for AGRs and more

generally containing segmental duplications. Notably, the patterns observed for *V. dahliae* are

also present in other *Verticillium* species. Thus, our analysis links 3D genome organization to

evolutionary features conserved throughout the *Verticillium* genus.

INTRODUCTION

The spatial organization of eukaryotic genomes is directly linked to their biological functions, although underlying mechanisms remain largely unclear. Many plant pathogenic fungi display distinct genome organization, commonly termed the 'two-speed genome', in which gene-poor and repeat-rich genomic regions contain genes that mediate host infection, display increased plasticity, and typically display features of heterochromatin[1,2]. These dynamic compartments are paramount for the coevolutionary 'arms-race' between pathogens and their hosts, potentially enabling the avoidance of host immune recognition and evolution of novel functions in pathogenicity[2–5].

In eukaryotic nuclei, physically separated genomic sites colocalize, while proximal sites may be separated through folding barriers into a three-dimensional (3D) genome structure that comprises various levels of organization[6,7]. Local 3D interactions shape chromosome structure into discrete genomic regions, commonly known as topologically associating domains (TADs); self-interacting genomic regions that are delineated by boundaries that display less interaction[8]. Although their function is still controversial, TADs have been associated with transcriptional regulation by governing the impact of regulatory sequences on nearby genes[8–10]. Other studies implicate TADs in genome replication by synchronizing origins of replication[11]. Interestingly, various studies point towards evolutionary conservation of TAD organization among related organisms[12–14]. While the genome clearly contains different chromatin states and 3D genome organization, their exact roles in genome function and evolution remain unclear.

In the fungi *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, and *Neurospora crassa*, the 3D genome organization is linked to heterochromatin distribution[15–17]. Determination of the 3D genome of the endophytic fungus *Epichloë festucae* has revealed that heterochromatic, repeat-rich regions frequently colocalize with TAD boundaries and can be

implicated in genome folding[18]. Moreover, *in planta* induced genes are enriched near those

regions[18]. These findings suggest an intimate link between the 3D genome, dynamic genomic

compartments, heterochromatin, and conditional gene expression.

The asexual soil-borne fungal plant pathogen *Verticillium dahliae* is a notorious broad

host-range vascular wilt pathogen[19]. The *V. dahliae* genome is characterized by the

occurrence of extensive large-scale genomic rearrangements that are associated with

segmental duplications that underwent substantial reciprocal gene losses and are enriched in

active transposable elements (TEs)[20–23]. This organization results in distinct dynamic genomic

compartments, previously termed lineage-specific regions, and presently referred to as

adaptive genomic regions (AGRs)[20–22,24]. These AGRs display unique chromatin

characteristics and are enriched for *in planta*-induced genes and other conditionally

responsive genes that contribute to environmental adaptation[20,23–25]. However, it presently

remains unclear whether and how the 3D genome affects the organization and evolution of *V.*

dahliae genome. Here, we explore the chromatin conformation of *V. dahliae* with DNA

proximity ligation followed by sequencing (Hi-C) to uncover the spatial organization of the

AGRs within the genome.

RESULTS

Topologically associating domains in the *Verticillium dahliae* genome

To uncover the 3D organization in *V. dahliae* strain JR2, we performed Hi-C in two highly correlated biological replicates (Fig. S1a-d), revealing that interaction strength generally negatively correlated with genomic distance (Fig. S1c). To query for the occurrence of TADs, the genome was divided into bins (average bin size ~4 kb) and interaction strengths between adjacent bins were determined. Given that a TAD is a self-interacting genomic region with sequences that physically interact more with each other than with sequences outside the TAD, bins that display a low insulation score weakly interact with neighboring bins and consequently were assigned as a TAD boundary region. Using this approach, we identified 353 TADs (mean size 102 kb) separated by 347 boundaries (mean size 4.7 kb) along the eight *V. dahliae* chromosomes (Fig. 1a,b,d, Fig. S1e,f). While 277 TADs (78.47%) and 308 boundaries (88.76%) localize in the core genome, 76 TADs (21.53%) and 39 boundaries (11.24%) localized in AGRs. Interestingly, the insulation of TADs is weaker in AGRs than in the core genome (Fig. 1c).

As observed previously[23–27], the core genome is generally enriched in H3K27ac and H3K4me2, while centromeres and TE-rich core regions are enriched in H3K9me3 and DNA methylation, and AGRs are enriched in H3K27me3 (Fig. 1d,e, Fig. S2). Such broad chromatin associations are maintained similarly on TADs and boundaries in core and AGRs (Fig. S3a,b), suggesting that chromatin characteristics mainly associate with the overall genome compartmentalization rather than with TAD organization. Nevertheless, we also observe that TADs and boundary regions differ in chromatin accessibility, histone modifications, and DNA methylation (Fig. 1e, Fig. S3b), and these observations support that TADs and boundaries differ in functionality and suggest that they may be further functionally dependent on AGR versus core compartments.

Figure 1. The *Verticillium dahliae* genome is organized in topological associating domains (TADs). (a) Hi-C contact matrix showing local interaction frequency, aggregated over predicted TADs (black bars) with 50 kb up- and down-stream sequence. The drop in intensity at boundaries at either side of the TADs indicates stronger interaction within TADs than with neighboring genomic regions. **(b)** Heatmap showing insulation scores centered over boundaries with 50 kb up- and down-stream sequence as rows, ordered on insulation score with weakest insulated boundaries on top. The top plot displays the average of the insulation scores in the heatmap below. **(c)** TADs in AGRs are weaker insulated when compared with core genome TADs. The X-axis indicates quintiles of boundaries, separated based on insulation scores. The Y-axis indicates Z-score and the -log10(p-value) color-scale after a permutation test for enrichment of boundaries in AGRs (10,000 iterations). The plot displays a linear regression (blue line) and confidence interval (light blue) as well as the R and p-value for the linear regression. **(d)** TAD distribution in *V. dahliae* strain JR2, exemplified by a section of chromosome 5. From top to bottom: Hi-C contact matrix depicting TADs as black triangles, open chromatin determined with ATAC-seq, histone modifications H3K4me2, H3K27ac, H3K27me3, and H3K9me3 normalized over a micrococcal nuclease digestion control, GC methylation, as well as gene and transposable element (TE) densities in 10 kb windows.

115 Adaptive genomic regions (AGRs) and the centromeric region are indicated in blue and black, respectively.
116 (e) Chromatin characteristics are differentially associated with TAD boundaries in the core genome and in
117 AGRs. On top, distribution of each chromatin feature centered for boundaries (dashed lines) with 50 kb up-
118 and down-stream sequence, for the core genome (dark line) and AGRs (light line). On the bottom the
119 corresponding heatmaps are shown.
120

Local TAD organization impacts transcriptional regulation

Given the chromatin differences between TADs and boundaries in AGRs and the enrichment

of AGRs in conditionally responsive genes[20,24,25], we hypothesized that transcriptional profiles

of core genes and those in AGRs differ between TADs and boundaries. Interestingly,

boundary regions in the core genome are significantly enriched in genes and consequently

depleted of TEs when compared with TADs (Fig. 2a). However, within AGRs we did not

observe differential enrichment of genes or TEs in TADs versus boundaries (Fig. 2a). To

assess the impact of TAD organization on gene expression, we integrated genomic,

transcriptomic, and chromatin data for all *V. dahliae* genes, and performed uniform manifold

approximation and projection (UMAP) for dimensional reduction[24,25]. As observed

previously, genes separated based on these characteristics show core and AGR groupings

(Fig. S3c)[24]. Additionally, when considering only boundary genes (Fig. S3c), genes in AGRs

are enriched for H3K27me3, while core genes are enriched for H3K27ac. Furthermore, there

was a clear separation based on transcriptional activity, with genes in core genome boundaries

generally displaying higher transcription levels, associated with increased H3K4me2 levels,

than genes in AGR boundaries (Fig. S3c). Interestingly, boundary insulation and expression

level positively correlated (Fig. 2b, Fig. S3c), indicating that genes in weakly insulated TAD

boundaries are generally lower expressed than genes in strongly insulated boundaries.

Overall, genes located within TAD boundaries are lower expressed than those in TADs, both

for the core genome and AGRs (Fig. 2d). However, the expression of genes within TAD

boundaries in the core genome is notably higher than the expression of genes within

142 boundaries in AGRs, whereas genes further away from boundaries in the core genome and

143 AGRs are similarly expressed (Fig. 2c).

145
146 **Figure 2. TAD organization affects transcription in *Verticillium dahliae*.** **(a)** Average density of genes
147 and transposable elements (TEs) per 1 kb window centred over boundaries with 50 kb up- and down-stream
148 sequence in the core genome and in adaptive genomic regions (AGRs) of *V. dahliae* strain JR2. The core
149 genome (dark line) and in AGRs (light line). **(b)** Genes in lowly insulated boundaries are lower expressed
150 than those in more highly insulated boundaries. Association between TAD boundary quintiles, separated on
151 insulation score, and transcription of genes located in TAD boundaries. The Y-axis depicts the Z-score,
152 colour of the datapoints indicates the -log10(p-value) after a permutation test (10,000 iterations) and size of
153 datapoints indicates mean transcription value (TPM) of represented genes. A linear regression (blue line),
154 with 95% confidence interval (light blue), between boundary quintiles is displayed. **(c)** Transcription values
155 for *V. dahliae* cultivated for 6 days in potato dextrose broth (PDB) and **(d)** absolute log2-fold change in
156 expression between cultivation in PDB or in Czapec-Dox medium (CZA), for all genes grouped based on
157 their distance to the closest boundary in the core genome (grey) or in AGRs (blue). Statistically significant
158 differences in average transcription level for the distance groups was compared to the group of genes
159 located in boundaries (distance 0) and determined by the Wilcoxon Rank-Sum test (* $p < 0.05$). **(e)** Linear
160 regression effect size of each TAD on differential gene expression between cultivation for 6 days in PDB or
161 in CZA. Mean effect size of each TAD is shown as a point, with 95% confidence interval, and TADs with a

162 significant effect (95% confidence interval is significantly different from 0) are shown in colour and
163 labelled by corresponding chromosome and TAD number, for TADs in the core genome (black labels) and
164 in AGRs (blue labels).
165

166    As TADs may govern differential gene expression, we hypothesized that differentially

expressed genes (DEGs) are enriched within TADs and depleted in boundaries. We

previously showed that DEGs are enriched in AGRs and in H3K27me3-rich regions in the

core genome[20,24,25], indicating that TADs may function as regulatory units both in the core

genome and in AGRs. However, we observed that only genes at distances of 5-15 kb and >25

kb from TAD boundaries in AGRs are significantly stronger differentially expressed than

genes located in AGR boundaries (Fig. 2d), suggesting that boundary genes are less

differentially expressed.

174    To investigate whether genes localizing within the same TAD in *V. dahliae* display

transcriptional co-regulation, we fitted a linear model in which differential expression of each

gene is predicted by TAD association. We identified 19 TADs with a significant effect on co-

transcription (Fig. 2e). Of these, 17 are associated with AGRs and only two with the core

genome (Fig. 2e), suggesting that transcriptional co-regulation of expression mainly occurs in

AGRs. In total, 68 out of 258 (26.4%) TADs in the core genome, while 64 out of 96 (66.7%)

TADs in AGRs contain more than five DEGs (Fig. S4B). Of these TADs, 13 out of 68

(19.1%) and 19 out 64 (29.7%) in the core genome and AGRs, respectively, contain more

than twice the number of genes that display co-regulation of differential transcription (Fig.

S3d). Thus, our results implies that TAD organization affects transcription and that, although

some TADs display transcriptional co-regulation of gene expression, this occurs only for a

subset of TADs that predominantly locate in AGRs.

TAD boundaries are conserved

To study TAD conservation within *V. dahliae*, we analysed Hi-C data of *V. dahliae* strain

VdLs17 that is 98% syntenic to strain JR2 (Fig. 3a)[21], and predicted 365 TADs (mean size 99

kb) and 357 boundaries (mean size 4.5 kb) (Fig. S4a-c). Notably, the TAD organization in

VdLs17 displays similar patterns of insulation scores, gene-enrichment, and TE-depletion as

in JR2, suggesting that TAD characteristics are conserved in *V. dahliae* (Fig. S4c-h).

Moreover, based on the distribution of TAD boundaries over syntenic regions between

VdLs17 and JR2 (n=342 and n=330 TADs, respectively), we observed a significant overlap

and correlation of insulation scores between boundaries of the two strains (n=225, p=9.6x10-

4, one-way Fisher exact test; Fig. 3a; Fig. S4g), and an overall overlap in TAD positions (Fig.

3a,b). Also, we observe that non-syntenic regions in VdLs17 are enriched for weak TAD

boundaries (z-score=2.3858, p=0.00001, permutation test after 10,000 iterations; Fig. S4h), a

characteristic of AGRs in *V. dahliae* strain JR2.

Genomic comparisons between *V. dahliae* strains have revealed extensive genomic

rearrangements and structural variations (SVs)[20–23]. However, as TAD boundaries are

conserved between *V. dahliae* strains JR2 and VdLs17, we hypothesized that boundaries may

lack such genomic variation. We used genome sequencing data of 42 *V. dahliae* strains[23,24] to

query the occurrence of single nucleotide variants (SNVs) and presence/absence

polymorphisms (PAVs) over TAD boundaries in *V. dahliae* strain JR2. Indeed, deletions,

duplications, inversions, and translocations occur more commonly in TADs than in

boundaries, indicating depletion of genomic variation from boundaries (Fig. 3c,d). One

possibility is that genomic variation at boundaries negatively impacts *V. dahliae* (i.e.

purifying selection). To assess this, we calculated the expected amount of SV breakpoints and

SV coverage occurring in boundaries and found that SVs occur less frequently than expected

in boundaries localized in syntenic regions (Fig. 3e). Interestingly, while we observed a

depletion of SNVs and of PAVs in boundaries in the core genome (Fig. 3c,d), boundaries in

AGRs showed increased PAV (Fig. S5c) combined with lower nucleotide diversity (Fig.

S5a,b). Moreover, SVs occur more commonly over boundaries in non-syntenic regions and in

AGRs, which agrees with previous observations that SVs occur frequently in non-syntenic

regions (Fig. 3e)[23]. Collectively, these findings indicate that TAD boundaries in the core

genome are strongly conserved, while boundaries in AGRs are evolutionary less stable.

Figure 3. Topological associating domain (TAD) organization is conserved in *Verticillium dahliae*. (a)
Top: *V. dahliae* strains JR2 and VdLs17 are highly similar as 97.84% and 98.30% of their respective
genomes are syntenic. Bottom: Most of the TAD boundaries overlap between JR2 and VdLs17. **(b)**
Syntenic block between JR2 chromosome 4 and VdLs17 chromosome 5 shows conserved distribution of

TADs and boundaries. Heatmaps represent contact matrixes of JR2 (top) and VdLs17 (bottom) with TADs
(black triangles). Genes and transposable elements (TEs) are displayed above and below. **(c)** Boundaries
are not enriched for genomic variation in a set of 42 *V. dahliae* strains. Nucleotide diversity, single
nucleotide variants (SNVs), presence/absence variation (PAVs). (d) Cumulative frequency plot of structural
variant (SV) breakpoints over distance from boundaries in the core genome (dashed line) and in AGRs
(solid line), overlaps with boundaries (distance = 0) are included. SVs are separated in deletions (DEL,
orange), duplications (DUP, yellow), inversions (INV, green) and translocations (TRA, blue) (Torres *et al.*,
2021). **(e)** TAD boundaries in AGRs and in the core genome contain more and fewer SVs than expected by
chance, respectively. SVs in boundaries in the core genome (open circles) and in AGRs (open triangles) are
indicated, as well as in boundaries in syntenic (solid circles) and non-syntenic (solid triangles) genomic
regions and in polymorphic TEs (grey circles). **(f)** Synteny breaks associated with transposable elements
affect TAD organization. Heatmaps represent contact matrixes of JR2 (top) and VdLs17 (bottom) with
TADs (black triangles), and TADs, genes and TEs are displayed in between. Syntenic regions are indicated
as grey blocks. A VdLTRE1 insertion in strain JR2 is indicated in yellow.

In *V. dahliae*, SVs often colocalize with polymorphic TEs that display PAV between

42 strains and are evolutionary young, scarcely methylated, and highly expressed[23]. As TE

activity may mediate the formation of SVs[22,23], we investigated whether polymorphic TEs

occur more frequently in TADs than in boundaries. We identified 36 polymorphic TEs

(21.8% of the total) that display PAV between *V. dahliae* strains JR2 and VdLs17. However,

we observed no overrepresentation nor depletion of polymorphic TEs in boundary regions

(Fig. 3e). Nevertheless, interestingly, some TE insertions in *V. dahliae* strain JR2 occur at

sites of boundary differences (Fig. 3f, Fig. S5d,e), suggesting that polymorphic TEs lead to

changes in TAD organization.

Conservation of TAD organization throughout the *Verticillium* genus

Given the conservation of TAD organization among *V. dahliae* strains, we investigated

conservation throughout the *Verticillium* genus based on synteny to *V. dahliae* strain JR2

(Fig. S5e) and calculated nucleotide conservation scores[28]. As expected, we observed higher

conservation scores for the core genome than for AGRs (Fig. 4a). Moreover, like *V. dahliae*

(Fig. 3c-e; Fig. S5f), we observed increased conservation scores for boundaries versus TADs

in the core genome, but not in AGRs (Fig. 4a), indicating that core boundaries are conserved

within the *Verticillium* genus.

Figure 4. TAD boundaries show signs of conservation in the *Verticillium* genus. (a) TAD boundaries
are more conserved than TADs. Boxplots display the conservation score of each TAD (grey) and boundary
(blue) in the core genome and in AGRs. P-values based on a one-way Wilcoxon rank sum test. **(b)**
Insulation score over TAD boundaries predicted by the sequence-based method, with 50 kb up- and down-
stream sequence, for each *Verticillium* species. Line plots display average signals over boundaries and
up/down-stream sequence, bottom plots display predicted boundaries in rows, ordered by insulation scores.

To assess conservation of TAD organization, we compared the sequence and position

of all TAD boundaries in *V. dahliae* strain JR2 to the genome sequences of the other

Verticillium species (Fig. S6a). We used previously generated Hi-C data[27] to assess whether

the insulation score of boundaries predicted based on sequence and position is lower than for

adjacent genomic regions, i.e. TADs (Fig. S6a). Employing this strategy to the genome of *V.*

dahliae VdLs17, we recovered 269 boundaries (Fig. S6b-d) that display significant positional

overlap with the boundaries as determined with the insulation method (z-score=27.1264,

p=9.99x10-5; Fig. 6b-e). Thus, we next used this approach for the other *Verticillium* spp. as

well. In general, the predicted boundaries in the *Verticillium* genus depict a drop in insulation

score with the adjacent genomic regions (Fig. 4b), indicating that we correctly assigned TAD

boundaries. The results show that boundaries of *V. dahliae* strain JR2 are more likely to be

shared with closer phylogenetic species (Fig. S6f). For instance, only 80 boundaries were

recovered in the more distantly related *V. albo-atrum*, whereas 254 boundaries were predicted

in *V. alfalfae* and 283 boundaries in *V. longisporum*, both close relatives of *V. dahliae* (Fig.

S6f). Additionally, we recovered full length TAD structures in syntenic regions of the other

Verticillium species, indicating high TAD conservation in the core genome (Fig. S6f).

Collectively, our results suggest that TADs and their boundaries are conserved among

Verticillium species consistent with phylogenetic distance.

Adaptive genomic regions physically colocalize

Besides local genome architecture, we assessed physical associations between distal genomic

regions. Making use of previously demonstrated centromeric interactions as references[27], we

identified 889 additional genomic regions that consistently colocalize. Interestingly, of these,

475 (53.4%) involve AGRs (Fig 5a,b, Table S1), which is a strong overrepresentation (chi-

squared test; p<0.05, Table S2) given that AGRs only represent 3.33 Mb (9.6%) of the 36.15

Mb total genome size[24]. Moreover, colocalization events among AGR regions comprise

nearly one-third of the non-centromeric interactions (28.1%), whereas AGR-core and core-

core colocalization events comprise 25.3% and 46.6%, respectively.

The colocalization of centromeres in *V. dahliae* strain JR2 correlates with CENH3

nucleosomes, *VdLTRE9* (LTR/Gypsy), and H3K9me3[27]. However, in our attempts to identify

epigenetic drivers of long-range interactions, we found no correlation between colocalizing

AGR regions and any of the histone modifications H3K9me3, H3K27me3, H3K4me2, and

H3K27ac, nor chromatin accessibility (Fig 5b). Given that all AGRs are involved in

individual bipartite colocalization events, and while AGRs share characteristics, they are

sequence diverse, it may not be surprising that long-ranger interactions do not have a simple

epigenome association (Fig 5b). We previously showed that *V. dahliae* evolution involved

large-scale segmental duplications[22,23]. Intriguingly, the colocalizing AGR regions are

associated with duplication events (Fig, 5c); of the 475 colocalization events that involve

AGR regions, 260 involve segmental duplications (Fig. 5c, Fig S7a, Table S3), which is a

significant enrichment not only genome-wide (z-test, $p<0.05$), but even within AGRs (z-test,

$p<0.05$). Moreover, whereas genome-wide 264 interactions were recorded that involve

segmental duplications, nearly all (260; 98.5%) concern AGRs.

Figure 5. Adaptive genomic regions physically colocalize in *Verticillium dahliae*. All circular plots
display in the outer track the eight chromosomes of *V. dahliae* with centromeres highlighted in yellow,
adaptive genomic regions (AGRs) in blue, and core regions in white. Long-range interactions in *V. dahliae*
strain JR2 (**a**) and VdLs17 (**d**) that exceed the average interaction strength of centromeres are shown as
edges. Edges for centromeric interactions are shown in yellow, AGR interactions in blue, and core
interactions in grey. Non-centromeric long range interactions in *V. dahliae* strains JR2 (**b**) and VdLs17 (**e**)
are shown as edges and the gene density (10 kb) is shown in black. For *V. dahliae* strain JR2, the inner
tracks depicting histone modification densities. From outside to inside: H3K27ac (green), H3K27me3
(yellow), H3K4me2 (purple), and H3K9m3 (blue). (**c** and **f**) Edges represent segmental duplications.

To assess conservation of AGR colocalization in *V. dahliae*, we similarly analyzed

HiC data of the VdLs17 strain. Interestingly, also in VdLs17, non-centromeric colocalization

events are enriched for AGR interactions that involve 452 out of 1,451 (31.2%) interactions

(chi-squared test; $p=8.5 \times 10^{-56}$; Fig. 5d,e). Moreover, we similarly observed that interacting

AGRs are strongly associated with segmental duplications (Fig. 5f, Fig S7b, Table 1).

Together, these results indicate that long-range interactions between AGRs represent a

conserved genome organization in *V. dahliae*.

We finally assessed whether the long-range colocalization patterns are conserved

throughout the *Verticillium* genus (Fig. 6, Table S3, Table S4). Interestingly, for several

Verticillium species we could demonstrate that AGRs are overrepresented in long-range

interactions (Table S5), namely in *V. albo-atrum*, *V. klebahnii*, *V. nubilum*, and *V.*

nonalfalfae. Interestingly, also in these species, segmental duplications are enriched in long-

range interactions. Moreover, although we were not able to demonstrate enrichment for

interactions between AGRs in *V. tricorpus*, *V. alfalfae*, *V. isaacii*, and *V. longisporum*, long-

range interactions predominantly occurred among segmental duplications in all species except

V. alfalfae (Fig. S9). Thus, long-range interactions are conserved across the *Verticillium*

genus and involve segmental duplications that are likely instrumental for AGR formation.

Figure 6. Adaptive genomic regions physically colocalize across the *Verticillium* genus. (**a**) Taxonomic relationship between ten *Verticillium* species. (**b**) Non-centromeric interactions occur preferentially between adaptive genomic regions (AGRs) in different *Verticillium* species. For all circular plots, outer track depicts centromeres in yellow, AGR regions in blue, and core regions in white. For every genome, non-centromeric interactions that exceed the average interaction strength of centromeres are shown. The right circular plots display segmental duplications for each genome.

DISCUSSION

Based on Hi-C analyses we uncovered the local and global 3D genome organization of *V. dahliae* and related species. The *V. dahliae* genome contains clear TADs that display reduced insulation in AGRs, as well as significantly enhanced co-regulation of gene expression compared with the core genome. Notably, TADs are conserved in the *Verticillium* genus and their boundaries generally lack structural variation. Intriguingly, we show that AGRs physically interact throughout the *Verticillium* genus, albeit not in an all-versus-all interaction like centromeres, but rather pairwise. These interactions are associated with segmental duplications that help to define AGRs. Collectively, our findings link 3D genome organization with genome function and evolution throughout the *Verticillium* genus.

TADs have been described for many eukaryotes, including fungi[11,15–18]. *V. dahliae* TADs are smaller (~100 kb) than typical mammalian TADs (~0.2-2.5 Mb)[8], yet similar to those in *Drosophila* and other fungi[11,16,18,29]. In metazoans, TAD boundaries are bound by CTCF (CCCTC-binding factor) proteins that act as insulator elements for cohesin-mediated loop extrusion, and are considered hallmarks of TADs[30,31]. Given that CTCF proteins have not been identified in filamentous fungi[32], annotation of fungal analogs as TADs may be considered controversial. However, TAD-like domains also occur in bacteria, known as chromosomal interaction domains (CIDs), and recently the global transcription repressor Rok was identified as a functional analog of the mammalian CTCF insulator elements[33]. A similar situation may be true for fungi, where additionally many other features are shared with metazoan TADs.

As DNA interacts more strongly within than between TADs, they separate genomes into discrete units that, in several organisms, coordinate DNA replication[11,34,35]. In addition, TADs have been implicated in transcriptional co-regulation[18,36–38], although a causal role in coordination of gene expression remains controversial[9]. Similar to studies on the fungi

Rhizophagus irregularis and *E. festucae*[18,38], we find that only few *V. dahliae* TADs display

transcriptional co-regulation. Nonetheless, it is still interesting that for this smaller subset of

TADs, they all reside in AGRs that are epigenetically distinct from the core genome,

involving lack of TE methylation, enrichment in H3K27me3, and accessible DNA[24].

Although in *V. dahliae*, similar to other plant-associated fungi[39–42], H3K27me3 plays a role in

transcriptional regulation, we also revealed that this modification does not act as the global

regulator of differential gene expression at AGRs[25], and additional factors must be involved in

transcriptional regulation.

TAD boundaries are typically conserved between close relatives[12,13,43–45], which we

similarly observe in *Verticillium*. However, in contrast to the core genome, TAD boundaries

in AGRs are less conserved, implying that TAD organization in the evolutionary younger and

dynamic AGRs still needs to settle or is less strictly defined. It is interesting to note that we

observed TE insertions near newly generated, or extensively rearranged, TAD boundaries in

AGRs. Cultivar-specific TAD boundaries in cotton were reported to generally harbor more

TEs than TAD boundaries that are shared between cultivars[46]. Moreover, *de novo* TE

insertions in humans generated new TAD boundaries[47]. TE activity is largely confined to

AGRs in *V. dahliae*[22,23], and may thus be involved in modulating TAD organization. How this

impacts longer-term trajectories of these regions remains to be determined.

Besides local interactions in TADs, the 3D genome displays long-distance interactions

within and between chromosomes. We previously reported centromere clustering in *V.*

dahliae[27], and now revealed long-range interactions among AGRs. In *N. crassa* such

interactions occur between heterochromatic regions, e.g. between H3K27me3 domains[16,48].

Thus, we hypothesized that H3K27me3-marked AGRs similarly colocalized in *V. dahliae*.

However, as not all AGRs interact, despite their enrichment in H3K27me3, and AGRs

interact in a pairwise fashion following the pattern of segmental duplications, we conclude

that we currently cannot assign a single epigenetic mark as a clear driver for colocalization. It

is tempting to speculate that physical clustering of AGRs in the nucleus involves membrane-

less nuclear bodies and liquid-liquid phase separation that allows spatial segregation of e.g.

transcription and DNA-repair[49,50]. Additionally, association with nuclear membranes could

cause spatial segregation of AGRs. In mammalian nuclei, heterochromatic regions are

associated with lamin proteins and additional anchor proteins to form lamina-associated

domains (LAD) at the nuclear membrane[51,52]. However, LAD proteins have not been found in

fungi[53], and more generally it is currently not clear what might drive these specific long-range

DNA interactions.

The 3D organization of chromosomes influences timing of replication[35,54,55] and may

lead to differential replication timing of AGRs and core genome. Such differences in

replication timing have been observed for H3K27me3-rich regions and have been associated

with chromosome instability in Z. tritici[41,56,57]. Such instability, and physical colocalization of

highly homologous sequences, may lead to decreased DNA separation efficiency during

mitosis and increased DNA double-strand breaks that may explain the increase in genomic

rearrangements in AGRs[20,24,58].

Altogether, we have uncovered a novel phenomenon that contributes to the divergence

of plastic regions and the core genome, by showing that their 3D organization differs with

clear impact on evolution and transcriptional regulation. This holistic view that combines

detailed genetic, epigenetic, and spatial information fosters further understanding of genome

function and evolution in fungi.

ACKNOWLEDGEMENTS

This work was supported by the Consejo Nacional de Ciencia y Tecnología, México to DET, by a PhD grant of the Research Council Earth and Life Sciences (project 831.15.002) to HMK, and by Human Frontier Science Program Postdoctoral Fellowship (HFSP, LT000627/2014-L), by USDA's National Institute of Food and Agriculture (award no. 2018-67013-28492) through the Plant Biotic Interactions Program, and by the National Science Foundation (award no. 1936800) Division of Molecular and Cellular Biosciences to DEC. Work in the laboratories of M.F.S and B.P.H.J.T. is supported by the Research Council Earth and Life Sciences (ALW) of the Netherlands Organization of Scientific Research (NWO). B.P.H.J.T acknowledges funding by the Alexander von Humboldt Foundation in the framework of an Alexander von Humboldt Professorship endowed by the German Federal Ministry of Education and Research is furthermore supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany´s Excellence Strategy – EXC 2048/1 – Project ID: 390686111.

AUTHOR CONTRIBUTIONS

M.F.S., D.E.C. and B.P.H.J.T. conceived the project. D.E.T., H.M.K., V.T., M.F.S. and B.P.H.J.T. designed the experiments. D.E.T., H.M.K., V.T., and G.L.F. carried out the experiments, D.E.T., H.M.K., V.T., G.L.F., D.E.C., M.F.S. and B.P.H.J.T. analyzed the data. D.E.T., H.M.K., V.T., M.F.S. and B.P.H.J.T. wrote the manuscript. All authors read and approved the final manuscript.

COMPETING INTERESTS

The authors declare no competing interests.

REFERENCES

1. Raffaele, S. & Kamoun, S. Genome evolution in filamentous plant pathogens: why bigger can be better. *Nature Reviews Microbiology* **10**, 417–430 (2012).

2. Torres, D. E., Oggenfuss, U., Croll, D. & Seidl, M. F. Genome evolution in fungal plant pathogens: looking beyond the two-speed genome model. *Fungal Biology Reviews* **34**, 136–143 (2020).

3. Seidl, M. F. & Thomma, B. P. H. J. Transposable elements direct the coevolution between plants and microbes. *Trends in Genetics* **33**, 842–851 (2017).

4. Croll, D. & McDonald, B. A. The accessory genome as a cradle for adaptive evolution in pathogens. *PLoS Pathogens* **8**, e1002608 (2012).

5. Frantzeskakis, L., Kusch, S. & Panstruga, R. The need for speed: compartmentalized genome evolution in filamentous phytopathogens. *Mol. Plant Pathol.* **20**, 3–7 (2019).

6. Lieberman-Aiden, E. *et al.* Comprehensive mapping of long-range interactions reveals folding principles of the human genome. *Science* **326**, 289–293 (2009).

7. Jerkovic, I. & Cavalli, G. Understanding 3D genome organization by multidisciplinary methods. *Nat. Rev. Mol Cell Biol.* 1–18 (2021).

8. Dixon, J. R. *et al.* Topological domains in mammalian genomes identified by analysis of chromatin interactions. *Nature* **485**, 376–380 (2012).

9. Cavalheiro, G. R., Pollex, T. & Furlong, E. E. M. To loop or not to loop: what is the role of TADs in enhancer function and gene regulation? *Curr. Opin. Genet. Develop.* **67**, 119–129 (2021).

10. Ghavi-Helm, Y. *et al.* Highly rearranged chromosomes reveal uncoupling between genome topology and gene expression. *Nat. Genet.* **51**, 1272–1282 (2019).

11. Eser, U. *et al.* Form and function of topologically associating genomic domains in budding yeast. *Proc. Natl. Acad. Sci. U. S. A.* **114**, E3061–E3070 (2017).

12. Harmston, N. *et al.* Topologically associating domains are ancient features that coincide

with Metazoan clusters of extreme noncoding conservation. *Nat. Commun.* **8**, 441 (2017).

13. Fudenberg, G. & Pollard, K. S. Chromatin features constrain structural variation across

evolutionary timescales. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 2175–2180 (2019).

14. Rowley, M. J. *et al.* Evolutionarily conserved principles predict 3D chromatin

organization. *Mol. Cell* **67**, 837-852.e7 (2017).

15. Mizuguchi, T. *et al.* Cohesin-dependent globules and heterochromatin shape 3D genome

architecture in *S. pombe*. *Nature* **516**, 432–435 (2014).

16. Galazka, J. M. *et al. Neurospora* chromosomes are organized by blocks of importin

alpha-dependent heterochromatin that are largely independent of H3K9me3. *Genome*

Res. **26**, 1069–1080 (2016).

17. Schalbetter, S. A., Fudenberg, G., Baxter, J., Pollard, K. S. & Neale, M. J. Principles of

meiotic chromosome assembly revealed in *S. cerevisiae*. *Nature Commun.* **10**, 1–12

(2019).

18. Winter, D. J. *et al.* Repeat elements organise 3D genome structure and mediate

transcription in the filamentous fungus *Epichloë festucae*. *PLoS Genetics* **14**, e1007467

(2018).

19. Fradin, E. F. & Thomma, B. P. H. J. Physiology and molecular aspects of Verticillium

wilt diseases caused by *V. dahliae* and *V. albo-atrum*. *Mol. Plant Pathol.* **7**, 71–86

(2006).

20. de Jonge, R. *et al.* Extensive chromosomal reshuffling drives evolution of virulence in an

asexual pathogen. *Genome Res.* **23**, 1271–1282 (2013).

21. Faino, L. *et al.* Single-molecule real-time sequencing combined with optical mapping

yields completely finished fungal genome. *mBio* **6**, (2015).

22. Faino, L. *et al.* Transposons passively and actively contribute to evolution of the two-
speed genome of a fungal pathogen. *Genome Res.* **26**, 1091–1100 (2016).

23. Torres, D. E., Thomma, B. P. H. J. & Seidl, M. F. Transposable elements contribute to
genome dynamics and gene expression variation in the fungal plant pathogen *Verticillium*
dahliae. *Gen. Biol. Evol* **13**, evab135 (2021).

24. Cook, D. E., Kramer, H. M., Torres, D. E., Seidl, M. F. & Thomma, B. P. H. J. A unique
chromatin profile defines adaptive genomic regions in a fungal plant pathogen. *eLife* **9**,
(2020).

25. Kramer, M., Seidl, M., Thomma, B. P. H. J. & Cook, D. Local rather than global
H3K27me3 dynamics associates with differential gene expression in *Verticillium dahliae*.
mBio **13**, e0356621 (2021).

26. Kramer, H. M., Cook, D. E., van den Berg, G. C. M., Seidl, M. F. & Thomma, B. P. H. J.
Three putative DNA methyltransferases of *Verticillium dahliae* differentially contribute
to DNA methylation that is dispensable for growth, development and virulence.
Epigenetics & Chromatin **14**, 1–15 (2021).

27. Seidl, M. F. *et al.* Repetitive elements contribute to the diversity and evolution of
centromeres in the fungal genus *Verticillium*. *mBio* **11**, e01714-20 (2020).

28. Siepel, A. *et al.* Evolutionarily conserved elements in vertebrate, insect, worm, and yeast
genomes. *Genome Res.* **15**, 1034–1050 (2005).

29. Sexton, T. *et al.* Three-dimensional folding and functional organization principles of the
Drosophila genome. *Cell* **148**, 458–472 (2012).

30. Kentepozidou, E. *et al.* Clustered CTCF binding is an evolutionary mechanism to
maintain topologically associating domains. *Genome Biology* **21**, 1–19 (2020).

31. Barutcu, A. R., Maass, P. G., Lewandowski, J. P., Weiner, C. L. & Rinn, J. L. A TAD

boundary is preserved upon deletion of the CTCF-rich Firre locus. *Nat. Commun.* **9**, 1–11

(2018).

32. Vietri Rudan, M. & Hadjur, S. Genetic tailors: CTCF and cohesin shape the genome

during evolution. *Trends Genet.* **31**, 651–660 (2015).

33. Dugar, G., Hofmann, A., Heermann, D. W. & Hamoen, L. W. A chromosomal loop

anchor mediates bacterial genome organization. *Nat. Genet.* **54**, 194–201 (2022).

34. Kolesnikova, T. D., Goncharov, F. P. & Zhimulev, I. F. Similarity in replication timing

between polytene and diploid cells is associated with the organization of the Drosophila

genome. *PLoS One* **13**, e0195207 (2018).

35. Pope, B. D. *et al.* Topologically associating domains are stable units of replication-timing

regulation. *Nature* **515**, 402–405 (2014).

36. Le Dily, F. & Beato, M. TADs as modular and dynamic units for gene regulation by

hormones. *FEBS letters* **589**, 2885–2892 (2015).

37. Jin, F. *et al.* A high-resolution map of the three-dimensional chromatin interactome in

human cells. *Nature* **503**, 290–294 (2013).

38. Yildirir, G. *et al.* Long reads and Hi C sequencing illuminate the two compartment

genome of the model arbuscular mycorrhizal symbiont *Rhizophagus irregularis*. *New*

Phytologist doi: 10.1111/nph.17842 (2021).

39. Connolly, L. R., Smith, K. M. & Freitag, M. The *Fusarium graminearum* histone H3

K27 methyltransferase KMT6 regulates development and expression of secondary

metabolite gene clusters. *PLoS Genetics* **9**, e1003916 (2013).

40. Soyer, J. L. *et al.* Epigenetic control of effector gene expression in the plant pathogenic

fungus *Leptosphaeria maculans*. *PLoS Genetics* **10**, e1004227 (2014).

41. Möller, M. *et al.* Destabilization of chromosome structure by histone H3 lysine 27

methylation. *PLoS Genetics* **15**, e1008093 (2019).

42. Zhang, W., Huang, J. & Cook, D. E. Histone modification dynamics at H3K27 are

associated with altered transcription of in planta induced genes in *Magnaporthe oryzae*.

PLoS Genetics **17**, e1009376 (2021).

43. Rao, S. S. P. *et al.* A 3D map of the human genome at kilobase resolution reveals

principles of chromatin looping. *Cell* **159**, 1665–1680 (2014).

44. McArthur, E. & Capra, J. A. Topologically associating domain boundaries that are stable

across diverse cell types are evolutionarily constrained and enriched for heritability. *Am.*

J. Hum. Genet. **108**, 269–283 (2021).

45. Krefting, J., Andrade-Navarro, M. A. & Ibn-Salem, J. Evolutionary stability of

topologically associating domains is associated with conserved gene regulation. *BMC*

Biol. **16**, 87 (2018).

46. Wang, M. *et al.* Comparative genome analyses highlight transposon-mediated genome

expansion and the evolutionary architecture of 3D genomic folding in cotton. *Molecular*

Biology and Evolution **38**, 3621–3636 (2021).

47. Zhang, Y. *et al.* Transcriptionally active HERV-H retrotransposons demarcate

topologically associating domains in human pluripotent stem cells. *Nature Genetics* **51**,

1380–1388 (2019).

48. Klocko, A. D. *et al.* Normal chromosome conformation depends on subtelomeric

facultative heterochromatin in *Neurospora crassa*. *Proc. Natl. Acad. Sci. U. S. A.* **113**,

15048–15053 (2016).

49. Sawyer, I. A. & Dundr, M. Nuclear bodies: Built to boost. *Journal of Cell Biology* **213**,

509–511 (2016).

50. Larson, A. G. *et al.* Liquid droplet formation by HP1α suggests a role for phase

separation in heterochromatin. *Nature* **547**, 236–240 (2017).

51. Harr, J. C. *et al.* Directed targeting of chromatin to the nuclear lamina is mediated by

chromatin state and A-type lamins. *J. Cell Biol.* **208**, 33–52 (2015).

52. Guelen, L. *et al.* Domain organization of human chromosomes revealed by mapping of

nuclear lamina interactions. *Nature* **453**, 948–951 (2008).

53. Koreny, L. & Field, M. C. Ancient eukaryotic origin and evolutionary plasticity of

nuclear lamina. *Genome Biol. Evol.* **8**, 2663–2671 (2016).

54. Gindin, Y., Valenzuela, M. S., Aladjem, M. I., Meltzer, P. S. & Bilke, S. A chromatin

structure-based model accurately predicts DNA replication timing in human cells. *Mol.*

Syst. Biol. **10**, 722 (2014).

55. Comoglio, F. *et al.* High-resolution profiling of Drosophila replication start sites reveals a

DNA shape and chromatin signature of metazoan origins. *Cell Rep.* **11**, 821–834 (2015).

56. Schotanus, K., Soyer, J. L. & Connolly, L. R. Histone modifications rather than the novel

regional centromeres of *Zymoseptoria tritici* distinguish core and accessory

chromosomes. *Epigenetics & Chromatin* 8:41 (2015).

57. Möller, M. & Stukenbrock, E. H. Evolution and genome architecture in fungal plant

pathogens. *Nat. Rev. Microbiol.* **15**, 771 (2017).

58. Huang, J. & Cook, D. E. The contribution of DNA repair pathways to genome editing

and evolution in filamentous pathogens. *FEMS Microbiol. Rev.* **46**, (2022).

59. Depotter, J. R. L. *et al.* The interspecific fungal hybrid *Verticillium longisporum* displays

subgenome-specific gene expression. *mBio* **12**, e0149621 (2021).

60. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina

sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

61. Li, H. & Durbin, R. Fast and accurate long-read alignment with Burrows–Wheeler

transform. *Bioinformatics* **26**, 589–595 (2010).

62. Wolff, J. *et al.* Galaxy HiCExplorer 3: a web server for reproducible Hi-C, capture Hi-C

and single-cell Hi-C data analysis, quality control and visualization. *Nucleic Acids*

Research **48**, W177–W184 (2020).

63. Imakaev, M. *et al.* Iterative correction of Hi-C data reveals hallmarks of chromosome

organization. *Nature Methods* **9**, 999–1003 (2012).

64. Yardımcı, G. G. *et al.* Measuring the reproducibility and quality of Hi-C data. *Genome*

Biology **20**, 1–19 (2019).

65. McInnes, L., Healy, J. & Melville, J. Umap: Uniform manifold approximation and

projection for dimension reduction. *arXiv* 1802.03426 (2018).

66. Hunter, J. D. Matplotlib: A 2D graphics environment. *Computing in Science &*

Engineering **9**, 90–95 (2007).

67. Waskom, M. *et al.* Mwaskom/Seaborn: V0. 8.1 (September 2017). *Zenodo* (2017).

68. van der Walt, S., Colbert, S. C. & Varoquaux, G. The NumPy array: a structure for

efficient numerical computation. *Computing in Science & Engineering* **13**, 22–30 (2011).

69. Gu, Z., Eils, R., Schlesner, M. & Ishaque, N. EnrichedHeatmap: an R/Bioconductor

package for comprehensive visualization of genomic signal associations. *BMC Genomics*

19, 1–7 (2018).

70. Team, R. C. R: A language and environment for statistical computing. (2013).

71. Kuzniar, A. *et al.* sv-callers: a highly portable parallel workflow for structural variant

detection in whole-genome sequence data. *PeerJ* **8**, e8214 (2020).

72. Goerner-Potvin, P. & Bourque, G. Computational tools to unmask transposable elements.

Nature Reviews Genetics **19**, 688–704 (2018).

604 73. Cameron, D. L., Di Stefano, L. & Papenfuss, A. T. Comprehensive evaluation and

605    characterisation of short read general-purpose structural variant calling software. *Nature*

606    *Commun.* **10**, 1–11 (2019).

607 74. Poplin, R. *et al.* A universal SNP and small-indel variant caller using deep neural

608    networks. *Nature Biotechnology* **36**, 983–987 (2018).

609 75. Stuart, T., Eichten, S. R., Cahn, J., Borevitz, J. & Lister, R. *Population scale mapping of*

610    *novel transposable element diversity reveals links to gene regulation and epigenomic*

611    *variation*. vol. 49 (2016).

612 76. Kurtz, S. *et al.* Versatile and open software for comparing large genomes. *Genome Biol.*

613    **5**, R12 (2004).

614 77. Lyons, E. *et al.* Finding and comparing syntenic regions among Arabidopsis and the

615    outgroups papaya, poplar, and grape: CoGe with rosids. *Plant Physiology* **148**, 1772–

616    1781 (2008).

617 78. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nature*

618    *Methods* **9**, 357–359 (2012).

619 79. Bertels, F., Silander, O. K., Pachkov, M., Rainey, P. B. & van Nimwegen, E. Automated

620    reconstruction of whole-genome phylogenies from short-sequence reads. *Mol. Biol. Evol.*

621    **31**, 1077–1088 (2014).

622 80. Armstrong, J. *et al.* Progressive Cactus is a multiple-genome aligner for the thousand-

623    genome era. *Nature* **587**, 246–251 (2020).

624 81. Hickey, G., Paten, B., Earl, D., Zerbino, D. & Haussler, D. HAL: a hierarchical format

625    for storing and analyzing multiple genome alignments. *Bioinformatics* **29**, 1341–1342

626    (2013).

82. Kruse, K., Hug, C. B. & Vaquerizas, J. M. FAN-C: a feature-rich framework for the

analysis and visualisation of chromosome conformation capture data. *Genome Biology*

21, 1–19 (2020).

83. Gel, B. *et al.* regioneR: an R/Bioconductor package for the association analysis of

genomic regions based on permutation tests. *Bioinformatics* **32**, 289–291 (2016).

84. Wickham, H. Programming with ggplot2. in *ggplot2* 241–253 (Springer, 2016).

ONLINE METHODS

Hi-C analysis and TAD prediction

Hi-C library preparation was performed with *V. dahliae* strains JR2 and VdLs17 as previously

described[27], and paired-end (2 × 150 bp) sequenced on the NextSeq500 platform at USEQ

(Utrecht, the Netherlands). Additional Hi-C datasets of *V. dahliae* strains JR2 and VdLs17, *V.*

albo-atrum strain PD747, *V. alfalfa* strain PD683, *V. isaacii* strain PD618, *V. klebahnii* strain

PD401, *V. longisporum* strains PD589 and VLB2, *V. nonalfalfae* strain T2, *V. nubilum* strain

397, *V. tricorpus* strain PD593, and *V. zaregamsianum* strain PD739 were previously

generated[27,59].

Sequenced read-pairs were quality-filtered and trimmed using Trimmomatic (v 0.36)

in paired end mode with default settings[60]. Filtered and trimmed reads were mapped to the

corresponding genomes[21,27] using Burrows-Wheeler aligner (BWA mem, settings: -A1 -B4 -

E50 -L0)[61]. Hi-C interaction matrices were built and analysed using HiCExplorer tools[62].

First, we used hicBuildMatrix to generate the interaction matrix based on the *in silico Dpn*II

restriction digested corresponding genome. Matrix resolution was reduced by merging 5

adjacent bins using hicMergeMatrixBins. For *V. dahliae* strains JR2 and VdLs17, replicates

were corrected separately according to the iterative correction and eigenvector decomposition

(ICE) method[63] using hicCorrectMatrix, and TADs were predicted using hicFindTADs

(settings: --delta 0.01). Correlation between replicates was determined by using a

reproducibility score based on a stratified cross-correlation using the HiCRep package[64].

To combine replicate matrices, resolutions of raw matrices were reduced by merging 5

adjacent bins using hicMergeMatrixBins, normalized between replicates using hicNormalize

(settings: --setToZeroThreshold 1), corrected separately according to the ICE method using

hicCorrectMatrix, and finally combined using hicSumMatrices[62]. For the other *Verticillium*

species, matrix resolution reduction and correction was performed as above, and

hicFindTADs was used to generate a table with per bin insulation scores.

Characterization of epigenetic profiles

Chromatin immunoprecipitation followed by sequencing (ChIP-seq) for H3K4me2,

H3K9me3, H3K27me3, and H3K27ac, and the assay for transposase-accessible chromatin

followed by sequencing (ATAC-seq) were performed for *V. dahliae* strain JR2 as described

previously[24,25,27]. ChIP datasets were normalized over MNase control samples.

We used the umap-learn implementation through the R/umap package. This

implementation make use of the python UMAP algorithm[65]. For the gene analysis, the

following variables were used: GC content, ATAC-seq, 5mC, H3K27ac, H3K27me3,

H3K9me3 and log2(PDB *in vitro* expression +1), with the following parameters

random_state=42, n_neighbors=50, n_components=2, min_dist=0.01, metric=cosine. The

resulting two-dimensional values from UMAP fit.transform were used for plotting and further

statistical analysis using Matplolib, Numbpy and Seaborn V0.8.1[66–68].

Characterization of transcriptional regulation

RNA sequencing of *V. dahliae* strain JR2 cultivated for six days in potato dextrose broth

(PDB) and Czapec-Dox medium (CZA) was previously performed[24,25]. Analyses of gene[21]

and TE presence[23,27] over TADs and TAD boundaries were performed using the

EnrichedHeatmap package in R[69,70]. To assess co-regulation of genes within TADs, we used

R to fit a linear model with log2 fold-change in expression of target genes between PDB and

CZA as the response variable and TAD membership as a predictor, similarly as previously

described[18].

Characterization of genomic variation

Structural variants (duplications, deletions, inversions and translocations), single nucleotide variants and polymorphic transposable elements were previously identified using paired-end sequencing reads of each 42 previously sequenced *V. dahliae* strains[23]. Briefly, structural variants were predicted using the 'sv-callers' workflow with few modifications that enabled parallel execution of multiple SV callers[71], an approach that is considered optimal as it exploits complementary information to predict SVs[72,73]. Single nucleotide variants were identified using the -HaplotypeCaller of the Genome Analysis Toolkit (GATK) v.4.0[74] and transposable element PAV was analyzed using TEPID v.2.0[75]. To investigate if SVs and polymorphic TEs co-localize with TAD boundaries, we summarized the overlap of each set of variants by their breakpoint frequency (start or ends ±1 bp of the feature) and coverage (number of bases covered) across the genome of *V. dahliae* strain JR2[13]. Similar to Fudenberg and Pollard (2019), we calculated the log10(observed/expected) of each feature representing the deviation from a uniform distribution across the genome, therefore accounting for the proportion of the genome covered by a specific genomic feature. Finally, we considered two scenarios: core genome vs AGRs, and syntenic regions between JR2 and VdLs17 versus non-syntenic regions. Syntenic regions between *V. dahliae* strains JR2 and VdLs17 were previously determined[22]. Briefly, whole-genome alignments between the eight chromosomes was performed using MUMmer 3.0 and GEvo[76,77], where only gene-coding regions were used as anchors between syntenic chromosomal regions.

To further expand our analysis of *V. dahliae* to the full genus, we used the recently available Hi-C-corrected genomes of all *Verticillium* species[27,59]. The phylogenetic tree was generated using Realphy v. 1.12 using a maximum likelihood inferred by RAxML[78,79]. We aligned the *Verticillium* genomes using ProgressiveCactus[80]. This approach allowed us to reduce the reference-bias and consider more accurate further analysis. We obtained the

specific MAF alignments on JR2 and syntenic regions using the HAL package[81]. To analyze

the nucleotide conservation throughout the genus, we used PhastCons, a hidden Markov

model-based method that estimates the probability that each nucleotide belongs to a

conserved element based on a multiple sequence alignment guided by stablished phylogenetic

relationships[28]. Briefly, for each independent JR2 chromosome we assumed a neutral

evolution model and correction for indels. For further analysis, we summarized the PhastCons

score over TADs and TAD boundaries in the core genome and in AGRs.

TAD boundary prediction throughout the *Verticillium* genus

The Hi-C datasets of the *Verticillium* species (excluding *V. dahliae*), were available with one

biological replicate. Therefore, we decided to predict TAD boundaries based on sequence

homology to boundaries in *V. dahliae* strain JR2. We first filtered the boundary sequences

that do not have a TE insertion and queried them to the *Verticillium* genomes using Blastn,

retaining those with >50% coverage that were contiguous in the same syntenic block. Finally,

we cross-referenced those putative TAD boundary regions with the previously calculated

insulation score for each independent species.

Statistical analysis and visualization

Hi-C matrix and TAD visualizations were performed using HiCExplorer and FAN-C[82].

Heatmap and enrichment visualization of insulation scores over boundaries, normalized

chromatin marks, structural and nucleotide variants, as well as the PhastCons score, were

performed using the R/EnrichedHeatmap v1.2 package[69]. Permutation tests were computed

using R/Bioconductor regionR v1.18.1 package[83] and performed with 10,000 iterations, using

overlaps between TAD boundaries divided by the insulation score quantiles and the

predefined AGRs, and circular randomization to maintain the order and distance of the

regions in the chromosomes. All statistical analyses and comparison tests were performed in

R v.3.6.3[70], and visualized with ggplot2[84].

Identification of significant colocalization events from Hi-C data

Expected-Observed counts were obtained from HiCExplorer (version 3.7) through the export

function with expected-observed nonzero from the corrected and summed matrices. Expected-

observed counts matrixes containing the interactions between all the merged bins were

filtered to keep only interaction between genetically distant regions. Two regions were

considered genetically distant if the bins belonged to different chromosomes or if two bins

were more than 20kb distant from each other. Next, the bins were annotated based on which

genomic compartment they belong to (centromere or AGR) and the remaining bins were

annotated as core genome. Centromere-centromere mean expected-observed count is

calculated and only bin to bin colocalization events with expected-observed counts above this

threshold are kept. Genomic bins which still have at least one interaction are considered as

genomic region involved in long range interaction. It needs to be noted that our analysis is

limited by the fact that we only assess those interactions that are stronger than the average

interaction strength that occurs between centromeres in each species but, given that the

constitution of centromeres differs between *Verticillium* spp.[27] this interaction strength may

consequently not be uniform between species. Visualization of the interaction and associated

genetic and epigenetic features was done in R with the circos package.

Identification of duplicated regions and association with long-range interactions

Self-whole-genome alignments of the genome assemblies of *Verticillium* species were

performed using MUMmer[76], regions that mapped elsewhere on the genome with a nucleotide

identity above 80% and above 1kb in size are considered duplicated. For each colocalization

event that associate two genetically distant regions, we verify if there is a duplicated region

that borders with or overlaps with both genetically distant regions. We consider the

colocalization event to be neighboring a duplicated region, if the average distance between the

colocalizing region and the duplicated region is below 50 kb.

We calculated the enrichment of colocalization events neighboring duplicated regions

through a permutation test. We simulated random interactions genome wide and calculated

the number of observed interactions neighboring a duplicated region. We repeated the process

100 times to generate a distribution of expected interactions neighboring duplicated regions

and ran a t-test versus the real observed value. This permutation test was performed using the

genome wide-duplicated regions distribution and AGR compartment specific.

Determination of adaptive genomic regions

Nucleotide sequences from the *V. dahliae* strain JR2 AGR compartment were aligned versus

each of the *Verticillium* genomes included in this work using ProgressiveCactus[80]. Syntenic

regions in the respective genomes are considered AGR regions in the respective organism.

Additionally, each *Verticillium* strain genome was aligned using ProgressiveCactus[80] with the

other *Verticillium* strain genomes included in this work. Genomic regions which were unique

and not found in any other genome are also considered AGR in that strain.