Adventitial macrophage accumulation impairs perivascular nerve function in mesenteric arteries with inflammatory bowel disease

Elizabeth A. Grunz[1], Benjamin W. Jones[1], Sidharth Sen[2], Katie Wilkenson[2], Trupti Joshi[2,3] and Erika M.
Boerman

[1]Department of Medical Pharmacology and Physiology, University of Missouri, Columbia, MO USA

[2]MU Institute for Data Science and Informatics, University of Missouri, Columbia, MO, USA

[3]Department of Health Management and Informatics and Christopher S Bond Life Science Center, University of
Missouri, Columbia, MO USA
8

9 *** Correspondence:**
10 Erika M Boerman
11 boermane@health.missouti.edu
12

13 **Keywords: perivascular sensory nerves, vascular inflammation, vasomotor function**

1. Abstract

Introduction: Inflammatory bowel disease (IBD) involves aberrant immune responses and is associated with both cardiovascular disease risk and altered intestinal blood flow. However, little is known about how IBD affects regulation of perivascular nerves that mediate blood flow. Previous work found perivascular nerve function is impaired in mesenteric arteries with IBD. The purpose of this study was to determine the mechanism of impaired perivascular nerve function.

Methods: RNA sequencing was performed on mesenteric arteries from IL10$^{-/-}$ mice treated with *H.hepaticus* to induce disease (IBD) or left non-gavaged (Control). For all other studies, Control and IBD mice received either saline or clodronate liposome injections to study the effect of macrophage depletion. Perivascular nerve function was assessed using pressure myography and electrical field stimulation. Fluorescent immunolabeling was used to label leukocyte populations and perivascular nerves.

Results: IBD was associated with increased in macrophage-associated gene expression, and immunolabeling showed accumulation of adventitial macrophages. Clodronate liposome injection eliminated adventitial macrophages, which reversed significant attenuation of sensory vasodilation, sympathetic vasoconstriction and sensory inhibition of sympathetic constriction in IBD. Acetylcholine-mediated dilation was impaired in IBD and restored after macrophage depletion, but sensory dilation remained nitric oxide independent regardless of disease and/or macrophage presence.

Conclusion: Altered neuro-immune signaling between macrophages and perivascular nerves in the arterial adventitia contributes to impaired vasodilation, particularly via dilatory sensory nerves. Targeting the adventitial macrophage population may help preserve intestinal blood flow in IBD patients.

2. Introduction

Inflammatory Bowel Diseases, such as Crohn's disease and ulcerative colitis, affect >3 million adults in the US, and prevalence increased 123% from 2007-2017(1). IBD is marked by chronic aberrant immune responses and both

intestinal and extraintestinal inflammation. Patients experience increased risk and incidence of cardiovascular

diseases including atherosclerosis(2), stroke(3), myocardial infarction(4) and heart failure(5) despite having lower

traditional risk factors (obesity, hypertension, hyperlipidemia, diabetes) than the general population(4). IBD

decreases intestinal perfusion and increases bowel ischemia (6, 7, 8), and blood flow measurements are used in

diagnosis and assessment of relapse risk(9). Collectively, these factors point to an important role for vascular

function in the pathogenesis and treatment of IBD, yet few studies have directly addressed changes in vascular

function with IBD.

Mesenteric arteries (MAs) direct blood to the intestines, and their function is critical to mobilizing ~25%

of cardiac output in and out of the gut(10). The perivascular sympathetic and sensory nerves enmeshed in the

adventitia directly constrict and dilate MAs, respectively, (11) to regulate perfusion. Perivascular sensory nerves

further facilitate dilation via feedback inhibition of sympathetic nerves(12, 13) and local axon reflexes(14). Despite

diminished vasodilator capacity(15) and blood flow(6, 8, 16) with IBD, the function of sensory nerves and their

neuroeffector signaling pathways in IBD is understudied. Pervious work in the lab showed significant defects in

both sensory vasodilation and sensory inhibition of sympathetic vasoconstriction with IBD (17). These deficits

coincide with modest changes in sensory neurotransmitter content, release and receptor function, but the specific

mechanism underlying the near-total loss of sensory vasodilation in IBD remains undefined.

The adventitia that houses perivascular nerves contains a dynamic mix of perivascular nerves, immune

cells, pericytes, and extracellular matrix that participate in paracrine signaling in both normal and pathological

function(18, 19). In healthy adventitia, immune cells are relatively quiescent, and sympathetic and sensory nerve

activity are balanced by reciprocal feedback(12, 13). Many cardiovascular and inflammatory diseases are

accompanied by an increase in the number and/or change in phenotype of adventitial(20) immune cells,

particularly macrophages. When activated by local signals, macrophages polarize into functional subpopulations

across a spectrum including classically activated M1-like, and alternatively activated M2-like phenotypes(21). In

IBD, macrophage accumulation and subtype disequilibrium in the colon wall contributes to disease pathogenesis

and severity(22, 23) where decreased blood flow is evident, but no studies have investigated adventitial

macrophages with respect to vascular function or blood flow in IBD. The role of perivascular sensory nerves in

these processes is also unclear. Using RNA sequencing, this study identified increased macrophage markers in

66   arteries with IBD and therefore tested whether macrophages and other leukocytes accumulate in the mesenteric

67   artery adventitia in the presence of established IBD. Given our previous findings of perivascular nerve

68   dysfunction (17), we therefore tested whether and how whole-body macrophage depletion improves perivascular

69   nerve function.

3. Materials and Methods

2.1 Animals and Artery Preparation

Male and female IL-10$^{-/-}$ mice (4–6 mo), originally obtained from the Jackson Laboratory (B6.129P2-IL-

10tm1Cgn/J, Strain #002251, RRID:IMSR_JAX:002251) were bred and housed at the University of Missouri in a

12:12 light:dark cycle and received standard chow and water ad libitum. At 2 and 4 days post-weaning, mice in

the IBD group were administered *H. hepaticus* by oral gavage as previously described(17) and developed disease

for 90 days as shown in **Figure 1A**. For studies using macrophage depletion, Control and IBD groups received

intraperitoneal injections of PBS- or clodronate liposomes (100 µl, 50 µl and 50 µl of 5mg/ml stock in PBS) at

days 111, 118 and 125, respectively, as shown in **Figure 1B**. For all experiments, mice were anesthetized via

intraperitoneal injection of ketamine-xylazine (25 mg/mL/2.5 mg/mL) followed by tissue dissection and

euthanasia via cardiac exsanguination. All experiments were performed in compliance with the Guide for the Care

and Use of Laboratory Animals (24) and were approved by the University of Missouri Animal Care and Use

Committee. Inflammation of the colon was determined post-hoc via blind analysis in H&E-stained intestine

sections using a previously-described scoring system (17, 25) ranging from 0-24. For all experiments, first or

second-order mesenteric arteries (MAs, 100-200 µm inner diameter) from the terminal ileum were isolated by

hand dissection in Ca^{2+} free physiological salt solution (PSS) [in mM: 140 NaCl, 5 KCl, 1 MgCl2, 10 HEPES, 10

glucose] at 4⁰C.

2.2 RNA sequencing

First and second-order mesenteric arteries from each mouse were pooled and placed into lysis buffer from the

RNAqueous Micro Kit with DNase from Ambion Inc. (Austin, TX, USA), followed by RNA extraction and

DNase treatment per manufacturer's guidelines. All samples were provided to the DNA Core at the University of

Missouri for RNA sequencing. Libraries were prepared using the TruSeq Stranded mRNA Library Kit according

to the manufacturer's instructions, with quantity and quality verified using the High Sensitivity NGS Fragment

Analysis Kit (Agilent Technologies). Libraries were pooled in equimolar amounts and sequenced with single end

75 base pair reads using an Illumina NextSeq 550. Resulting Bcl files were converted to Fastq format using

Illumina software, and trimming was performed using Trimomatic (26) with a threshold of 36. Reads were

aligned to the GRCm38.p5 Mus musculus annotated Genome assembly from Ensembl. Transcript abundance

(expressed in FPKM) was generated using the Cufflinks tool from the Tuxedo suite (27) as well as genome

annotation files downloaded from the same source. Differential gene expression analysis was performed using the

Cuffdiff tool(27). Differentially expressed transcripts were defined as those that had a fold change with an

adjusted $P < 0.05$ via Wald test and by Benjamini–Hochberg correction. Clustering, principal component analysis

(PCA) and heatmap generation were performed on $\ln(x)$-transformed data using ClustVis software(28). The

heatmap was generated using row centering via unit variance scaling. Rows and columns were clustered using

correlation distance and average linkage. Principal component analysis used singular value decomposition with

imputation. Relevant GO terms were determined based on the list of differentially expressed genes using the GO

Enrichment Analyzer web tool (http://geneontology.org/).

2.3 Pressure Myography

Unbranched Second-order arteries 500-1000 µm in length were transferred to a cannulation chamber with

platinum electrodes (Warner Instruments), cannulated onto glass pipettes, and tied with two 11-0 sutures on each

end. After transferring the rig to microscope stage, arteries were pressurized to 70 mmHg and heated to 37°C. For

all functional studies, arteries were superfused with PSS (above) containing 2 mM calcium. Images were acquired

on an inverted Olympus BX61W1 microscope using a 10X objective and a CCD camera (Basler, Inc) coupled to

edge-tracking software (LabView, National Instruments). Perivascular nerves were stimulated in an electrical

field as previously described (17). Briefly, electrical pulses (70 V, 2 ms) were delivered using a Grass S88

stimulator and stimulus isolation unit at 1-16 Hz until a stable diameter was achieved (<30 s). Sympathetic

constrictions were measured in the presence and absence of capsaicin (10 μM), which blocks sensory nerves.

Sensory vasodilation was measured in vessels preconstricted with phenylephrine (1 μM) and the sympathetic

nerve blocker guanethidine (10 μM). To determine the endothelium dependence of sensory vasodilation,

stimulations were repeated in the presence of the nitric oxide inhibitor N(G)-Nitro-L-arginine methyl ester (L-

NAME, 100 μM). In all experiments, maximum diameters were determined by incubating vessels in Ca^{2+} free

PSS containing 10 μM sodium nitroprusside.

2.4 Immunofluorescence

Isolated mesenteric arteries cleared of perivascular adipose were pinned in a Sylgard-coated 24-well plate for

immunolabeling. Arteries were fixed in 4% paraformaldehyde, blocked and permeabilized with phosphate

buffered saline (PBS) containing 1% bovine serum albumin and 0.1% Triton X-100, and incubated overnight in

primary antibodies (details in **Table 1**). Immune cells were labeled as follows: leukocytes with CD45, M1-like

macrophages with CD68, M2-like macrophages with CD163, antigen presenting cells with MCHII. Perivascular

nerves were labeled as follows: all nerves with PGP 9.5, sympathetic nerves with tyrosine hydroxylase, sensory

nerves with CGRP and substance P. Tissues were then blocked again, incubated in secondary antibodies, and

mounted on slides using ProLong Gold with or without DAPI. Each experiment included no primary controls that

demonstrated a lack of nonspecific fluorescence. All slides were imaged on a Leica TCS SP8 laser scanning

confocal microscope at 25X with 1 μm z-sections. Each fluorophore was imaged sequentially using a separate

laser line for excitation to prevent cross-excitation. Representative images were prepared by generating maximum

z-projections through one vessel wall. Immune cells were counted manually post-hoc. Nerve density was

quantified from maximum projections in Image J (29) using a previously-described procedure (17) to measure the

fluorescence area fraction of each antibody.

2.5 Chemicals, Reagents and Tools

All key reagents, chemicals and drugs are listed in **Table 2** with their vendor, catalog number, and

concentration(s) used. As described above, antibodies are outlined in **Table 1**.

 2.6 Statistical Analysis

All data are reported as means ± SEM, with significance at $P < 0.05$. n values represent the number of biological

replicates per group used in each experiment unless otherwise noted. Statistical analysis was completed using

Graphpad Prism 9. All concentration-response and electrical field stimulation data were analyzed by two-way

repeated measures ANOVA followed by Bonferroni post hoc test. Disease scoring, colon length and body weights

were analyzed by unpaired t tests. Fluorescence imaging analysis for immune cell counts and nerve density were

analyzed via nested t-test to account for the use of multiple images from each biological replicate.

4. Results

4.1 Increased expression of macrophage marker genes in mesenteric arteries with IBD

RNA sequencing of mesenteric arteries identified a total of 228 genes with significantly differential expression in

IBD vs Control, 152 with increased expression (**Figure 2A, 3A, Supplemental Table 1**) and 76 with decreased

expression (**Figure 2B, 3A**). Hierarchical heatmap creation also showed that individual Control and IBD samples

segregated together based on differential gene expression patterns (**Figure 2A**). Principal component analysis

determined that PC1 and PC2 combined account for 73.9% of variability in gene expression. PC1 and PC2 were

also used to look for any apparent patterns that clearly differentiate males from females in this group of genes,

with none uncovered. The top 30 genes with upregulated expression in arteries from IBD mice included a number

of macrophage marker genes including *cd163* (CD163), *cd68*, *itgam* (CD11b), *adgre1* (F4/80), *csf1r* (CD115) and

mrc1 (CD206). Additional genes found in the top 30 are not macrophage markers but associated with macrophage

function: *ms4a4a*, *ccl9*, *siglec1* (CD169), *clec10a* (CD301). All significantly up- and downregulated genes are

shown in in **Figure 2**, and the complete list of differentially expressed genes with fold changes, P and Q values is

in **Supplemental Table 1.** Based on these data, we conducted additional experiments looking at changes in the

adventitial macrophage population in IBD and its role in perivascular nerve function/dysfunction.

4.2 IBD induction and macrophage depletion in IL10^{-/-} mice

In the present study, all mice were treated with either PBS liposomes (macrophages intact) or clodronate

liposomes (macrophages depleted). In mice treated with PBS liposomes, consistent with our previous studies[17],

IBD induction with *H. hepaticus* gavage led to a significant increase in colon inflammation scores (**Figure 4**).

There was a non-significant trend toward decreased body weight, and no changes in colon weight or length 90

days post-gavage. Clodronate liposome treatment was associated with decreased body weight in Control and IBD

groups, with a significantly greater decrease in IBD-CLOD mice (**Figure 4**). Clodronate decreased colon weight

to a similar extent in Control and IBD groups compared to PBS treatment, but it had no effect on colon length or

inflammation scores in either group compared to PBS treatment (**Figure 4**).

4.3 IBD increases, clodronate decreases adventitial immune cells

To characterize the adventitial leukocyte and macrophage populations in Control and IBD arteries with and

without macrophage depletion, we conducted two sets of confocal immunolabeling studies counting the number

of cell labeled for: CD45 (all leukocytes), MCHII (antigen presenting cells), CD68 (M1-like macrophages) and

CD163 (M2-like macrophages). All markers were significantly increased in IBD-PBS arteries vs CONT-PBS

arteries (**Figures 5-6**). Macrophage depletion with clodronate led to significant attenuation of CD45+ cells in both

CONT-CLOD and IBD-CLOD groups, although significantly more CD45+ cells persisted post-clodronate in

CONT-CLOD. Clodronate treatment resulted in near-total elimination of MHCII+, CD68+ and CD163+ cells in

both CONT-CLOD and IBD-CLOD groups, indicating successful depletion of macrophages in the artery

adventitia (**Figures 5-6**). This is consistent with control flow cytometry experiments showing similar levels of

macrophage depletion in the spleens of the same experimental mice (data not shown).

4.4 Macrophage depletion improves perivascular nerve dysfunction in mesenteric arteries with IBD

4.4.1 Macrophage depletion restores impaired sympathetic vasoconstriction and sensory inhibition of sympathetic vasodilation with IBD

Pressure myography and electrical field stimulation were used to assess sympathetic vasoconstriction in the presence (PSS) and absence (capsaicin) of sensory neurotransmission in CONT-PBS, CONT-CLOD, IBD-PBS and IBD-CLOD groups. Consistent with previous findings, sympathetic constriction was decreased in mesenteric arteries from IBD-PBS vs CONT-PBS mice. Despite impaired sympathetic constriction with IBD, the presence of sensory neurotransmission failed to decrease sympathetic vasoconstriction to even a small degree (**Figure 7A,C**). Macrophage depletion had opposing effects in Control and IBD groups. CONT-CLOD arteries exhibited decreased sympathetic vasoconstriction compared to CONT-PBS while retaining the ability of sensory nerves to oppose sympathetic vasoconstriction (i.e., capsaicin still increased sympathetic constriction in CONT-CLOD arteries) (**Figure 7A,B**). In contrast, macrophage depletion increased sympathetic constriction in IBD-CLOD vs IBD-PBS (**Figure 7C,D**), restoring them to CONT-PBS levels. Similarly, macrophage depletion restored sensory inhibition of sympathetic inhibition (**Figure 7C,D**), with IBD-CLOD constrictions in PSS and capsaicin becoming indistinguishable from CONT-PBS (**Figure 7A,D**).

4.4.2 Macrophage depletion restores impaired sensory vasodilation in mesenteric arteries with IBD

Consistent with previous studies in our group (17), we found that IBD was associated with significantly attenuated perivascular sensory vasodilation, with a ~75% decrease in maximum achievable vasodilation in IBD-PBS vs CONT-PBS arteries (**Figure 8**). Macrophage depletion did not alter sensory vasodilation in CONT-CLOD arteries. In contrast, macrophage depletion increased sensory vasodilation in IBD-CLOD arteries, restoring dilations to Control levels.

4.4.3 Macrophage depletion does not alter the endothelium-independence of sensory vasodilation

Because sensory vasodilation has been reported to include both endothelial dependent and independent mechanisms, we repeated sensory vasodilation measurements in isolated arteries before (PSS) and after

incubation in the nitric oxide synthase inhibitor L-NAME (100 µM) (**Figure 9**). Although some significant

differences in vasodilation were observed at specific stimulation frequencies, statistical analysis revealed that L-

NAME treatment was not a significant source of variation in sensory vasodilation arteries from CONT-PBS,

CONT-CLOD, IBD-PBS or IBD-CLOD groups.

4.5 Macrophage depletion improves endothelium-mediated vasodilation in mesenteric arteries with IBD

Because our previous studies found that IBD desensitized endothelium-dependent vasodilation of mesenteric

arteries to acetylcholine (Ach) (17), we tested whether macrophage depletion affected ACh-mediated

vasodilation. Consistent with the previous study, mesenteric arteries from IBD-PBS mice displayed a significant

decrease in maximum ACh dilation and sensitivity (mean EC50s $10^{-7.1}$ µM vs $10^{-6.9}$ µM, respectively) compared to

CONT-PBS mice (**Figure 10**). Macrophage depletion did not alter ACh concentration-response curves in arteries

CONT-CLOD vs CONT-PBS mice. In contrast, macrophage depletion increased ACh potency ad sensitivity in

IBD-CLOD vs IBD-PBS mice (**Figure 10**).

4.6 Effect of macrophage depletion on perivascular nerve density

Because macrophage depletion was associated with significant improvements in sympathetic vasodilation and

sensory vasodilation, we used quantitative immunofluorescence to test whether clodronate treatment altered

perivascular nerve density. Analysis revealed no differences in total perivascular nerve density (PGP9.5+) of

sympathetic nerve density (tyrosine hydroxylase, TH+) between CONT-PBS, CONT-CLOD, IBD-PBS and IBD-

CLOD groups (**Figure 11A,B**). Sensory nerve density, measured with both CGRP+ and SP+ labelling was not

different in CONT-PBS vs IBD-PBS arteries (**Figure 11C-D**). Macrophage depletion did not alter CGRP+ or SP+

labeling in CONT-PBS vs CONT-CLOD arteries or IBD-PBS vs IBD-CLOD arteries. However, both CGRP+ and

SP+ labeling was significantly increased in IBD-CLOD vs CONT-PBS arteries (**Figure 11C-D**).

5. Discussion

5.1 Summary

The present study used an established IL10$^{-/-}$ + *H. hepaticus* mouse model of IBD to define the mechanisms by which perivascular nerve function in mesenteric arteries is significantly impaired with IBD, likely contributing to the restricted intestinal blood flow often observed in patients. RNA sequencing and analysis of differentially expressed genes from IBD vs Control arteries identified adventitial leukocytes and, more specifically, macrophages as a potential driver of perivascular nerve dysfunction in mesenteric arteries with IBD. Quantitative analysis of confocally-imaged immunolabeled arteries showed that leukocytes, including a significant population of CD68+ and CD163+ macrophages, accumulate in the arterial adventitia near perivascular nerves with IBD. Further, this macrophage population was successfully eliminated following intraperitoneal injection of clodronate liposomes. Next, clodronate or PBS liposome treatment was utilized to determine the effect of macrophage depletion in perivascular nerve function in isolated mesenteric arteries. Using pressure myography and electrical field stimulation to define sympathetic and sensory nerve function, our key findings are that elimination of adventitial macrophages was associated with restoration of (1) sensory vasodilation, (2) sensory inhibition of sympathetic vasoconstriction, and (3) sympathetic vasoconstriction. Sensory vasodilation in this model and its improvements following macrophage depletion appear to be mostly endothelium independent, and they also occur in the absence of major changes in nerve density. Based on the current data, we suggest that impaired sensory vasodilation in mesenteric arteries during IBD occurs due to aberrant adventitial neuro-immune signaling between perivascular nerves and the nearby expanded macrophage population.

5.2 Macrophage accumulation in the vascular adventitia with IBD

Adventitial macrophage accumulation and/or polarization has been described in both conduit and resistance arteries from a wide variety of diseases, including autoimmune diseases (see review (30)), but this is the first study to measure the adventitial macrophage population with IBD or investigate its effect on perivascular nerve function in IBD. RNA sequencing of intact mesenteric arteries identified multiple macrophage markers and macrophage-associated genes among the genes with the most increased expression with IBD. Examination of GO

terms associated with differentially expressed genes also pointed to leukocytes, macrophages and associated

inflammation as a potential target in vascular dysfunction with IBD (**Figure 2-3, Supplemental Table 1**). Based

on this data, and because macrophages have been identified as the most abundant cells in the adventitia in health

and disease (31), we pursued macrophage infiltration as a potential cause of perivascular nerve dysfunction. In

RNA sequencing, IBD- upregulated genes included pan-macrophage markers, genes associated with both

classical (M1-like) and alternative activation (M2-like), and genes associated with macrophage chemotaxis.

Upregulated pan macrophage markers included *adgre1, itgam* and *csf1r,* which have each been identified as

putative macrophage markers in the vascular adventitia (31). Upregulated genes associated with an M1-like

phenotype included *ccl9*, *ccl5* and *cxcl9* (32). Immunolabeling for MHCII and CD68 further indicated an increase

in total and M1-like macrophages (**Figure 5-6**). Upregulated genes associated with M2-like phenotype included

cd163, *mrc1, ms4a4a* (33), with immunolabeling for CD163 also increased in IBD (**Figure 5-6**). Collectively,

these data support a broad expansion of arterial macrophages of multiple phenotypes in IBD. Although a more

nuanced characterization of adventitial macrophage phenotype is relevant to both IBD and vascular inflammation,

the major purpose of the present study was to functionally define the relationship between the overall adventitial

macrophage population and perivascular nerve function in IBD. An important focus for ongoing work will be to

determine whether macrophage expansion in the adventitia with IBD represents expansion of the resident

population or recruitment of new monocyte-derived macrophages, as both populations are present in the healthy

adventitia (31). In support of expanded resident macrophage population, we found that many genes with increased

expression in arteries with IBD (*ms4a4a, ms4a6c, il1r1, cd36,* **Figures 2-3**, **Supplemental Table 1***)* overlap with

genes enriched in peripheral nervous system (PNS) macrophages, (34), which are self-renewing. The overlap

between the gene signatures of IBD adventitia and PNS macrophages is also interesting because both macrophage

populations reside in close proximity to a nerve microenvironment. Supporting an expanded population of bone

marrow derived macrophages, we found increased expression of *s100a9* in mesenteric arteries, which is

associated with early monocyte to macrophage differentiation (31).

5.3 Adventitial nerve-macrophage signaling in IBD

Consistent with previous work in our laboratory (17), this study found that IBD was associated with drastically

impaired sensory vasodilation (**Figure 8A,C**) and impaired inhibition of sympathetic vasoconstriction (**Figure**

7A,C), despite depressed sympathetic vasoconstriction in IBD (**Figure 8A,C**). Surprisingly, all three vasomotor

functions were completely restored following macrophage depletion with clodronate liposomes (**Figures 7-8**).

These data highlight that the net result of adventitial macrophage signaling across all phenotypes is impaired

perivascular nerve function. Most work on neuro-immune interactions within and outside the vasculature has

focused on sympathetic nerves, and the positive correlation between sympathetic outflow and inflammation

throughout the body is well-defined in many diseases. In IBD, studies in human intestine have demonstrated a

correlation between increased MHCII+ immune cell presence and decreased enteric nerves function (35) and

density (36). Counterintuitively, our study found that IBD was associated with increased adventitial macrophages

(**Figure 5-6**) but decreased sympathetic vasoconstriction, which was restored following macrophage depletion

(**Figure 7**); sympathetic nerve density remained unchanged (**Figure 11**). Few studies have investigated the

macrophage-nerve relationship in the mesenteric vasculature. Studies in a rat model of hypertension found that

CD163+ macrophage accumulation in the mesenteric artery adventitia is linked to increased blood pressure

through macrophage-derived oxidate stress acting on ganglionic neuronal alpha-2 adrenergic receptors (37). In

that study, clodronate-based macrophage depletion acted to lower blood pressure by preserving neuronal alpha-2

receptor function. Here, increased CD163+ macrophage population in the mesenteric artery adventitia (**Figures**

5-6) is associate with decreased sympathetic nerve function, suggesting that macrophages may mediate vascular

dysfunction in distinct ways during IBD versus hypertension and other cardiovascular diseases.

Macrophage accumulation appears to have a much greater and more detrimental effect on perivascular

sensory nerves compared to sympathetic nerves. In earlier studies, we found that despite impaired sensory

vasodilation with IBD, there was not a significant decrease in the content or release of the sensory

neurotransmitters CGRP and substance P. Similarly, the function of CGRP receptors, the primary vasodilatory

peptide, released was unaltered with IBD. Norepinephrine sensitivity was also unchanged with IBD (17).

Considering the seemingly preserved neurotransmission and receptor function with unchanged nerve density

(**Figure 11**), what is happening to the released neurotransmitter? We propose that rather than acting downstream

to affect vasodilation, released CGRP and substance P bind to receptors expressed on the expanded population of

closely-approximated macrophages. Thus, sensory transmitter released from nerves would mediate immune-

mediated effects in place of the vasodilation observed in arteries with a healthy adventitial macrophage

population. This possibility is supported by multiple areas of research. Both CGRP and substance P can affect

macrophage phenotype and the release of inflammatory mediators in vitro (38, 39, 40), although the effects of

each neuropeptide on perivascular macrophages is unknown. Local macrophage activation by neuronally released

neurotransmitter was recently defined in the intestinal wall. In a series of experiments, Gabanyi et al (41)

characterized a population of intestinal macrophages expressing β2 adrenergic receptors that are uniquely located

near nerves and activated by norepinephrine released from extrinsic nerves to induce M2 polarization. This

concept is consistent with our findings of depressed sympathetic vasoconstriction (**Figure 5**) and increased gene

and protein indicators of M2 polarization (**Figures 2,3,5,6**) with IBD. The gene expression profiling from the

Gabanyi study further found that the CGRP receptor protein RAMP1 (receptor activity modifying protein 1) was

the only other neurotransmitter receptor with increased expression in the muscularis macrophage population,

although follow-up imaging studies were not performed. Given that perivascular nerves on mesenteric arteries

extend into the intestinal wall (42), they may similarly affect the local macrophage population that express

neurotransmitter receptors.

5.4 Role of endothelium and nitric oxide in perivascular nerve dysfunction

The role of the endothelium in both sensory vasodilation and IBD pathogenesis is unclear and somewhat

controversial. In both health and disease, sensory vasodilation has been described as mostly endothelium

dependent (43, 44), partially endothelium dependent (45, 46) and endothelium independent (47, 48). Previous

work in mouse mesenteric arteries supports partial endothelial dependence as well as increased endothelial

dependence with aging (46), suggesting that the mechanism of sensory vasodilation in mesenteric arteries can

change based on pathology. In the present study, we found that blocking nitric oxide synthase with L-NAME had

a negligible effect on sensory vasodilation in vessels from Control and IBD mice with intact macrophages

(**Figure 9 A,C**). L-NAME also had no effect on sensory vasodilation in Control or IBD mice after macrophage

depletion (**Figure 9 B,D**), suggesting that nitric oxide generation is not an important target for restoring sensory

vasodilation in this model. Independent of sensory nerve function, endothelial dysfunction of varying severity and

mechanism has been demonstrated in murine (49) and human (50) studies of IBD. Here, we found a small but

significant decrease in acetylcholine-mediated dilation in mesenteric arteries with IBD (**Figure 10**). Macrophage

depletion restored acetylcholine responses in arteries from IBD mice, suggesting that macrophage-mediated

endothelial dysfunction contributes to vascular dysfunction in IBD independent of perivascular nerve function.

The presence of some endothelial dysfunction in this model is consistent with the IBD-related increase in

expression of the chemokine *ccl5* and its receptor *ccr1*, which are linked to endothelial dysfunction and leukocyte

transmigration into arteries (51). Collectively, our data suggest that "inside-out" vascular dysfunction arising from

the endothelium is less significant in IBD than "outside-in" mechanisms initiated within the adventitia.

5.5 Limitations

This study clearly demonstrates the presence of significant macrophage-associated perivascular nerve

dysfunction, but there are also limitations to the study that warrant further discussion. It is important to note that

no mouse model of IBD can currently recapitulate all aspects of human IBD. The IL10$^{-/-}$ model is widely used and

accepted (52) but has limiting factors. The use of the IL10$^{-/-}$ model may affect the immune responses observed, as

immune cells cannot produce and respond to IL10 in the same way as human IBD patients. In addition, the

severity of IBD in IL10$^{-/-}$ is quite variable across studies, and as disease severity varies with microbiome

differences arising from mouse strain, vendor and specific housing conditions (53). Clodronate studies also have

limitations. The use of clodronate to deplete macrophages may have indeterminate effects on vascular function

independent of the elimination of adventitial macrophages, as it depletes macrophages throughout the body. We

also halve doses 2 and 3 of clodronate, as repeating the original dose led to ~25% mortality in the control group

(consistent with published studies (54, 55)) and up to 50% mortality in the IBD group. No studies using

clodronate in IL10$^{-/-}$ mice have reported mortality. Macrophage phenotyping is also a limitation in this study, as

the leukocyte phenotyping used for immunofluorescence was relatively rudimentary. While it effectively shows

accumulations of very broad populations of cells, it remains a non-definitive classification of macrophage

subtypes. Follow-up studies with flow cytometry and/or single cell sequencing will be needed to more rigorously

define the adventitial immune cell population with IBD.

Data availability statement

369   The original contributions presented in the study are included in the article/supplementary material, further

370   inquiries can be directed to the corresponding author.

Author contributions

373   EB conceived and designed the experiments. EB, EG and BJ performed the experiments. EB, BJ, TJ, KW and SS

374   analyzed the data. All authors approved the final version of the manuscript.

Funding

Conflict of interest

380   The authors declare that the research was conducted in the absence of any commercial or financial relationships

381   that could be considered a potential conflict of interest.

383 **Table 1. Primary antibody information**

Target Antigen	Species	Vendor	Catalog Number	Concentration	RRID
Calcitonin gene-related peptide (CGRP)	rabbit	MilliporeSigma	PC205L	1:250	AB_2068524
Substance P (SP)	mouse	Abcam	ab14184	1:250	AB_300971
Tyrosine hydroxylase (TH)	mouse	MilliporeSigma	T2928	1:250	AB_477569
Protein Gene Product 9.5 (PGP 9.5)	rabbit	ThermoFisher	PA5-32567	1:500	AB_2550032
CD45	rat	Biorad	MCA1388	1:250	AB_321729
CD68	mouse	ThermoFisher	14-0681-82	1:250	AB_2572857
CD163	rabbit	Bioss	Bs-2527R	1:250	AB_10856166
Major Histocompatibility Complex class II (MHCII)	mouse	ThermoFisher	14-5321-85	1:250	AB_467561

384 **Table 2. Drug information**

Name	Vendor	Catalog Number	Concentration	Vehicle
Phenylephrine hydrochloride (PE)	MilliporeSigma	P6126	1 µM	Water
Guanethidine sulfate (GE)	BOC Sciences	645-43-2	10 µM	Water
Sodium nitroferricyanide (nitroprusside, SNP)	MilliporeSigma	228710	10 µM	Water
Capsaicin (CAP)	MilliporeSigma	M2028	10 µM	Ethanol
Nω-Nitro-L-arginine methyl ester hydrochloride (L-NAME)	MilliporeSigma	N5751	100 µM	Water
Clodronate liposomes	Liposoma	C-005	5mg/ml stock	PBS
PBS liposomes	Liposoma	P-005	--	PBS

Figure 1. Experimental IBD model and timelines for experiments. Figures depict experimental timelines for

studies A) without macrophage depletion and B) with macrophage depletion via clodronate liposome injection.

Note that in all experiments, Control animals are non-gavaged $IL10^{-/-}$ mice, while IBD groups are $IL10^{-/-}$ mice

gavaged with *H. hepaticus* after weaning.

Figure 2. Summary of up- and downregulated genes in MAs from IBD vs Control mice. Data are fold

changes in expression for IBD vs Control showing only genes with Q values < 0.05. RNA sequencing identified

A) Genes with significantly increased expression and B) Genes with significantly decreased expression in

mesenteric arteries from IBD vs Control mice. A complete table of fold-change expression, P- and Q-values for

each differentially expressed gene can be found in Supplementary Table 1.

Figure 3. Heatmap, principal component analysis, and GO enrichment of differentially expressed genes

from RNA sequencing of mesenteric arteries. A) Clustered heatmap of differentially expressed genes in

mesenteric arteries from Control and IBD mice. Red colors indicate genes with increased expression in IBD vs

Control, while blue colors represent genes with decreased expression. All differences have P < 0.05. B) Principal

component analysis of differentially expressed genes from Control and IBD mice. C) Top 20 significantly

enriched GO biological processes in mesenteric arteries from IBD vs Control mice. All data originated from RNA

sequencing of N=5 samples each from Control and IBD groups.

Figure 4. IBD and macrophage depletion affect body weight, colon size and colon inflammation. Data are

mean ± SE for A) body weight, B) colon weight, C) colon length and D) histological disease score of H&E

stained colon sections in Control (CONT) and IBD mice treated with either PBS or clodronate (CLOD)

liposomes. N=4-8 mice per measurement in each group. * = P<0.05 vs CONT-PBS; # = P<0.05 vs IBD-PBS.

Figure 5. IBD increases and clodronate decreases adventitial macrophages and leukocytes. Images are

representative maximum z-projections through one artery wall showing nuclei leukocyte, M1-like macrophages

and perivascular nerves in mesenteric arteries in each of 4 groups: Control-PBS, Control-CLOD, IBD-PBS and

IBD-CLOD. Panels for each group depict two separate image sets- Top row (left to right): Nuclei (DAPI, blue),

leukocytes (CD45, magenta), M1-like macrophages (CD68, green), perivascular nerves (PGP9.5, red), merged

image; Bottom row (left to right): nuclei (DAPI, blue), antigen presenting cells (MHC II, red), M2-like

macrophages (CD163, green), merged image. Images are 185 x 185 µm.

Figure 6. Quantitation of immunofluorescence for adventitial immune cells. Data are mean ± SE number of

leukocytes (CD45+), antigen-presenting cells (MHCII+), M1-like macrophages (CD68+) and M2-like

macrophages (CD68+) per field of view (FOV, 185 x 185 µm) of confocal z-projections through the mesenteric

artery adventitia of vessels from Control and IBD mice treated with PBS or clodronate (CLOD) liposomes. N=12-

20 images from 4 mice per group. * = $P<0.05$ vs Control-PBS; # = $P<0.05$ vs IBD-PBS.

Figure 7. Macrophage depletion improves sensory inhibition of sympathetic vasodilation. Data are mean ±

SE for %constrictions to electrical field stimulation (1-16 Hz) before (PSS) and after sensory nerve blockade with

capsaicin (CAP, 10 µM) in A) CONT-PBS: Control mice treated with PBS liposomes, B) CONT-CLOD: Control

mice treated with clodronate liposomes, C) IBD-PBS: IBD mice treated with PBS liposomes, and D) IBD-CLOD:

IBD mice treated with clodronate liposomes. Dashed line indicates the max sympathetic constriction reached in

CONT-PBS arteries in PSS. N = 4-5 per group. * = $P< 0.05$ vs PSS

Figure 8. Macrophage depletion restores sensory vasodilation in mesenteric arteries with IBD. Data are

mean ± SE % maximum dilations elicited by sensory nerves via electrical field stimulation (1-16 Hz) in arteries

from Control (blue) and IBD (red) mice treated with PBS liposomes (solid lines) or clodronate liposomes (dashed

lines). N=4-5 per group. * = $P<0.05$ vs IBD-PBS.

Figure 9. Sensory vasodilation is mostly endothelium-independent and unaffected by macrophage

depletion. Data mean ± SE % maximum dilations elicited by sensory nerves via electrical field stimulation (1-16

Hz) in arteries before (PSS) and after incubation in the nitric oxide synthase inhibitor L-NAME (100 μ) from A)

CONT-PBS: Control mice treated with PBS liposomes, B) CONT-CLOD: Control mice treated with clodronate

liposomes, C) IBD-PBS: IBD mice treated with PBS liposomes, and D) IBD-CLOD: IBD mice treated with

clodronate liposomes. Dashed line indicates the max sensory vasodilation reached in CONT-PBS arteries in PSS

N=4-5 per group. * = $P < 0.05$ vs equivalent stimulation in PSS.

Figure 10. Macrophage depletion improves attenuated ACh dilation in mesenteric arteries with IBD. Data

are Data mean ± SE % maximum dilations to increasing concentrations of Acetylcholine (ACh) in preconstricted

(phenylephrine, 10μM) mesenteric arteries from Control (blue) and IBD (red) mice treated with PBS liposomes

(solid lines) or clodronate liposomes (dashed lines). N=4-5 per group. * = $P < 0.05$ vs IBD-PBS.

Figure 11. Macrophages depletion increases fluorescence area of sensory neurotransmitters but not the

total perivascular nerves or perivascular sympathetic nerves. Data are mean ± SE fluorescence area A) all

perivascular nerves (PGP 9.5), B) perivascular sympathetic nerves (tyrosine hydroxylase, TH), and C-D) sensory

neurotransmitters calcitonin gene related peptide (CGRP) and substance P (SP) in of confocal z-projections

through the mesenteric artery adventitia of vessels from Control (blue, circles markers) and IBD (red, squares)

mice treated with PBS (open bars), or clodronate (CLOD, striped bars) liposomes. N = 10-25 images from 3-4

mice per group.* = $P < 0.05$ vs CONT-PBS.

References

1.   Ye Y, Manne S, Treem WR, Bennett D. Prevalence of Inflammatory Bowel Disease in Pediatric and
Adult Populations: Recent Estimates From Large National Databases in the United States, 2007-2016. Inflamm
Bowel Dis. 2020;26(4):619-25.

2.   Papa A, Danese S, Urgesi R, Grillo A, Guglielmo S, Roberto I, et al. Early atherosclerosis in patients with
inflammatory bowel disease. Eur Rev Med Pharmacol Sci. 2006;10(1):7-11.

3.   Miehsler W, Reinisch W, Valic E, Osterode W, Tillinger W, Feichtenschlager T, et al. Is inflammatory
bowel disease an independent and disease specific risk factor for thromboembolism? Gut. 2004;53(4):542-8.

4.   Aniwan S, Pardi DS, Tremaine WJ, Loftus EV, Jr. Increased Risk of Acute Myocardial Infarction and
Heart Failure in Patients With Inflammatory Bowel Diseases. Clin Gastroenterol Hepatol. 2018;16(10):1607-15
e1.

5.   Kristensen SL, Ahlehoff O, Lindhardsen J, Erichsen R, Lamberts M, Khalid U, et al. Inflammatory bowel
disease is associated with an increased risk of hospitalization for heart failure: a danish nationwide cohort study.
Circ Heart Fail. 2014;7(5):717-22.

6.   Hultén L, Lindhagen J, Lundgren O, Fasth S, Ahrén C. Regional intestinal blood flow in ulcerative colitis
and Crohn's disease. Gastroenterology. 1977;72(3):388-96.

7.   Harris NR, Carter PR, Lee S, Watts MN, Zhang S, Grisham MB. Association between blood flow and
inflammatory state in a T-cell transfer model of inflammatory bowel disease in mice. Inflamm Bowel Dis.
2010;16(5):776-82.

8.   Thornton M, Solomon MJ. Crohn's disease: in defense of a microvascular aetiology. Int J Colorectal Dis.
2002;17(5):287-97.

9.   Ludwig D, Wiener S, Brüning A, Schwarting K, Jantschek G, Stange EF. Mesenteric blood flow is
related to disease activity and risk of relapse in Crohn's disease: a prospective follow-up study. Am J
Gastroenterol. 1999;94(10):2942-50.

10.   Ralevic V. Splanchnic circulatory physiology. Hepatogastroenterology. 1999;46 Suppl 2:1409-13.

11.   Westcott EB, Segal SS. Perivascular innervation: A multiplicity of roles in vasomotor control and
myoendothelial signaling. Microcirculation. 2013.

12.   Coffa FP, Kotecha N. Modulation of sympathetic nerve activity by perivascular sensory nerves in the
arterioles of the guinea-pig small intestine. J Auton Nerv Syst. 1999;77(2-3):125-32.

13.   Kawasaki H. Regulation of vascular function by perivascular calcitonin gene-related peptide-containing
nerves. Jpn J Pharmacol. 2002;88(1):39-43.

14.   Yaprak M. The Axon Reflex. Neuroanatomy. 2008;7:17-9.

15.   Hatoum OA, Binion DG, Otterson MF, Gutterman DD. Acquired microvascular dysfunction in
inflammatory bowel disease: Loss of nitric oxide-mediated vasodilation. Gastroenterology. 2003;125(1):58-69.

16.   Koch TR. Inflammatory bowel disease activity--the role of blood flow. Am J Gastroenterol.
1998;93(5):841-2.

17.   Norton CE, Grunz-Borgmann EA, Hart ML, Jones BW, Franklin CL, Boerman EM. Role of perivascular
nerve and sensory neurotransmitter dysfunction in inflammatory bowel disease. American journal of physiology
Heart and circulatory physiology. 2021;320(5):H1887-H902.

18.   Strupp M, Mandala M, Lopez-Escamez JA. Peripheral vestibular disorders: an update. Curr Opin Neurol.
2019;32(1):165-73.

19.   Nosalski R, Guzik TJ. Perivascular adipose tissue inflammation in vascular disease. Br J Pharmacol.
2017;174(20):3496-513.

20.   Maiellaro K, Taylor WR. The role of the adventitia in vascular inflammation. Cardiovasc Res.
2007;75(4):640-8.

21.   Mantovani A, Garlanda C, Locati M. Macrophage diversity and polarization in atherosclerosis: a question
of balance. Arterioscler Thromb Vasc Biol. 2009;29(10):1419-23.

22.   Mueller PA, Zhu L, Tavori H, Huynh KT, Giunzioni I, Stafford JM, et al. Response by Mueller et al to
Letter Regarding Article, "Deletion of Macrophage Low-Density Lipoprotein Receptor-Related Protein 1 (LRP1)
Accelerates Atherosclerosis Regression and Increases C-C Chemokine Receptor Type 7 (CCR7) Expression in
Plaque Macrophages". Circulation. 2019;139(16):1983-4.

23.   Seyedizade SS, Afshari K, Bayat S, Rahmani F, Momtaz S, Rezaei N, et al. Current Status of M1 and M2
Macrophages Pathway as Drug Targets for Inflammatory Bowel Disease. Arch Immunol Ther Exp (Warsz).
2020;68(2):10.

24.   Guide for the Care and Use of Laboratory Animals. Council NR, editor: National Academic Press; 1996.
140 p.

25.   Alvarado CG, Kocsis AG, Hart ML, Crim MJ, Myles MH, Franklin CL. Pathogenicity of Helicobacter
ganmani in mice susceptible and resistant to infection with H. hepaticus. Comp Med. 2015;65(1):15-22.

26.   Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data.
Bioinformatics. 2014;30(15):2114-20.

27.   Trapnell C, Roberts A, Goff L, Pertea G, Kim D, Kelley DR, et al. Differential gene and transcript
expression analysis of RNA-seq experiments with TopHat and Cufflinks. Nat Protoc. 2012;7(3):562-78.

28.   Metsalu T, Vilo J. ClustVis: a web tool for visualizing clustering of multivariate data using Principal
Component Analysis and heatmap. Nucleic Acids Res. 2015;43(W1):W566-70.

29.   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an open-source
platform for biological-image analysis. Nat Methods. 2012;9(7):676-82.

30.   Shirai T, Hilhorst M, Harrison DG, Goronzy JJ, Weyand CM. Macrophages in vascular inflammation--
From atherosclerosis to vasculitis. Autoimmunity. 2015;48(3):139-51.

31.   Weinberger T, Esfandyari D, Messerer D, Percin G, Schleifer C, Thaler R, et al. Ontogeny of arterial
macrophages defines their functions in homeostasis and inflammation. Nat Commun. 2020;11(1):4549.

32.   Orecchioni M, Ghosheh Y, Pramod AB, Ley K. Macrophage Polarization: Different Gene Signatures in
M1(LPS+) vs. Classically and M2(LPS-) vs. Alternatively Activated Macrophages. Front Immunol.
2019;10:1084.

33.   Sanyal R, Polyak MJ, Zuccolo J, Puri M, Deng L, Roberts L, et al. MS4A4A: a novel cell surface marker
for M2 macrophages and plasma cells. Immunol Cell Biol. 2017;95(7):611-9.

34.   Wang PL, Yim AKY, Kim KW, Avey D, Czepielewski RS, Colonna M, et al. Peripheral nerve resident
macrophages share tissue-specific programming and features of activated microglia. Nat Commun.
2020;11(1):2552.

35.   Geboes K. Immunopathological studies of the small intestinal intramural nervous system and of
intramural vessels in Crohn's disease. Verh K Acad Geneeskd Belg. 1993;55(4):267-301; discussion -3.

36.   Kyosola K, Penttila O, Salaspuro M. Rectal mucosal adrenergic innervation and enterochromaffin cells in
ulcerative colitis and irritable colon. Scand J Gastroenterol. 1977;12(3):363-7.

37.   Mui RK, Fernandes RN, Garver HG, Van Rooijen N, Galligan JJ. Macrophage-dependent impairment of
alpha(2)-adrenergic autoreceptor inhibition of Ca(2+) channels in sympathetic neurons from DOCA-salt but not
high-fat diet-induced hypertensive rats. American journal of physiology Heart and circulatory physiology.
2018;314(4):H863-H77.

38.   Baliu-Pique M, Jusek G, Holzmann B. Neuroimmunological communication via CGRP promotes the
development of a regulatory phenotype in TLR4-stimulated macrophages. Eur J Immunol. 2014;44(12):3708-16.

39. Liu J, Chen M, Wang X. Calcitonin gene-related peptide inhibits lipopolysaccharide-induced interleukin-
12 release from mouse peritoneal macrophages, mediated by the cAMP pathway. Immunology. 2000;101(1):61-7.

40. Mashaghi A, Marmalidou A, Tehrani M, Grace PM, Pothoulakis C, Dana R. Neuropeptide substance P
and the immune response. Cell Mol Life Sci. 2016;73(22):4249-64.

41. Gabanyi I, Muller PA, Feighery L, Oliveira TY, Costa-Pinto FA, Mucida D. Neuro-immune Interactions
Drive Tissue Programming in Intestinal Macrophages. Cell. 2016;164(3):378-91.

42. Phillips RJ, Powley TL. Innervation of the gastrointestinal tract: patterns of aging. Auton Neurosci.
2007;136(1-2):1-19.

43. Lu R, Zhu HQ, Peng J, Li NS, Li YJ. Endothelium-dependent vasorelaxation and the expression of
calcitonin gene-related peptide in aged rats. Neuropeptides. 2002;36(6):407-12.

44. Moll-Kaufmann C, Sumanovski LT, Sieber CC. Neurally-mediated vasodilatation in normal and portal
hypertensive rats: role of nitric oxide and calcitonin gene-related peptide. J Hepatol. 1998;28(6):1031-6.

45. Sullivan JC, Davison CA. Effect of age on electrical field stimulation (EFS)-induced endothelium-
dependent vasodilation in male and female rats. Cardiovasc Res. 2001;50(1):137-44.

46. Boerman EM, Segal SS. Depressed perivascular sensory innervation of mouse mesenteric arteries with
advanced age. J Physiol. 2016;594(8):2323-38.

47. Edvinsson L, Ahnstedt H, Larsen R, Sheykhzade M. Differential localization and characterization of
functional calcitonin gene-related peptide receptors in human subcutaneous arteries. Acta Physiol (Oxf).
2014;210(4):811-22.

48. Wisskirchen FM, Gray DW, Marshall I. Receptors mediating CGRP-induced relaxation in the rat isolated
thoracic aorta and porcine isolated coronary artery differentiated by h(alpha) CGRP(8-37). Br J Pharmacol.
1999;128(2):283-92.

49. Wu H, Hu T, Zhang L, Xia X, Liu X, Zhu Q, et al. Abdominal Aortic Endothelial Dysfunction Occurs in
Female Mice With Dextran Sodium Sulfate-Induced Chronic Colitis Independently of Reactive Oxygen Species
Formation. Front Cardiovasc Med. 2022;9:871335.

50. Cibor D, Domagala-Rodacka R, Rodacki T, Jurczyszyn A, Mach T, Owczarek D. Endothelial dysfunction
in inflammatory bowel diseases: Pathogenesis, assessment and implications. World J Gastroenterol.
2016;22(3):1067-77.

51. Schober A. Chemokines in vascular dysfunction and remodeling. Arterioscler Thromb Vasc Biol.
2008;28(11):1950-9.

52.   Keubler LM, Buettner M, Hager C, Bleich A. A Multihit Model: Colitis Lessons from the Interleukin-10-
deficient Mouse. Inflamm Bowel Dis. 2015;21(8):1967-75.

53.   Hart ML, Ericsson AC, Franklin CL. Differing Complex Microbiota Alter Disease Severity of the IL-10(-
/-) Mouse Model of Inflammatory Bowel Disease. Front Microbiol. 2017;8:792.

54.   Li Z, Xu X, Feng X, Murphy PM. The Macrophage-depleting Agent Clodronate Promotes Durable
Hematopoietic Chimerism and Donor-specific Skin Allograft Tolerance in Mice. Sci Rep. 2016;6:22143.

55.   Ma Y, Li Y, Jiang L, Wang L, Jiang Z, Wang Y, et al. Macrophage depletion reduced brain injury
following middle cerebral artery occlusion in mice. J Neuroinflammation. 2016;13:38.

A

Age-matched C56BL6/J; no gavage = **CONTROL**

IL-10$^{-/-}$ + H.hepaticus gavage = **IBD**

Begin Experiments

Birth *Wean*

Develop Disease

Day

0   21 24 26   111

B

IL10$^{-/-}$ - H.hepaticus = CONTROL

IL10$^{-/-}$ + H.hepaticus = **IBD**

1. CONT-PBS

2. CONT-CLOD

3. IBD-PBS

4. IBD-CLOD

Develop Disease   *Mø Depletion*

Begin Experiments

Birth   Wean   **Gavage**   **Liposomes**

Day

0   21   24 26   111 118 125 132

A

B

Control ● IBD ●

PC1 (58.4%)
PC2 (15.5%)

C

inflammatory response
leukocyte migration
leukocyte cell-cell adhesion
positive regulation of cell death
positive regulation of immune response
blood vessel morphogenesis
adaptive immune response
granulocyte activation
negative regulation of immune system process
peptide secretion
blood vessel morphogenesis
adaptive immune response
macrophage activation
granulocyte activation
negative regulation of immune system process
peptide secretion
cellular extravasation
regulation of TNF superfamily cytokine production
positive regulation of macrophage chemotaxis
regulation of T cell apoptotic process

-log10(P)