RESEARCH ARTICLE

Logic-Based Modeling of Inflammatory Macrophage Crosstalk with Glomerular Endothelial Cells in Diabetic Kidney Disease

Krutika Patidar[1] and Ashlee N. Ford Versypt[1,2,3,4,*]

[1]Department of Chemical and Biological Engineering, University at Buffalo, The State University of New York, Buffalo, NY, 14260, USA
[2]Department of Biomedical Engineering, University at Buffalo, The State University of New York, Buffalo, NY, 14260, USA
[3]Institute for Artificial Intelligence and Data Science, University at Buffalo, The State University of New York, Buffalo, NY, 14260, USA
[4]Department of Pharmaceutical Sciences, University at Buffalo, The State University of New York, Buffalo, NY, 14215, USA
[*]Correspondence: ashleefv@buffalo.edu

ABSTRACT   Diabetic kidney disease is a complication in one out of three patients with diabetes. Aberrant glucose metabolism in diabetes leads to structural and functional damage in glomerular tissue and a systemic inflammatory immune response. Complex cellular signaling is at the core of metabolic and functional derangement. Unfortunately, the mechanism underlying the role of inflammation in glomerular endothelial cell dysfunction during diabetic kidney disease is not fully understood. Mathematical models in systems biology allow the integration of experimental evidence and cellular signaling networks to understand mechanisms involved in disease progression. This study developed a logic-based ordinary differential equations model to study inflammatory crosstalk between macrophages and glomerular endothelial cells during diabetic kidney disease progression using a protein signaling network stimulated with glucose and lipopolysaccharide. This modeling approach reduced the biological parameters needed to study signaling networks. The model was fitted to and validated against available biochemical data from *in vitro* experiments. The model identified mechanisms for dysregulated signaling in macrophages and glomerular endothelial cells during diabetic kidney disease. In addition, the influence of signaling interactions on glomerular endothelial cell morphology through selective knockdown and downregulation was investigated. Simulation results showed that partial knockdown of VEGF receptor 1, PLC-γ, adherens junction proteins, and calcium partially recovered the intercellular gap width between glomerular endothelial cells. These findings contribute to understanding signaling and molecular perturbations that affect the glomerular endothelial cells in the early stage of diabetic kidney disease.

NEW & NOTEWORTHY   The work provides a novel analysis of signaling crosstalk between macrophages and glomerular endothelial cells in the early stage of diabetic kidney disease. A logic-based mathematical modeling approach identified vital signaling molecules and interactions that regulate glucose-mediated inflammation in the glomerular endothelial cells and cause endothelial dysfunction in the diabetic kidney. Simulated interactions among vascular endothelial growth factor receptor 1, nitric oxide, calcium, and junction proteins significantly affect the intercellular gap between glomerular endothelial cells.

KEYWORDS   logic-based differential equations; protein signaling network; diabetic nephropathy; inflammation; glomerular endothelial cell

INTRODUCTION

Diabetic kidney disease (DKD) is a major microvascular complication in the kidney that affects about 30% of patients with type 1 diabetes and 40% of patients with type 2 diabetes (1). DKD causes reduced kidney function and may progress into end-stage renal disease (ESRD). Reports suggest that approximately 12–55% of cases of ESRD are due to diabetes (2). This imposes a high burden on the cost of living and quality of life. The high mortality in diabetic patients is often associated with DKD complications and progression to ESRD (3). Although diabetes management is one of the preventative treatments for DKD (4), the rising number of cases of type 1 and type 2 diabetes continues to increase the DKD prevalence (1). There is an urgent need

to understand the pathophysiology of DKD better to design effective and preventative treatment strategies that can mitigate
early signs of microvascular damage and ultimately reduce DKD prevalence in diabetic patients (3).

Ongoing research suggests that DKD is caused by many complex and interlinked processes associated with high blood
glucose and poor glycemic control. The hyperglycemic microenvironment in type 2 diabetes has been linked to metabolic
dysfunction in immune cells and inflammation (5–7). The microenvironment influences the activation of pro-inflammatory
stimuli, such as exogenous lipopolysaccharide (LPS), interferon-gamma (IFN-γ), and tumor necrosis factor-alpha (TNF-α)
(7, 8), which in turn increase glucose uptake in immune cells (9). In the early stage of DKD, immune cells, including
macrophages, tend to polarize to a pro-inflammatory phenotype over an anti-inflammatory phenotype (9, 10). The synergistic
action of pro-inflammatory mediators and increased glucose uptake contribute to dysregulated immune response and signaling
(9). This dysregulated signaling enhances the expression of pro-inflammatory molecules, such as interleukin (IL)-6, TNF-α,
IL-1β, and reactive oxygen species (ROS), which affect downstream macrophage response and activity (9, 10). Inflammatory
molecules can recruit leukocytes to sites of inflammation and initiate the process of leukocyte transmigration (11, 12). These
circulating pro-inflammatory molecules infiltrate the glomerular interstitium (Fig. 1a) in the diabetic kidney (13) and cause
microvascular leakage by upregulation and redistribution of cell-cell adhesion molecules (8, 11, 14, 15). Recruitment of
inflammatory macrophages and their mediators is a critical event in the pathophysiology of the early stage of DKD (13). There
is still an incomplete understanding of the highly complex and diverse molecular interactions in the diabetic kidney (8). Hence,
a deeper analysis of glucose-mediated inflammatory mechanisms and pathways in the diabetic kidney can contribute to a more
comprehensive understanding of DKD development and progression.

Glomerular endothelial dysfunction is a pathological phenomenon that characterizes the early stages of DKD development
and progression. Glomerular endothelial cells (GECs) are early targets of diabetes-dependent inflammation and injury (2).
Glomerular endothelial activation is accompanied by changes in the expression of leukocytes, cytokine production, and loss of
barrier integrity (16). Numerous studies suggest that activation of GECs in early DKD can dysregulate signaling pathways,
promote intracellular hyperglycemia, elevate advanced glycation end products (AGEs) and their receptors, affect cell-cell
adhesion, increase glomerular permeability, and promote the recruitment of inflammatory molecules (17). LPS-stimulated
pro-inflammatory macrophages are involved in the activation of GECs *in vitro* (18). The inflammatory mediators and immune
cells disrupt vascular endothelial growth factor (VEGF) expression, which plays a central role in maintaining the vascular
function of GECs and nitric oxide (NO) homeostasis (19, 20). Imbalanced VEGF expression causes uncoupling of endothelial
nitric oxide synthase (eNOS), imbalance of NO (19), and loss of endothelial barrier integrity (11, 12). The uncoupling of eNOS
causes eNOS to produce NO as a product and ROS as a by-product, which leads to excessive oxidation of NO and reduces NO
bioavailability in GECs (21). Hence, the uncoupling of eNOS is a crucial indicator of GEC activation and dysfunction. NO
bioavailability within GECs has been associated with glomerular endothelial permeability (22, 23). An association between
NO bioavailability and endothelial cell junction integrity has been suggested (11, 24). In the kidneys, hyperglycemia and
inflammatory stimuli induce intercellular junction disruption and enhanced microvascular leakage in the glomeruli. The loss
of intercellular junction integrity leads to increased paracellular permeability, leukocyte transmigration, and dysregulated
intracellular signals (11, 15). Many experimental studies have associated VEGF-mediated tyrosine phosphorylation of junction
proteins with increased permeability (25, 26), decreased transendothelial electrical resistance (27, 28), and regulating the
passage of solutes/macromolecules in kidney (27–29). Stimulation with the inflammatory cytokine IL-6 affected junction protein
VE-cadherin, and phosphorylation increased solute permeability and reduced transendothelial electrical resistance in GECs
(12). Thus, the intercellular gap width changes are considered an early marker of GEC activation and dysfunction during DKD
progression (Fig. 1a). About 50% of the filtration through the glomerulus is dependent on GECs and the endothelial cell surface
layer, making GECs an essential contributor to the glomerular permeability of the kidney (30). Transcellular and paracellular
routes may restrict the transport of water, solutes, or macromolecules through GECs. The fenestrations are transcellular holes
located throughout GECs and are among the main contributors to the glomerular filtration barrier (31, 32). Despite some
experimental studies on GECs in healthy and diseased states, a direct mechanistic link between macrophage-dependent GEC
activation and dysfunction in the early stage of DKD is not fully understood (24). This study investigated the early pathological
events under hyperglycemic and inflammatory conditions leading to GEC activation and signaling dysregulation.

In the present study, we built a protein signaling network (PSN) of interconnected pathways between two cell types—GECs
and macrophages—to study glucose-mediated inflammatory mechanisms in DKD. A PSN depicts the relationship between
interacting proteins using directional arrows (33). The PSN used here was assembled manually using published literature for
selected species and pathways in the PSN and curated experimental data; we have also explored using artificial intelligence-
enabled text mining for automated PSN assembly in other work (34). Here, we used a mathematical approach to integrate
published experimental evidence and the PSN. Ordinary differential equations (ODEs) are often used to model PSNs and other
biochemical networks (35–38), but ODE-based models rely on extensive data, parameter values, and information on reaction
kinetics. Alternatively, modelers have used logic-based modeling, fuzzy logic modeling, and extreme pathways analysis to
reduce the burden of kinetics information needed to study complex mammalian signaling networks (33, 39). These techniques

Figure 1: (a) Visualization of immune-mediated signaling dysregulation and dysfunction in the glomerulus. Infiltrating macrophage cells interact with glomerular endothelial cells, which experience a widening of the intercellular gaps between endothelial cells, leading to the loss of solute and macromolecules from the blood. Created with BioRender.com. (b) The multi-cellular protein signaling network of crosstalk between macrophages (left) and glomerular endothelial cells (right) is stimulated with glucose (GLU) and lipopolysaccharide (LPS), a pro-inflammatory stimulus. Green ovals are input nodes, blue ovals are output nodes, and white ovals are regulatory nodes. Black arrows are activating interactions, a red line with a flat-head arrow is an inhibiting interaction, and red circles indicate logic *AND* gates. An *OR* logic rule connects two or more edges to a subsequent node throughout the network unless indicated otherwise by an *AND* logic gate. The subscript "ec" denotes an intracellular species expressed in endothelial cells. IL-6, TNF-α, IL-1β, and VEGF-A are protein levels expressed in extracellular space. ROS, ROS$_{ec}$, VEGF-A$_{mRNA}$, and NO are expressed within the cells. The Gap Width node denotes a fractional change in the intercellular gap between GECs. The pJunction node represents the phosphorylated junction protein levels. Abbreviations are defined in Table 1.

require few or no parameters. For instance, a discrete logic model (Boolean model) successfully showed the changes in endothelial cell behavior and signaling due to extracellular microenvironments (40). A logic-based model (LBM) was used to gain a qualitative understanding of species expression, signaling pathways, and missing interactions involved in T cell activation (41). Furthermore, pathway analysis of differentially expressed genes and profiles in early DKD has pointed to critical pathophysiological factors in DKD (42). However, each approach has limitations, including a lack of prediction capability for continuous dynamics (time course) in Boolean models, unrealistic timescales in fuzzy logic analysis, and extensive data collection requirements for pathway analysis.

Here, the emphasis was on predicting continuous signaling dynamics while incorporating the prior mechanistic knowledge of intracellular reaction details and the limited available kinetic data. Thus, we considered mathematical methods that combine ODEs and LBMs to leverage the advantages of each approach. Previously, a hybrid modeling method used an ODE model to explain pathway interactions and a qualitative logic-based model to define cell-cell interactions within a tissue environment (43). We adopted a modeling technique that combines continuous dynamic ODE models with qualitative LBMs (39), termed a logic-based ordinary differential equations (LBODEs) model. Several research studies incorporated such transformation of logic-based models into systems of first-order ODEs (44, 45). One study transformed a discrete Boolean model into ODEs using non-normalized Hill functions in the Odefy toolbox to study ligand concentrations and binding affinities for the activation of T cells (44). Logic-based formalism was also used to reverse engineer complex biochemical networks from data and prior knowledge using metaheuristic optimization strategies, as seen in CellNOpt software (46, 47). Netflux is another user-friendly software for developing dynamic logic-based mathematical models of biological networks (39). Kraeutler et al. presented the advantages of using LBODEs, mainly when biochemical parameters are limited or unavailable; LBODEs were shown to identify unexpected network relationships, to generate experimentally testable predictions, and to agree with findings from a validated biochemical model (39). The Netflux software is readily available and easy to use for building a PSN and generating

Table 1: Chemical species abbreviations and definitions used in the study.

Abbreviation	Definition
AGE	advanced glycation end product
AKT	serine/threonine-specific protein kinases
Ca	calcium
eNOS	endothelial nitric oxide synthase
Gap Width	intercellular gap width between GECs
GLU	glucose
IL	interleukin
LPS	lipopolysaccharide
NADPH	nicotinamide adenine dinucleotide phosphate
NFκB	nuclear factor kappa B
NO	nitric oxide
ONOO	peroxynitrite
PI3K	phosphoinositide 3-kinases
pJunction	phosphorylated junction protein
PLC-γ	phospholipase C gamma
RAGE	receptor of advanced glycation end product
ROS	reactive oxygen species
TLR	toll-like receptor
TNF-α	tumor necrosis factor-alpha
VEGF	vascular endothelial growth factor
VEGFR	vascular endothelial growth factor receptor

the corresponding set of LBODEs that comprise the mathematical model for a biochemical system. Such tested software
standardizes the methods used to obtain biological insights, which may not be evident through discrete models (44). We used
Netflux to apply the LBODEs modeling approach to our PSN between macrophages and GECs stimulated under different
conditions to analyze the effects of the signaling crosstalk between these cell types. We predicted species dynamics, network
relationships, and critical drivers of macrophage-dependent inflammation and glomerular endothelial dysfunction in the early
stage of DKD.

METHODS

Protein signaling network assembly

We formulated a PSN for crosstalk between GECs and macrophages (Fig. 1b) by manually assembling information from
literature studies (8, 9, 19, 21, 40, 48–54) and pathway databases (55, 56). The PSN captures the signaling pathways and
molecules (Table 1) that cause inflammation and dysfunction in two cell types: macrophages and GECs. The multi-cellular PSN
has 30 species nodes and 40 interaction edges for reactions (Fig. 1b). The green ovals are input nodes or stimuli, the white ovals
are regulatory nodes, and the blue ovals are model response output nodes. The network has two stimuli: glucose (GLU) and
LPS. LPS is used as a representative systemic inflammatory stimulus like that triggered by diabetes. The *in vitro* data for model
fitting used LPS as an exogenous inflammatory stimulus as it is known to induce a systemic immune response in humans (57).
The network is stimulated with three different treatment conditions: GLU only, LPS only, and both GLU and LPS. Based on the
logic-based formalism, two or more incoming arrows to a node are connected by an *OR* logic rule except where a red circle is
placed denoting an *AND* logic rule (Fig. 1b). An *OR* logic rule between two input nodes is used when either input can activate
(black arrow) or inhibit (red flat-head arrow) the output node. The *OR* logic combines any number of input nodes acting on the
same output; the corresponding reaction rules are listed independently but are joined by the *OR* operator in the LBODEs. An
AND logic rule between two input nodes is used when both inputs are needed to activate the output node; a single reaction rule
is defined for an *AND* logic rule. Fig. 2 shows representative examples of the interaction types between nodes in the PSN
from Fig. 1b. Considering the complex nature of network interactions, we included signaling pathways involved in two main
events in early DKD. These events include the release of pro-inflammatory molecules due to the infiltration of inflammatory
macrophages and macrophage-dependent GEC activation and dysfunction. The rationale behind selecting these pathways and
molecules is discussed further.

Figure 2: Logic-based formalism for a protein signaling network (PSN). The species interactions (a)–(d) are used as examples to explain logic-based reaction rules for the interaction edges in the PSN (Fig. 1). Each of the example descriptions is followed by its corresponding reaction rule. (a) LPS activates TLR4: LPS $\Rightarrow$ TLR4. (b) NO inhibits Ca: !NO $\Rightarrow$ Ca, and Ca activates NO: Ca $\Rightarrow$ NO. (c) VEGFR1 or VEGFR2 activate PI3K$_{ec}$ using an *OR* logic rule: VEGFR1 $\Rightarrow$ PI3K$_{ec}$ *OR* VEGFR2 $\Rightarrow$ PI3K$_{ec}$. (d) ROS$_{ec}$ and NO activate ONOO using an *AND* logic rule: ROS$_{ec}$ & NO $\Rightarrow$ ONOO. A black arrow is an activating interaction, a red line with a flat-head arrow is an inhibiting interaction, and a red circle is a logic *AND* gate. The subscript "ec" denotes an intracellular species expressed in endothelial cells. Abbreviations are defined in Table 1.

In diabetes, immune cells like macrophages fail to metabolize glucose or regulate the efficient transport of glucose, which leads to intracellular hyperglycemia (58, 59). Studies showed that the hyperglycemic microenvironment initiates a pro-inflammatory signaling response downstream, mediated by the activation of LPS and toll-like receptors (TLR) such as TLR4 (9, 10). In cell culture experiments, bacterial LPS is a representative pathogen-associated molecular pattern (PAMP) that allows mammalian cells to recognize bacterial invasion and trigger innate immune responses (9). In cultured macrophages, LPS stimulates a robust inflammatory response, which initiates glucose-mediated inflammation in macrophages (9). Hence, LPS is considered one of the stimuli in our PSN.

LPS transduction signal is carried by activation of PI3K and AKT kinases, which mediate macrophage response, energy consumption, and energy metabolism (60). AKT is a primary substrate of PI3K and is associated with regulating inflammatory and energetic metabolic responses. Therefore, we speculate that PI3K and AKT are crucial in macrophage polarization in the PSN (9).

Advanced glycation is one of the major pathways involved in the development and progression of different diabetic complications, including DKD (54). This protein glycation leads to advanced glycation end products (AGE), which activate and modify intracellular pathways through receptor-mediated signaling (54). The expression of AGE receptors (RAGE) is enhanced in cells like macrophages and endothelial cells during diabetes and inflammation. AGE and RAGE interaction causes oxidative stress and activation of nuclear factor kappa B (NFκB) (54). The glycated proteins release ROS via activation of NADPH oxidase (49, 61). Moreover, aberrant PI3K-AKT signaling increases ROS levels through either direct modulation of mitochondrial bioenergetics or indirect production of ROS as a metabolic by-product (62). Hence, NADPH- and PI3K-dependent ROS are considered in the PSN to gain a better perspective on mediators of ROS.

Further, transcription factor NFκB contributes to the structural and functional changes observed in DKD, including pro-inflammatory macrophage polarization (54). NFκB is responsible for the expression of pro-inflammatory cytokines, like IL-6, TNF-α, and IL-1β, and other inflammatory markers, growth factors including VEGF-A, and adhesion molecules (8).

The macrophage-dependent VEGF-A expression is associated with the direct activation of GECs in the PSN. VEGF expression modulates the crosstalk between endothelial cells and macrophages (63). VEGF and its isoforms are critical to GECs. Excess VEGF contributes to DKD development and progression through disrupted signaling in GECs, whereas low VEGF levels can result in loss of capillaries and endothelial cells in the glomerulus (64). VEGF expression promotes endothelial cell development through NO modulation (64). Studies suggested that glucose indirectly regulates VEGF-A and the signaling of its receptors (VEGFR1 and VEGFR2) in GECs (63, 65, 66). VEGFR1 is a positive regulator of inflammation and immune cell recruitment via PLC-γ and PI3K pathways (51, 67, 68). The downstream signaling of VEGFR1 has yet to be fully understood, mainly due to its mild biological activity in culture (51). VEGF-A mediates signal transduction for cell proliferation, survival, migration, and enhanced vascular permeability via VEGFR2 (20). VEGFR2-dependent signaling promotes eNOS activity, which produces NO, an important vasodilator in GECs (20). VEGF-A and VEGFR2 activity reduced in the GECs during diabetes (69). This phenomenon leads to perturbations in downstream signaling, eNOS uncoupling, and reduced NO bioavailability (70). As a result, excess ROS in GECs can react with available NO to form reactive nitrogen species like peroxynitrite (ONOO) (64, 71). In the PSN, the VEGF-dependent signaling pathway accounts for VEGF-A receptors VEGFR1 and VEGFR2, PLC-γ, eNOS, NO, ROS, and reactive nitrogen species expression. The VEGF-dependent NO bioavailability is connected with GEC dysfunction.

Cell-cell junction proteins hold adjacent GECs together and modulate the passage of macromolecules, proteins, or infiltrating

macrophages through the GEC layer. For activated GECs, the phosphorylation of these junction proteins, such as VE-cadherin,
is regulated by calcium (Ca) expression via the PLC-γ pathway and NO expression (23, 72). The phosphorylation of junction
proteins like VE-cadherin causes a loss of barrier integrity in GECs and triggers the opening of the intercellular junctions.
Intercellular junction protein is shown as a pJunction node in the PSN. We posit that the junction opening or closing is directly
proportional to the size of the intercellular gap between GECs, shown as the Gap Width node in the network.

In addition to the literature-based prior knowledge discussed above that is incorporated into the PSN, the following list
summarizes additional assumptions and restrictions in scope for the PSN:

• Literature suggests a higher polarization of the pro-inflammatory phenotype of macrophage cells during early stages
of DKD (9). We considered a pro-inflammatory phenotype of macrophages expressing pro-inflammatory species and
ignored anti-inflammatory species for simplicity.

• GECs comprise a negatively charged matrix, the glycocalyx, which also contributes to endothelial permeability (2). We
only focused on the endothelial cell layer for GEC signaling and dysfunction. Glycocalyx signaling and characteristics
were not part of this study.

• AGEs are mediators and drivers of diabetic complications (18, 73). We assumed that glucose directly regulates AGE
activation in the PSN. The mechanism regulating glucose-dependent AGE activation is not part of this study.

• The interaction kinetics across different endothelial cell lines are assumed to be the same, but protein levels of species may
differ across cell lines (20). This has been previously considered in other models where data from different endothelial
cell lines were curated due to the lack of studies on a specific cell line.

• The protein signaling network considers only two types of cells: macrophages and GECs. These cells are stimulated by
GLU and LPS, which mimic hyperglycemic and inflammatory responses during the early stages of DKD, respectively.
Using a fixed physiological range allowed us to extract protein expression data from studies performed under similar
treatment conditions. We assumed that full activation of GLU stimulus (i.e., $W_{\text{GLU}} = 1$) is analogous to a 25–30 mM
hyperglycemic range. Full activation of LPS ($W_{\text{LPS}} = 1$) is the same as an LPS treatment of 0.1 µg/ml in cultured cells.

206 Logic-based ordinary differential equations model

We simulated the dynamic influences of stimuli through the PSN using an LBODEs model. The LBODEs modeling technique
combines ODEs that are continuous functions of time with qualitative logic-based Boolean up- or down-regulation (i.e.,
activation or inhibition) using normalized Hill functions (saturating sigmoidal terms) for the logic-based modeling portion (39).
The LBODEs model equations were generated by the Netflux software package (74) after we supplied the necessary interaction
rules and species list from the PSN that we assembled (Tables 2 and 3). The Netflux package was used to generate the LBODEs
model equations and default parameters, which were exported to MATLAB (75). MATLAB was used to simulate, fit, and
validate the model with further customization beyond Netflux for parameter estimation and other analyses. The LBODEs were
solved using the `ode23s` function in MATLAB.

Each LBODE defining the rate of change of normalized activity of a species A with index i follows the same general
form (39):

$$\frac{dA}{dt} = \frac{\overbrace{y_{\text{max}_A} s(A)}^{\text{term 1}} \overbrace{-A}^{\text{term 2}}}{\tau_A} \tag{1}$$

where term 1 is the net activation rate (detailed further below), term 2 is the first-order de-activation rate, and τ_A is the
characteristic time constant for the species. The initial condition is y_{0_A}. The net activation rate is scaled by y_{max_A}, the maximum
normalized activation of species A. Note that when discussing the species parameters more generally, we use the notation τ_i,
y_{0_i}, and y_{max_i}.

The quantity $s(A)$ is the source of A from activating or inhibiting interactions that act alone or in combination through *OR*
or *AND* logic operators. The activation or inhibition of output node species A is based on an interaction rule computed using
a normalized-Hill function dependent on the signal from input node X. The activation function $f_{\text{act}_j}(X)$ for how species X
activates reaction j that produces A is

$$f_{\text{act}_j}(X) = W_j \frac{\beta_j X^{n_j}}{K_j^{n_j} + X^{n_j}}, \tag{2}$$

225 and the inhibition function $f_{\text{inhib}_j}(X)$ for how species X inhibits reaction j that produces A is

$$f_{\text{inhib}_j}(X) = W_j - f_{\text{act}_j}(X) \tag{3}$$

226 where W_j is the weight of reaction j, n_j is the Hill coefficient for the reaction, EC_{50_j} is the half effect,

$$\beta_j = \frac{EC_{50_j}^{n_j} - 1}{2EC_{50_j}^{n_j} - 1}, \tag{4}$$

227 and

$$K_j = (\beta_j - 1)^{1/n_j}. \tag{5}$$

228 Multiple input signals to the same output node are automatically combined using an *OR* logical operator unless indicated
229 explicitly by an *AND* logical operator. Mathematically, $OR(a, b)$ is an *OR* logical operator for input a and b:

$$OR(a, b) = a + b - ab \tag{6}$$

230 $AND(a, b)$ is an *AND* logical operator for input a and b:

$$AND(a, b) = ab \tag{7}$$

231 We next illustrate the development of a representative subset of the equations (*Eqs. 8–11*) for the nodes and interactions
232 illustrated in Fig. 2. The complete set of interaction rules is provided in Table 2, and the corresponding equations are listed in
233 *Eqs. A1–A30*.

234 A representative single activation interaction rule is LPS $\Rightarrow$ TLR4 (Fig. 2a). The corresponding LBODE for tracking TLR4
235 is

$$\frac{d\text{TLR4}}{dt} = \frac{y_{\max_{\text{TLR4}}} f_{\text{act}_3}(\text{LPS}) - \text{TLR4}}{\tau_{\text{TLR4}}} \tag{8}$$

236 where the source $s(\text{TLR4})$ is the activation of TLR4 by LPS in reaction rule $j = 3$ (Table 2), which follows *Eq. 2*.

237 A representative single inhibition interaction rule is !NO $\Rightarrow$ Ca (Fig. 2b), part of the negative feedback loop between NO
238 and Ca. The corresponding LBODE for tracking Ca is

$$\frac{d\text{Ca}}{dt} = \frac{y_{\max_{\text{Ca}}} f_{\text{inhib}_{36}}(\text{NO}) - \text{Ca}}{\tau_{\text{Ca}}} \tag{9}$$

239 where the source $s(\text{Ca})$ is the inhibition of Ca by NO in reaction rule $j = 36$ (Table 2), which follows *Eq. 3*.

240 If a species is activated by two or more species, an *AND* or an *OR* logic operator is applied to combine the signals. A
241 representative *OR* logic rule combining two activating signals is VEGFR1 $\Rightarrow$ PI3K$_{\text{ec}}$ *OR* VEGFR2 $\Rightarrow$ PI3K$_{\text{ec}}$ (Fig. 2c). The
242 corresponding LBODE for tracking PI3K$_{\text{ec}}$ is

$$\frac{d\text{PI3K}_{\text{ec}}}{dt} = \frac{y_{\max_{\text{PI3K}_{\text{ec}}}} OR\left(f_{\text{act}_{24}}(\text{VEGFR1}), f_{\text{act}_{23}}(\text{VEGFR2})\right) - \text{PI3K}_{\text{ec}}}{\tau_{\text{PI3K}_{\text{ec}}}} \tag{10}$$

243 where the source $s(\text{PI3K}_{\text{ec}})$ is the *OR* function (*Eq. 6*) combining activation of reaction rules $j = 23$ and $j = 24$ (Table 2),
244 which follow *Eq. 2* individually.

245 A representative *AND* logic rule combining two activating signals is ROS$_{\text{ec}}$ & NO $\Rightarrow$ ONOO (Fig. 2d). The corresponding
246 LBODE for tracking ONOO is

$$\frac{d\text{ONOO}}{dt} = \frac{y_{\max_{\text{ONOO}}} AND\left(f_{\text{act}_{35}}(\text{NO}), f_{\text{act}_{35}}(\text{ROS}_{\text{ec}})\right) - \text{ONOO}}{\tau_{\text{ONOO}}} \tag{11}$$

247 where the source $s(\text{ONOO})$ is the *AND* function (*Eq. 7*) combining activation by both ROS$_{\text{ec}}$ and NO in reaction rule $j = 35$
248 (Table 2), which follow *Eq. 2* individually.

Model parameter defaults and constraints

250 The LBODEs model has parameters that can be divided into two categories: reaction parameters (W_j, EC_{50_j}, and n_j) and species
251 parameters (τ_i, y_{0_i}, and $y_{\max_i}$). The default values for these parameters, as generated by Netflux, are $W_j = 1$, $EC_{50_j} = 0.5$,
252 $n_j = 1.4$, $\tau_i = 1$, $y_{0_i} = 0$, and $y_{\max_i} = 1$.

253 Reaction parameters include reaction weight W_j, half maximal effective concentration or half effect EC_{50_j}, and Hill
254 coefficient n_j. The reaction parameters govern the interactions between species in the network. The parameter definitions,

Table 2: List of optimal reaction parameters for the LBODEs model organized by reaction index j. Reaction weight (W_j), Hill coefficient (n_j), and half-maximal effect (EC_{50_j}) are shown for respective reaction rules or interactions. Reaction rule A $\Rightarrow$ C denotes input A activates output C. Reaction rule !A $\Rightarrow$ C denotes input A inhibits output C. Reaction rule A & B $\Rightarrow$ C denotes the *AND* logic operator between input A and input B to give output C. Default values are $W_j = 1$, $n_j = 1.4$, and $EC_{50_j} = 0.5$. Estimated values of sensitive parameters are displayed with three significant figures. $W_j = 0$ for $\Rightarrow$ GLU and $\Rightarrow$ LPS reaction rules represent inactive states. W_j is set to 1 based on the active treatment condition for $j = 1$ and/or $j = 2$. W_1 and W_2 are referred as W_{GLU} and W_{LPS}, respectively. The source terms for GLU ($s(\text{GLU})$) and LPS ($s(\text{LPS})$) are only dependent on W_j and independent of n_j and EC_{50_j}. Abbreviations are defined in Table 1.

Index j	Reaction Rule	W_j	n_j	EC_{50_j}
1	$\Rightarrow$ GLU	0	-	-
2	$\Rightarrow$ LPS	0	-	-
3	LPS $\Rightarrow$ TLR4	1	2.70	0.419
4	GLU $\Rightarrow$ AGE	1	1.44	0.5
5	AGE $\Rightarrow$ RAGE	1	2.71	0.474
6	RAGE $\Rightarrow$ NADPH	0.944	2.69	0.470
7	TLR4 & ROS $\Rightarrow$ NFκB	1	2.69	0.5
8	TLR4 $\Rightarrow$ PI3K	0.950	2.69	0.422
9	NADPH $\Rightarrow$ ROS	0.943	2.63	0.545
10	PI3K $\Rightarrow$ AKT	1	2.69	0.5
11	PI3K $\Rightarrow$ ROS	0.950	2.70	0.419
12	NFκB$_{\text{ec}}$ $\Rightarrow$ TNF-α	0.999	3.96	0.839
13	AKT $\Rightarrow$ NFκB	1	2.70	0.5
14	NFκB $\Rightarrow$ IL-6	0.950	2.70	0.420
15	NFκB $\Rightarrow$ TNF-α	0.950	2.70	0.419
16	NFκB $\Rightarrow$ VEGF-A$_{\text{mRNA}}$	0.949	2.70	0.422
17	VEGF-A$_{\text{mRNA}}$ $\Rightarrow$ VEGF-A	0.948	2.77	0.594
18	NFκB $\Rightarrow$ IL-1β	0.950	2.69	0.421
19	VEGF-A $\Rightarrow$ VEGFR1	1	2.71	0.5
20	VEGF-A $\Rightarrow$ VEGFR2	1	2.72	0.5
21	AGE $\Rightarrow$ RAGE$_{\text{ec}}$	1	1.55	0.838
22	RAGE$_{\text{ec}}$ $\Rightarrow$ NADPH$_{\text{ec}}$	0.937	1.59	0.838
23	VEGFR2 $\Rightarrow$ PI3K$_{\text{ec}}$	1	2.72	0.5
24	VEGFR1 $\Rightarrow$ PI3K$_{\text{ec}}$	1	2.72	0.5
25	NADPH$_{\text{ec}}$ $\Rightarrow$ ROS$_{\text{ec}}$	0.946	3.91	0.838
26	PI3K$_{\text{ec}}$ $\Rightarrow$ AKT$_{\text{ec}}$	1	2.92	0.688
27	AKT$_{\text{ec}}$ $\Rightarrow$ eNOS	1	3.12	0.576
28	VEGFR1 $\Rightarrow$ PLC-γ	1	1.4	0.5
29	PLC-γ $\Rightarrow$ NFκB$_{\text{ec}}$	1	1.4	0.5
30	ROS$_{\text{ec}}$ $\Rightarrow$ NFκB$_{\text{ec}}$	1	2.70	0.0100
31	NFκB$_{\text{ec}}$ $\Rightarrow$ IL-6	0.986	2.69	0.471
32	NFκB$_{\text{ec}}$ $\Rightarrow$ IL-1β	0.962	2.70	0.391
33	eNOS $\Rightarrow$ NO	1	1.45	0.156
34	eNOS $\Rightarrow$ ROS$_{\text{ec}}$	1	3.66	0.832
35	ROS$_{\text{ec}}$ & NO $\Rightarrow$ ONOO	1	1.4	0.5
36	!NO $\Rightarrow$ Ca	1	2.68	0.5
37	PLC-γ $\Rightarrow$ Ca	1	1.4	0.5
38	Ca $\Rightarrow$ pJunction	1	1.4	0.5
39	pJunction $\Rightarrow$ Gap Width	1	1.4	0.5
40	Ca $\Rightarrow$ NO	1	1.4	0.5

Table 3: List of optimal species parameter τ_i for the LBODEs model organized by species index i. The time constant (τ_i) values in units of hours for the species used in the model are shown for each treatment condition. The default value of τ_i is 1 hour. Estimated values of sensitive parameters are displayed with three significant figures. The default values of other species parameters are the initial value ($y_{0_i} = 0$) and maximal value ($y_{\max_i} = 1$) for all species unless stated otherwise. $y_{0_i} = 1$ for eNOS, NO, VEGF-A$_{\mathrm{mRNA}}$, and VEGF-A species for the GLU only treatment condition. Abbreviations are defined in Table 1.

Index i	Species	GLU only	LPS only	GLU and LPS
1	GLU	1	1	1
2	LPS	5.05	0.800	1.00
3	AGE	1	1	1
4	VEGFR1	1	1	1
5	VEGFR2	1	1	1
6	VEGF-A$_{\mathrm{mRNA}}$	3.38	0.944	1.47
7	RAGE$_{\mathrm{ec}}$	1	1	1
8	RAGE	1	1	1
9	TLR4	5.03	0.800	1.00
10	NADPH	1	1	1
11	NADPH$_{\mathrm{ec}}$	1	1	1
12	ROS$_{\mathrm{ec}}$	9.96	1.06	5.15
13	ROS	5.88	0.800	1.00
14	PI3K	5.03	0.800	1.00
15	AKT	1	1	1
16	PI3K$_{\mathrm{ec}}$	6.59	0.977	1.00
17	AKT$_{\mathrm{ec}}$	5.75	0.984	1.00
18	NFκB$_{\mathrm{ec}}$	1	1	1
19	NFκB	5.06	0.800	1.00
20	NO	4.35	1.00	6.80
21	ONOO	1	1	1
22	eNOS	1.91	0.981	1.00
23	IL-6	4.65	0.800	1.00
24	TNF-α	9.89	0.800	1.00
25	IL-1β	4.90	1.00	4.94
26	PLC-γ	1	1	1
27	VEGF-A	3.14	0.971	1.10
28	pJunction	1	1	1
29	Ca	1	1	1
30	Gap Width	1	1	1

[255] default values, and constraint ranges are used as defined previously (39, 76). For a given interaction, the reaction weight
[256] W_j denotes the fraction of activation or inhibition of output species, and EC_{50_j} is the fraction of input required to induce a
[257] half-maximal effect on the output (39). EC_{50_j} value below 0.5 indicates that less than 50% of input species is required to
[258] achieve maximum output activity. Correspondingly, EC_{50_j} value above 0.5 indicates that more than 50% of input species
[259] is required to achieve maximum output activity. The Hill coefficient n_j is a shape factor in the LBODEs model (77). These
[260] reaction parameters define the normalized Hill activation and inhibition functions (*Eqs. 2–3*). These functions are constrained
[261] as $f(0) = 0$, $f(1) = W_j$, and $f(0.5) = 0.5W_j$ in Netflux (39, 74). Each parameter's lower and upper bounds are based on prior
[262] knowledge, as in Cao et al. (76). The reaction weight constraints are default: $0 \leq W_j \leq 1$. The constraints on EC_{50_j} and n_j are
[263] selected to satisfy $\beta_j > 1$ (*Eqs. 4–5*) (76), giving $0.001 < EC_{50_j} < 0.84$ and $1.4 \leq n_j \leq 4$.

[264] The species parameters include time constant τ_i (constrained $0.01 \leq \tau_i \leq 10$), initial value y_{0_i}, and maximal fractional
[265] activation of species y_{max} to describe the activity of a species in the network. The time constant (τ_i) parameter is an arbitrary
[266] parameter that modulates the signal propagation during early or late events and provides a rough estimate of the local dynamic
[267] behavior of the network (41). y_{max_i} is set to 1 by default for the full activity of a species. y_{max_i} can be below or above a value
[268] of 1 to denote species knock-down or over-expression, respectively (39). y_{0_i} is set to 0 for all species except when species
[269] downregulation is predicted. In those cases, y_{0_i} is set to 1.

[270] $W_j = 0$ for a reaction j denotes an inactive state. The source terms for the LBODEs model inputs GLU ($s(\text{GLU})$) and LPS
[271] ($s(\text{LPS})$) are only dependent on reaction weight W_j and independent of n_j and EC_{50_j}. W_1 and W_2 are referred as W_{GLU} and
[272] W_{LPS}, respectively, hereafter. Values of W_{GLU} or $W_{\text{LPS}} = 1$ mean the corresponding input is active.

[273] Data selection and normalization

[274] The data for model fitting (parameter estimation) and validation were curated from published experimental studies based on the
[275] following inclusion criteria:

[276] • co-culture *in vitro* studies (18) were used to study cell-cell interaction between macrophages and GECs with some
[277]   exceptions (9, 10, 29, 78–84) for individual cell culture studies on either macrophages or endothelial cells,

[278] • bone-marrow derived macrophages (BMDM) from diabetic subjects and the RAW 264.7 murine macrophage cell line
[279]   were most commonly considered (9, 10, 18, 78, 85),

[280] • cells were stimulated with GLU concentration 5–30 mM and LPS concentration 0–0.1 µg/ml,

[281] • protein concentration from bar graph or data tables at several time points (3, 6, 12, 24, and 48 hours) for three treatment
[282]   conditions GLU only, LPS only, and both GLU and LPS were collected where available (e.g., Fig. 3).

[283] Co-culture experimental systems have gained popularity in the field, especially for studying and engineering complex
[284] multi-cellular synthetic systems and mimicking tissue structures to some extent (86). We used protein concentrations from
[285] individual cell culture studies on either macrophages or endothelial cells for IL-6, TNF-α, VEGF-A, ROS, eNOS, NO, and
[286] ROS_{ec} when data from co-culture systems were unavailable (9, 10, 78, 79). We extracted data from figures or tables in these
[287] publications (10, 18, 78–81) for calibration and from these publications (9, 29, 81–85) for validation using WebPlotDigitizer, a
[288] web-based data extraction software (87). VEGF-A activity under LPS only stimulus was validated using pooled data from
[289] (81, 83). VE-cadherin activity under LPS only stimulus was validated using pooled data from (29, 82). We used ImageJ software
[290] (88) to quantify Western blot images from published papers when needed. Fig. 3 concisely represents a sample bar graph of
[291] protein concentration at different glucose or LPS treatment conditions.

[292] We normalized the curated data for two reasons: (1) to estimate model parameters by fitting model output to data and (2)
[293] to pool data from multiple studies. It has been previously shown that it is essential to normalize the data and simulations in
[294] the same way to ensure uniformity between model output and data (89). Our model output ranges between 0 and 1. Thus, we
[295] normalized the data in this range. Data normalization involves normalizing a data point with respect to the highest value in the
[296] series, but this can overemphasize data outliers and underweight minor and recurring differences in the series (33). Instead,
[297] we used a multi-step non-linear data normalization technique (33) using the open-source software package CellNOptR (47).
[298] To perform data normalization in CellNOptR, the data were first stored as comma-separated values in a MIDAS-formatted
[299] file (Fig. 3). Then, the `normaliseCNOlist` function from the CellNOptR package was used to perform data normalization in
[300] R (90). This function performs the normalization using the following steps (illustrated in Fig. S1):

[301] 1. The data values are transformed to a fold change for the same experimental treatment condition at time 0.

[302] 2. The fold change is transformed using a Hill function similar to *Eq. 2* (91).

3. a penalty for noise in the data (if any) is multiplied by the transformed value to compute the normalized data value.

4. If the Hill-transformed fold changes after noise penalty have negative values, the normalized data are rescaled using the
minimum and maximum values in the normalized data list.

We used the no-noise option, which skips step 3. All experimental data values were considered within a large dynamic range
$[0,\infty)$, which is the default option for the `normaliseCNOlist` function. More detailed information on this normalization
technique can be found in (33) and CellNOptR reference manual (91).

Figure 3: Sample bar graph (left) of protein concentration representing activation from baseline at different glucose (GLU) or
lipopolysaccharide (LPS) treatment conditions as found in published studies. The mean values and upper bounds were obtained
using WebPlotDigitizer. These values were stored in a MIDAS-formatted comma-separated values (CSV) file, as shown in
the table (right). TR:GLU and TR:LPS represent GLU and LPS treatment, respectively, where 0 represents inactive and 1
represents active treatment. DA:TNFa and DV:TNFa represent the time (hour) and concentration (pg/ml) of TNF-α protein,
respectively. Normalized values in the table were obtained using the method proposed by Saez-Rodriguez et al. (33). Baseline
values (black bar) were used for each treatment condition at the start of the experiment (time 0) in the table (bold text). The
values at 48 hours correspond to the respective treatment conditions in the bar graph. For instance, the data obtained from the
blue, red, black, and purple bars corresponds to the respective colored rows in the table. TR: treatment. DA: data attribute. DV:
data value. TNF: tumor necrosis factor.

Parameter estimation and uncertainty quantification

The LBODEs model was first simulated with default parameters for treatment conditions: $W_{\text{GLU}} = 1$ and $W_{\text{LPS}} = 0$ for GLU
only, $W_{\text{GLU}} = 0$ and $W_{\text{LPS}} = 1$ for LPS only, and $W_{\text{GLU}} = W_{\text{LPS}} = 1$ for both GLU and LPS. However, the model predictions
based on default parameters were only qualitative and not fitted to the data. The model parameters are not available in the
literature, and not all parameters have biophysical definitions. Therefore, default parameters were used as a starting point.
We performed structural identifiability and observability analyses to minimize uncertainty due to non-identifiable parameters
in our model (92, 93). Assessing the model's structural identifiability and observability helps detect structural issues before
model calibration (94, 95). We also performed a global sensitivity analysis using Sobol's variance-based method (96) to
identify the most influential parameters affecting the model response and reduce the parameter space for model calibration. We
followed several recommendations from Linden et al. (97) regarding analyses before parameter estimation and best practices for
uncertainty quantification. We then conducted parameter estimation on the sensitive parameters. We quantified the uncertainty
associated with the estimated parameter values and how this propagates to model prediction uncertainty. Finally, we used local
sensitivity analysis on the optimized and validated model as a measure of the effects of local perturbations representing *in silico*
knockdowns on model predictions for all species in the PSN. Each of these procedures is detailed in the remainder of this
section.
The structural identifiability and observability analyses for the LBODEs model were performed before model calibration
using the STRIKE-GOLDD software package in MATLAB (98). The structural identifiability analysis evaluates the possibility
of determining the parameter values from output measurements. In contrast, the observability analysis describes the ability
to infer the dynamic variables of the model from the model output (94). The structural identifiability is entirely defined by
the model equations and input-output mapping and disregards the limitation of data quantity or quality (93–95, 99). The full
input-state-parameter observability (FISPO) algorithm in STRIKE-GOLDD was used to perform a joint structural identifiability

analysis of model parameters (W_j, n_j, and EC_{50_j} for reactions j and τ_i for species i) and observability analysis of all dynamic
variables. The observability of the model variables (states, parameters, and inputs) was determined by calculating the rank of a
generalized observability-identifiability matrix using Lie derivatives (100). The FISPO analysis yielded a full rank matrix for
each computed Lie derivative, which suggests that the LBODEs model is observable and that unknown parameters (W_j, n_j,
EC_{50_j}, and τ_i) are structurally identifiable when the initial states are known. Further details on using the STRIKE-GOLDD
software package and the FISPO algorithm are available elsewhere (94, 98, 100).

We used Sobol's variance-based method to perform a global sensitivity analysis of the model parameters (96). The UQLab
toolbox (101) was used to perform the global sensitivity analysis (102) in MATLAB. We assessed the effect of reaction weight
(W_j), Hill coefficient (n_j), half effect (EC_{50_j}), and time constant (τ_i) on the model response. First, we sampled the parameter
from ranges specified in a previous study on uncertainty quantification of an LBODEs model also developed using Netflux (76).
For this analysis, W_j, n_j, EC_{50_j}, and τ_i were constrained in the ranges of [0.8, 1], [1.4, 4], [0.4, 0.6], and [0.5, 10], respectively.
A Monte Carlo estimator in the UQLab sensitivity analysis toolbox was used to sample from the parameters in respective
ranges. The sensitivity indices were computed using Sobol pseudo-random sampling (101, 102). The sample size for Monte
Carlo estimation was set to 1000. The sample size was chosen based on the optimum computation time for a global sensitivity
analysis on 144 parameters. The Sobol sensitivity analysis computes sensitivity indices for scalar output quantities (97). We
chose the predicted responses of 8 of the 10 output nodes in Fig. 1b (IL-6, TNF-α, IL-1β, VEGF-A, ROS, eNOS, NO, and
ROS$_{ec}$) at 48 hours as our scalar quantities of interest. We computed and plotted the first-order (Supplemental Fig. S2) and
total Sobol sensitivity indices (Supplemental Fig. S3) for the quantities of interest organized by each type of parameter. We
selected the most sensitive parameters as those that had a total Sobol sensitivity index above a given threshold or cut-off value.
Identifying a single cut-off absolute value for all the parameters (W_j, n_j, EC_{50_j}, and τ_i) was challenging. Therefore, we assumed
a different threshold value for each parameter set as in Linden et al. (97). The total order Sobol sensitivity index within 10% of
the maximum sensitivity index was fixed as the threshold for each parameter set. A total of 82 parameters (from the 204 total
for the LBODEs model: the three reaction parameters W_j, n_j, and EC_{50_j} for the 38 downstream reactions in Table 2 and the
one species parameter τ_i for each of the three treatment conditions for the 30 species in Table 3) were identified as sensitive and
were marked for parameter estimation. The non-sensitive parameters were set to their default values. More information on using
the UQLab toolbox for global sensitivity analysis can be found in the user manual (102).

We estimated the sensitive parameters by calibrating the LBODEs model to the normalized curated datasets for each
treatment condition. The sensitive parameters were optimized using a multi-start non-linear least squares parameter optimization
method to avoid converging to a local minimum (103). We sampled 100 sets of parameter start values using the Latin hypercube
sampling technique (104). Each parameter was sampled uniformly across the following ranges: [0.9, 1], [1.4, 4], [0.001, 0.84],
and [0.01, 10] for W_j, n_j, EC_{50_j}, and τ_i, respectively. The Latin hypercube sampling technique combined these parameter
samples to cover the parameter space efficiently. We used the Latin hypercube sampling routine in MATLAB as implemented in
functions provided online (105) using published methods (104, 106). Each of the sampled sets of parameter start values was
used as initial guesses in the `fmincon` nonlinear programming solver in MATLAB, which returns an optimized parameter set.
`fmincon` minimized the objective function, which was the sum of squared error (SSE) between model predictions and published
data (Fig. 3). The SSE was computed between observed responses from data at available time points and model-predicted
responses for the following output nodes in Fig. 1b: IL-6, TNF-α, IL-1β, VEGF-A$_{mRNA}$, ROS, eNOS, NO, and ROS$_{ec}$. First,
we performed a multi-start optimization of sensitive W_j, n_j, EC_{50_j}, and τ_i parameters for the GLU only treatment condition,
where $W_{GLU} = 1$ and $W_{LPS} = 0$. For the GLU only condition, initial values (y_{0_i}) of VEGF-A, NO, and eNOS were set to 1 to
depict relative downregulation. It is relevant to note that the parameter optimization was carried out sequentially in the order
of occurrence of species nodes from top to bottom in the network to avoid bias from multiple data points. For the GLU only
condition, model parameters governing the expression of ROS, ROS$_{ec}$, NO, eNOS, and VEGF-A were fitted against data before
fitting the parameters for IL-6, TNF-α, and IL-1β response. The optimized reaction parameters (W_j, n_j, and EC_{50_j}) were fixed
across different treatment conditions (GLU only, LPS only, both GLU and LPS). The sensitive τ_i parameters were re-calibrated
for the LPS only and both GLU and LPS treatment conditions. The re-calibration of τ_i values was necessary for different
treatment conditions as the strength of the input signal affects rapid or delayed species response. The multi-start parameter
estimation procedure generated 100 sets of fitted parameters for each treatment condition and an optimal (best) parameter set
that gave the smallest SSE between model predictions and data. From the 100 sets of fitted parameters for each treatment
condition, a set was labeled as part of the acceptable subset of parameters if the SSE for the set was within 20% of the smallest
SSE. The means of the parameters within the acceptable subset were defined as the nominal parameter set for each condition.
These nominal parameters are reported in Tables 2 and 3. Estimated values of sensitive parameters are displayed with three
significant figures to distinguish them from the non-sensitive parameters at their default values.

We quantified the uncertainty associated with model predictions using Monte Carlo ensemble simulation, also known
as sampling-based uncertainty propagation (97, 107). This approach repeatedly simulates a model using samples from the
posterior distributions of the parameters as inputs to obtain a mapping to results (107), producing a posterior distribution of

model predictions for each treatment case. To obtain a measure of the uncertainty of the parameters, we used the function
`randsample(population, numSamples)` in MATLAB to return `numSamples` values sampled uniformly at random from
the values in the vector `population`. With `numSamples` at 5000, we called `randsample` independently for each sensitive
parameter. For each parameter, we specified `population` as the values of that parameter in the acceptable parameters subset for
each treatment condition that resulted from the multi-start parameter optimization approach described in the previous paragraph.
The resulting 5000 values for each parameter were considered as the posterior distribution for the parameter. The posterior
distributions for the parameters are shown along with the nominal parameter values (means of acceptable parameters subset) in
Supplemental Fig. S4. We stored the values returned from `randsample` for all the parameters by their sample number. For a
given sample number, each parameter was a sample from its posterior distribution, independent of the other parameter values.
Together, these parameter values at the same sample number comprised a set of parameters that were run through the LBODEs
model to obtain model predictions. The Monte Carlo ensemble simulation involved repeatedly running the LBODEs model
for each of the 5000 combinations of parameter sets organized by sample number. The resulting posterior distributions of
model predictions were used to calculate the 95% equal-tail credible interval at each time point for each response. The 95%
equal-tail credible interval is bound by a lower limit representing the 2.5th percentile of the posterior distribution and upper
limit representing the 97.5th percentile of the posterior distribution (108). In other words, a 95% equal-tail credible interval
suggests that there is a 95% probability that a true estimate is within this interval, and all values inside the interval are more
likely to represent the parameter than the values outside the interval (108, 109). The use of 95% equal-tail credible intervals
provides better interpretability of these estimates. Showing model results with credible intervals from posterior distributions
of model predictions has been widely adopted in systems biology (97, 99, 109, 110). The statistical analysis of the posterior
distribution of parameters and model predictions through credible intervals and visual checks allowed us to understand the
extent of uncertainty in the estimated parameters (Supplemental Figs. S4–S6).

The robustness of reaction rules and species in the model to perturbations for plausible *in silico* knockdowns were analyzed
using a local sensitivity analysis with finite difference approximation:

$$S_{i,k} = \frac{\partial Y_i}{\partial P_k} \approx \frac{\Delta Y_i}{\Delta P_k} \tag{12}$$

408 where $S_{i,k}$ is a sensitivity coefficient for a given output species i and parameter k, $Y_i(P)$ is the output for species i at the nominal
409 parameter values P, $\Delta Y_i = Y_i(P + \Delta P_k) - Y_i(P)$ is the change in output species i calculated at the perturbed parameter value,
410 and ΔP_k is a small perturbation in parameter k.

RESULTS

Model calibration

The simulation results are shown in Figs. 4–6 for the LBODEs model calibrated to the normalized biochemical data for
three different treatment conditions, (1) active GLU stimulus, (2) active LPS stimulus, and (3) active GLU and LPS stimuli,
respectively. Based on the published co-culture experimental studies, the model was simulated for short-term exposure (48
hours). For each treatment condition, most fitted simulations for each output response captured the observed time-dependent
data very well. The posterior distributions and means of acceptable reaction parameters (W_j, n_j, EC_{50_j}) are plotted at 12
hours for respective reaction index j and rule as reported in Table 2 in Supplemental Fig. S4. Supplemental Fig. S5 shows the
posterior distributions and means of acceptable time constant (τ_i) parameter values for different treatment conditions at 12
hours. Further, the distributions of prediction posteriors for IL-6, TNF-α, IL-1β, VEGF-A$_{mRNA}$, ROS, eNOS, NO, ROS$_{ec}$,
VEGF-A, and Gap Width at 12 hours for each condition are reported in Supplemental Fig. S6. The distribution of predicted
responses captures the prediction uncertainty at 12 hours for the respective output species. We observed a low prediction
uncertainty in most output responses, as demonstrated by narrow 95% credible intervals. Also, most observed data were within
the 95% credible intervals of the model predictions.

Under fixed GLU treatment, the upregulated pro-inflammatory cytokines (IL-6, TNF-α, IL-1β) and ROS activity reached a
steady state around 24 hours. As the responses showed the widest credible intervals around 12 hours (Fig. 4c–e), the posterior
distributions of the prediction responses for the 10 output nodes in Fig. 1b are shown at 12 hours (Supplemental Fig. S6). The
fitted IL-1β response had relatively high prediction uncertainty, as observed from the wide 95% credible interval (Fig. 4c).
More experimental evidence at different time points could reduce this prediction uncertainty. Glucose-mediated downregulation
of VEGF-A$_{mRNA}$ (Fig. 4d) caused the decrease in eNOS and NO activity (Fig. 4f–g). Although the PSN model considers a
simplified VEGF-dependent eNOS/NO pathway, it efficiently captured the glucose-mediated NO downregulation. Under high
glucose conditions, we identified that the degree of influence of Ca on NO, i.e., the reaction weight for reaction rule Ca $\Rightarrow$ NO
($j = 40$), also contributed to 100% downregulation of NO activity.

Figure 4: PSN responses to glucose (GLU) for 48 hours ($W_{\text{GLU}} = 1$ and $W_{\text{LPS}} = 0$) for calibration to biochemical data available from cell culture experiments in Tessaro et al. (10), Li et al. (18), de Souza et al. (78), Nakagawa et al. (79), Kasai et al. (80), and Yang et al. (81). Dynamic LBODEs model predicted responses are compared to normalized protein levels of (a) IL-6, (b) TNF-α, (c) IL-1β, (d) VEGF-A$_{\text{mRNA}}$ levels, (e) ROS fluorescence, (f) eNOS, (g) NO, and (h) ROS$_{\text{ec}}$. The model predictions at nominal parameter values are shown as black solid curves, data are shown as orange points, and the 95% credible intervals for the calibration are shown as blue-shaded regions. The subfigure titles provide the sources for the data in the corresponding plots. The subscript "ec" denotes intracellular expression in glomerular endothelial cells. Abbreviations are defined in Table 1.

Figure 5: PSN responses to lipopolysaccharide (LPS) for 48 hours ($W_{GLU} = 0$ and $W_{LPS} = 1$) for calibration to biochemical data available from cell culture experiments in Tessaro et al. (10), Li et al. (18), de Souza et al. (78), Nakagawa et al. (79), Kasai et al. (80), and Yang et al. (81). Dynamic LBODEs model predicted responses are compared to normalized protein levels of (a) IL-6, (b) TNF-α, (c) IL-1β, (d) VEGF-A$_{mRNA}$ levels, (e) ROS fluorescence, (f) eNOS, (g) NO, and (h) ROS$_{ec}$. The model predictions at nominal parameter values are shown as black solid curves, data are shown as orange points, and the 95% credible intervals for the calibration are shown as blue-shaded regions. The subfigure titles provide the sources for the data in the corresponding plots. The subscript "ec" denotes intracellular expression in glomerular endothelial cells. Abbreviations are defined in Table 1.

Figure 6: PSN responses to both glucose (GLU) and lipopolysaccharide (LPS) for 48 hours ($W_{\text{GLU}} = 1$ and $W_{\text{LPS}} = 1$) for calibration to biochemical data available from cell culture experiments in Tessaro et al. (10), Li et al. (18), de Souza et al. (78), Nakagawa et al. (79), Kasai et al. (80), and Yang et al. (81). Dynamic LBODEs model predicted responses are compared to normalized protein levels of (a) IL-6, (b) TNF-α, (c) IL-1β, (d) VEGF-A$_{\text{mRNA}}$ levels, (e) ROS fluorescence, (f) eNOS, (g) NO, and (h) ROS$_{\text{ec}}$. The model predictions at nominal parameter values are shown as black solid curves, data are shown as orange points, and the 95% credible intervals for the calibration are shown as blue-shaded regions. The subfigure titles provide the sources for the data in the corresponding plots. The subscript "ec" denotes intracellular expression in glomerular endothelial cells. Abbreviations are defined in Table 1.

Calibrating time constant parameters for each treatment condition separately (Table 3) allowed us to capture the earlier
activation of IL-6, TNF-α, and IL-1β cytokines for LPS treatment (Fig. 5a–c). LPS-mediated output responses achieved
50% of maximum activation as early as 6 hours when compared to GLU only treatment, which demonstrates the difference
between the strength of glucose (Fig. 4) and LPS (Fig. 5) stimuli. Under LPS treatment, a relatively higher ROS_{ec} activity
was observed (Fig. 5h compared to Fig. 4h) and mainly mediated by NO uncoupling from its pathway. The reaction weight of
eNOS $\Rightarrow ROS_{ec}$ ($j = 34$) was relatively higher than for $NADPH_{ec} \Rightarrow ROS_{ec}$ ($j = 25$) (Table 2). LPS-mediated VEGF-A_{mRNA}
upregulation (Fig. 5d) initiated the increase in eNOS and NO activity (Fig. 5f–g). ROS expression was upregulated due to the
PI3K-dependent pathway in macrophages (Fig. 5e) and the eNOS-dependent pathway in GECs (Fig. 5h). Also, note that the
fitted IL-1β response had smaller prediction uncertainty (narrow 95% credible interval) compared to the GLU only case due to
an additional data point for the LPS only case (Fig. 5c compared to Fig. 4c).
When both GLU and LPS stimuli were active, LPS stimulus dominated the expression of growth factors, cytokines, and ROS,
as observed from their early activation, except for IL-1β (Fig. 6c), NO (Fig. 6g), and ROS_{ec} (Fig. 6h). The fitted VEGF-A_{mRNA}
activity (Fig. 6d) increased after an initial delay compared to the LPS only case, possibly due to the combined effects countering
that of GLU only (decrease from baseline). NO upregulation (Fig. 6g) was also delayed due to the combined negative effect
of GLU and the positive effect of LPS on NO. The fitted IL-1β and NO response had the highest uncertainty in estimates as
observed from the wide 95% credible intervals (Fig. 6c,g), which could be resolved with more comprehensive experimental
data. The broad posterior distribution of uncertainty in the time constant value for NO (Supplemental Fig. S5) is the source for
prediction uncertainty or wide credible intervals. The posterior distributions of predictions for IL-1β and NO with both GLU
and LPS showed a wide range of predicted values at 12 hours, unlike other species (Supplemental Fig. S6).

Model validation

The predicted activity for output responses compared well against most of the validation data not used in the calibration
(Figs. 7–9). For GLU only, the predicted ROS_{ec} activity did not capture the validation dataset well. Although VEGF-A activity
was not fitted to experimental data (instead, VEGF-A (mRA) activity was fitted), the predicted VEGF-A activity agrees with
the validation dataset for GLU treatment (Fig. 7d). The predicted VEGF-A activity for the other treatment conditions did not
match as well with observed data (Fig. 8d,Fig. 9d) with a consistent offset. The estimated W_j for VEGF-$A_{mRNA} \Rightarrow$ VEGF-A
protein conversion ($j = 17$) was less than 1 (Table 2), which explains why the VEGF-A activity reaches a steady state below 1.
Due to the lack of quantitative data for intercellular gap width, we validated the pJunction node against data for the expression
of phosphorylated VE-cadherin, a type of junction protein, instead (Fig. 8e). The change in intercellular Gap Width in the
PSN depends on the phosphorylation of cell-cell junction proteins such as VE-cadherin, which results in the weakening of
endothelial junction integrity (24, 111). The predicted phosphorylated VE-cadherin activity (pJunction node) agreed quite well
with the validation data in the LPS only case (Fig. 8e).

Figure 7: PSN responses to glucose (GLU) for 48 hours ($W_{\text{GLU}} = 1$ and $W_{\text{LPS}} = 0$) for validation against biochemical data available from cell culture experiments in Ayala et al. (9), He et al. (84), Yang et al. (81), Seo et al. (29), and Ramanathan et al. (85). Dynamic LBODEs model predicted responses are compared for selected species against available normalized protein levels of (a) IL-6, (b) TNF-α, (c) IL-1β protein levels, (d) VEGF-A levels, (e) ROS$_{\text{ec}}$, and (f) NO. The model predictions at nominal parameter values are shown as black solid curves, data are shown as orange points, and the 95% credible intervals for the validation are shown as purple-shaded regions. The subfigure titles provide the sources for the data in the corresponding plots. The subscript "ec" denotes intracellular expression in glomerular endothelial cells. Abbreviations are defined in Table 1.

Figure 8: PSN responses to lipopolysaccharide (LPS) for 48 hours ($W_{GLU} = 0$ and $W_{LPS} = 1$) for validation against biochemical data available from cell culture experiments in Ayala et al. (9), He et al. (84), Yang et al. (81), Itaya et al. (83), Seo et al. (29), Du et al. (82), and Ramanathan et al. (85) upon stimulation with lipopolysaccharide (LPS) for 48 hours ($W_{GLU} = 0$ and $W_{LPS} = 1$). Dynamic LBODEs model predicted responses are compared for selected species against available normalized protein levels of (a) IL-6, (b) TNF-α, (c) IL-1β protein levels, (d) VEGF-A levels, (e) phosphorylated VE-cadherin (representing the pJunction node in the network), and (f) NO. The model predictions at nominal parameter values are shown as black solid curves, data are shown as orange points, and the 95% credible intervals for the validation are shown as purple-shaded regions. The subfigure titles provide the sources for the data in the corresponding plots. Abbreviations are defined in Table 1.

Figure 9: PSN responses to both glucose (GLU) and lipopolysaccharide (LPS) for 48 hours ($W_{GLU} = 1$ and $W_{LPS} = 1$) for validation against biochemical data available from cell culture experiments in Ayala et al. (9) and Yang et al. (81) upon stimulation with both glucose (GLU) and lipopolysaccharide (LPS) for 48 hours ($W_{GLU} = 1$ and $W_{LPS} = 1$). Dynamic LBODEs model predicted responses are compared for selected species against available normalized protein levels of (a) IL-6, (b) TNF-α, and (c) VEGF-A. The model predictions at nominal parameter values are shown as black solid curves, data are shown as orange points, and the 95% credible intervals for the validation are shown as purple-shaded regions. The subfigure titles provide the sources for the data in the corresponding plots. Abbreviations are defined in Table 1.

PSN regulatory species dynamics

We plotted the dynamics of the regulatory species in the PSN that were not used in model fitting (white regulatory nodes in Fig. 1b) for the three different treatment conditions (Figs. 10–12). The species dynamics show the behaviors of signaling molecules, transcription factors, and receptors for which dynamic experimental data that met our inclusion criteria were unavailable. The model predicted a short-term upregulation of VEGFR1, VEGFR2, PLC-γ, PI3K$_{ec}$, and AKT$_{ec}$ that returned to baseline (Fig. 10d) following VEGF-A downregulation under GLU treatment. The model predicted that an increase in Gap Width was directly related to Ca and pJunction activity. In the absence of GLU treatment, the predicted activity of AGE, RAGE, RAGE$_{ec}$, NADPH, and NADPH$_{ec}$ remained at baseline (Fig. 11a,e).

Figure 10: PSN responses to glucose (GLU) for 48 hours ($W_{\text{GLU}} = 1$ and $W_{\text{LPS}} = 0$) showing dynamic LBODEs model predictions for all regulatory species at nominal parameter values. (a) Regulatory species activity associated with GLU only stimulation. (b) Regulatory species activity associated with LPS only stimulation. (c) Regulatory species activity observed in glomerular endothelial cells. (d) Regulatory species activity associated with VEGF-A activation. (e) Regulatory species activity associated with Gap Width change. The subscript "ec" denotes intracellular expression in the glomerular endothelial cells. Abbreviations are defined in Table 1.

The model does not consider the competitive binding of VEGF-A to VEGFR1- and VEGFR2 receptors (Fig. 12d). This was based on the calculated total order sensitivities (Supplemental Fig. S3) for the following reaction rules that were relatively insensitive: VEGF-A $\Rightarrow$ VEGFR1 ($j = 19$), VEGF-A $\Rightarrow$ VEGFR2 ($j = 20$), VEGFR1 $\Rightarrow$ PI3K$_{ec}$ ($j = 24$), and VEGFR2 $\Rightarrow$ PI3K$_{ec}$ ($j = 23$). The strength of these interactions (W_j) was equal to 1 (Table 2). However, we consider different VEGFR1- and VEGFR2-mediated downstream signaling in the PSN. VEGFR1 promotes downstream pro-inflammatory signaling via the PLC-γ pathway, whereas VEGFR2 promotes higher kinase activity via PI3K/AKT activation. We also observed that glucose promoted PI3K$_{ec}$ activity for longer (24 hours) when compared to PLC-γ activity (12 hours) (Fig. 10d). Under combined treatment, PI3K$_{ec}$ kinase activity also preceded PLC-γ activity (Fig. 12d).

Figure 11: PSN responses to lipopolysaccharide (LPS) for 48 hours ($W_{GLU} = 0$ and $W_{LPS} = 1$) showing dynamic LBODEs model predictions for all regulatory species at nominal parameter values. (a) Regulatory species activity associated with GLU only stimulation. (b) Regulatory species activity associated with LPS only stimulation. (c) Regulatory species activity observed in glomerular endothelial cells. (d) Regulatory species activity associated with VEGF-A activation. (e) Regulatory species activity associated with Gap Width change. The subscript "ec" denotes intracellular expression within glomerular endothelial cells in the network. Abbreviations are defined in Table 1.

Model predictions under partial activation or knockdown

The calibrated and validated LBODEs model was used to make predictions under other combinations of stimuli beyond those tested experimentally in the literature. The model simulations shown previously were performed with combinations of fully active stimuli. Here, we predicted output responses under partially active stimuli. To modulate the strength of GLU and LPS stimuli, we varied the reaction weights (W_j) for GLU and LPS. The other reaction parameters are set at nominal values unless otherwise indicated (Table 2). We used *Eq. 12* to analyze the robustness of reaction rules and species in the model to perturbations for plausible *in silico* knockdowns, specifically in response to one-at-a-time changes in W_j for all reactions and y_{max_i} for all species.

First, we simulated the effects of a partially active GLU stimulus ($0 \leq W_{GLU} \leq 1$) to explore ranges of hyperglycemia in the early stage of DKD while LPS was not active ($W_{LPS} = 0$) mimicking a non-inflammatory state (Fig. 13). We considered the nominal species parameters for this analysis to be those for GLU only treatment (Table 3). We observed that 50% of the maximum glucose activity ($W_{GLU} \geq 0.25$) was sufficient to activate and express IL-6 cytokine (Fig. 13a). Low glucose activity was insufficient for maintaining upregulated IL-6 levels, and IL-6 activity returned to the baseline unless GLU was fully activated. The half effect (EC_{50}) for $NADPH_{ec} \Rightarrow ROS_{ec}$ ($j = 25$) is 0.838; this interaction rule upstream of IL-6 limits IL-6 to reach its maximal activity when $W_{GLU} = 0.75$. The intracellular levels of ROS were lower in GECs when stimulated with low glucose levels (below 0.5) (Fig. 13b). We observed no recovery in the intercellular gap width when exposed to all non-zero glucose levels ($0.25 \leq W_{GLU} \leq 1$) (Fig. 13d). We believe other intervention strategies besides glucose control may be needed to potentially reverse the changes in gap width and loss in barrier integrity.

Next, the macrophages and GECs were also stimulated with varying levels of LPS stimulus ($0 \leq W_{LPS} \leq 1$) when glucose $W_{GLU} = 1$ was set as high (Fig. 14). This condition mimics the early phase of DKD development, where high glucose levels cause metabolic dysregulation in pro-inflammatory species and the environment. The nominal species parameters were based on GLU only treatment (Table 3). Increasing levels of LPS activity had minimal effect on pro-inflammatory cytokines (Fig. 14a–b) but a substantial effect on the PLC-γ upregulation (Fig. 14c). Note that $y_{0_i} = 1$ for NO for the GLU only treatment at the nominal parameter values (Fig. 4), and $y_{0_i} = 0$ for NO for the cases of LPS only and both GLU and LPS treatments. When considering the partial activation of LPS, we selected $y_{0_i} = 1$ for NO to compare to the GLU only treatment dynamics (Fig. 4g)

Figure 12: PSN responses to both glucose (GLU) and lipopolysaccharide (LPS) for 48 hours (W_{GLU} = 1 and W_{LPS} = 1) showing dynamic LBODEs model predictions for all regulatory species at nominal parameter values. (a) Regulatory species activity associated with GLU only stimulation. (b) Regulatory species activity associated with LPS only stimulation. (c) Regulatory species activity observed in glomerular endothelial cells. (d) Regulatory species activity associated with VEGF-A activation. (e) Regulatory species activity associated with Gap Width change. The subscript "ec" denotes intracellular expression within glomerular endothelial cells in the network. Abbreviations are defined in Table 1.

Figure 13: LBODEs model predictions under partially active glucose (GLU) stimulus to explore ranges of hyperglycemia in the early stage of diabetic kidney disease. W_{GLU} varied between 0 and 1. W_{LPS} was set to 0 to mimic a non-inflammatory state. All other parameters were at their nominal values for treatment with only GLU. Abbreviations are defined in Table 1.

506 as $W_{GLU} = 1$ in all the cases shown. In contrast, different levels of partial LPS activation were considered. We observed
507 that glucose-dependent downregulation of NO activity recovered to the baseline level as LPS activity increased above 50%
508 ($W_{LPS} \geq 0.5$) (Fig. 14d).

Figure 14: LBODEs model predictions under partially active lipopolysaccharide (LPS) and fully active glucose (GLU) stimulus
($W_{GLU} = 1$) to mimic gradual rise in inflammation in the early stage of diabetic kidney disease. W_{LPS} was varied between 0 and
1. All other parameters were at their nominal values for treatment with only GLU. Abbreviations are defined in Table 1.

509 We identified influential interactions (reaction rules) in the PSN that were sensitive to *in silico* knockdowns by reducing the
reaction weight (W_j) of all interactions and performing a local sensitivity analysis on the LBODEs model in response to these
perturbations. We also repeated the process to determine the influential species by reducing the y_{max_i} parameter of all species in
the PSN. We perturbed W_j and y_{max_i} by reducing their optimal (nominal) values one-at-a-time by 15%. The optimal values
for W_j are available in Table 2, and nominally y_{max_i} for each species i. $W_j < 1$ denotes a reduced strength of the interaction
(j). $y_{max_i} < 1$ denotes the knockdown of a species (i) in the PSN. The sensitivity indices were calculated using *Eq. 12* for
each interaction and species upon local perturbation in W_j and y_{max_i}, respectively. We used heatmaps to display the resulting
sensitivity indices at 48 hours (Supplemental Fig. S7). We considered the nominal parameters for this analysis to be those for
the treatment with both GLU and LPS (Tables 2 and 3). A magnitude of sensitivity index greater than 0.1 was used as a criterion
to identify sensitive interactions and species from Supplemental Fig. S7 corresponding to a specific row labeled on the y-axis.
In Fig. 15, we observed the predicted effects on Gap Width from targeting these sensitive interactions and species using an *in*
silico knockdown test. The strengths (W_j) of the targeted interactions and the maximal activities (y_{max_i}) of the targeted species
were reduced by 15% (Fig. 15a–b) and 50% (Fig. 15c–d) from the nominal values. For all examined interactions and species, a
15% reduction in parameter values positively correlated with the decrease in GEC Gap Width compared to the no inhibition
case (no perturbations) (Fig. 15a–b). We found that 50% inhibition of the following reactions reduced the GEC Gap Width
substantially (Fig. 15c): VEGF-A$_{mRNA}$ $\Rightarrow$ VEGF-A ($j = 17$), VEGF-A $\Rightarrow$ VEGFR1 ($j = 19$), VEGFR1 $\Rightarrow$ PLC-γ ($j = 28$),
PLC-γ $\Rightarrow$ Ca ($j = 37$), Ca $\Rightarrow$ pJunction ($j = 38$), and pJunction $\Rightarrow$ Gap Width ($j = 39$). Targeting VEGFR1, NFκB, NO,
PLC-γ, VEGF-A, pJunction, and Ca species in the network substantially recovered the GEC Gap Width (Fig. 15d). Among
the sensitive reactions and species identified in Supplemental Fig. S7, 50% inhibition of NFκB $\Rightarrow$ VEGF-A$_{mRNA}$ reaction
($j = 16$) and VEGF-A$_{mRNA}$ did not reduce the Gap Width or even follow the trajectory for the case of no inhibition. Instead, the
predicted values of Gap Width plateau at 1 in these cases. We postulate that 50% or more inhibition of NFκB $\Rightarrow$ VEGF-A$_{mRNA}$
reaction or of VEGF-A$_{mRNA}$ results in an imbalance in NO and Ca activity due to the negative feedback loop between NO and
Ca. This imbalance can explain the increase in Gap Width slightly compared to the value without inhibition (Fig. 15c–d).

Figure 15: LBODEs model predictions of the normalized Gap Width between GECs under *in silico* knockdowns. Reaction weight W_j of each locally sensitive interaction was reduced by (a) 15% and (c) 50% from its nominal value. Maximum activity y_{max_i} of each locally sensitive species was reduced by (b) 15% and (d) 50% from its nominal value. All other parameters were at their nominal values for treatment with both GLU and LPS. The solid black curve represents the normalized Gap Width without any changes in reaction or species parameters from their nominal values. Panels a and c share a legend, and panels b and d share a legend. Abbreviations are defined in Table 1.

DISCUSSION

In this work, we predicted the continuous signaling dynamics of proteins in macrophages and GECs using an LBODEs modeling technique and prior knowledge of intracellular and intercellular pathways compiled into the PSN. We fitted and validated the model using published *in vitro* data. We demonstrated the effect of hyperglycemia and inflammation on signaling pathways and protein expression.

Glucose- and LPS-mediated upregulation of pro-inflammatory cytokines further augments the phenotypic switch towards the pro-inflammatory state in macrophages in the early stage of DKD (Fig. 7a–c). The model demonstrated this pro-inflammatory phenotype in macrophages with an increased expression of pro-inflammatory species, such as IL-6, TNF-α, IL-1β, and ROS. The consistent secretion of these inflammatory molecules leads to a chronic shift in macrophage phenotype (112). The combined effect of glucose and LPS affected the intensity and early activation of pro-inflammatory cytokines, oxidative stress, and growth factor activity (Figures 6 and 9). The model predicted a slight decrease in TNF-α expression after 6 hours (Fig. 8b). Multiple reactions and reaction parameters were responsible for TNF-α activity. NFκB transcription factor in macrophages and NFκB$_{ec}$ transcription factor in GECs were responsible for direct activation of TNF-α in the network. The differences in the optimal reaction weight (W_j) and half-maximal effect (EC_{50_j}) parameters (Table 2) for NFκB⇒TNF-α and NFκB$_{ec}$⇒TNF-α can explain the shift in TNF-α activity. The model predictions demonstrated the effect of high glucose on early signs of glomerular endothelial activation through reduced VEGF-mediated kinase activity (Fig. 10d), reduced NO bioavailability (Fig. 7f) due to uncoupling of NO from its pathway, increased inflammation (Fig. 7), and loss of intercellular junction integrity (Fig. 10e). The model predicted that VEGF-A primarily regulated endothelial cell survival, kinase activity, NO homeostasis, and permeability, as seen by early PI3K$_{ec}$ activation (113) (Fig. 12d), followed by leukocyte migration and inflammation mediated by VEGFR1/PLC-γ activation (52) (Fig. 12d). Besides VEGF-A, GECs can also be activated by angiopoietins, chemokine ligands, endothelin-1, and pro-inflammatory cytokines (IL-6, TNF-α, IL-1β) upon cell-cell communication with glomerular cells like podocytes (114). The proposed PSN can be extended to include other sources of GEC activation and interaction with glomerular cells. We demonstrated the effects of VEGF-A and its receptors on cell-cell junction opening. From a molecular standpoint, the endothelial barrier is sealed by cell-cell junction proteins, among which VE-cadherin serves as a cornerstone (115). The endothelial barrier is maintained by VE-cadherin and associated catenins in a non-phosphorylated state (26, 72, 116). The modulated phosphorylation of these junction proteins allows for the selective opening or closing of the intercellular barrier (11). The relationship between VEGF-dependent NO and calcium and phosphorylated junction proteins was captured as a

negative feedback loop in the network. A negative feedback loop between NO and Ca allowed for better information processing and cell behavior regulation in the PSN. We observed that lower than healthy NO levels in GECs affected barrier integrity reversibly (40) (Fig. 10e). High glucose and inflammation caused a prolonged and consistent disruption of the cell-cell barrier integrity in GECs (Fig. 12e).

Regulating glucose levels and inflammation to baseline values effectively reduced cytokine expression and oxidative stress and restored NO bioavailability (Figures 13 and 14). We showed that targeting the mechanistic pathway that links VEGF-A, Ca, and phosphorylated junctions is relevant to understanding the early signs of glomerular endothelial activation and dysfunction. We identified that NFκB transcription factor in macrophages, VEGFR1, PLC-γ, Ca, and junction proteins on GECs are potential targets to control intercellular Gap Width (Fig. 15d). Further *in silico* experiments can suggest the minimum percentage of protein knockdown or reaction inhibition required to maintain intercellular GEC Gap Width. We also observed that excessive knockdown of a protein or a reaction could result in an undesirable effect on Gap Width due to the complex nature of intracellular signaling. Although we modeled the change in Gap Width using a general form of the normalized-Hill type ODE, we obtained a reasonable and qualitative prediction of change in Gap Width (Fig. 15). Our understanding of modeling the Gap Width change is limited by literature, *in vitro* data, and other mechanistic models. Further improvements to the model can be made as new information becomes available. Our predictive findings suggest that a multi-targeted approach is necessary to modulate the early markers of DKD development and progression. Multi-targeted strategies may involve glucose control (Fig. 13), attenuating inflammation (Fig. 14), reducing macrophage infiltration and transmigration via PLC-γ, and maintaining GEC barrier integrity. The potential targets identified by the model can be further analyzed and validated through experimental studies and quantitative systems pharmacology models.

Future extensions of this work could include translating the LBODEs model developed here to study protein dysregulation, GEC activation, and dysfunction in diabetic mice using *in vivo* data. This would enable the study of the effects of systemic changes in glucose concentration and long-term exposure to glucose on structural and functional changes in GECs *in vivo*. The structure of GECs consists of transcellular holes, known as fenestrations, which support the glomerular filtration barrier (117, 118). Alterations in the size and density of GEC fenestrations have been recently associated with the disruption in glomerular filtration and the progression of DKD (117). Recent literature provides more understanding of damaged fenestrated endothelium like GECs in disease conditions (117–120) and suggests that—besides VEGF, NO, and calcium—other proteins such as rho-associated protein kinases and myosin light chain kinases are involved in regulating the fenestration structure in endothelial cells (22, 121–123). We aim to extend the proposed PSN by incorporating these additional hypothesized mechanisms responsible for regulating GEC fenestration structure and permeability.

The quantitative agreement between normalized model prediction and observed data shows the usefulness of the LBODEs framework in understanding complex disease mechanisms. Furthermore, the LBODEs framework provides testable hypotheses on the most sensitive mechanisms that lead to GEC activation and dysfunction in DKD. As the modeling framework is composed of ordinary differential equations, it is suitable to be used with many commercially available tools for analysis of dynamical systems and easy to integrate with other mechanistic models in systems biology (39, 124, 125). Throughout this study, we have depicted how these valuable and practical methods for identifiability, parameter sensitivity, and parameter uncertainty analysis can be integrated with the LBODEs dynamical system.

We also acknowledge some limitations of the LBODEs framework compared to other biochemical models. The species behavior is quantified as a fractional activation or inhibition rather than absolute quantities such as concentration. The LBODEs framework is limited to analyses with normalized protein levels. The presented LBODEs framework represents all protein-protein interactions using activating or inhibiting Hill functions, but other mathematical functions can be used as appropriate. The model predictions are limited to the structure of the proposed PSN, but additional processes, proteins, and pathways can be added. Despite careful consideration when pooling data from multiple experimental studies, there were some discrepancies across cell culture studies, especially when measuring oxidative stress and nitric oxide levels. ROS species are often short-lived and highly reactive, making them challenging to measure. Usually, special assays are needed to detect specific forms of oxidative stress and nitric oxide. Such discrepancies could explain the differences in observed normalized data across studies, and they limit model validation (Fig. 7e). Furthermore, in the absence of GEC-specific experimental studies, we obtained protein concentrations from similar endothelial cell types. We believe the observed uncertainty in prediction can be resolved as more comprehensive dynamic and cell-specific data becomes available.

Despite such limitations, the present model motivates new experiments and testable hypotheses that can better elucidate the early signs of DKD progression. The LBODEs modeling framework enables quantifying protein behavior for large-scale and complex signaling networks, especially when experimental validation is limited. The mechanisms identified from the LBODEs model can be used to develop a less complex mechanistic model for which kinetic and cell-specific parameters are available. Our proposed *in silico* exploration of signal transduction and protein expression could support pre-clinical research to test new mechanisms for therapeutic intervention or improve on existing strategies.

CONCLUSIONS

In this study, we presented a multi-cellular signaling LBODEs model to understand the role of diabetes and immune response in the progression of microvascular complications in the diabetic kidney. We created a signaling network of crosstalk between macrophages and GECs stimulated by two inputs—high GLU and a pro-inflammatory stimulus LPS. The signaling network was modeled using continuous ODEs and logic-based interaction rules, where interaction between species was formulated using normalized Hill functions. The model was fitted and validated against published data from *in vitro* experiments performed on murine macrophage cell lines and endothelial cell lines. Our modeling approach combined the advantages of quantitative biochemical models and qualitative LBMs. These pathways and interactions between macrophages and GECs in the early stage of DKD have not been mathematically studied before. In the presence of high GLU and a pro-inflammatory LPS stimulus, we observed upregulation of pro-inflammatory cytokines (IL-6, TNF-α, and IL-1β), oxidative stress, VEGF-A, and intercellular GEC gap width. We demonstrated the role of oxidative stress in macrophages in regulating NFκB transcription factor and NO bioavailability in the downstream signaling response. We found that partial knockdown of VEGFR1, PLC-γ, pJunction, and Ca have the potential to partially recover the intercellular GEC barrier integrity. The potential protein and interaction targets can be further tested through modeling or experimentation. Moreover, the proposed PSN and model can be translated to study histological changes in GECs *in vivo*. Through our signaling network and LBODEs model, we have studied glucose-mediated inflammatory pathways in the diabetic kidney to support an understanding of DKD development and progression.

APPENDIX

Model equations

The following LBODEs (*Eqs. A1–A30*) govern the PSN interactions. The species abbreviations are defined in Table 1. The parameter values are available in Tables 2 and 3. Other model term definitions are available in *Eqs. 1–7*.

$$\frac{d\text{GLU}}{dt} = \frac{y_{\max_{\text{GLU}}} W_{\text{GLU}} - \text{GLU}}{\tau_{\text{GLU}}} \tag{A1}$$

$$\frac{d\text{LPS}}{dt} = \frac{y_{\max_{\text{LPS}}} W_{\text{LPS}} - \text{LPS}}{\tau_{\text{LPS}}} \tag{A2}$$

$$\frac{d\text{AGE}}{dt} = \frac{y_{\max_{\text{AGE}}} f_{\text{act}_4}(\text{GLU}) - \text{AGE}}{\tau_{\text{AGE}}} \tag{A3}$$

$$\frac{d\text{VEGFR1}}{dt} = \frac{y_{\max_{\text{VEGFR1}}} f_{\text{act}_{19}}(\text{VEGF-A}) - \text{VEGFR1}}{\tau_{\text{VEGFR1}}} \tag{A4}$$

$$\frac{d\text{VEGFR2}}{dt} = \frac{y_{\max_{\text{VEGFR2}}} f_{\text{act}_{20}}(\text{VEGF-A}) - \text{VEGFR2}}{\tau_{\text{VEGFR2}}} \tag{A5}$$

$$\frac{d\text{VEGF-A}_{\text{mRNA}}}{dt} = \frac{y_{\max_{\text{VEGF-A}_{\text{mRNA}}}} f_{\text{act}_{16}}(\text{NF}\kappa\text{B}) - \text{VEGF-A}_{\text{mRNA}}}{\tau_{\text{VEGF-A}_{\text{mRNA}}}} \tag{A6}$$

$$\frac{d\text{RAGE}}{dt} = \frac{y_{\max_{\text{RAGE}}} f_{\text{act}_5}(\text{AGE}) - \text{RAGE}}{\tau_{\text{RAGE}}} \tag{A7}$$

$$\frac{d\text{RAGE}_{\text{ec}}}{dt} = \frac{y_{\max_{\text{RAGE}_{\text{ec}}}} f_{\text{act}_{21}}(\text{AGE}) - \text{RAGE}_{\text{ec}}}{\tau_{\text{RAGE}_{\text{ec}}}} \tag{A8}$$

$$\frac{d\text{TLR4}}{dt} = \frac{y_{\max_{\text{TLR4}}} f_{\text{act}_3}(\text{LPS}) - \text{TLR4}}{\tau_{\text{TLR4}}} \tag{A9}$$

$$\frac{d\text{NADPH}}{dt} = \frac{y_{\max_{\text{NADPH}}} f_{\text{act}_6}(\text{RAGE}) - \text{NADPH}}{\tau_{\text{NADPH}}} \tag{A10}$$

$$\frac{d\text{NADPH}_{\text{ec}}}{dt} = \frac{y_{\max_{\text{NADPH}_{\text{ec}}}} f_{\text{act}_{22}}(\text{RAGE}_{\text{ec}}) - \text{NADPH}_{\text{ec}}}{\tau_{\text{NADPH}_{\text{ec}}}} \tag{A11}$$

$$\frac{d\text{ROS}}{dt} = \frac{y_{\max_{\text{ROS}}} OR\left(f_{\text{act}_{11}}(\text{PI3K}), f_{\text{act}_9}(\text{NADPH})\right) - \text{ROS}}{\tau_{\text{ROS}}} \tag{A12}$$

$$\frac{d\text{ROS}_{\text{ec}}}{dt} = \frac{y_{\text{max}_{\text{ROS}_{\text{ec}}}} OR(f_{\text{act}_{34}}(\text{eNOS}), f_{\text{act}_{25}}(\text{NADPH}_{\text{ec}})) - \text{ROS}_{\text{ec}}}{\tau_{\text{ROS}_{\text{ec}}}} \tag{A13}$$

$$\frac{d\text{PI3K}}{dt} = \frac{y_{\text{max}_{\text{PI3K}}} f_{\text{act}_8}(\text{TLR4}) - \text{PI3K}}{\tau_{\text{PI3K}}} \tag{A14}$$

$$\frac{d\text{AKT}}{dt} = \frac{y_{\text{max}_{\text{AKT}}} f_{\text{act}_{10}}(\text{PI3K}) - \text{AKT}}{\tau_{\text{AKT}}} \tag{A15}$$

$$\frac{d\text{PI3K}_{\text{ec}}}{dt} = \frac{y_{\text{max}_{\text{PI3K}_{\text{ec}}}} OR(f_{\text{act}_{24}}(\text{VEGFR1}), f_{\text{act}_{23}}(\text{VEGFR2})) - \text{PI3K}_{\text{ec}}}{\tau_{\text{PI3K}_{\text{ec}}}} \tag{A16}$$

$$\frac{d\text{AKT}_{\text{ec}}}{dt} = \frac{y_{\text{max}_{\text{AKT}_{\text{ec}}}} f_{\text{act}_{26}}(\text{PI3K}_{\text{ec}}) - \text{AKT}_{\text{ec}}}{\tau_{\text{AKT}_{\text{ec}}}} \tag{A17}$$

$$\frac{d\text{NF}\kappa\text{B}}{dt} = \frac{y_{\text{max}_{\text{NF}\kappa\text{B}}} OR(AND(f_{\text{act}_7}(\text{TLR4}), f_{\text{act}_7}(\text{ROS})), f_{\text{act}_{13}}(\text{AKT})) - \text{NF}\kappa\text{B}}{\tau_{\text{NF}\kappa\text{B}}} \tag{A18}$$

$$\frac{d\text{NF}\kappa\text{B}_{\text{ec}}}{dt} = \frac{y_{\text{max}_{\text{NF}\kappa\text{B}_{\text{ec}}}} OR(f_{\text{act}_{29}}(\text{PLC-}\gamma), f_{\text{act}_{30}}(\text{ROS}_{\text{ec}})) - \text{NF}\kappa\text{B}_{\text{ec}}}{\tau_{\text{NF}\kappa\text{B}_{\text{ec}}}} \tag{A19}$$

$$\frac{d\text{NO}}{dt} = \frac{y_{\text{max}_{\text{NO}}} OR(f_{\text{act}_{33}}(\text{eNOS}), f_{\text{act}_{40}}(\text{Ca})) - \text{NO}}{\tau_{\text{NO}}} \tag{A20}$$

$$\frac{d\text{ONOO}}{dt} = \frac{y_{\text{max}_{\text{ONOO}}} AND(f_{\text{act}_{35}}(\text{NO}), f_{\text{act}_{35}}(\text{ROS}_{\text{ec}})) - \text{ONOO}}{\tau_{\text{ONOO}}} \tag{A21}$$

$$\frac{d\text{eNOS}}{dt} = \frac{y_{\text{max}_{\text{eNOS}}} f_{\text{act}_{27}}(\text{AKT}_{\text{ec}}) - \text{eNOS}}{\tau_{\text{eNOS}}} \tag{A22}$$

$$\frac{d\text{IL-6}}{dt} = \frac{y_{\text{max}_{\text{IL-6}}} OR(f_{\text{act}_{14}}(\text{NF}\kappa\text{B}), f_{\text{act}_{31}}(\text{NF}\kappa\text{B}_{\text{ec}})) - \text{IL-6}}{\tau_{\text{IL-6}}} \tag{A23}$$

$$\frac{d\text{TNF-}\alpha}{dt} = \frac{y_{\text{max}_{\text{TNF-}\alpha}} OR(f_{\text{act}_{15}}(\text{NF}\kappa\text{B}), f_{\text{act}_{12}}(\text{NF}\kappa\text{B}_{\text{ec}})) - \text{TNF-}\alpha}{\tau_{\text{TNF-}\alpha}} \tag{A24}$$

$$\frac{d\text{IL-}\beta}{dt} = \frac{y_{\text{max}_{\text{IL-1}\beta}} OR(f_{\text{act}_{18}}(\text{NF}\kappa\text{B}), f_{\text{act}_{32}}(\text{NF}\kappa\text{B}_{\text{ec}})) - \text{IL-1}\beta}{\tau_{\text{IL-1}\beta}} \tag{A25}$$

$$\frac{d\text{PLC-}\gamma}{dt} = \frac{y_{\text{max}_{\text{PLC-}\gamma}} f_{\text{act}_{28}}(\text{VEGFR1}) - \text{PLC-}\gamma}{\tau_{\text{PLC-}\gamma}} \tag{A26}$$

$$\frac{d\text{VEGF-A}}{dt} = \frac{y_{\text{max}_{\text{VEGF-A}}} f_{\text{act}_{17}}(\text{VEGF-A}_{\text{mRNA}}) - \text{VEGF-A}}{\tau_{\text{VEGF-A}}} \tag{A27}$$

$$\frac{d\text{Ca}}{dt} = \frac{y_{\text{max}_{\text{Ca}}} OR(f_{\text{inhib}_{36}}(\text{NO}), f_{\text{act}_{37}}(\text{PLC-}\gamma)) - \text{Ca}}{\tau_{\text{Ca}}} \tag{A28}$$

$$\frac{d\text{pJunction}}{dt} = \frac{y_{\text{max}_{\text{pJunction}}} f_{\text{act}_{38}}(\text{Ca}) - \text{pJunction}}{\tau_{\text{pJunction}}} \tag{A29}$$

$$\frac{d\text{Gap Width}}{dt} = \frac{y_{\text{max}_{\text{Gap Width}}} f_{\text{act}_{39}}(\text{pJunction}) - \text{Gap Width}}{\tau_{\text{Gap Width}}} \tag{A30}$$

SUPPLEMENTAL MATERIAL

Supplemental Figs. S1–S7.

DATA AVAILABILITY

No primary data were generated in this study. The supporting MATLAB code, including parameter values, scripts for plotting, data, and documentation, can be accessed through an open-source GitHub repository (126) available at this link: https://github.com/ashleefv/KidneyImmuneLBM

ACKNOWLEDGMENTS

We thank lab members for their thorough feedback on this manuscript. Preprint (127) is available at https://doi.org/10.1101/2023.04.04.535594.

GRANTS

This work was supported by National Institutes of Health grant R35GM133763 and National Science Foundation CAREER grant 2133411. The content is solely the responsibility of the authors and does not necessarily represent the official views of the funding agencies.

DISCLOSURES

The authors have nothing to disclose.

AUTHOR CONTRIBUTIONS

ANFV and KP conceived and designed research, KP performed experiments, KP analyzed data, ANFV and KP interpreted results of experiments, ANFV and KP prepared figures, KP drafted the manuscript, ANFV and KP edited and revised manuscript, and ANFV and KP approved the final version of the manuscript.

REFERENCES

1. **Centers for Disease Control and Prevention**. National diabetes statistics report; 2020. [1 Nov. 2023]. https://www.cdc.gov/diabetes/data/statistics-report/index.html.

2. **Yang J, Liu Z**. Mechanistic pathogenesis of endothelial dysfunction in diabetic nephropathy and retinopathy. *Front Endocrinol* 13:816400. 2022.

3. **Thomas HY, Ford Versypt AN**. Pathophysiology of mesangial expansion in diabetic nephropathy: mesangial structure, glomerular biomechanics, and biochemical signaling and regulation. *J Biol Eng* 161:19. 2022.

4. **Thomas HY, Ford Versypt AN**. Collagen deposition in diabetic kidney disease boosts intercellular signaling: a mathematical model. *IFAC PapersOnLine* 55:68-73. 2022.

5. **Nikolajczyk BS, Jagannathan-Bogdan M, Shin H, Gyurko R**. State of the union between metabolism and the immune system in type 2 diabetes. *Genes Immun* 124:239-50. 2011.

6. **Sell H, Habich C, Eckel J**. Adaptive immunity in obesity and insulin resistance. *Nat Rev Endocrinol* 812:709-16. 2012.

7. **Tsalamandris S, Antonopoulos AS, Oikonomou E, Papamikroulis GA, Vogiatzi G, Papaioannou S, et al.** The role of inflammation in diabetes: current concepts and future perspectives. *Eur Cardiol* 141:50-9. 2019.

8. **Navarro-González JF, Mora-Fernández C, Muros de Fuentes M, García-Pérez J**. Inflammatory molecules and pathways in the pathogenesis of diabetic nephropathy. *Nat Rev Nephrol* 7:327-40. 2011.

9. **Ayala TS, Tessaro FHG, Jannuzzi GP, Bella LM, Ferreira KS, Martins JO**. High glucose environments interfere with bone marrow-derived macrophage inflammatory mediator release, the TLR4 pathway and glucose metabolism. *Sci Rep* 9:11447. 2019.

10. **Tessaro FHG, Ayala TS, Bella LM, Martins JO**. Macrophages from a type 1 diabetes mouse model present dysregulated PI3K/AKT, ERK 1/2 and SAPK/JNK levels. *Immunobiology* 225:151879. 2019.

11. **Sluiter TJ, van Buul JD, Huveneers S, Quax PHA, de Vries MR**. Endothelial barrier function and leukocyte transmigration in atherosclerosis. *Biomedicines* 94:328. 2021.

12. **Yang YC, Fu H, Zhang B, Wu YB**. Interleukin-6 downregulates the expression of vascular endothelial-cadherin and increases permeability in renal glomerular endothelial cells via the trans-signaling pathway. *Inflammation* 456:2544-58. 2022.

13. **Rayego-Mateos S, Morgado-Pascual J, Opazo-Rios L, Guerrero-Hue M, García-Caballero C, Vázquez-Carballo C, et al.** Pathogenic pathways and therapeutic approaches targeting inflammation in diabetic nephropathy. *Int J Mol Sci* 21:3798. 2020.

14. **Alicic RZ, Rooney MT, Tuttle KR**. Diabetic kidney disease: Challenges, progress, and possibilities. *Clin J Am Soc Nephrol* 1212:2032-45. 2017.

15. **Bazzoni G, Dejana E**. Endothelial cell-to-cell junctions: molecular organization and role in vascular homeostasis. *Physiol Rev* 843:869-901. 2004.

16. **Hunt BJ, Jurd KM**. Endothelial cell activation. A central pathophysiological process. *BMJ* 3167141:1328-9. 1998.

17. **Johnson-Léger C, Imhof B**. Forging the endothelium during inflammation: pushing at a half-open door? *Cell Tissue Res* 314:93-105. 2003.

18. **Li Y, Yu W, He M, Yuan F**. The effects of M1/M2 macrophages on the mRNA expression profile of diabetic glomerular endothelial cells. *Nephron* 145:1-11. 2021.

19. **Takahashi T, Harris RC**. Role of endothelial nitric oxide synthase in diabetic nephropathy: lessons from diabetic eNOS knockout mice. *J Diabetes Res* 2014:590541. 2014.

20. **Wu Q, Finley S**. Mathematical model predicts effective strategies to inhibit VEGF-eNOS signaling. *J Clin Med* 9:1255. 2020.

21. **Zhao HJ, Wang S, Cheng H, Zhang MZ, Takahashi T, Fogo AB, et al.** Endothelial nitric oxide synthase deficiency produces accelerated nephropathy in diabetic mice. *J Am Soc Nephrol* 17:2664-9. 2006.

22. **Shu A, Du Q, Chen J, Gao Y, Zhu Y, Lv G, et al.** Catalpol ameliorates endothelial dysfunction and inflammation in diabetic nephropathy via suppression of RAGE/RhoA/ROCK signaling pathway. *Chem Biol Interact* 348:109625. 2021.

23. **Bates DO**. Vascular endothelial growth factors and vascular permeability. *Cardiovasc Res* 87:262-71. 2010.

24. **Di Lorenzo A, Lin MI, Murata T, Landskroner-Eiger S, Schleicher M, Kothiya M, et al.** eNOS-derived nitric oxide regulates endothelial barrier function through VE-cadherin and Rho GTPases. *J Cell Sci* 126:5541-52. 2013.

25. **Yamaguchi I, Tchao BN, Burger ML, Yamada M, Hyodo T, Giampietro C, et al.** Vascular endothelial cadherin modulates renal interstitial fibrosis. *Nephron Exp Nephrol* 1201:e20-31. 2011.

26. **Esser S, Lampugnani MG, Corada M, Dejana E, Risau W**. Vascular endothelial growth factor induces VE-cadherin tyrosine phosphorylation in endothelial cells. *J Cell Sci* 111:1853-65. 1998.

27. **Satchell SC, Anderson KL, Mathieson PW**. Angiopoietin 1 and vascular endothelial growth factor modulate human glomerular endothelial cell barrier properties. *J Am Soc Nephrol* 153. 2004.

28. **Li L, Liu Q, Shang T, Song W, Xu D, Allen TD, et al.** Aberrant activation of Notch1 signaling in glomerular endothelium induces albuminuria. *Circ Res* 1285:602-18. 2021.

29. **Seo JA, Sayyed ND, Lee YJ, Jeon HY, Kim EB, Hong SH, et al.** Midazolam ameliorates hyperglycemia-induced glomerular endothelial dysfunction by inhibiting transglutaminase 2 in diabetes. *Int J Mol Sci* 23:753. 2022.

30. **Satchell S, Tooke J**. What is the mechanism of microalbuminuria in diabetes: A role for the glomerular endothelium? *Diabetologia* 51:714-25. 2008.

31. **Satchell SC, Braet F**. Glomerular endothelial cell fenestrations: an integral component of the glomerular filtration barrier. *Am J Physiol Renal Physiol* 296:F947-56. 2009.

32. **Denzer L, Muranyi W, Schroten H, Schwerk C**. The role of plvap in endothelial cells. *Cell Tissue Res* 3922:393-412. 2023.

33. **Saez-Rodriguez J, Alexopoulos LG, Epperlein J, Samaga R, Lauffenburger DA, Klamt S, et al.** Discrete logic modelling as a means to link protein signalling networks with functional analysis of mammalian signal transduction. *Mol Syst Biol* 5:331. 2009.

34. **Patidar K, Deng JH, Mitchell CS, Ford Versypt AN**. Cross-domain text mining of pathophysiological processes associated with diabetic kidney disease. *Int J Mol Sci* 258:4503. 2024.

35. **Ford Versypt AN, Harrell GK, McPeak AN**. A pharmacokinetic/pharmacodynamic model of ACE inhibition of the renin-angiotensin system for normal and impaired renal function. *Comput Chem Eng* 104:311-22. 2017.

36. **Pilvankar MR, Higgins MA, Ford Versypt AN**. Mathematical model for glucose dependence of the local renin–angiotensin system in podocytes. *Bull Math Biol* 804:880-905. 2018.

37. **Pilvankar MR, Yong HL, Ford Versypt AN**. A glucose-dependent pharmacokinetic/pharmacodynamic model of ACE inhibition in kidney cells. *Processes* 73:131. 2019.

38. **Ford Versypt AN**. Multiscale modeling in disease. *Curr Opin Syst Biol* 27:100340. 2021.

39. **Kraeutler MJ, Soltis AR, Saucerman JJ**. Modeling cardiac β-adrenergic signaling with normalized-Hill differential equations: comparison with a biochemical model. *BMC Syst Biol* 4:157. 2010.

40. **Weinstein N, Mendoza L, Gitler I, Klapp J**. A network model to explore the effect of the micro-environment on endothelial cell behavior during angiogenesis. *Front Physiol* 8:960. 2017.

41. **Saez-Rodriguez J, Simeoni L, Lindquist JA, Hemenway R, Bommhardt U, Arndt B, et al.** A logical model provides insights into T cell receptor signaling. *PLoS Comput Biol* 3:e163. 2007.

42. **Fu J, Akat K, Sun Z, Zhang W, Schlondorff D, Liu H, et al.** Single-cell RNA profiling of glomerular cells shows dynamic changes in experimental diabetic kidney disease. *J Am Soc Nephrol* 30:533-45. 2019.

43. **Selvaggio G, Cristellon S, Marchetti L**. A novel hybrid logic-ODE modeling approach to overcome knowledge gaps. *Front Mol Biosci* 8:760077. 2021.

44. **Wittmann D, Krumsiek J, Saez-Rodriguez J, Lauffenburger D, Klamt S, Theis F**. Transforming Boolean models to continuous models: methodology and application to T-cell receptor signaling. *BMC Syst Biol* 3:98. 2009.

45. **Krumsiek J, Pölsterl S, Wittmann D, Theis F**. Odefy – from discrete to continuous models. *BMC Bioinf* 11:233. 2010.

46. **Henriques D, Rocha M, Saez-Rodriguez J, Banga J**. Reverse engineering of logic-based differential equation models using a mixed-integer dynamic optimisation approach. *Bioinformatics* 31:2999—3007. 2015.

47. **Terfve C, Cokelaer T, Henriques D, MacNamara A, Goncalves E, Morris MK, et al.** Cellnoptr: A flexible toolkit to train protein signaling networks to data using multiple logic formalisms. *BMC Syst Biol* 6:133. 2012.

48. **Afanasév I**. Signaling of reactive oxygen and nitrogen species in diabetes mellitus. *Oxid Med Cell Longevity* 3:361-73. 2010.

49. **Rendra E, Riabov V, Mossel DM, Sevastyanova T, Harmsen MC, Kzhyshkowska J**. Reactive oxygen species (ROS) in macrophage activation and function in diabetes. *Immunobiology* 224:242-53. 2018.

50. **Durán W, Breslin J, Sanchez F**. The NO cascade, eNOS location, and microvascular permeability. *Cardiovasc Res* 87:254-61. 2010.

51. **Shibuya M**. Vascular endothelial growth factor receptor-1 (VEGFR-1/Flt-1): a dual regulator for angiogenesis. *Angiogenesis* 9:225-30. 2006.

52. **Shibuya M**. Structure and dual function of vascular endothelial growth factor receptor-1 (Flt-1). *Int J Biochem Cell Biol* 33:409-20. 2001.

53. **Dalal P, Muller W, Sullivan D**. Endothelial cell calcium signaling during barrier function and inflammation. *Am J Pathol* 190:535-42. 2019.

54. **Singh VP, Bali A, Singh N, Jaggi AS**. Advanced glycation end products and diabetic complications. *Korean J Physiol Pharmacol* 18:1-14. 2014.

55. **Kanehisa M, Goto S**. Kegg: Kyoto encyclopedia of genes and genomes. *Nucleic Acids Res* 28:27-30. 2000.

56. **Pico AR, Kelder T, van Iersel M P, Hanspers K, Conklin BR, C E**. WikiPathways: pathway editing for the people. *PLoS Biol* 6:1-4. 2008.

57. **Brooks D, Barr LC, Wiscombe S, McAuley DF, Simpson AJ, Rostron AJ**. Human lipopolysaccharide models provide mechanistic and therapeutic insights into systemic and pulmonary inflammation. *Eur Respir J* 56:1901298. 2020.

58. **Lee H, Kim MJ, Lee IK, Hong CW, Jeon JH**. Impact of hyperglycemia on immune cell function: a comprehensive review. *Diabetol Int* 154:745-60. 2024.

59. **Witcoski Jr L, de Lima JD, Somensi AG, de Souza Santos LB, Paschoal GL, Uada TS, et al.** Metabolic reprogramming of macrophages in the context of type 2 diabetes. *Eur J Med Res* 291:497. 2024.

60. **Kanehisa M**. Post-Genome Informatics. Kyoto University, Japan: Oxford University Press; 2000.

61. **Paravicini TM, Touyz RM**. NADPH oxidases, reactive oxygen species, and hypertension clinical implications and therapeutic possibilities. *Diabetes Care* 31:S170-80. 2008.

62. **Koundouros N, Poulogiannis G**. Phosphoinositide 3-kinase/Akt signaling and redox metabolism in cancer. *Front Oncol* 8:160. 2018.

63. **Carranza K, Veron D, Cercado A, Bautista N, Pozo W, Tufro A, et al.** Cellular and molecular aspects of diabetic nephropathy; the role of VEGF-A. *Nefrologia* 35:131-8. 2015.

64. **Nakagawa T, Sato W, Kosugi T, Johnson RJ**. Uncoupling of VEGF with endothelial NO as a potential mechanism for abnormal angiogenesis in the diabetic nephropathy. *J Diabetes Res* 2013:184539. 2013.

65. **Warren CM, Ziyad S, Briot A, Der A, Iruela-Arispe ML**. A ligand-independent VEGFR2 signaling pathway limits angiogenic responses in diabetes. *Sci Signaling* 7:ra1. 2014.

66. **Yang KS, Lim JH, Kim TW, Kim MY, Kim Y, Chung S, et al.** Vascular endothelial growth factor-receptor 1 inhibition aggravates diabetic nephropathy through eNOS signaling pathway in db/db mice. *PloS One* 9:e94540. 2014.

67. **Imoukhuede PI, Dokun AO, Annnex BH, Popel AS**. Endothelial cell-by-cell profiling reveals the temporal dynamics of VEGFR1 and VEGFR2 membrane localization after murine hindlimb ischemia. *Am J Physiol Heart Circ Physiol* 304:H1085-96. 2013.

68. **Weddell JC, Imoukhuede PI**. VEGFR1 promotes cell migration and proliferation through PLCγ and PI3K pathways. *NPJ Syst Biol Appl* 4:1. 2018.

69. **Feliers D, Chen X, Akis N, Choudhury GG, Madaio M, Kasinath BS**. VEGF regulation of endothelial nitric oxide synthase in glomerular endothelial cells. *Kidney Int* 68:1648-59. 2005.

70. **Pinter E, Haigh J, Nagy A, Madri JA**. Hyperglycemia-induced vasculopathy in the murine conceptus is mediated via reductions of VEGF-A expression and VEGF receptor activation. *Am J Pathol* 158:1199-206. 2001.

71. **Ferraro M**. Vascular endothelial growth factor: basic science and clinical progress. *Endocr Rev* 25:581-611. 2004.

72. **Stan R, Tse D, Deharvengt S, Smits N, Xu Y, Luciano M, et al.** The diaphragms of fenestrated endothelia: gatekeepers of vascular permeability and blood composition. *Dev Cell* 23:1203-18. 2012.

73. **Wang RS, Saadatpour A, Albert R**. Boolean modeling in systems biology: an overview of methodology and applications. *Phys Biol* 9:055001. 2012.

74. **Saucerman J**. Netflux; 2015. [18 Sept. 2024]. https://github.com/saucermanlab/Netflux.

75. **MathWorks**. MATLAB (R2021b); 2021. [4 Sept. 2024]. https://www.mathworks.com.

76. **Cao S, Aboelkassem Y, Wang A, Valdez-Jasso D, Saucerman JJ, Omens JH, et al.** Quantification of model and data uncertainty in a network analysis of cardiac myocyte mechanosignalling. *Phil Trans R Soc A* 378:20190336. 2020.

77. **Prinz H**. Hill coefficients, dose-response curves and allosteric mechanisms. *J Chem Biol* 3:37-44. 2009.

78. **de Souza LF, Barreto F, da Silva EG, Andrades ME, Guimarães ELM, Behr GA, et al.** Regulation of LPS stimulated ROS production in peritoneal macrophages from alloxan-induced diabetic rats: involvement of high glucose and PPARγ. *Life Sci* 81:153-9. 2007.

79. **Nakagawa T, Sato W, Sautin YY, Glushakova O, Croker B, Atkinson MA, et al.** Uncoupling of vascular endothelial growth factor with nitric oxide as a mechanism for diabetic vasculopathy. *J Am Soc Nephrol* 17:736-45. 2006.

80. **Kasai N, Sugimoto K, Horiba N, Suda T**. Effect of D-glucose on nitric oxide release from glomerular endothelial cells. *Diabetes Metab Res Rev* 17:217-22. 2001.

81. **Yang Z, Mo X, Gong Q, Pan Q, Yang X, Cai W, et al.** Critical effect of VEGF in the process of endothelial cell apoptosis induced by high glucose. *Apoptosis* 13:1331-43. 2008.

82. **Du L, Dong F, Guo L, Yinglong H, Yi F, Liu J, et al.** Interleukin-1β increases permeability and upregulates the expression of vascular endothelial-cadherin in human renal glomerular endothelial cells. *Mol Med Rep* 11:3708-14. 2015.

83. **Itaya H, Imaizumi T, Yoshida H, Koyama M, Suzuki S, Satoh K**. Expression of vascular endothelial growth factor in human monocyte/macrophages stimulated with lipopolysaccharide. *Thromb Haemostasis* 85:171-6. 2001.

84. **He S, Hu Q, Xu X, Niu Y, Chen Y, Lu Y, et al.** Advanced glycation end products enhance m1 macrophage polarization by activating the MAPK pathway. *Biochem Biophys Res Commun* 525:334-40. 2020.

85. **Ramanathan M, Giladi A, Leibovich JS**. Regulation of vascular endothelial growth factor gene expression in murine macrophages by nitric oxide and hypoxia. *Exp Biol Med* 228:697-705. 2003.

86. **Goers L, Freemont P, Polizzi K**. Co-culture systems and technologies: taking synthetic biology to the next level. *J R Soc Interface* 11:96. 2014.

87. **Rohatgi A**. WebPlotdigitizer: Version 4.4; 2020. [4 Sept. 2024]. https://automeris.io/WebPlotDigitizer.

88. **Abramoff MD, Magalhaes PJ, Ram SJ**. Image processing with ImageJ. *Biophotonics Int* 11:36-42. 2004.

89. **Makaryan S, Finley S**. Enhancing network activation in natural killer cells: predictions from in silico modeling. *Integr Biol* 12:109-21. 2020.

90. **RStudio Team**. RStudio: Integrated Development Environment in R; 2020. [4 Sept. 2024]. http://www.rstudio.com.

91. **Cokelaer T, Eduati F, MacNamara A, Schrier S, Terfve C, Gjerga E, et al.**. Training of Boolean logic models of signalling networks using prior knowledge networks and perturbation data; 2022. [10 Sept. 2024]. https://www.bioconductor.org/packages/release/bioc/manuals/CellNOptR/man/CellNOptR.pdf.

92. **Villaverde AF, Raimúndez E, Hasenauer J, Banga JR**. A comparison of methods for quantifying prediction uncertainty in systems biology. *IFAC-PapersOnLine* 5226:45-51. 2019.

93. **Fröhlich F, Theis FJ, Hasenauer J**. Uncertainty analysis for non-identifiable dynamical systems: profile likelihoods, bootstrapping and more. In: **Mendes P, Dada JO, Smallbone K**, editors. Computational Methods in Systems Biology. Springer; 2014. p. 61-72.

94. **Villaverde AF, Tsiantis N, Banga JR**. Full observability and estimation of unknown inputs, states and parameters of nonlinear biological models. *J R Soc Interface* 16:20190043. 2019.

95. **Hass H, Kreutz C, Timmer J, Kaschek D**. Fast integration-based prediction bands for ordinary differential equation models. *Bioinformatics* 32:1204-10. 2015.

96. **Saltelli A, Ratto M, Andres T, Campolongo F, Cariboni J, Gatelli D, et al.** Variance-based methods. In: Global Sensitivity Analysis. The Primer. Hoboken, NJ: John Wiley & Sons; 2008. p. 155-82.

97. **Linden NJ, Kramer B, Rangamani P**. Bayesian parameter estimation for dynamical models in systems biology. *PLoS Comput Biol* 1810:1-48. 2022.

98. **Díaz-Seoane S, Rey Barreiro X, Villaverde AF**. STRIKE-GOLDD 4.0: user-friendly, efficient analysis of structural identifiability and observability. *Bioinformatics* 391:btac748. 2022.

99. **Ciupe S, Tuncer N**. Identifiability of parameters in mathematical models of SARS-CoV-2 infections in humans. *Sci Rep* 12:14637. 2022.

100. **Villaverde AF**. STRIKE-GOLDD v4.0 User Manual; 2022. [4 Sept. 2024]. http://afvillaverde.webs.uvigo.gal.

101. **Marelli S, Sudret B**. UQLab: a framework for uncertainty quantification in MATLAB. *2nd Int Conf Vulnerability Risk Anal Manage (ICVRAM 2014)*:2554—2563. 2014.

102. **Marelli S, Lamas C, Konakli K, Mylonas C, Wiederkehr P, Sudret B**. UQLab user manual – Sensitivity analysis; 2022. [4 Sept. 2024]. https://www.uqlab.com/sensitivity-user-manual.

103. **Mitra ED, Hlavacek WS**. Parameter estimation and uncertainty quantification for systems biology models. *Curr Opin Syst Biol* 18:9-18. 2019.

104. **Marino S, Hogue I, Ray C, Kirschner D**. A methodology for performing global uncertainty and sensitivity analysis in systems biology. *J Theor Biol* 254:178-96. 2008.

105. **Kirschner D**. Uncertainty and sensitivity functions and implementation; 2008. [30 Nov. 2024]. http://malthus.micro.med.umich.edu/lab/usadata/.

106. **Stein M**. Large sample properties of simulations using latin hypercube sampling. *Technometrics* 29:143-51. 1987.

107. **Helton JC, Johnson JD, Sallaberry CJ, Storlie CB**. Survey of sampling-based methods for uncertainty and sensitivity analysis. *Reliab Eng Syst Saf* 91:1175-209. 2006.

108. **Hespanhol L, Vallio C, Saragiotto B, Costa L**. Understanding and interpreting confidence and credible intervals around effect estimates. *Braz J Phys Ther* 23:290-301. 2018.

109. **Huber HA, Georgia SK, Finley SD**. Systematic Bayesian posterior analysis guided by Kullback-Leibler divergence facilitates hypothesis formation. *J Theor Biol* 558:111341. 2023.

110. **Mortlock RD, Georgia SK, Finley SD**. Dynamic regulation of JAK-STAT signaling through the prolactin receptor predicted by computational modeling. *Cell Mol Bioeng* 141:15-30. 2021.

111. **Gavard J**. Endothelial permeability and VE-cadherin: a wacky comradeship. *Cell Adh Migr* 7:455-61. 2014.

112. **Rex J, Albrecht U, Ehlting C, Thomas M, Zanger U, Sawodny O, et al.** Model-based characterization of inflammatory gene expression patterns of activated macrophages. *PLoS Comput Biol* 12:e1005018. 2016.

113. **Zhang QX, Nakhaei-Nejad M, Haddad G, Wang X, Loutzenhiser R, Murray AG**. Glomerular endothelial PI3 kinase-a couples to VEGFR2, but is not required for eNOS activation. *Am J Physiol Renal Physiol* 3016:F1242-50. 2011.

114. **Li M, Armelloni S, Mattinzoli D, Ikehata M, Chatziantoniou C, Alfieri C, et al.** Crosstalk mechanisms between glomerular endothelial cells and podocytes in renal diseases and kidney transplantation. *Kidney Res Clin Pract* 431:47-62. 2024.

115. **Sidibé A, Imhof BA**. VE-Cadherin phosphorylation decides: Vascular permeability or diapedesis. *Nat Immunol* 153:215-7. 2014.

116. **Thomas KA**. Angiogenesis. In: Encyclopedia of Cell Biology. Academic Press; 2016. p. 102-16.

117. **Finch NC, Fawaz SS, Neal CR, Butler MJ, Lee VK, Salmon AJ, et al.** Reduced glomerular filtration in diabetes is attributable to loss of density and increased resistance of glomerular endothelial cell fenestrations. *J Am Soc Nephrol* 336. 2022.

118. **Finch NC, Neal CR, Welsh GI, Foster RR, Satchell SC**. The unique structural and functional characteristics of glomerular endothelial cell fenestrations and their potential as a therapeutic target in kidney disease. *Am J Physiol Renal Physiol* 3254:F465-78. 2023.

119. **Szafranska K, Kruse LD, Holte CF, McCourt P, Zapotoczny B**. The whole story about fenestrations in LSEC. *Front Physiol* 12:735573. 2021.

120. **DeLeve LD**. Liver sinusoidal endothelial cells in hepatic fibrosis. *Hepatology* 615. 2015.

121. **Zapotoczny B, Braet F, Kus E, Ginda-Mäkelä K, Klejevskaja B, Campagna R, et al.** Actin-spectrin scaffold supports open fenestrae in liver sinusoidal endothelial cells. *Traffic* 2012:932-42. 2019.

122. **Zapotoczny B, Szafranska K, Lekka M, Ahluwalia BS, McCourt P**. Tuning of liver sieve: the interplay between actin and myosin regulatory light chain regulates fenestration size and number in murine liver sinusoidal endothelial cells. *Int J Mol Sci* 2317. 2022.

123. **Esser S, Wolburg K, Wolburg H, Breier G, Kurzchalia T, Risau W**. Vascular endothelial growth factor induces endothelial fenestrations in vitro. *J Cell Biol* 1404:947-59. 1998.

124. **Rikard SM, Athey TL, Nelson AR, Christiansen SLM, Lee J, Holmes JW, et al.** Multiscale coupling of an agent-based model of tissue fibrosis and a logic-based model of intracellular signaling. *Front Physiol* 10. 2019.

125. **Chudasama VL, Ovacik MA, Abernethy DR, Mager DE**. Logic-based and cellular pharmacodynamic modeling of bortezomib responses in u266 human myeloma cells. *J Pharmacol Exp Ther* 3543:448-58. 2015.

126. **Patidar K, Ford Versypt AN**. KidneyImmuneLBM; 2024. [9 Feb. 2024]. https://github.com/ashleefv/KidneyImmuneLBM.

127. **Patidar K, Ford Versypt AN**. Logic-based modeling of inflammatory macrophage crosstalk with glomerular endothelial cells in diabetic kidney disease. *bioRxiv* Preprint:2023.04.04.535594v3. 2024.