Telling mutualistic and antagonistic ecological networks apart by learning their multiscale structure

Running headline: Multiscale network structure and interaction type

Benoît Pichon[1,2,*] (ORCID : 0009-0007-6206-5482)

Rémy Le Goff [2] (ORCID : 0009-0002-7827-7096)

Hélène Morlon[2, #] (ORCID : 0000-0002-3195-7521

Benoît Perez-Lamarque[2,3, #] (ORCID : 0000-0001-7112-7197)

[1] ISEM, Univ Montpellier, CNRS, EPHE, IRD, F-34095 Montpellier, France

[2] Institut de biologie de l'École normale supérieure (IBENS), École normale supérieure, CNRS, INSERM, Université PSL, 46 rue d'Ulm, 75 005 Paris, France

[3] Institut de Systématique, Évolution, Biodiversité (ISYEB), Muséum national d'histoire naturelle, CNRS, Sorbonne Université, EPHE, UA, CP39, 57 rue Cuvier 75 005 Paris, France

* Corresponding author: benoit.pichon0@gmail.com

\# B.P.L and H.M. jointly supervised this work.

Acknowledgments

We thank I. Overcast, J. Voznica, L. Aristide, S. Lambert, I. Quintero, C. Fruciano, J. Clavel, A. Silva, for comments on the first version of the manuscript. We also thank S. Chaffron, C. Bowler, C. Nef, and E. Thébault for fruitful discussions.

The authors declare no conflict of interest.

Author contributions

All authors conceived the ideas and designed methodology; B.P performed research. B.P, H.M and B.P.L led the writing of the manuscript. All authors contributed critically to the drafts and gave final approval for publication.

Data availability

The code used for the analyses is available on Zenodo: https://doi.org/10.5281/zenodo.7786771

Abstract

1. Characterizing and understanding the processes that shape the structure of ecological
networks, which represent who interacts with whom in a community, has many
implications in ecology, evolutionary biology, and conservation. A highly debated
question is whether and how the structure of a bipartite ecological network differs
between antagonistic (e.g., herbivory) and mutualistic (e.g., pollination) interaction
types.

2. Here, we tackle this question by using a multiscale characterization of network
structure, machine learning tools, and a large database of empirical and simulated
bipartite networks.

3. Contrary to previous studies focusing on global structural metrics, such as nestedness
and modularity, which concluded that antagonistic and mutualistic networks cannot
be told apart from only their structure, we find that they can be told apart with a
multiscale characterization of their structure. Mesoscale structures, such as motif
frequencies, appear particularly informative, with an over-representation of densely
connected motifs in antagonistic networks, and of motifs with asymmetrical
specialization in mutualistic networks. These characteristics can be used to predict
interaction types with relatively good confidence. We also find structural uniqueness
linked to specific ecologies (e.g., pollination, parasitism) beyond the classical
mutualism/antagonism dichotomy.

4. Our results clarify structural differences between antagonistic and mutualistic
networks and highlight machine learning as a promising approach for characterizing
interaction types in systems where it is not directly observable.

Keywords: ecological interactions, network structure, motif frequency, interaction
classification, machine learning

1. Introduction

Species in ecosystems engage in multiple types of antagonistic or mutualistic interactions, such as predation, parasitism, pollination, seed dispersal, or mycorrhizal symbioses (Bronstein, 2015; Thomas et al., 2005). Such interactions between two groups of potentially taxonomically distant species are often represented by bipartite networks (Bascompte et al., 2003; Bascompte & Jordano, 2007; Fontaine et al., 2011). Studying the structure of these interaction networks is fundamental for unraveling the different processes behind their ecological organization (Bastolla et al., 2009; Fontaine et al., 2011; Suweis et al., 2013; Thompson, 2005), understanding their evolutionary dynamics, and prioritizing conservation efforts (Bastolla et al., 2009; Olesen et al., 2007; Rezende, Lavabre, et al., 2007).

Early studies on bipartite ecological networks, focused mainly on pollination and herbivory networks, suggested that mutualistic and antagonistic networks have different structures (Bascompte et al., 2003; Thébault & Fontaine, 2008, 2010). Antagonistic networks tend to have a modular structure, where species within modules preferentially interact with each other and rarely interact with species from other modules (Krause et al., 2003; Newman, 2006b). Conversely, mutualistic networks tend to have a nested structure, where specialist species mainly interact with generalists, and generalists form a core of interactions (Jordano et al., 2002; Rohr et al., 2014; Bascompte et al., 2003). Such structural differences have been observed in both terrestrial (Fontaine et al., 2011) and marine networks (Guimarães et al., 2007). However, several studies have challenged the generality of the modular/nested dichotomy (Chagnon, 2016; Fontoura et al., 2020; Michalska-Smith & Allesina, 2019; Pilosof et al., 2014). The dichotomy seems to apply only to highly connected networks (*i.e.* with a high proportion of realized interactions), whereas networks with lower connectance are often

simultaneously modular and nested (Fortuna et al., 2010). In addition, networks tend to be

more modular when their interactions are more intimate, *i.e.* when the degree of biological

integration between interacting individuals is high, regardless of interaction type (Fontaine et

al., 2011; Guimarães et al., 2007; Pires & Guimarães, 2013). Recently, using a large and diverse

dataset of empirical networks with different types of ecology (*e.g.* ant-plant, bacteria-phage,

host-parasitoid,…), Michalska-Smith and Allesina (2019) showed that the structure of

ecological networks is extremely heterogeneous and that global measures of network structure

including nestedness and modularity are not sufficient to discriminate antagonistic and

mutualistic networks (Michalska-Smith & Allesina, 2019). Song and Saavedra (2020) suggested

that the confounding effect of environmental conditions indeed prevents telling antagonistic

and mutualistic networks apart from such global metrics unless environmental differences are

accounted for (Song & Saavedra, 2020).

Macro-scale metrics of network structure, like nestedness and modularity, are not the

only way to characterize network structure. The nestedness/modularity paradigm, as the main

method to characterize bipartite network structure, has recently been questioned (Simmons,

Cirtwill, et al., 2019). The authors highlighted the limits of macro-scale metrics for capturing

structural differences between bipartite networks and showed the usefulness of motifs (a small

subset of species interactions exhibiting varying topologies), which capture the mesoscale

structure (Milo et al., 2002). Motifs have been informative in various contexts, like the

assessment of network stability or resistance to biological invasions (Losapio et al., 2021;

Stouffer & Bascompte, 2010; Vitali et al., 2022). Another way to characterize the structure of

interaction networks is through the spectral density of the Laplacian graph. Laplacian graphs

have been used in other fields, for instance, to compare structural properties of species phylogenies (Lewitus & Morlon, 2016) or brain networks (de Lange et al., 2014), but only marginally to study the structure of ecological networks (Michalska-Smith & Allesina, 2019). If such approaches can be informative to identify structural components that differ between antagonistic and mutualistic networks, this will both improve our understanding of the processes involved in the evolution and maintenance of these networks and offer the possibility to predict interaction type in systems where interactions cannot be directly observed. For instance, many widespread fungal endophytes have ecologies that are either unknown or controversial (Chagnon et al., 2016; Selosse et al., 2018). More generally, networks involving microorganisms, such as bacteria in soils (Freilich et al., 2010) and plankton in the oceans (Chaffron et al., 2021; Lima-Mendez et al., 2015), have often been constructed using presence/absence data rather than direct observations, and the type of a large proportion of these interactions remains undetermined (Bjorbækmo et al., 2020; Lima-Mendez et al., 2015).

Here, we use a variety of network structural metrics combined with different machine learning tools to (i) investigate if there are structural components of networks, and which ones, that are distinct in mutualistic *versus* antagonistic networks, (ii) assess whether unsupervised and supervised classification approaches can robustly predict interspecific interaction type from network structure, and (iii) evaluate whether currently available simulation tools for bipartite networks can help improve the accuracy of the classifiers. We compare the predictive power of the different structural metrics and classification methods. Finally, we discuss the implications of our findings for the understanding of natural communities.

2. Materials and methods

Empirical interaction networks

We analyzed 343 local or regional scales bipartite interaction networks with well-

known interaction types. We obtained these networks by downloading the networks from

(Michalska-Smith & Allesina, 2019), among which some come from the Web of Life database

(web-of-life.es (Fortuna et al., 2014)). We kept networks containing at least 10 species in each

guild, as the structure of small networks is difficult to characterize (Olesen et al., 2007), in

particular in terms of motif composition (preliminary results not shown). In addition, we

gathered 9 mycorrhizal networks that also fit this size criterion: 6 were collected from

individual papers (Jacquemyn et al., 2011, 2015; Martos et al., 2012; Montesinos-Navarro et al.,

2012; Öpik et al., 2009; Xing et al., 2019) and 3 from the database generated by (Põlme et al.,

2018). Among the 343 networks, 216 were mutualistic and 127 antagonistic. They comprised 9

different types of ecology: bacteria-phage (antagonistic; n=17), seed dispersal (mutualistic;

n=29), host-parasite (antagonistic; n=77), host-parasitoid (antagonistic; n=10), plant-

mycorrhiza (mutualistic; n=9), ant-plant (mutualistic; n=3), herbivory (antagonistic; n=23),

anemone-fish (mutualistic; n=1), and pollination (mutualistic; n=174). Each network was

converted into a binary matrix, taking the value 1 if species i interacts with species j, and 0

otherwise, *i.e.* we only focused on the presence/absence of interactions, not on their

frequencies. The guild with the largest (resp. lowest) number of species was in rows (resp.

columns).

Comparing the structure of mutualistic *versus* antagonistic networks

We characterized the structure of ecological networks in three ways focusing on different network scales (Fig. 1).

First, we characterized the macro-scale structure of the networks using 5 global metrics: the total size (number of species in both guilds), the absolute difference in the number of species in the two guilds, the connectance, the nestedness, and the modularity. We quantified nestedness using the NODFc index that corrects for network size ("NODFc" function from the R-package maxnodf, version 1.0.0 (Hoeppke & Simmons, 2021)). We quantified modularity using the function *computeModules* from the R-package bipartite (version 2.16) (Dormann et al., 2008) with the "Beckett" method.

Second, we characterized the mesoscale structure of the networks by performing bipartite motif analyses that depict the patterns of interactions between subsets of species (called motifs; Fig. 1) (Simmons, Sweering, et al., 2019). For each network, we computed the frequencies of the 44 possible motifs involving between 2 and 6 nodes using the function *mcount* from the bmotif R-package (version 2.0.2) (Simmons et al., 2018) with the normalization procedure "normalise_sum" (*i.e.* which computes the frequency of motifs among the total number of motifs). The function *mcount* does not allow counting motifs with more than 6 nodes, and limiting the size to 6 seemed reasonable given the size of our smaller networks (10 species): the frequencies of larger motifs is likely to be more important in larger networks, independently of the type of interactions.

Third, we used a multi-scale approach stemming from graph theory that integrates the

whole structural information of a network, from micro- to macro-scales: the spectral density

of the Laplacian graph (Newman, 2006a). We computed the normalized Laplacian graph of

each network (Newman, 2006a), defined as $L_{norm} = D^{-1/2} L D^{-1/2}$, where D is the degree matrix (a

diagonal matrix counting the number of interactions for each species) and L is the Laplacian

graph, defined as the difference between the adjacency matrix (a square binary matrix whose

elements represent the interactions between species of the network) and the degree matrix

(Fig. S2A). The normalized Laplacian graph is a symmetric and positive-definite matrix whose

eigenvalues are constrained between 0 and 2 (Banerjee & Jost, 2008), which makes them

comparable between networks of different sizes, and symmetrical around 1. There are as many

eigenvalues equal to 0 (or 2) as there are connected components (*i.e.* perfect modules of species

that are not connected with any species from other modules), and we can thus expect a

modular network to have more eigenvalues close to 0 (and 2) than a nested network. To be

able to efficiently compare Laplacian graphs across networks of different sizes, we converted

the eigenvalues into a spectral density using a Gaussian convolution (Lewitus & Morlon,

2016)(Fig. S2A). To do so, we adapted the *spectR* function from the RPANDA R-package

(version 1.9) (Morlon et al., 2016) for bipartite networks and used a bandwidth of 0.025 for the

Gaussian convolution. We tested the influence of the bandwidth value (from 0.001 to 0.091) on

the resulting spectral densities. We characterized each network by extracting its spectral

density values at 100 points uniformly distributed between 0 and 1.

The type of ecology (*e.g.,* pollination, host-parasite) can impact network structure

(Chagnon et al., 2016; Guimarães et al., 2007). To assess whether there are structural differences

between mutualistic and antagonistic networks beyond ecology, we controlled for the

confounding effect of ecology by fitting linear mixed models (function *lmer* from the R-package

lme4 (version 1.1-27)) with the type of interaction (*i.e.* mutualistic or antagonistic) as a fixed

effect, the type of ecology as a random effect, and one structural metric as the response variable

(Type II Wald test). In addition, for each structural metric, we fitted linear models with only

the type of ecology or the type of interaction as fixed effects and evaluated the best-fitting

model by comparing their Akaike information criterion (AIC).

Unsupervised classification

202   To visualize structural differences across antagonistic *versus* mutualistic networks, we

203   used principal component analyses (PCA), with networks characterized either in terms of their

204   global metrics, motif frequencies, or Laplacian spectral densities. PCAs were performed with

205   the functions *PCA* and *fviz_pca_ind* from the FactoMineR and factoextra R-packages (versions

206   2.4 and 1.0.7 respectively) (Lê et al., 2008).

207   For each type of characterization, we then used unsupervised K-means classification

on the two main principal components projections. K-means were performed with the function

kmeans with 2 clusters, and we performed chi-squared tests to investigate whether the two

clusters tend to separate mutualistic from antagonistic networks. To quantify the ability to

separate antagonistic and mutualistic networks, we used a measure of classification accuracy

computed as the F-score with the function *F1_score* from the R-package MLmetrics. The F-

score is the harmonic mean of the precision (*e.g.,* the frequency of true mutualistic among

predicted mutualistic networks) and the recall (*e.g.,* the frequency of predicted mutualistic

among true mutualistic networks).

Supervised classification

216
Next, we used supervised classifiers to classify antagonistic *versus* mutualistic

networks based on their structure. As input for the supervised classifier, for each interaction

network, we used a set of structural variables, either its global metrics, its motif frequencies,

its Laplacian spectral density, or any combination of these sets of variables. As an output, the

supervised classifier provides an interaction type: "mutualistic" or "antagonistic". Supervised

classifiers are based on supervised learning: the classifiers have first to be trained with a

random subset of the interaction networks (the "training set") in order to optimize its internal

parameters; once trained, it can classify the other networks (the "test set") from the input

variables. We measured the accuracy of the approach using both the percentages of mutualistic

and antagonistic networks from the test set that are correctly classified by the classifier and the

F-scores of the classification of both types of networks.

We first investigated the accuracy of linear classifiers using logit and Lasso regressions.

In Lasso regression, a constraint on the estimated model coefficients can make the regression

with a high number of variables more reliable. Logit regressions were computed with the *glm*

function, whereas the Lasso regressions were performed with the function *cv.glmnet* from the

glmnet R-package (version 4.1-2) (Friedman et al., 2010). We found that training the classifiers

with 80% of the interaction networks and testing it on the remaining 20% provided a good

trade-off between the size of the test set and the accuracy of the classification.

Second, we used artificial neural networks (ANN) implemented in the *neuralnet*

function from the R-package neuralnet (version 1.44.2) (Günther & Fritsch, 2010). We

optimized their configuration (number of layers and number of neurons in each layer)

according to the input variables. We selected a 3 layers network, constituted of the input and

output layers and an intermediate layer with 5, 20, and 60 neurons for global metrics, motif

frequencies, and spectral density inputs respectively. Preliminary analyses showed that these

choices provided a good compromise between classification accuracy and overfitting (results

not shown). We also selected a non-linear logistic transformation, and the maximal number of

steps was fixed at 10^8.

We independently trained and tested the classifiers on empirical networks using the

different sets of structural input variables. For each configuration, the ANN trainings were

replicated 50 times with different training and test sets, and we reported the average

percentages of correct classifications among antagonistic and mutualistic networks and their

respective F-scores.

Sensitivity analyses on the supervised classification using the motif frequencies

We selected the most accurate classification approach (supervised classification using

ANN on motif frequencies; see Results) and carried out a series of additional sensitivity

analyses: we (i) developed a measure of classification confidence, (ii) assessed the impact of

overfitting and the over-representation of mutualistic *versus* antagonistic networks and of

specific types of ecology (e.g. pollination) in the training set, (iii) tested the specificity of our

approach, *sensu* (Michalska-Smith & Allesina, 2019) (iv) investigated the effect of the types of

ecology on our classification, and finally (iv) analyzed which motif complexity is required to

accurately classify networks.

First, because the training sets were limited in size (we only had a total of 343 empirical

networks), we introduced a measure of the classification confidence for each empirical

network. We trained and tested the ANN classifier on motif frequencies for 10,000

independent training and test sets of empirical networks: each empirical network was

therefore classified on average 2,000 times with different training sets. We then computed the

percentage of classifications that fell in each category, which gives an idea of classification

confidence: we considered that a given network was classified as mutualistic (resp.

antagonistic) with "high confidence" if it was classified as mutualistic (resp. antagonistic) in

more than 75% of the classifications. Conversely, a given network was classified as mutualistic

(resp. antagonistic) with "low confidence" if it was classified as mutualistic (resp. antagonistic)

in more than 50% of the classifications but in less than 75% of them.

Second, we measured overfitting by comparing the percentages of correct

classifications obtained on the training *versus* test sets as a function of the number of neurons

in the hidden layer. In addition, we assessed the impact on our results of the over-

representation of mutualistic *versus* antagonistic networks in the training set (more than 60%

of the networks were mutualistic): we trained the classifier with the same number of

antagonistic and mutualistic networks (from 5 to 125 networks for each type of interaction)

and compared the percentages of correct classifications with the ones obtained with the same-

size training sets composed of networks randomly chosen in the two interaction types. In each

condition, we replicated the classification 50 times with different training sets. Similarly, we

assessed the impact on our results of the over or under-representation of some types of ecology

in the training set (e.g. the pollination networks represent ~50% of the complete database). To

do so, we trained the classifier with the same number of networks per type of ecology (from 5

to 20 networks for each type of ecology; if a type of ecology had fewer networks than the

number per type of ecology, all networks were included in the training set) and compared the

percentages of correct classifications with the ones obtained with same-size training sets

composed of networks chosen randomly across ecology types.

Third, we tested the specificity of our approach, *sensu* (Michalska-Smith & Allesina,

2019): we applied perturbations to the empirical networks in order to investigate whether the

classification accuracies we obtained were not driven only by differences in network size,

connectance, and/or degree distribution between antagonistic and mutualistic networks (Fig.

2A). We used three different randomization strategies, all of which maintained network size

constant, and tested our classification approach on each randomized dataset. First, we

randomized the networks while keeping their connectance fixed using the *shuffle.web* function

of the **bipartite** R-package. Second, we randomized the networks while keeping their degree

distributions fixed using the r2table algorithm implemented in the *null.model* function of the

bipartite R-package. Third, we randomized the networks while keeping both the connectance

and their degree distributions fixed using the *permatswap* function with quasiswap method

from the **vegan** R-package (v2.5-7). For each of these 3 randomization strategies, we

randomized 200 times each network, and for each randomization, we classified them using

ANN on motif frequencies with 50 different training and testing sets.

Fourth, we investigated if the classifiers were strictly learning to separate antagonistic

from mutualistic networks or if they also learned the potential structural singularities of each

type of ecology from our database (*i.e.*, data-linkage). We performed two different analyses to

control for the signal from the type of ecology. First, we removed one by one each type of

ecology from the training test (*e.g.* pollination networks), we trained the classifiers on 50

independent training sets constituted by 80% of the remaining networks, and we tested the

classification on the networks from the removed type of ecology (*e.g.* are pollination networks

classified among mutualistic networks when no pollination network has been used in the

training?). If the classifiers learn to separate antagonistic from mutualistic networks regardless

of ecology, the classification should perform as well as the original classification. If they learn

to classify ecologies rather than the type of interaction, then it should perform very badly.

Second, we investigated whether the structural differences between mutualistic and

antagonistic networks learned by our ANN classifier were explained only by structural

differences specific to each type of ecology (*e.g.,* pollination *versus* host-parasite). To do so, we

shuffled the "mutualistic" or "antagonistic" nature of each type of ecology while keeping 5

(*resp.* 4) different types of ecology as "mutualistic" (*resp.* "antagonistic"). In such shuffled

datasets, "antagonistic" networks could for example correspond to all pollination, host-

parasitoid, mycorrhizal and anemone-fish networks, whereas "mutualistic" networks could

correspond to all bacteria-phage, herbivory, host-parasite, ant-plant, and seed dispersal

networks. If the structural differences from the type of ecology alone drive the ANN

classifications, we expect similar accuracies with these shuffled datasets compared with the

original dataset. We performed 300 of such shuffling, and for each shuffled dataset, we

classified "mutualistic" *versus* "antagonistic" networks using ANN on motif frequencies with

50 different training and testing sets. In this analysis, we noticed cases when most networks

were classified in one type of interaction (the most represented interaction in terms of number

of networks); in this case, the percentage of correct classifications will appear very high for this

interaction type, and very low for the other. Therefore, in addition to the mean accuracy of

antagonistic and mutualistic networks, we measured the difference in accuracy. A large

difference reflects a loss of the ANN ability to classify properly.

Finally, we investigated which level of motif complexity (size) is required to

distinguish mutualistic from antagonistic networks. To do so, we measured the classification

accuracy obtained when training the ANN on two subsets of all motifs, corresponding to the

removal of the 6-nodes motifs, and then the removal of both 5 and 6-nodes motifs.

Simulations of bipartite interaction networks

The accuracy of supervised classifiers often depends on the size of the available training

set (i.e., data for which the classification is known), which in the case of bipartite interaction

networks is rather small (n=343 here). A possible approach to increase the training set is to use

simulated data. We explored this possibility with the recently-developed BipartiteEvol model

(Maliet et al., 2020), an eco-evolutionary model that simulates the evolution of interacting

individuals within two clades (as detailed in Supplementary Note 1).

3. Results

Structure of antagonistic versus mutualistic networks

We found differences in macro-scale structures between antagonistic and mutualistic

networks.. For instance, nestedness was significantly higher in mutualistic networks (mean:

$2.16 \pm$ s.d. 0.71) than in antagonistic ones (1.94 ± 1.09; Wilcoxon-Mann-Whitney test: $W = 8673$,

p-value < 0.001), whereas antagonistic networks were significantly more connected ($0.23 \pm$

0.17) than mutualistic ones (0.15 ± 0.09; Wilcoxon-Mann-Whitney test: $W = 17709$, p-value $<$

0.001). The difference in terms of modularity was not significant (Wilcoxon-Mann-Whitney

test: W = 12559, p-value = 0.19) (Fig. 2A). In addition, we found that mutualistic networks were

composed of more species compared to antagonistic ones (76.1± 92.3 for antagonistic networks

and 82.7 ± 72.7 for mutualistic ones; Wilcoxon-Mann-Whitney test: W = 11124, p-value = 0.003).

These differences were however better explained by specific ecologies rather than the overall

interaction type. Specifically, when controlling for the different types of ecology (*e.g.,*

pollination, herbivory, seed dispersal...), antagonistic and mutualistic networks were not

significantly different in terms of their nestedness (Wald test: χ2 = 0.060, p-value = 0.81),

connectance (Wald test: χ2 = 0. 41, p-value = 0.52), modularity (Wald test: χ2 = 0.42, p-value =

0. 52), the total number of species (Wald test: χ2 = 0. 29, p-value = 0.59), and absolute difference

in the number of species per guild (Wald test: χ2 = 0. 95, p-value = 0.33). Model comparisons

confirmed that differences in terms of global metrics were better explained by the different

types of ecology than by the antagonistic/mutualistic type of interactions (Table S1). For

instance, compared to other types of ecology, bacteria-phage networks were more connected

but generally less nested, while herbivory networks were more nested (Fig. S1, Table S2 A-C).

Next, using the spectral density of the Laplacian graph (Fig. S2A) on perfectly modular

or nested networks revealed that modular networks have rather smooth densities with high

peaks whereas nested networks have noisier densities with several small peaks (Fig. S2B-E).

We found that mutualistic networks were enriched in eigenvalues near 1, while antagonistic

networks had significantly more spectral eigenvalues near 0 (Fig. 2B), which was mainly

driven by herbivory but not host-parasite networks (Fig. S3).

Finally, using motif frequencies, we found, following the motif denomination of

(Simmons, Sweering, et al., 2019), that antagonistic networks were significantly enriched in

densely connected motifs (such as motif 12 and motifs 31 to 37), whereas mutualistic networks

were enriched in motifs with asymmetrical specialization, wherein specialist species tend to

interact with the most generalist species in the motif (such as motifs 10, 19, 20, 25, 28 Fig. 2C).

In total, 24 out of the 44 motifs differed significantly in frequency between antagonistic and

mutualistic networks (p<0.01). When controlling for the type of ecology, 4 motifs were

significantly different in frequency between antagonistic and mutualistic networks (motifs 17,

25, 28, and 44). 31 out of 44 motifs differed significantly in frequency across the different types

of ecology (Fig. S4C): for example, seed-dispersal and host-parasite networks had relatively

high frequencies of 4- to 6-nodes motifs, in particular motifs 29 to 43, which may be linked to

their high connectance (Figs. S1, S4., Table S2A).

Unsupervised classifications of empirical networks

A principal component analysis (PCA) on the global metrics did not identify distinct clusters

for mutualistic and antagonistic networks (Fig. 3A). Consequently, an unsupervised K-means

classification on the principal component axes was not sufficient to confidently classify the

networks (F-scores of 0.63 and 0.55 for mutualistic and antagonistic networks respectively)

(Fig. 3A). Similarly, networks with different types of ecology (*e.g.* seed dispersal, herbivory,

ant-plant…) were largely mixed on the PCA projection (Fig. S5) and did not systematically fell

in one of the two clusters of the K-means classification (Table S3A).

PCAs performed on spectral density values (Fig. 3B) or motif frequencies (Fig. 3C) were

better at separating antagonistic *versus* mutualistic empirical networks than when using the

global metrics, but only slightly so (Fig. 3A). Antagonistic networks appeared to be spread on

the projection, while mutualistic networks were more clustered, which is potentially linked to

the high number of pollination networks in the database (Figs. S6-7). Unsupervised K-means

classifications on the principal component axes were not sufficient to confidently classify

mutualistic and antagonistic networks (Fig. 3B-C). Networks with different types of ecology

were largely mixed between the two clusters from the K-means classification (Tables S3 B-C),

suggesting that unsupervised classifications do not perform well at classifying ecological

networks, regardless of the metric used to characterize their structure.

Supervised classifications of empirical networks

We found that the two linear classifiers (Logit and Lasso regressions) mainly classified

empirical networks as mutualistic, regardless of their interaction type (Table S4). Artificial

neural networks (ANN) performed much better, in particular when using motif frequencies.

Motif frequencies led to better classifications than the spectral density or the global metrics,

and adding other variables to motif frequencies did not improve the classifications (Table S4).

The ANN based on motif frequencies classified an average of 65% of the antagonistic networks

(resp. 84% of the mutualistic networks) classified as antagonistic (resp. mutualistic), resulting

in an F-score of 0.68 and 0.82 for antagonistic and mutualistic networks respectively (Table S4).

Irrespectively of the bandwidth used to smooth the Laplacian spectrum, using the ANN with

the spectral density led to lower accuracies (Tables S4-5). We therefore used ANN on motif

frequencies for the remaining analyses.

Supervised classification of the empirical networks using motif frequencies

Using our measure of classification accuracy with ANN on motif frequencies, we found

that 69% (52% and 17% with high and low confidence respectively; F-score=0.74) of the

antagonistic networks and 89% (76% and 13% with high and low confidence respectively; F-

score=0.86) of the mutualistic networks were correctly classified (Table 1). In total, 81% (67%

and 14% with high and low confidence respectively) of the networks were correctly classified

(Table 1). The percentage of correct classifications varied according to the type of ecology.

Indeed, most bacteria-phage, seed dispersal, and pollination networks were correctly

classified by this approach (*e.g.* 23/29 of the seed dispersal networks were correctly classified,

among which ~85% with high confidence). Conversely, high intimacy mutualistic networks

(*i.e.* ant-plant and mycorrhizal networks) presented frequent misclassification (only 7 out of 12

networks (~60%) were classified as mutualistic), herbivory networks (which are low intimacy

antagonistic networks) were mainly classified as mutualistic (13/23, *i.e.* 57%), and 50% (5/10)

of the host-parasitoid networks were also misclassified (Table 1).

Robustness of the classification using motif frequencies

Overfitting had a limiting impact on the result just presented (Fig. S8). In addition,

controlling for the overrepresentation of mutualistic network in the database resulted in

similar mean percentage of correct classifications (Fig. S9A). Similarly, equalizing the number

of networks per type of ecology in the training set did not significantly affect the overall

percentages of correct classifications (Fig. S9B). Finally, we found lower classification

accuracies when randomizing the networks compared to the original classifications (Fig. S16).

Yet, classifications were not completely random: for instance, 56% (mean F-score = 0.57) and

77% (mean F-score = 0.76) of the antagonistic and mutualistic networks respectively were

correctly classified using randomized networks with fixed network size and connectance (Fig.

S16), suggesting that differences in network size, connectance, and/or degree distributions

between mutualistic and antagonistic networks partially contribute to our ability to tell

mutualistic and antagonistic ecological networks apart from their structure but fail to explain

the high classification accuracy we report. Besides, we found that removing a given type of

ecology during the ANN training decreased the percentage of correct classifications but

remained quite good (Table S7). For instance, 65% of the seed dispersal and 63% of the ant-

plant networks were correctly classified when none of these networks were used during

training, compared to 79% and 66% respectively when they were included (Table S7; Table 1).

Also, artificially shuffling the mutualistic or antagonistic nature of each type of ecology

decreased the mean percentages of correct classifications by about 10%, and the F-scores by

about 0.08, while increasing the difference in percentages of correct classifications and F-scores

between antagonistic and mutualistic networks by 30% and 0.3 respectively, which reflects a

major loss of the classification ability (Fig. S17). Hence, although the ANN classifiers partially

learned the singularities of the networks linked to ecology (*i.e.* data linkage), they also learned

the antagonistic/mutualistic dichotomy beyond ecology.

Finally, we found that the classification accuracy increased when including more

complex motifs (Table S9). The difference was most notable between analyses carried with

motifs of size at most 4 and of size at most 5 or 6 (Table S9), and rather insignificant between

analyses carried with motifs of size at most 5 and at most 6. This suggests that complex motifs

(with at least 5 nodes) are needed to distinguish between mutualistic and antagonistic

networks, but that adding even larger motifs would not significantly increase classification

accuracy.

Ability of simulated networks to improve the accuracy of classifiers

Using simulated networks did not improved our ability to distinguish apart

mutualistic from antagonistic networks both when using K-means classifications (results not

shown) and ANN (Table S8B). Mutualistic networks, in particular, were often wrongly

classified as antagonistic. When classifying simulated BipartiteEvol networks from the test

sets, 70% (F-score =0.71) of antagonistic networks and 58% (F-score = 0.57) of mutualistic ones

were accurately classified when using motif frequencies and ANN (Table S8A), suggesting

difficulties in separating antagonistic *versus* mutualistic networks simulated with

BipartiteEvol using ANN. When comparing the structure of simulated and empirical

networks, we found differences in global metrics, motif composition, and density of eigen-

values that may explain why simulations failed to improve the accuracy of classification

(Supplementary Note 2).

4. Discussion

In this study, we showed that discriminating antagonistic *versus* mutualistic ecological

networks based on their structure is overall possible. Supervised machine learning classifiers

such as artificial neural networks trained on motif frequencies reached an average of 81%

correct classifications (F-score of 0.74 and 0.86 for antagonistic and mutualistic networks

respectively). Structural differences between antagonistic and mutualistic networks can be

captured at the network mesoscale through differences in some motif frequencies, whereas the

macro-scale modular/nested dichotomy does not seem sufficient. Beyond the classical

dichotomy between mutualism/antagonism, we emphasize that the different types of ecology

have a important structural imprint on the structure of ecological networks.

Using motif frequencies and ANN, we were able to correctly classify around 81% of

the empirical interaction networks. This percentage of successful classifications is higher than

the success rate achieved by the only previous attempt we are aware of to classify networks

with ANN based exclusively on their structure (Michalska-Smith & Allesina, 2019), and

achieved without accounting for environmental conditions (as in Song & Saavedra, 2020). We

also demonstrated the relative specificity of our approach (*sensu* Michalska-Smith & Allesina,

2019): when randomizing the network structures, we obtained lower classification accuracies,

suggesting that motif frequencies, which depict patterns of species interactions at the meso-

scale, capture structural differences between mutualistic and antagonistic networks that are

not only due to differences in network size, connectance, or degree distributions. This

therefore indicates that there are significant structural differences at the network meso-scale,

which cannot be fully captured when looking at macro- or -micro-scales alone using global

metrics and/or Laplacian spectral densities.

Our analyses confirm that global metrics, including the modular/nested dichotomy, are

not sufficient to discriminate between mutualistic and antagonistic network structures

(Fontoura et al., 2020; Michalska-Smith & Allesina, 2019). We did not find a significant

difference in modularity values between mutualistic and antagonistic networks. We did find

that empirical mutualistic networks tend to be more nested and less connected than

antagonistic networks, as previously suggested by (Bascompte et al., 2003; Thébault &

Fontaine, 2010), but this dichotomy is not consistent enough (*i.e.,* there are too many

exceptions) to provide a robust classification criterion. There are several empirical examples

of exceptions to the nested/modular dichotomy in the literature, such as a nested structure in

host-parasite antagonistic networks (Pilosof et al., 2014). Within our database, some

pollination and mycorrhizal networks are known to be modular (Chagnon et al., 2015;

Jacquemyn et al., 2015; Martos et al., 2012; Olesen et al., 2007; Perez-Lamarque et al., 2022; Xing

et al., 2019), while they assemble mainly mutualistic interactions.

Why motifs performed better than the spectral density of the Laplacian graph is

unclear, as this latter representation is supposed to integrate all scales. Using a constant

bandwidth value to convert the Laplacian graph spectrum into a density may result in

information loss when comparing networks of different sizes. Also, transforming the discrete

Laplacian spectrum into a continuous density may be problematic for small networks with

only a few eigenvalues.

Motifs focus on precise structural patterns of interspecific interactions that can be

missed when summarizing all the information in a single global value at the network macro-

scale (Simmons, Cirtwill, et al., 2019). Densely connected motifs (*e.g.* motifs 12 and 31 to 37)

are significantly more frequent in antagonistic networks, while mutualistic networks are

characterized by motifs with asymmetrical specialization (*e.g.,* motifs 10, 19, 20, 25, 28). In

mutualistic networks, the emergence of motifs with asymmetrical specialization, which often

parallels the emergence of nestedness, has been related to three different types of explanations.

First, a pattern of asymmetric specialization may be partly explained from trait

complementarity and convergence. While generalist species can establish interactions with

broad range of mutualistic partners, trait complementarity generates specialization that

constrains the possible interactions of specialist species (Rezende, Jordano, et al., 2007;

Thompson, 2005). Second, such a pattern can emerge from differences in species abundances

(neutrality hypothesis): if interactions between individuals occur at random, rare specialist

species will be more likely to interact with abundant generalist species than with another rare

specialist (Vázquez et al., 2009). It has been proposed that a combination of trait

complementarity and abundance mechanisms best explains the observed structure of

mutualistic networks (Santamaría & Rodríguez-Gironés, 2007; Vázquez et al., 2009). Finally, it

has also been suggested that asymmetric specialization reduces interspecific competition

(Bastolla et al., 2009; Thompson, 2005), therefore enhancing the number of coexisting species

(Bascompte et al., 2006; Bastolla et al., 2009) and widening the conditions over which

mutualists coexist (Rohr et al., 2014), which in the end increases the persistence and resilience

of mutualistic networks to perturbations (Okuyama & Holland, 2008; Thébault & Fontaine,

2010). On the contrary, we expect antagonistic partners to engage in eco-evolutionary arm

races, which may result in modules formed by highly specialized within-module interactions

(i.e. densely connected motifs, which tend to be more frequent in antagonistic networks)

(Maliet et al., 2020; Thompson, 2005). Such modular structures would increase the resilience

of antagonistic communities (Thébault & Fontaine, 2010). Differences in motif frequencies can

also be influenced by sampling biases, including missing links or preferential detection of

some species. Future works will be required to identify the ecological processes and/or

sampling effects that generate such structural differences between mutualistic and

antagonistic networks (Brimacombe et al., 2023).

Mutualistic and antagonistic networks cover a wide range of ecologies (*e.g.* pollination,

parasitism, herbivory) and our analyses showed that the type of ecology matters in the

supervised classification, a phenomenon referred to as data linkage: networks with different

ecologies tend to have their own structural signatures beyond the mutualistic/antagonistic

dichotomy, and the ANN classifier partially learns to identify each type of ecology. In

particular, we showed that a better classification of networks of a particular type of ecology is

achieved by using networks from the same ecology during the training of the classifiers. This

is consistent with the result from linear mixed models that, after controlling for the type of

ecology, the only significant difference between mutualistic and antagonistic networks is the

frequency of a few motifs (motifs 17, 25, 28 and 44). These results on the importance of the type

of ecology corroborate previous studies that found, for instance, significant structural

differences between pollination and seed dispersal networks (Mello et al., 2011) or between

mutualistic plant-fungus and plant-animal networks (Toju et al., 2015). This impact of the type

of ecology may also explain the low classification accuracy of some networks, like the

herbivory networks, which share similar structural properties with the pollination networks

in both spectral density and motifs frequency. Nevertheless, we also demonstrated by

shuffling the dataset that the high classification accuracy we obtained is achieved by learning

structural differences between antagonistic and mutualistic networks beyond the type of

ecology. Beyond the singularities of the type of ecology, there are structural differences

between mutualistic and antagonistic networks that can be captured using our meso-scale

approach and ANNs.

The accuracy of supervised classification approaches strongly depends on the size and

quality of the dataset used for training. We used one of the largest publicly available databases

of empirical networks, but it contained only 343 networks in total. In addition, it had a strong

unbalanced representation of the different types of ecology, with 80% of the mutualistic

networks represented by pollination networks and 60% of the antagonistic networks

represented by host-parasite networks. There was also a strong imbalance in the

representation of interaction intimacy, which is known to impact network structure (Fontaine

et al., 2011; Guimarães et al., 2007; Hembry et al., 2018; Pires & Guimarães, 2013): mutualistic

networks were mainly low intimacy networks, except ant-plants (n=3) or mycorrhizal

networks (n=9), and antagonistic networks were mainly high intimacy networks, except

herbivory networks (n=23). As a consequence, only ~60% of the high-intimacy mutualistic

networks and ~40% of the low-intimacy antagonistic networks were correctly classified. These

results confirm that interaction intimacy may confer a particular structure to ecological

networks (Fontaine et al., 2011; Guimarães et al., 2007). Increasing the number of empirical

networks, especially mutualistic networks with high intimacy (*e.g.* symbiotic networks) and

antagonistic networks with low intimacy (*e.g.* trophic networks) would improve further our

ability to classify ecological networks by better representing the large structural heterogeneity

of ecological networks (Michalska-Smith & Allesina, 2019).

The accuracy of supervised classification approaches can also strongly depend on the

type, architecture, and tuning of the classifier. For example, the overfitting we noticed in the

ANN (which did not seem to negatively impact our results) could potentially be avoided by

using other regularization technics, such as dropout (Srivastava et al., 2014). We chose to use

simple, few-layers neural networks implemented in R for their availability and ease of usage

to the large community of R users, as well as because our dataset for training was rather small.

However, several other machine learning tools exist and could be tested, although they may

require larger training sets; graph neural networks in particular have been specifically

designed to learn information from graphs such as ecological networks (Wu et al., 2021).

One approach to better train the ANNs would be to increase the size of the training

data thanks to synthetic data, obtained through either simulations or data augmentation. We

investigated the possibility to train from simulated data with BipartiteEvol, an individual-

based model that mimics the eco-evolutionary emergence of bipartite interaction networks

(Maliet et al., 2020). However, ANN classifiers trained on BipartiteEvol simulations were not

able to discriminate mutualistic and antagonistic empirical networks, suggesting that the

model somehow fails at capturing important structural differences between antagonistic and

mutualistic networks. (Maliet et al., 2020) showed that the model produces realistic networks,

which we confirm here, and is able to reproduce major global differences observed between

mutualistic and antagonistic networks. However, we find that BipartiteEvol simulations tend

to have higher frequencies of complex motifs and a lower connectance compared to empirical

networks. Consequently, their projection only covers a small proportion of the space occupied

by empirical networks. One possible explanation is the continuum between mutualism and

antagonism in many interactions found in natural communities (Montesinos-Navarro et al.,

2017). For instance, it has been shown that parrot-plant networks can combine the two types

of interactions (Montesinos-Navarro et al., 2017). Similarly, pollination or mycorrhizal

networks often contain cheating species that use antagonistic strategies and induce structural

changes within the interaction networks (Genini et al., 2010; Klironomos, 2003; Perez-

Lamarque et al., 2020). Interactions can also shift from mutualism to antagonism on short

timescales, depending on some external conditions (Canestrari et al., 2014; Sachs et al., 2011).

By contrast, BipartiteEvol models the emergence of a network in strict antagonistic or

mutualistic communities. This calls for future efforts to improve the BipartiteEvol model or to

use simulations under other generative models (*e.g.,* (Cai et al., 2020; Thebault & Fontaine,

2010). Alternatively, data augmentation methods could also be tested to generate synthetic

networks by randomly subsampling, mixing and rewiring empirical networks; but whether

such approaches produce realistic interaction networks would need to be tested first.

Ultimately, supervised classifying approaches could be particularly useful to predict

the type of undetermined interactions. For instance, many plant-fungus interaction networks

are assembled from high-throughput sequencing data, but we currently ignore the mutualistic

versus the antagonistic type of many of these endophytic interactions (Chagnon et al., 2016;

Selosse et al., 2018). Similarly, an increasing number of networks are built from co-occurrence

data generated with high-throughput sequencings, such as the ones between microscopic

planktonic species sampled by the *Tara Oceans* expeditions (Chaffron et al., 2021; Lima-

Mendez et al., 2015). The ecology of the majority of species in these networks is currently

undetermined (Bjorbækmo et al., 2020; Lima-Mendez et al., 2015; Strydom et al., 2021).

However, a preliminary analysis of bipartite networks built from co-occurrence data in the

Tara Oceans data revealed that these networks were structurally very different from the

empirical and simulated networks we used here (results not shown). Classifying these

networks would thus require using a specific training dataset composed of microbial networks

with known interaction types (as discussed in Supplementary Note 3).

Conclusion

Mutualistic and antagonistic networks have structural differences that can be

particularly well captured by motif frequencies. Antagonistic networks are enriched in

densely connected motifs, while mutualistic networks are enriched in motifs with

asymmetrical specialization. Besides this dichotomy, there are structural differences across

networks with distinct ecologies. These structural differences can be used to classify bipartite

interaction networks with supervised machine learning with pretty good accuracy. These

performances could be further improved by increasing the number of empirical networks used

for training, diversifying the type of ecology sampled, and improving currently available

models to simulate networks with realistic structures. While more work is needed to predict

interaction types for co-occurrence networks, our analyses provide promising results on the

ability to predict interaction types from network structure.

Table 1. The type of interaction of the empirical ecological networks can be predicted based on their

motif frequencies using artificial neural network classifiers.

The following results were obtained by training the artificial neural networks (ANN) with 10,000

different training and test sets, using motif frequencies as the input structural variables. From those

10,000 classifications, we calculated the percentage of antagonistic *versus* mutualistic classifications for

each network: a network was predicted as a mutualistic (resp. antagonistic) network with high

confidence (highlighted in grey) if it was classified as a mutualistic (resp. antagonistic) network in more

than 75% of the classifications. It was predicted as a mutualistic (resp. antagonistic) network with low

confidence if it was classified as a mutualistic (resp. antagonistic) network in more than 50% but less

than 75% of the classifications. We therefore obtained a predicted type of interaction for each of the 343

empirical networks.

The second column indicates the number of interaction networks in our database gathered for each

ecology of interaction. Then, we indicate the number (and the percentage) of networks classified as

either mutualistic or antagonistic for each ecology of interaction and at the bottom, we show the global

contingency table for all mutualistic and antagonistic networks. We also indicate the F-score associated

with the global contingency table for each antagonistic or mutualistic type of interaction. Rows are

ordered based on the % of networks classified as mutualistic within each type of ecology. In the end, 88

antagonistic and 191 mutualistic networks are correctly classified, i.e., 279 of the 343 empirical networks

(81%).

Network ecology	Number of networks	Prediction			
		Antagonistic network (high confidence)	**Antagonistic network (low confidence)**	**Mutualistic network (high confidence)**	**Mutualistic network (low confidence)**
Anemone-Fish (mutualistic)	1	0 (0%)	0 (0%)	1 (100%)	0 (0%)
Pollination (mutualistic)	174	7 (4%)	7 (4%)	139 (80%)	21 (12%)
Seed dispersal (mutualistic)	29	6 (21%)	0 (0%)	20 (69%)	3 (10%)
Ant-Plant (mutualistic)	3	0 (0%)	1 (33%)	1 (33%)	1 (33%)
Herbivory (antagonistic)	23	4 (17.5%)	6 (26%)	9 (39%)	4 (17.5%)
Mycorrhiza (mutualistic)	9	3 (33.5%)	1 (11%)	3 (33.5%)	2 (22%)
Host-Parasitoid (antagonistic)	10	4 (40%)	1 (10%)	4 (40%)	1 (10%)
Bacteria-Phage (antagonistic)	17	12 (70%)	1 (6%)	3 (18%)	1 (6%)
Host-Parasitism (antagonistic)	77	46 (60%)	14 (18%)	10 (13%)	7 (9%)
Antagonistic networks	127	**66 (52%)**	**22 (17%)**	26 (21%)	13 (10%)
Mutualistic networks	216	16 (7%)	9 (4%)	**164 (76%)**	**27 (13%)**
Correct classifications (%) F-score		88 (69%) F-score = 0.74		191 (88%) F-score = 0.86	

Figures:

 Figure 1: Multi-scale structure of bipartite networks

 The structure of bipartite ecological networks, represented here by an ant-plant network

 (Blüthgen et al., 2004) can be studied at the macro-, meso-, and micro-scale. The Laplacian

 spectral density corresponds to a multi-scale approach from the macro to the micro-scale,

 whereas global metrics and motif frequencies focus on macro- and meso-scales, respectively.

 Blue nodes represent plant species, brown nodes ant species, and grey edges interspecific

 interactions.

Figure 2. Differences between antagonistic and mutualistic networks across scales of network

structure.

(A) Connectance, nestedness (NODFc), and modularity (Newman's index) of antagonistic and

mutualistic empirical networks. Stars indicate the statistical significance of the difference between

antagonistic and mutualistic networks, as measured by a Wilcoxon-Mann-Whitney test (*** = p-value <

0.01, ns = non-significant).

(B) Densities of eigenvalues from the Laplacian spectral density that differ significantly between

antagonistic and mutualistic networks (Wilcoxon-Mann-Whitney test: p-value<0.01). The

discontinuities in the x-axis are due to the filtering of eigenvalues densities that differ significantly

between antagonistic and mutualistic networks.

(C) Frequencies of the motifs that differ significantly between antagonistic and mutualistic networks

(Wilcoxon-Mann-Whitney test: p-value<0.01). Each motif is illustrated at the top of each boxplot

(following the denomination of Simmons *et al.* 2018).

Mutualistic networks are shown in green and antagonistic ones in yellow. Boxplots present the median

surrounded by the first and third quartiles, and whiskers extend to the extreme values but no further

than 1.5 of the interquartile range.

Figure 3. Mesoscale network structure as a scale to discriminate mutualistic *from* antagonistic

networks.

Projection of the 343 empirical networks on the two principal components (PC1 and PC2) obtained using

principal coordinate analysis (PCA) on (A) global metrics (*i.e.* nestedness, modularity, connectance, and

network size), (B) spectral density of the Laplacian graph and (C) motif frequencies. The percentage of

explained variance is indicated on each axis.

Each associated table corresponds to the contingency table of unsupervised K-means classification

based on the global metrics (A), spectral density of the Laplacian graph (B), and motifs frequencies (C).

Chi-squared test associated with the K-means classification is indicated.

References

Banerjee, A., & Jost, J. (2008). On the spectrum of the normalized graph Laplacian. *Linear Algebra and*

Its Applications, *428*(11–12), 3015–3022. https://doi.org/10.1016/j.laa.2008.01.029

Bascompte, J., & Jordano, P. (2007). Plant-animal mutualistic networks: The architecture of

biodiversity. *Annual Review of Ecology, Evolution, and Systematics*, *38*(1), 567–593.

https://doi.org/10.1146/annurev.ecolsys.38.091206.095818

Bascompte, J., Jordano, P., Melian, C. J., & Olesen, J. M. (2003). The nested assembly of plant-animal

mutualistic networks. *Proceedings of the National Academy of Sciences*, *100*(16), 9383–9387.

https://doi.org/10.1073/pnas.1633576100

Bascompte, J., Jordano, P., & Olesen, J. M. (2006). Asymmetric coevolutionary networks facilitate

biodiversity maintenance. *Science*, *312*(5772), 431–433.

https://doi.org/10.1126/science.1123412

Bastolla, U., Fortuna, M. A., Pascual-García, A., Ferrera, A., Luque, B., & Bascompte, J. (2009). The

architecture of mutualistic networks minimizes competition and increases biodiversity.

Nature, *458*(7241), 1018–1020. https://doi.org/10.1038/nature07950

Bjorbækmo, M. F. M., Evenstad, A., Røsæg, L. L., Krabberød, A. K., & Logares, R. (2020). The planktonic

protist interactome: Where do we stand after a century of research? *The ISME Journal*, *14*(2),

544–559. https://doi.org/10.1038/s41396-019-0542-5

Blanchet, F. G., Cazelles, K., & Gravel, D. (2020). Co-occurrence is not evidence of ecological

interactions. *Ecology Letters*, *23*(7), 1050–1063. https://doi.org/10.1111/ele.13525

Brimacombe, C., Bodner, K., Michalska-Smith, M., Poisot, T., & Fortin, M.-J. (2023). Shortcomings of

reusing species interaction networks created by different sets of researchers. *PLOS Biology*,

21(4), e3002068. https://doi.org/10.1371/journal.pbio.3002068

Bronstein, J. L. (Ed.). (2015). *Mutualism*. Oxford University Press.

https://doi.org/10.1093/acprof:oso/9780199675654.001.0001

Cai, W., Snyder, J., Hastings, A., & D'Souza, R. M. (2020). Mutualistic networks emerging from adaptive

niche-based interactions. *Nature Communications*, *11*(1), 5470.

https://doi.org/10.1038/s41467-020-19154-5

Canestrari, D., Bolopo, D., Turlings, T. C. J., Roder, G., Marcos, J. M., & Baglione, V. (2014). From

parasitism to mutualism: Unexpected interactions etween a cuckoo and Its host. *Science*,

343(6177), 1350–1352. https://doi.org/10.1126/science.1249008

Chaffron, S., Delage, E., Budinich, M., Vintache, D., Henry, N., Nef, C., Ardyna, M., Zayed, A. A., Junger,

P. C., Galand, P. E., Lovejoy, C., Murray, A. E., Sarmento, H., null, null, Acinas, S. G., Babin, M.,

Iudicone, D., Jaillon, O., Karsenti, E., … Eveillard, D. (2021). Environmental vulnerability of the

global ocean epipelagic plankton community interactome. *Science Advances*, *7*(35), eabg1921.

https://doi.org/10.1126/sciadv.abg1921

Chagnon, P.-L. (2016). Seeing networks for what they are in mycorrhizal ecology. *Fungal Ecology*, *24*,

148–154. https://doi.org/10.1016/j.funeco.2016.05.004

Chagnon, P.-L., Bradley, R. L., & Klironomos, J. N. (2015). Trait-based partner selection drives

mycorrhizal network assembly. *Oikos*, *124*(12), 1609–1616.

https://doi.org/10.1111/oik.01987

Chagnon, P.-L., U'Ren, J. M., Miadlikowska, J., Lutzoni, F., & Arnold, A. E. (2016). Interaction type

influences ecological network structure more than local abiotic conditions: Evidence from

endophytic and endolichenic fungi at a continental scale. *Oecologia*, *180*(1), 181–191.

https://doi.org/10.1007/s00442-015-3457-5

de Lange, S. C., de Reus, M. A., & van den Heuvel, M. P. (2014). The Laplacian spectrum of neural

networks. *Frontiers in Computational Neuroscience*, *7*.

https://doi.org/10.3389/fncom.2013.00189

Dormann, C. F., Gruber, B., & Fründ, J. (2008). *Introducing the bipartite Package: Analysing Ecological*

Networks. *8*, 5.

Fontaine, C., Guimarães, P. R., Kéfi, S., Loeuille, N., Memmott, J., van der Putten, W. H., van Veen, F. J.

F., & Thébault, E. (2011). The ecological and evolutionary implications of merging different

types of networks: Merging networks with different interaction types. *Ecology Letters*, *14*(11),

1170–1181. https://doi.org/10.1111/j.1461-0248.2011.01688.x

Fontoura, L., Cantor, M., Longo, G. O., Bender, M. G., Bonaldo, R. M., & Floeter, S. R. (2020). The

macroecology of reef fish agonistic behaviour. *Ecography*, ecog.05079.

https://doi.org/10.1111/ecog.05079

Fortuna, M. A., Ortega, R., & Bascompte, J. (2014). The Web of Life. *arXiv:1403.2575 [q-Bio]*.

http://arxiv.org/abs/1403.2575

Fortuna, M. A., Stouffer, D. B., Olesen, J. M., Jordano, P., Mouillot, D., Krasnov, B. R., Poulin, R., &

Bascompte, J. (2010). Nestedness versus modularity in ecological networks: Two sides of the

same coin? *Journal of Animal Ecology*. https://doi.org/10.1111/j.1365-2656.2010.01688.x

Freilich, S., Kreimer, A., Meilijson, I., Gophna, U., Sharan, R., & Ruppin, E. (2010). The large-scale

organization of the bacterial network of ecological co-occurrence interactions. *Nucleic Acids*

Research, *38*(12), 3857–3868. https://doi.org/10.1093/nar/gkq118

Friedman, J., Hastie, T., & Tibshirani, R. (2010). Regularization paths for generalized linear models via

coordinate descent. *Journal of Statistical Software*, *33*(1).

https://doi.org/10.18637/jss.v033.i01

Genini, J., Morellato, L. P. C., Guimarães, P. R., & Olesen, J. M. (2010). Cheaters in mutualism networks.

Biology Letters, *6*(4), 494–497. https://doi.org/10.1098/rsbl.2009.1021

Guimarães, P. R., Rico-Gray, V., Oliveira, P. S., Izzo, T. J., dos Reis, S. F., & Thompson, J. N. (2007).

Interaction intimacy affects structure and coevolutionary dynamics in mutualistic networks.

Current Biology, *17*(20), 1797–1803. https://doi.org/10.1016/j.cub.2007.09.059

Günther, F., & Fritsch, S. (2010). neuralnet: Training of Neural Networks. *The R Journal*, *2*(1), 30.

https://doi.org/10.32614/RJ-2010-006

Hembry, D. H., Raimundo, R. L. G., Newman, E. A., Atkinson, L., Guo, C., Guimarães, P. R., & Gillespie,

R. G. (2018). Does biological intimacy shape ecological network structure? A test using a brood

pollination mutualism on continental and oceanic islands. *Journal of Animal Ecology*, *87*(4),

1160–1171. https://doi.org/10.1111/1365-2656.12841

Hoeppke, C., & Simmons, B. (2021). Maxnodf: An R package for fair and fast comparisons of nestedness

between networks. *Methods in Ecology and Evolution*, *12*. https://doi.org/10.1111/2041-

210x.13545

Jacquemyn, H., Brys, R., Waud, M., Busschaert, P., & Lievens, B. (2015). Mycorrhizal networks and

coexistence in species-rich orchid communities. *New Phytologist*, *206*(3), 1127–1134.

https://doi.org/10.1111/nph.13281

Jacquemyn, H., Merckx, V., Brys, R., Tyteca, D., Cammue, B. P. A., Honnay, O., & Lievens, B. (2011).

Analysis of network architecture reveals phylogenetic constraints on mycorrhizal specificity in

the   genus   Orchis   (Orchidaceae).   *New   Phytologist*,   *192*(2),   518–528.

https://doi.org/10.1111/j.1469-8137.2011.03796.x

Jordano, P., Bascompte, J., & Olesen, J. M. (2002). Invariant properties in coevolutionary networks of

plant-animal interactions: Invariant properties in coevolutionary networks. *Ecology Letters*,

6(1), 69–81. https://doi.org/10.1046/j.1461-0248.2003.00403.x

Klironomos, J. N. (2003). Variation in plant response to native and exotic arbuscular mycorrhizal fungi.

Ecology, *84*(9), 2292–2301. https://doi.org/10.1890/02-0413

Krause, A. E., Frank, K. A., Mason, D. M., Ulanowicz, R. E., & Taylor, W. W. (2003). Compartments

revealed   in   food-web   structure.   *Nature*,   *426*(6964),   282–285.

https://doi.org/10.1038/nature02115

Lê, S., Josse, J., & Husson, F. (2008). FactoMineR : an R package for multivariate analysis. *Journal of

Statistical Software*, *25*(1). https://doi.org/10.18637/jss.v025.i01

Lewitus, E., & Morlon, H. (2016). Characterizing and comparing phylogenies from their Laplacian

spectrum. *Systematic Biology*, *65*(3), 495–507. https://doi.org/10.1093/sysbio/syv116

Lima-Mendez, G., Faust, K., Henry, N., Decelle, J., Colin, S., Carcillo, F., Chaffron, S., Ignacio-Espinosa, J.

C., Roux, S., Vincent, F., Bittner, L., Darzi, Y., Wang, J., Audic, S., Berline, L., Bontempi, G.,

Cabello, A. M., Coppola, L., Cornejo-Castillo, F. M., … Raes, J. (2015). Determinants of

community structure in the global plankton interactome. *Science*, *348*(6237), 1262073–

1262073. https://doi.org/10.1126/science.1262073

Losapio, G., Schöb, C., Staniczenko, P. P. A., Carrara, F., Palamara, G. M., De Moraes, C. M., Mescher,

M. C., Brooker, R. W., Butterfield, B. J., Callaway, R. M., Cavieres, L. A., Kikvidze, Z., Lortie, C. J.,

Michalet, R., Pugnaire, F. I., & Bascompte, J. (2021). Network motifs involving both competition

and facilitation predict biodiversity in alpine plant communities. *Proceedings of the National*

Academy of Sciences, *118*(6), e2005759118. https://doi.org/10.1073/pnas.2005759118

Maliet, O., Loeuille, N., & Morlon, H. (2020). An individual-based model for the eco-evolutionary

emergence of bipartite interaction networks. *Ecology Letters*, ele.13592.

https://doi.org/10.1111/ele.13592

Martos, F., Munoz, F., Pailler, T., Kottke, I., Gonneau, C., & Selosse, M.-A. (2012). The role of epiphytism

in architecture and evolutionary constraint within mycorrhizal networks of tropical orchids.

Molecular Ecology, *21*(20), 5098–5109. https://doi.org/10.1111/j.1365-294X.2012.05692.x

Mello, M. A. R., Marquitti, F. M. D., Guimarães, P. R., Kalko, E. K. V., Jordano, P., & de Aguiar, M. A. M.

(2011). The missing part of seed dispersal networks: Structure and robustness of bat-fruit

interactions. *PLoS ONE*, *6*(2), e17395. https://doi.org/10.1371/journal.pone.0017395

Michalska-Smith, M. J., & Allesina, S. (2019). Telling ecological networks apart by their structure: A

computational challenge. *PLOS Computational Biology*, *15*(6), e1007076.

https://doi.org/10.1371/journal.pcbi.1007076

Milo, R., Shen-Orr, S., Itzkovitz, S., Kashtan, N., Chklovskii, D., & Alon, U. (2002). Network motifs: Simple

building blocks of complex networks. *Science*, *298*(5594), 824–827.

Montesinos-Navarro, A., Hiraldo, F., Tella, J. L., & Blanco, G. (2017). Network structure embracing
mutualism–antagonism continuums increases community robustness. *Nature Ecology &*
Evolution, *1*(11), 1661–1669. https://doi.org/10.1038/s41559-017-0320-6

Montesinos-Navarro, A., Segarra-Moragues, J. G., Valiente-Banuet, A., & Verdú, M. (2012). Plant
facilitation occurs between species differing in their associated arbuscular mycorrhizal fungi.
New Phytologist, *196*(3), 835–844. https://doi.org/10.1111/j.1469-8137.2012.04290.x

Morlon, H., Lewitus, E., Condamine, F. L., Manceau, M., Clavel, J., & Drury, J. (2016). RPANDA : an R
package for macroevolutionary analyses on phylogenetic trees. *Methods in Ecology and*
Evolution, *7*(5), 589–597. https://doi.org/10.1111/2041-210X.12526

Newman, M. E. J. (2006a). Finding community structure in networks using the eigenvectors of matrices.
Physical Review E, *74*(3), 036104. https://doi.org/10.1103/PhysRevE.74.036104

Newman, M. E. J. (2006b). Modularity and community structure in networks. *Proceedings of the*
National   Academy   of   Sciences,   *103*(23),   8577–8582.
https://doi.org/10.1073/pnas.0601602103

Okuyama, T., & Holland, J. N. (2008). Network structural properties mediate the stability of mutualistic
communities.   *Ecology   Letters*,   *11*(3),   208–216.   https://doi.org/10.1111/j.1461-
0248.2007.01137.x

Olesen, J. M., Bascompte, J., Dupont, Y. L., & Jordano, P. (2007). The modularity of pollination
networks. *Proceedings of the National Academy of Sciences*, *104*(50), 19891–19896.
https://doi.org/10.1073/pnas.0706375104

Öpik, M., Metsis, M., Daniell, T. J., Zobel, M., & Moora, M. (2009). Large-scale parallel 454 sequencing
reveals host ecological group specificity of arbuscular mycorrhizal fungi in a boreonemoral
forest. *New Phytologist*, *184*(2), 424–437. https://doi.org/10.1111/j.1469-8137.2009.02920.x

Perez-Lamarque, B., Petrolli, R., Strullu-Derrien, C., Strasberg, D., Morlon, H., Selosse, M.-A., & Martos,
F. (2022). Structure and specialization of mycorrhizal networks in phylogenetically diverse

tropical communities. *Environmental Microbiome*, *17*(1), 38. https://doi.org/10.1186/s40793-
022-00434-0

Perez-Lamarque, B., Selosse, M., Öpik, M., Morlon, H., & Martos, F. (2020). Cheating in arbuscular
mycorrhizal mutualism: A network and phylogenetic analysis of mycoheterotrophy. *New*
Phytologist, *226*(6), 1822–1835. https://doi.org/10.1111/nph.16474

Pilosof, S., Fortuna, M. A., Cosson, J.-F., Galan, M., Kittipong, C., Ribas, A., Segal, E., Krasnov, B. R.,
Morand, S., & Bascompte, J. (2014). Host–parasite network structure is associated with
community-level immunogenetic diversity. *Nature Communications*, *5*(1), 5172.
https://doi.org/10.1038/ncomms6172

Pires, M. M., & Guimarães, P. R. (2013). Interaction intimacy organizes networks of antagonistic
interactions in different ways. *Journal of The Royal Society Interface*, *10*(78), 20120649.
https://doi.org/10.1098/rsif.2012.0649

Põlme, S., Bahram, M., Jacquemyn, H., Kennedy, P., Kohout, P., Moora, M., Oja, J., Öpik, M., Pecoraro,
L., & Tedersoo, L. (2018). Host preference and network properties in biotrophic plant-fungal
associations. *New Phytologist*, *217*(3), 1230–1239. https://doi.org/10.1111/nph.14895

Rezende, E. L., Jordano, P., & Bascompte, J. (2007). Effects of phenotypic complementarity and
phylogeny on the nested structure of mutualistic networks. *Oikos*, *116*(11), 1919–1929.
https://doi.org/10.1111/j.0030-1299.2007.16029.x

Rezende, E. L., Lavabre, J. E., Guimarães, P. R., Jordano, P., & Bascompte, J. (2007). Non-random
coextinctions in phylogenetically structured mutualistic networks. *Nature*, *448*(7156), 925–
928. https://doi.org/10.1038/nature05956

Rohr, R. P., Saavedra, S., & Bascompte, J. (2014). On the structural stability of mutualistic systems.
Science, *345*(6195), 1253497–1253497. https://doi.org/10.1126/science.1253497

Sachs, J. L., Skophammer, R. G., & Regus, J. U. (2011). Evolutionary transitions in bacterial symbiosis.
Proceedings of the National Academy of Sciences, *108*(Supplement_2), 10800–10807.
https://doi.org/10.1073/pnas.1100304108

Santamaría, L., & Rodríguez-Gironés, M. A. (2007). Linkage Rules for Plant–Pollinator Networks: Trait

Complementarity   or   Exploitation   Barriers?   *PLoS   Biology*,   *5*(2),   e31.

https://doi.org/10.1371/journal.pbio.0050031

Selosse, M., Schneider-Maunoury, L., & Martos, F. (2018). Time to re-think fungal ecology? Fungal

ecological   niches   are   often   prejudged.   *New   Phytologist*,   *217*,   968–972.

https://doi.org/10.1111/nph.14983

Simmons, B. I., Cirtwill, A. R., Baker, N. J., Wauchope, H. S., Dicks, L. V., Stouffer, D. B., & Sutherland,

W. J. (2019). Motifs in bipartite ecological networks: Uncovering indirect interactions. *Oikos*,

128(2), 154–170. https://doi.org/10.1111/oik.05670

Simmons, B. I., Sweering, M. J. M., Schillinger, M., Dicks, L. V., Sutherland, W. J., & Clemente, R. D.

(2018). *bmotif: A package for motif analyses of bipartite networks* [Preprint]. Ecology.

https://doi.org/10.1101/302356

Simmons, B. I., Sweering, M. J. M., Schillinger, M., Dicks, L. V., Sutherland, W. J., & Di Clemente, R.

(2019). bmotif: A package for motif analyses of bipartite networks. *Methods in Ecology and*

Evolution, *10*(5), 695–701. https://doi.org/10.1111/2041-210X.13149

Song, C., & Saavedra, S. (2020). Telling ecological networks apart by their structure: An environment-

dependent   approach.   *PLOS   Computational   Biology*,   *16*(4),   e1007787.

https://doi.org/10.1371/journal.pcbi.1007787

Srivastava, N., Hinton, G., Krizhevsky, A., Sutskever, I., & Salakhutdinov, R. (2014). Dropout: A simple

way to prevent neural networks from overfitting. *Journal of Machine Learning Research*,

15(56), 1929–1958.

Stouffer, D. B., & Bascompte, J. (2010). Understanding food-web persistence from local to global scales.

Ecology Letters, *13*(2), 154–161. https://doi.org/10.1111/j.1461-0248.2009.01407.x

Strydom, T., Catchen, M. D., Banville, F., Caron, D., Dansereau, G., Desjardins-Proulx, P., Forero-Muñoz,

N. R., Higino, G., Mercier, B., Gonzalez, A., Gravel, D., Pollock, L., & Poisot, T. (2021). A roadmap

towards predicting species interaction networks (across space and time). *Philosophical*

Transactions of the Royal Society B: Biological Sciences, *376*(1837), 20210063.

https://doi.org/10.1098/rstb.2021.0063

Suweis, S., Simini, F., Banavar, J. R., & Maritan, A. (2013). Emergence of structural and dynamical

properties of ecological mutualistic networks. *Nature*, *500*(7463), 449–452.

https://doi.org/10.1038/nature12438

Thébault, E., & Fontaine, C. (2008). Does asymmetric specialization differ between mutualistic and

trophic networks? *Oikos*, *117*(4), 555–563. https://doi.org/10.1111/j.0030-1299.2008.16485.x

Thébault, E., & Fontaine, C. (2010). Stability of ecological communities and the architecture of

mutualistic and trophic networks. *Science*, *329*(5993), 853–856.

https://doi.org/10.1126/science.1188321

Thebault, E., & Fontaine, C. (2010). Stability of ecological communities and the architecture of

mutualistic and trophic networks. *Science*, *329*(5993), 853–856.

https://doi.org/10.1126/science.1188321

Thomas, F., Renaud, F., & Guegan, J.-F. (Eds.). (2005). *Parasitism and Ecosystems*. Oxford University

Press. https://doi.org/10.1093/acprof:oso/9780198529873.001.0001

Thompson, J. N. (2005). *The Geographic Mosaic of Coevolution*. University of Chicago Press.

https://doi.org/10.7208/9780226118697

Toju, H., Guimarães, P. R., Olesen, J. M., & Thompson, J. N. (2015). Below-ground plant–fungus network

topology is not congruent with above-ground plant–animal network topology. *Science*

Advances, *1*(9), e1500291. https://doi.org/10.1126/sciadv.1500291

Vázquez, D. P., Chacoff, N. P., & Cagnolo, L. (2009). Evaluating multiple determinants of the structure

of plant–animal mutualistic networks. *Ecology*, *90*(8), 2039–2046. https://doi.org/10.1890/08-

1837.1

Vitali, A., Sasal, Y., Vázquez, D. P., Miguel, M. F., & Rodríguez-Cabal, M. A. (2022). The disruption of a

keystone interaction erodes pollination and seed dispersal networks. *Ecology*, *103*(1).

https://doi.org/10.1002/ecy.3547

Wu, Z., Pan, S., Chen, F., Long, G., Zhang, C., & Yu, P. S. (2021). A comprehensive survey on graph neural

networks. *IEEE Transactions on Neural Networks and Learning Systems*, *32*(1), 4–24.

https://doi.org/10.1109/TNNLS.2020.2978386

Xing, X., Jacquemyn, H., Gai, X., Gao, Y., Liu, Q., Zhao, Z., & Guo, S. (2019). The impact of life form on

the architecture of orchid mycorrhizal networks in tropical forest. *Oikos*, *128*(9), 1254–1264.

https://doi.org/10.1111/oik.06363

1010