# Telling mutualistic and antagonistic ecological networks apart by learning their multiscale structure

Running headline: Multiscale network structure and interaction type

Benoît Pichon[1,2,*] (ORCID : 0009-0007-6206-5482)

Rémy Le Goff [2] (ORCID : 0009-0002-7827-7096)

Hélène Morlon[2, #] (ORCID : 0000-0002-3195-7521)

Benoît Perez-Lamarque[2,3, #] (ORCID : 0000-0001-7112-7197)

[1] ISEM, Univ Montpellier, CNRS, EPHE, IRD, F-34095 Montpellier, France

[2] Institut de biologie de l'École normale supérieure (IBENS), École normale supérieure, CNRS, INSERM, Université PSL, 46 rue d'Ulm, 75 005 Paris, France

[3] Institut de Systématique, Évolution, Biodiversité (ISYEB), Muséum national d'histoire naturelle, CNRS, Sorbonne Université, EPHE, UA, CP39, 57 rue Cuvier 75 005 Paris, France

*Corresponding author: benoit.pichon0@gmail.com*

[#] B.P.L and H.M. jointly supervised this work.

## Acknowledgments

We thank I. Overcast, J. Voznica, L. Aristide, S. Lambert, I. Quintero, C. Fruciano, J. Clavel, A. Silva, for comments on the first version of the manuscript. We also thank S. Chaffron, C. Bowler, C. Nef, and E. Thébault for fruitful discussions. We thank the three reviewers for their comments that improved the manuscript.

The authors declare no conflict of interest.

## Author contributions

All authors conceived the ideas and designed methodology; B.P performed research. B.P, H.M and B.P.L led the writing of the manuscript. All authors contributed critically to the drafts and gave final approval for publication.

## Data availability

The code used for the analyses is available on Zenodo: https://doi.org/10.5281/zenodo.7786771

# Abstract

1.  Characterizing and understanding the processes that shape the structure of ecological networks, which represent who interacts with whom in a community, has many implications in ecology, evolutionary biology, and conservation. A highly debated question is whether and how the structure of a bipartite ecological network differs between antagonistic (e.g., herbivory) and mutualistic (e.g., pollination) interaction types.

2.  Here, we tackle this question by using a multiscale characterization of network structure, machine learning tools, and a large database of empirical and simulated bipartite networks.

3.  Contrary to previous studies focusing on global structural metrics, such as nestedness and modularity, which concluded that antagonistic and mutualistic networks cannot be told apart from only their structure, we find that they can be told apart by combining a meso-scale characterization of their structure and supervised machine learning. Motif frequencies appear particularly informative, with an over-representation of densely connected motifs in antagonistic networks and of motifs with asymmetrical specialization in mutualistic networks. These characteristics can be used to predict interaction types with relatively good confidence. Beyond this classical mutualism/antagonism dichotomy, we also find significant structural uniqueness linked to specific ecologies (e.g., pollination, parasitism).

4.  Our results clarify structural differences between antagonistic and mutualistic networks and suggest the investigation of the structural uniqueness of specific ecologies as a promising approach for characterizing interactions beyond the coarse antagonistic/mutualistic dichotomy.

**Keywords:** ecological interactions, network structure, motif frequency, interaction classification, machine learning

# 1. Introduction

Species in ecosystems engage in multiple types of antagonistic or mutualistic interactions, such as predation, parasitism, pollination, seed dispersal, or mycorrhizal symbioses (Bronstein, 2015; Thomas et al., 2005). Such interactions between two groups of potentially taxonomically distant species are often represented by bipartite networks (Bascompte et al., 2003; Bascompte & Jordano, 2007; Fontaine et al., 2011). Studying the structure of these interaction networks is fundamental for unraveling the different processes behind their ecological organization (Bastolla et al., 2009; Fontaine et al., 2011; Suweis et al., 2013; Thompson, 2005), understanding their evolutionary dynamics, and prioritizing conservation efforts (Bastolla et al., 2009; Olesen et al., 2007; Rezende, Lavabre, et al., 2007).

Early studies on bipartite ecological networks, focused mainly on pollination and herbivory networks, suggested that mutualistic and antagonistic networks have different structures (Bascompte et al., 2003; Thébault & Fontaine, 2008, 2010). Antagonistic networks tend to have a modular structure, where species within modules preferentially interact with each other and rarely interact with species from other modules (Krause et al., 2003; Newman, 2006b). Conversely, mutualistic networks tend to have a nested structure, where specialist species mainly interact with generalists, and generalists form a core of interactions (Jordano et al., 2002; Rohr et al., 2014; Bascompte et al., 2003). Such structural differences have been observed in both terrestrial (Fontaine et al., 2011) and marine networks (Guimarães et al., 2007). However, several studies have challenged the generality of the modular/nested dichotomy (Chagnon, 2016; Fontoura et al., 2020; Michalska-Smith & Allesina, 2019; Pilosof et al., 2014). The dichotomy seems to apply only to highly connected networks (*i.e.* with a high proportion of realized interactions), whereas networks with lower connectance are often

simultaneously modular and nested (Fortuna et al., 2010). Recently, using a large and diverse

dataset of empirical networks with different types of ecology (*e.g.* ant-plant, bacteria-phage,

host-parasitoid,…), Michalska-Smith and Allesina (2019) showed that the structure of

ecological networks is extremely heterogeneous and that global measures of network structure

including nestedness and modularity are not sufficient to discriminate antagonistic and

mutualistic networks (Michalska-Smith & Allesina, 2019). Song and Saavedra (2020) suggested

that the confounding effect of environmental conditions indeed prevents telling antagonistic

and mutualistic networks apart from such global metrics unless environmental differences are

accounted for (Song & Saavedra, 2020).

Beyond the mutualistic/antagonistic dichotomy, the type of network ecology (*e.g.*, seed-

dispersal compared to pollination) may significantly impact network structure. In particular,

networks tend to be more modular when their interactions are more intimate, *i.e.* when the

degree of biological integration between interacting individuals is high, regardless of

interaction type (Fontaine et al., 2011; Guimarães et al., 2007; Hembry et al., 2018; Pires &

Guimarães, 2013). For instance, within mutualistic networks, intimate ant-plant networks

were found to be more modular than pollination or seed dispersal networks, the latter forming

the least intimate interactions resulting in networks with low specialization (Mello et al., 2011).

Within ant-plant networks, obligate symbiotic networks were found to be more specialized

and modular than nectar-mediated facultative ones (Blüthgen et al., 2007; Dáttilo et al., 2013).

More specific mutualistic associations between plants and mycorrhizal fungi also result in

more modular mycorrhizal networks (Perez-Lamarque et al. 2022). These few examples

illustrate that beyond the mutualistic/antagonistic dichotomy, the traits and ecologies of the

interacting species leave an imprint on the  network structures (Lewinsohn et al., 2006).

Macro-scale metrics of network structure, like nestedness and modularity, are not the only way to characterize network structure. The nestedness/modularity paradigm, as the main method to characterize bipartite network structure, has recently been questioned. Simmons, Cirtwill, et al., (2019) highlighted the limits of macro-scale metrics for capturing structural differences between bipartite networks and showed the usefulness of motifs (a small subset of species interactions exhibiting varying topologies), which capture the mesoscale structure (Milo et al., 2002). Motifs have been informative in various contexts, like the assessment of network stability, the resistance to biological invasions (Losapio et al., 2021; Milo et al., 2002; Stouffer & Bascompte, 2010; Vitali et al., 2022), or the characterization of the singular positions of functional groups (Lanuza et al., 2023). Another way to characterize the structure of interaction networks is through the spectral densities of the Laplacian graph representing each interaction network. Indeed, the eigenvectors of the Laplacian graph inform on species-level properties and can be used to identify keystone species within networks (Estrada, 2007), whereas the eigenvalues integrate information about the overall network structure, and have been linked to nestedness (Staniczenko et al., 2013) and modularity (Shen & Cheng, 2010). The spectral density of Laplacian graphs have been used in other fields, for instance, to compare structural properties of species phylogenies (Lewitus & Morlon, 2016) or brain networks (de Lange et al., 2014), but only marginally to study the structure of ecological networks (Michalska-Smith & Allesina, 2019). Since spectral densities of the Laplacian graph integrate the overall network structure, they may be informative to identify structural components that differ between antagonistic and mutualistic networks.

Here, we use a variety of network structural metrics combined with different machine

learning tools to (i) investigate if there are structural components of networks, and which ones,

that are distinct in mutualistic *versus* antagonistic networks, (ii) assess whether unsupervised

and supervised classification approaches can robustly predict interspecific interaction type

from network structure, (iii) evaluate whether currently available simulation tools for bipartite

networks can help improve the accuracy of the classifiers, and (iv) assess the structural signal

from specific ecologies (*e.g.,* seed-dispersal, pollination, parasitism). We compare the

predictive power of the different structural metrics and classification methods. Finally, we

discuss the implications of our findings for the understanding of natural communities.

# 2. Materials and methods

**Empirical interaction networks**

We analyzed 343 local or regional scales bipartite interaction networks with well-

known interaction types. We obtained these networks by downloading the networks from

(Michalska-Smith & Allesina, 2019), among which some come from the Web of Life database

(web-of-life.es (Fortuna et al., 2014)). We kept networks containing at least 10 species in each

guild, as the structure of small networks is difficult to characterize (Olesen et al., 2007), in

particular in terms of motif composition (preliminary results not shown). In addition, we

gathered 9 mycorrhizal networks that also fit this size criterion: 6 were collected from

individual papers (Jacquemyn et al., 2011, 2015; Martos et al., 2012; Montesinos-Navarro et al.,

2012; Öpik et al., 2009; Xing et al., 2019) and 3 from the database generated by (Põlme et al.,

2018). Among the 343 networks, 216 were mutualistic and 127 antagonistic. They comprised 9

different types of ecology: bacteria-phage (antagonistic; n=17), seed dispersal (mutualistic;

n=29), host-parasite (antagonistic; n=77), host-parasitoid (antagonistic; n=10), plant-

mycorrhiza (mutualistic; n=9), ant-plant (mutualistic; n=3), herbivory (antagonistic; n=23),

anemone-fish (mutualistic; n=1), and pollination (mutualistic; n=174). Each network was

converted into a binary matrix, taking the value 1 if species $i$ interacts with species $j$, and 0

otherwise, *i.e.* we only focused on the presence/absence of interactions, not on their

frequencies. The guild with the largest (resp. lowest) number of species was in rows (resp.

columns).

**Comparing the structure of mutualistic *versus* antagonistic networks**

Compared to the two previous studies comparing network structures between

mutualistic and antagonistic networks (Michalska-Smith & Allesina, 2019; Song & Saavedra,

2020), we aimed at separately measuring the ability of different scales of network organization

(macro-, meso-, and micro-scales; Fig. 1) to distinguish the structure of mutualistic *versus*

antagonistic networks. We therefore considered the characterization of the structure of

ecological networks by focusing on three different network scales.

First, we characterized the macro-scale structure of the networks using 5 global metrics:

the total size (number of species in both guilds), the absolute difference in the number of

species in the two guilds, the connectance, the nestedness, and the modularity. We quantified

nestedness using the NODFc index that corrects for network size ("NODFc" function from the

R-package maxnodf, version 1.0.0 (Hoeppke & Simmons, 2021)). We quantified modularity

using the function *computeModules* from the R-package bipartite (version 2.16) (Dormann et

al., 2008) with the "Beckett" method. Preliminary analyses indicated that including the total

size and the absolute difference significantly improve the classification accuracies (not shown).

Second, we characterized the mesoscale structure of the networks by performing bipartite

motif analyses that depict the patterns of interactions between subsets of species (called motifs;

Fig. 1) (Simmons, Sweering, et al., 2019). For each network, we computed the frequencies of

the 44 possible motifs involving between 2 and 6 nodes using the function *mcount* from the

bmotif R-package (version 2.0.2) (Simmons et al., 2018) with the normalization procedure

"normalise_sum" (*i.e.* which computes the frequency of motifs among the total number of

motifs). The function *mcount* does not allow counting motifs with more than 6 nodes, and

limiting the size to 6 seemed reasonable given the size of our smaller networks (10 species):

the frequencies of larger motifs is likely to be more important in larger networks,

independently of the type of interactions.

Third, we used a multi-scale approach stemming from graph theory that integrates the

whole structural information of a network, from micro- to macro-scales: the spectral density

of the Laplacian graph (Newman, 2006a). We computed the normalized Laplacian graph of

each network (Newman, 2006a), defined as $L_{norm} = D^{-1/2} L D^{-1/2}$, where D is the degree matrix (a

diagonal matrix counting the number of interactions for each species) and L is the Laplacian

graph, defined as the difference between the adjacency matrix (a square binary matrix whose

elements represent the interactions between species of the network) and the degree matrix

(Fig. S2A). The normalized Laplacian graph is a symmetric and positive-definite matrix whose

eigenvalues are constrained between 0 and 2 (Banerjee & Jost, 2008), which makes them

comparable between networks of different sizes, and symmetrical around 1. There are as many

eigenvalues equal to 0 (or 2) as there are connected components (*i.e.* perfect modules of species

that are not connected with any species from other modules), and we can thus expect a

modular network to have more eigenvalues close to 0 (and 2) than a nested network. To be

able to efficiently compare Laplacian graphs across networks of different sizes, we converted

the eigenvalues into a spectral density using a Gaussian convolution (Lewitus & Morlon,

2016)(Fig. S2A). To do so, we adapted the *spectR* function from the RPANDA R-package

(version 1.9) (Morlon et al., 2016) for bipartite networks and used a bandwidth of 0.025 for the

Gaussian convolution. We tested the influence of the bandwidth value (from 0.001 to 0.091) on

the resulting spectral densities. We characterized each network by extracting its spectral

density values at 100 points uniformly distributed between 0 and 1.

To assess whether there are structural differences between mutualistic and antagonistic

networks beyond the different types of ecology (*e.g.,* pollination, seed-dispersal, herbivory…),

we controlled for their confounding effects by fitting linear mixed models (function *lmer* from

the R-package lme4 (version 1.1-27)) with the type of interaction (*i.e.* mutualistic or

antagonistic) as a fixed effect, the type of ecology (*e.g.,* pollination, host-parasite…) as a

random effect, and one structural metric as the response variable (Type II Wald test). In

addition, for each structural metric, we fitted linear models with only the type of ecology or

the type of interaction as fixed effects and evaluated the best-fitting model by comparing their

Akaike information criterion (AIC).

**Unsupervised classification**

220   To visualize structural differences across antagonistic *versus* mutualistic networks, we

221 used principal component analyses (PCA), with networks characterized either in terms of their

222 global metrics, motif frequencies, or Laplacian spectral densities. PCAs were performed with

223 the functions *PCA* and *fviz_pca_ind* from the FactoMineR and factoextra R-packages (versions

224 2.4 and 1.0.7 respectively) (Lê et al., 2008).

225   For each type of characterization, we then used unsupervised K-means classification

226 on the two main principal components projections. K-means were performed with the function

227 *kmeans* with 2 clusters, and we performed chi-squared tests to investigate whether the two

228 clusters tend to separate mutualistic from antagonistic networks. To quantify the ability to

229 separate antagonistic and mutualistic networks, we used a measure of classification accuracy

230 computed as the F-score with the function *F1_score* from the R-package MLmetrics. The F-

231 score is the harmonic mean of the precision (*e.g.,* the frequency of true mutualistic among

232 predicted mutualistic networks) and the recall (*e.g.,* the frequency of predicted mutualistic

233 among true mutualistic networks).

234 **Supervised classification**

235   Next, we used supervised classifiers to classify antagonistic *versus* mutualistic

236 networks based on their structure. As input for the supervised classifier, for each interaction

237 network, we used a set of structural variables, either its global metrics, its motif frequencies,

238 its Laplacian spectral density, or any combination of these sets of variables. As an output, the

239 supervised classifier provides an interaction type: "mutualistic" or "antagonistic". Supervised

240 classifiers are based on supervised learning: the classifiers have first to be trained with a

241 random subset of the interaction networks (the "training set") in order to optimize its internal

242 parameters; once trained, it can classify the other networks (the "test set") from the input

variables. We measured the accuracy of the approach using both the percentages of mutualistic

and antagonistic networks from the test set that are correctly classified by the classifier and the

F-scores of the classification of both types of networks.

We first investigated the accuracy of linear classifiers using logit and Lasso regressions.

In Lasso regression, a constraint on the estimated model coefficients can make the regression

with a high number of variables more reliable. Logit regressions were computed with the *glm*

function, whereas the Lasso regressions were performed with the function *cv.glmnet* from the

glmnet R-package (version 4.1-2) (Friedman et al., 2010). We found that training the classifiers

with 80% of the interaction networks and testing it on the remaining 20% provided a good

trade-off between the size of the test set and the accuracy of the classification.

Second, we used artificial neural networks (ANN) implemented in the *neuralnet*

function from the R-package neuralnet (version 1.44.2) (Günther & Fritsch, 2010). We

optimized their configuration (number of layers and number of neurons in each layer)

according to the input variables. We selected a 3 layers network, constituted of the input and

output layers and an intermediate layer with 5, 20, and 60 neurons for global metrics, motif

frequencies, and spectral density inputs respectively. Preliminary analyses showed that these

choices provided a good compromise between classification accuracy and overfitting (results

not shown). We also selected a non-linear logistic transformation, and the maximal number of

steps was fixed at $10^8$.

We independently trained and tested the classifiers on empirical networks using the

different sets of structural input variables. For each configuration, the ANN trainings were

replicated 50 times with different training and test sets, and we reported the average

percentages of correct classifications among antagonistic and mutualistic networks and their

respective F-scores.

**Sensitivity analyses on the supervised classification using the motif frequencies**

We selected the most accurate classification approach (supervised classification using

ANN on motif frequencies; see Results) and carried out a series of additional sensitivity

analyses: we (i) developed a measure of classification confidence, (ii) assessed the impact of

overfitting and the over-representation of mutualistic *versus* antagonistic networks and of

specific types of ecology (e.g. pollination) in the training set, (iii) tested the specificity of our

approach, *sensu* (Michalska-Smith & Allesina, 2019), (iv) investigated the effect of the types of

ecology on our classification, and finally (v) analyzed which motif complexity is required to

accurately classify networks.

First, because the training sets were limited in size (we only had a total of 343 empirical

networks), we introduced a measure of the classification confidence for each empirical

network. We trained and tested the ANN classifier on motif frequencies for 10,000

independent training and test sets of empirical networks: each empirical network was

therefore classified on average 2,000 times with different training sets. We then computed the

percentage of classifications that fell in each category, which gives an idea of classification

confidence: we considered that a given network was classified as mutualistic (resp.

antagonistic) with "high confidence" if it was classified as mutualistic (resp. antagonistic) in

more than 75% of the classifications. Conversely, a given network was classified as mutualistic

(resp. antagonistic) with "low confidence" if it was classified as mutualistic (resp. antagonistic)

in more than 50% of the classifications but in less than 75% of them.

Second, we measured overfitting by comparing the percentages of correct

classifications obtained on the training *versus* test sets as a function of the number of neurons

in the hidden layer. In addition, we assessed the impact on our results of the over-

representation of mutualistic *versus* antagonistic networks in the training set (more than 60%

of the networks were mutualistic): we trained the classifier with the same number of

antagonistic and mutualistic networks (from 5 to 125 networks for each type of interaction)

and compared the percentages of correct classifications with the ones obtained with the same-

size training sets composed of networks randomly chosen in the two interaction types. In each

condition, we replicated the classification 50 times with different training sets. Similarly, we

assessed the impact on our results of the over or under-representation of some types of ecology

in the training set (e.g. the pollination networks represent ~50% of the complete database). To

do so, we trained the classifier with the same number of networks per type of ecology (from 5

to 20 networks for each type of ecology; if a type of ecology had fewer networks than the

number per type of ecology, all networks were included in the training set) and compared the

percentages of correct classifications with the ones obtained with same-size training sets

composed of networks chosen randomly across ecology types.

Third, we tested the specificity of our approach *sensu* Michalska-Smith & Allesina

(2019) by evaluating the classification accuracy after randomizing the network structures. To

do so, we applied perturbations to the empirical networks in order to investigate whether the

classification accuracies we obtained were not driven only by differences in network size,

connectance, or degree distribution between antagonistic and mutualistic networks (Fig. 2A).

We used two different randomization strategies, all of which maintained network size

constant, and tested our classification approach on each randomized dataset. First, we

randomized the networks while keeping their connectance fixed using the *shuffle.web* function

of the **bipartite** R-package. Second, we randomized the networks while keeping their degree

distributions fixed using the r2table algorithm implemented in the *null.model* function of the

**bipartite** R-package. For each of these 2 randomization strategies, we trained the ANN using

80% of empirical networks, randomized the 20% remaining networks, and tested the ANN on

these randomized networks. We replicated this randomization strategy so that each empirical

network was randomized 200 times. In addition, we measured the effect of each

randomization strategy on the network structure by quantifying the structural similarity

between the original and randomized empirical networks. We first performed a PCA on the

motif frequencies of empirical networks, and we projected randomized ones on this PCA.

Then, we quantified the distance of each empirical network to the PCA-centroid of its

corresponding randomized networks having either fixed connectance or degree distribution;

a smaller distance indicates that the randomization strategy only slightly changes network

structure. In addition, we computed the overlap between the volumes occupied by empirical

networks and the randomized ones in the PCA using Jaccard similarity; a higher overlap

indicates that randomized networks have kept similar motif frequencies (*hypervolume_set* in

the R package **hypervolume**, v3.1.3).

Fourth, we investigated if the classifiers were strictly learning to separate antagonistic

from mutualistic networks or if they also learned the potential structural singularities of each

type of ecology from our database (*i.e.,* data-linkage). We performed two different analyses to

control for the signal from the type of ecology. First, we removed one by one each type of

ecology from the training test (*e.g.* pollination networks), we trained the classifiers on 50

independent training sets constituted by 80% of the remaining networks, and we tested the

classification on the networks from the removed type of ecology (*e.g.* are pollination networks

classified among mutualistic networks when no pollination network has been used in the

training?). If the classifiers learn to separate antagonistic from mutualistic networks regardless

of the type of ecology, the classification for the removed type of ecology should perform as

well as the original classification (*e.g.* pollination networks should be correctly classified as

mutualistic networks although no pollination network has been used in the training). If they

learn to classify ecologies rather than the type of interaction, then it should perform very badly

(*e.g.* many pollination networks should be wrongly classified as antagonistic networks as no

pollination network has been used in the training). Second, we investigated whether the

structural differences between mutualistic and antagonistic networks learned by our ANN

classifier were explained only by structural differences specific to each type of ecology (*e.g.,*

pollination *versus* host-parasite). To do so, we shuffled the labels "mutualistic" or

"antagonistic" of each type of ecology. During this shuffling process, the only constraint we

imposed was keeping 5 (*resp.* 4) different types of ecology as "mutualistic" (*resp.*

"antagonistic"), as in the original dataset. A shuffled dataset could for instance correspond to

"antagonistic" networks represented by pollination, host-parasitoid, mycorrhizal, and

anemone-fish networks, whereas "mutualistic" networks would be represented by all

bacteria-phage, herbivory, host-parasite, ant-plant, and seed dispersal networks. If the

structural differences from the type of ecology alone drive the ANN classifications, we expect

similar accuracies with these shuffled datasets compared with the original dataset. We

performed 200 of such shuffling, and for each shuffled dataset, we classified "mutualistic"

*versus* "antagonistic" networks using ANN on motif frequencies with 50 different training and

testing sets. In this analysis, we obtained cases where most networks were classified in one

type of interaction (the most represented interaction in terms of number of networks); in this

case, the percentage of correct classifications will appear very high for this interaction type,

and very low for the other. Therefore, in addition to the mean accuracy of antagonistic and

mutualistic networks, we measured the difference in accuracy. A large difference reflects a loss

of the ANN ability to classify properly.

Finally, we investigated which level of motif complexity (size) is required to

distinguish mutualistic from antagonistic networks. To do so, we measured the classification

accuracy obtained when training the ANN on two subsets of all motifs, corresponding to the

removal of the 6-nodes motifs, and then the removal of both 5 and 6-nodes motifs.

**Simulations of bipartite interaction networks**

The accuracy of supervised classifiers often depends on the size of the available training

set (i.e., data for which the classification is known), which in the case of bipartite interaction

networks is rather small (n=343 here). A possible approach to increase the training set is to use

simulated data. We explored this possibility with the recently-developed BipartiteEvol model

(Maliet et al., 2020), an eco-evolutionary model that simulates the evolution of interacting

individuals within two clades (as detailed in Supplementary Note 1).

# 3. Results

**Structure of antagonistic versus mutualistic networks**

We found differences in macro-scale structures between antagonistic and mutualistic

networks. For instance, nestedness was significantly higher in mutualistic networks (mean:

2.16 ± s.d. 0.71) than in antagonistic ones (1.94 ± 1.09; Wilcoxon-Mann-Whitney test: W = 8673,

p-value < 0.001), whereas antagonistic networks were significantly more connected (0.23 ±

0.17) than mutualistic ones (0.15 ± 0.09; Wilcoxon-Mann-Whitney test: W = 17709, p-value <

0.001). The difference in terms of modularity was not significant (Wilcoxon-Mann-Whitney

test: W = 12559, p-value = 0.19) (Fig. 2A). In addition, we found that mutualistic networks were

composed of more species compared to antagonistic ones (76.1± 92.3 for antagonistic networks

and 82.7 ± 72.7 for mutualistic ones; Wilcoxon-Mann-Whitney test: W = 11124, p-value = 0.003).

These differences were however better explained by specific ecologies rather than the overall

interaction type. Specifically, when controlling for the different types of ecology (*e.g.,*

pollination, herbivory, seed dispersal...), antagonistic and mutualistic networks were not

significantly different in terms of their nestedness (Wald test: $\chi2 = 0.060$, p-value = 0.81),

connectance (Wald test: $\chi2 = 0.41$, p-value = 0.52), modularity (Wald test: $\chi2 = 0.42$, p-value =

0.52), the total number of species (Wald test: $\chi2 = 0.29$, p-value = 0.59), and absolute difference

in the number of species per guild (Wald test: $\chi2 = 0.95$, p-value = 0.33). Model comparisons

confirmed that differences in terms of global metrics were better explained by the different

types of ecology than by the antagonistic/mutualistic type of interactions (Table S1). For

instance, compared to other types of ecology, bacteria-phage networks were more connected

but generally less nested, while herbivory networks were more nested (Fig. S1, Table S2 A-C).

Next, using the spectral density of the Laplacian graph (Fig. S2A) on perfectly modular

or nested networks revealed that modular networks have rather smooth densities with high

peaks whereas nested networks have noisier densities with several small peaks (Fig. S2B-E).

We found that mutualistic networks were enriched in eigenvalues near 1, while antagonistic

networks had significantly more spectral eigenvalues near 0 (Fig. 2B), which was mainly

driven by herbivory but not host-parasite networks (Fig. S3).

Finally, using motif frequencies, we found, following the motif denomination of

(Simmons, Sweering, et al., 2019), that antagonistic networks were significantly enriched in

densely connected motifs (such as motif 12 and motifs 31 to 37), whereas mutualistic networks

were enriched in motifs with asymmetrical specialization, wherein specialist species tend to

interact with the most generalist species in the motif (such as motifs 10, 19, 20, 25, 28 Fig. 2C).

In total, 24 out of the 44 motifs differed significantly in frequency between antagonistic and

mutualistic networks (p<0.01). When controlling for the type of ecology (*e.g.,* pollination *versus*

herbivory), 4 motifs were significantly different in frequency between antagonistic and

mutualistic networks (motifs 17, 25, 28, and 44). 31 out of 44 motifs differed significantly in

frequency across the different types of ecology (Fig. S4C): for example, seed-dispersal and

host-parasite networks had relatively high frequencies of 4- to 6-nodes motifs, in particular

motifs 29 to 43, which may be linked to their high connectance (Figs. S1, S4., Table S2A).

**Unsupervised classifications of empirical networks**

A principal component analysis (PCA) on the global metrics did not identify distinct clusters

for mutualistic and antagonistic networks (Fig. 3A). Consequently, an unsupervised K-means

classification on the principal component axes was not sufficient to confidently classify the

networks (F-scores of 0.63 and 0.55 for mutualistic and antagonistic networks respectively)

(Fig. 3A). Similarly, networks with different types of ecology (*e.g.* seed dispersal, herbivory,

ant-plant…) were largely mixed on the PCA projection (Fig. S5) and did not systematically fell

in one of the two clusters of the K-means classification (Table S3A).

PCAs performed on spectral density values (Fig. 3B) or motif frequencies (Fig. 3C) were

better at separating antagonistic *versus* mutualistic empirical networks than when using the

global metrics, but only slightly so (Fig. 3A). Antagonistic networks appeared to be spread on

the projection, while mutualistic networks were more clustered, which is potentially linked to

the high number of pollination networks in the database (Figs. S6-7). Unsupervised K-means

classifications on the principal component axes were not sufficient to confidently classify

mutualistic and antagonistic networks (Fig. 3B-C). Networks with different types of ecology

were largely mixed between the two clusters from the K-means classification (Tables S3 B-C),

suggesting that unsupervised classifications do not perform well at classifying ecological

networks, regardless of the metric used to characterize their structure.

## Supervised classifications of empirical networks

We found that the two linear classifiers (Logit and Lasso regressions) mainly classified

empirical networks as mutualistic, regardless of their interaction type (Table S4). Artificial

neural networks (ANN) performed much better, in particular when using motif frequencies.

Motif frequencies led to better classifications than the spectral density or the global metrics,

and adding other variables to motif frequencies did not improve the classifications (Table S4).

The ANN based on motif frequencies classified an average of 65% of the antagonistic networks

(resp. 84% of the mutualistic networks) classified as antagonistic (resp. mutualistic), resulting

in an F-score of 0.68 and 0.82 for antagonistic and mutualistic networks respectively (Table S4).

Irrespectively of the bandwidth used to smooth the Laplacian spectrum, using the ANN with

the spectral density led to lower accuracies (Tables S4-5). We therefore used ANN on motif

frequencies for the remaining analyses.

**Supervised classification of the empirical networks using motif frequencies**

Using our measure of classification accuracy with ANN on motif frequencies, we found

that 69% (52% and 17% with high and low confidence respectively; F-score=0.74) of the

antagonistic networks and 89% (76% and 13% with high and low confidence respectively; F-

score=0.86) of the mutualistic networks were correctly classified (Table 1). In total, 81% (67%

and 14% with high and low confidence respectively) of the networks were correctly classified

(Table 1). The percentage of correct classifications varied according to the type of ecology.

Indeed, most bacteria-phage, seed dispersal, and pollination networks were correctly

classified by this approach (*e.g.* 23/29 of the seed dispersal networks were correctly classified,

among which ~85% with high confidence). Conversely, high intimacy mutualistic networks

(*i.e.* ant-plant and mycorrhizal networks) presented frequent misclassification (only 7 out of 12

networks (~60%) were classified as mutualistic), herbivory networks (which are low intimacy

antagonistic networks) were mainly classified as mutualistic (13/23, *i.e.* 57%), and 50% (5/10)

of the host-parasitoid networks were also misclassified (Table 1, Fig. S18).

**Robustness of the classification using motif frequencies**

Overfitting had a limiting impact on the previous results (Fig. S8). In addition,

controlling for the overrepresentation of mutualistic networks in the database resulted in

similar mean percentage of correct classifications (Fig. S9A). Similarly, equalizing the number

of networks per type of ecology in the training set did not significantly affect the overall

percentages of correct classifications (Fig. S9B). Finally, we found lower classification

accuracies when randomizing the empirical networks compared to the original classifications

(Fig. S16a,b). Classifications were arbitrary when randomizing networks with fixed network

size and connectance (mean F-scores = 0.39 and 0.53 for antagonistic and mutualistic networks

respectively). When randomizing networks with fixed network size and degree distributions,

classifications were not completely arbitrary (mean F-scores = 0.39 and 0.73 for antagonistic

and mutualistic networks respectively (Fig. S16a,b). Such a deviation from strict random

classification (50/50) can be explained as follows: mutualistic networks have a lower

connectance (Fig. 2A) and are therefore more constrained during the network randomization

than antagonistic networks, which results in motif frequencies closer to the original ones (Fig.

S16c). Indeed, when projecting all empirical and randomized networks on a PCA, empirical

mutualistic networks are significantly closer to their randomized counterparts (mean

distance=8.9) compared to antagonistic networks (mean distance =14.2; Fig. S16d), and the

overlap between the volumes occupied by empirical and randomized networks is higher for

mutualistic networks than for antagonistic ones (Fig. S16e). In any case, the F-scores

significantly dropped by 0.35 and 0.13 for antagonistic and mutualistic networks respectively

when randomizing the networks, which suggests that antagonistic and mutualistic networks

could not be told apart with high accuracy once randomized. Thus, differences in network

size, connectance, or degree distributions between mutualistic and antagonistic networks may

partially contribute to our ability to tell mutualistic and antagonistic ecological networks apart

from their structure but fail to explain the high classification accuracy we report. Besides, we

found that removing a given type of ecology during the ANN training decreased the

percentage of correct classifications for this removed type of ecology but remained quite good

in some cases (Table S7). For instance, 65% of the seed dispersal and 63% of the ant-plant

networks were correctly classified when none of these networks were used during training,

compared to 79% and 66% respectively when they were included (Table S7; Table 1). Also,

artificially shuffling the mutualistic or antagonistic nature of each type of ecology decreased

the mean percentages of correct classifications by about 10%, and the F-scores by about 0.08,

while increasing the difference in percentages of correct classifications and F-scores between

antagonistic and mutualistic networks by 30% and 0.3 respectively (Fig. S17). This major loss

of the classification ability suggests that although the ANN classifiers partially learned the

singularities of the networks linked to ecology (*i.e.* data linkage), they also learned the

antagonistic/mutualistic dichotomy beyond ecology.

Finally, we found that the classification accuracy increased when including more

complex motifs (Table S9). The difference was most notable between analyses carried with

motifs of size at most 4 and of size at most 5 or 6 (Table S9), and rather insignificant between

analyses carried with motifs of size at most 5 and at most 6. This suggests that complex motifs

(with at least 5 nodes) are needed to distinguish between mutualistic and antagonistic

networks, but that adding even larger motifs would not significantly increase classification

accuracy.

**Ability of simulated networks to improve the accuracy of classifiers**

Using simulated networks did not improved our ability to distinguish mutualistic from

antagonistic networks both when using K-means classifications (results not shown) and ANN

(Table S8B). Mutualistic networks, in particular, were often wrongly classified as antagonistic.

When classifying simulated BipartiteEvol networks from the test sets, 70% (F-score =0.71) of

antagonistic networks and 58% (F-score = 0.57) of mutualistic ones were accurately classified

when using motif frequencies and ANN (Table S8A), suggesting difficulties in separating

antagonistic *versus* mutualistic networks simulated with BipartiteEvol using ANN. When

comparing the structure of simulated and empirical networks, we found differences in global

metrics, motif composition, and density of eigen-values that may explain why simulations

failed to improve the accuracy of classification (Supplementary Note 2).

# 4. Discussion

In this study, we showed that discriminating antagonistic *versus* mutualistic ecological networks based on their structure is overall possible with supervised machine learning. Artificial neural networks trained on motif frequencies reached an average of 81% correct classifications (F-scores of 0.74 and 0.86 for antagonistic and mutualistic networks respectively). Structural differences between antagonistic and mutualistic networks can be captured at the network mesoscale through differences in some motif frequencies, whereas the macro-scale modular/nested dichotomy does not seem sufficient. Beyond the classical dichotomy between mutualism/antagonism, we emphasize that the different types of ecology have an important structural imprint on the structure of ecological networks.

Using motif frequencies and ANN, we were able to correctly classify around 81% of the empirical interaction networks. This percentage of successful classifications is higher than the success rate achieved by the only previous attempt we are aware of to classify networks with ANN based exclusively on their structure (Michalska-Smith & Allesina, 2019), and achieved without accounting for environmental conditions (as in Song & Saavedra, 2020). We also demonstrated the specificity of our approach (*sensu* Michalska-Smith & Allesina, 2019): when randomizing sufficiently the network structures, we obtained lower classification accuracies, suggesting that motif frequencies, which depict patterns of species interactions at the meso-scale, capture structural differences between mutualistic and antagonistic networks that are not only due to differences in network size, connectance, or degree distributions. This therefore indicates that there are significant structural differences at the network meso-scale, which cannot be fully captured when looking at macro-scale alone using global metrics or that

can be blurred when using integrated measures of network structure such as the Laplacian

spectral densities.

Our analyses confirm that global metrics, including the modular/nested dichotomy, are

not sufficient to discriminate between mutualistic and antagonistic network structures

(Fontoura et al., 2020; Michalska-Smith & Allesina, 2019). We did not find a significant

difference in modularity values between mutualistic and antagonistic networks. We did find

that empirical mutualistic networks tend to be more nested and less connected than

antagonistic networks, as previously suggested by (Bascompte et al., 2003; Thébault &

Fontaine, 2010), but this dichotomy is not consistent enough (*i.e.,* there are too many exceptions

and their variances are too large) to provide a robust classification criterion. In addition, it is

worth mentioning that these global metrics are largely sensitive to the sampling effort (Nielsen

& Bascompte, 2007) and may be less reliable in poorly sampled networks. In addition, there

are several empirical examples of exceptions to the nested/modular dichotomy in the

literature, such as a nested structure in host-parasite antagonistic networks (Pilosof et al.,

2014). Within our database, some pollination and mycorrhizal networks are known to be

modular (Chagnon et al., 2015; Jacquemyn et al., 2015; Martos et al., 2012; Olesen et al., 2007;

Perez-Lamarque et al., 2022; Xing et al., 2019), while they assemble mainly mutualistic

interactions.

Why motifs performed better than the spectral density of the Laplacian graph is

unclear, as this latter representation is supposed to integrate all scales. Using a constant

bandwidth value to convert the Laplacian graph spectrum into a density may result in

information loss when comparing networks of different sizes. Also, transforming the discrete

Laplacian spectrum into a continuous density may be problematic for small networks with

only a few eigenvalues.

Motifs focus on precise structural patterns of interspecific interactions that can be

missed when summarizing all the information in a single global value at the network macro-

scale (Simmons, Cirtwill, et al., 2019). Although motif frequencies are highly variable across

mutualistic and antagonistic networks making it impossible to accurately classify using

unsupervised classification techniques (*e.g.,* PCA), there seems to exist slight differences in

motif frequencies that are distinguishable using supervised machine learning. In particular,

densely connected motifs (*e.g.* motifs 12 and 31 to 37) are significantly more frequent in

antagonistic networks, while mutualistic networks are characterized by motifs with

asymmetrical specialization (*e.g.,* motifs 10, 19, 20, 25, 28). In mutualistic networks, the

emergence of motifs with asymmetrical specialization, which often parallels the emergence of

nestedness, has been related to three different types of explanations. First, a pattern of

asymmetric specialization may be partly explained from trait complementarity and

convergence.  While generalist species can establish interactions with broad range of

mutualistic partners, trait complementarity generates specialization that constrains the

possible interactions of specialist species  (Rezende, Jordano, et al., 2007; Thompson, 2005).

Second, such a pattern can emerge from differences in species abundances (neutrality

hypothesis): if interactions between individuals occur at random, rare specialist species will

be more likely to interact with abundant generalist species than with another rare specialist

(Vázquez et al., 2009). It has been proposed that a combination of trait complementarity and

abundance mechanisms best explains the observed structure of mutualistic networks

(Santamaría & Rodríguez-Gironés, 2007; Vázquez et al., 2009). Finally, it has also been

suggested that asymmetric specialization reduces interspecific competition (Bastolla et al.,

2009; Thompson, 2005), therefore enhancing the number of coexisting species (Bascompte et

al., 2006; Bastolla et al., 2009) and widening the conditions over which mutualists coexist (Rohr

et al., 2014), which in the end increases the persistence and resilience of mutualistic networks

to perturbations (Okuyama & Holland, 2008; Thébault & Fontaine, 2010). On the contrary, we

expect antagonistic partners to engage in eco-evolutionary arm races, which may result in

modules formed by highly specialized within-module interactions (i.e. densely connected

motifs, which tend to be more frequent in antagonistic networks) (Maliet et al., 2020;

Thompson, 2005). Such modular structures would increase the resilience of antagonistic

communities (Thébault & Fontaine, 2010). Differences in motif frequencies can also be

influenced by sampling biases, including missing links, preferential detection of some species,

and spatial and temporal limits of the sampling. Future works will be required to identify the

ecological processes and/or sampling effects that generate such structural differences between

mutualistic and antagonistic networks (Brimacombe et al., 2023).

Mutualistic and antagonistic networks cover a wide range of ecologies (*e.g.* pollination,

parasitism, herbivory) and our analyses showed that the type of ecology matters in the

supervised classification, a phenomenon referred to as data linkage: networks with different

ecologies tend to have their own structural signatures beyond the mutualistic/antagonistic

dichotomy, and the ANN classifier partially learns to identify the structural uniqueness of

each type of ecology. In particular, we showed that a better classification of networks of a

particular type of ecology is achieved by using networks from the same ecology during the

training of the classifiers. This is consistent with the result from linear mixed models that, after

controlling for the type of ecology, the only significant difference between mutualistic and

antagonistic networks is the frequency of a few motifs (motifs 17, 25, 28 and 44). These results

on the importance of the type of ecology corroborate previous studies that found, for instance,

significant structural differences between pollination and seed dispersal networks (Mello et

al., 2011), between mutualistic plant-fungus and plant-animal networks (Toju et al., 2015), or

across mycorrhizal networks formed by different fungal lineages (Perez-Lamarque et al.,

2022). Further works are needed to better identify the species traits and/or ecological factors

responsible for the structural uniqueness of the different types of network ecology. This impact

of the type of ecology may also explain the low classification accuracy of some networks, like

the herbivory networks, which share similar structural properties with the pollination

networks in both spectral density and motifs frequency. Nevertheless, we also demonstrated

by shuffling the dataset that the high classification accuracy we obtained is achieved by

learning structural differences between antagonistic and mutualistic networks beyond the

structural patterns specific to each type of ecology. Beyond the structural singularities of the

type of ecology, there are structural differences between mutualistic and antagonistic

networks that can be captured using our meso-scale approach and ANNs.

The accuracy of supervised classification approaches strongly depends on the size and

quality of the dataset used for training. We used one of the largest publicly available databases

of empirical networks, but it contained only 343 networks in total. In addition, it had a strong

unbalanced representation of the different types of ecology, with 80% of the mutualistic

networks represented by pollination networks and 60% of the antagonistic networks

represented by host-parasite networks. There was also a strong imbalance in the

representation of interaction intimacy, which is known to impact network structure (Fontaine

et al., 2011; Guimarães et al., 2007; Hembry et al., 2018; Pires & Guimarães, 2013): mutualistic

networks were mainly low intimacy networks, except ant-plants (n=3) or mycorrhizal

networks (n=9), and antagonistic networks were mainly high intimacy networks, except

herbivory networks (n=23). As a consequence, only ~60% of the high-intimacy mutualistic

networks and ~40% of the low-intimacy antagonistic networks were correctly classified. These

results confirm that interaction intimacy may confer a particular structure to ecological

networks (Fontaine et al., 2011; Guimarães et al., 2007). Increasing the number of empirical

networks, especially mutualistic networks with high intimacy (*e.g.* symbiotic networks) and

antagonistic networks with low intimacy (*e.g.* trophic networks) would improve further our

ability to classify ecological networks by better representing the large structural heterogeneity

of ecological networks (Michalska-Smith & Allesina, 2019). In addition, using information on

interaction strengths may foster our classification accuracy, as it contains relevant structural

information (*e.g.*, on nestedness, Staniczenko et al., 2013). We could not test this properly as

only 69 out of 343 networks in our database contained interaction strengths but the ongoing

sampling of more and more weighted networks paves the way for the inclusion of interaction

strengths into classification approaches in the future.

The accuracy of supervised classification approaches can also strongly depend on the

type, architecture, and tuning of the classifier. For example, the overfitting we noticed in the

ANN (which did not seem to negatively impact our results) could potentially be avoided by

using other regularization technics, such as dropout (Srivastava et al., 2014). We chose to use

simple, few-layers neural networks implemented in R for their availability and ease of usage

to the large community of R users, as well as because our dataset for training was rather small.

However, several other machine learning tools exist and could be tested, although they may

require larger training sets; graph neural networks in particular have been specifically

designed to learn information from graphs such as ecological networks (Wu et al., 2021).

658           One approach to better train the ANNs would be to increase the size of the training

data thanks to synthetic data, obtained through either simulations or data augmentation. We

investigated the possibility to train from simulated data with BipartiteEvol, an individual-

based model that mimics the eco-evolutionary emergence of bipartite interaction networks

(Maliet et al., 2020). However, ANN classifiers trained on BipartiteEvol simulations were not

able to discriminate mutualistic and antagonistic empirical networks, suggesting that the

model somehow fails at capturing important structural differences between antagonistic and

mutualistic networks. (Maliet et al., 2020) showed that the model produces realistic networks,

which we confirm here, and is able to reproduce major global differences observed between

mutualistic and antagonistic networks. However, we find that BipartiteEvol simulations tend

to have higher frequencies of complex motifs and a lower connectance compared to empirical

networks. Consequently, their projection only covers a small proportion of the space occupied

by empirical networks. We cannot exclude that using different parameters in our simulations

would provide more heterogeneous simulated networks that better match the high variability

observed in empirical networks. Alternatively, one possible explanation is that many bipartite

interaction networks belong to a continuum between antagonism and mutualism rather than

a strict dichotomy. Indeed, such a continuum between mutualism and antagonism has been

observed in many interactions found in natural communities (Montesinos-Navarro et al.,

2017). For instance, it has been shown that parrot-plant networks can combine the two types

of interactions (Montesinos-Navarro et al., 2017). Similarly, pollination or mycorrhizal

networks often contain cheating species that use antagonistic strategies and induce structural

changes within the interaction networks (Genini et al., 2010; Klironomos, 2003; Perez-

Lamarque et al., 2020). Interactions can also shift from mutualism to antagonism on short

timescales, depending on some external conditions (Canestrari et al., 2014; Sachs et al., 2011).

By contrast, BipartiteEvol models the emergence of a network in strict antagonistic or

mutualistic communities. This calls for future efforts to improve the BipartiteEvol model or to

use simulations under other generative models (*e.g.,* (Cai et al., 2020; Thebault & Fontaine,

2010). Alternatively, data augmentation methods could also be tested to generate synthetic

networks by randomly subsampling, mixing and rewiring empirical networks; but whether

such approaches produce realistic interaction networks would need to be tested first.

Ultimately, supervised classifying approaches could be particularly useful to predict

the type of undetermined interactions. For instance, many plant-fungus interaction networks

are assembled from high-throughput sequencing data, but we currently ignore the mutualistic

*versus* the antagonistic type of many of these endophytic interactions (Chagnon et al., 2016;

Selosse et al., 2018). Similarly, an increasing number of networks are built from co-occurrence

data generated with high-throughput sequencings, such as the ones between microscopic

planktonic species sampled by the *Tara Oceans* expeditions (Chaffron et al., 2021; Lima-

Mendez et al., 2015). The ecology of the majority of species in these networks is currently

undetermined (Bjorbækmo et al., 2020; Lima-Mendez et al., 2015; Strydom et al., 2021).

However, a preliminary analysis of bipartite networks built from co-occurrence data in the

*Tara Oceans* data revealed that these networks were structurally very different from the

empirical and simulated networks we used here (results not shown). Classifying these

networks would thus require using a specific training dataset composed of microbial networks

with known interaction types (as discussed in Supplementary Note 3). Similarly, while our

approach is strictly limited to bipartite networks, the increasing availability of multipartite

networks, embracing different ecologies (Domínguez-García & Kéfi, 2024; Pocock et al., 2012),

and the recent development of generative models for multipartite networks (Hale et al., 2023)

suggest that a classification approach to more complex networks may now be possible.

Embracing the challenge of telling apart interaction types in multipartite networks from their

structure would also allow to evaluate differences in structure between simulated and

empirical multipartite networks.

**Conclusion**

Mutualistic and antagonistic networks have structural differences that can be

particularly well captured by motif frequencies with supervised machine learning.

Antagonistic networks are enriched in densely connected motifs, while mutualistic networks

are enriched in motifs with asymmetrical specialization. Besides this dichotomy, there are

structural differences across networks with distinct ecologies. These structural differences can

be used to classify bipartite interaction networks with supervised machine learning with

pretty good accuracy. These performances could be further improved by increasing the

number of empirical networks used for training, diversifying the type of ecology sampled, and

improving currently available models to simulate networks with realistic structures. While

more work is needed to predict interaction types for co-occurrence networks, our analyses

provide promising results on the ability to predict interaction types from network structure.

**Table 1. The type of interaction of the empirical ecological networks can be predicted based on their**

**motif frequencies using artificial neural network classifiers.**

The following results were obtained by training the artificial neural networks (ANN) with 10,000

different training and test sets, using motif frequencies as the input structural variables. From those

10,000 classifications, we calculated the percentage of antagonistic *versus* mutualistic classifications for

each network: a network was predicted as a mutualistic (resp. antagonistic) network with high

confidence (highlighted in grey) if it was classified as a mutualistic (resp. antagonistic) network in more

than 75% of the classifications. It was predicted as a mutualistic (resp. antagonistic) network with low

confidence if it was classified as a mutualistic (resp. antagonistic) network in more than 50% but less

than 75% of the classifications. We therefore obtained a predicted type of interaction for each of the 343

empirical networks. The distributions of these classification confidence for all networks are given in Fig.

S18.

The second column indicates the number of interaction networks in our database gathered for each

ecology of interaction. Then, we indicate the number (and the percentage) of networks classified as

either mutualistic or antagonistic for each ecology of interaction and at the bottom, we show the global

contingency table for all mutualistic and antagonistic networks. We also indicate the F-score associated

with the global contingency table for each antagonistic or mutualistic type of interaction. Rows are

ordered based on the % of networks classified as mutualistic within each type of ecology. In the end, 88

antagonistic and 191 mutualistic networks are correctly classified, i.e., 279 of the 343 empirical networks

(81%).

| Network ecology | Number of networks | Prediction | | | |
|---|---|---|---|---|---|
| | | Antagonistic network (high confidence) | Antagonistic network (low confidence) | Mutualistic network (high confidence) | Mutualistic network (low confidence) |
| Anemone-Fish (mutualistic) | 1 | 0 (0%) | 0 (0%) | 1 (100%) | 0 (0%) |
| Pollination (mutualistic) | 174 | 7 (4%) | 7 (4%) | 139 (80%) | 21 (12%) |
| Seed dispersal (mutualistic) | 29 | 6 (21%) | 0 (0%) | 20 (69%) | 3 (10%) |
| Ant-Plant (mutualistic) | 3 | 0 (0%) | 1 (33%) | 1 (33%) | 1 (33%) |
| Herbivory (antagonistic) | 23 | 4 (17.5%) | 6 (26%) | 9 (39%) | 4 (17.5%) |
| Mycorrhiza (mutualistic) | 9 | 3 (33.5%) | 1 (11%) | 3 (33.5%) | 2 (22%) |
| Host-Parasitoid (antagonistic) | 10 | 4 (40%) | 1 (10%) | 4 (40%) | 1 (10%) |
| Bacteria-Phage (antagonistic) | 17 | 12 (70%) | 1 (6%) | 3 (18%) | 1 (6%) |
| Host-Parasitism (antagonistic) | 77 | 46 (60%) | 14 (18%) | 10 (13%) | 7 (9%) |
| **Antagonistic networks** | 127 | **66** **(52%)** | **22** **(17%)** | **26** **(21%)** | **13** **(10%)** |
| **Mutualistic networks** | 216 | **16** **(7%)** | **9** **(4%)** | **164** **(76%)** | **27** **(13%)** |
| **Correct classifications (%)** **F-score** | | **88 (69%)** **F-score = 0.74** | | **191 (88%)** **F-score = 0.86** | |

**Figures:**

**Figure 1: Multi-scale structure of bipartite networks**

The structure of bipartite ecological networks, represented here by an ant-plant network

(Blüthgen et al., 2004) can be studied at the macro-, meso-, and micro-scale. The Laplacian

spectral density corresponds to a multi-scale approach from the macro to the micro-scale,

whereas global metrics and motif frequencies focus on macro- and meso-scales, respectively.

Blue nodes represent plant species, brown nodes ant species, and grey edges interspecific

interactions.

**Figure 2. Differences between antagonistic and mutualistic networks across scales of network**

**structure.**

(A) Connectance, nestedness (NODFc), and modularity (Newman's index) of antagonistic and

mutualistic empirical networks. Stars indicate the statistical significance of the difference between

antagonistic and mutualistic networks, as measured by a Wilcoxon-Mann-Whitney test (*** = p-value <

0.01, ns = non-significant).

(B) Densities of eigenvalues from the Laplacian spectral density that differ significantly between

antagonistic and mutualistic networks (Wilcoxon-Mann-Whitney test: p-value<0.01). The

discontinuities in the x-axis are due to the filtering of eigenvalues densities that differ significantly

between antagonistic and mutualistic networks.

(C) Frequencies of the motifs that differ significantly between antagonistic and mutualistic networks

(Wilcoxon-Mann-Whitney test: p-value<0.01). Each motif is illustrated at the top of each boxplot

(following the denomination of Simmons *et al.* 2018).

Mutualistic networks are shown in green and antagonistic ones in yellow. Boxplots present the median

surrounded by the first and third quartiles, and whiskers extend to the extreme values but no further

than 1.5 of the interquartile range.

**Figure 3. Mesoscale network structure as a scale to discriminate mutualistic *from* antagonistic**

**networks.**

Projection of the 343 empirical networks on the two principal components (PC1 and PC2) obtained using

principal coordinate analysis (PCA) on (A) global metrics (*i.e.* nestedness, modularity, connectance, and

network size), (B) spectral density of the Laplacian graph and (C) motif frequencies. The percentage of

explained variance is indicated on each axis.

Each associated table corresponds to the contingency table of unsupervised K-means classification

based on the global metrics (A), spectral density of the Laplacian graph (B), and motifs frequencies (C).

Chi-squared test associated with the K-means classification is indicated.

| Unsupervised classification | Mutualistic | Antagonistic |
|---|---|---|
| Cluster 1 | 97 | 85 |
| Cluster 2 | 119 | 42 |
| F-scores | **0.63** | **0.55** |

$\chi^2 = 15.572$, p-value<0.001

| Unsupervised classification | Mutualistic | Antagonistic |
|---|---|---|
| Cluster 1 | 37 | 52 |
| Cluster 2 | 179 | 75 |
| F-scores | **0.76** | **0.48** |

$\chi^2 = 23.607$, p-value < 0.001

| Unsupervised classification | Mutualistic | Antagonistic |
|---|---|---|
| Cluster 1 | 13 | 49 |
| Cluster 2 | 203 | 78 |
| F-scores | **0.81** | **0.51** |

$\chi^2 = 55.093$, df = 1, p-value <0.001

**A** Global metrics

**B** Laplacian graph

**C** Motif frequencies

● Mutualistic networks  ● Antagonistic networks

# References

Banerjee, A., & Jost, J. (2008). On the spectrum of the normalized graph Laplacian. *Linear Algebra and*

*Its Applications*, *428*(11–12), 3015–3022. https://doi.org/10.1016/j.laa.2008.01.029

Bascompte, J., & Jordano, P. (2007). Plant-animal mutualistic networks: The architecture of

biodiversity. *Annual Review of Ecology, Evolution, and Systematics*, *38*(1), 567–593.

https://doi.org/10.1146/annurev.ecolsys.38.091206.095818

Bascompte, J., Jordano, P., Melian, C. J., & Olesen, J. M. (2003). The nested assembly of plant-animal

mutualistic networks. *Proceedings of the National Academy of Sciences*, *100*(16), 9383–9387.

https://doi.org/10.1073/pnas.1633576100

Bascompte, J., Jordano, P., & Olesen, J. M. (2006). Asymmetric coevolutionary networks facilitate

biodiversity maintenance. *Science*, *312*(5772), 431–433.

https://doi.org/10.1126/science.1123412

Bastolla, U., Fortuna, M. A., Pascual-García, A., Ferrera, A., Luque, B., & Bascompte, J. (2009). The

architecture of mutualistic networks minimizes competition and increases biodiversity.

*Nature*, *458*(7241), 1018–1020. https://doi.org/10.1038/nature07950

Bjorbækmo, M. F. M., Evenstad, A., Røsæg, L. L., Krabberød, A. K., & Logares, R. (2020). The planktonic

protist interactome: Where do we stand after a century of research? *The ISME Journal*, *14*(2),

544–559. https://doi.org/10.1038/s41396-019-0542-5

Blüthgen, N., Menzel, F., Hovestadt, T., Fiala, B., & Blüthgen, N. (2007). Specialization, Constraints, and

Conflicting Interests in Mutualistic Networks. *Current Biology*, *17*(4), 341–346.

https://doi.org/10.1016/j.cub.2006.12.039

Brimacombe, C., Bodner, K., Michalska-Smith, M., Poisot, T., & Fortin, M.-J. (2023). Shortcomings of

reusing species interaction networks created by different sets of researchers. *PLOS Biology*,

*21*(4), e3002068. https://doi.org/10.1371/journal.pbio.3002068

Bronstein, J. L. (Ed.). (2015). *Mutualism*. Oxford University Press. https://doi.org/10.1093/acprof:oso/9780199675654.001.0001

Cai, W., Snyder, J., Hastings, A., & D'Souza, R. M. (2020). Mutualistic networks emerging from adaptive niche-based interactions. *Nature Communications*, *11*(1), 5470. https://doi.org/10.1038/s41467-020-19154-5

Canestrari, D., Bolopo, D., Turlings, T. C. J., Roder, G., Marcos, J. M., & Baglione, V. (2014). From parasitism to mutualism: Unexpected interactions etween a cuckoo and Its host. *Science*, *343*(6177), 1350–1352. https://doi.org/10.1126/science.1249008

Chaffron, S., Delage, E., Budinich, M., Vintache, D., Henry, N., Nef, C., Ardyna, M., Zayed, A. A., Junger, P. C., Galand, P. E., Lovejoy, C., Murray, A. E., Sarmento, H., null, null, Acinas, S. G., Babin, M., Iudicone, D., Jaillon, O., Karsenti, E., … Eveillard, D. (2021). Environmental vulnerability of the global ocean epipelagic plankton community interactome. *Science Advances*, *7*(35), eabg1921. https://doi.org/10.1126/sciadv.abg1921

Chagnon, P.-L. (2016). Seeing networks for what they are in mycorrhizal ecology. *Fungal Ecology*, *24*, 148–154. https://doi.org/10.1016/j.funeco.2016.05.004

Chagnon, P.-L., Bradley, R. L., & Klironomos, J. N. (2015). Trait-based partner selection drives mycorrhizal network assembly. *Oikos*, *124*(12), 1609–1616. https://doi.org/10.1111/oik.01987

Chagnon, P.-L., U'Ren, J. M., Miadlikowska, J., Lutzoni, F., & Arnold, A. E. (2016). Interaction type influences ecological network structure more than local abiotic conditions: Evidence from endophytic and endolichenic fungi at a continental scale. *Oecologia*, *180*(1), 181–191. https://doi.org/10.1007/s00442-015-3457-5

Dáttilo, W., Izzo, T. J., Vasconcelos, H. L., & Rico-Gray, V. (2013). Strength of the modular pattern in Amazonian symbiotic ant–plant networks. *Arthropod-Plant Interactions*, *7*(4), 455–461. https://doi.org/10.1007/s11829-013-9256-1

de Lange, S. C., de Reus, M. A., & van den Heuvel, M. P. (2014). The Laplacian spectrum of neural

networks. *Frontiers in Computational Neuroscience*, *7*. https://doi.org/10.3389/fncom.2013.00189

https://doi.org/10.3389/fncom.2013.00189

Domínguez-García, V., & Kéfi, S. (2024). The structure and robustness of ecological networks with two

interaction types. *PLOS Computational Biology*, *20*(1), e1011770.

https://doi.org/10.1371/journal.pcbi.1011770

Dormann, C. F., Gruber, B., & Fründ, J. (2008). *Introducing the bipartite Package: Analysing Ecological*

*Networks*. *8*, 5.

Estrada, E. (2007). Characterization of topological keystone species. *Ecological Complexity*, *4*(1–2), 48–

57. https://doi.org/10.1016/j.ecocom.2007.02.018

Fontaine, C., Guimarães, P. R., Kéfi, S., Loeuille, N., Memmott, J., van der Putten, W. H., van Veen, F. J.

F., & Thébault, E. (2011). The ecological and evolutionary implications of merging different

types of networks: Merging networks with different interaction types. *Ecology Letters*, *14*(11),

1170–1181. https://doi.org/10.1111/j.1461-0248.2011.01688.x

Fontoura, L., Cantor, M., Longo, G. O., Bender, M. G., Bonaldo, R. M., & Floeter, S. R. (2020). The

macroecology of reef fish agonistic behaviour. *Ecography*, ecog.05079.

https://doi.org/10.1111/ecog.05079

Fortuna, M. A., Ortega, R., & Bascompte, J. (2014). The Web of Life. *arXiv:1403.2575 [q-Bio]*.

http://arxiv.org/abs/1403.2575

Fortuna, M. A., Stouffer, D. B., Olesen, J. M., Jordano, P., Mouillot, D., Krasnov, B. R., Poulin, R., &

Bascompte, J. (2010). Nestedness versus modularity in ecological networks: Two sides of the

same coin? *Journal of Animal Ecology*. https://doi.org/10.1111/j.1365-2656.2010.01688.x

Friedman, J., Hastie, T., & Tibshirani, R. (2010). Regularization paths for generalized linear models via

coordinate descent. *Journal of Statistical Software*, *33*(1).

https://doi.org/10.18637/jss.v033.i01

Genini, J., Morellato, L. P. C., Guimarães, P. R., & Olesen, J. M. (2010). Cheaters in mutualism networks.

*Biology Letters*, *6*(4), 494–497. https://doi.org/10.1098/rsbl.2009.1021

Guimarães, P. R., Rico-Gray, V., Oliveira, P. S., Izzo, T. J., dos Reis, S. F., & Thompson, J. N. (2007).

Interaction intimacy affects structure and coevolutionary dynamics in mutualistic networks.

*Current Biology*, *17*(20), 1797–1803. https://doi.org/10.1016/j.cub.2007.09.059

Günther, F., & Fritsch, S. (2010). neuralnet: Training of Neural Networks. *The R Journal*, *2*(1), 30.

https://doi.org/10.32614/RJ-2010-006

Hale, K. R. S., Thébault, E., & Valdovinos, F. S. (2023). *A general trait-based model for multiplex*

*ecological        networks*        (p.        2023.08.08.552546).        bioRxiv.

https://doi.org/10.1101/2023.08.08.552546

Hembry, D. H., Raimundo, R. L. G., Newman, E. A., Atkinson, L., Guo, C., Guimarães, P. R., & Gillespie,

R. G. (2018). Does biological intimacy shape ecological network structure? A test using a brood

pollination mutualism on continental and oceanic islands. *Journal of Animal Ecology*, *87*(4),

1160–1171. https://doi.org/10.1111/1365-2656.12841

Hoeppke, C., & Simmons, B. (2021). Maxnodf: An R package for fair and fast comparisons of nestedness

between networks. *Methods in Ecology and Evolution*, *12*. https://doi.org/10.1111/2041-

210x.13545

Jacquemyn, H., Brys, R., Waud, M., Busschaert, P., & Lievens, B. (2015). Mycorrhizal networks and

coexistence in species-rich orchid communities. *New Phytologist*, *206*(3), 1127–1134.

https://doi.org/10.1111/nph.13281

Jacquemyn, H., Merckx, V., Brys, R., Tyteca, D., Cammue, B. P. A., Honnay, O., & Lievens, B. (2011).

Analysis of network architecture reveals phylogenetic constraints on mycorrhizal specificity in

the    genus    Orchis    (Orchidaceae).    *New    Phytologist*,    *192*(2),    518–528.

https://doi.org/10.1111/j.1469-8137.2011.03796.x

Jordano, P., Bascompte, J., & Olesen, J. M. (2002). Invariant properties in coevolutionary networks of

plant-animal interactions: Invariant properties in coevolutionary networks. *Ecology Letters*,

*6*(1), 69–81. https://doi.org/10.1046/j.1461-0248.2003.00403.x

Klironomos, J. N. (2003). Variation in plant response to native and exotic arbuscular mycorrhizal fungi.

*Ecology*, *84*(9), 2292–2301. https://doi.org/10.1890/02-0413

Krause, A. E., Frank, K. A., Mason, D. M., Ulanowicz, R. E., & Taylor, W. W. (2003). Compartments

revealed in food-web structure. *Nature*, *426*(6964), 282–285.

https://doi.org/10.1038/nature02115

Lanuza, J. B., Allen-Perkins, A., & Bartomeus, I. (2023). The non-random assembly of network motifs in

plant–pollinator networks. *Journal of Animal Ecology*, *92*(3), 760–773.

https://doi.org/10.1111/1365-2656.13889

Lê, S., Josse, J., & Husson, F. (2008). FactoMineR : an R package for multivariate analysis. *Journal of*

*Statistical Software*, *25*(1). https://doi.org/10.18637/jss.v025.i01

Lewinsohn, T. M., Inácio Prado, P., Jordano, P., Bascompte, J., & M. Olesen, J. (2006). Structure in

plant–animal interaction assemblages. *Oikos*, *113*(1), 174–184.

https://doi.org/10.1111/j.0030-1299.2006.14583.x

Lewitus, E., & Morlon, H. (2016). Characterizing and comparing phylogenies from their Laplacian

spectrum. *Systematic Biology*, *65*(3), 495–507. https://doi.org/10.1093/sysbio/syv116

Lima-Mendez, G., Faust, K., Henry, N., Decelle, J., Colin, S., Carcillo, F., Chaffron, S., Ignacio-Espinosa, J.

C., Roux, S., Vincent, F., Bittner, L., Darzi, Y., Wang, J., Audic, S., Berline, L., Bontempi, G.,

Cabello, A. M., Coppola, L., Cornejo-Castillo, F. M., … Raes, J. (2015). Determinants of

community structure in the global plankton interactome. *Science*, *348*(6237), 1262073–

1262073. https://doi.org/10.1126/science.1262073

Losapio, G., Schöb, C., Staniczenko, P. P. A., Carrara, F., Palamara, G. M., De Moraes, C. M., Mescher,

M. C., Brooker, R. W., Butterfield, B. J., Callaway, R. M., Cavieres, L. A., Kikvidze, Z., Lortie, C. J.,

Michalet, R., Pugnaire, F. I., & Bascompte, J. (2021). Network motifs involving both competition

and facilitation predict biodiversity in alpine plant communities. *Proceedings of the National*

*Academy of Sciences*, *118*(6), e2005759118. https://doi.org/10.1073/pnas.2005759118

Maliet, O., Loeuille, N., & Morlon, H. (2020). An individual-based model for the eco-evolutionary

emergence of bipartite interaction networks. *Ecology Letters*, ele.13592.

https://doi.org/10.1111/ele.13592

Martos, F., Munoz, F., Pailler, T., Kottke, I., Gonneau, C., & Selosse, M.-A. (2012). The role of epiphytism

in architecture and evolutionary constraint within mycorrhizal networks of tropical orchids.

*Molecular Ecology*, *21*(20), 5098–5109. https://doi.org/10.1111/j.1365-294X.2012.05692.x

Mello, M. A. R., Marquitti, F. M. D., Guimarães, P. R., Kalko, E. K. V., Jordano, P., & de Aguiar, M. A. M.

(2011). The missing part of seed dispersal networks: Structure and robustness of bat-fruit

interactions. *PLoS ONE*, *6*(2), e17395. https://doi.org/10.1371/journal.pone.0017395

Michalska-Smith, M. J., & Allesina, S. (2019). Telling ecological networks apart by their structure: A

computational challenge. *PLOS Computational Biology*, *15*(6), e1007076.

https://doi.org/10.1371/journal.pcbi.1007076

Milo, R., Shen-Orr, S., Itzkovitz, S., Kashtan, N., Chklovskii, D., & Alon, U. (2002). Network motifs: Simple

building blocks of complex networks. *Science*, *298*(5594), 824–827.

Montesinos-Navarro, A., Hiraldo, F., Tella, J. L., & Blanco, G. (2017). Network structure embracing

mutualism–antagonism continuums increases community robustness. *Nature Ecology &*

*Evolution*, *1*(11), 1661–1669. https://doi.org/10.1038/s41559-017-0320-6

Montesinos-Navarro, A., Segarra-Moragues, J. G., Valiente-Banuet, A., & Verdú, M. (2012). Plant

facilitation occurs between species differing in their associated arbuscular mycorrhizal fungi.

*New Phytologist*, *196*(3), 835–844. https://doi.org/10.1111/j.1469-8137.2012.04290.x

Morlon, H., Lewitus, E., Condamine, F. L., Manceau, M., Clavel, J., & Drury, J. (2016). RPANDA : an R

package for macroevolutionary analyses on phylogenetic trees. *Methods in Ecology and*

*Evolution*, *7*(5), 589–597. https://doi.org/10.1111/2041-210X.12526

Newman, M. E. J. (2006a). Finding community structure in networks using the eigenvectors of matrices.

*Physical Review E*, *74*(3), 036104. https://doi.org/10.1103/PhysRevE.74.036104

Newman, M. E. J. (2006b). Modularity and community structure in networks. *Proceedings of the*

*National Academy of Sciences*, *103*(23), 8577–8582.

https://doi.org/10.1073/pnas.0601602103

Nielsen, A., & Bascompte, J. (2007). Ecological Networks, Nestedness and Sampling Effort. *Journal of*

*Ecology*, *95*(5), 1134–1141.

Okuyama, T., & Holland, J. N. (2008). Network structural properties mediate the stability of mutualistic

communities. *Ecology Letters*, *11*(3), 208–216. https://doi.org/10.1111/j.1461-

0248.2007.01137.x

Olesen, J. M., Bascompte, J., Dupont, Y. L., & Jordano, P. (2007). The modularity of pollination

networks. *Proceedings of the National Academy of Sciences*, *104*(50), 19891–19896.

https://doi.org/10.1073/pnas.0706375104

Öpik, M., Metsis, M., Daniell, T. J., Zobel, M., & Moora, M. (2009). Large-scale parallel 454 sequencing

reveals host ecological group specificity of arbuscular mycorrhizal fungi in a boreonemoral

forest. *New Phytologist*, *184*(2), 424–437. https://doi.org/10.1111/j.1469-8137.2009.02920.x

Perez-Lamarque, B., Petrolli, R., Strullu-Derrien, C., Strasberg, D., Morlon, H., Selosse, M.-A., & Martos,

F. (2022). Structure and specialization of mycorrhizal networks in phylogenetically diverse

tropical communities. *Environmental Microbiome*, *17*(1), 38. https://doi.org/10.1186/s40793-

022-00434-0

Perez-Lamarque, B., Selosse, M., Öpik, M., Morlon, H., & Martos, F. (2020). Cheating in arbuscular

mycorrhizal mutualism: A network and phylogenetic analysis of mycoheterotrophy. *New*

*Phytologist*, *226*(6), 1822–1835. https://doi.org/10.1111/nph.16474

Pilosof, S., Fortuna, M. A., Cosson, J.-F., Galan, M., Kittipong, C., Ribas, A., Segal, E., Krasnov, B. R.,

Morand, S., & Bascompte, J. (2014). Host–parasite network structure is associated with

community-level immunogenetic diversity. *Nature Communications*, *5*(1), 5172.

https://doi.org/10.1038/ncomms6172

Pires, M. M., & Guimarães, P. R. (2013). Interaction intimacy organizes networks of antagonistic

interactions in different ways. *Journal of The Royal Society Interface*, *10*(78), 20120649.

https://doi.org/10.1098/rsif.2012.0649

Pocock, M. J. O., Evans, D. M., & Memmott, J. (2012). The robustness and restoration of a network of

ecological networks. *Science*, *335*(6071), 973–977. https://doi.org/10.1126/science.1214915

Põlme, S., Bahram, M., Jacquemyn, H., Kennedy, P., Kohout, P., Moora, M., Oja, J., Öpik, M., Pecoraro,

L., & Tedersoo, L. (2018). Host preference and network properties in biotrophic plant-fungal

associations. *New Phytologist*, *217*(3), 1230–1239. https://doi.org/10.1111/nph.14895

Rezende, E. L., Jordano, P., & Bascompte, J. (2007). Effects of phenotypic complementarity and

phylogeny on the nested structure of mutualistic networks. *Oikos*, *116*(11), 1919–1929.

https://doi.org/10.1111/j.0030-1299.2007.16029.x

Rezende, E. L., Lavabre, J. E., Guimarães, P. R., Jordano, P., & Bascompte, J. (2007). Non-random

coextinctions in phylogenetically structured mutualistic networks. *Nature*, *448*(7156), 925–

928. https://doi.org/10.1038/nature05956

Rohr, R. P., Saavedra, S., & Bascompte, J. (2014). On the structural stability of mutualistic systems.

*Science*, *345*(6195), 1253497–1253497. https://doi.org/10.1126/science.1253497

Sachs, J. L., Skophammer, R. G., & Regus, J. U. (2011). Evolutionary transitions in bacterial symbiosis.

*Proceedings of the National Academy of Sciences*, *108*(Supplement_2), 10800–10807.

https://doi.org/10.1073/pnas.1100304108

Santamaría, L., & Rodríguez-Gironés, M. A. (2007). Linkage Rules for Plant–Pollinator Networks: Trait

Complementarity or Exploitation Barriers? *PLoS Biology*, *5*(2), e31.

https://doi.org/10.1371/journal.pbio.0050031

Selosse, M., Schneider-Maunoury, L., & Martos, F. (2018). Time to re-think fungal ecology? Fungal

ecological niches are often prejudged. *New Phytologist*, *217*, 968–972.

https://doi.org/10.1111/nph.14983

Shen, H.-W., & Cheng, X.-Q. (2010). Spectral methods for the detection of network community

structure: A comparative analysis. *Journal of Statistical Mechanics: Theory and Experiment*,

*2010*(10), P10020.

Simmons, B. I., Cirtwill, A. R., Baker, N. J., Wauchope, H. S., Dicks, L. V., Stouffer, D. B., & Sutherland,

W. J. (2019). Motifs in bipartite ecological networks: Uncovering indirect interactions. *Oikos*,

*128*(2), 154–170. https://doi.org/10.1111/oik.05670

Simmons, B. I., Sweering, M. J. M., Schillinger, M., Dicks, L. V., Sutherland, W. J., & Clemente, R. D.

(2018). *bmotif: A package for motif analyses of bipartite networks* [Preprint]. Ecology.

https://doi.org/10.1101/302356

Simmons, B. I., Sweering, M. J. M., Schillinger, M., Dicks, L. V., Sutherland, W. J., & Di Clemente, R.

(2019). bmotif: A package for motif analyses of bipartite networks. *Methods in Ecology and*

*Evolution*, *10*(5), 695–701. https://doi.org/10.1111/2041-210X.13149

Song, C., & Saavedra, S. (2020). Telling ecological networks apart by their structure: An environment-

dependent approach. *PLOS Computational Biology*, *16*(4), e1007787.

https://doi.org/10.1371/journal.pcbi.1007787

Srivastava, N., Hinton, G., Krizhevsky, A., Sutskever, I., & Salakhutdinov, R. (2014). Dropout: A simple

way to prevent neural networks from overfitting. *Journal of Machine Learning Research*,

*15*(56), 1929–1958.

Staniczenko, P. P. A., Kopp, J. C., & Allesina, S. (2013). The ghost of nestedness in ecological networks.

*Nature Communications*, *4*(1), 1391. https://doi.org/10.1038/ncomms2422

Stouffer, D. B., & Bascompte, J. (2010). Understanding food-web persistence from local to global scales.

*Ecology Letters*, *13*(2), 154–161. https://doi.org/10.1111/j.1461-0248.2009.01407.x

Strydom, T., Catchen, M. D., Banville, F., Caron, D., Dansereau, G., Desjardins-Proulx, P., Forero-Muñoz,

N. R., Higino, G., Mercier, B., Gonzalez, A., Gravel, D., Pollock, L., & Poisot, T. (2021). A roadmap

towards predicting species interaction networks (across space and time). *Philosophical*

*Transactions of the Royal Society B: Biological Sciences*, *376*(1837), 20210063.

https://doi.org/10.1098/rstb.2021.0063

Suweis, S., Simini, F., Banavar, J. R., & Maritan, A. (2013). Emergence of structural and dynamical

properties of ecological mutualistic networks. *Nature*, *500*(7463), 449–452.

https://doi.org/10.1038/nature12438

Thébault, E., & Fontaine, C. (2008). Does asymmetric specialization differ between mutualistic and

trophic networks? *Oikos*, *117*(4), 555–563. https://doi.org/10.1111/j.0030-1299.2008.16485.x

Thébault, E., & Fontaine, C. (2010). Stability of ecological communities and the architecture of

mutualistic     and     trophic     networks.     *Science*,     *329*(5993),     853–856.

https://doi.org/10.1126/science.1188321

Thebault, E., & Fontaine, C. (2010). Stability of ecological communities and the architecture of

mutualistic     and     trophic     networks.     *Science*,     *329*(5993),     853–856.

https://doi.org/10.1126/science.1188321

Thomas, F., Renaud, F., & Guegan, J.-F. (Eds.). (2005). *Parasitism and Ecosystems*. Oxford University

Press. https://doi.org/10.1093/acprof:oso/9780198529873.001.0001

Thompson, J. N. (2005). *The Geographic Mosaic of Coevolution*. University of Chicago Press.

https://doi.org/10.7208/9780226118697

Toju, H., Guimarães, P. R., Olesen, J. M., & Thompson, J. N. (2015). Below-ground plant–fungus network

topology is not congruent with above-ground plant–animal network topology. *Science*

*Advances*, *1*(9), e1500291. https://doi.org/10.1126/sciadv.1500291

Vázquez, D. P., Chacoff, N. P., & Cagnolo, L. (2009). Evaluating multiple determinants of the structure

of plant–animal mutualistic networks. *Ecology*, *90*(8), 2039–2046. https://doi.org/10.1890/08-

1837.1

Vitali, A., Sasal, Y., Vázquez, D. P., Miguel, M. F., & Rodríguez-Cabal, M. A. (2022). The disruption of a

keystone interaction erodes pollination and seed dispersal networks. *Ecology*, *103*(1).

https://doi.org/10.1002/ecy.3547

Wu, Z., Pan, S., Chen, F., Long, G., Zhang, C., & Yu, P. S. (2021). A comprehensive survey on graph neural

networks. *IEEE Transactions on Neural Networks and Learning Systems*, *32*(1), 4–24.

https://doi.org/10.1109/TNNLS.2020.2978386

Xing, X., Jacquemyn, H., Gai, X., Gao, Y., Liu, Q., Zhao, Z., & Guo, S. (2019). The impact of life form on

the architecture of orchid mycorrhizal networks in tropical forest. *Oikos*, *128*(9), 1254–1264.

https://doi.org/10.1111/oik.06363

1107