Uncoupled evolution of the Polycomb system and deep origin of non-canonical
PRC1

Bastiaan de Potter[1,†], Maximilian W. D. Raas[1,2], Michael F. Seidl[1], C. Peter Verrijzer[3#], Berend
Snel[1#*]

[1]Theoretical Biology and Bioinformatics, Department of Biology, Science Faculty, Utrecht
University, Utrecht, Netherlands

[2,†]Hubrecht institute, Royal Netherlands Academy of Arts and Sciences, Utrecht, Netherlands

[3]Department of Biochemistry, Erasmus University Medical Center, Rotterdam, Netherlands

[†]Present address

[#]Equal contribution

[*]Corresponding author

Email: b.snel@uu.nl

Keywords: Polycomb, Evolution, PRC1, PRC2, Comparative genomics

Abstract

Polycomb group (PcG) proteins modulate chromatin states to silence gene transcription in plants and animals. Most PcG proteins function as part of distinct multi-subunit Polycomb repressive complexes (PRCs). Gene repression by the Polycomb system involves chromatin compaction by canonical PRC1 (cPRC1), mono-ubiquitylation of histone H2A (H2Aub1) by non-canonical PRC1 (ncPRC1) and tri-methylation of histone H3K27 (H3K27me3) by PRC2. Prevalent models for Polycomb repression emphasize a tight functional coupling between PRC1 and PRC2. However, whether this paradigm indeed reflects the evolution and functioning of the Polycomb system remains unclear. Here, we examined the relationship between cPRC1, ncPRC1 and PRC2 through a comprehensive analysis of their presence and evolution across the entire eukaryotic tree of life. We show that both PRC1 and PRC2 were present in the Last Eukaryotic Common Ancestor (LECA), but that their subsequent evolution is uncoupled. The identification of orthologs for ncPRC1-defining subunits in unicellular relatives of animals and of fungi suggests that the origin of ncPRC1 predates that of cPRC1, and we develop a scenario for the evolution of cPRC1 from ncPRC1. Our results demonstrate the independent evolution and function of PRC1 and PRC2 and show that crosstalk between these complexes is a secondary development in evolution.

Main Text

Introduction

Packaging of the eukaryotic genome into chromatin is crucial for its maintenance, replication and expression. The nucleosome, comprising 147 bp stretches of DNA wrapped in a left-handed superhelical turn around a histone octamer, is the fundamental repeating unit of chromatin. Histones and additional structural chromatin proteins impede access of DNA-binding proteins, such as transcription factors, thereby generating a fundamental level of gene regulation. Consequently, modulation of chromatin structure through post-translational histone modifications, nucleosome re-positioning and higher-order folding are central to eukaryotic gene regulation (Bracken et al. 2019). Polycomb group (PcG) proteins are conserved chromatin regulators that repress gene transcription to maintain cellular identity in plants and animals (Schuettengruber et al. 2017; Kassis et al. 2017; Baile et al. 2022). The *PcG* genes were first identified in *Drosophila* as repressors that are crucial to prevent mis-expression of homeotic (*Hox*) genes in inappropriate parts of the body (Schuettengruber et al. 2017; Kassis et al. 2017; Baile et al. 2022). This function is reflected by the name Polycomb, which refers to the extra sex combs appearing on the second and third pair of legs of male flies, due to de-repression of the *sex combs reduced* gene. Subsequent research showed that, in addition to silencing of *Hox* genes, PcG proteins are involved in the regulation of cell proliferation, stem cell pluripotency and oncogenesis (Bracken et al. 2019; Hennig and Derkacheva 2009; Kim and Kingston 2022; Piunti and Shilatifard 2021).

PcG proteins function as part of two main classes of multi-protein complexes, named Polycomb Repressive Complex (PRC) 1 and 2, which change chromatin structure across multiple scales (Baile et al. 2022; Kang et al. 2022; Kim and Kingston 2022; Piunti and Shilatifard 2021; Schuettengruber et al. 2017; Loubiere et al. 2019). The PRC2 methyltransferases mediate mono-, di- and tri-methylation of H3K27 (H3K27me3), whereas the PRC1s mediate chromatin compaction and mono-ubiquitylation of H2A (H2Aub1) on Lys 118, 119 or 120 in *Drosophila*, mammals or *Arabidopsis*, respectively (**Fig.1A**). The catalytic core of PRC2 comprises the methyltransferase EZH associated with EED, SUZ12 and RBBP (or their paralogs) (Yu et al. 2019; Laugesen et al.

2019). EZH proteins contain a catalytic SET domain that is responsible for methylating H3K27,

while EED is a WD40 repeat protein that allosterically activates EZH upon binding to H3K27me3,

thus creating a positive feedback loop (Yu et al. 2019; Laugesen et al. 2019; Glancy et al. 2021).

The C-terminal VEFS-box of SUZ12 further stabilizes the interaction between EED and EZH. The

WD40 repeat protein RBBP, enhances chromatin association through direct binding to histones.

Selective association of the PRC2 core with accessory proteins generates unique subassemblies

with different functional properties (Glancy et al. 2021; Baile et al. 2022; Laugesen et al. 2019; Yu

et al. 2019; Bieluszewski et al. 2021).

The PRC1 class can be subdivided in canonical (cPRC1) and non-canonical PRC1

(ncPRC1) (Baile et al. 2022; Kang et al. 2022; Loubiere et al. 2019; Piunti and Shilatifard 2021).

Both are assembled around a RING-RING heterodimer, which in animals comprise a RING1 protein

paired with a PcG Finger (PCGF) protein. Notably, plant RING1 and PCGF orthologs can form

homo- as well as heterodimers (Baile et al. 2022). RING1 and PCGF proteins share a similar

structural organization characterized by an N-terminal zf-RING domain and a C-terminal RAWUL

domain (**Fig.1B**). cPRC1 is defined by the association of a Chromobox protein (CBX) and

Polyhomeotic homolog (PHC) subunit, which are absent in ncPRC1 (Gao et al. 2012). Additionally,

SCM is a PcG protein that is more loosely associated with cPRC1. The presence of a RYBP/YAF

subunit defines the ncPRC1s, which through association with additional partner proteins form

multiple variant complexes. Pertinently, most or all of H2Aub1 is deposited by ncPRC1 rather than

cPRC1 (Tamburri et al. 2022; Barbour et al. 2020; Lagarou et al. 2008).

The relative importance of the different PRC activities and their order of recruitment to

chromatin remains controversial (Bracken et al. 2019; Kim and Kingston 2022; Loubiere et al. 2019;

Piunti and Shilatifard 2021; Tamburri et al. 2022). While there is compelling evidence for the

essential function of H3K27me3 for Polycomb repression (Sankar et al. 2022; Pengelly et al. 2013),

the role of H2Aub1 is more ambiguous (Barbour et al. 2020). Although H2Aub1 is essential for

animal viability, mutational studies established that PRC1 rather than its enzymatic activity is

essential for transcriptional repression of canonical Polycomb target genes (Bonnet et al. 2022;

Illingworth et al. 2015; Pengelly et al. 2015; Tsuboi et al. 2018). Moreover, cPRC1 mediates long-

range chromatin interactions via polymerization of the SAM domains within PH, which is

independent of H2Aub1 (Isono et al. 2013; Boyle et al. 2020; Seif et al. 2020). Thus, it appears that

chromatin compaction is essential for Polycomb repression in animals, while H2Aub1 stimulates

this process but is not absolutely essential. Intriguingly, while the core subunits RING1 and PCGF

of PRC1 are well-conserved in plants, the nc- and cPRC1 accessory subunits appear to be absent.

This implies an alternate scenario for the potential formation of higher-order Polycomb structures

in plants (Baile et al. 2022). The binding of CBX proteins in cPRC1 to H3K27me3, and conversely,

the binding of PRC2-accessory proteins to H2Aub1, provides cross-talk between PRC1 and PRC2

(Glancy et al. 2021; Tamburri et al. 2022; Blackledge et al. 2020; Tamburri et al. 2020). These

observations prompted the formulation of hierarchical recruitment models through "writing" and

"reading" of histone modifications by PRC1 and PRC2 **(Fig.1A)** (Bracken et al. 2019; Kim and

Kingston 2022; Loubiere et al. 2019; Piunti and Shilatifard 2021; Tamburri et al. 2022). At odds with

such scenarios, there are accumulating examples of largely uncoupled gene repression by PRC1

and PRC2 (Bonnet et al. 2022; Cohen et al. 2021; Leeb et al. 2010; Pengelly et al. 2015; Sijm et

al. 2022; Zepeda-Martinez et al. 2020; Chiacchiera et al. 2016; Rossi et al. 2016). Moreover, it is

becoming increasingly clear that in mammals, as previously established in *Drosophila*, DNA-

binding plays a major role in the genomic targeting of PRCs (Bracken et al. 2019; Glancy et al.

2021; Loubiere et al. 2019; Piunti and Shilatifard 2021; Kassis et al. 2017). In summary, PRC

recruitment involves a combination of cooperative- and redundant mechanisms that may differ

depending on the cellular- or genomic context. However, the functional- and evolutionary coupling

between PRC1 and PRC2 remains an unresolved question.

Our knowledge of the function and evolution of the Polycomb system is primarily based on

studies on animals and plants. This has limited our understanding of its role in developmental gene

regulation in other eukaryotic lineages (Gahan et al. 2020; Huang et al. 2017, 2019; Berke and

Snel 2015). Recent research has started to expand the diversity of eukaryotes in which Polycomb

repression is analyzed. For example, studies in ciliates have shown that PRC2-mediated

trimethylation of histones is involved in the silencing of transposons (Xu et al. 2021; Wang et al. 2022; Déléris et al. 2021). In addition, a phylogenetic screening has shown that PRC2 is conserved across major eukaryotic groups and is likely present in the last eukaryotic common ancestor (LECA) (Sharaf et al. 2022). However, PRC1 has not been studied as extensively in non-animal lineages, and therefore, whether PRC1 and PRC2 co-evolved remains unresolved.

In the present study we used highly sensitive and comprehensive phylogenetic profiling to analyze the evolutionary history of PRC1 and PRC2 across the available scope of genomes covering the eukaryotic tree of life. Our analysis suggests that both PRC1 and PRC2 were present in LECA. Importantly, we found that while their intra-complex subunits evolved cohesively, PRC1 and PRC2 evolve independently. Furthermore, we discovered previously unreported ncPRC1-defining subunits in the relatives of animals and fungi, indicating that the differentiation of ncPRC1 occurred before cPRC1. These findings provide a foundation for future research on the biological functions of these proteins in a wider range of eukaryotes and in areas where PRC1 is understudied.

Results

Conservation of PRC1 subunits throughout eukaryotes traces its presence to LECA

To study the evolution of PRC1 beyond animals and plants, we traced the presence of PRC1 core subunits, RING1 and PCGF, in a diverse collection of 178 eukaryotes that covers all known major eukaryotic groups **(SI Data and Methods and External Dataset S1)**. Both RING1 and PCGF consist of an N-terminal zf-RING domain and a C-terminal RAWUL domain **(Fig.1B)**. These protein domains are the structural and functional units that are responsible for PRC1 functionality. As such, the conservation of specific protein domains across different species can be a strong indication that the protein is performing the same or similar functions in those species. To study the domain conservation of RING1 and PCGF proteins across our major eukaryotic groups,

we utilized Pfam (v.35) domain profile hidden-Markov models (HMMs) to annotate the sequences

included in the phylogenetic trees (Mistry et al. 2021). These profiles, however sometimes failed to

detect zf-RING or RAWUL domains in protein sequences that clustered in potential orthologous

groups, while their presence was evident after manual inspection of sequence alignments

combined with AlphaFold2 structural predictions (Varadi et al. 2022; Mirdita et al. 2022; Jumper et

al. 2021). To overcome the lack of sensitivity in default Pfam profiles (v.35), we manually curated

HMM profiles for the zf-RING, and RAWUL domain, by combining sequence alignments with

structural predictions by AlphaFold2 (Varadi et al. 2022; Jumper et al. 2021). A detailed description

of our approach and curated profiles are presented in **SI Data and Methods and External Dataset**

S2.

This approach enabled us to identify 69 and 67 species with high-confidence orthologs for

RING1 and PCGF, respectively. We considered an ortholog of high-confidence if it clustered

monophyletically with the other sequences and contained both the zf-RING and RAWUL domain

as predicted by our sequence models and AlphaFold2 **(Fig.2, SI Fig.S1A,B, Fig.S2)**. Additionally,

we detected 15 species with a putative RING1- and ten species with putative PCGF orthologs,

containing a zf-RING-, but lacking a recognizable RAWUL domain (**SI Fig.S1A,B, Fig.S2)**. The

phylogenetic trees of RING1 and PCGF contained multiple groups of sequences from species that

are part of all major eukaryotic groups in which the deep duplication nodes displayed large species

overlap, which is a defining feature of orthologous groups in a gene phylogeny. The subsequent

speciation node can thus be identified with high reliability as a LECA presence **(Fig.2, 3A, S1A,B)**.

The increased sensitivity of our approach is illustrated by the detection of a previously unknown

RING1 ortholog in the diatom (Stramenopile) *Phaeodactylum tricornutum*. In addition, we were able

to detect a PCGF ortholog in *Arabidopsis thaliana* that was missed in a recent study (Grau-Bové et

al. 2022), most likely because the canonical Pfam model (v.35) is not sensitive enough to detect

the RAWUL domain. Importantly, we identified RING1 and PCGF orthologs in all unicellular

eukaryotes that are currently placed proximal to the root of the eukaryotic tree of life, such as

Discoba, Metamonada, Malawimonadida, CRuMs, Hemimastigophora, and Provora **(Fig.2,**
S1A,B).

Although the eukaryotic group of fungi is generally believed to have lost PRC1
(Schuettengruber et al. 2017), we uncovered PRC1 orthologs in multiple deep branching fungal
species. For instance, we observed PRC1 (either RING1 or PCGF) in the deep branching fungi
Rozella allomycis and *Gonapodya prolifera* **(Fig.2,3B)**. Moreover, we confirmed the presence of
PRC1 orthologs in the fungus *Spizellomyces punctatus* (Grau-Bové et al. 2022). Collectively, our
phylogenetic analysis reveals that the PRC1 core was already present in LECA and remained
highly conserved throughout eukaryotic life **(Fig.2, SI Fig.S2)**.

Figure 1. Schematic representation of the core subunits of PRC1 and PRC2 and their enzymatic activities. (A)
Overview of histone modifications by PRC1 and PRC2, and cross-talk via recognition of each other's histone
marks (B) Schematic representation of the core subunits of PRC1 and PRC2. Shown are the major conserved
domains for each protein based on structural information in humans. In EZH, we did not depict the additional
SBD, EBD, BAM, SAL, and SRM domain. RAWUL, RING finger- and WD40-associated ubiquitin-like; MCSS,
motif connecting SANT1 and SANT2; SANT, SWI3, ADA2, N-CoR and TFIIIB DNA-binding domain; CxC,
cysteine-rich domain; SET, Su(var) 3–9, enhancer of zeste, trithorax domain; WD, WD-40 domain; WDB, WD-
40 binding domain; Zn, Zn-finger region; VEFS, conserved among VRN2-EMF2-FIS2-SU(Z)12.

Evidence for the presence of PRC2 in LECA

To reconstruct the evolution of PRC2 we traced the functional core of PRC2, consisting of

EZH, EED, SUZ12 and RBBP (Glancy et al. 2021; Bieluszewski et al. 2021), across our diverse

set of eukaryotes. Similar to PRC1, we studied the domain conservation of these subunits. Because

Pfam domain profiles (v.35) again failed to detect key domains in a substantial number of orthologs,

we curated our HMM profiles by combining sequence alignments with AlphaFold2 structural

predictions **(Fig.1B, SI Data and Methods and External Dataset S2)**. Within our selection of 178

eukaryotes, we detected 94 species with orthologs for EZH, 91 with EED, 87 with SUZ12, and 178

with RBBP **(Fig.2, SI Fig.S1C-E)**. In accordance with our phylogenetic analysis of PRC1, the

phylogenetic trees of PRC2 subunits also exhibited numerous orthologous groups with extensive

species overlap. This observation thus similarly enables the reliable inference of the speciation

node and the presence in LECA for the PRC2 subunits. EZH serves as the key enzymatic subunit

and is a crucial indicator of the presence of a functional PRC2 (Kasinath et al. 2018). In our

phylogenetic profile, EZH, EED, and SUZ12 exhibited strong co-occurrence. However, the co-

occurrence of RBBP with other PRC2 was less prominent, owing to its involvement in various other

protein assemblies (Suganuma et al. 2008). The phylogenetic trees of EZH and RBBP had the

highest support values throughout all major eukaryotic groups **(SI Fig.S1C-E)**. In contrast, the

phylogenomic delineation of EED orthologs was more ambiguous, due to their low sequence

conservation which results in trees with lower resolution as well as difficulties in findings homologs

and annotating the tree with protein domains **(SI Fig.S1D)**. For instance, PRC2 was recently

experimentally identified in the ciliates *Paramecium tetraulia* and *Tetrahymena thermophila* (Xu et

al. 2021; Wang et al. 2022), but the EED sequences in these organisms were undetectable by

standard sequence-based homology methods. By performing a HMM search with a query based

on the profile of the sequence alignment of these two sequences however, we retrieved EED

orthologs in most ciliates, but not in the Alveolates *Stentor coeruleus* and *Colponemida vietnamica*

(Fig.3C).

Figure 2. The evolutionary history of the core subunits of PRC1 and PRC2 can be traced back to LECA
(Fig.S1A-E). Shown is a summary of the phylogenetic profiles of PRC1 and PRC2 in all major eukaryotic
groups included in our study **(SI Fig.S2)**. Detected presences are presented as fractions of the species
included in that group illustrated as a pie chart. For groups in which we only detected putative orthologs, only
the outer line of the pie chart was colored.

Like EED, SUZ12 was also found to be highly divergent in various species, e.g., a recent

study demonstrated that MES-3 in *Caenorhabditis elegans* is a diverged SUZ12 ortholog rather

than a *Caenorhabditis elegans* specific addition to PRC2 (Snel et al. 2022). The N-terminus of

SUZ12 in animals contains five motifs: a zinc-finger binding (ZnB), WD-domain binding 1 (WDB1),

C2 domain, zinc finger (Zn), and WD-domain binding 2 (WDB2) **(Fig.1B)** (Chen et al. 2018; Glancy

et al. 2021; Kasinath et al. 2018). All these motifs are highly conserved throughout animals and

plants. Pertinently, the C-terminal VEFS box of SUZ12 is crucial for the formation of the PRC2

catalytic lobe (Chen et al. 2018; Glancy et al. 2021; Kasinath et al. 2018). Therefore, we used the

VEFS-box as the basis for our inference analysis to systematically detect SUZ12 orthologs. This

approach uncovered a strong correlation between SUZ12 orthologs and the phylogenetic profiles

of the other PRC2 subunits **(SI Fig.S2)**. Note that examples of SUZ12-like orthologs without a

VEFS-box were recently described, e.g., *Paramecium tetraulia* has an adapted SUZ12-like ortholog

without a VEFS-box (Wang et al. 2022). Consequently, we could not detect these SUZ12-like

orthologs in our phylogenetic inference **(Fig.3C)**. In addition, a previous study reported that in the

yeast *Cryptococcus neoformans,* SUZ12 is substituted by a structurally unrelated protein, Bnd1

(Dumesic et al. 2015). Indeed, we initially missed a *Cryptococcus neoformans* SUZ12 ortholog in

the yeast *Cryptococcus neoformans*. However, by combining HHpred searches with AlphaFold2

structural predictions, we found that, like MES-3 in *Caenorhabditis elegans,* Bnd1 is most likely a

highly diverged SUZ12 ortholog **(SI Fig.S3)**. Further inferred absences of subunits predicted by our

analysis might thus have been substituted by other, currently unknown, subunits, or were not

detected due to high sequence divergence (e.g., EED, SUZ12). Consistent with the notion that

EZH, EED and SUZ12 comprise the crucial functional core of PRC2, these subunits typically co-

occur across the eukaryotic tree of life, reflecting the close and ancient biochemical

interdependence of EZH, EED and SUZ12.

Figure 3. Phylogenetic profiles of the core subunits of PRC1 and PRC2. (A) Phylogenetic profiles of species
in major eukaryotic groups proximal to the eukaryotic root, solidifying their presence in LECA. (B)
Representation of the presence of PRC1 and PRC2 core subunits in fungi, a group that was previously
believed to have lost PRC1 completely. (C) Phylogenetic profile of ciliates, highlighting the detection of highly
diverged EED sequences.

Intra-complex evolution of PRC1 and PRC2 is cohesive while inter-complex evolution is
uncoupled

The evolution of RING1 and PCGF is largely coupled, i.e. these core PRC1 subunits were mostly lost or retained together during eukaryotic evolution **(Fig.4A)**. Likewise, the genes encoding the core of PRC2 were typically lost or retained together. This suggests that both PRC1 and PRC2 function and evolve as cohesive units. To obtain a quantitative measure of their cohesiveness throughout eukaryotic evolution, we calculated the average Pearson's correlation coefficient between the phylogenetic profiles of the subunits of PRC1 and PRC2 **(Fig.4B)**. We found that the subunits of the PRC2 core (EZH, EED, and SUZ12) had a high average correlation ($r = 0.73$), as did the core subunits of PRC1 (RING1 and PCGF, $r = 0.7$) **(SI Fig.S4)**. This confirms that these subunits tend to evolve together within their resident complexes. As expected given its participation in multiple additional chromatin regulating pathways, RBBP only correlated weakly with other PRC2 subunits.

These observations raise the question whether the evolution of PRC1 and PRC2 might be coupled too. To address this question, we scored the presence of PRC1 or PRC2 only if all defining core subunits were present (i.e., both RING1 and PCGF for PRC1, and EZH, EED and SUZ12 for PRC2). Our analysis of 178 eukaryotes, selected to cover all known major eukaryotic groups, revealed that 40 species retained both PRC1 and PRC2, 12 lost PRC2 but retained PRC1, 17 retained only PRC2, whereas 52 lost both complexes. The Polycomb status of the remaining 57 eukaryotes remains unclear because they contained sequences without a recognizable RAWUL domain or harbored apparently partial PRCs. Thus, ~33% of species we analyzed contains remnants of the Polycomb system but lacks an unambiguous complete PRC1 or PRC2. About 22% of eukaryotes retained both PRC1 and PRC2, ~16% of eukaryotes retained either PRC1 or PRC2, without a clear preference for one over the other, while ~29% lost both. Thus, a substantial portion (~42%) of organisms that contain (part of) the Polycomb system use only PRC1 or only PRC2. We expect that with increasingly sensitive detection, the degree of uncoupling between PRC1 and

PRC2 might turn-out to be even higher than presented in this study. The presence of either only

PRC1 or only PRC2 was particularly noticeable in Amoebozoans, fungi, Cryptista, Haptista,

Rhodophyta, Glaucophyta, Chloroplastida, and SAR, but we were unable to determine a consistent

trend of either complex being more likely to be lost or retained **(Fig.4A)**. In conclusion, while PRC1

and PRC2 were both present in LECA, their evolution throughout eukaryotic life has been largely

uncoupled. Concurrent conservation of both complexes was most frequent in land plants and

animals, while other major taxa displayed a more diverse pattern of conservation for PRC1 and

PRC2 (**Fig.4A**). In contrast to their high intra-complex correlations, PRC1 and PRC2 subunits only

displayed a low degree of inter-PRC correlation ($r = 0.4$). Collectively, these results reflect great

evolutionary flexibility of the Polycomb system.

Figure 4. Intra-complex evolution of PRC1 and PRC2 is cohesive, while inter-complex evolution is largely
uncoupled. (A) Phylogenetic profiles of PRC1 and PRC2 core subunits in all independent species included in
our analysis. The profiles show the overall cohesive loss- or retention of intra-complex subunits, and the largely
uncoupled evolution of PRC1 and PRC2. RBBP was not included because it does not represent the presence
or absence of the PRC2. Data was split into two panels and subunits are colored as follows: RING (blue),
PCGF (Pink), EZH (Red), Orange (EED), SUZ12 (Green). Asterisks highlight species where either PRC1 or
PRC2 is uncoupled. (B) Heatmap of the correlations between the subunits. Intra-complex subunits have a
*higher correlation than inter-complex subunits. Full names of all species are provided in **SI Fig.S2**.*

Structural similarity between RYBP and CBX suggests a common ancestor

Both PRC1 and PRC2 form subcomplexes through the incorporation of accessory subunits

or through the combinatorial assembly of paralogous components (Baile et al. 2022; Kang et al.

2022; Piunti and Shilatifard 2021; Kim and Kingston 2022). ncPRC1 and cPRC1 are distinguished

by the mutually exclusive binding of either CBX (cPRC1) or RYBP (ncPRC1) (Gao et al. 2012).

CBX harbors a chromodomain at its N-terminus and a CBX7_C domain at its C-terminus, while

RYBP contains a zf-RanBP domain at its N-terminus and a YAF2 domain at its C-terminus (**Fig.5A**).

Interestingly, both the CBX7_C- and YAF2 domain form a ß-hairpin of around 16 amino acids. From

an evolutionary perspective, mutual exclusivity (such as between RYBP and CBX) is often found

between paralogs that define variant complexes (Szklarczyk et al. 2008; Ori et al. 2016). Therefore,

we sought to investigate if CBX and RYBP might actually be distant paralogs by determining

whether there is structural or highly diverged sequence homology between them. Whereas we did

not detect full length structural- or sequence homology between CBX and RYBP, a structural

alignment of their C-terminal ß-hairpins using the TM-align tool (Zhang 2005) revealed that they

superimpose almost perfectly (RMSD = 2.35 Å) **(Fig.5A, SI External Dataset S3)**. Moreover, a

sensitive sequence profile-vs-profile (HHpred) (Hildebrand et al. 2009) search with the C-terminal

motifs of RYBP and CBX8 revealed that they were reciprocal best hits **(Fig.5B, SI External Dataset**

S3). Based on this evidence, we conclude that the domains CBX7_C and YAF are likely to be

distant paralogs. Thus, to avoid any confusion resulting from the use of multiple names referring to

the same domain, we will use ß-HP throughout the remainder of this article. Combined, the

structural similarity and the reciprocal best hits of their ß-hairpin domains, provide evidence that

CBX and RYBP might be switching paralogs in PRC1.

Figure 5. Structural homology and reciprocal best hits suggests that the ß-hairpin of RYBP and CBX are
orthologous. (A) Schematic representation of RYBP/YAF and CBX. Both the C-terminal YAF domain and
CBX7_C domain consist of a ß-hairpin. (B) Structural alignment of the ß-hairpin of RYBP, and CBX. The
structures superimpose almost perfectly (RMSD = 2.35). (C) Output of the YAF2 C-terminal ß-hairpin as the
best hit of a sensitive sequence profile-vs-profile (HHpred) search with the ß-hairpin of CBX.

Evolutionary evidence for the occurrence of ncPRC1 prior to cPRC1

To further study the emergence of ncPRC1 and cPRC1, we investigated the evolutionary

origin of their signature subunits RYBP, CBX, and PHC. Recently, a putative ortholog of RYBP,

containing only the N-terminal zf-RanBP motif, was identified in the Choanoflagellate *Salpingoeca*

rosetta, which is a close unicellular relative of animals (Gahan et al. 2020). Additionally, we

identified full length orthologs of RYBP in two Choanoflagellata, *Salpingoeca kvevrii* and

Diaphanoeca grandis. These orthologs contain both the N-terminal zf-RanBP motif as well as the

C-terminal ß-hairpin, suggesting that these are *bona fide* RYBP orthologs **(Fig.6A, SI External**

Dataset S4,5). We also detected a putative ortholog containing only the N-terminal zf-RanBP motif

in the nucleariid *Parvularia atlantis*, which is a close relative of fungi **(Fig.6A)**. In contrast, we were

not able to detect orthologs of CBX outside of animals (**SI External Dataset S4)**. Thus, it appears

that RYBP originated before CBX. Next, we performed a phylogenetic analysis of SAM domains

which are present in the PRC1-associated proteins PHC, SCML, SCMH, L3MBTL, SFMBT and

MBTD (**SI Data and Methods External Dataset S4).** The SAM domains clustered together in a

monophyletic group, reflecting a shared ancestry **(SI Fig.S1F)**, but could be separated further into

two distinctive groups; one that consisted mostly of proteins containing an MBT domain, and one

without such a domain **(Fig.6B)**. We therefore next performed a phylogenetic analysis of MBT

domains with the aim to find orthologs outside of animals that would imply an earlier origin for these

proteins (**SI Data and Methods and External Dataset S4)**. Our analysis revealed that the MBT

domains of the PRC1 accessory subunits were part of an orthologous group. Strikingly, we

identified two sequences in the relatives of animals and fungi that contained both a potential SAM-

and a MBT domain: one filasterean, *Capsaspora owczarzaki*, and the nucleariid *Parvularia atlantis*

(SI Fig.S1G). Although these sequences most likely encode orthologs of the L3MBTL/SCML PRC1

accessory subunits, the SAM- and MBT domains are in reverse order compared to their animal

counterparts **(Fig.7)**. Adding the *Capsaspora owczarzaki*, and *Parvularia atlantis* SAM domains to

our earlier inference analysis did not yield a monophyletic clustering with the SAM domains of

SCML, SCMH, L3MBTL, and PHC **(SI Fig.S1H)**. This negative result might be due the limited

phylogenetic signal of the relatively small SAM domain (65-70 amino acids). Thus, the orthology of

the filasterean, *Capsaspora owczarzaki*, and the nucleariid *Parvularia atlantis* SAM domains

remains unclear.

In summary, our collective results suggest that the ancestor of animals and fungi already

harbored RYBP and a L3MBTL-associated ortholog **(Fig.7)**. We propose a scenario in which RYBP

was lost in fungi and Filasterea, but retained in Nucleariidae, Choanoflagellata and animals.

Furthermore, we hypothesize that CBX arose via a gene duplication of RYBP at the root of animals,

after which the zf-RanBP was replaced by a chromo-domain **(Fig.7)**. Similarly, our data suggests

that the SAM domain containing accessory subunits of PRC1 most likely originated from a singular

gene rearrangement event of an ancestral L3MBTL-associated ortholog. The occurrence of RYBP

and L3MBTL-associated proteins in these early branching lineages suggests that ncPRC1 defining

subunits, as we know them in animals, originated before cPRC1 accessory subunits.

Figure 6. Phylogenetic history and domain organization of RYBP and SAM-domain proteins. (A) Domain
organizations of the identified RYBP orthologs in the relatives of animals and fungi. (B) Schematic
*representation of the two clusters in the SAM domain phylogeny **(SI Fig.S1F)** and the full-length domain*
annotations of the identified proteins.

Discussion

Here, we present a systematic characterization of the Polycomb system throughout the

eukaryotic tree of life. Phylogenetic and structural analysis of 178 eukaryotes representing all

known major eukaryotic groups revealed broad conservation of both PRC1 and PRC2. Our results

imply that PRC1 and PRC2 were both present in LECA, but that their subsequent evolution is

uncoupled. Sensitive analyses allowed us to identify highly diverged orthologs, which previously

escaped detection. Contrary to the general believe that fungi have lost PRC1, we uncovered PRC1

orthologs in multiple deep branching fungal species. Furthermore, we identified orthologs that had

previously gone undetected because of their significant sequence divergence, such as EED in

ciliates and SUZ12 orthologs in C*ryptococcus neoformans* and *Caenorhabditis elegans* (Snel et al.

2022). We expect that increasingly sensitive techniques will uncover additional species harboring

divergent Polycomb orthologs or other, currently unknown, subunits. Thus, the evolutionary

conservation and flexible development of the Polycomb system might be even more prevalent than

presented here. Finally, we identified orthologs of ncPRC1-signature domains, but not cPRC1-
specific domains, in unicellular relatives of animals and fungi. Based on the distant homology
between the C-terminal domains of RYBP and CBX, we propose that these subunits might be
switching paralogs. Moreover, we conclude that the origin of ncPRC1 predates that of cPRC1,
strengthening earlier hypotheses that ncPRC1 accessory subunits originated before cPRC1
(Schuettengruber et al. 2017).

Ever since their discovery, the relationship between PRC1 and PRC2 has been debated.
The binding of CBX subunits of cPRC1 to H3K27me3, and the recognition of H2AK119ub1 by
PRC2.2-associated factors JARID2 or AEBP2, has inspired multiple hierarchical or interdependent
recruitment models (Baile et al. 2022; Kang et al. 2022; Kim and Kingston 2022; Piunti and
Shilatifard 2021; Schuettengruber et al. 2017; Loubiere et al. 2019; Blackledge et al. 2020;
Tamburri et al. 2020). However, in spite of cooperative cross-talk, *in vivo* studies showed that there
is no crucial dependency between PRC1 and PRC2 that is essential for repression of canonical
Polycomb targets (Bonnet et al. 2022; Cohen et al. 2021; Leeb et al. 2010; Pengelly et al. 2015;
Sijm et al. 2022; Zepeda-Martinez et al. 2020). Rather, PRC1 and PRC2 function in largely
redundant pathways, thereby ensuring robust gene silencing. Our analysis showed that a
substantial portion (~42%) of organisms that contain (part of) the Polycomb system use only PRC1
or only PRC2. These results provide compelling evolutionary support for the predominantly
independent function of PRC1 and PRC2.

Figure 7. ncPRC1 evolved prior to cPRC1. The ancestor of animals and fungi most likely already had a distinct
ncPRC1 functionality, while cPRC1 originated later in the root of animals. The upper left panel presents a
schematic overview of the identified L3MBTL-associated ortholog. Shown is furthermore an evolutionary
scenario of the origin and evolution of ncPRC1 and cPRC1 accessory subunits. Our scenario suggests that
RYBP was lost in fungi and Filasterea, and retained in Nuclearidae, Choanoflagellata and animals, while CBX
arose via a gene duplication of RYBP at the root of animals, after which the zf-RanBP got replaced by a
chromo-domain. SAM domain containing accessory subunits of PRC1 likely originated from diversification of
the ancestral L3MBTL-associated ortholog, after duplications.

While the central function of H3K27me3 for Polycomb repression in animals is unequivocal

(Sankar et al. 2022; Pengelly et al. 2015), the role of H2Aub1 turned-out to be more enigmatic.

During mouse and *Drosophila* development, H2Aub1 contributes, but is not absolutely required for

the silencing of Polycomb targets (Bonnet et al. 2022; Boyle et al. 2020; Illingworth et al. 2015;

Pengelly et al. 2015; Tsuboi et al. 2018). Rather, it appears that higher order chromatin organization

by cPRC1, which is independent of H2Aub1, is crucial for Polycomb function in animals (Bonnet et

al. 2022; Illingworth et al. 2015; Pengelly et al. 2015; Tsuboi et al. 2018). Supporting this notion,

phylogenetic profiling of histone modifications revealed broad conservation of H3K27me3, while

H2Aub1 is mainly restricted to animals and plants (Grau-Bové et al. 2022). Thus, while PRC1 is

highly conserved and can be traced back to LECA, H2Aub1 is not. This implies that the primary

function of PRC1 does not depend on H2Aub1. Collectively, these observations argue that

biochemical cross-talk between PRC1 and PRC2 is a secondary development during the evolution

of animals.

In agreement with our phylogenetic analysis, functional studies in organisms other than

animals revealed that the Polycomb system is highly plastic and employs diverse mechanisms of

action. The Polycomb system in plants is the result of an independent evolutionary trajectory,

yielding unique PRCs, which are absent in animals. E.g., plants lack cPRC1 but contain a wide

variety of plant-specific PRC1 and PRC2 accessory subunits. One of the plant-specific PRC2-

associated proteins, like-heterochromatin protein 1 (LHP1) contains a chromo domain and chromo-

shadow domain that bind to H3K27me3 (Libault et al. 2005; Gaudin et al. 2001). Thus, LHP1 might

mediate a positive feedback loop that stimulates the formation of H3K27me3 domains (Baile et al.

2022). The EZH ortholog in ciliates, Ezl1, is a dual-specificity methyltrasferase that mediates both

H3K27me3 and H3K9me3 (Frapporti et al. 2019). Ezl1 binds a PRC2 core associated with the RNA

interference effector Ptiwi09, and RING-finger proteins Rnf1 and Rnf2 (Wang et al. 2022; Xu et al.

2021; Miró-Pina et al. 2022). Thus, the Ezl1 complex appears to resemble a combination of PRC2

and PRC1. However, Rnf1 and Rnf2 are not orthologous to the canonical RING1 proteins found in

PRC1 and are not required for H2Aub1, but they do contribute to histone methylation by the Ezl1

complex. Ptiwi09 mediates small-RNA-guided recruitment of the Ezl1-PRC2 complex to

transposons, where it deposits H3K27me3 and H3K9me3, triggering transposon elimination (Wang

et al. 2022; Xu et al. 2021). Examples of adaptation in the absence of PRC1 **(Fig.3B)** are provided

by the filamentous fungus *Neurospora crassa* and the yeast *Cryptococcus neoformans*. In

Neurospora crassa, the BAH- and PHD-domain protein EPR-1 acts as a H3K27me2/3 reader.

EPR1 is a crucial protein involved in H3K27me3-directed silencing and mediates the formation of

nuclear foci resembling Polycomb bodies (Wiles et al. 2020). The *Cryptococcus neoformans* PRC2-

associated chromo-domain protein Ccc1 recognizes H3K27me3, which is crucial for its localization

to repressive subtelomeric domains (Dumesic et al. 2015). Thus, in the absence of PRC1, other

proteins recognize H3K27me3 to effectuate transcriptional silencing. Finally, it should be noted that

some Polycomb factors have been repurposed for additional, chromatin-independent functions.

E.g., *Drosophila* PSC uses a diptera-specific domain to bind and regulate cyclin B, thereby

controlling cell cycle progression **(Fig.S5)** (Mohd-Sarip et al. 2012). Mammalian SCML2B regulates

the G1/S checkpoint by binding and modulating the CDK/Cyclin/p21 complex (Lecona et al. 2013).

These examples from diverse eukaryotes emphasize the remarkable evolutionary flexibility of the

Polycomb system.

461   What was the original function of the Polycomb system? There is accumulating evidence

that the primary purpose of PRC2 in unicellular eukaryotes is the silencing of transposable

elements to protect genome integrity (Shaver et al. 2010; Xu et al. 2021; Montgomery et al. 2020;

Hisanaga et al. 2022). Similarly, in addition to gene control, plant PcG proteins ensure

transcriptional repression of repetitive genomic elements and transposable elements (Fuchs et al.

2008; Montgomery et al. 2020; Shaver et al. 2010; Widiez et al. 2014; Deleris et al. 2012; Moreno-

Romero et al. 2016; Schubert 2019). Collectively, these observations give rise to the hypothesis

that gene-selective transcriptional repression by the Polycomb system evolved from an initial

function in silencing of transposal elements (Déléris et al. 2021). Additional experimental studies

will be necessary to establish the ancestral functions of PRC1 and PRC2, and delineate how these

evolved towards their role in gene regulation in animals and plants. The findings presented here

provide unique insights into the evolutionary development of the Polycomb system and its function

in genome regulation. Additionally, it provides a framework for experimental analysis of the

Polycomb system in diverse eukaryotic organisms. We anticipate that these future studies will shed

light on the connection between genome evolution and functional adaptation of PRC1 and PRC2.

Materials and Methods

Proteome database

479   To study the occurrences of PRC1 and PRC2 genes across the eukaryotic tree of life, a

reference dataset of predicted eukaryote proteomes comprising 178 diverse eukaryotic species

was assembled. A detailed description of the included species and how the data was collected is

described in **SI Data and Methods and External Dataset S1**. Briefly, species were selected based

on multiple criteria; i) we attempted to include representatives of all major eukaryotic groups

encompassing most currently known eukaryotic diversity; ii) we weighed the species against

genome quality statistics (BUSCO); if multiple proteomes or different representatives of a single

species were available, the most complete one was selected; and iii), in clades with many closely

related, high quality genome assemblies, we selected important (model) organisms.

Homolog detection and phylogenetic inferences of core subunits PRC1 and PRC2

To study the evolutionary history of the PRC1 and PRC2 subunits, a general homolog

detection and phylogenetic inference method was initially applied and iterated upon if needed.

Briefly, eukaryotic homologs were identified using pairwise searches with BLASTP v.2.12.0+

(Altschul et al. 1990) against our local proteome database with full length *Homo sapiens* sequences

as queries (**SI External Dataset S6**). At most 500 significant hits (default: 1e-5) were aligned using

MAFFT v.7.490 (settings: genafpair, maxiterate = 1000) and processed with trimalAl v1.4.rev15 [gt

= 0.1]) (Capella-Gutierrez et al. 2009; Katoh and Standley 2013) **(SI External Dataset S7)**.

Phylogenetic trees were inferred with IQ-TREE v.2.1.4 (ModelFinder, ultrafast bootstrap = 1,000)

(Nguyen et al. 2015), and subsequently visualized and annotated using iTOL **(SI Fig.S1A-H,**

External Dataset S7) (Letunic and Bork 2021). To manually delineate orthologous groups from the

phylogenetic trees, all sequences in the trees were annotated with independent function predictions

(eggnog) and predicted domains based on HMM searches with the 'hmmsearch' tool from the

HMMER package (http://hmmer.org/, HMMER 3.3.2), with profiles from Pfam (v.35) (Huerta-Cepas

et al. 2019; Mistry et al. 2021). Delineation of the orthologous group was based on i) support values

(>90); ii) consistency of domain annotations; and iii) consistency of independent function

predictions. If all three sufficed, all sequences in the group were considered to constitute an

orthologous group.

In some cases, interpretation of the phylogenetic trees was hampered by highly diverged

sequences, incomplete genomes, low support values, or low sensitivity of Pfam profiles (v.35) that

were unable to detect protein domains. Subsequent manual inspection of homologues by

AlphaFold2 structures through ColabFold and multiple sequence alignments revealed the likely

presence of known, yet highly diverged, proteins domains (Varadi et al. 2022; Mirdita et al. 2022;

Jumper et al. 2021). In these cases, custom profiles for domains were created with the 'hmmbuild'

command from the HMMER package (http://hmmer.org/, HMMER 3.3.2) **(SI External Dataset S2)**.

These custom profiles were used to annotate sequences included in the respective phylogenetic

trees, which enabled the detection of proteins domains that were previously reported to be absent,

and therefore supported our orthologous inference. For some analyses, we also performed HMM

searches with full length profiles. The specific analyses for these subunits, and supporting data,

are described in **SI Data and Methods and External Dataset S4**. Fasta sequence files of our

orthologous groups are provided in **SI External Dataset S8**. AlphaFold2 predicted structures used

in this study are provided in **SI External Dataset S5**.

Structural analysis of CBX and RYBP

To detect structural similarity between the C-terminal regions of CBX and RYBP, the

pdb_selres.py script from pdb-tools (https://wenmr.science.uu.nl/pdbtools/) was used to cut out the

ß-sheet motifs. Accordingly, we used TMalign tool (https://zhanggroup.org/TM-align/) to perform a

structural alignment (Zhang 2005) **(SI External Dataset S3)**. Additionally, a single blast was

performed   with   the   ß-sheet   motifs   using   the   MPI   bioinformatics   toolkit

(https://toolkit.tuebingen.mpg.de/) and the MSA was used as a query to perform a HH search

(https://toolkit.tuebingen.mpg.de/) against the Euk_Homo_sapiens_04_Jul_2017 proteome, with

PDB_mmCIF70_12_Aug,   Pfam-A_v35,   and   SMART_v6.0   as   structural/domain   databases

(Hildebrand et al. 2009; Zimmermann et al. 2018; Gabler et al. 2020) **(SI External Dataset S3)**.

Correlation plot quantification and visualization

To study whether PRC1 and PRC2 co-evolved, a correlation plot of the phylogenetic

profiles was created. For this correlation plot, putative orthologs were excluded, i.e. they did not

contribute to the presences nor the absences. Phylogenetic profiles were created by setting a '1'

for presence and '0' for absence. To then evaluate the co-occurrence of PRC1 and PRC2, the

similarity of their phylogenetic profiles was quantified by using the Pearson correlation coefficient

which for binary data is equivalent to the phi coefficient. Genes were clustered based on their

phylogenetic profiles using average linkage and the resulting heatmap was visualized with ggplot

in R version 4.1.2 (Wickham 2009).

Competing Interest Statement: The authors declare no conflict of interest.

Acknowledgments

We thank Eelco Tromer, Jolien van Hooff and Carlos Santana-Molina for advice in compiling the

eukaryotic proteome database. We thank Carlos Santana-Molina for sharing scripts to remove

splice variants from the proteome database. This work was supported by the Netherlands

Organisation for Scientific Research (NWO-Vici 016.160.638, to B.S.).

Author Contributions: Author contributions: B.d.P., M.F.S., P.V., and B.S. designed research;

B.d.P. performed research; B.d.P., analyzed data; M.W.D.R., developed the methods for, and

created, the proteome database; and B.d.P, M.W.D.R., M.F.S., P.V., and B.S. wrote the paper.

All authors approved the final version of the manuscript.

Data Availability

The data supporting the conclusions made in this article are available via its online Supplementary

information, or deposited at the figshare depository as described in the Supplementary information.

Scripts to remove splice variants from the proteome database are available upon reasonable

request to the corresponding author.

References

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. *J*
Mol Biol **215**: 403–410.

Baile F, Gómez-Zambrano Á, Calonje M. 2022. Roles of Polycomb complexes in regulating gene
expression and chromatin structure in plants. *Plant Commun* **3**: 100267.
https://www.sciencedirect.com/science/article/pii/S2590346221001814.

Barbour H, Daou S, Hendzel M, Affar EB. 2020. Polycomb group-mediated histone H2A
monoubiquitination in epigenome regulation and nuclear processes. *Nat Commun* **11**:
5947.

Berke L, Snel B. 2015. The plant Polycomb repressive complex 1 (PRC1) existed in the ancestor of
seed plants and has a complex duplication history. *BMC Evol Biol* **15**: 44.

Bieluszewski T, Xiao J, Yang Y, Wagner D. 2021. PRC2 activity, recruitment, and silencing: a
comparative perspective. *Trends Plant Sci* **26**: 1186–1198.

Blackledge NP, Fursova NA, Kelley JR, Huseyin MK, Feldmann A, Klose RJ. 2020. PRC1 Catalytic
Activity Is Central to Polycomb System Function. *Mol Cell* **77**: 857-874.e9.

Bonnet J, Boichenko I, Kalb R, le Jeune M, Maltseva S, Pieropan M, Finkl K, Fierz B, Müller J.
2022. PR-DUB preserves Polycomb repression by preventing excessive accumulation of
H2Aub1, an antagonist of chromatin compaction. *Genes Dev* **36**: 1046–1061.

Boyle S, Flyamer IM, Williamson I, Sengupta D, Bickmore WA, Illingworth RS. 2020. A central role
for canonical PRC1 in shaping the 3D nuclear landscape. *Genes Dev* **34**: 931–949.

Bracken AP, Brien GL, Verrijzer CP. 2019. Dangerous liaisons: interplay between SWI/SNF, NuRD,
and Polycomb in chromatin regulation and cancer. *Genes Dev* **33**: 936–959.

Capella-Gutierrez S, Silla-Martinez JM, Gabaldon T. 2009. trimAl: a tool for automated alignment
trimming in large-scale phylogenetic analyses. *Bioinformatics* **25**: 1972–1973.

Chen S, Jiao L, Shubbar M, Yang X, Liu X. 2018. Unique Structural Platforms of Suz12 Dictate
Distinct Classes of PRC2 for Chromatin Binding. *Mol Cell* **69**: 840-852.e5.

Chiacchiera F, Rossi A, Jammula S, Piunti A, Scelfo A, Ordóñez-Morán P, Huelsken J, Koseki H,
Pasini D. 2016. Polycomb Complex PRC1 Preserves Intestinal Stem Cell Identity by
Sustaining Wnt/β-Catenin Transcriptional Activity. *Cell Stem Cell* **18**: 91–103.

Cohen I, Bar C, Liu H, Valdes VJ, Zhao D, Galbo PM, Silva JM, Koseki H, Zheng D, Ezhkova E. 2021.
Polycomb complexes redundantly maintain epidermal stem cell identity during
development. *Genes Dev* **35**: 354–366.

Déléris A, Berger F, Duharcourt S. 2021. Role of Polycomb in the control of transposable
elements. *Trends Genet* **37**: 882–889.

Deleris A, Stroud H, Bernatavichute Y, Johnson E, Klein G, Schubert D, Jacobsen SE. 2012. Loss of
the DNA Methyltransferase MET1 Induces H3K9 Hypermethylation at PcG Target Genes
and Redistribution of H3K27 Trimethylation to Transposons in Arabidopsis thaliana. *PLoS*
Genet **8**: e1003062.

Dumesic PA, Homer CM, Moresco JJ, Pack LR, Shanle EK, Coyle SM, Strahl BD, Fujimori DG, Yates
JR, Madhani HD. 2015. Product Binding Enforces the Genomic Specificity of a Yeast
Polycomb Repressive Complex. *Cell* **160**: 204–218.

Frapporti A, Miró Pina C, Arnaiz O, Holoch D, Kawaguchi T, Humbert A, Eleftheriou E, Lombard B,
Loew D, Sperling L, et al. 2019. The Polycomb protein Ezl1 mediates H3K9 and H3K27
methylation to repress transposable elements in Paramecium. *Nat Commun* **10**: 2710.

Fuchs J, Jovtchev G, Schubert I. 2008. The chromosomal distribution of histone methylation
marks in gymnosperms differs from that of angiosperms. *Chromosome Res* **16**: 891–8.

Gabler F, Nam S, Till S, Mirdita M, Steinegger M, Söding J, Lupas AN, Alva V. 2020. Protein
Sequence Analysis Using the MPI Bioinformatics Toolkit. *Curr Protoc Bioinformatics* **72**.

Gahan JM, Rentzsch F, Schnitzler CE. 2020. The genetic basis for PRC1 complex diversity
emerged early in animal evolution. *Proceedings of the National Academy of Sciences* **117**:
22880–22889.

Gao Z, Zhang J, Bonasio R, Strino F, Sawai A, Parisi F, Kluger Y, Reinberg D. 2012. PCGF homologs,
CBX proteins, and RYBP define functionally distinct PRC1 family complexes. *Mol Cell* **45**:
344–56.

Gaudin V, Libault M, Pouteau S, Juul T, Zhao G, Lefebvre D, Grandjean O. 2001. Mutations in LIKE
HETEROCHROMATIN PROTEIN 1 affect flowering time and plant architecture in
Arabidopsis. *Development* **128**: 4847–58.

Glancy E, Ciferri C, Bracken AP. 2021. Structural basis for PRC2 engagement with chromatin. *Curr*
Opin Struct Biol **67**: 135–144.

Grau-Bové X, Navarrete C, Chiva C, Pribasnig T, Antó M, Torruella G, Galindo LJ, Lang BF, Moreira
D, López-Garcia P, et al. 2022. A phylogenetic and proteomic reconstruction of eukaryotic
chromatin evolution. *Nat Ecol Evol* **6**: 1007–1023.

Hennig L, Derkacheva M. 2009. Diversity of Polycomb group complexes in plants: same rules,
different players? *Trends Genet* **25**: 414–23.

Hildebrand A, Remmert M, Biegert A, Söding J. 2009. Fast and accurate automatic structure
prediction with HHpred. *Proteins: Structure, Function, and Bioinformatics* **77**: 128–132.

Hisanaga T, Romani F, Wu S, Kowar T, Lintermann R, Jamge B, Montgomery SA, Axelsson E,
Dierschke T, Bowman JL, et al. 2022. Transposons repressed by H3K27me3 were co-opted
as cis-regulatory elements of H3K27me3 controlled protein coding genes during evolution
of plants. *bioRxiv* 2022.10.24.513474.
http://biorxiv.org/content/early/2022/10/24/2022.10.24.513474.abstract.

Huang Y, Chen D-H, Liu B-Y, Shen W-H, Ruan Y. 2017. Conservation and diversification of
polycomb repressive complex 2 (PRC2) proteins in the green lineage. *Brief Funct Genomics*
16: 106–119.

Huang Y, Jiang L, Liu B-Y, Tan C-F, Chen D-H, Shen W-H, Ruan Y. 2019. Evolution and
conservation of polycomb repressive complex 1 core components and putative associated
factors in the green lineage. *BMC Genomics* **20**: 533.

Huerta-Cepas J, Szklarczyk D, Heller D, Hernández-Plaza A, Forslund SK, Cook H, Mende DR,
Letunic I, Rattei T, Jensen LJ, et al. 2019. eggNOG 5.0: a hierarchical, functionally and
phylogenetically annotated orthology resource based on 5090 organisms and 2502 viruses.
Nucleic Acids Res **47**: D309–D314.

Illingworth RS, Moffat M, Mann AR, Read D, Hunter CJ, Pradeepa MM, Adams IR, Bickmore WA.
2015. The E3 ubiquitin ligase activity of RING1B is not essential for early mouse
development. *Genes Dev* **29**: 1897–1902.

Isono K, Endo TA, Ku M, Yamada D, Suzuki R, Sharif J, Ishikura T, Toyoda T, Bernstein BE, Koseki
H. 2013. SAM Domain Polymerization Links Subnuclear Clustering of PRC1 to Gene
Silencing. *Dev Cell* **26**: 565–577.

Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K, Bates R,
Žídek A, Potapenko A, et al. 2021. Highly accurate protein structure prediction with
AlphaFold. *Nature* **596**: 583–589.

Kang H, Cabrera JR, Zee BM, Kang HA, Jobe JM, Hegarty MB, Barry AE, Glotov A, Schwartz YB,
Kuroda MI. 2022. Variant Polycomb complexes in Drosophila consistent with ancient
functional diversity. *Sci Adv* **8**: eadd0103.

Kasinath V, Faini M, Poepsel S, Reif D, Feng XA, Stjepanovic G, Aebersold R, Nogales E. 2018.
Structures of human PRC2 with its cofactors AEBP2 and JARID2. *Science (1979)* **359**: 940–
944.

Kassis JA, Kennison JA, Tamkun JW. 2017. Polycomb and Trithorax Group Genes in Drosophila.
Genetics **206**: 1699–1725.

Katoh K, Standley DM. 2013. MAFFT Multiple Sequence Alignment Software Version 7:
Improvements in Performance and Usability. *Mol Biol Evol* **30**: 772–780.

Kim JJ, Kingston RE. 2022. Context-specific Polycomb mechanisms in development. *Nat Rev
Genet* **23**: 680–695. https://doi.org/10.1038/s41576-022-00499-0.

Lagarou A, Mohd-Sarip A, Moshkin YM, Chalkley GE, Bezstarosti K, Demmers JAA, Verrijzer CP.
2008. dKDM2 couples histone H2A ubiquitylation to histone H3 demethylation during
Polycomb group silencing. *Genes Dev* **22**: 2799–2810.

Laugesen A, Højfeldt JW, Helin K. 2019. Molecular Mechanisms Directing PRC2 Recruitment and
H3K27 Methylation. *Mol Cell* **74**: 8–18.

Lecona E, Rojas LA, Bonasio R, Johnston A, Fernández-Capetillo O, Reinberg D. 2013. Polycomb
Protein SCML2 Regulates the Cell Cycle by Binding and Modulating CDK/CYCLIN/p21
Complexes. *PLoS Biol* **11**: e1001737.

Leeb M, Pasini D, Novatchkova M, Jaritz M, Helin K, Wutz A. 2010. Polycomb complexes act
redundantly to repress genomic repeats and genes. *Genes Dev* **24**: 265–276.

Letunic I, Bork P. 2021. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree
display and annotation. *Nucleic Acids Res* **49**: W293–W296.

Libault M, Tessadori F, Germann S, Snijder B, Fransz P, Gaudin V. 2005. The Arabidopsis LHP1
protein is a component of euchromatin. *Planta* **222**: 910–25.

Loubiere V, Martinez A-M, Cavalli G. 2019. Cell Fate and Developmental Regulation Dynamics by
Polycomb Proteins and 3D Genome Architecture. *BioEssays* **41**: 1800222.

Mirdita M, Schütze K, Moriwaki Y, Heo L, Ovchinnikov S, Steinegger M. 2022. ColabFold: making
protein folding accessible to all. *Nat Methods* **19**: 679–682.

Miró-Pina C, Charmant O, Kawaguchi T, Holoch D, Michaud A, Cohen I, Humbert A, Jaszczyszyn
Y, Chevreux G, del Maestro L, et al. 2022. Paramecium Polycomb repressive complex 2
physically interacts with the small RNA-binding PIWI protein to repress transposable
elements. *Dev Cell* **57**: 1037-1052.e8.

Mistry J, Chuguransky S, Williams L, Qureshi M, Salazar GA, Sonnhammer ELL, Tosatto SCE,
Paladin L, Raj S, Richardson LJ, et al. 2021. Pfam: The protein families database in 2021.
Nucleic Acids Res **49**: D412–D419.

Mohd-Sarip A, Lagarou A, Doyen CM, van der Knaap JA, Aslan Ü, Bezstarosti K, Yassin Y, Brock
HW, Demmers JAA, Verrijzer CP. 2012. Transcription-independent function of Polycomb
group protein PSC in cell cycle control. *Science* **336**: 744–7.

Montgomery SA, Tanizawa Y, Galik B, Wang N, Ito T, Mochizuki T, Akimcheva S, Bowman JL,
Cognat V, Maréchal-Drouard L, et al. 2020. Chromatin Organization in Early Land Plants
Reveals an Ancestral Association between H3K27me3, Transposons, and Constitutive
Heterochromatin. *Current Biology* **30**: 573-588.e7.

Moreno-Romero J, Jiang H, Santos-González J, Köhler C. 2016. Parental epigenetic asymmetry of
<scp>PRC</scp> 2-mediated histone modifications in the *Arabidopsis* endosperm. *EMBO J*
35: 1298–1311.

Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. 2015. IQ-TREE: A Fast and Effective
Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. *Mol Biol Evol* **32**:
268–274.

Ori A, Iskar M, Buczak K, Kastritis P, Parca L, Andrés-Pons A, Singer S, Bork P, Beck M. 2016.
Spatiotemporal variation of mammalian protein complex stoichiometries. *Genome Biol* **17**:
47.

Pengelly AR, Copur Ö, Jäckle H, Herzig A, Müller J. 2013. A Histone Mutant Reproduces the
Phenotype Caused by Loss of Histone-Modifying Factor Polycomb. *Science (1979)* **339**:
698–699.

Pengelly AR, Kalb R, Finkl K, Müller J. 2015. Transcriptional repression by PRC1 in the absence of
H2A monoubiquitylation. *Genes Dev* **29**: 1487–1492.

Piunti A, Shilatifard A. 2021. The roles of Polycomb repressive complexes in mammalian
development and cancer. *Nat Rev Mol Cell Biol* **22**: 326–345.
https://doi.org/10.1038/s41580-021-00341-1.

Rossi A, Ferrari KJ, Piunti A, Jammula S, Chiacchiera F, Mazzarella L, Scelfo A, Pelicci PG, Pasini D.
2016. Maintenance of leukemic cell identity by the activity of the Polycomb complex PRC1
in mice. *Sci Adv* **2**: e1600972.

Sankar A, Mohammad F, Sundaramurthy AK, Wang H, Lerdrup M, Tatar T, Helin K. 2022. Histone
editing elucidates the functional roles of H3K27 methylation and acetylation in mammals.
Nat Genet **54**: 754–760.

Schubert D. 2019. Evolution of Polycomb-group function in the green lineage. *F1000Res* **8**: 268.

Schuettengruber B, Bourbon H-M, di Croce L, Cavalli G. 2017. Genome Regulation by Polycomb
and Trithorax: 70 Years and Counting. *Cell* **171**: 34–57.

Seif E, Kang JJ, Sasseville C, Senkovich O, Kaltashov A, Boulier EL, Kapur I, Kim CA, Francis NJ.
2020. Phase separation by the polyhomeotic sterile alpha motif compartmentalizes
Polycomb Group proteins and enhances their activity. *Nat Commun* **11**: 5609.

Sharaf A, Vijayanathan M, Oborník M, Mozgová I. 2022. Phylogenetic profiling resolves early
emergence of PRC2 and illuminates its functional core. *Life Sci Alliance* **5**: e202101271.

Shaver S, Casas-Mollano JA, Cerny RL, Cerutti H. 2010. Origin of the polycomb repressive
complex 2 and gene silencing by an E(z) homolog in the unicellular alga Chlamydomonas.
Epigenetics **5**: 301–12.

Sijm A, Atlasi Y, van der Knaap JA, Wolf van der Meer J, Chalkley GE, Bezstarosti K, Dekkers
DHW, Doff WAS, Ozgur Z, van IJcken WFJ, et al. 2022. USP7 regulates the ncPRC1 Polycomb
axis to stimulate genomic H2AK119ub1 deposition uncoupled from H3K27me3. *Sci Adv* **8**.

Simon JA, Kingston RE. 2009. Mechanisms of Polycomb gene silencing: knowns and unknowns.
Nat Rev Mol Cell Biol **10**: 697–708.

Snel B, van den Heuvel S, Seidl MF. 2022. Caenorhabditis elegans MES-3 is a highly divergent
ortholog of the canonical PRC2 component SUZ12. *iScience* **25**: 104633.

Suganuma T, Pattenden SG, Workman JL. 2008. Diverse functions of WD40 repeat proteins in
histone recognition: Figure 1. *Genes Dev* **22**: 1265–1268.

Szklarczyk R, Huynen MA, Snel B. 2008. Complex fate of paralogs. *BMC Evol Biol* **8**: 337.

Tamburri S, Conway E, Pasini D. 2022. Polycomb-dependent histone H2A ubiquitination links
developmental disorders with cancer. *Trends in Genetics* **38**: 333–352.

Tamburri S, Lavarone E, Fernández-Pérez D, Conway E, Zanotti M, Manganaro D, Pasini D. 2020.
Histone H2AK119 Mono-Ubiquitination Is Essential for Polycomb-Mediated Transcriptional
Repression. *Mol Cell* **77**: 840-856.e5.

Tsuboi M, Kishi Y, Yokozeki W, Koseki H, Hirabayashi Y, Gotoh Y. 2018. Ubiquitination-
Independent Repression of PRC1 Targets during Neuronal Fate Restriction in the
Developing Mouse Neocortex. *Dev Cell* **47**: 758-772.e5.

Varadi M, Anyango S, Deshpande M, Nair S, Natassia C, Yordanova G, Yuan D, Stroe O, Wood G,
Laydon A, et al. 2022. AlphaFold Protein Structure Database: massively expanding the
structural coverage of protein-sequence space with high-accuracy models. *Nucleic Acids*
Res **50**: D439–D444.

Wang C, Solberg T, Maurer-Alcalá XX, Swart EC, Gao F, Nowacki M. 2022. A small RNA-guided
PRC2 complex eliminates DNA as an extreme form of transposon silencing. *Cell Rep* **40**:
111263.

Wickham H. 2009. *ggplot2*. Springer New York, New York, NY.

Widiez T, Symeonidi A, Luo C, Lam E, Lawton M, Rensing SA. 2014. The chromatin landscape of
the moss *Physcomitrella patens* and its dynamics during development and drought stress.
The Plant Journal **79**: 67–81.

Wiles ET, McNaught KJ, Kaur G, Selker JML, Ormsby T, Aravind L, Selker EU. 2020. Evolutionarily
ancient BAH–PHD protein mediates Polycomb silencing. *Proceedings of the National*
Academy of Sciences **117**: 11614–11623.

Xu J, Zhao X, Mao F, Basrur V, Ueberheide B, Chait BT, Allis CD, Taverna SD, Gao S, Wang W, et
al. 2021. A Polycomb repressive complex is required for RNAi-mediated heterochromatin
formation and dynamic distribution of nuclear bodies. *Nucleic Acids Res* **49**: 5407–5425.

Yu J-R, Lee C-H, Oksuz O, Stafford JM, Reinberg D. 2019. PRC2 is high maintenance. *Genes Dev*
33: 903–935.

Zepeda-Martinez JA, Pribitzer C, Wang J, Bsteh D, Golumbeanu S, Zhao Q, Burkard TR, Reichholf
B, Rhie SK, Jude J, et al. 2020. Parallel PRC2/cPRC1 and vPRC1 pathways silence lineage-
specific genes and maintain self-renewal in mouse embryonic stem cells. *Sci Adv* **6**.

Zhang Y. 2005. TM-align: a protein structure alignment algorithm based on the TM-score.
Nucleic Acids Res **33**: 2302–2309.

Zimmermann L, Stephens A, Nam S-Z, Rau D, Kübler J, Lozajic M, Gabler F, Söding J, Lupas AN,
Alva V. 2018. A Completely Reimplemented MPI Bioinformatics Toolkit with a New HHpred
Server at its Core. *J Mol Biol* **430**: 2237–2243.