Agent-Based Modelling shows that Operation of Fisher's Principle does not explain

Zero Heritability in Human Offspring Sex Ratio

Kaitlyn T. Harper[1*], Morgan J. Sidari[1*], and Brendan P. Zietsch[1]

*Authors contributed equally

[1]Centre for Psychology and Evolution, School of Psychology, University of Queensland, Brisbane, Australia

Author Note

Kaitlyn T. Harper ⓘ https://orcid.org/0000-0001-9597-872X

Morgan J. Sidari ⓘ https://orcid.org/0000-0001-5315-4546

Brendan P. Zietsch ⓘ https://orcid.org/0000-0003-0274-6140

We have no conflicts of interest to disclose.

Corresponding author email: kaitlyn.harper@uq.net.au

Data availability statement: All R code and results are available at https://osf.io/cbpxr/?view_only=5c437beb09534d4e8b701045e16faf01.

24   **Abstract**

The evolutionary basis of the ratio of male to female births has been a focus of research for

over a century. Fisher's principle suggests that offspring sex ratios remain roughly equal due

to negative frequency-dependent balancing selection: individuals with alleles that produce the

rarer sex have greater fitness, increasing these alleles' frequency until the sex ratio is

balanced. However, recent analysis of Swedish births since 1932 reveals that offspring sex

ratio is not heritable, challenging Fisher's principle's application to humans. Here, we test the

hypothesis that Fisher's principle uniquely depletes offspring sex ratio heritability at

equilibrium. We used agent-based modelling to simulate Fisher's principle over 5000

generations and found that Fisher's principle at equilibrium does not eliminate genetic

variation. These findings, along with evidence of non-heritable offspring sex ratios in

humans, suggest Fisher's principle is not a suitable explanatory framework for human

offspring sex ratio.

Key words: offspring sex ratio, Fisher's principle, negative frequency-dependent balancing

selection, agent-based model, genetic variation, sex allocation

Agent-based Modelling shows that Operation of Fisher's Principle does not explain

41   **Zero Heritability in Human Offspring Sex Ratio**

Offspring sex ratio – a parent's ratio of male to female births – has garnered interest

and debate for more than a century. It is also the subject of Fisher's principle (Fisher, 1930),

which traces back to Darwin (1871) and Dusing (cited in Edwards, 1998) and is sometimes

referred to as Dusing-Fisher theory. Fisher's principle states that in sexually reproducing

species, the offspring sex ratio will stabilise around 1:1 due to sexual selection favouring

whichever sex is less common. This process is an example of frequency-dependent balancing

selection: all else being equal, individuals genetically predisposed to producing the scarce sex

will tend to have greater fitness (more grandchildren), so those individuals' alleles will

increase in frequency until the population sex ratio rebalances and the advantage of the allele

disappears. According to the theory, this process keeps the sex ratio approximately even

(under the condition that male and female offspring require equal parental investment).

Fisher's principle has been the primary framework by which human offspring sex

ratio has been understood (Hardy, 2002; West, 2009). Recently, though, using data from the

whole Swedish population born 1932 or later (3,543,243 individuals and their 4,753,269

children), Zietsch et al. (2020) showed that offspring sex ratio is not heritable in humans.

That is, genetic differences do not influence the sex ratio of individuals' offspring, and

therefore variation in offspring sex ratio cannot be inherited. Even within individuals there

was no consistency at all in the sex of subsequent offspring, suggesting that offspring sex was

determined by random Mendelian segregation of sex chromosomes. These observations are

incompatible with the operation of Fisher's principle – offspring sex ratio must be heritable to

respond to selection as Fisher described.

If Fisher's principle cannot operate in humans, how can it serve as a framework for

understanding human sex ratio? Some have argued that Fisher's principle makes no

prediction about the heritability of offspring sex ratio at equilibrium (i.e. a stable state

reached through the operation of a process) and therefore can still serve as a valid

explanatory framework (Lehtonen, 2021; Orzack & Hardy, 2021). The simplest explanation

under this view is that the operation of Fisher's principle has eliminated the genetic variation

in offspring sex ratio, since it must have been heritable for Fisher's principle to operate, but is

no longer heritable. While negative frequency-dependent selection is typically expected to

actively maintain genetic variation (Brisson, 2018; Clarke, 1979; Zietsch et al.,

2021)(Asmussen & Basnayake, 1990; Trotter & Spencer, 2007; Villanea et al., 2015),

Fisher's principle is a unique process, and it has not been established what effect the

operation of Fisher's principle has on genetic variation in offspring sex ratio. Given the

complex underlying mechanisms of offspring sex ratio evolution, which are not always

obvious, it is important to investigate this specific mechanism. Here, we used agent-based

modelling to simulate the operation of Fisher's principle and its effects on genetic variation

and heritability in offspring sex ratio.

79   **Method**

We created an agent-based model in R to test whether the operation of Fisher's

principle depletes genetic variation in offspring sex ratio (RCore, 2016; RStudio, 2022). The

model was comprised of male and female agents (2000 total), each possessing a small

genome of 20 loci which influenced their tendency to produce male or female offspring. We

enabled agents to reproduce over 5000 generations, allowing us to demonstrate Fisher's

principle and observe its long-term effects at equilibrium. We decided to use 20 loci to ensure

that offspring sex ratio was modelled as a quantitative trait (as opposed to a discrete trait with

one or two loci). Computational load was also considered when deciding on number of loci,

number of generations, and population size. In the following sections, we describe how

agents were created and the rules by which they reproduced. All code and results can be

accessed at https://osf.io/cbpxr/?view_only=5c437beb09534d4e8b701045e16faf01.

Creating Initial Generation

The initial population of 2000 agents was generated, 40% were males. Each agent had

20 loci, each with two alleles, that influenced offspring sex ratio. Initial allele values were

generated as either 0 or 1 (low variant or high variant), which occurred with a probability that

reflected the initial population sex ratio. In the reported model, the initial population sex ratio

was 40% male, so the allele values were randomised with a 0.4 probability (such that genetic

scores reflected a 40% probability of producing a male offspring). Genotype was then

calculated as *allele 1 + allele 2*, thereby giving agents genotype values of 0, 1, or 2 at each

locus. Genetic score for offspring sex ratio was calculated by summing genotypes for all loci.

In the reported model there were 20 loci influencing offspring sex ratio, so possible genetic

scores ranged between 0 and 40.

Partner Selection

The following operations were performed five times to create five 'mating rounds'.

Agents were paired randomly, with some surplus-sex agents missing out on partnering and

reproducing for that round. Mating success for the surplus sex therefore varied randomly

between agents (ranging from 0-5), a result of the number of times they were randomly

excluded from the mating round. Partners' data were then linked with the agents' data so that

values for both parents were available to determine offspring attributes.

Determining Offspring Sex

The following process emulated a genetic predisposition toward male or female

offspring with additional environmental variance, such that offspring sex was determined

80% by genes and 20% by environmental factors (to simulate a largely but not completely

heritable trait, like most biological traits). For each offspring, the parents' genetic scores were

averaged and transformed to a proportion of the highest theoretical genetic score (e.g. in the

reported model the maximum genetic score was 40, so a score of 10 would be transformed to

0.25). This proportionate genetic score was then multiplied by 0.8, and an additional random

value between 0 and 0.2 was added to represent 20% environmental influence. The resulting

score was used as the probability in a random binomial outcome where 0 = female, and 1 =

male; higher scores were more likely to produce male offspring. Outcome measures were

recorded for the current generation at this point.

Reproduction

Alleles were then inherited through independent assortment. That is, at each locus, an

offspring inherited one allele from each parent, determined randomly. The population matrix

was then truncated (i.e. all parent information removed), leaving the offspring to become the

next generation, and the process described thus far was repeated. In the reported model, each

simulation was run for 5000 generations.

Outcome Variables and Analysis

We ran the model described above 50 times and averaged the simulation results for

each generation. The outcome variables of interest were population sex ratio, genetic score,

heritability, and genetic variance. Offspring sex ratio was recorded as the proportion of male

offspring produced. Genetic score was recorded as the mean of the raw genetic scores. Given

that our data was simulated, and we had complete genome information for each agent, we

used simpler calculations of genetic variance and heritability than those normally used in the

broader quantitative genetics literature. To calculate heritability, i.e. the proportion of

phenotypic variance accounted for by genetic variance, we used the r^2 of genetic score

predicting offspring sex ratio (i.e. the proportion of the agent's offspring that were male). To

calculate genetic variance, we multiplied the total phenotypic variance (variance in offspring

sex ratio) by the heritability, giving us the variance in offspring sex ratio that is explained by

genetic scores. We used correlations to assess trends in outcome variables over generations.

For ease of presentation in the main text, we have reported our model of Fisher's principle for

1000 generations. Extended results (with 5000 generations) can be found in the

Supplementary Material.

143   **Results**

144   **Sex ratio and genetic scores**

145   To determine whether Fisher's principle operated as expected in our model, we

needed to establish whether genetic scores, and consequently sex ratio at birth, changed over

time. Results demonstrated that the genetic scores diverged based on agent gender, with male

agents having higher scores than female agents (see Figure 1A). This result was to be

expected, given that a male is more likely to have been produced by male-making alleles, and

thus will have inherited more male-making alleles themselves; likewise, females will have

inherited more female-making alleles. We ran a simulation where the initial genetic scores for

each sex reflected this natural separation between male and female genetic scores, and all

results were comparable.

154   In addition to the divergence of genetic scores by sex, the mean genetic score also

increased over time (from the initial mean genetic score which was biased towards female-

making alleles), such that alleles for having male offspring (i.e. the rarer sex) became more

prevalent (Figure 1A). The rate of this increase slowed until around generation 500 when the

process reached an equilibrium at a 1:1 predisposition for offspring sex (at the midpoint of

possible genetic scores – 20). The resulting population sex ratio also approached a 1:1

equilibrium (0.5 male) over the generations (see Figure 1B). The observed pattern of results

thus far is precisely as Fisher's principle would predict, where alleles leading to the less

common sex increased in frequency until the sex ratio reached equilibrium at 1:1, at which

point these alleles no longer conferred a benefit.

Genetic variation and heritability

To address arguments that offspring sex ratio might no longer be heritable due to the

operation of Fisher's principle (Lehtonen, 2021; Orzack & Hardy, 2021), we next sought to

test whether genetic variation and heritability were eliminated over generations.

Although heritability and genetic variance appear to be stable across generations at

first glance (see Figure 1C and Figure 1D), correlations revealed that they each subtly

decreased over time ($r = -0.36$, $p < .001$, for both analyses). Upon further exploration of the

data, this decrease appears to be due to fixation of some alleles due to drift. An allele can fix

at zero or 100% frequency due to chance fluctuations over many generations (i.e. drift), but

once fixed it cannot change frequency (without mutation); that is, allelic variation can be lost

but not replenished in our model, and so the occasional chance fixations will cause the

genetic variation to decrease slightly over time. In analyses described in the Supplementary

Material, we demonstrate that the slight decreases in heritability detected in our model of

Fisher's principle are slightly less than those observed under no selection at all (and far less

than observed under directional and stabilising selection). In summary, the operation of

Fisher's principle does not deplete genetic variation, but also does not entirely buffer the loss

of genetic variation due to drift.

Figure 1. A Genetic score, **B** population sex ratio, **C** heritability, and **D** genetic variance over 1000

generations.

Note. Grey lines represent individual results from 50 simulations. Coloured lines represent the mean across

all simulations at each generation. In Figure B, the blue line represents the mean male genetic score and the

red line represents the mean female genetic score. In Figure A and Figure B, the dotted horizontal line

represents the midpoint of offspring sex ratio (1:1)

Discussion

In the broader field of quantitative genetics, the maintenance of genetic variation

throughout various forms of selection has been extensively studied (Walsh et al., 2018). The

existing literature highlights the intricate and often unpredictable nature of genetic variation

under different evolutionary pressures. Our research provides a focused examination of

genetic variation under a specific case of negative frequency-dependent balancing selection:

Fisher's principle. Widely accepted as the foundation for explaining the 1:1 human offspring

sex ratio, Fisher's principle has been recently contradicted by findings that offspring sex ratio

is not heritable in humans (Zietsch et al., 2020). Some have argued that zero heritability at

equilibrium is not ruled out by Fisher's principle (Lehtonen, 2021; Orzack & Hardy, 2021),

yet it is uncertain how the heritability necessary for the operation of Fisher's principle could

have been eliminated. With broader literature suggesting that negative frequency-dependent

balancing selection is expected to maintain genetic variation (Asmussen & Basnayake, 1990;

Trotter & Spencer, 2007; Villanea et al., 2015), our study aimed to determine whether

Fisher's principle could be an exceptional case where a form of negative frequency-

dependent selection actively depletes genetic variation over time.

In the first generation of our model, males were the scarcer sex and the population's

alleles reflected this with a bias toward producing females. Over many generations, the sex

ratio and genetic scores of the population approached equilibrium at a 1:1 offspring sex ratio.

This pattern of results demonstrates Fisher's principle as expected, such that alleles that

produce the scarcer sex will increase until they are no longer beneficial (once a 1:1

population sex ratio is reached). At this point, there is no selection for these alleles, which is

consistent with our findings that population sex ratio and genetic scores remained stable after

this point. While our model demonstrates that Fisher's principle is operationally sound, it

does not support arguments that it could have eliminated offspring sex ratio heritability over

time. Fisher's principle, in our simulations, maintained heritability for longer than directional

and stabilising selection and slightly longer than genetic drift (no selection). Previous

modelling shows that negative frequency-dependent selection maintains genetic variation

entirely, even counteracting drift (Villanea et al., 2015), so it is unclear whether factors of

model complexity such as population size and number of alleles affected the influence of drift

in our model.

218   Regardless, Fisher's principle cannot account for the current observable state of no

heritability in offspring sex ratio while there remains abundant heritability in virtually all

variable traits that have been studied, both those under selection and no selection (Polderman

et al., 2015). Given our results, Fisher's principle would predict offspring sex ratio to be more

heritable than most traits; instead, offspring sex ratio is basically unique among variable traits

in being non-heritable. Additionally, anticipating the possible argument that offspring sex

ratio heritability may be unobservable, we note that substantial heritability in offspring sex

ratio has been detected across various species (Premoli et al., 1996; Varandas et al., 1997).

226   Our findings suggest that the operation of Fisher's principle cannot be reconciled with

227   the current non-heritable nature of offspring sex ratio, thus challenging the principle's

228   relevance for understanding human offspring sex ratio. We suggest that sex allocation theory

229   in humans needs to be revised, given the foundational status of Fisher's principle in the field,

230   as described by West (2009, p.14):

231   *'Fisher's theory of equal investment provides the basic null model for sex allocation*

232   *theory, but it is also the foundation for all subsequent theoretical developments.'*

Hardy (2002, p.2, emphases added) also acknowledges that Fisher's principle underpins

extensive subsequent research:

'Darwin, Düsing, Fisher, and Shaw and Mohler established the fundamental principle

that members of the minority sex tend to have higher fitness than members of the

majority sex. They also outlined how various ecological, demographic and genetic

variables might affect the details of sex-allocation strategies by modifying both the

*constraints and the fitness functions. **Modern sex-allocation research is devoted***

***largely to the exploration of such effects**, which connect sex ratios to many other*

aspects of the biologies of many species'

Our model is a simplified representation of the real world, and as with any model,

there are limitations. Discrete generations in our model are simpler than true mating

environments, and our genomes of 20 loci do not represent human genetic architecture.

Nevertheless, we have no reason to expect that complexities involving linkage disequilibrium

or minor allele frequencies would substantively affect our findings, especially since Fisher's

description of his principle does not implicate any of these factors. We also did not

investigate factors such as group size and environmental sex ratio fluctuations. For instance,

some might argue that in small groups, ongoing sex-specific mortality could lead to

continuous selection and faster loss of genetic variation. However, this perspective overlooks

that environmental fluctuations would not alter the genetic predisposition for offspring sex

ratio in the surviving individuals. Therefore, while these changes might temporarily affect the

sex ratio, the population would return to its natural equilibrium in the next generation. While

differential mortality between sexes during parental care might influence selection on

offspring sex ratio (Kahn et al., 2015), no specific set of circumstances has been proposed

that would result in a continuous reduction of genetic variance over time.

Our findings, combined with recent evidence that offspring sex ratio is not heritable in

humans (Zietsch et al., 2020), suggest that Fisher's principle is not an appropriate explanatory

framework for variation in the offspring sex ratio in humans. We anticipate various

arguments that Fisher's principle may have operated to equilibrium in humans, at which point

a number of mechanisms could have depleted genetic variance in offspring sex ratio, for

example an invading genetic variant coding for a 1:1 offspring sex ratio. We accept these

possibilities and encourage the development of new theoretical models, but we caution the

use of Fisher's principle as an assumed starting point for new theory, given the lack of

evidence for its operation in humans. Moreover, the inconsistency of offspring sex ratio

within individuals (Zietsch et al.) suggests a random Mendelian process determines the

offspring sex ratio in humans, although it is unclear what individual or group level

evolutionary processes could have led to this outcome. Regardless, such a mechanism would

be far removed from Fisher's principle as originally proposed. We anticipate that new

theoretical models will be necessary to reconstruct our understanding of sex ratio evolution in

humans, and we look forward to seeing progress in this field.

273   References

274   Asmussen, M. A., & Basnayake, E. (1990). Frequency-dependent selection: the high potential
275   for permanent genetic variation in the diallelic, pairwise interaction model. *Genetics*,
276   *125*(1), 215-230. https://doi.org/10.1093/genetics/125.1.215
277   Brisson, D. (2018). Negative Frequency-Dependent Selection Is Frequently Confounding
278   [Perspective]. *Frontiers in Ecology and Evolution*, *6*.
279   https://doi.org/10.3389/fevo.2018.00010
280   Clarke, B. C. (1979). The evolution of genetic diversity. *Proceedings of the Royal Society of
281   London. Series B. Biological Sciences*, *205*(1161), 453-474.
282   https://doi.org/10.1098/rspb.1979.0079
283   Darwin, C. R. (1871). *The descent of man, and selection in relation to sex*. John Murray.
284   Edwards, A. W. F. (1998). Natural selection and the sex ratio: Fisher's sources. *The American
285   Naturalist*, *151*(6), 564-569. https://doi.org/10.1086/286141
286   Fisher, R. A. (1930). *The genetical theory of natural selection*. Oxford University Press.
287   Hardy, I. (2002). *Sex ratios : concepts and research methods*. Cambridge University Press.
288   Kahn, A. T., Jennions, M. D., & Kokko, H. (2015). Sex allocation, juvenile mortality and the
289   costs imposed by offspring on parents and siblings. *Journal of Evolutionary Biology*,
290   *28*(2), 428-437.
291   Lehtonen, J. (2021). Fisher's principle remains a plausible explanation for human sex ratio
292   evolution. *Proceedings of the Royal Society B: Biological Sciences*.
293   Orzack, S. H., & Hardy, I. (2021). Does the lack of heritability of human sex ratios require a
294   rethink of sex ratio theory? No: a response to Zietsch et al. 2020. *Proceedings of the
295   Royal Society B: Biological Sciences*.
296   Polderman, T. J. C., Benyamin, B., de Leeuw, C. A., Sullivan, P. F., van Bochoven, A.,
297   Visscher, P. M., & Posthuma, D. (2015). Meta-analysis of the heritability of human
298   traits based on fifty years of twin studies [Analysis]. *Nat Genet*, *47*(7), 702-709.
299   https://doi.org/10.1038/ng.3285

300   http://www.nature.com/ng/journal/v47/n7/abs/ng.3285.html#supplementary-information
301   Premoli, M. C., Sella, G., & Berra, G. P. (1996). Heritable variation of sex ratio in a
302   polychaete worm. *Journal of Evolutionary Biology*, *9*(6), 845-854.
303   https://doi.org/10.1046/j.1420-9101.1996.9060845.x
304   RCore, T. (2016). R: A language and environment for statistical computing. R Foundation for
305   Statistical Computing, Vienna, Austria. *URL http://www*. *R-project. org*.
306   RStudio. (2022). *RStudio: Integrated Development for R. *. In *RStudio BPC*.
307   Trotter, M. V., & Spencer, H. G. (2007). Frequency-Dependent Selection and the
308   Maintenance of Genetic Variation: Exploring the Parameter Space of the Multiallelic
309   Pairwise Interaction Model. *Genetics*, *176*(3), 1729-1740.
310   https://doi.org/10.1534/genetics.107.073072
311   Varandas, F. R., Sampaio, M. C., & Carvalho, A. B. (1997). Heritability of sexual proportion
312   in experimental sex-ratio populations of Drosophila mediopunctata. *Heredity*, *79*(1),
313   104-112. https://doi.org/10.1038/hdy.1997.128
314   Villanea, F. A., Safi, K. N., & Busch, J. W. (2015). A General Model of Negative Frequency
315   Dependent Selection Explains Global Patterns of Human ABO Polymorphism. *PLoS
316   ONE*, *10*(5), e0125003. https://doi.org/10.1371/journal.pone.0125003
317   Walsh, B., Lynch, M., Walsh, B., & Lynch, M. (2018). Maintenance of Quantitative Genetic
318   Variation. In *Evolution and Selection of Quantitative Traits* (pp. 0). Oxford University
319   Press. https://doi.org/10.1093/oso/9780198830870.003.0028
320   West, S. A. (2009). *Sex allocation*. Princeton University Press.

Zietsch, B. P., Walum, H., Lichtenstein, P., Verweij, K. J., & Kuja-Halkola, R. (2021). When theory cannot explain data, the theory needs rethinking. Invited replies to: Orzack SH, Hardy ICW. 2021, and Lehtonen J. 2021. *Proceedings of the Royal Society B*, *288*(1947), 20210304.

Zietsch, B. P., Walum, H., Lichtenstein, P., Verweij, K. J. H., & Kuja-Halkola, R. (2020). No genetic contribution to variation in human offspring sex ratio: a total population study of 4.7 million births. *Proc Biol Sci*, *287*(1921), 20192849-20192849. https://doi.org/10.1098/rspb.2019.2849