Agent-Based Modelling shows that Operation of Fisher's Principle does not explain

Zero Heritability in Human Offspring Sex Ratio

Kaitlyn T. Harper[1*], Morgan J. Sidari[1*], and Brendan P. Zietsch[1]

*Authors contributed equally

[1]Centre for Psychology and Evolution, School of Psychology, University of Queensland, Brisbane, Australia

Author Note

Kaitlyn T. Harper https://orcid.org/0000-0001-9597-872X

Morgan J. Sidari https://orcid.org/0000-0001-5315-4546

Brendan P. Zietsch https://orcid.org/0000-0003-0274-6140

We have no conflicts of interest to disclose.

Corresponding author email: kaitlyn.harper@uq.net.au

Data availability statement: All R code, results, and supplementary material are available at https://osf.io/cbpxr/?view_only=5c437beb09534d4e8b701045e16faf01.

Abstract

The ratio of male to female births has garnered interest and debate for more than a century. According to Fisher's principle, offspring sex ratio remains approximately equal through a process of negative frequency-dependent balancing selection. That is, individuals whose alleles predispose them to produce the rarer sex will have greater fitness, causing these alleles to increase in frequency until the population sex ratio is even again and the allele is no longer advantageous. Recently, though, analysis of all Swedish births since 1932 revealed that offspring sex ratio is not heritable, demonstrating that Fisher's principle cannot operate in humans. Some have argued that Fisher's principle makes no prediction about the heritability of offspring sex ratio at equilibrium and therefore can still serve as a valid explanatory framework. We used agent-based modelling to simulate the operation of Fisher's principle and its effects on genetic variation in offspring sex ratio over 5000 generations. Our simulations demonstrate that the operation of Fisher's principle at equilibrium does not actively eliminate heritability, contrary to implicit assumptions in previous arguments. These results, combined with recent evidence that offspring sex ratio is not heritable in humans, suggest that Fisher's principle is not a valid explanatory framework for human offspring sex ratio.

Key words: offspring sex ratio, Fisher's principle, negative frequency-dependent balancing selection, agent-based model, genetic variation, sex allocation

46   **Agent-Based Modelling shows that Operation of Fisher's Principle does not explain**

47   **Zero Heritability in Human Offspring Sex Ratio**

Offspring sex ratio – a parent's ratio of male to female births – has garnered interest

and debate for more than a century. It is also the subject of Fisher's principle (Fisher, 1930),

which traces back to Darwin (1871) and Dusing (cited in Edwards, 1998) and is sometimes

referred to as Dusing-Fisher theory. Fisher's principle states that in sexually reproducing

species, the offspring sex ratio will stabilise around 1:1 due to sexual selection favouring

whichever sex is less common. This process is an example of frequency-dependent balancing

selection: all else being equal, individuals genetically predisposed to producing the scarce sex

will tend to have greater fitness (more grandchildren), so those individuals' alleles will

increase in frequency until the population sex ratio rebalances and the advantage of the allele

disappears. According to the theory, this process keeps the sex ratio approximately even

(under the condition that male and female offspring require equal parental investment).

Fisher's principle has been the primary framework by which human offspring sex

ratio has been understood (Hardy, 2002; West, 2009). Recently, though, using data from the

whole Swedish population born 1932 or later (3,543,243 individuals and their 4,753,269

children), Zietsch et al. (2020) showed that offspring sex ratio is not heritable in humans.

That is, genetic differences do not influence the sex ratio of individuals' offspring, and

therefore variation in offspring sex ratio cannot be inherited. Even within individuals there

was no consistency at all in the sex of subsequent offspring, suggesting that offspring sex was

determined by random Mendelian segregation of sex chromosomes. These observations are

incompatible with the operation of Fisher's principle – offspring sex ratio must be heritable to

respond to selection as Fisher described.

If Fisher's principle cannot operate in humans, how can it serve as a framework for

understanding human sex ratio? Some have argued that Fisher's principle makes no

prediction about the heritability of offspring sex ratio at equilibrium (i.e. a stable state

reached through the operation of a process) and therefore can still serve as a valid

explanatory framework (Lehtonen, 2021; Orzack & Hardy, 2021). This position implies that

the operation of Fisher's principle has eliminated the genetic variation in offspring sex ratio

(since it must have been heritable for Fisher's principle to operate but is no longer heritable).

However, it is not clear why this elimination would be expected under Fisher's principle or

how it would occur. In fact, negative frequency-dependent selection (of which Fisher's

principle is an example) is expected to actively maintain genetic variation, not eliminate it

(Brisson, 2018; Clarke, 1979; Zietsch et al., 2021). It is therefore difficult to see how zero

heritability is compatible with Fisher's principle as an explanation of human offspring sex

ratio.

The expectation that negative frequency-dependent selection actively maintains

genetic variation is supported by modelling where a given trait is influenced by variants at a

limited number of genetic loci (Asmussen & Basnayake, 1990; Trotter & Spencer, 2007;

Villanea et al., 2015). To our knowledge there has been no modelling of the effect of Fisher's

principle per se on genetic variation (and subsequent heritability) in offspring sex ratio. Given

the complex underlying mechanisms of offspring sex ratio evolution, which are not always

obvious, it is important to investigate this specific mechanism. Here, we used agent-based

modelling to simulate the operation of Fisher's principle and its effects on genetic variation

and heritability in offspring sex ratio.

91   **Method**

We created an agent-based model to test whether the operation of Fisher's principle depletes

genetic variation in offspring sex ratio. This model was comprised of male and female agents,

each possessing a small genome of 20 loci which influenced their tendency to produce male

or female offspring. By simulating a population of 2000 agents to reproduce over 5000

generations, we were able to see Fisher's principle influence the population sex ratio, as well

as observe how selection affected agents' genetic predisposition to have male or female

offspring and whether the genetic variation and heritability of this trait were depleted over

generations. We decided to use 20 loci to ensure that offspring sex ratio was modelled as a

quantitative trait (as opposed to a discrete trait with one or two loci). We simulated 5000

generations to capture any long-term effects occurring at sex ratio equilibrium.

Computational load was also considered when deciding on number of loci, number of

generations, and population size. We arbitrarily labelled the scarce sex as male, but the

results are identical the other way around. In the following sections, we describe how agents

were created and the rules by which they reproduced. All code and results can be accessed at

https://osf.io/cbpxr/?view_only=5c437beb09534d4e8b701045e16faf01. All modelling was

performed in R in R studio version 1.3.959 (RCore, 2016; RStudio, 2022).

Creating Initial Generation

An initial population of 2000 agents was generated, 40% were males. In the reported

model each agent had 20 loci, each with two alleles, that influenced offspring sex ratio. Initial

allele values were generated as either 0 or 1 (low variant or high variant), which occurred

with a probability that reflected the initial population sex ratio. In the reported model, the

initial population sex ratio was 40% male, so the allele values were randomised with a 0.4

probability (such that genetic scores reflected a 40% probability of producing a male

offspring). Genotype was then calculated as *allele 1 + allele 2*, thereby giving agents

genotype values of 0, 1, or 2 at each locus. Genetic score for offspring sex ratio was

calculated by summing genotypes for all loci. In the reported model there were 20 loci

influencing offspring sex ratio, so possible genetic scores ranged between 0 and 40.

Partner Selection

The following operations were performed five times to create five 'offspring rounds'.

Agents were paired randomly, with some surplus-sex agents missing out on partnering and

reproducing for that round. Mating success for the surplus sex therefore varied randomly

between agents (ranging from 0-5), a result of the number of times they were randomly

excluded from the mating round. Partners' data were then linked with the agents' data so that

values for both parents were available to determine offspring attributes.

Determining Offspring Sex

The following process emulated a genetic predisposition toward male or female

offspring with additional environmental variance, such that offspring sex was determined

80% by genes and 20% by environmental factors (to simulate a largely but not completely

heritable trait, like most biological traits). For each offspring, the parents' genetic scores were

averaged and transformed to a proportion of the highest theoretical genetic score (e.g. in the

reported model the maximum genetic score was 40, so a score of 10 would be transformed to

0.25). This proportionate genetic score was then multiplied by 0.8, and an additional random

value between 0 and 0.2 was added to represent 20% environmental influence. The resulting

score was used as the probability in a random binomial outcome where 0 = female, and 1 =

male; higher scores were more likely to produce male offspring.

Reproduction

Alleles were then inherited through independent assortment. That is, at each locus, an

offspring inherited one allele from each parent, determined randomly. Outcome measures

were recorded for the current generation at this point. The population matrix was then

truncated (i.e. all parent information removed), leaving the offspring to become the next

generation, and the process described thus far was repeated. In the reported model, each

simulation was run for 5000 generations.

Outcome Variables and Analysis

We ran the model described above 50 times and averaged the simulation results for

each generation. The outcome variables of interest were population sex ratio, genetic score,

heritability, and genetic variance. Offspring sex ratio was recorded as the proportion of male

offspring produced. Genetic score was recorded as the mean of the raw genetic scores. Given

that our data was simulated, and we had complete genome information for each agent, we

used simpler calculations of genetic variance and heritability than those normally used in the

broader quantitative genetics literature. To calculate heritability, i.e. the proportion of

phenotypic variance accounted for by genetic variance, we used the r^2 of genetic score

predicting offspring sex ratio (i.e. the proportion of the agent's offspring that were male). To

calculate genetic variance, we multiplied the total phenotypic variance (variance in offspring

sex ratio) by the heritability, giving us the variance in offspring sex ratio that is explained by

genetic scores. We used correlations to assess trends in outcome variables over generations.

For ease of presentation in the main text, we have reported our model of Fisher's principle for

1000 generations. Extended results (with 5000 generations) can be found in the

Supplementary Material.

Results

Sex ratio and genetic scores

To determine whether Fisher's principle operated as expected in our model, we

needed to establish whether genetic scores, and consequently sex ratio at birth, changed over

time. Results demonstrated that the genetic scores diverged based on agent gender, with male

agents having higher scores than female agents (see Figure 1A). This result was to be

expected, given that a male is more likely to have been produced by male-making alleles, and

thus will have inherited more male-making alleles themselves; likewise, females will have

inherited more female-making alleles. We ran a simulation where the initial genetic scores for

each sex reflected this natural separation between male and female genetic scores, and all

results were comparable.

In addition to the divergence of genetic scores by sex, the mean genetic score also

increased over time (from the initial mean genetic score which was biased towards female-

making alleles), such that alleles for having male offspring (i.e. the rarer sex) became more

prevalent (Figure 1A). The rate of this increase slowed until around generation 500 when the

process reached an equilibrium at a 1:1 predisposition for offspring sex (at the midpoint of

possible genetic scores – 20). The resulting population sex ratio also approached a 1:1

equilibrium (0.5 male) over the generations (see Figure 1B). The observed pattern of results

thus far is precisely as Fisher's principle would predict, where alleles leading to the less

common sex increased in frequency until the sex ratio reached equilibrium at 1:1, at which

point these alleles no longer conferred a benefit.

Genetic variation and heritability

To address arguments that offspring sex ratio might no longer be heritable due to the

operation of Fisher's principle (Lehtonen, 2021; Orzack & Hardy, 2021), we next sought to

test whether genetic variation and heritability were eliminated over generations.

Although heritability and genetic variance appear to be stable across generations at

first glance (see Figure 1C and Figure 1D), correlations revealed that they each subtly

decreased over time ($r = -0.36$, $p < .001$, for both analyses). Upon further exploration of the

data, this decrease appears to be due to fixation of some alleles due to drift. An allele can fix

at zero or 100% frequency due to chance fluctuations over many generations (i.e. drift), but

once fixed it cannot change frequency (without mutation); that is, allelic variation can be lost

but not replenished in our model, and so the occasional chance fixations will cause the

genetic variation to decrease slightly over time. In analyses described in the Supplementary

Material, we demonstrate that the slight decreases in heritability detected in our model of

Fisher's principle are slightly less than those observed under no selection at all (and far less

than observed under directional and stabilising selection). In summary, the operation of

Fisher's principle does not deplete genetic variation, but also does not entirely buffer the loss

of genetic variation due to drift.

Figure 1. A Genetic score, **B** population sex ratio, **C** heritability, and **D** genetic variance over 1000

generations.

Note. Grey lines represent individual results from 50 simulations. Coloured lines represent the mean across

all simulations at each generation. In Figure B, the blue line represents the mean male genetic score and the

red line represents the mean female genetic score. In Figure A and Figure B, the dotted horizontal line

represents the midpoint of offspring sex ratio (1:1)

Discussion

For more than a century, Fisher's principle has been widely accepted as the key

explanatory framework for the 1:1 offspring sex ratio in humans (Hardy, 2002; West, 2009).

However, a recent study demonstrated that offspring sex ratio is not heritable in humans

(Zietsch et al., 2020), which is inconsistent with the current operation of Fisher's principle

and challenges Fisher's principle as a framework for understanding human offspring sex

ratio. It has been suggested that Fisher's principle is consistent with zero heritability of

offspring sex ratio at equilibrium, and therefore can still be used to understand the evolution

of human offspring sex ratio (Lehtonen, 2021; Orzack & Hardy, 2021). This position implies

that Fisher's principle operating in the past has eliminated the heritability of offspring-sex

ratio over time – yet it is unclear how Fisher's principle would lead to this elimination

(Zietsch et al., 2021). Previous models in limited contexts have demonstrated that negative

frequency-dependent balancing selection, of which Fisher's principle is an example, is

expected to maintain (rather than deplete) genetic variance (Asmussen & Basnayake, 1990;

Trotter & Spencer, 2007; Villanea et al., 2015). However, given the unique complexities of

sex ratio evolution, we developed a specific computational model of Fisher's principle to

assess whether it could have caused the elimination of genetic variation and heritability of

offspring sex ratio over evolutionary time.

In the first generation of our model, males were the scarcer sex and the population's

alleles reflected this with a bias toward producing females. Over many generations, the sex

ratio and genetic scores of the population approached equilibrium at a 1:1 offspring sex ratio.

This pattern of results demonstrates Fisher's principle as expected, such that alleles that

produce the scarcer sex will increase until they are no longer beneficial (once a 1:1

population sex ratio is reached). At this point, there is no selection for these alleles, which is

consistent with our findings that population sex ratio and genetic scores remained stable after

reaching equilibrium. While our model demonstrates that Fisher's principle is operationally

sound, it does not support arguments that it could have eliminated offspring sex ratio

heritability over time. We conclude this because Fisher's principle, in our simulations,

maintained heritability for longer than directional and stabilising selection and was

comparable to genetic drift (no selection). Thus, Fisher's principle cannot account for the

current observable state of no heritability in offspring sex ratio while there remains abundant

heritability in virtually all variable traits that have been studied (both those under selection

and no selection).

237   Our findings suggest that the operation of Fisher's principle cannot be reconciled with

238   the current non-heritable nature of offspring sex ratio, thus challenging the principle's

239   relevance for understanding human offspring sex ratio. We suggest that ideas of how

240   offspring sex ratio has evolved in humans need to be overhauled, given that Fisher's principle

241   has had a foundational and lasting influence in the field, as described by West (2009, p.14):

242   *'Fisher's theory of equal investment provides the basic null model for sex allocation*

243   *theory, but it is also the foundation for all subsequent theoretical developments.'*

244   Hardy (2002, p.2, emphases added) also acknowledges that Fisher's principle underpins

245   extensive subsequent research:

246   *'Darwin, Düsing, Fisher, and Shaw and Mohler established the fundamental principle*

247   *that members of the minority sex tend to have higher fitness than members of the*

248   *majority sex. They also outlined how various ecological, demographic and genetic*

249   *variables might affect the details of sex-allocation strategies by modifying both the*

250   *constraints and the fitness functions. **Modern sex-allocation research is devoted***

251   ***largely to the exploration of such effects**, which connect sex ratios to many other*

252   *aspects of the biologies of many species'*

Our model is a simplified representation of the real world, and as with any model,

there are limitations. For example, the discrete generations in our model are a simplified

representation of mating environments, which in reality have overlapping generations.

Additionally, our agents and their genomes of 20 loci cannot comprehensively represent true

human genetic architecture. However, we have no reason to expect that complexities

involving linkage disequilibrium or minor allele frequencies would substantively affect our

findings, especially since Fisher's description of his principle does not implicate any of these

factors. We also did not consider factors such as group size and environmental fluctuations in

sex ratio. For instance, some might argue that ongoing environmental fluctuations in

population sex ratio in small groups, where one or the other sex temporarily experiences high

mortality, could lead to continuous selection and therefore a faster loss of genetic variation.

However, this perspective overlooks the fact that environmental fluctuations would not alter

the existing genetic predisposition for offspring sex ratio in the surviving individuals.

Therefore, while these changes might temporarily affect the sex ratio, the population would

return to its natural equilibrium in the next generation. While differential mortality between

sexes during parental care might influence selection on offspring sex ratio (Kahn et al.,

2015), no specific set of circumstances has been proposed that would result in a continuous

reduction of genetic variance over time.

Additionally, while our model demonstrates that Fisher's principle maintains genetic

variation in comparison to other forms of selection, genetic variation still slightly reduced,

though the rate was slower than under no selection at all. While some previous models of

negative frequency-dependent selection did not allow for the effects of drift (Asmussen &

Basnayake, 1990; Trotter & Spencer, 2007), one study found that negative frequency-

dependent selection preserved heterozygosity, even counteracting drift (Villanea et al., 2015).

Drift may have had more of an effect in our model due to a range of factors related to model

complexity, such as population size and number of alleles; for example, the fact that our

model used 20 loci instead of one (as in Villanea et al., 2015) would mean that the balancing

selection on any one allele is much lower and thus more vulnerable to being lost to drift.

Regardless, could the slight decrease in heritability in our model explain the observed

zero heritability of offspring sex ratio that we currently observe? This possibility does not

make sense in the context of pervasive heritability of human traits (Polderman et al., 2015),

most of which would be either under selection that would deplete heritability or under no

selection at all. Given our results, Fisher's principle would predict offspring sex ratio to be

more heritable than most traits; instead, offspring sex ratio is basically unique among variable

traits in being non-heritable.

Our models' findings, combined with recent evidence that offspring sex ratio is not

heritable in humans (Zietsch et al., 2020), suggest that Fisher's principle is not a valid

explanatory framework for variation in the offspring sex ratio in humans. Moreover, the

inconsistency of offspring sex ratio within people (Zietsch et al., 2020) suggests that there is

no predisposition for offspring sex at an individual level. Instead, a random Mendelian

process may determine the offspring sex ratio in humans, although it is unclear what

evolutionary processes could have led to this outcome. Orzack and Hardy (2021) proposed

that natural selection for a 1:1 sex ratio could be linked to the Mendelian process of sex

determination, but such a mechanism would be far removed from Fisher's principle as

originally proposed. It remains unclear whether individual or group selection for a 1:1

offspring sex ratio could lead to fixation of a mechanism that produces a random sex ratio at

the individual level. We encourage readers to explore the previous papers on this issue for

more theoretical discussion of these possibilities (Lehtonen, 2021; Orzack & Hardy, 2021;

Zietsch et al., 2021; Zietsch et al., 2020). We anticipate that new theoretical models will be

necessary to reconstruct our understanding of sex ratio evolution in humans, and we look

forward to seeing progress in this field.

305   References

306   Asmussen, M. A., & Basnayake, E. (1990). Frequency-dependent selection: the high potential
307   for permanent genetic variation in the diallelic, pairwise interaction model. *Genetics,*
308   *125*(1), 215-230. https://doi.org/10.1093/genetics/125.1.215

309   Brisson, D. (2018, 2018-February-21). Negative Frequency-Dependent Selection Is
310   Frequently Confounding [Perspective]. *Frontiers in Ecology and Evolution, 6.*
311   https://doi.org/10.3389/fevo.2018.00010

312   Clarke, B. C. (1979). The evolution of genetic diversity. *Proceedings of the Royal Society of*
313   *London. Series B. Biological Sciences, 205*(1161), 453-474.
314   https://doi.org/10.1098/rspb.1979.0079

315   Darwin, C. R. (1871). *The descent of man, and selection in relation to sex*. John Murray.

316   Edwards, A. W. F. (1998, 1998/06/01). Natural selection and the sex ratio: Fisher's sources.
317   *The American Naturalist, 151*(6), 564-569. https://doi.org/10.1086/286141

318   Fisher, R. A. (1930). *The genetical theory of natural selection*. Oxford University Press.

319   Hardy, I. (2002). *Sex ratios : concepts and research methods*. Cambridge University Press.

320   Kahn, A. T., Jennions, M. D., & Kokko, H. (2015). Sex allocation, juvenile mortality and the
321   costs imposed by offspring on parents and siblings. *Journal of Evolutionary Biology,*
322   *28*(2), 428-437.

323   Lehtonen, J. (2021). Fisher's principle remains a plausible explanation for human sex ratio
324   evolution. *Proceedings of the Royal Society B: Biological Sciences*.

325   Orzack, S. H., & Hardy, I. (2021). Does the lack of heritability of human sex ratios require a
326   rethink of sex ratio theory? No: a response to Zietsch et al. 2020. *Proceedings of the*
327   *Royal Society B: Biological Sciences*.

328   Polderman, T. J. C., Benyamin, B., de Leeuw, C. A., Sullivan, P. F., van Bochoven, A.,
329   Visscher, P. M., & Posthuma, D. (2015, 07//print). Meta-analysis of the heritability of
330   human traits based on fifty years of twin studies [Analysis]. *Nat Genet, 47*(7), 702-
331   709. https://doi.org/10.1038/ng.3285

332   RCore, T. (2016). R: A language and environment for statistical computing. R Foundation for
333   Statistical Computing, Vienna, Austria. *URL http://www. R-project. org*.

334   RStudio. (2022). *RStudio: Integrated Development for R. . In RStudio BPC*.

335   Trotter, M. V., & Spencer, H. G. (2007). Frequency-Dependent Selection and the
336   Maintenance of Genetic Variation: Exploring the Parameter Space of the Multiallelic
337   Pairwise Interaction Model. *Genetics, 176*(3), 1729-1740.
338   https://doi.org/10.1534/genetics.107.073072

339   Villanea, F. A., Safi, K. N., & Busch, J. W. (2015). A General Model of Negative Frequency
340   Dependent Selection Explains Global Patterns of Human ABO Polymorphism. *PLoS*
341   *ONE, 10*(5), e0125003. https://doi.org/10.1371/journal.pone.0125003

West, S. A. (2009). *Sex allocation*. Princeton University Press.

Zietsch, B. P., Walum, H., Lichtenstein, P., Verweij, K. J., & Kuja-Halkola, R. (2021). When
theory cannot explain data, the theory needs rethinking. Invited replies to: Orzack SH,
Hardy ICW. 2021, and Lehtonen J. 2021. *Proceedings of the Royal Society B,*
288(1947), 20210304.

Zietsch, B. P., Walum, H., Lichtenstein, P., Verweij, K. J. H., & Kuja-Halkola, R. (2020). No
genetic contribution to variation in human offspring sex ratio: a total population study
of 4.7 million births. *Proc Biol Sci, 287*(1921), 20192849-20192849.
https://doi.org/10.1098/rspb.2019.2849

352