Title: Whole-genome sequencing confirms multiple species of Galapagos giant tortoises

Authors: Stephen J. Gaughran[1*], Rachel Gray[2], Menna Jones[3], Nicole Fusco[4,5], Alexander

Ochoa[4], Joshua M. Miller[6], Nikos Poulakakis[7,8,9], Kevin de Queiroz[10], Adalgisa Caccone[4],

Evelyn L. Jensen[2*]

[1] Department of Ecology & Evolutionary Biology, Princeton University, Princeton, NJ, USA

08544

[2] School of Natural and Environmental Sciences, Newcastle University, Newcastle Upon Tyne,

UK

[3] Department of Earth Sciences, Oxford University, Oxford, UK

[4] Department of Ecology & Evolutionary Biology, Yale University, New Haven, CT, USA

06511

[5] Department of Ecology & Evolutionary Biology, University of Connecticut - Stamford,

Stamford, CT, 06901

[6] Department of Biological Sciences, MacEwan University, Edmonton, AB, Canada

[7] Department of Biology, School of Sciences and Engineering, University of Crete, Heraklion,

Greece

[8] The Natural History Museum of Crete, School of Sciences and Engineering, University of

Crete, Heraklion, Greece

[9] Institute of Molecular Biology and Biotechnology, Foundation for Research and Technology-

Hellas, Heraklion, Greece

[10] Department of Vertebrate Zoology, National Museum of Natural History, Smithsonian

Institution, Washington, DC 20560-0162 USA

*SJG: sjgaughran@princeton.edu; ELJ: evelyn.jensen@newcastle.ac.uk

Running title: Galapagos giant tortoise species delimitation

Keywords: Galapagos giant tortoise, species, taxonomy, multispecies coalescent, genomics

Abstract

Galapagos giant tortoises are endemic to the Galapagos Archipelago, where they are found in

isolated populations. While these populations are widely considered distinguishable in

morphology, behavior, and genetics, the recent divergence of these taxa has made their status as

species controversial. Here, we apply multispecies coalescent methods for species delimitation to

whole genome resequencing data from 38 tortoises across all 13 extant taxa to assess support for

delimiting these taxa as species. In contrast to previous studies based solely on divergence time,

we find strong evidence to reject the hypothesis that all Galapagos giant tortoises belong to a single

species. Instead, a conservative interpretation of model-based and divergence-based results

indicates that these taxa form a species complex consisting of a minimum of 9 species, with some

analyses supporting as many as 13 species. There is mixed support for the species status of taxa

living on the same island, with some methods delimiting them as separate species and others

suggesting multiple populations of a single species per island. These results make clear that

Galapagos giant tortoise taxa represent different stages in the process of speciation, with some taxa

further along in that evolutionary process than others. A better understanding of the more complex

parts of that process is urgently needed, given the threatened status of Galapagos giant tortoises.

Lay Summary

Species delimitation is a challenging problem in evolutionary biology, but one that is central to the

field. Distinguishing species can affect conservation management practices, from conservation

status assessments to strategies for breeding programs. More fundamentally, understanding species

boundaries affects our ability to assess biodiversity and to study evolutionary processes. The

Galapagos Archipelago presents several radiations of closely related taxa that inspired Charles

Darwin to develop his theory of evolution by natural selection and later led to foundational case

studies in speciation. The Galapagos giant tortoises were one such inspiration. Nearly two

centuries later, there is still an ongoing debate about the taxonomic status of these tortoises, with

opinions on their status ranging from barely differentiated populations to separate species. Here,

we present the first genomic species delimitation of Galapagos giant tortoises and provide

convincing evidence that this group is a complex consisting of between 9 and 13 species. These

results provide valuable guidance to conservation stakeholders in the Galapagos, while also adding

an important case study to the delimitation of island species.

Introduction

Speciation is a complex biological process driven at least in part by ecological context such

as physical barriers to gene flow, adaptation to local environments, and population-specific

demographic dynamics. In recently diverged lineages, the relative roles of the evolutionary forces

responsible for divergence can be challenging to describe accurately because of shared ancestral

polymorphisms in the descendant lineages and introgression due to ongoing gene flow (Shaffer &

Thomson 2007). Adding to these difficulties are the small founding populations that characterize

the origin of some species, especially in island settings, which can speed up the divergence of

lineages through the rapid loss or fixation of alleles (Kimura & Ohta 1969). The iconic radiation

of Galapagos giant tortoises (a clade within the genus *Chelonoidis*) is a compelling example of the

complexities of species delimitation in a case of recent diversification: their divergence has been

molded by a combination of vicariance and colonization events (Caccone *et al*. 1999, 2002;

Poulakakis *et al*. 2012, 2020) brought about by both natural and anthropogenic environmental

changes and making accurate species delimitation challenging.

The Galapagos Islands, the conceptual home of the theory of evolution by natural selection,

provide a crucible for the study of speciation and taxonomic complexity. Within the Galapagos

Archipelago there are many examples that show clear evidence of recently diverged but genetically

distinct species, including finches (Grant & Grant 2003), lava lizards (Benavides *et al.* 2009),

iguanas (MacLeod *et al.* 2015), mockingbirds (Arbogast *et al.* 2006), moths (Schmitz *et al.* 2007),

and tortoises (Caccone *et al.* 1999). This archipelago therefore provides a model system to

understand the complex realities of speciation, allowing us to explore multiple lines of molecular,

morphological, and ecological evidence when proposing species delimitations.

There are 16 Galapagos giant tortoise taxa, three of which (*niger*, *abingdonii*, and an

unnamed taxon from Santa Fé Island) are recently extinct (Figure 1). Morphological, behavioral,

and genetic differences have been documented among the taxa (Gaughran *et al.* 2018, Jensen *et*

al. 2021, Chiari 2021, Hunter *et al.* 2013). The most striking morphological difference is carapace

shape, which ranges from a domed shape to a saddleback shape with an elevated anterior carapace

opening. These carapace shapes have a genetic underpinning, as evidenced by a clear phylogenetic

signal (Jensen *et al.* 2022) and a strong inheritance in juveniles of different taxa raised in a common

environment (Pritchard 1996). The taxa also differ in coloration on the head and neck, as well as

in limb length (reviewed in Chiari 2021) and aggressive behaviors (Schafer & Krekorian 1983).

Mitochondrial, microsatellite, and genomic data all suggest that geographically distinct

populations represent genetically differentiated lineages (e.g., Caccone *et al.* 2002, Beheregaray

et al. 2003, Gaughran *et al.* 2018, Miller *et al.* 2018).

Galapagos giant tortoises are descended from individuals that likely floated from

continental South America along the Humboldt Current to the newly formed Galapagos Islands

around 3 million years ago (Ma; Caccone *et al.* 2002). As more volcanic islands emerged over

time, tortoises migrated from existing islands in the east to newer ones in the west (Poulakakis *et*

al. 2020). Divergence dates from mitochondrial DNA suggest that the progressive colonization

from the first colonized island likely occurred after 2 Ma and continued up to the emergence of

the youngest island to the west, Fernandina, around 60,000 years ago (Geist *et al.* 2014). The

layout of the archipelago has changed dramatically over this time period due to volcanic emergence

and subsidence, and sea level changes. During the Pleistocene the central archipelago had a larger

landmass that has since fragmented into some of the islands seen today (Geist *et al.* 2014). Thus,

the diversification of Galapagos giant tortoises has been triggered by a combination of dispersal

to isolated islands and vicariance as landmasses fragmented (Parent *et al.* 2008). In addition, within

the last 200 years, humans have moved tortoises between islands (Caccone *et al.* 2002, Poulakakis

et al. 2008) and land use change has brought previously isolated lineages into contact (Russello *et*

al. 2005), resulting in low levels of recent gene flow between previously isolated populations.

Because of this recent and dynamic evolutionary background, the taxonomy of the

Galapagos giant tortoises has been the subject of almost endless debate since the time of Darwin

(e.g., Darwin 1839, Günther 1877, Van Denburgh 1907, Pritchard 1996, Zug 1998, Caccone *et al.*

1999, Kehlmaier *et al.* 2021). Most taxonomic proposals have recognized these lineages as either

a single species with 2-14 subspecies, or as many as 14 separate species (reviewed in Pritchard

1996). Following work by Caccone *et al.* (1999), they have been listed as multiple species based

on genetic data accumulated over the last several decades (Rhodin *et al.* 2017). More recently,

however, two publications have advocated collapsing all taxa into a single species with multiple

subspecies (Loire *et al.* 2013, Kehlmaier *et al.* 2021), which was adopted by the IUCN Tortoise

and Freshwater Turtle Specialist Group in 2021 (Rhodin *et al.* 2021).

To address this question, we turned to whole genome resequencing data and methods that

quantitatively assess the degree to which taxa are independently evolving under a multispecies

coalescent (MSC) framework (Zhang *et al.* 2011, Jackson *et al.* 2017, Morales *et al.* 2018, Mays

et al. 2019, Leaché *et al.* 2019, Marshall *et al.* 2021). Here, we apply several of these methods of

species delimitation to the Galapagos giant tortoises, and thereby provide an assessment of

taxonomic considerations in this endangered clade.

Methods

<u>Terminology, study design and data</u>

The aim of this paper is to assess genomic evidence for the distinctiveness and taxonomic

status of Galapagos giant tortoise taxa using phylogenetic and coalescent frameworks. To remain

agnostic about the taxonomic status of the various tortoise populations prior to presenting our

results, we avoid designating the taxa as *species* or *subspecies* and refer to them using only the

epithets (e.g., *phantasticus* rather than *Chelonoidis phantasticus* or *Chelonoidis niger*

phantasticus). We mapped existing Illumina short-read data (NCBI Bioproject PRJNA761229;

Jensen *et al.* 2021, Jensen *et al.* 2022) to the Pinta Island Galapagos giant tortoise reference

genome (NCBI assembly ASM359739v1, Quesada *et al.* 2019). Samples in this short-read data

set included two individuals of *phantasticus*, and three individuals each of *guntheri*, *microphyes*,

vandenburghi, *vicina*, *becki* (PBL population), *becki* (PBR population), *darwini*, *chathamensis*,

donfaustoi, *porteri*, *duncanensis*, and *hoodensis*, for a total of 38 Galapagos giant tortoise

individuals plus one outgroup (Chaco tortoise, *C. chilensis*) individual. We called variant sites for

each individual using the BCFtools variant calling pipeline (Li *et al.* 2009) and created a consensus

fasta file for each individual from the VCF. We then generated a random set of 1kb loci separated

by 100kb. We filtered, masked, and generated phased haplotypes for each locus. Details on these

terminology choices, study design, and data are in the Supplemental Methods.

<u>SNAPP tree generation</u>

To assess the phylogenetic relationships among the extant populations of Galapagos giant

tortoises, we used SNAPP v1.5.2 (Bryant *et al.* 2012) and MODEL_SELECTION v1.5.3 (Baele

et al. 2012), as implemented in BEAST2 v2.6.7 (Bouckaert *et al.* 2014), to test 21 different

phylogenetic models (Supplemental Table S1) on a dataset comprising 1,000 SNPs from 38

Galapagos giant tortoise individuals, plus the *C. chilensis* individual as an outgroup. We then

applied the priors from the best model to an extended dataset comprising 5,000 SNPs for

phylogenetic reconstruction and downstream species delimitation comparisons. A more detailed

description of these methods is presented in Supplemental Methods.

 Species delimitation models

We implemented the Bayesian species delimitation model in BPP v4.0 (Flouri *et al.* 2018)

to test hypotheses about the number of species in our data set. We ran the species delimitation

analyses with 50, 200, 500, or 1000 phased loci, with each locus a randomly selected alignment of

1000 bp (see Supplemental Methods). We ran BPP species delimitation with the SNAPP guide

tree (i.e., "A10" analysis) and without a guide tree (i.e., "A11" analysis), repeating each run three

times with different random seeds to assess convergence among runs. Because the Galapagos

Islands are geologically young (<3.5 Ma) and the deepest divergence time among Galapagos giant

tortoise lineages is estimated to be between 2 and 3 Ma (Caccone *et al*. 2002), we chose a small

but diffuse prior for divergence time ($\tau \sim$ IG(3, 0.001)), which corresponds to a mean of 83,333

generations (approximately 2.08 million years). In addition, given the low estimates of effective

population size for Galapagos giant tortoises (Garrick *et al*. 2015, Jensen *et al*. 2021) and because

the carrying capacity must be low for large-bodied terrestrial vertebrates on these semi-desert

islands, we used a prior scaled effective population size of $\theta \sim$ IG(3, 0.001), which is equivalent

to N_e=20,833. However, because there is uncertainty about historical population size, we also re-

ran the analyses with scaled effective population size priors that were larger ($\theta \sim$ IG(3, 0.005),

N_e=104,167) or smaller ($\theta \sim$ IG(3, 0.0001), N_e=2083), keeping the priors for τ the same. In each

of run, we used a burn-in of 100,000, a sampling frequency of 2, and collected 300,000 samples.

To assess if the chosen loci and included taxa affected our analyses, we ran the above

analyses with a different set of 50, 200, 500, and 830 loci and including the Chaco tortoise. The

inclusion of this outgroup allowed the model to explore a species delimitation of two (i.e.,

Galapagos giant tortoises and Chaco tortoise). Because the divergence time estimate of the Chaco

tortoise and Galapagos giant tortoises is around 12 Ma (Caccone *et al.* 1999, Kehlmaier *et al.*

2017), we used a divergence time prior of $\tau \sim IG(3, 0.01)$. Finally, we re-ran the analyses with 50

unphased loci to ensure that our phasing method was not affecting the results.

<u>Genealogical divergence index (*gdi*) from BPP</u>

The use of species delimitation models in BPP is thought to be a robust method for

sympatric species delimitation, but there are concerns that using this MSC-based model selection

for species delimitation may delimit populations rather than species, especially if the taxa are

allopatric (Jackson *et al.* 2017, Leaché *et al.* 2019). Therefore, we also calculated the *gdi*, a

distance-based method with heuristic cutoffs for species delimitation (Jackson *et al.* 2017). The

gdi is calculated from the divergence time in coalescent units ($2\tau/\theta$) and reflects the probability

that two sequences from a purported species coalesce before the divergence time with a sister

species (τ) (Leaché *et al.* 2019). As such, it is a coalescent-based measure of genetic divergence,

as opposed to a sequence-based measure. Using the effective population size and divergence time,

the *gdi* can be calculated in a pairwise manner for sister species *a* and *b*, and can range from 0

(panmixia) to 1 (strong divergence). Based on empirical data and simulations, Jackson *et al.* (2017)

proposed a general heuristic for delimiting species based on the *gdi*. Namely, *gdi* < 0.2 indicates a

single species, and *gdi* > 0.7 indicates different species. Values for *gdi* between these cutoffs

indicate ambiguous delimitation.

To calculate these indices, we used BPP for parameter estimation of τ and θ, under the

guide tree and fixed MSC model ("A00") function in BPP. For these analyses, we used the 830

phased loci available in the data set with the Chaco outgroup. Using the phylogeny from our

SNAPP analysis as a fixed guide tree, we estimated τ for each node and θ for each branch and tip,

by running the MCMC with a burn-in of 100,000, a sample frequency of 2, and a total sample of

200,000. We used the same loci and priors as in the species delimitation modeling. We then used

the MCMC output to create posterior distributions of τ (for each node) and θ (for each tip), which

we then used to calculate *gdi*. Using posterior distributions allowed us to calculate 95% credibility

intervals. Following the suggestion of Leaché *et al*. 2019, we also successively collapsed each

node of the phylogeny, re-ran the A00 parameter estimation model, and calculated *gdi* for each

collapsed taxon.

<u>PHRAPL</u>

We then assessed the presence of gene flow between all pairs of lineages within each of

the two main clades of the tree (domed and saddleback) and tested for a collapse event using

PHRAPL (Jackson *et al*. 2017) implemented in R version 4.2 (R Core Team, 2022). PHRAPL

works by estimating the probability that a set of gene trees are observed under a given model by

calculating the frequency at which they are observed in a distribution of expected tree topologies,

weighting the probability by Akaike information criterion (AIC). In this way, PHRAPL can

determine the most likely demographic history. Three demographic models for each pairwise

comparison were constructed to test varying divergence and gene flow scenarios between

populations of Galapagos giant tortoises (Supplemental Figure S1). Model 1 was a two species

isolation only model, which tested for divergence between two populations with no ongoing

migration. Model 2 was also a two-species model and tested for constant symmetrical migration

between the two populations. The final model (Model 3) was the same as Model 2 but we removed

the divergence between the two populations using the *setCollapseZero* function in PHRAPL. This

single species model therefore allowed for constant symmetrical gene flow between the

populations with no divergence events. Input gene trees were generated for each of the 830 phased

loci in RaxML (v 8.2.12, Stamatakis 2014) with 20 replicate searches, rapid hill-climbing and the

GTRGAMMA model. The Chaco tortoise was included as an outgroup to root the tree. PHRAPL

is a demographic model and requires individuals to be assigned to a population prior to model

construction. Here, individuals were assigned to a population based on geographic location, as

detailed in Figure 1. In this way, the tortoises on each island were represented by a single

population except that six geographically separate populations were recognized on Isabela Island

and two on Santa Cruz Island. each island lineage was assigned to a distinct population, with the

exception of Isabela Island having six populations, based on the six geographically separated

lineages, and Santa Cruz island having two populations. Gene trees were subsampled at random

with replacement 100 times, sampling 2 individuals per lineage in each replicate. The outgroup

was not included in the subsampling and so was not included in the PHRAPL runs. Simulation of

100,000 gene trees was conducted using a grid of parameter values for divergence time (t) and

migration (m; Supplemental Table S2). The initial divergence within the Galapagos tortoise

radiation is estimated to be within the last 2 Ma, but the species pairs we compare here are more

recent (Poulakakis *et al.* 2020). We set the parameters of our grid search to capture this by limiting

the maximum divergence (t) to 1 Ma. Migration rates were equal to 4Nm, where Nm is the number

of migrants per generation, with the lowest migration being equivalent to one migrant every 800

years and the highest equivalent to one migrant every 25 years. This range of values was designed

to capture the putatively complex historical gene flow among populations. Akaike weights (wAIC)

were used to compare models and calculate model probabilities ranging from 0 (low support) to 1

(high support). To present the best supported hypothesis overall for each pairwise comparison, the

summed wAIC of the two-species models (Models 1 and 2) was compared with the wAIC of the

single species model (Model 3). We interpreted a summed wAIC of > 0.9 across the two-species

models as strong support for a two species hypothesis, a wAIC > 0.9 of the one species model as

strong support for a single species hypothesis, and all other scenarios as ambiguous. In addition to

identifying the top model chosen by PHRAPL, we also calculated the *gdi* value between taxon

pairs following the approach taken in Jackson *et al.* 2017 and using the *CalculateGdi* function in

PHRAPL. The *gdi* value from PHRAPL differs from that from BPP in that it is calculated using

the model averaged divergence rate (*t*), migration rate into population 1 (*M1*), and migration into

population 2 (*M2*). Because our models did not estimate the direction of migration, *M1=M2* in our

analyses. We interpreted the *gdi* values using the same thresholds described above.

Results

<u>SNAPP</u>

Of the 21 SNAPP models tested, Model 6 ($\alpha = 5$, $\beta = 150$, $\lambda = 39$) presented the greatest

marginal likelihood estimate (−14,448.23), with a Bayes factor equivalent to −299.07 relative to

the default, reference model (*i.e.*, Model 7; Table S1). Using the framework of Kass & Raftery

(1995), the Bayes factor in support for Model 6 is "decisive" compared to the reference model.

The final tree derived from the parameters used for Model 6 (combined posterior effective sample

size = 1,085; cladogram shown in Figure 2, phylogram with scaled branch lengths in Supplemental

Figure S2) has two primary sub-clades, which reflect the two carapace morphologies: the larger

clade contains the nine taxa with mostly domed carapaces, whilst the smaller clade contains the

four taxa with mostly saddleback carapaces. Most nodes (82%) are highly supported, with

posterior probabilities ≥ 0.95 (Figure 2).

<u>Species delimitation models in BPP</u>

When we ran an "unguided" species delimitation (A11 analysis in BPP) with our most

realistic prior of $\theta \sim$ IG(3, 0.001), the analysis supported 13 species with P>0.98 when using 50,

200, 500, or 1000 phased loci across all runs (Supplemental Table S3). When a smaller θ prior was

used, 13 species were almost always supported with P>0.99 (Supplemental Table S3). When using

a slightly larger θ prior, the analysis generally supported 12 or 13 species when 200, 500, and 1000

loci were used, but as few as 9 species when only 50 loci were used (Supplemental Table S3). In

cases where there was support for fewer than 13 species, the analysis typically supported the

collapse of two or more taxa on Isabela Island into one species. In four out of 45 runs, the A11

analysis supported a single species of Galapagos giant tortoise (Supplemental Tables S3 and S5),

but these runs likely represent poor mixing (see Discussion). When the Chaco tortoise outgroup

was included, most runs supported more than 10 species (Supplemental Table S4). Importantly, a

two-species model (with Galapagos giant tortoises as one species and the Chaco tortoise as another

species) was never supported. However, runs with the Chaco tortoise did not always converge on

the same distributions of posterior support when started with different random seeds

(Supplemental Table S4). Results were nearly identical when we used the unphased loci

(Supplemental Tables S5-S6) and running the "A10" analysis with the guide tree (Supplemental

Tables S7-S8). Using our most realistic priors, therefore, the BPP analyses supported delimitation

of 13 species (Figure 2).

gdi calculation from BPP parameter estimates

Overall, the median _gdi_ estimated in BPP ranged from 0.023 (_phantasticus_) to 0.534

(_hoodensis_; Figure 2, Table S11). None of the taxa exceeded a _gdi_ of 0.7, which has been proposed

as a heuristic for strongly delimited species. Five taxa (_becki (PBR)_, _donfaustoi_, _hoodensis_,

duncanensis, and *chathamensis*) had *gdi* in the ambiguous delimitation range of $0.2 < gdi < 0.7$.

The remaining eight taxa had *gdi* below 0.2. Estimates of *gdi* were effectively identical across

priors (Supplemental Tables S12–S13). In many cases, the *gdi* differed dramatically between sister

taxa, emphasizing the asymmetry commonly observed in this statistic.

When taxa do not meet the heuristic *gdi* threshold for species delimitation, Leaché *et al.*

(2019) recommended progressively collapsing taxon pairs, rerunning the MCMC and calculating

the *gdi* of the new groups. If the collapsed taxon represents a better-supported species, the

expectation is that the *gdi* would increase. This occurred when the taxa on central and southern

Isabela Island were collapsed (*gdi*=0.226, Figure 3A and Supplemental Table S16) and when those

on Santa Cruz Island were collapsed (*gdi*=0.291, Figure 3C and Supplemental Table S14). In all

other cases, however, the collapsed taxa had *gdi* lower than the separated taxa (Figure 3B and 3D,

Supplemental Tables S14-S18). Overall, the *gdi* estimates provide ambiguous support for

delimiting most taxa on different islands as separate species but favor treating some populations

on the same island as conspecific (Figure 2).

<u>PHRAPL</u>

For the PHRAPL analysis we focus on results of pairwise comparisons for populations within the

two primary clades (Figure 2). For the saddleback clade, there are four lineages spread across four

separate islands and so all pairwise comparisons were included here. Within the domed clade,

Isabela and Santa Cruz islands host five and two lineages respectively, while Santiago has a single

living lineage. For this clade we have presented the results in Figure 2 as within Santa Cruz Island

for its two lineages, within central and southern Isabela Island for its four lineages that form a

clade and within the clade of Santiago and its sister taxa, *becki (PBR)* and *becki (PBL)* on northern

Isabela Island. Results for model wAIC and *gdi* values for all possible pairwise comparisons within

the saddleback and domed morphology clades are presented in Supplemental Table S20. Within

the saddleback clade, all pairwise comparisons supported a two-species model when wAIC values

were summed, supporting four species across this clade. The *gdi* values in this clade were generally

in the ambiguous range of 0.2<*gdi*<0.7, except for the *gdi* of *hoodensis* and *phantasticus*

(*gdi*=0.179; Figure 2 and Table S20). For the Santa Cruz Island tortoises (*porteri* and *donfaustoi*),

PHRAPL supported a two-species model with a wAIC > 0.9, but the *gdi* was 0.181 (Figure 2).

Similarly, within the central and southern Isabela Island clade there was full support for two-

species models in pairwise comparisons, yet mixed support from the *gdi* values across pairwise

comparisons of the four lineages. The pairwise comparisons between taxa on Santa Cruz and

Isabela Islands also all favored a two-species model, and all *gdi* estimates fell between 0.2 and 0.7

(Figure 2). Similarly, all comparisons between the Santiago lineage (*darwini*) and the taxa on

Isabela and Santa Cruz Islands strongly supported a two-species model, with five out of eight *gdi*

comparisons above 0.2. In total, the species delimitation models in PHRAPL support the

delimitation of 13 species, while the *gdi* estimates from PHRAPL suggest between 7 and 9 species,

depending on the interpretation of heuristic thresholds.

Discussion

Despite considerable progress in achieving a more unified concept of species (Mayden

1997; de Queiroz 1998, 2007; Hey 2006), the empirical application of such a concept presents

challenges, particularly in cases of recent and/or incomplete divergence (e.g., de Queiroz 2005a,

Carstens *et al*. 2013, Jackson *et al*. 2017). The Galapagos giant tortoises represent such a case.

Here, we build on decades of research on the molecular evolution of these tortoises and provide

genomic evidence that the Galapagos giant tortoise complex consists of multiple distinct species.

Our phylogenetic analysis using SNAPP supported the topology previously inferred from

genome-wide sequence data (Jensen *et al.* 2022), except for the relationships among the most

recently diverged taxa (*guntheri*, *vandenburghi* and *microphyes*) and the placement of the Santa

Cruz Island taxa. Given this strong support, we used this nuclear phylogeny as a guide tree in

subsequent analyses. Notably, the topology of these trees inferred from nuclear loci deviates in

some ways from mitochondrial trees for these taxa (Poulakakis *et al.* 2012, 2020), an incongruence

that highlights the complex and rapid evolutionary history of Galapagos giant tortoises.

Our first line of evidence for the distinctiveness of Galapagos giant tortoise species is the

model-based species delimitation analysis in BPP. This analysis supported 13 species of Galapagos

giant tortoises across different numbers of loci when our most realistic priors were applied. Across

different priors we observe stronger support for 13 species as more loci are used, which may

indicate that the greater information content from more loci provides better biological resolution.

Theory and simulations suggest that BPP will increasingly favor a two-species model when

hundreds or thousands of loci are used, even under high rates of migration (Leaché *et al.* 2019).

Under our most realistic prior set, however, even 50 loci strongly support the delimitation of 13

species both with and without a guide tree.

In cases where fewer than 13 species were supported, the model favored collapsing two or

more of the taxa in a clade of four taxa on Isabela Island (i.e., *guntheri*, *microphyes*, *vandenburghi*,

and *vicina*). These taxa are geographically adjacent to each other on the central and southern

volcanoes of Isabela Island, which suggests that recent divergence or gene flow is a factor. The

delimitation models sometimes collapsed taxa that were not sister taxa on our phylogeny,

emphasizing the finding from this and other studies that the phylogeny of taxa on Isabela Island is

complex (Poulakakis *et al.* 2012, Jensen *et al.* 2022) and likely needs genomic data from a larger

number of individuals to be resolved.

In four runs of our A11 analysis including only Galapagos giant tortoise individuals the

analysis supported a single species of Galapagos giant tortoise. These runs are likely the result of

poor mixing, which is a known issue in the reversible-jump Markov Chain Monte Carlo (rjMCMC)

algorithm of BPP (Yang and Rannala 2010, Flouri *et al.* 2018). Mixing can be especially

problematic when sampling the fully resolved or fully collapsed tree, where the algorithm can get

"stuck" (Giarla *et al.* 2014). That we never find support for a single Galapagos giant tortoise

species when the Chaco tortoise individual is added to the analysis nor when we provide a guide

tree further supports the conclusion that the single-species runs of the A11 analysis are the result

of poor mixing.

We also modeled species delimitation in PHRAPL, comparing support for models of each

taxonomic pair as one species, two species in isolation, and two species with gene flow. We found

two-species models favored in all cases (Figure 2, Supplemental Table S20). The migration rates

chosen accounted for migration occurring up to 800 generations ago and there was mixed support

across populations for the two-species model that allowed migration vs. no migration. Jackson *et*

al. (2017) described the process of model selection in PHRAPL to be less accurate for isolation

only models (here, Model 1) with recent divergence times ($t<2$). Given that we tailored the

divergence values to be <1 to suit the evolutionary time scale of the Galapagos giant tortoise

lineages, there may be some bias in results where an isolation-with-migration model is selected

over the isolation-only model. This is a limitation of PHRAPL in that it has difficulty

distinguishing incomplete lineage sorting from gene flow, and so will favor the isolation-with-

migration model over the model of recent isolation. Furthermore, PHRAPL is more accurate with

a larger dataset (Jackson *et al.* 2017). The sample size for each taxon in our data set is small (n=2–

3 individuals per taxon). Because PHRAPL uses subsampling of individuals to generate expected

tree topologies, having a larger sample size that better represents the whole population could help

disentangle the history of the unresolved populations. Current work is underway to increase the

number of reference sequences for extant populations.

The species delimitation model in BPP has been criticized as oversplitting species and

recovering population structure rather than species (Jackson *et al.* 2017), as has species

delimitation modeling in PHRAPL (Leaché *et al.* 2019). Some have argued that the *gdi* better

reflects species differences by explicitly incorporating information about divergence time and

either population size (when calculated from BPP parameter estimates) or migration rates (when

calculated in PHRAPL) (Jackson *et al.* 2017, Leaché *et al.* 2019). Using the *gdi* thresholds of

gdi<0.2 for single species and *gdi*>0.7 for two species, we found ambiguous support (0.2<*gdi*<0.7)

in the BPP *gdi* estimates for *becki (PBR)*, *donfaustoi*, *hoodensis*, *chathamensis*, and *duncanensis*.

All other taxa had a *gdi*<0.2, creating multiple scenarios in which one sister taxon had *gdi*>0.2 and

the other taxon had *gdi*<0.2. In PHRAPL, the *gdi* estimates for most taxa were also in the

ambiguous range of 0.2<*gdi*<0.7, with some taxon pairs falling below 0.2 (i.e., *phantasticus* and

hoodensis, *porteri* and *donfaustoi*, and some of the domed taxa on Isabela and Santiago Islands).

Although these thresholds were proposed in the literature from a small sample of species,

work in other natural populations has shown that well-accepted species can fall into the ambiguous

range of 0.2–0.7, including horned lizards (Leaché *et al.* 2021), penguins (Mays *et al.* 2019), and

flying lizards (Reilly *et al.* 2022). Furthermore, there appears to be significant variation in *gdi*

within clades and between them. For example, in a survey of bird species the median *gdi* was 0.346

(1st quartile: 0.012; 3rd quartile: 0.742), while the surveyed mammalian species had a median *gdi*

value of 0.799 (1st quartile: 0.716; 3rd quartile: 0.955) (Jackson *et al*. 2017). This suggests that

appropriate heuristic cutoffs for *gdi* may differ among taxa, given that divergence times,

population sizes and life histories differ substantially across these clades. Likewise, island

radiations may also have a different underlying distribution of *gdi* given their shallow divergence

times and propensity for rapid adaptation. Before a *gdi* heuristic can be confidently applied to

Galapagos giant tortoise species, more thorough surveys of *gdi* among testudine species and across

island radiations are needed.

When sister taxa are not species, Leaché *et al*. (2019) found that collapsing the taxa and

recalculating *gdi* results in a larger *gdi*, signaling better support for the resulting species. This

behavior of *gdi* can therefore be used to interpret the *gdi* without relying as heavily on heuristic

thresholds. When we replicate this iterative process in BPP with our taxon set, we find that this

pattern occurs when we collapse all central and southern Isabela Island taxa (i.e., *guntheri*,

microphyes, *vandenburghi*, *vicina*) and when we collapse the Santa Cruz taxa (*porteri* and

donfaustoi). All other collapses, however, result in estimates of *gdi* that are smaller rather than

larger than the *gdi* of one or both sister taxa (Figure 3).

Applying population genetic principles to the *gdi* equation reveals why this occurs.

Because *gdi* is calculated from the ratio of τ to θ, it will remain constant when these values change

proportionally (e.g., the *gdi* will be 0.33 when τ is 0.001 and θ is 0.005, and when τ is 0.002 and θ

is 0.01). When sister taxa are collapsed, both τ and θ will increase. By definition, in recent

radiations τ will increase only marginally at each successive node. On the other hand, θ will be

inflated when calculated from a highly structured sample (e.g., differentiated species) because the

number of segregating sites is expected to be higher in a combined sample. Thus, the lower *gdi*

values that we found for successive collapsing of taxa from different islands further supports the

idea that these taxa do not belong to a single species.

The analyses presented in this study represent the most thorough attempt to date to tackle

the question of species delimitation in Galapagos giant tortoises using genetic data. Several prior

attempts to delimit Galapagos giant tortoise species focused on population genetic clustering,

mitochondrial monophyly, and amount of sequence divergence (e.g., Caccone *et al.* 1999,

Poulakakis *et al.* 2015, Loire *et al*. 2013, Kehlmaier *et al*. 2021). While these types of information

can provide some evidence concerning species boundaries, they lack a strong foundation in

modern population genetic theory as applied to species delimitation.

One such study claimed that genomic differentiation did not exist among *becki*,

vandenburghi, and *porteri* (Loire *et al*. 2013), but a re-analysis of the data with appropriate filtering

showed clear genomic differentiation of the three taxa (see Supplemental Figure S6 in Gaughran

et al. 2018). Another study comparing only mitochondrial DNA data between Galapagos giant

tortoises and extinct Caribbean giant tortoises argued that the recent divergence times of Galapagos

giant tortoise taxa relative to other tortoise species disqualified them from species status

(Kehlmaier *et al.* 2021). This view of species delimitation, however, ignores the complex reality

of speciation (see Donoghue 1985, Mallet 1995, Hey 2006). In addition, it ignores the fact that

some clades, such as those that colonized islands or other new environments, may be subject to

different demographic and selective forces that are associated with adaptive radiations. Outside of

tortoises, such a view would require a drastic re-delimitation of recently radiated species, with a

disproportionate effect on small, endangered island populations.

To date, all published genetic evidence, including the data from Loire *et al*. (2013),

supports the idea that Galapagos tortoise taxa are genetically distinct populations. Given the

statistical power of genome-wide SNPs to accurately detect differentiation (Gaughran *et al.* 2018),

this pattern of genetic distinctiveness appears unlikely to be overturned. Moving forward, genomic

discussions of Galapagos giant tortoise taxonomy should recognize that the preponderance of

evidence supports several genetically distinct taxa. However, whether these genetically distinct

populations deserve to be considered species is a more nuanced question.

The unified species concept (de Queiroz 2005b, 2007) provides one way to understand this

question. It highlights the idea of an independently evolving metapopulation lineage as the

keystone property of every species concept, with other properties (e.g., morphological

distinctiveness, genetic divergence, reproductive isolation) representing lines of evidence that a

taxon is an independently evolving lineage. The unified species concept thereby provides a

conceptual framework for evaluating candidate species that are morphologically similar, recently

diverged, or continuing to hybridize. The evidence we present here, combined with decades of

work documenting the distinctiveness of these taxa across multiple axes, shows how the Galapagos

giant tortoise taxa meet many criteria discussed as important aspects of modern species concepts.

As described above, the diverse methods we apply here are not magic solutions to species

delimitation, and each is the subject of ongoing debate. Still, interpreting our results holistically

provides some clarification on species delimitation in Galapagos giant tortoises and highlights

some areas of the taxonomy that remain difficult to resolve with our current data. Importantly, our

results largely refute the single-species model of Galapagos giant tortoise taxonomy that was

recently adopted by the IUCN Turtle Taxonomy Working Group (Rhodin *et al.* 2021). Instead, our

modeling in BPP delimits at least 12 species and our modeling in PHRAPL delimits 13 species.

On the other hand, the *gdi* results from BPP suggest 9 species, while those from PHRAPL suggest

between 7 and 9 species (Figure 2). We find significant support for the species status of some taxa,

especially the cases in which there is a single taxon per island. On the other hand, we find mixed

support for taxa inhabiting the same island as different species (i.e., Isabela Island and Santa Cruz

Island). Although model selection in BPP and PHRAPL supports the delimitation of species within

islands, the pattern of *gdi* results suggests that Santa Cruz Island may be home to a single species

and that the taxa of central and southern Isabela Island may also be a single species. Future work,

incorporating more samples for each taxon, will likely resolve the ambiguous delimitations by

making clear if the populations on Santa Cruz Island and central/southern Isabela Island have been

most affected by constant migration, secondary contact after divergence, or other demographic

processes.

Fundamentally, the taxonomic designation of Galapagos giant tortoises is both a scientific

and a philosophical question, and one that is deserving of genuine debate in the literature. Still,

this specific debate must be informed by decades of broader debates on species definitions and

methods of species delimitation. The rich literature around this topic highlights the fact that

speciation is a complex process: it can proceed at different rates and under different circumstances,

and our own temporally-biased observations mean that we necessarily study taxa at different stages

of this process. Galapagos giant tortoise taxa are evidently at different stages of lineage separation

and divergence and therefore offer an exciting system in which to study both species boundaries

and the process of speciation.

Figure 1. Galapagos giant tortoises are found on seven islands and form at least 13 genetically distinct taxa. Purple shapes indicate current ranges of populations. Carapace morphology (domed = gray, saddleback = white) is shown for each population, with "intermediate" shell shape indicated with overlapping icons. Icons with an "X" indicate extinct tortoise populations. Isabela Island and Santa Cruz Island are both home to multiple genetically distinct, allopatric populations. Inset map shows the names of islands with recently extant Galapagos giant tortoise taxa, with geological ages of each island in million years below.

Figure 2. Multiple lines of evidence confirm the Galapagos giant tortoise species complex consists of multiple genetically divergent, independently evolving species. On the left, the cladogram generated from genome-wide SNPs in SNAPP, with branch posterior support values. Within-clade PHRAPL pairwise *gdi* values are indicated on the tree by colored lines connecting lineages. In the center, support for species delimitation models in PHRAPL and BPP. On the right, a boxplot of the *gdi* values estimated from the posterior distributions of τ and θ generated by BPP for each Galapagos giant tortoise taxon (red: **gdi** > 0.7, meeting the heuristic threshold for strongly delimited species; blue: **gdi** < 0.2, below the heuristic threshold for a single species; orange: 0.2 < *gdi* < 0.7, which is considered ambiguous species delimitation). Model-based species delimitation in BPP and PHRAPL model selection supported 13 species (not shown).

Figure 3. **Estimate _gdi_ for progressively collapsed taxa shows support for some delimited species but not others.** _gdi_ calculated for each taxon, including when sister taxa are collapsed following the method of Leaché _et al._ (2019). Blue boxes are single named taxa; green boxes are two or more taxa collapsed into a single "species." A) Isabela Island clade with progressive taxonomic collapse, showing a higher _gdi_ above 0.2 when the four taxa from central and southern Isabela Island are collapsed. B) Santiago Island and northern Isabela Island clade with progressive taxonomic collapse, showing lower _gdi_ when taxa are collapsed. C) Santa Cruz Island clade with progressive taxonomic collapse, showing a higher _gdi_ above 0.2 when the two taxa are collapsed. D) Saddleback clade with progressive taxonomic collapse, showing lower _gdi_ when taxa are collapsed.

References

Baele, G., Lemey, P., Bedford, T., Rambaut, A., Suchard, M. A., & Alekseyenko, A. V. (2012). Improving the accuracy of demographic and molecular clock model comparison while accommodating phylogenetic uncertainty. *Molecular Biology and Evolution*, *29*(9), 2157-2167.

Benavides, E., Baum, R., Snell, H. M., Snell, H. L., & Sites, Jr, J. W. (2009). Island biogeography of Galápagos lava lizards (Tropiduridae: Microlophus): species diversity and colonization of the archipelago. *Evolution*, *63*(6), 1606-1626.

Bouckaert, R., Heled, J., Kühnert, D., Vaughan, T., Wu, C. H., Xie, D., ... & Drummond, A. J. (2014). BEAST 2: a software platform for Bayesian evolutionary analysis. *PLoS Computational Biology*, *10*(4), e1003537.

Bryant, D., Bouckaert, R., Felsenstein, J., Rosenberg, N. A., & RoyChoudhury, A. (2012). Inferring species trees directly from biallelic genetic markers: bypassing gene trees in a full coalescent analysis. *Molecular Biology and Evolution*, *29*(8), 1917-1932.

Caccone, A., Gibbs, J. P., Ketmaier, V., Suatoni, E., & Powell, J. R. (1999). Origin and evolutionary relationships of giant Galápagos tortoises. *Proceedings of the National Academy of Sciences*, *96*(23), 13223-13228.

Caccone, A., Gentile, G., Gibbs, J. P., Fritts, T. H., Snell, H. L., Betts, J., & Powell, J. R. (2002). Phylogeography and history of giant Galápagos tortoises. *Evolution*, *56*(10), 2052-2066.

Carstens, B. C., Pelletier, T. A., Reid, N. M., & Satler, J. D. (2013). How to fail at species delimitation. *Molecular Ecology*, *22*(17), 4369-4383.

Chiari, Y. (2021). Morphology. In J. P. Gibbs, L. Cayot, and W. Tapia A (eds.). *Galapagos Giant Tortoises*. Academic Press.

de Queiroz, K. (1998). The general lineage concept of species, species criteria, and the process of

speciation. In D. J. Howard and S. H. Berlocher (eds). *Endless forms: species and*

speciation. Oxford University Press

de Queiroz, K. (2005a). Different species problems and their resolution. *BioEssays*, *27*(12),

1263-1269.

de Queiroz, K. (2005b). A unified concept of species and its consequences for the future of

taxonomy. *Proceedings of the California Academy of Sciences*.

de Queiroz, K. (2007). Species concepts and species delimitation. *Systematic Biology*, *56*(6),

879-886.

Flouri, T., Jiao, X., Rannala, B., & Yang, Z. (2018). Species tree inference with BPP using genomic

sequences and the multispecies coalescent. *Molecular Biology and Evolution*, *35*(10),

2585-2593.

Garrick, R. C., Kajdacsi, B., Russello, M. A., Benavides, E., Hyseni, C., Gibbs, J. P., ... & Caccone,

A. (2015). Naturally rare versus newly rare: demographic inferences on two timescales

inform conservation of Galápagos giant tortoises. *Ecology and Evolution*, *5*(3), 676-694.

Gaughran, S. J., Quinzin, M. C., Miller, J. M., Garrick, R. C., Edwards, D. L., Russello, M. A., ...

& Caccone, A. (2018). Theory, practice, and conservation in the age of genomics: The

Galápagos giant tortoise as a case study. *Evolutionary Applications*, *11*(7), 1084-1093.

Geist, D. J., H. Snell, H. Snell, C. Goddard, and M. D. Kurz. (2014). A paleogeographic model

of the Galápagos islands and biogeographical and evolutionary implications. Pp. 145-166

in K. S. Harpp, E. Mittelstaedt, N. d'Ozouville, and D. W. Graham (eds). *The Galápagos:*

a natural laboratory for the earth science. John Wiley & Sons, Inc.

Giarla, T. C., Voss, R. S., & Jansa, S. A. (2014). Hidden diversity in the Andes: comparison of

species delimitation methods in montane marsupials. *Molecular Phylogenetics and*

Evolution, *70*, 137-151.

Hey, J. (2006). On the failure of modern species concepts. *Trends in Ecology & Evolution*, *21*(8),

447-450.

Hunter, E. A., Gibbs, J. P., Cayot, L. J., & Tapia, W. (2013). Equivalency of Galápagos giant

tortoises used as ecological replacement species to restore ecosystem

functions. *Conservation Biology*, *27*(4), 701-709.

Jackson, N. D., Carstens, B. C., Morales, A. E., & O'Meara, B. C. (2017). Species delimitation

with gene flow. *Systematic Biology*, *66*(5), 799-812.

Jensen, E. L., Gaughran, S. J., Garrick, R. C., Russello, M. A., & Caccone, A. (2021).

Demographic history and patterns of molecular evolution from whole genome sequencing

in the radiation of Galapagos giant tortoises. *Molecular Ecology*, *30*(23), 6325-6339.

Jensen, E. L., Gaughran, S. J., Fusco, N. A., Poulakakis, N., Tapia, W., Sevilla, C., ... &

Caccone, A. (2022). The Galapagos giant tortoise *Chelonoidis phantasticus* is not extinct.

Communications Biology, *5*(1), 546.

Kass, R. E., & Raftery, A. E. (1995). Bayes factors. *Journal of the American Statistical Association*,

90(430), 773-795.

Kehlmaier, C., Albury, N. A., Steadman, D. W., Graciá, E., Franz, R., & Fritz, U. (2021).

Ancient mitogenomics elucidates diversity of extinct West Indian tortoises. *Scientific*

Reports, *11*(1), 3224.

Kimura, M., & Ohta, T. (1969). The average number of generations until fixation of a mutant

gene in a finite population. *Genetics*, *61*(3), 763.

Leaché, A. D., Zhu, T., Rannala, B., & Yang, Z. (2019). The spectre of too many species.

Systematic Biology, *68*(1), 168-181.

Leaché, A. D., Davis, H. R., Singhal, S., Fujita, M. K., Lahti, M. E., & Zamudio, K. R. (2021).

Phylogenomic assessment of biodiversity using a reference-based taxonomy: an example

with Horned Lizards (Phrynosoma). *Frontiers in Ecology and Evolution*, *9*, 678110.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., ... & 1000 Genome Project

Data Processing Subgroup. (2009). The sequence alignment/map format and SAMtools.

Bioinformatics, *25*(16), 2078-2079.

Loire, E., Chiari, Y., Bernard, A., Cahais, V., Romiguier, J., Nabholz, B., ... & Galtier, N.

(2013). Population genomics of the endangered giant Galápagos tortoise. *Genome*

Biology, *14*, 1-11.

Mallet, J. (1995). A species definition for the modern synthesis. *Trends in Ecology & Evolution*,

10(7), 294-299.

Mays Jr, H. L., Oehler, D. A., Morrison, K. W., Morales, A. E., Lycans, A., Perdue, J., ... &

Weakley, L. A. (2019). Phylogeography, population structure, and species delimitation in

rockhopper penguins (*Eudyptes chrysocome* and *Eudyptes moseleyi*). *Journal of*

Heredity, *110*(7), 801-817.

Mayden, R. L. (1997). A hierarchy of species concepts: the denouement in the saga of the

species problem. pp. 381–423 in M. F. Claridge, H. A. Dawah & M. R. Wilson (eds.).

Species: the Units of Diversity. Chapman & Hall.

Parent, C. E., Caccone, A., & Petren, K. (2008). Colonization and diversification of Galápagos

terrestrial fauna: a phylogenetic and biogeographical synthesis. *Philosophical*

Transactions of the Royal Society B: Biological Sciences, *363*(1508), 3347-3361.

Poulakakis, N., Glaberman, S., Russello, M., Beheregaray, L. B., Ciofi, C., Powell, J. R., &

Caccone, A. (2008). Historical DNA analysis reveals living descendants of an extinct

species of Galápagos tortoise. *Proceedings of the National Academy of Sciences*, *105*(40),

15464-15469.

Poulakakis, N., Russello, M., Geist, D., & Caccone, A. (2012). Unravelling the peculiarities of

island life: vicariance, dispersal and the diversification of the extinct and extant giant

Galápagos tortoises. *Molecular Ecology*, *21*(1), 160-173.

Poulakakis, N., Edwards, D. L., Chiari, Y., Garrick, R. C., Russello, M. A., Benavides, E., ... &

Caccone, A. (2015). Description of a new Galápagos giant tortoise species (Chelonoidis;

Testudines: Testudinidae) from Cerro Fatal on Santa Cruz Island. *PLoS One*, *10*(10),

e0138779.

Poulakakis, N., Miller, J. M., Jensen, E. L., Beheregaray, L. B., Russello, M. A., Glaberman, S.,

... & Caccone, A. (2020). Colonization history of Galapagos giant tortoises: Insights from

mitogenomes support the progression rule. *Journal of Zoological Systematics and*

Evolutionary Research, *58*(4), 1262-1275.

Pritchard, P. C. (1996). *The Galápagos tortoises: nomenclature and survival status*. Chelonian

Research Foundation.

Quesada, V., Freitas-Rodríguez, S., Miller, J., Pérez-Silva, J. G., Jiang, Z. F., Tapia, W., ... &

López-Otín, C. (2019). Giant tortoise genomes provide insights into longevity and age-

related disease. *Nature ecology & evolution*, *3*(1), 87-95.

R Core Team (2022). R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Reilly, S. B., Stubbs, A. L., Arida, E., Karin, B. R., Arifin, U., Kaiser, H., ... & McGuire, J. A.

(2022). Phylogenomic analysis reveals dispersal-driven speciation and divergence with

gene flow in Lesser Sunda flying lizards (Genus *Draco*). *Systematic Biology*, *71*(1), 221-

241.

Rhodin, A.G.J., Iverson, J.B., Bour, R., Fritz, U., Georges, A., Shaffer, H.B., and van Dijk, P.P.

(2021). Turtles of the World: Annotated Checklist and Atlas of Taxonomy, Synonymy,

Distribution, and Conservation Status (9th Ed.). In Rhodin, A.G.J., Iverson, J.B., van Dijk,

P.P., Stanford, C.B., Goode, E.V., Buhlmann, K.A., and Mittermeier, R.A. (Eds.).

Conservation Biology of Freshwater Turtles and Tortoises: A Compilation Project of the

IUCN/SSC Tortoise and Freshwater Turtle Specialist Group. *Chelonian Research*

Monographs 8:1–472. doi:10.3854/crm.8.checklist.atlas.v9.2021.

Ruffley, M., Smith, M. L., Espíndola, A., Carstens, B. C., Sullivan, J., & Tank, D. C. (2018).

Combining allele frequency and tree-based approaches improves phylogeographic

inference from natural history collections. *Molecular Ecology*, *27*(4), 1012-1024.

Russello, M. A., Glaberman, S., Gibbs, J. P., Marquez, C., Powell, J. R., & Caccone, A. (2005).

A cryptic taxon of Galápagos tortoise in conservation peril. *Biology Letters*, *1*(3), 287-

290.

Schafer, S. F., & Krekorian, C. N. (1983). Agonistic behavior of the Galapagos tortoise,

Geochelone elephantopus, with emphasis on its relationship to saddle-backed shell shape.

Herpetologica, 448-456.

Smith, M. L., & Carstens, B. C. (2020). Process-based species delimitation leads to identification

of more biologically relevant species. *Evolution*, *74*(2), 216-229.

Stamatakis A. (2014). RAxML version 8: a tool for phylogenetic analysis and post-analysis of

large phylogenies. *Bioinformatics.* 30(9):1312-3

Yang, Z., & Rannala, B. (2010). Bayesian species delimitation using multilocus sequence

data. *Proceedings of the National Academy of Sciences*, *107*(20), 9264-9269.

Acknowledgements

We thank the Galápagos National Park Directorate for their collaboration. This work was

supported by the Galapagos Conservancy (AWD0008098), RE:wild (AWD0008746), the Turtle

Conservancy, and the Oak Foundation. SJG was supported by the National Science Foundation

under Grant No. 2010918. We are grateful for input from Adam Leaché, Tomas Flouri, Ariadna

Morales, and Bridgett vonHoldt in the development of this manuscript. This research made use of

the Rocket High Performance Computing service at Newcastle University, the Yale Center for

Research Computing, and Princeton Research Computing.

Author Contributions

SJG, RG, AC, and ELJ designed the study. SJG, RG, AO, MJ, AC, and ELJ carried out the methodology. SJG, RG, MJ, NF, AO, JMM, NP, KdQ, and ELJ contributed to the investigation. SJG, RG, AO, MJ, NF, and ELJ visualized the results. SJG, AC, and ELJ acquired funding for the work. Project administration and supervision was provided by SJG, AC, and ELJ. SJG, RG, AO, MJ, AC, and ELJ wrote the original draft, and all authors contributed to reviewing and editing subsequent drafts.

Data Accessibility

The whole genome resequencing data from Jensen *et al*. (2022) is available under NCBI BioProject PRJNA761229. Code for running BPP and PHRAPL, as well as alignments for loci used in these analyses, are available at: https://github.com/sjgaughran/tortoise-species-delimitation. Output files from SNAPP and BPP are available at: https://doi.org/10.5061/dryad.63xsj3v84

Conflict of Interest

The authors declare no conflicts of interest.

Supplemental Methods

<u>Terminology, study design and data</u>

We use several terms to refer to the demographic and evolutionary units of tortoises living

in the Galapagos. We use the term *population* in the evolutionary genetics sense of a group of

interbreeding individuals existing together in time and space (e.g., Hedrick 2009). We use the term

lineage in a slightly broader sense, to expand the time component of *population*, thereby including

ancestor-descendant relationships. We use the systematic term *taxon* to refer broadly to an

evolutionary unit (e.g., lineage) without specifying whether that unit is a species or a subspecies.

Because our goal is to evaluate the status of Galapagos giant tortoise taxa as species, we avoid

using the terms *species* and *subspecies* as *a priori* descriptors. To maintain this agnosticism, we

use only the epithet to refer to each taxon (e.g., *phantasticus* rather than *Chelonoidis phantasticus*

or *Chelonoidis niger phantasticus*).

We designed a whole-genome study of population distinctiveness and species delimitation,

taking advantage of existing illumina short-read whole genome resequencing data (Jensen *et al.*

2021, Jensen *et al*. 2022), which are available under NCBI Bioproject PRJNA761229. The samples

in our data set represent all extant taxa, which have previously been classified as populations,

species, or subspecies. The samples were originally selected for sequencing as representatives of

the genetic and geographic populations (Figure 1) that have previously been shown to exist using

microsatellite (Ciofi *et al.* 2002, Ciofi *et al.* 2006) and reduced representation SNP (Miller *et al.*

2018) data. Previous genetic research suggests that the *becki* taxon consists of two geographically

and genetically distinct populations, referred to in the literature as "PBL" (Piedras Blancas) and

"PBR" (Puerto Bravo) (Garrick *et al.* 2014). A prior phylogeny constructed from these genome

data in Jensen *et al*. (2022) showed the named taxa to be monophyletic, with the exception of

becki. Within *becki*, PBR individuals formed a clade sister to *darwini* and PBL individuals did not

form a clade, with one sample closest to PBR and *darwini* and the other two closest to other taxa

from Isabela. We have therefore chosen to test *becki (PBL)* and *becki (PBR)* as potentially distinct

taxa, and have retained the potentially admixed PBL individual to avoid biasing our analyses.

We mapped these reads to a reference genome for *abingdonii* (Quesada et al. 2019) using

bwa-mem (Li 2013). Jensen *et al*. (2021) previously investigated the potential for reference bias

in using the *abingdonii* assembly and found no evidence for such a bias. We called SNPs using the

BCFtools variant calling pipeline (Li et al. 2009). We then filtered the VCF for bi-allelic SNPs

that had a minimum GQ of 25, a minimum map quality score of 25, and at least 2 reads for each

allele in the genotype. We then created a consensus fasta file for each individual using BCFtools

and masked each consensus sequence for all missing genotypes from the VCF to ensure that

missing data were not erroneously assigned to the reference allele. We also masked every fasta

with a mask of repetitive regions downloaded from the UCSC Genome Browser and with a

mappable regions mask generated by Jensen *et al*. (2021). From these filtered and masked fastas,

we generated a random set of 1kb loci separated by 100kb. We retained loci that had less than 10%

missing data, GC content between 30% and 70%, and at least one variable site. Because BPP

requires phased data when using more than ~100 loci, we phased each locus using PHASE v. 2.1.1

(Stephens *et al*. 2001, Stephens and Scheet 2005). We then used several methods to delimit species

under different phylogenetic and coalescent models.

<u>SNAPP Tree Generation</u>

We used SNAPP v1.5.2 (Bryant *et al.* 2012) and MODEL_SELECTION v1.5.3 (Baele *et*

al. 2012), as implemented in BEAST2 v2.6.7 (Bouckaert *et al.* 2014), to generate a guide tree

reflecting the phylogenetic relationships among the extant populations of Galapagos giant tortoises

for downstream species delimitation comparisons. We first tested 21 different phylogenetic

models on a dataset comprising 1,000 random, unlinked SNPs—and corresponding genotypes—

from 39 samples. We obtained this dataset from a previous study conducted by Jensen et al. (2022)

and it included one Chaco tortoise (*C. chilensis*) sample to serve as an outgroup to the Galapagos

giant tortoises. We ensured the Galapagos giant tortoises presented genetic variation across loci

and we allowed no missing data.

We grouped the samples into populations, assumed forward and backward mutation rates

equivalent to the unit ($u = v = 1$), and used combinations of shape ($\alpha = 5, 12, 30$), scale ($\beta = 50,$

$60, 70, 80, 110, 150$), and speciation rate ($\lambda = 0.01, 10, 39$) parameters as priors to constitute the

models to be tested (see Table S1). Path sampling runs for each model consisted of 24 steps,

100,000 MCMC generations sampled every 100 generations, and a 50% burn-in. We compared

the resulting marginal likelihood estimates from each run and selected the best model via Bayes

factor delimitation (Kass & Raftery 1995) after using the default model ($\alpha = 12, \beta = 110, \lambda = 0.01$)

as a reference.

For the best model, we then used an additional set of 4,000 random, unlinked biallelic SNPs

(5,000 SNPs total) for final phylogenetic reconstruction. For this dataset, we performed four

independent runs consisting of 2,000,000 MCMC generations sampled every 1,000 generations

and a 10% burn-in. We used Tracer v1.7.2 (Rambaut *et al.* 2018) to assess statistical convergence

across runs and LogCombiner v2.6.7 (Drummond & Rambaut 2007) to summarize the posterior

trees.

<u>Divergence time from BPP</u>

We can calculate the absolute divergence time in generations from τ, the rate-scaled

divergence time in BPP, as:

$T = \tau/\mu$

Where μ is the per-generation mutation rate. Although the per-generation mutation rate in

Galapagos giant tortoises is unknown, we use 6×10^{-9} reflecting the *de novo* mutation rate recently

measured in the painted turtle (Bergeron *et al*. 2023).

831   **Supplemental Figures and Tables**

Model 1: Two species, no gene flow   **Model 2:** Two species, constant gene flow   **Model 3:** One species, constant gene flow

Population 1   Population 2   Population 1   Population 2   Population 1   Population 2

Figure S1. The three models used to test one and two species models under differing gene flow

scenarios in PHRAPL. The grey arrow indicates bi-directional gene flow between the two

populations.

Figure S2. Phylogeny with scaled branch lengths generated from 5,000 genome-wide SNPs in

SNAPP from the best supported model ($\alpha = 5$, $\beta = 150$, $\lambda = 39$) with branch posterior support

values indicated. The scale bar indicates the number of nucleotide substitutions per site.

Figure S3. Galapagos giant tortoise species ranges when the Santa Cruz Island taxa (*porteri* and *donfaustoi*) are collapsed and the central and southern Isabela Island (*microphyes*, *vandenburghi*, *guntheri*, and *vicina*) are collapsed. Purple shapes indicate current ranges of populations. Blue dotted lines surround taxa that are collapsed into a single species according to some of our analyses. Carapace morphology (domed = gray, saddleback = white) is shown for each population, with "intermediate" shell shape indicated with overlapping icons. Icons with an "X" indicate extinct tortoise populations. Inset map shows the names of islands with recently extant Galapagos giant tortoise taxa, with geological ages of each island in million years below.

861 **Table S1.** Summary statistics from 21 different phylogenetic models tested on 1,000 random,

862 unlinked biallelic SNPs—and corresponding genotypes—from 39 tortoises arranged into

863 populations. Each run consisted of 24 steps, 100,000 MCMC generations sampled every 100

864 generations, and a 50% burn-in. The default, reference model for Bayes factor delimitation is

865 shown with an asterisk (*).

Model	α	β	λ	Effective Sample Size	Marginal Likelihood	Rank	Bayes Factor
1	5	50	0.01	1,330.47	−14,556.42	8	−82.70
2	5	50	10	1,994.26	−14,471.91	4	−251.73
3	5	50	39	1,204.93	−14,456.56	2	−282.42
4	5	150	0.01	1,816.32	−14,552.45	7	−90.64
5	5	150	10	1,808.11	−14,463.95	3	−267.65
6	5	150	39	1,574.37	−14,448.23	1	−299.07
7*	12	110	0.01	1,749.29	−14,597.77	9	−
8	12	110	10	2,007.71	−14,509.09	6	−177.35
9	12	110	39	1,430.58	−14,496.23	5	−203.09
10	30	60	0.01	2,476.20	−15,036.56	21	+877.57
11	30	60	10	2,939.10	−14,929.72	18	+663.89
12	30	60	39	2,235.82	−14,917.57	17	+639.60
13	30	70	0.01	2,597.08	−14,985.58	20	+775.62
14	30	70	10	2,258.89	−14,882.03	16	+568.52
15	30	70	39	2,420.92	−14,872.60	15	+549.67
16	30	80	0.01	2,347.20	−14,941.95	19	+688.35
17	30	80	10	2,904.94	−14,841.68	14	+487.81
18	30	80	39	2,179.97	−14,831.56	13	+467.57
19	30	150	0.01	2,060.26	−14,764.49	12	+333.44
20	30	150	10	1,920.77	−14,675.53	11	+155.51
21	30	150	39	2,119.72	−14,663.75	10	+131.95

871   **Table S2.** Parameter values for migration (*m*) and divergence (*t*) used for the grid search within

872   PHRAPL. The maximum divergence time is equivalent to 1Ma.

m	0.005	0.02	0.08	0.12	0.25	0.4	0.8
t	0.01	0.05	0.1	0.25	0.5	0.75	1

892   **Table S3.** Posterior output from the A11 analysis species delimitation model in BPP using phased

893   loci from the 38 Galapagos giant tortoise individuals. Support for collapsed taxa as species is

894   included in the final column. The "Medium θ" prior is the most realistic prior.

θ prior	No. loci	Seed	Posterior for number of species	Posteriors for collapsed taxonomic groups or notes
Small θ	50 loci	Seed 1	P[13]=0.997	
		Seed 2	P[13]=0.998	
		Seed 3	P[13]=0.999	
	200 loci	Seed 1	P[13]=1.000	
		Seed 2	P[13]=1.000	
		Seed 3	P[13]=1.000	
	500 loci	Seed 1	P[13]=1.000	
		Seed 2	P[13]=1.000	
		Seed 3	P[1]=1.000	single_species=1.000
	1000 loci	Seed 1	P[7]=0.162 P[6]=0.079 P[1]=0.752	single_species=0.752 *darwini+beck*(PBR)=0.248 *porteri+donfaustoi*=0.243 *phantasticus+hoodensis*=0.243 *guntheri+vandenburghi+* *microphyes+vicina*=0.162 *guntheri+vandenburghi+* *microphyes+vicina+* *becki (PBL)*=0.162
		Seed 2	P[13]=1.000	
		Seed 3	P[13]=1.000	
Medium θ	50 loci	Seed 1	P[13]=0.988 P[12]=0.012	*guntheri+vandenburghi*=0.010
		Seed 2	P[13]=0.991	
		Seed 3	P[13]=0.992	
	200 loci	Seed 1	P[13]=1.000	
		Seed 2	P[13]=1.000	
		Seed 3	P[13]=1.000	
	500 loci	Seed 1	P[13]=1.000	
		Seed 2	P[13]=1.000	
		Seed 3	P[13]=1.000	
	1000 loci	Seed 1	P[13]=1.000	
		Seed 2	P[13]=1.000	
		Seed 3	P[13]=1.000	
Large θ	50 loci	Seed 1	P[13]=0.282	*guntheri+vandenburghi*=0.259

			P[12]=0.413 P[11]=0.261	*becki (PBR)+darwini*=0.175 *guntheri+microphyes*=0.161 *vandenburghi+vicina*=0.079 *guntheri+vandenburghi+ microphyes*=0.077 *porteri+donfaustoi*=0.068 *guntheri+vandenburghi+ vicina*=0.068 *guntheri+vicina*=0.023
		Seed 2	P[13]=0.192 P[12]=0.304 P[11]=0.297 P[10]=0.155 P[9]=0.048	*guntheri+vandenburghi*=0.197 *becki (PBR)+darwini*=0.176 *guntheri+vandenburghi+ microphyes*=0.161 *guntheri+vandenburghi+ microphyes+vicina*=0.148 *guntheri+microphyes*=0.112 *porteri+donfaustoi*=0.094 *vicina+vandenburghi*=0.058 *vicina+vandenburghi+ guntheri*=0.052 *vicina+guntheri*=0.036
		Seed 3	P[13]=0.060 P[12]=0.118 P[11]=0.146 P[10]=0.541 P[9]=0.127	*guntheri+vandenburghi+ microphyes+vicina*=0.632 *becki (PBR)+darwini*=0.160 *guntheri+microphyes*=0.079 *guntheri+vandenburghi+ microphyes*=0.077 *porteri+donfaustoi*=0.073 *guntheri+vandenburghi*=0.064 *vicina+vandenburghi*=0.038 *vicina+vandenburghi+ guntheri*=0.032 *vicina+guntheri*=0.010
	200 loci	Seed 1	P[13]=0.924 P[12]=0.076	*guntheri+vandenburghi*=0.073
		Seed 2	P[13]=0.897 P[12]=0.103	*guntheri+vandenburghi*=0.103
		Seed 3	P[13]=0.957 P[12]]=0.043	*guntheri+vandenburghi*=0.043
	500 loci	Seed 1	P[13]=0.258 P[12]=0.484 P[11]=0.053 P[10]=0.204	*guntheri+microphyes*=0.524 *guntheri+microphyes+ vandenburghi+vicina*=0.204 *vandenburghi+vicina*=0.040 *guntheri+microphyes+ vandenburghi*=0.012
		Seed 2	P[13]=1.000	

		Seed 3	P[1]=1.000	single_species=1.000
	1000 loci	Seed 1	P[13]=1.000	
		Seed 2	P[13]=1.000	
		Seed 3	P[11]=0.710 P[10]=0.290	*guntheri+microphyes+ vandenburghi* =0.710 *guntheri+microphyes+ vandenburghi+vicina*=0.290

915 **Table S4.** Posterior output from the A11 analysis species delimitation model in BPP using phased

916 loci from the 38 Galapagos giant tortoise individuals and the Chaco tortoise individual. Support

917 for collapsed taxa as species is included in the final column. The "Medium θ" prior is the most

918 realistic prior.

θ prior	No. loci	Seed	Posterior for number of species	Posteriors for collapsed taxonomic groups or notes
Small θ	50 loci	Seed 1	P[12]=0.993	*guntheri+microphyes+ vandenburghi*=0.993
		Seed 2	P[13]=0.273 P[12]=0.716	*guntheri+microphyes+ vandenburghi*=0.716 *guntheri + vandenburghi*=0.270
		Seed 3	P[12]=0.949 P[11]=0.051	*guntheri+microphyes+ vandenburghi*=0.995 *phantasticus+becki (PBL)*=0.045
	200 loci	Seed 1	P[14]=0.809 P[12]=0.187	*guntheri+microphyes+ vandenburghi*=0.187
		Seed 2	P[14]=1.000	
		Seed 3	P[12]=0.790 P[11]=0.210	*guntheri+microphyes+ vandenburghi*=0.790 *guntheri+microphyes+ vandenburghi+vicina*=0.210
	500 loci	Seed 1	P[14]=1.000	
		Seed 2	P[14]=0.723 P[12]=0.121 P[11]=0.025 P[10]=0.025 P[2]=0.095	all_single_species=0.095 *guntheri+vandenburghi+ microphyes*=0.121 *guntheri+vandenburghi+ microphyes+vicina*=0.051 *porteri+donfaustoi*=0.026
		Seed 3	P[14]=1.000	
	830 loci	Seed 1	P[14]=1.000	
		Seed 2	P[14]=1.000	
		Seed 3	P[14]=1.000	
Medium θ	50 loci	Seed 1	P[14]=0.018 P[13]=0.335 P[12]=0.646	*guntheri+microphyes+ vandenburghi*=0.351 *guntheri+vandenburghi*=0.316 *guntheri+vandenburghi+ vicina*=0.289 *vandenburghi+vicina*=0.018
		Seed 2	P[14]=0.012	*guntheri+microphyes+*

			P[13]=0.159 P[12]=0.785 P[11]=0.045	*vandenburghi*=0.537 *guntheri+vandenburghi+* *vicina*=0.226 *vandenburghi+vicina*=0.079 *guntheri+vandenburghi*=0.063 *guntheri+microphyes+* *vandenburghi+vicina*=0.045 *guntheri+microphyes*=0.041 *microphyes+vandenburghi*=0.012
		Seed 3	P[14]=0.016 P[13]=0.216 P[12]=0.767	*guntheri+microphyes+* *vandenburghi*=0.717 *guntheri+vandenburghi*=0.177 *guntheri+vandenburghi+* *vicina*=0.051 *guntheri+vandenburghi*=0.040
	200 loci	Seed 1	P[14]=0.992	
		Seed 2	P[14]=0.961 P[13]=0.039	*guntheri+vandenburghi*=0.039
		Seed 3	P[14]=0.524 P[13]=0.076 P[12]=0.352 P[10]=0.042	*guntheri+microphyes+* *vandenburghi*=0.352 *guntheri+vandenburghi*=0.072 *guntheri+microphyes+* *vandenburghi+vicina*=0.048 *porteri+donfaustoi*=0.042
	500 loci	Seed 1	P[14]=0.810 P[13]=0.017 P[12]=0.053 P[11]=0.056 P[9]=0.014 P[8]=0.048	*guntheri+microphyes+* *vandenburghi+vicina+* *becki (PBL)*=0.064 *porteri+donfaustoi*=0.062 *guntheri+microphyes+* *vandenburghi+vicina*=0.056 *guntheri+microphyes+* *vandenburghi*=0.053 *darwini+becki (PBR)*=0.048 *guntheri+microphyes*=0.017
		Seed 2	P[14]=0.892 P[13]=0.025 P[12]=0.050 P[11]=0.033	*guntheri+microphyes+* *vandenburghi*=0.050 *guntheri+microphyes+* *vandenburghi+vicina*=0.033 *guntheri+microphyes*=0.025
		Seed 3	P[14]=1.000	
	830 loci	Seed 1	P[14]=1.000	
		Seed 2	P[14]=1.000	
		Seed 3	P[14]=1.000	
Large θ	50 loci	Seed 1	P[12]=0.044 P[11]=0.388 P[10]=0.549	*guntheri+microphyes+* *vandenburghi+vicina*=0.896 *porteri+donfaustoi*=0.607

			P[9]=0.022	*guntheri+microphyes+ vandenburghi*=0.098 *becki (PBL)+becki (PBR)*=0.031
		Seed 2	P[12]=0.081 P[11]=0.442 P[10]=0.464 P[9]=0.011	*guntheri+microphyes+ vandenburghi+vicina*=0.671 *porteri+donfaustoi*=0.690 *guntheri+microphyes+ vandenburghi*=0.285 *becki (PBL)+becki (PBR)*=0.042 *guntheri+vicina+ vandenburghi*=0.023 *guntheri+microphyes*=0.020 *vandenburghi+*vicina=0.017
		Seed 3	P[12]=0.013 P[11]=0.342 P[10]=0.613 P[9]=0.032	*guntheri+microphyes+ vandenburghi+vicina*=0.872 *porteri+donfaustoi*=0.737 *guntheri+microphyes+ vandenburghi*=0.111 *becki (PBL)+becki (PBR)*=0.048 *microphyes+guntheri*=0.010
	200 loci	Seed 1	P[11]=0.683 P[10]=0.248 P[9]=0.070	*guntheri+microphyes+ vandenburghi+vicina*=1.000 *porteri+donfaustoi*=0.317 *darwini+becki (PBR)*=0.070
		Seed 2	P[12]=0.690 P[11]=0.253 P[10]=0.057	*guntheri+microphyes+ vandenburghi+vicina*=0.310 *guntheri+microphyes+ vandenburghi* =0.690 *porteri+donfaustoi*=0.057
		Seed 3	P[11]=1.000	*guntheri+microphyes+ vandenburghi+vicina*=1.000
	500 loci	Seed 1	P[14]=1.000	
		Seed 2	P[11]=1.000	*guntheri+microphyes+ vandenburghi+vicina*=1.000
		Seed 3	P[11]=1.000	*guntheri+microphyes+ vandenburghi+vicina*=1.000
	830 loci	Seed 1	P[11]=1.000	*guntheri+microphyes+ vandenburghi+vicina*=1.000
		Seed 2	P[14]=1.000	
		Seed 3	P[11]=1.000	*guntheri+microphyes+ vandenburghi+vicina*=1.000

922   **Table S5.** Posterior output from the A11 analysis species delimitation model in BPP using

923   unphased loci from the 38 Galapagos giant tortoise individuals. Support for collapsed taxa as

924   species is included in the final column. The "Medium θ" prior is the most realistic prior.

θ prior	No. loci	Seed	Posterior for number of species	Posteriors for collapsed taxonomic groups or notes
Small θ	50 loci	Seed 1	P[13]=0.986 P[12]=0.014	*porteri+donfaustoi*=0.013
		Seed 2	P[13]=0.997	
		Seed 3	P[13]=0.995	
Medium θ	50 loci	Seed 1	P[13]=0.992	
		Seed 2	P[13]=0.993	
		Seed 3	P[1]=1.000	single_species=1.000
Large θ	50 loci	Seed 1	P[13]=0.577 P[12]=0.357 P[11]=0.063	*guntheri+vandenburghi*=0.228 *darwini+becki (PBR)*=0.056 *microphyes+guntheri*=0.037 *vandenburghi+vicina*=0.036 *guntheri+vicina*=0.036 *porteri+donfaustoi*=0.026 *guntheri+vandenburghi+ vicina*=0.025
		Seed 2	P[13]=0.517 P[12]=0.334 P[11]=0.066 P[10]=0.071 P[9]=0.012	*guntheri+vandenburghi*=0.232 *darwini+becki (PBR)*=0.066 *guntheri+vandenburghi+ microphyes+vicina*=0.075 *microphyes+guntheri*=0.050 *porteri+donfaustoi*=0.031 *guntheri+vicina*=0.030 *guntheri+vandenburghi+ vicina*=0.029 *vandenburghi+vicina*=0.027
		Seed 3	P[13]=0.493 P[12]=0.391 P[11]=0.106 P[10]=0.010	*guntheri+vandenburghi*=0.248 *darwini+becki (PBR)*=0.092 *microphyes+guntheri*=0.064 *guntheri+vicina*=0.039 *guntheri+vandenburghi+ vicina*=0.039 *vandenburghi+vicina*=0.036 *porteri+donfaustoi*=0.024 *guntheri+vandenburghi+ microphyes* =0.022

928 **Table S6.** Posterior output from the A11 analysis species delimitation model in BPP using

929 unphased loci from the 38 Galapagos giant tortoise individuals and the Chaco tortoise individual.

930 Support for collapsed taxa as species is included in the final column. The "Medium θ" prior is the

931 most realistic prior.

932
933

θ prior	No. loci	Seed	Posterior for number of species	Posteriors for collapsed taxonomic groups or notes
Small θ	50 loci	Seed 1	P[13]=0.036 P[12]=0.964	*guntheri+vandenburghi+vicina*=0.964 *guntheri+vandenburghi*=0.036
		Seed 2	P[13]=0.173 P[12]=0.826	*guntheri+vandenburghi+vicina*=0.449 *guntheri+vandenburghi+microphyes*=0.378 *guntheri+vandenburghi*=0.173
		Seed 3	P[12]=0.987	*guntheri+vandenburghi+microphyes*=0.628 *guntheri+vandenburghi+vicina*=0.359
Medium θ	50 loci	Seed 1	P[13]=0.401 P[12]=0.589	*guntheri+vandenburghi+microphyes*=0.118 *guntheri+vandenburghi+vicina*=0.472 *guntheri+vandenburghi*=0.400
		Seed 2	P[14]=0.010 P[13]=0.157 P[12]=0.832	*guntheri+vandenburghi+microphyes*=0.735 *guntheri+vandenburghi*=0.146 *guntheri+vandenburghi+vicina*=0.097 *guntheri+microphyes*=0.011
		Seed 3	P[14]=0.010 P[13]=0.110 P[12]=0.879	*guntheri+vandenburghi+vicina*=0.573 *guntheri+vandenburghi+microphyes*=0.301 *guntheri+vandenburghi*=0.080 *guntheri+microphyes*=0.024 *vandenburghi+microphyes*=0.011
Large θ	50 loci	Seed 1	P[11]=0.241 P[10]=0.634	*guntheri+vandenburghi+microphyes+vicina*=0.980

			P[9]=0.123	*porteri+donfaustoi*=0.623 *becki (PBL)+becki (PBR)*=0.232 *darwini+becki (PBR)*=0.042
		Seed 2	P[12]=0.022 P[11]=0.385 P[10]=0.430 P[9]=0.163	*guntheri+vandenburghi+* *microphyes+vicina*=0.841 *porteri+donfaustoi*=0.536 *becki (PBL)+becki (PBR)*=0.344 *guntheri+vandenburghi+* *microphyes* =0.159 *darwini+becki (PBR)*=0.011
		Seed 3	P[11]=0.235 P[10]=0.497 P[9]=0.263	*guntheri+vandenburghi+* *microphyes+vicina*=0.997 *porteri+donfaustoi*=0.624 *becki (PBL)+becki (PBR)*=0.349 *darwini+becki (PBL)+becki* *(PBR)*=0.019 *darwini+becki (PBR)*=0.019

949 **Table S7.** Posterior output from the A10 analysis species delimitation model in BPP using phased

950 loci from the 38 Galapagos giant tortoise individuals. Nodes in the guide tree with a posterior

951 probability <0.95 were counted as a single species. The "Medium θ" prior is the most realistic

952 prior. A maximum of 13 species (i.e., each Galapagos giant tortoise taxon) was possible.

θ prior	No. loci	Seed	Number of species with >0.95 posterior support
Small θ	50 loci	Seed 1	13
		Seed 2	13
		Seed 3	13
	200 loci	Seed 1	13
		Seed 2	13
		Seed 3	13
	500 loci	Seed 1	13
		Seed 2	13
		Seed 3	13
	1000 loci	Seed 1	13
		Seed 2	13
		Seed 3	13
Medium θ	50 loci	Seed 1	13
		Seed 2	13
		Seed 3	13
	200 loci	Seed 1	13
		Seed 2	13
		Seed 3	13
	500 loci	Seed 1	13
		Seed 2	13
		Seed 3	13
	1000 loci	Seed 1	13
		Seed 2	13
		Seed 3	13
Large θ	50 loci	Seed 1	8
		Seed 2	9
		Seed 3	9
	200 loci	Seed 1	11
		Seed 2	10
		Seed 3	11
	500 loci	Seed 1	12
		Seed 2	12
		Seed 3	12
	1000 loci	Seed 1	12

		Seed 2	13
		Seed 3	13

Table S8. Posterior output from the A10 analysis species delimitation model in BPP using phased

loci from the 38 Galapagos giant tortoise individuals and the Chaco tortoise individual. Nodes in

the guide tree with a posterior probability <0.95 were counted as a single species. The "Medium

θ" prior is the most realistic prior. A maximum of 14 species (i.e., each Galapagos giant tortoise

taxon and the Chaco tortoise) was possible.

θ prior	No. loci	Seed	Number of species with >0.95 posterior support
Small θ	50 loci	Seed 1	10
		Seed 2	11
		Seed 3	12
	200 loci	Seed 1	14
		Seed 2	13
		Seed 3	11
	500 loci	Seed 1	10
		Seed 2	12
		Seed 3	10
	830 loci	Seed 1	14
		Seed 2	10
		Seed 3	14
Medium θ	50 loci	Seed 1	12
		Seed 2	12
		Seed 3	12
	200 loci	Seed 1	14
		Seed 2	14
		Seed 3	12
	500 loci	Seed 1	14
		Seed 2	14
		Seed 3	14
	830 loci	Seed 1	14
		Seed 2	14
		Seed 3	14
Large θ	50 loci	Seed 1	9
		Seed 2	10
		Seed 3	10
	200 loci	Seed 1	12

		Seed 2	12
		Seed 3	11
	500 loci	Seed 1	11
		Seed 2	13
		Seed 3	12
	830 loci	Seed 1	13
		Seed 2	11
		Seed 3	12

978 **Table S9**. Divergence time estimates in generations from BPP, with prior $\theta \sim$ IG(3, 0.001) and

979 prior $\tau \sim$ IG(3, 0.001). Median and 95% credible interval calculated from the posterior distribution

980 of τ. Divergence time calculated using a per-generation mutation rate of 6×10^{-9}.

Taxon	Median divergence time in generations (95% credible interval)
darwini	2667 (2000–3333)
becki (PBR)	2667 (2000–3333)
guntheri	1333 (1167–1667)
microphyes	1333 (1167–1667)
vandenburghi	1500 (1333–1667)
vicina	2333 (2000–2667)
becki (PBL)	3833 (3167–4667)
donfaustoi	3167 (2500–4000)
porteri	3167 (2500–4000)
hoodensis	4500 (3500–5500)
phantasticus	4500 (3500–5500)
duncanensis	5667 (4833–6833)
chathamensis	6333 (5500–7833)
C. chilensis	368,500 (352,667–392,167)

987 **Table S10**. Effective population size estimates from BPP, with prior $\theta \sim$ IG(3, 0.001) and prior τ

988 $\sim$ IG(3, 0.001) and an assumed mutation rate of 6.0 x 10^{-9}. Median and 95% credible interval

989 calculated from the posterior distribution of θ. Effective population size calculated using a per-

990 generation mutation rate of 6×10^{-9}.

Taxon	Median effective population size (95% credible interval)
darwini	8208 (6000–10,667)
becki (PBR)	5458 (4083–6958)
guntheri	6416 (4833–8833)
microphyes	3917 (3125–5083)
vandenburghi	12,125 (8583–19,042)
vicina	27,333 (25,542–29,208)
becki (PBL)	19,750 (15,208–25,875)
donfaustoi	5083 (3958–6625)
porteri	45,125 (26,083–94,333)
hoodensis	2917 (2333–3542)
phantasticus	94,042 (45,208–223,375)
duncanensis	5750 (4875–7125)
chathamensis	9333 (7833–11,375)
C. chilensis	77,917 (71,292–85,292)

996   **Table S11**. Estimates of *gdi* for each taxon, calculated from the posterior distributions of θ and τ

997   in BPP, with prior $\theta \sim$ IG(3, 0.001) and prior $\tau \sim$ IG(3, 0.001).

Taxon	Median *gdi* (95% credible interval)
darwini	0.149 (0.140–0.159)
becki (PBR)	0.218 (0.187–0.247)
guntheri	0.104 (0.077–0.131)
microphyes	0.162 (0.132–0.194)
vandenburghi	0.063 (0.040–0.087)
vicina	0.107 (0.082–0.134)
becki (PBL)	0.094 (0.075–0.115)
donfaustoi	0.266 (0.231–0.278)
porteri	0.034 (0.016–0.058)
hoodensis	0.534 (0.497–0.572)
phantasticus	0.023 (0.010–0.049)
duncanensis	0.423 (0.393–0.455)
chathamensis	0.262 (0.236–0.290)

1000

1001

1002

1003

1004

1005

1006 **Table S12**. Estimates of *gdi* for each taxon, calculated from the posterior distributions of θ and τ

1007 in BPP, with prior $\theta \sim$ IG(3, 0.005) and prior $\tau \sim$ IG(3, 0.001).

Taxon	Median *gdi* (95% credible interval)
darwini	0.137 (0.111–0.166)
becki (PBR)	0.197 (0.168–0.227)
guntheri	0.083 (0.060–0.108)
microphyes	0.134 (0.106–0.160)
vandenburghi	0.048 (0.029–0.069)
vicina	0.092 (0.070–0.116)
becki (PBL)	0.091 (0.072–0.112)
donfaustoi	0.235 (0.223–0.271)
porteri	0.028 (0.008–0.049)
hoodensis	0.515 (0.477–0.551)
phantasticus	0.028 (0.010–0.052)
duncanensis	0.429 (0.398–0.460)
chathamensis	0.268 (0.241–0.296)

1008

1009

1010

1011

1012

1013

1014

1015

1016 **Table S13**. Estimates of *gdi* for each taxon, calculated from the posterior distributions of θ and τ

1017 in BPP, with prior $\theta \sim$ IG(3, 0.0001) and prior $\tau \sim$ IG(3, 0.001).

Taxon	Median *gdi* (95% credible interval)
darwini	0.152 (0.124–0.180)
becki (PBR)	0.223 (0.193–0.255)
guntheri	0.109 (0.077–0.143)
microphyes	0.171 (0.133–0.210)
vandenburghi	0.064 (0.041–0.090)
vicina	0.107 (0.081–0.135)
becki (PBL)	0.094 (0.074–0.115)
donfaustoi	0.271 (0.233–0.309)
porteri	0.032 (0.012–0.058)
hoodensis	0.537 (0.502–0.573)
phantasticus	0.025 (0.013–0.046)
duncanensis	0.413 (0.381–0.446)
chathamensis	0.256 (0.228–0.283)

1018

1019

1020

1021

1022

1023

1024

1025 **Table S14**. Estimates of *gdi* for each taxon after the first round of sister taxa were collapsed. The

1026 *gdi* was calculated from the posterior distributions of θ and τ of the collapsed taxa in BPP, with

1027 prior $\theta \sim$ IG(3, 0.001) and prior $\tau \sim$ IG(3, 0.001).

Taxon	Median *gdi* (95% credible interval)
darwini+becki (PBR)	0.161 (0.144–0.180)
guntheri+microphyes	0.044 (0.029–0.062)
vandenburghi	0.158 (0.127–0.190)
vicina	0.105 (0.080–0.131)
becki (PBL)	0.093 (0.074–0.113)
donfaustoi+porteri	0.291 (0.268–0.314)
hoodensis+phantasticus	0.190 (0.170–0.211)
duncanensis	0.427 (0.396–0.460)
chathamensis	0.263 (0.235–0.291)

1028

1029

1030

1031

1032

1033

1034

1035

1036

1037

1038

Table S15. Estimates of *gdi* for each taxon after the second round of sister taxa were collapsed.

The *gdi* was calculated from the posterior distributions of θ and τ of the collapsed taxa in BPP,

with prior $\theta \sim$ IG(3, 0.001) and prior $\tau \sim$ IG(3, 0.001).

Taxon	Median *gdi* (95% credible interval)
darwini+becki (PBR)+becki (PBL)	0.096 (0.086–0.106)
guntheri+microphyes+vandenburghi	0.094 (0.078–0.111)
vicina	0.102 (0.077–0.129)
donfaustoi+porteri	0.295 (0.273–0.318)
hoodensis+ phantasticus+chathamensis	0.129 (0.115–0.145)
duncanensis	0.423 (0.392–0.456)

1042

1043

Table S16. Estimates of *gdi* for each taxon after the third round of sister taxa were collapsed. The

gdi was calculated from the posterior distributions of θ and τ of the collapsed taxa in BPP, with

prior $\theta \sim$ IG(3, 0.001) and prior $\tau \sim$ IG(3, 0.001).

Taxon	Median *gdi* (95% credible interval)
darwini+becki (PBR)+becki (PBL)	0.099 (0.089–0.110)

guntheri+microphyes+vandenburghi+vicina	0.226 (0.211–0.243)
donfaustoi+porteri	0.298 (0.275–0.322)
hoodensis+ phantasticus+duncanensis+ chathamensis	0.144 (0.134–0.156)

1047

1048 **Table S17**. Estimates of *gdi* for each taxon after the fourth round of sister taxa were collapsed.

1049 The *gdi* was calculated from the posterior distributions of θ and τ of the collapsed taxa in BPP,

1050 with prior $\theta \sim$ IG(3, 0.001) and prior $\tau \sim$ IG(3, 0.001).

Taxon	Median *gdi* (95% credible interval)
darwini+becki (PBR)+becki (PBL)+ guntheri+microphyes+vandenburghi+vicina	0.113 (0.103–0.124)
donfaustoi+porteri	0.287 (0.264–0.310)
hoodensis+phantasticus+duncanensis+ chathamensis	0.144 (0.132–0.155)

1051

1052

1053 **Table S18.** Estimates of *gdi* for each taxon after the fifth round of sister taxa were collapsed. The

1054 *gdi* was calculated from the posterior distributions of θ and τ of the collapsed taxa in BPP, with

1055 prior $\theta \sim$ IG(3, 0.001) and prior $\tau \sim$ IG(3, 0.001).

Taxon	Median *gdi* (95% credible interval)
darwini+becki (PBR)+becki (PBL)+ guntheri+microphyes+vandenburghi+vicina+ donfaustoi+porteri	0.110 (0.102–0.118)
hoodensis+ phantasticus+duncanensis+ chathamensis	0.142 (0.132–0.153)

1056

1057

1058

1059

1060

1061 **Table S19**. Estimates of *gdi* for each taxon after the sixth round of sister taxa were collapsed. The

1062 *gdi* was calculated from the posterior distributions of θ and τ of the collapsed taxa in BPP, with

1063 prior $\theta \sim$ IG(3, 0.001) and prior $\tau \sim$ IG(3, 0.001).

Taxon	Median *gdi* (95% credible interval)
darwini+becki (PBR)+becki (PBL)+ guntheri+microphyes+vandenburghi+vicina+ donfaustoi+porteri+ hoodensis+ phantasticus+duncanensis+chathamensis	0.988 (0.984–0.991)
C. chilensis	0.92 (0.901–0.939)

1064

1065

Table S20: Pairwise comparisons analyzed in PHRAPL across the two main clades of the phylogenetic tree, along with the wAIC for each of the three PHRAPL models. Models with a high Akaike weight (wAIC) > 0.9 for a model set are highlighted in gray. The cumulative sum of wAIC for all two species models is given for each comparison, with wAIC > 0.9 also highlighted in gray. Mean *gdi* < 0.2 are shown in bold.

Taxon 1	Taxon 2	wAIC Model 1 2 species, no gene flow	wAIC Model 2 2 species, constant gene flow	wAIC Model 3 1 species	Sum wAIC of 2 Species Models	Mean *gdi*
Within Saddleback Clade						
duncanensis	*hoodensis*	0.984	0.016	0.000	1.000	0.387
chathamensis	*hoodensis*	0.465	0.535	0.000	1.000	0.384
chathamensis	*duncanensis*	0.722	0.278	0.000	1.000	0.217
hoodensis	*phantasticus*	0.539	0.461	0.000	1.000	**0.179**
duncanensis	*phantasticus*	0.022	0.978	0.000	1.000	0.319

chathamensis	*phantasticus*	0.000	1.000	0.000	1.000	0.319

Within Isabela Island (Domed Clade)

becki(PBL)	*becki*(PBR)	0.916	0.084	0.000	1.000	**0.182**
vicina	*guntheri*	0.227	0.773	0.000	1.000	0.447
vicina	*vandenburghi*	0.000	1.000	0.000	1.000	0.304
vicina	*microphyes*	0.005	0.995	0.000	1.000	0.345
vicina	*becki*(PBL)	0.127	0.873	0.000	1.000	**0.179**
vicina	*becki*(PBR)	0.000	1.000	0.000	1.000	0.307
guntheri	*vandenburghi*	0.001	0.999	0.000	1.000	0.455
guntheri	*microphyes*	0.000	1.000	0.000	1.000	0.460
guntheri	*becki*(PBL)	0.702	0.298	0.000	1.000	0.404

guntheri	*becki*(PBR)	0.000	1.000	0.000	1.000	0.536
vandenburghi	*microphyes*	0.790	0.210	0.000	1.000	**0.180**
vandenburghi	*becki*(PBL)	0.000	1.000	0.000	1.000	0.280
vandenburghi	*becki*(PBR)	0.000	1.000	0.000	1.000	0.316
microphyes	*becki*(PBL)	0.000	1.000	0.000	1.000	0.314
microphyes	*becki*(PBR)	0.006	0.994	0.000	1.000	0.344
Within Santa Cruz Island (Domed Clade)						
donfaustoi	*porteri*	0.489	0.511	0.000	1.000	**0.181**
Santa Cruz Island with Isabela Island (Domed Clade)						
donfaustoi	*becki*(PBL)	0.745	0.255	0.000	1.000	0.386

donfaustoi	*becki*(PBR)	0.004	0.996	0.000	1.000	0.458
donfaustoi	*vandenburghi*	0.665	0.335	0.000	1.000	0.386
donfaustoi	*microphyes*	0.000	1.000	0.000	1.000	0.484
donfaustoi	*guntheri*	0.000	1.000	0.000	1.000	0.276
donfaustoi	*vicina*	0.488	0.512	0.000	1.000	0.400
porteri	*becki*(PBL)	0.000	1.000	0.000	1.000	0.324
porteri	*becki*(PBR)	0.653	0.347	0.000	1.000	0.384
porteri	*vandenburghi*	0.665	0.335	0.000	1.000	0.386
porteri	*microphyes*	0.831	0.169	0.000	1.000	0.386
porteri	*guntheri*	0.000	1.000	0.000	1.000	0.281

porteri	*vicina*	0.000	1.000	0.000	1.000	0.344
Isabela and Santa Cruz Islands compared with Santiago Island (Domed Clade)						
darwini	*becki*(PBL)	0.004	0.996	0.000	1.000	0.278
darwini	*becki*(PBR)	0.000	1.000	0.000	1.000	0.295
darwini	*donfaustoi*	0.597	0.403	0.000	1.000	**0.181**
darwini	*porteri*	0.203	0.797	0.000	1.00	**0.181**
darwini	*guntheri*	0.188	0.812	0.000	1.000	**0.178**
darwini	*microphyes*	0.000	1.000	0.000	1.000	0.375
darwini	*vandenburghi*	0.000	1.000	0.000	1.000	0.372
darwini	*vicina*	0.000	1.000	0.000	1.000	0.318

1070

<u>**Supplemental References**</u>

Austin, C. C., Rittmeyer, E. N., Richards, S. J., & Zug, G. R. (2010). Phylogeny, historical

biogeography and body size evolution in Pacific Island Crocodile skinks Tribolonotus

(Squamata; Scincidae). *Molecular Phylogenetics and Evolution*, *57*(1), 227-236.

Baele, G., Lemey, P., Bedford, T., Rambaut, A., Suchard, M. A., & Alekseyenko, A. V. (2012).

Improving the accuracy of demographic and molecular clock model comparison while

accommodating phylogenetic uncertainty. *Molecular Biology and Evolution*, *29*(9), 2157-

2167.

Bergeron, L. A., Besenbacher, S., Zheng, J., Li, P., Bertelsen, M. F., Quintard, B., ... & Zhang,

G. (2023). Evolution of the germline mutation rate across vertebrates. *Nature*, *615*(7951),

285-291.

Bouckaert, R., Heled, J., Kühnert, D., Vaughan, T., Wu, C. H., Xie, D., ... & Drummond, A. J.

(2014). BEAST 2: a software platform for Bayesian evolutionary analysis. *PLoS*

Computational Biology, *10*(4), e1003537.

Bryant, D., Bouckaert, R., Felsenstein, J., Rosenberg, N. A., & RoyChoudhury, A. (2012).

Inferring species trees directly from biallelic genetic markers: bypassing gene trees in a

full coalescent analysis. *Molecular Biology and Evolution*, *29*(8), 1917-1932.

Ciofi, C., Milinkovitch, M. C., Gibbs, J. P., Caccone, A., & Powell, J. R. (2002). Microsatellite

analysis of genetic divergence among populations of giant Galápagos tortoises. *Molecular*

Ecology, *11*(11), 2265-2283.

Ciofi, C., Wilson, G. A., Beheregaray, L. B., Marquez, C., Gibbs, J. P., Tapia, W., ... & Powell, J.

R. (2006). Phylogeographic history and gene flow among giant Galápagos tortoises on

southern Isabela Island. *Genetics*, *172*(3), 1727-1744.

Drummond, A. J., & Rambaut, A. (2007). BEAST: Bayesian evolutionary analysis by sampling

trees. *BMC Evolutionary Biology*, *7*, 214.

Garrick, R. C., Benavides, E., Russello, M. A., Hyseni, C., Edwards, D. L., Gibbs, J. P., ... &

Caccone, A. (2014). Lineage fusion in Galápagos giant tortoises. *Molecular Ecology*,

23(21), 5276-5290.

Hedrick, P. W. (2009). *Genetics of populations*. Jones & Bartlett Publishers.

Iannucci, A., Benazzo, A., Natali, C., Arida, E. A., Zein, M. S. A., Jessop, T. S., ... & Ciofi, C.

(2021). Population structure, genomic diversity and demographic history of Komodo

dragons inferred from whole-genome sequencing. *Molecular Ecology*, *30*(23), 6309-

6324.

Jensen, E. L., Gaughran, S. J., Garrick, R. C., Russello, M. A., & Caccone, A. (2021).

Demographic history and patterns of molecular evolution from whole genome sequencing

in the radiation of Galapagos giant tortoises. *Molecular Ecology*, *30*(23), 6325-6339.

Jensen, E. L., Gaughran, S. J., Fusco, N. A., Poulakakis, N., Tapia, W., Sevilla, C., ... &

Caccone, A. (2022). The Galapagos giant tortoise *Chelonoidis phantasticus* is not extinct.

Communications Biology, *5*(1), 546.

Kass, R. E., & Raftery, A. E. (1995). Bayes factors. *Journal of the American Statistical*

Association, *90*(430), 773-795.

Leaché, A. D., Zhu, T., Rannala, B., & Yang, Z. (2019). The spectre of too many species. *Systematic Biology*, *68*(1), 168-181.

Li, H. (2013). Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. *arXiv preprint arXiv:1303.3997*.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., ... & 1000 Genome Project Data Processing Subgroup. (2009). The sequence alignment/map format and SAMtools. *Bioinformatics*, *25*(16), 2078-2079.

Miller, J. M., Quinzin, M. C., Edwards, D. L., Eaton, D. A., Jensen, E. L., Russello, M. A., ... & Caccone, A. (2018). Genome-wide assessment of diversity and divergence among extant Galapagos giant tortoise species. *Journal of Heredity*, *109*(6), 611-619.

Quesada, V., Freitas-Rodríguez, S., Miller, J., Pérez-Silva, J. G., Jiang, Z. F., Tapia, W., ... & López-Otín, C. (2019). Giant tortoise genomes provide insights into longevity and age-related disease. *Nature Ecology & Evolution*, *3*(1), 87-95.

Rambaut, A., Drummond, A. J., Xie, D., Baele, G., & Suchard, M. A. (2018). Posterior summarization in Bayesian phylogenetics using Tracer 1.7. *Systematic Biology*, *67*(5), 901-904.

Rannala, B. and Yang, Z. (2020). Species Delimitation. In Scornavacca, C., Delsuc, F., and Galtier, N. (eds.). pp. 5.5:1– 5.5:18 in *Phylogenetics in the Genomic Era*. No commercial publisher | Authors open access book. https://hal.inria.fr/PGE

Stephens, M., Smith, N. J., & Donnelly, P. (2001). A new statistical method for haplotype reconstruction from population data. *The American Journal of Human Genetics*, *68*(4), 978-989.

1134   Stephens, M., & Scheet, P. (2005). Accounting for decay of linkage disequilibrium in haplotype

1135   inference and missing-data imputation. *The American Journal of Human Genetics*, *76*(3),

1136   449-462.

1137

1138