Inhibition of microtubule detyrosination by parthenolide facilitates functional CNS axon regeneration

Marco Leibinger[1,2*], Charlotte Zeitler[1,2*], Miriam Paulat[2], Philipp Gobrecht[1,2], Alexander Hilla[2], Anastasia Andreadaki[1,2], Rainer Guthoff[3] and Dietmar Fischer[1,2]

*both authors have contributed equally

[1]Center for Pharmacology, Institute II, Medical Faculty and University of Cologne, Paul-Schallück-Straße 8, 50937, Cologne, Germany

[2]Department of Cell Physiology, Ruhr University of Bochum, Universitätsstraße 150, 44780 Bochum, Germany

[3]Eye Hospital, Heinrich Heine University Düsseldorf, Moorenstrasse 5, 40225 Düsseldorf, Germany

Lead Contact:

Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, Dietmar Fischer, dietmar.fischer@uni-koeln.de

Correspondence and proofs:

Dietmar Fischer, PharmD, Ph.D., Center of Pharmacology, Institute II, Medical Faculty and University of Cologne, Paul-Schallück-Straße 8, 50937, Cologne, Germany

Acknowledgments: We thank Dr. Günter Gisselmann, Anastasia Andreadaki, Christopher Brennsohn, Davina Stoesser, and Kessy Brzozowski for their technical support. The German Research Foundation supported this work.

Abstract

Injured axons in the central nervous system (CNS) usually fail to regenerate, causing permanent disabilities. However, the knockdown of PTEN or treatment of neurons with hyper-IL-6 (hIL-6) transforms neurons into a regenerative state, allowing them to regenerate axons in the injured optic nerve and spinal cord. Transneuronal delivery of hIL-6 to the injured brain stem neurons even enables functional recovery after severe spinal cord injury. Here we demonstrate that the beneficial hIL-6 and PTEN knockout effects on axon growth are limited by the induction of tubulin detyrosination in axonal growth cones. Hence, cotreatment with parthenolide, a compound blocking microtubule detyrosination, synergistically accelerates neurite growth of cultured murine and primary RGCs isolated from adult human eyes. Systemic application of the prodrug dimethylamino-parthenolide (DMAPT) enables axon regeneration in the injured optic nerve and spinal cord. Moreover, combinatorial treatment further improves hIL-6 induced axon regeneration and locomotor recovery after severe SCI. Thus, DMAPT facilitates functional CNS regeneration and reduces the limiting effects of pro-regenerative treatments, making it a promising drug candidate for treating CNS injuries.

Significance statement

Injured axons in the CNS usually fail to regenerate, causing permanent disabilities. No clinically approved drugs are currently available to improve or accelerate axon regeneration. We demonstrate that pro-regenerative treatments' induced tubulin detyrosination in axonal growth cones limits axon growth. Hence, cotreatment with parthenolide, a compound blocking detyrosination, synergistically accelerates neurite growth of primary murine and, demonstrated for the first time, <u>human</u> retinal ganglion cells. Furthermore, systemic application of the parthenolide's prodrug DMAPT accelerates hIL-6-mediated optic nerve regeneration and locomotor recovery after severe spinal cord injury. Thus, DMAPT facilitates functional CNS regeneration and reduces the limiting effects of pro-regenerative treatments, making it a potential (adjuvant) drug candidate for treating CNS injuries.

Introduction

Traumatic CNS injuries often lead to irreversible impairments, such as blindness or loss of motor and sensory function, due to the inability of severed axons to regrow. This regenerative failure is mainly caused by an insufficient intrinsic growth capacity of mature CNS neurons and the inhibitory environment for axonal growth cones at the injury site [15, 45, 54].

Although there is currently no clinically applicable treatment available to human patients, various strategies exist to promote axon regeneration in experimental models of CNS injury [15, 54, 64]. For example, targeting the inhibitory environment at the lesion site by neutralizing myelin-derived factors or chondroitin sulfate proteoglycans (CSPGs) led to axonal regeneration and, in some cases, induced functional recovery after incomplete spinal cord lesions[5, 7, 34, 47]. On the other hand, it became evident that not only disinhibition but also strategies to stimulate cell-intrinsic signaling pathways are essential to promote robust axon growth. Among these, lens injury-induced expression and release of the IL-6-type cytokines ciliary neurotrophic factor (CNTF) and leukemia inhibitory factor (LIF) from retinal astrocytes transforms retinal ganglion cells (RGCs) into a regenerative state and enables moderate axon growth in the injured optic nerve [16, 38, 49, 63]. The designer cytokine hyper-IL-6 (hIL-6), a fusion protein of IL-6 and the soluble IL-6 receptor α (IL-6Rα), can enormously increase these effects. In contrast to natural cytokines, hIL-6 directly binds to the gp130 receptor and does not rely on previous interaction with cytokine-specific alpha-receptors [18, 28, 35, 60]. This enables a more potent activation of downstream signaling pathways, such as the Janus kinase/signal transducer and activator of transcription 3 (JAK/STAT3) and phosphoinositide 3-kinase/protein kinase B (PI3K/AKT) pathways [35]. As a result, AAV-mediated hIL-6 expression in RGCs elicits a more vigorous optic nerve regeneration over long distances than most other single treatments [14, 35]. Similarly, hIL-6 expression in corticospinal neurons simultaneously promotes axon regeneration of multiple descending tracts and partially restores motor function after a complete spinal cord crush [40, 60].

Another way to stimulate CNS axon regeneration is to directly activate the PI3K/AKT pathway by knockout phosphatase and tensin homolog (PTEN[-/-]) [43, 53, 66]. This approach stimulated axon regeneration in the corticospinal tract even when applied one year after injury and induced functional recovery of voluntary skilled movements after incomplete spinal cord injury or pyramidotomy [12, 13, 42, 43]. PTEN[-/-] or hIL-6-mediated AKT activation causes phosphorylation of several substrates, including inactivating phosphorylation of glycogen synthase kinase 3 (GSK3) [58, 59]. The specific GSK3β knockout (GSK3β[-/-]) in RGCs facilitates optic nerve regeneration by reducing collapsin response mediator protein 2 (CRMP2) phosphorylation in axons, while constitutively active GSK3 (GSK3[S/A]) compromises PTEN[-/-] and cytokine-induced regeneration, underlining the relevance of the AKT/GSK3/CRMP2 pathway for CNS regeneration [36, 37].

Besides CRMP2, GSK3 phosphorylates microtubule-associated protein 1B (MAP1B). Phosphorylated MAP1B keeps microtubules in a dynamic state by blocking the removal of the C-terminal tyrosine from α-tubulin subunits (detyrosination) in axonal growth cones [20, 22-24, 46, 52]. In addition, inhibition of tubulin detyrosination accelerates peripheral nerve regeneration [20]. This was achieved by local or systemic application of the sesquiterpene lactone parthenolide, an inhibitor of VASH1/2, which mediates detyrosination [1, 19, 21, 51].

The current study addressed whether parthenolide affects axon regeneration in the CNS. It shows that it markedly accelerates neurite outgrowth of primary CNS neurons by reducing tubulin detyrosination in axonal tips. Moreover, intraperitoneal application of its prodrug dimethylamino-parthenolide (DMAPT) facilitates axon regeneration after optic nerve and spinal cord injuries *in vivo*. Furthermore, hIL-6 and PTEN[-/-] were found to increase tubulin detyrosination. At the same time, parthenolide antagonized this effect, thereby further enhancing hIL-6-induced optic nerve and spinal cord regeneration and functional recovery of hindlimb locomotion after severe spinal cord injury. The effects of parthenolide on detyrosination and neurite outgrowth were reproducible in human adult RGCs, suggesting it is a promising drug candidate

in patients suffering CNS injuries. In particular, parthenolide is an ideal cotreatment to accelerate axon regeneration and functional recovery in combination with other treatment strategies.

Materials and Methods

Mouse strains

Male and female C57BL/6 and 129/Ola mice were used for all experiments. All mice were housed under the same conditions for at least 10 days before the experiments and maintained on a 12 h light/dark cycle with *ad libitum* access to food and water. All experimental procedures were approved by the local animal care committee (LANUV Recklinghausen) and were conducted in compliance with federal and state guidelines for animal experiments in Germany.

Human eyes

The use of human eyes was approved by the ethics committee of Heinrich Heine University Düsseldorf and performed by the ethical standards as laid down in the 1964 Declaration of Helsinki and its later amendments or comparable ethical standards.

Surgical procedures

Optic nerve crush and intravitreal injections

Adult C57BL/6 mice were anesthetized by intraperitoneal injections of ketamine (120 mg/kg) and xylazine (16 mg/kg). For western blot and cell culture experiments, RGCs were transduced 3 weeks before tissue isolation by intravitreal injection of 2 µl AAV2-hIL-6. For *in vivo* regeneration experiments, the left optic nerve was intraorbitally crushed 1 mm behind the eyeball for 10 s using a jeweler's forceps (Hermle), as described previously (Leibinger et al. 2009). Mice received intravitreal injections of 2 µl AAV2-hIL-6 directly after optic nerve crush. Two days before tissue isolation, regenerating axons were anterogradely labeled by intravitreal injection of 2 µl Alexa Fluor 555-conjugated cholera toxin β subunit (CTB, 0.5% in PBS; Invitrogen).

Intracortical viral injections

Adult 129/Ola mice were placed in a stereotactic frame and continuously supplied with 1.5 - 2 % isoflurane for anesthesia. A midline incision was made over the skull to open the skin and reveal bregma. A micro drill with a 0.5 mm bit was used to open a 2 x 2 mm window on each

side of the skull to expose the sensorimotor cortex. AAV2-hIL-6 was injected into the cortex layer V to transduce corticospinal neurons. Four injections of 500 nl each were applied into the left cortex through a glass capillary attached to a nanoliter injector (Nanoject III, Drummond) at the following coordinates: 1.5 mm lateral, 0.6 mm deep, and 0.5 mm anterior; 0.0 mm, 0.5 mm, and 1.0 mm posterior to bregma. At each injection site, five pulses of 100 nl were applied at a rate of 10 nl/s. A pause of 10 s was made after each pulse to allow the distribution of the virus solution. The needle was left in place for 1 min before moving to the next site.

Complete spinal cord crush

A complete spinal cord crush was performed as described previously [40, 61]. Adult 129/Ola mice were continuously supplied with 1.5 - 2 % isoflurane for anesthesia. A midline incision of ~1.5 cm was made over the thoracic vertebrae. The fat and muscle tissue were cleared from thoracic vertebrae 7 and 8 (T7, T8). While holding onto T7 with forceps, a laminectomy was performed at T8 to expose the spinal cord. The complete spinal cord was then crushed for 2 s with forceps that had been filed to a width of 0.15 mm for the last 5 mm of the tips to generate a homogeneously thin lesion site [66]. After surgery, muscle layers were sutured with 5.0 resorbable sutures, and the skin was secured with wound clips.

DHT injection

To deplete serotonergic inputs to the spinal cord, mice received bilateral intracerebroventricular injections of 30 µg of the serotonin neurotoxin 5, 7-dihydroxytryptamine (DHT, Biomol). DHT was dissolved in 0.5 µl of 0.2 % ascorbic acid in saline, as described previously [40]. To this end, mice were anesthetized and fixed under a stereotactic frame described above for intra-cortical injections. In addition, the tip of a glass micropipette attached to a nanoject III (Drummond) was positioned at the following coordinates: 0.6 mm posterior, 1.6 mm lateral to bregma, and 2 mm deep from the cortical surface. DHT was injected at the same rate as AAV2 injections, and the pipette was left for 1 minute before the withdrawal. Thirty minutes before the

DHT injection, the monoamine uptake inhibitor desipramine (Sigma) was intraperitoneally administered at 25 mg/kg to prevent DHT uptake into noradrenergic neurons.

Preparation of AAV2

AAV plasmids carrying cDNA for hIL-6 downstream of a CMV promoter were co-transfected with pAAV-RC (Stratagene) encoding the AAV genes rep and cap, and the helper plasmid (Stratagene) encoding E24, E4, and VA into AAV-293 cells (Stratagene) for recombinant AAV2 generation. In addition, purifying virus particles was performed as described previously [35, 40, 65].

Western blot assays

For protein lysate preparation, optic nerves from adult C57BL/6 mice, either untreated or 3 weeks after intraperitoneal AAV2-hIL-6 injection, were dissected and homogenized in lysis buffer (20 mM Tris/HCl pH 7.5, 10 mM KCl, 250 mM sucrose, 10 mM NaF, 1 mM DTT, 0.1 mM Na3VO4, 1% Triton X-100, 0.1% SDS) with protease inhibitors (Calbiochem, Darmstadt, Germany) and phosphatase inhibitors (Roche, Basel, Switzerland) using 5 sonication pulses at 40% power (Bandelin Sonoplus, Berlin, Germany). Lysates were cleared by centrifugation in an Eppendorf tabletop centrifuge at 4150 × g for 10 min at 4°C. Proteins were separated by sodium-dodecyl-sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) using Mini TGX gels (10%, Bio-Rad, Hercules, USA) according to standard protocols and transferred to nitrocellulose membranes (0.2 μm, Bio-Rad, Hercules, USA). Blots were blocked in 5% dried milk in phosphate-buffered saline with 0.05% Tween-20 (PBS-T) (Sigma, St Louis, USA) and incubated with antibodies against S21-phosphorylated GSK3α (1:2000; Cell Signaling Technology, RRID: AB_11166651), S9-phosphorylated GSK3β (1:1000; Cell Signaling Technology, RRID:AB_2115196), total GSK3α/β (1:1000; Cell Signaling Technology, RRID: AB_10547140), T514-phosphorylated CRMP2 (1:1000; Abcam, RRID:AB_942229), total CRMP2 (1:1000; Cell Signaling Technology, RRID:AB_2094339), detyrosinated tubulin (1:1000; Millipore, RRID:AB_177350) and βIII-tubulin (1:2000; BioLegend,

RRID:AB_2313773) at 4°C overnight. All antibodies were diluted in PBS-T containing 5% BSA (Sigma). A conformation-specific HRP-conjugated anti-rabbit IgG (1:4000; Cell Signaling Technology) was used as a secondary antibody. Antigen-antibody complexes were visualized using enhanced chemiluminescence substrate (Bio-Rad) on a FluorChem E detection system (ProteinSimple). Western blots were repeated at least three times to verify the results. Band intensities were quantified relative to respective loading controls by using ImageJ software.

Dissociated mouse retinal cell cultures

Adult C57BL/6 mice were used for all cell culture experiments. Some experiments included animals that had received intravitreal AAV2-hIL-6 injections 3 weeks before dissociation. Retinal cultures were prepared as described previously [25]. In brief, retinae were dissected from the eyecups and digested in Dulbecco's Modified Eagle medium (DMEM; Thermo Fisher) containing papain (10 U/ml; Worthington) and L-cysteine (0.2 µg/ml; Sigma) at 37°C for 30 minutes. After digestion, retinae were washed with DMEM, triturated, and centrifuged in 50 ml DMEM (500 g, 7 minutes). Cell pellets were resuspended in a medium containing B27 supplement (1:50; Invitrogen) and penicillin/streptomycin (1:50; Merck) and passed through a cell strainer (40 µm; Greiner Bio-One). Cells were treated with parthenolide (Sigma; 0.25-5.0 nM in dimethyl sulfoxide), recombinant rat CNTF (Peprotech; 200 ng/ml), DMAPT (Abcam; 0.25-5.0 nM in dimethyl sulfoxide), or a combination of parthenolide or CNTF. 300 µl cell suspension was added to each well on 4-well plates (Nunc), which were coated with poly-D-lysine (0.1 mg/ml, molecular weight 70,000-150,000 Da; Sigma) and laminin (20 µg/ml; Sigma). Retinal cells were cultured at 37 °C and 5 % CO_2.

After 4 days in culture, retinal cells were fixed with 4 % paraformaldehyde (PFA; Sigma). Cells were stained with primary antibodies against βIII-tubulin (1:2000; BioLegend, RRID:AB_2313773), detyrosinated tubulin (1:2000; Millipore), and phosphorylated (Tyr705) STAT3 (1:200; Cell Signaling Technology, RRID:AB_2491009). Secondary antibodies included donkey-anti-mouse and anti-rabbit antibodies conjugated to Alexa Fluor 488 or 594 (1:1000; ThermoFisher).

For studying neurite growth, all RGC neurites were photographed under a fluorescent micro-scope (200x; Axio Observer.D1, Zeiss), and neurite length was determined using ImageJ soft-ware. To evaluate microtubule detyrosination, at least 30 axonal tips per well were photo-graphed (400x; Axio Observer.D1, Zeiss) and analyzed using ImageJ software. Axon tips were defined as the last 15 µm of βIII-tubulin positive neurites. For the quantification of STAT3 phos-phorylation, at least 30 RGCs per well were evaluated (400x; Axio Observer.D1, Zeiss).

Quantification of pCRMP2 staining intensity in murine RGC cultures

To quantify pCRMP2 immunofluorescence, retinal cells were cultured in either 0.5 nM par-thenolide or the vehicle. After 4 d in culture, cells were fixed and immunostained against βIII-tubulin (1:1,000; BioLegend; RRID:AB_2313773) and phospho(T514)-CRMP2 (1:2,000; Abcam; RRID:AB_942229). The staining intensity of pCRMP2 in βIII-tubulin–positive RGCs were analyzed using ImageJ software and calculated according to the following formula: Cor-rected total cell fluorescence = integrated density − (area of selected cell × mean fluorescence of background reading). Data represent means ± SEM of three independent experiments. In each experiment, 80 RGCs per group were analyzed in two replicate wells.

Dissociated human retinal cell cultures

Primary human retinal cultures were generated in two independent experiments with retinae isolated from two adult human patients' enucleated eyes.

The retinae were dissected from the eyecups, and the vitreous body was removed. Afterward, the retinae were cut into 8 pieces and digested separately, each in 10 ml Dulbecco's Modified Eagle medium (DMEM; Thermo Fisher) containing papain (10 U/ml; Worthington) and L-cys-teine (0.2 µg/ml; Sigma) at 37 °C for 45 minutes. After digestion, retinae were washed with DMEM, triturated, and centrifuged in 50 ml DMEM (900 g, 5 minutes). Cell pellets were resus-pended in a medium containing B27 supplement (1:50; Invitrogen) and penicillin/streptomycin (1:50; Merck) and passed through a cell strainer (40 µm; Greiner Bio-One). Retinal cultures were incubated with parthenolide (Sigma; 0.25-5.0 nM in dimethyl sulfoxide) or the vehicle.

Some groups received recombinant rat CNTF (Peprotech; 200 ng/ml), either solely or in combination with parthenolide. Cells were cultured at 37 °C and 5 % CO_2 on 4-well plates (Nunc) (300 µl/well; 4 wells/group), which were coated with poly-D-lysine (0.1 mg/ml, molecular weight 70,000-150,000 Da; Sigma). After 4 days in culture, retinal cells were fixed with 4 % paraformaldehyde (PFA; Sigma) and stained with primary antibodies against βIII-tubulin (1:2000; BioLegend) and detyrosinated tubulin (1:2000; Millipore). Secondary antibodies included donkey-anti-mouse and anti-rabbit antibodies conjugated to Alexa Fluor 488 or 594 (1:1000; ThermoFisher). For studying neurite growth, and the percentage of detyrosinated axon tips, cultures were analyzed in the same way as described above for murine RGCs.

Intraperitoneal injections of DMAPT

Due to its high oral bioavailability of ~70% [27] and the ability to cross the blood-brain barrier [29], DMAPT can be delivered to the CNS via a systemic route of administration. Mice were given daily intraperitoneal injections of either DMAPT (0.2, 2, or 20 µg/kg; Abcam; dissolved in DMSO and further diluted in sterile PBS) or vehicle (DMSO diluted in sterile PBS). All spinal cord regeneration experiments were performed using the 2 µg/kg dosage. Treatment started directly after the optic nerve or spinal cord crush and was continued until the day of perfusion. Intraperitoneal injections were performed under blinded conditions.

Basso Mouse Scale (BMS)

We scored mice according to the Basso Mouse Scale to assess locomotor behavior after spinal cord injury [3]. To this end, each mouse was placed separately into a round open field of 1 m in diameter and observed by two testers for 4 minutes. Scoring was based on different parameters, including ankle movement, paw placement, stepping pattern, coordination, trunk instability, and tail position, with a minimum score of 0 (no movement) to a maximum of 9 (normal locomotion). Before testing, mice were acclimated to the open field environment. BMS tests were performed before the injury or 1, 3, and 7 days after the injury, and then weekly. As quality

criteria for lesion completeness, we used a BMS score of 0 on day 1 after injury and the absence of BDA or serotonin staining in transverse spinal cord sections (as described below). Mice that did not meet these criteria were excluded from the experiment. BMS tests were performed under blinded conditions.

Immunohistochemistry

Animals were anesthetized and transcardially perfused with cold PBS, followed by 4 % PFA in PBS. Brains, spinal cords, and eyes with attached optic nerves were isolated, post-fixed overnight in 4 % PFA at 4 °C, and subsequently transferred to 30 % sucrose for at least 4 h for eyes and optic nerves or 5 days for brains and spinal cords.

Tissues were embedded in the KP-cryo compound (Klinipath) and frozen at -20 °C (retina, optic nerve, spinal cord) or on dry ice (brain). Sections were cut using a CM3050 S cryostat (Leica Biosystems). Optic nerves were cut into longitudinal sections (14 µm). For spinal cords of mice subjected to T8 crush, a segment from 3 mm rostral to 8 mm caudal from the injury site was cut into sagittal sections (20 µm). Spinal cord tissue, just rostral and caudal from this segment, was used for transverse sections (20 µm) to evaluate lesion completeness by identifying potentially spared axons. Tissue from the medullary pyramids ~1 mm above the pyramidal decussation was cut into coronal sections (20 µm). Sections were thaw-mounted onto charged glass slides (Superfrost Plus, VWR) and stored at -20 °C until further use.

The retinal, spinal cord and brain sections were thawed, washed with PBS, and permeabilized for 10 min in methanol (Sigma). Sections were stained with primary antibodies against βIII-tubulin (1:1000; BioLegend, RRID: AB_10063408), phosphorylated STAT3 (1:200; Cell Signaling Technology, RRID:AB_2491009), serotonin (1:5000; Immunostar, RRID:AB_572262, RRID:AB_572263), GFP (1:500; Novus Biologicals, RRID:AB_10128178), NeuN (1:500; Abcam, RRID:AB_2532109), GFAP (1:500; Abcam, RRID:AB_305808), or CS-56 (1:70,000; Sigma, RRID:AB_476879). Secondary antibodies included donkey-anti-rabbit, anti-goat, and anti-mouse antibodies conjugated to Alexa Fluor 488, 594 (1:1000; ThermoFisher), or 405 (1:500; Jackson ImmunoResearch).

Retinal wholemounts were prepared without prior perfusion, fixed in 4 % PFA for 30 min, and permeabilized in 2 % Triton X-100 in PBS for 1 h. RGCs were stained using a primary antibody against βIII-tubulin (1:1000; BioLegend) and an anti-mouse secondary antibody conjugated to Alexa Fluor 488 (1:1000; ThermoFisher).

Sections and wholemounts were coverslipped with Mowiol (Sigma), and pictures were taken using a slide scanner (VS120; Olympus) or a confocal laser scanning microscope (SP8; Leica). Tissue analysis was performed under blinded conditions.

Quantification of regenerated axons in the optic nerve

Pictures of 6 longitudinal sections per animal were taken using a slide scanner (VS120; Olympus). The number of CTB-labelled axons extending 0.25, 0.5, and 0.75 mm (for DMAPT experiments) or 0.5, 1, 1.5, 2, and 2.5mm beyond the lesion site (for hIL-6 + DMAPT experiments) was quantified and normalized to the width of the optic nerve at the respective measuring point. Each experimental group included 6-7 mice. Eyes with lens injury were excluded from the experiment.

Quantification of RGCs in retinal wholemounts

Retinal wholemounts were divided into four quadrants. In each quadrant, 4 non-overlapping pictures were taken using a fluorescent microscope (400x; Axio Observer.D1, Zeiss), proceeding from the center to the periphery. The number of βIII-tubulin positive RGCs was determined and normalized to an area of 1 mm^2. Values were averaged per retina and across all animals. Each experimental group included 5-9 retinae. Eyes with lens injury were excluded from the experiment.

Quantification of RpST axon regeneration

Sagittal and coronal spinal cord sections were stained against serotonin and imaged with a slide scanner (20x; VS120; Olympus). A grid with lines spaced every 500 μm was aligned to each image to count the number of serotonin-positive RpST axons at 500 μm intervals from 0

to 8 mm caudal from the lesion site. In addition, every other spinal cord section was analyzed, and the number of serotonin-positive axons was extrapolated to the total number of sections per spinal cord. Animals with spared serotonin-positive axons in transverse spinal cord sections > 8 mm distal from the lesion site were excluded from the experiment. Each experimental group included 7-10 animals.

Quantification of spinal cord lesion sizes and CSPG staining

To evaluate the lesion size after SCC, coronal spinal cord sections containing the T8 lesion site were stained against GFAP, and pictures were taken using a fluorescence microscope (Olympus VS120). The lesion site, defined by the GFAP-positive astrocyte scar border, was traced and measured using ImageJ to obtain the total area. Furthermore, the width at the lesion center and both lateral edges were measured and averaged to determine the lesion width. The same sections were co-stained against CS-56 to visualize CSPGs at the injury site. Using ImageJ, the total area occupied by CSPGs was measured by tracing the CS-56 staining. Lesion sizes and CSPG staining was analyzed in three sections per animal.

Statistics

Significances of intergroup differences were evaluated using Student's t-test analysis of variance (ANOVA) followed by Tukey or Holm-Sidak post hoc tests using the Sigma STAT3.1 software (Systat Software).

Results

Parthenolide and CNTF synergistically promote neurite growth of adult RGCs.

In contrast to adult RGCs, sensory neurons usually show spontaneous neurite growth [35, 39, 41, 49]. Moreover, parthenolide-induced inhibition of tubulin detyrosination in axonal growth cones markedly accelerates the regeneration of sensory neurons in culture and *in vivo* [20, 22]. To investigate whether parthenolide has similar effects on CNS neurons, we cultured murine RGCs in the presence of increasing compound concentrations for 4 days. As in sensory neurons [20], parthenolide significantly enhanced neurite extension concentration-dependently with maximum effects at 0.5 and 1 nM (Fig. 1 A, B). Expectedly RGC survival remained unaffected (Fig. 1 C) since RGCs do not undergo axotomy-induced cell death as early as 4 days after culturing. Moreover, cotreatment of parthenolide with CNTF showed a significant additive effect (Fig. 1 D, E), resulting in a 3.5-fold higher average neurite length than vehicle-treated controls (Fig. 1 D, E). Although parthenolide reportedly reduces JAK/STAT3 signaling at micromolar concentrations [55, 56], the nanomolar concentrations used in our experiments did not alter CNTF-induced STAT3 phosphorylation in RGCs (Fig. 1 F, G). Conversely, CNTF did not measurably affect parthenolide's inhibitory effect on microtubule detyrosination in axon tips (Fig. 1 H, I). We also tested parthenolide on human RGCs. To this end, we prepared and cultured primary RGCs from adult human eyes. Parthenolide and CNTF significantly enhanced the average neurite length at similar concentrations as in mouse RGCs, while cotreatment enhanced neurite growth sixfold compared to controls after 4 days (Fig. 1 J, K). Parthenolide also reduced microtubule detyrosination in axonal growth cones (Fig.1 M, N) without affecting the number of RGCs per well (Fig. 1 L).

Parthenolide counteracts hIL-6 and PTEN[-/-]-induced microtubule detyrosination

Due to its direct binding of the signal-transducing gp130 receptor, hIL-6 promotes neurite growth of RGCs more efficaciously than CNTF and, contrary to CNTF, also measurably induces PI3K/AKT signaling in these neurons [35]. However, AKT inactivates GSK3 (Fig. 2 A),

which promotes CRMP2 activity (Leibinger et al., 2017, 2019). While enhanced CRMP2 activity is favorable for axon growth [36, 44], GSK3 inhibition might also elevate tubulin detyrosination via reduced MAP1B activation (Fig. 2 A). However, as shown in the previous experiments (Fig. 1), reducing microtubule detyrosination promotes neurite growth, suggesting that this effect of AKT/GSK3 might limit optimal axonal growth. To test these possibilities, we intravitreally injected AAV2-hIL-6 into mice 3 weeks before tissue isolation. Subsequent western blot analysis of respective optic nerve lysates indeed revealed increased inhibitory phosphorylation of both GSK3 isoforms (pGSK3α, pGSK3β) by hIL-6 compared to untreated controls, without affecting total GSK3 expression (Fig. 2 B-E). Moreover, while hIL-6 reduced inhibitory phosphorylation of CRMP2 at threonine 514 (T514) without changing total CRMP2 levels (Fig. 2 B, F, G), it expectedly elevated tubulin detyrosination (Fig. 2 B, H). Since parthenolide specifically acts on tubulin detyrosination without influencing CRMP2 activity (supplementary Fig. 1 A, B), it should be suitable to counteract an hIL-6-mediated increase in axonal tubulin detyrosination. To test this hypothesis, we cultured RGCs from AAV-hIL-6-treated mice or respective controls for 4 days without or presence of different parthenolide concentrations. Confirmingly, microtubule detyrosination in axonal tips was significantly higher in hIL-6-treated cultures than in controls, and parthenolide cotreatment reversed this effect (Fig. 2 I, J). Furthermore, the increased detyrosination in hIL-6-treated neurons also affected the effective parthenolide concentration on neurite extension in these cultures. Compared to the control cultures, where parthenolide showed the most substantial increase between 0.5 and 1 nM, followed by a decrease at 5 nM (Fig. 1 A, B), neurite growth of hIL-6-stimulated RGCs was increased even at higher parthenolide concentrations (up to 5 nM) (Fig. 2 K, L). The hIL-6-induced STAT3 phosphorylation remained unaffected by any parthenolide concentration applied (supplementary Fig. 2 A, B). AKT is also activated by PTEN[-/-], another regeneration-promoting approach [4, 37, 43, 53]. To investigate microtubule detyrosination in PTEN[-/-] RGC neurites, we prepared RGC cultures from PTEN-floxed mice after intravitreal AAV2-Cre injection. As expected, PTEN[-/-] enhanced tubulin detyrosination in axon tips (Fig. 3 A, B, C). This effect was counteracted by par-

thenolide, which further improved neurite outgrowth of PTEN$^{-/-}$-RGCs concentration-dependently (Fig. 3 B, C, D, E). Moreover, activation of the mTOR pathway, a significant contributor to the PTEN$^{-/-}$-mediated effects on axon growth, was not compromised by parthenolide, reflected by unaltered activation of ribosomal protein S6 in these RGC cultures (supplementary Fig. 2 C, D). Thus, parthenolide counteracts AKT/GSK3-induced microtubule detyrosination and synergistically enhances hIL-6, or PTEN$^{-/-}$-stimulated neurite growth.

Dimethylamino-parthenolide accelerates optic nerve regeneration

To test whether parthenolide can accelerate CNS regeneration *in vivo*, we used its water-soluble prodrug dimethylamino-parthenolide (DMAPT), which can cross the blood-brain barrier [27, 29, 50]. To this end, adult mice were subjected to an optic nerve crush (ONC) and received daily intraperitoneal injections of either vehicle or DMAPT (2 µg/kg) (Fig. 4 A). The number of CTB-labeled, regenerated axons was quantified in optic nerve sections 2 weeks post-injury. Compared to vehicle-treated controls, more axons reached distances of up to 0.5 mm distal to the lesion site after DMAPT application, indicating a slight but significant effect on regeneration (Fig. 4 B-E). At the same time, the RGC survival was not affected (Fig. 4 F, G). We then repeated these experiments in combination with intravitreally applied AAV2-hIL-6 (Fig. 4 H). Two weeks after injury, retinal cross-sections revealed robust phosphorylation of STAT3 in AAV2-hIL-6-treated RGCs, which remained unaffected by DMAPT treatment (Fig. 4 I). As reported previously [35], hIL-6 treatment alone enabled substantial regeneration with CTB-labeled axons reaching distances of up to 1.5 mm 2 weeks after injury (Fig. 4 J-O). Notably, additional DMAPT application at 2 µg/kg increased the number of axons at 1.5 mm fivefold, with some axons reaching distances of 2.5 mm within the same time after injury (Fig. 4 J, K, N, O). As observed in RGC cultures, DMAPT acted dose-dependently, with less pronounced effects at lower (0.2 µg/kg) or higher (20 µg/kg) dosages (supplementary Fig. 3 A, B). Thus, the systemic application of DMAPT sufficiently enhances and accelerates the effect of hIL-6 on optic nerve regeneration.

Systemic DMAPT application accelerates raphespinal regeneration after spinal cord crush

Hyper-IL-6-mediated axonal regeneration of raphespinal fibers enables functional recovery after a severe spinal cord crush (SCC) [40]. Therefore, we investigated whether systemic DMAPT treatment also affects raphespinal regeneration or functional recovery. To this end, we performed a complete SCC at the thoracic vertebra 8 (T8) level, followed by daily intraperitoneal injections of either vehicle or DMAPT at 2 µg/kg (Fig. 5 A). Hindlimb movement was analyzed using the Basso Mouse Scale (BMS) [3] over 8 weeks after injury. One day after SCC, BMS scores dropped to 0, indicating the absence of hindlimb movement (Fig. 5 B, C). After 2 weeks, most vehicle-treated animals recovered from the injury-induced spinal shock and reestablished typical extensive ankle movement (BMS: 2) (Fig. 5 B, C; supplementary video 1). In contrast, DMAPT treatment slightly but significantly improved functional recovery. Most of these animals showed plantar hind paw placement as early as 2 weeks after injury, occasionally accompanied by bodyweight support (BMS: 3) (Fig. 5 B, C; supplementary video 2). For this reason, the spinal cord tissue of some animals was analyzed for serotonergic fiber regeneration at this early time. The absence of serotonergic fibers at ~10 mm distal to the lesion site verified the lesion completeness in each animal included in this study to rule out the presence of spared raphespinal axons (Fig. 5 D, supplementary Fig. 4 A-H). Vehicle-treated control animals showed only typical short-distance sprouting of serotonergic axons less than 2 mm distal to the lesion site (Fig. 5 E-G, K). At the same time, DMAPT treatment increased the number and length of axons, with the longest axons reaching 4 mm (Fig. 5 E, H-K). When analyzed 8 weeks after SCC (Fig. 6 A), DMAPT further accelerated raphespinal regeneration, with the longest axons reaching >7 mm (Fig. 6 B-M). However, despite the more pronounced anatomical regeneration, the BMS score was only slightly further increased compared to 14 days after SCC (Fig. 6 N). In addition, the lesion size and CSPG expression at the glial scar remained unaffected by DMAPT treatment (supplementary Fig. 5 A-D). Thus, systemic DMAPT application accelerates raphespinal regeneration and some functional recovery after a severe spinal cord injury.

Raphespinal regeneration contributes to DMAPT-mediated functional recovery

We prepared additional mice to investigate whether the DMAPT-mediated effects on functional recovery depend on serotonergic fiber regeneration, as previously shown for hIL-6 [40]. To this end, we performed SCC with either vehicle or DMAPT treatment. After the DMAPT group had reached maximum functional recovery, we injected the serotonergic neurotoxin 5,7-dihydroxytryptamine (DHT), which depletes raphe neurons as early as one day after injection, as shown previously [32, 40]. Expectedly, DHT treatment reduced the number of serotonergic brainstem raphe neurons and their axons without affecting other neuronal populations (Fig. 7 A-J; supplementary Fig. 6 A, B). This did not affect BMS scores of vehicle-treated controls (Fig. 7 K, L; supplementary video 3) but immediately abolished DMAPT-induced gain of motor function (Fig. 7 K, L; supplementary video 4), suggesting an essential contribution of regenerated serotonergic axons.

DMAPT accelerates hIL-6-induced functional recovery after complete spinal cord crush.

Intracortical injection of AAV2-hIL-6 promotes axon regeneration and functional recovery after SCC [40]. To test for synergistic effects, as seen in the optic nerve (Fig. 4), we combined hIL-6 and DMAPT treatment. To this end, we subjected wild-type mice to SCC and injected AAV2-hIL-6 into the sensorimotor cortex. Mice received daily intraperitoneal injections of either vehicle or DMAPT for 8 weeks (Fig. 8 A). As determined by immunostaining, AAV2-hIL-6-mediated transduction of layer V cortical motorneurons induced robust phosphorylation of STAT3 (supplementary Fig. 7 A-E). Consistent with [40], intracortical AAV2-hIL-6 also transneuronally stimulated medullary raphe neurons (supplementary Fig. 7 I-M). At the same time, DMAPT treatment did not affect STAT3 phosphorylation in cortical (supplementary Fig. 7 B, F, G, H) or raphespinal (supplementary Fig. 7 J, N, O, P) neurons at applied doses used in this study. Consistent with previous results [40], AAV2-hIL-6 treatment restored plantar hind paw placement and stepping with lift-off, followed by forwarding limb advancement and re-establishment

of bodyweight support, reaching a maximum BMS score of 4-5 (Fig. 8 B-C). Interestingly, combinatorial treatment with DMAPT accelerated functional recovery, allowing plantar steps in most animals already 14 d after SCC. These mice reached the maximum effect already at 21 days after injury, while animals solely treated with hIL-6 needed 42 days (Fig. 8 B-C; supplementary video 5, 6). Moreover, additional DMAPT treatment also reduced the variance, with fewer weakly responding animals, resulting in a higher average final score of 4 compared to 3.5 in the hIL-6 control group.

DMAPT affects RpST fiber regeneration after spinal cord crush

Regarding fiber regeneration, additional DMAPT treatment further enhanced the hIL6-induced regeneration of serotonergic fibers 8 weeks after SCC (Fig. 8 D-K). Compared to AAV2-hIL-6 therapy, which led to robust axonal regeneration up to 7 mm distal from the lesion site (comparable to [40]) (Fig. 8, D-G, K), additional DMAPT treatment significantly increased the number of serotonergic axons at this distance, with some axons reaching even 8 mm (Fig. 8 D, H-K).

Discussion

Here we demonstrate that the systemic application of DMAPT facilitates axon regeneration in the injured optic nerve and accelerates functional recovery after severe spinal cord injury. Parthenolide's effects on CNS regeneration in vivo were confirmed in cell culture using primary RGCs from adult mice and humans. As on sensory neurons (preprint server) [21], parthenolide reduces microtubule detyrosination in axonal tips, which increases their dynamics and promotes axon extension [20]. As for sensory neurons, this effect is likely mediated by the inhibition of VASH1/VASH2, the enzymes catalyzing the detyrosination [21]. Consistently, the effective concentration range and doses of parthenolide/DMAPT to accelerate axon regeneration of RGC, serotonergic fibers in the spinal cord, and functional recovery was similar to sensory neurons and regeneration in the injured peripheral nerve [21]. Parthenolide did not affect CRMP2 or STAT3 activity at the nanomolar concentrations used in this study, both critical for CNS axon regeneration [35, 37] and suggesting the reduction of microtubule detyrosination causes the axon growth-promoting effect.

Systemic DMAPT treatment significantly promotes axon extension in the optic nerve and regeneration of serotonergic fibers beyond the injury site. The selective elimination of serotonergic fibers by DHT [8, 11, 32] abolishes the recovered locomotion, demonstrating that the improved regeneration of serotonergic fibers is essential for recovery after SCC. These data are consistent with previously published data, showing a similar effect on AAV-hIL-6 mediated recovery [40]. Despite the fact that DMAPT-mediated functional improvement depends on the regeneration of serotonergic fibers, we cannot exclude the possibility that spinal tracts other than the raphespinal tract contributed as well. The fact that DMAPT improved axon regeneration in the optic nerve and the spinal cord also implies that sufficient amounts of the drug reached the axonal growth cones, where the drug affects the microtubule detyrosination [20]. It is likely that an oral application of DMAPT, which reportedly is >60% bioavailable [21, 27, 57], has similar effects on nerve regeneration, bolstering the potential as a drug to treat axonal CNS damage.

Another significant finding of the current study is that pro-regenerative hIL-6 and PTEN$^{-/-}$ treatments induce microtubule detyrosination in RGC axons. Given that a parthenolide-mediated reduction of microtubule detyrosination promotes axon growth, these observations suggest that this effect of hIL-6 or PTEN$^{-/-}$ compromises their full axon regeneration-promoting potential. hIL-6 and PTEN$^{-/-}$ induced microtubule detyrosination is likely caused by inhibitory GSK3 phosphorylation via increased PI3K/AKT activity. As a result, MAP1B activity is decreased, and the axonal detyrosination rate is elevated [26, 30, 36]. However, CNTF does not affect detyrosination due to its lower AKT activation than hIL-6 [35, 48, 49]. At the same time, hIL-6 and PTEN$^{-/-}$ mediated GSK3 inhibition confers disinhibitory effects toward myelin-associated molecules because less CRMP2 is inhibited [36, 37, 44]. Thus, the disinhibitory effect of GSK3 inhibition promotes regeneration, while the detyrosinating effect on microtubules rather compromises it. Therefore, by counteracting detyrosination, parthenolide/DMAPT cotreatment further increases the axon growth-promoting effect of hIL-6. Consistently, effective parthenolide concentrations reduce detyrosination levels in hIL-6-treated and PTEN$^{-/-}$ RGCs, and combined treatment leads to more substantial neurite growth than each treatment alone. This notion is also supported by the fact that PTEN$^{-/-}$ and hIL-6-stimulated neurons sustain parthenolide-induced additive elevation of axon growth even at higher parthenolide concentrations, which already diminish outgrowth on naïve or CNTF-stimulated RGCs. The level of detyrosinated tubulin at the maximal applied parthenolide concentration in hIL-6 treated RGCs was still similar to untreated control RGCs. The synergistic effects of DMAPT and hIL-6 were also reproduced *in vivo*, in the optic nerve and spinal cord. Thus, parthenolide/DMAPT neutralizes the hIL-6-mediated limiting effect of decreased microtubule dynamics and synergistically accelerates CNS regeneration.

Our findings also indicate that DMAPT mainly accelerates axon growth and has little effect on the intrinsic regenerative state of neurons. Therefore, the RGC regeneration-promoting effects of parthenolide on its own are only moderate but synergistically increase hIL-6-mediated axon extension. Thus, DMAPT enabled substantial regeneration of serotonergic RpST after complete SCC. Furthermore, the length of regenerating RpST axons after SCC further increased

from 2 to 8 wpc, while the functional recovery was already almost maximal in the early phase. This observation might be based on the different possibilities of achieving active recovery. On the one hand, descending supraspinal axons might need to grow over long distances to reestablish functional connections within the lumbar central pattern generator (CPG), a network of neurons responsible for generating stereotyped hindlimb movement [10, 31, 62]. On the other hand, it is also possible that severed axons only need to overcome short distances of a few spinal segments and then connect to existing propriospinal neurons, which might serve as a relay station to the CPG [2, 9, 10]. An observation accounting for the latter possibility is increased axonal sprouting in the region of the lumbar CPG (~6-7 mm distal to the lesion) at 8 wpc, whereas the substantial increase in functional recovery occurred already earlier (~2-3 wpc) before the axons had reached the CPG. Thus, DMAPT-induced long-distance RpST regeneration might not be essential for its beneficial effects on functional recovery, whereas DHT experiments showed that RpST regeneration, in general, is crucial.

In conclusion, systemically applied DMAPT measurably accelerates CNS regeneration and functional recovery and overcomes the limiting effect of AKT/GSK3 on elevating microtubule detyrosination. Whether DMAPT treatment can also improve functional recovery in more clinically relevant injury models, such as contusion injuries, or even more effective in combination with other regeneration-promoting strategies [6, 17, 33] will be investigated in the future. Finally, the axon growth-promoting effects on primary adult human CNS neurons underline the potential of parthenolide/DMAPT as a possible future treatment strategy for CNS injury in humans.

Fig. 1: Parthenolide and CNTF synergistically promote neurite growth of murine and primary adult human RGCs

A) Images of βIII-tubulin (tubulin) positive RGCs cultured for 4 days in the presence of either vehicle (-) or parthenolide (par; 0.5 nM or 5 nM). Scale bar: 15 µm

B) Quantification of neurite growth of cultures described in **A** with par concentrations between 0.25 and 5 nM. Data were normalized to untreated controls with an average neurite length of 4.6 µm per RGC and represent means ± SEM of six independent experiments.

C) Quantification of RGC numbers per well in cultures described in **A**.

D) Images of βIII-tubulin positive RGCs cultured for 4 days in the presence of vehicle (-) or par (0.5 nM), either without (con) or in combination with CNTF (200 ng/ml). Scale bar: 50 µm.

E) Quantification of neurite growth in cultures described in **D**. Data were normalized to un-treated controls with an average neurite length of 9.6 µm per RGC and represent means ± SEM of seven independent experiments.

F) Representative images of βIII-tubulin (tubulin; green) positive RGCs and phosphorylated STAT3 (pSTAT3, red) in cultures described in **D**. Insets show higher magnifications of selected cells, indicated by white arrows. CNTF, but not par, induced STAT3 phosphorylation. Scale bars: 15 µm

G) Quantification of pSTAT3 positive RGCs in cultures described in **D**. Only CNTF, but not par, affected pSTAT3 levels. Data represent means ± SEM of three independent experiments.

H) Representative images of RGC neurites from cultures described in **D**, immunohistochemically stained for βIII-tubulin (green) and detyrosinated α-tubulin (detyr, red). White arrows indicate the last 15 µm from axon tips after respective treatments. Scale bar: 15 µm.

I) Percentages of detyrosinated tubulin-positive (detyr$^+$) axon tips. Par but not CNTF reduced detyrosination. Data represent means ± SEM of three independent experiments.

J) βIII-tubulin (tubulin) stained human RGCs isolated from the eyes of adult human patients and cultured for 4 days in the presence or absence of CNTF (200 ng/ml) and/or parthenolide (par, 0.5 nM). Scale bar: 100 µm.

K) Quantification of the average neurite length per RGC in cultures depicted in **J**. Par was applied at indicated concentrations, either alone or combined with CNTF. Data were normalized to untreated controls with an average neurite length of 16.3 µm per RGC and represent means ± SEM of four technical replicates, each from two independent experiments with individual human eyes (n=8).

L) Quantification of RGCs per well in cultures described in **J**, **K**.

M) Images of axon tips from RGCs as described in **J**. Cells were cultured in the absence (-) or presence of par (0.5 nM) and then immunostained for detyrosinated α-tubulin (detyr, red) and βIII-tubulin (tubulin, green). White arrows indicate detyrosinated tubulin-positive or negative neurite tips in control and the par-treated groups. Scale bar: 10 µm.

N) Quantification of axon tips containing detyrosinated tubulin from RGCs described in **J**. Thirty neurite tips per group were analyzed in three technical replicates from one- two independent experiments with individual human eyes (n=3-6).

Significances of intergroup differences in **B**, **C**, **E**, **G**, **I**, **K**, and **N** were analyzed using a one-way (**B**, **C**), two-way (**E, G, I**, **K**, **N**) analysis of variance (ANOVA) followed by the Tukey or Holm-Sidak post hoc test. P-values indicate statistical significance compared to the untreated (black p-values) or the CNTF-treated (red p-values) groups. n.s.=non-significant. Dots in B, C, E, G, and I represent values from at least three independent experiments. Dots in K, L, and N, represent values from technical replicates, each from two independent experiments.

Fig. 2: hIL-6 elevates microtubule detyrosination

A) Schematic drawing illustrating the effect of hIL-6 on the PI3K/AKT/GSK3 signaling pathway. hIL-6 binding to gp130 activates phosphatidylinositol 3-kinase (PI3K), converting phosphatidylinositol (4, 5)-bisphosphate (PIP$_2$) to phosphatidylinositol (3, 4, 5)-trisphosphate (PIP$_3$). PIP$_3$ stimulates AKT. Subsequent effects on GSK3, CRMP2, and microtubule detyrosination are shown in **B**-**L**.

B) Western blots of optic nerve lysates from either untreated mice (con) or animals that had received intravitreal AAV2-hIL-6 injections 3 weeks earlier. Retinal hIL-6 expression elevated inhibitory phosphorylation of GSK3α and GSK3β, while total GSK3 levels remained unaffected. Inhibitory CRMP2 phosphorylation was reduced without altering total CRMP2 levels, while detyrosinated tubulin levels were increased. βIII-tubulin (tubulin) served as a loading control.

C-H) Densitometric quantification of western blots shown in **B** relative to βIII-tubulin and normalized to the untreated control (con). Data represent means ± SEM of samples from at least three animals per group.

I) Representative images of axon tips from dissociated RGCs from mice as described in **B**. RGCs were cultured for 4 days in the presence of either vehicle (-) or parthenolide (par; 5 nM) and were immunohistochemically stained for βIII-tubulin (tubulin, green) and detyrosinated α-

tubulin (detyr, red)**.** White arrows indicate the last 15 μm of detyrosinated or negative axon tips after respective treatments. Scale bar: 15 μm

J) Quantification of the percentages of detyrosinated tubulin-positive (detyr$^+$) axon tips. Par was applied at indicated concentrations. hIL-6 treatment increased tubulin detyrosination, while par blocked this effect. Data represent means ± SEM of 3 technical replicates from 2 independent experiments. P-values refer to untreated control.

K) Representative images of βIII-tubulin (tubulin) positive RGCs from cultures as described in **I**, cultured in the presence or absence of parthenolide (par; 0.5 nM or 5 nM). Scale bar: 50 μm

L) Quantification of the average neurite length per RGC in cultures depicted in **K**. Par was applied at indicated concentrations in combination with hIL-6. Data were normalized to un-treated controls with an average neurite length of 8.4 μm per RGC and represent means ± SEM of three technical replicates from two independent experiments.

Significances of intergroup differences were evaluated using Student's t-test for quantifications shown in **C-H**. A one-way analysis of variance (ANOVA) followed by a Holm-Sidak or Tukey post hoc test is shown in **J** and **L**, respectively. P-values indicate statistical significance com-pared to the untreated (black p-values) or the vehicle-treated hIL-6 (red p-values) groups. n.s.=non-significant. Dots represent values from single animals in **C-H** and three technical rep-licates of two independent experiments, respectively, in **J** and **L**.

Fig. 3: PTEN$^{-/-}$ increases microtubule detyrosination

A) Schematic drawing illustrating the effect of PTEN$^{-/-}$ on the PI3K/AKT/GSK3 signaling pathway. PTEN$^{-/-}$ promotes the conversion of phosphatidylinositol (4, 5)-bisphosphate (PIP$_2$) to phosphatidylinositol (3, 4, 5)-trisphosphate (PIP$_3$). PIP$_3$ stimulates AKT and thus induces inhibitory GSK3 phosphorylation. As a result, the phosphorylation of CRMP2 and MAP1B is reduced. The effect on tubulin detyrosination is illustrated in **B-E**.

B) Representative images of axon tips from dissociated RGCs of PTEN-floxed mice. Animals were either untreated (con) or had received intravitreal injections of AAV2-Cre 3 weeks before dissociation (PTEN$^{-/-}$). RGCs were cultured for 4 days in the presence of either vehicle (-) or parthenolide (par; 5 nM) and were immunohistochemically stained for βIII-tubulin (tubulin, green) and detyrosinated α-tubulin (detyr, red). White arrows indicate the last 15 µm of axon tips. Scale bar: 15 µm

C) Quantification of the percentages of detyrosinated tubulin-positive (detyr $^+$) axon tips. Par was applied at indicated concentrations. PTEN$^{-/-}$ increased tubulin detyrosination, while par blocked this effect. Data represent means ± SEM of 3 technical replicates from 2 independent experiments.

D) Representative images of βIII-tubulin (tubulin) positive RGCs from cultures as described in **B**, cultured in the presence or absence of parthenolide (par; 0.5 nM or 5 nM). Scale bar: 50 µm

E) Quantification of the average neurite length per RGC in cultures depicted in **D**. Par was applied at indicated concentrations in combination with PTEN$^{-/-}$. Data were normalized to untreated controls with an average neurite length of 1.04 µm per RGC and represent means ± SEM of three technical replicates from two independent experiments.

The significance of intergroup differences was evaluated through a one-way analysis of variance (ANOVA) followed by a Holm-Sidak or Tukey post hoc test for data shown in **C** and **E**, respectively. P-values indicate statistical significance compared to the untreated (black p-values) or the vehicle-treated PTEN$^{-/-}$ (red p-values) groups. n.s. = non-significant. Dots represent values from three technical replicates of two independent experiments.

Fig. 4: DMAPT promotes optic nerve regeneration and enhances the effect of hIL-6

A) Timeline showing surgical interventions and intraperitoneal (i.p.) drug application for experiments presented in **B**-**G**.

B) Longitudinal optic nerve sections containing CTB (white) traced axons from mice 2 weeks after optic nerve crush (ONC) and daily repeated intraperitoneal vehicle (veh) or DMAPT injections, as described in **A**. Red asterisks indicate the lesion sites. Scale bar: 100 µm.

C) Quantification of regenerating axons at indicated distances distal to the lesion site in optic nerves as depicted in **B**, showing significantly enhanced optic nerve regeneration after DMAPT

treatment compared to controls. Values represent the mean ± SEM of 6-7 animals per group (veh: n=6, dmapt: n=7).

D, E) Higher magnification of dashed boxes as indicated in **B**. Scale bar: 50 µm.

F) Confocal scans of retinal wholemounts from mice either uninjured (con) or subjected to ONC 14 days before tissue isolation. Animals received daily intraperitoneal injections of vehicle (veh) or DMAPT. RGCs were visualized by immunohistochemical βIII-tubulin staining (tubulin). Scale bar: 20 µm.

G) Quantification of RGC density in retinal wholemounts, as described in **E**. DMAPT, did not affect RGC survival in uninjured mice or after ONC. Values represent the mean ± SEM of 5-9 animals per group (veh, con: n=5; dmapt, con: n=5; veh ONC: n=7; dmapt, ONC: n=9).

H) Timeline showing surgical interventions and intraperitoneal (i.p.) drug application for experiments presented in **I-K** and **Supplementary Fig. 3**.

I) Immunostained retinal cross-sections prepared from mice that had received an intravitreal injection of AAV2-hIL-6-GFP simultaneously to optic nerve crush (ONC) and then daily intraperitoneal injections of vehicle or DMAPT (2 µg/kg) for 2 weeks. Sections were immunostained for phosphorylated STAT3 (pSTAT3, red). Transduction with AAV-2 hIL-6 is visualized by GFP (green) staining. hIL-6-induced STAT3-phosphorylation was not affected by DMAPT cotreatment. Scale bar: 20 µm.

J) Longitudinal optic nerve sections containing CTB (white) traced axons from mice described in **A**. Red asterisks indicate the lesion sites. Scale bar: 100 µm.

K) Quantification of regenerating axons distal to the lesion site at indicated distances in optic nerves from mice that received AAV2-hIL-6 combined with vehicle or DMAPT treatment, as shown in **H** and **I**. DMAPT significantly enhanced hIL-6-induced optic nerve regeneration. Values represent the mean ± SEM of 6-7 animals per group (veh: n=6; dmapt: n=7).

L-O) Higher magnification of dashed boxes as indicated in **J**. Scale bar: 50 µm.

Significances of intergroup differences in **C**, **G**, and **K** were evaluated using a two-way analysis of variance (ANOVA) with a Tukey post hoc test. P-values indicate statistical significance. Dots represent values from single animals.

Figure 5: DMAPT accelerates RpST regeneration and functional recovery after spinal cord crush

A) Timeline showing surgical interventions, BMS testing, and intraperitoneal (i.p.) drug application for experiments presented in **B**-**K**.

B) Representative photos showing open field hindlimb movement of mice at 1, 7, and 14 days post-injury (dpi). Before, mice had received spinal cord crush (SCC) and daily intraperitoneal injections of either DMAPT (dmapt) or the vehicle (veh).

C) BMS score of animals as described in **A** and **B** over 2 weeks after SCC. DMAPT significantly improved hindlimb movement's functional recovery, resulting in plantar paw placement in most animals 2 weeks after spinal cord crush. Values represent the mean ± SEM of 13-17 animals per group (veh: n=13; dmapt n=17).

D) Immunohistochemical staining of transverse spinal cord sections from mice treated as described in **A** and **B** 2 weeks after injury. Serotonergic (5-HT) axons are present in the raphespinal tract proximal but not distal to the lesion site, indicating lesion completeness. Scale bar: 200 μm

E) Coronal thoracic spinal cord sections isolated from mice 2 weeks after SCC and daily intraperitoneal injections of either dmapt or the veh. Raphespinal tract (RpST) axons were stained

using an anti-serotonin antibody (5-HT, white). Only dmapt but not vehicle-treated mice showed regeneration of serotonergic axons > 2 mm beyond the lesion site (red arrow). Scale bar: 500 µm.

F-J) Higher magnifications of dashed boxes as indicated in **E**. Scale bar: 50 µm.

K) Quantification of regenerating 5-HT-positive axons from animals described in **A** and **E** at indicated distances beyond the lesion. Values represent the mean ± SEM of 7-8 mice per group (veh: n=8; dmapt: n=7).

Significances of intergroup differences in **C** and **K** were evaluated using a two-way analysis of variance (ANOVA) with a Tukey post-hoc-test. P-values indicate statistical significance.

Figure 6: Spinal cord regeneration and functional recovery after long-term DMAPT treatment

A) Timeline showing surgical interventions and intraperitoneal (i.p.) drug application for experiments presented in **B**-**N**.

B) Coronal thoracic spinal cord sections isolated from mice 8 weeks after spinal cord crush (SCC) and daily intraperitoneal injections of either DMAPT (dmapt) or the vehicle (veh). Raphespinal tract (RpST) axons were stained using an anti-serotonin antibody (5-HT, white). In contrast to the vehicle-treated control, dmapt enabled RpST regeneration over long distances beyond the lesion site (red arrow). Scale bar: 500 µm.

C-L) Higher magnification of dashed boxes indicated in **B**. Scale bar: 50 µm.

M) Quantification of regenerating 5-HT-positive axons as described in **A** and **B** at indicated distances beyond the lesion. Values represent the mean ± SEM of 7-10 animals per group (veh: n=7; DMAPT: n=10).

N) BMS score of animals as described in **A** and **B**, over 8 weeks after spinal cord injury. DMAPT significantly improved functional recovery of hindlimb movement, reaching a plateau 2 weeks after SCC. Values represent the mean ± SEM of 7-10 animals per group (veh: n=7; DMAPT: n=10).

Significances of intergroup differences in **M** and **N** were evaluated using a two-way analysis of variance (ANOVA) with a Tukey post-hoc-test. P-values indicate statistical significance.

Figure 7: DMAPT-mediated functional recovery is RpST-dependent

A) Validation of DHT-mediated depletion of serotonergic (5-HT, red) neurons in raphe nuclei (NRM = nucleus raphe magnus, NRPa=nucleus raphe pallidus, NRO=nucleus raphe obscurus) located above the pyramidal tracts in coronal medullary sections from mice as depicted in **K, L** 1 week after intracerebroventricular 5, 7-dihydroxytryptamine (DHT) injection, or untreated controls (con). Scale bar: 100 µm.

B-G) Higher magnification of indicated areas in images, as shown in A. Scale bar: 50 µm.

H) Quantification of serotonin (5-HT) positive raphe neurons per section in medullary immunostainings for untreated (con: n=5) and DHT-injected (DHT: n=8) mice as described in **A**.

I) Thoracic spinal cord cross-sections from animals, as described in **A**. Raphespinal axons were visualized by serotonin (5-HT) staining. Scale bar: 500 µm.

J) Quantification of 5-HT staining as sown in **I** for untreated (con: n=5) and DHT-injected (DHT, n=8) mice.

K) Representative images of vehicle- (veh) or DMAPT- (dmapt) treated mice at 7 or 21 d post-injury (dpi) and 1 day after DHT treatment.

L) BMS score of animals treated as described in **(K)** at indicated time points after spinal cord crush and DHT treatment. Values represent means ± SEM of 7-8 animals per group (veh: n = 8; dmapt: n=7), showing the average left and right hind paws score.

Significances of intergroup differences were evaluated using a two-way analysis of variance (ANOVA) with a Holm Sidak post hoc test **(H, L)** or Student's t-test **(J)**. The significance between group means is indicated by p-values. In **L,** comparisons are shown within the vehicle group (black), within the dmapt group (red), and between groups (blue).

Figure 8: DMAPT treatment accelerates hIL-6-induced RpST regeneration and functional recovery after spinal cord crush despite increased CST dieback

A) Timeline showing surgical interventions and intraperitoneal (i.p.) drug application for experiments presented in **B-K** and **Fig. 11 A-B**.

B) Representative photos showing open field hindlimb movement of mice at 1, 14, 35, and 56 days post-injury (dpi) of animals as described in **A**.

C) BMS scores of animals as described in **A** over 8 weeks after SCC. DMAPT treatment accelerated functional recovery of hindlimb movement in hIL6-treated mice. Values represent the mean ± SEM of 10 animals per group (veh: n = 10; DMAPT: n = 10).

D) Sagittal thoracic spinal cord sections isolated from mice described in **A**, 8 weeks after spinal cord crush (SCC), followed by intracortical AAV2-hIL-6 injection and daily intraperitoneal injection of either vehicle (veh) or DMAPT (dmapt). Raphespinal tract axons were stained using an anti-serotonin antibody (5-HT, white). DMAPT treatment did not significantly affect axon regeneration compared to vehicle-treated controls. Scale bar: 250 μm

E-J) Higher magnifications of dashed boxes as indicated in **D**. Scale bar: 100 μm

K) Quantification of regenerated 5-HT positive axons from animals described in **A** at indicated distances beyond the lesion site. DMAPT did not significantly enhance hIL-6-mediated raphespinal axon regeneration further. Values represent the mean ± SEM of 7-10 animals per group (veh: n = 7; DMAPT: n = 10).

Values represent means ± SEM of 7 or 10 (veh: n = 7; dmapt: n = 10) animals per group. Dots represent values for single animals. Significances of intergroup differences in **C**, **K**, and **M** were evaluated using a two-way analysis of variance (ANOVA) with a Tukey post-hoc- test. P-values indicate statistical significance.

Supplementary Figure 1: Parthenolide does not affect CRMP2 phosphorylation

A) Representative photographs of dissociated RGCs stained for T514-phosphorylated CRMP2 (pCRMP2; red) and βIII-tubulin (tubulin; green) after 4 d in culture. RGCs were isolated from WT mice and either incubated with 0.5 nM parthenolide (par) or the vehicle (veh). Scale bar: 50 μm.

B) Quantification of pCRMP2 staining intensities of cultured RGCs as described in **A**. Integrated density of pCRMP2 staining was measured and normalized against values of the vehicle-treated control. Values represent means ± SEM of three independent experiments per group. In each experiment, 80 RGCs per group were analyzed in two replicate wells. Statistical analysis using Student's t-test revealed no significant differences between the two groups.

Supplementary Figure 2: Parthenolide does not affect STAT3 or mTOR signaling

A) Representative images of βIII-tubulin positive RGCs (green) and phosphorylated STAT3 (pSTAT3, red) from cultures as described in **Fig. 2 I-L**. hIL-6, but not par, induced STAT3 phosphorylation. Scale bar: 20 µm

B) Quantification of pSTAT3 positive RGCs in cultures described in **C**. Data represent means ± SEM of 3 technical replicates from 2 independent experiments.

C) Representative images of βIII-tubulin positive RGCs (tubulin; green) and phosphorylated S6 (pS6; red) from cultures as described in Fig. **3 B-E**. PTEN⁻/⁻, but not par, induced S6 phosphorylation. Scale bar: 20 µm

D) Quantification of pS6 positive RGCs in cultures as described in C. Data represents means ± SEM of 3 technical replicates from 2 independent experiments.

Significances of intergroup differences in **B** and **D** were evaluated by a two-way analysis of variance (ANOVA) followed by a Tukey post hoc test. P-values indicate statistical significance. n.s.=non-significant. Dots represent values from three technical replicates of two independent experiments.

Supplementary Figure 3: Optic nerve regeneration after AAV2-hIL-6 application and DMAPT treatment at suboptimal concentrations

A) Quantification of regenerating axons distal to the lesion site at indicated distances in optic nerves from mice that received AAV2-hIL-6 delivery combined with DMAPT treatment, as shown in **6 A**. DMAPT was applied at 0.2 µg/kg. Regeneration was slightly enhanced compared to the vehicle-treated group (dashed line, as depicted in **Fig. 4 C, D**). Values represent the mean ± SEM of 4 animals.

B) Quantification of regenerating axons distal to the lesion site at indicated distances in optic nerves from mice that received AAV2-hIL-6 delivery combined with DMAPT treatment, as shown in **6 A**. DMAPT was applied at 20 µg/kg. Regeneration was not improved compared to the vehicle-treated group (dashed line, as depicted in **Fig. 4 C, D**). Values represent the mean ± SEM of 5 animals.

Significances of intergroup differences in **A** and **B** were evaluated using a two-way analysis of variance (ANOVA) with a Tukey post hoc test. P-values indicate statistical significance. Dots represent values from single animals.

Supplementary Figure 4: Verification of spinal cord lesion completeness

A) Schematic drawing showing the location of spinal cord cross sections proximal and distal to the T8 crush to identify potentially spared axons from the raphespinal tract (red).

B) Schematic drawing illustrating the expected pattern of serotonergic axons (red) on proximal spinal cord cross sections and an absence of axons on distal sections in case of a complete lesion.

C-H) Images of spinal cord cross sections proximal and distal to the crush site of animals as described in **Fig. 5 A**, **Fig. 6 A,** and **Fig. 8 A**. All animals in these experiments showed the expected pattern of serotonin-positive axons (5-HT, white) in the thoracic spinal cord > 3 mm proximal to the lesion site. No spared serotonergic axons were observed in the lumbar spinal cord > 8 mm distal to the lesion site, verifying the completeness of the crush injury. Scale bars: 500 μm.

Supplementary Figure 5: DMAPT does not affect lesion size or CSPG release after SCC

A) Representative coronal spinal cord sections showing the T8 lesion site 8 weeks after SCC and daily intraperitoneal application of vehicle (veh) or DMAPT, as described in **Fig. 6 A**. Lesion sites and CSPGs were visualized by GFAP or CS-56 staining, respectively. Scale bar: 200 µm**.**

B-C) Quantification of the lesion area (**B**) and the lesion width (**C**) on spinal cord sections described in **A**. **D)** Quantification of the CS-56-positive (CS-56[+]) (CSPGs) area on spinal cord sections described in **A**. Statistical analyses in **B-D** were performed using a Student's t-test.

n.s. = non-significant. Data represent means±SEM of 6 (veh) or 5 (DMAPT) animals per group.

Dots represent values for single animals.

Supplementary Figure 6: DHT specifically depletes serotonergic neurons

A) Schematic of the Allen Brian Atlas illustrating the location of raphe nuclei (NRM= nucleus raphe magnus, NRPa = nucleus raphe pallidus) above the pyramidal tracts (Py) in a coronal view of the medulla. The white box indicates the location of images in B.

B) Sections from an untreated animal (con) and a mouse 1 day after intracerebroventricular 5,7-dihydroxytryptamine (DHT) injection. Neurons are immunostained for neuN (green), while Raphe neurons are visualized by serotonin (5-HT, red) immunostaining. Dapi (blue) labels all nuclei. DHT treatment depletes serotonergic neurons with only very few 5-HT positive neurons remaining (white arrows), while other neuN positive neurons are unaffected. Scale bar: 250 µm.

Supplementary Figure 7: DMAPT does not affect hIL-6-mediated STAT3 activation

A) Schematic of a coronal brain section with the AAV2-hIL-6 injection sites (dashed box) shown in **B.**

B) Section of layer V sensorimotor cortex isolated from mice 8 weeks after spinal cord crush (SCC), intracortical AAV2-hIL-6 injection, and subsequent daily intraperitoneal injections of either vehicle (veh) or DMAPT (dmapt). GFP (green) was co-expressed by AAV2-hIL-6. The tissue was immunohistochemically stained for phosphorylated STAT3 (pSTAT3, red). DMAPT treatment did not affect STAT3 phosphorylation. Scale bar: 50 μm

C-H) Higher magnification of dashed white boxes indicated in **B**. Scalebar: 50 μm.

I) Schematic drawing of a coronal medulla section showing transneuronal stimulation (STAT3 phosphorylation) of neurons in raphe nuclei (RN) with hIL-6 secreted from collateral sprouts from the pyramidal tract (PY).

J) Section of the medulla showing mice's medullary raphe nuclei (RN) as described in **B**. The tissue was immunohistochemically stained for pSTAT3 (red) and 5-HT (green). STAT3 phosphorylation indicates transneuronal stimulation of raphe neurons with hIL-6 secreted from pyramidal axon sprouts (PY). DMAPT treatment did not affect pSTAT3 levels. Scale bar: 50 μm.

K-P) Higher magnification of dashed white boxes indicated in **J**. Scalebar: 100 μm.

Literature

Aillaud C, Bosc C, Peris L, Bosson A, Heemeryck P, Dijk JV, Friec JL, Boulan B, Vossier F, Sanman LEet al (2017) Vasohibins/SVBP are tubulin carboxypeptidases (TCPs) that regulate neuron differentiation. Doi 10.1126/science.aao4165

Anderson MA, O'Shea TM, Burda JE, Ao Y, Barlatey SL, Bernstein AM, Kim JH, James ND, Rogers A, Kato Bet al (2018) Required growth facilitators propel axon regeneration across complete spinal cord injury. Nature 561: 396-400 Doi 10.1038/s41586-018-0467-6

Basso DM, Fisher LC, Anderson AJ, Jakeman LB, McTigue DM, Popovich PG (2006) Basso Mouse Scale for locomotion detects differences in recovery after spinal cord injury in five common mouse strains. J Neurotrauma 23: 635-659 Doi 10.1089/neu.2006.23.635

Bei F, Lee HHC, Liu X, Gunner G, Jin H, Ma L, Wang C, Hou L, Hensch TK, Frank Eet al (2016) Restoration of Visual Function by Enhancing Conduction in Regenerated Axons. Cell 164: 219-232 Doi 10.1016/j.cell.2015.11.036

Bradbury EJ, Moon LD, Popat RJ, King VR, Bennett GS, Patel PN, Fawcett JW, McMahon SB (2002) Chondroitinase ABC promotes functional recovery after spinal cord injury. Nature 416: 636-640 Doi 10.1038/416636a

Burnside ER, De Winter F, Didangelos A, James ND, Andreica EC, Layard-Horsfall H, Muir EM, Verhaagen J, Bradbury EJ (2018) Immune-evasive gene switch enables regulated delivery of chondroitinase after spinal cord injury. Brain 141: 2362-2381 Doi 10.1093/brain/awy158

Cafferty WB, Strittmatter SM (2006) The Nogo-Nogo receptor pathway limits a spectrum of adult CNS axonal growth. J Neurosci 26: 12242-12250 Doi 10.1523/JNEUROSCI.3827-06.2006

Capela JP, Lautenschlager M, Dirnagl U, Bastos ML, Carvalho F, Meisel A (2008) 5,7-Dihydroxitryptamine toxicity to serotonergic neurons in serum free raphe cultures. Eur J Pharmacol 588: 232-238 Doi 10.1016/j.ejphar.2008.04.046

Chen B, Li Y, Yu B, Zhang Z, Brommer B, Williams PR, Liu Y, Hegarty SV, Zhou S, Zhu Jet al (2018) Reactivation of Dormant Relay Pathways in Injured Spinal Cord by KCC2 Manipulations. Cell 174: 521-535.e513 Doi 10.1016/j.cell.2018.06.005

Courtine G, Song B, Roy RR, Zhong H, Herrmann JE, Ao Y, Qi J, Edgerton VR, Sofroniew MV (2008) Recovery of supraspinal control of stepping via indirect propriospinal relay connections after spinal cord injury. Nat Med 14: 69-74 Doi 10.1038/nm1682

Cummings KJ, Commons KG, Fan KC, Li A, Nattie EE (2009) Severe spontaneous bradycardia associated with respiratory disruptions in rat pups with fewer brain stem 5-HT neurons. Am J Physiol Regul Integr Comp Physiol 296: R1783-1796 Doi 10.1152/ajpregu.00122.2009

Danilov CA, Steward O (2015) Conditional genetic deletion of PTEN after a spinal cord injury enhances regenerative growth of CST axons and motor function recovery in mice. Exp Neurol 266: 147-160 Doi 10.1016/j.expneurol.2015.02.012

Du K, Zheng S, Zhang Q, Li S, Gao X, Wang J, Jiang L, Liu K (2015) Pten Deletion Promotes Regrowth of Corticospinal Tract Axons 1 Year after Spinal Cord Injury. J Neurosci 35: 9754-9763 Doi 10.1523/JNEUROSCI.3637-14.2015

Fischer D (2017) Hyper-IL-6: a potent and efficacious stimulator of RGC regeneration. Eye (Lond) 31: 173-178 Doi 10.1038/eye.2016.234

Fischer D, Leibinger M (2012) Promoting optic nerve regeneration. Prog Retin Eye Res 31: 688-701 Doi 10.1016/j.preteyeres.2012.06.005

Fischer D, Pavlidis M, Thanos S (2000) Cataractogenic lens injury prevents traumatic ganglion cell death and promotes axonal regeneration both in vivo and in culture. Invest Ophthalmol Vis Sci 41: 3943-3954

Fischer D, Petkova V, Thanos S, Benowitz LI (2004) Switching mature retinal ganglion cells to a robust growth state in vivo: gene expression and synergy with RhoA inactivation. J Neurosci 24: 8726-8740 Doi 10.1523/JNEUROSCI.2774-04.2004

Fischer M, Goldschmitt J, Peschel C, Brakenhoff JP, Kallen KJ, Wollmer A, Grotzinger J, Rose-John S (1997) I. A bioactive designer cytokine for human hematopoietic progenitor cell expansion. Nat Biotechnol 15: 142-145

Fonrose X, Ausseil F, Soleilhac E, Masson V, David B, Pouny I, Cintrat J-C, Rousseau B, Barette C, Massiot G et al (2007) Parthenolide Inhibits Tubulin Carboxypeptidase Activity. Doi 10.1158/0008-5472.CAN-06-3732

Gobrecht P, Andreadaki A, Diekmann H, Heskamp A, Leibinger M, Fischer D (2016) Promotion of Functional Nerve Regeneration by Inhibition of Microtubule Detyrosination. J Neurosci 36: 3890-3902 Doi 10.1523/JNEUROSCI.4486-15.2016

Gobrecht P, Gebel J, Hilla A, Gisselmann G, Fischer D (2022) VASH1/2 inhibition accelerates functional recovery of injured nerves. BioRxiv: Doi 10.1101/2022.09.20.507919

Gobrecht P, Leibinger M, Andreadaki A, Fischer D (2014) Sustained GSK3 activity markedly facilitates nerve regeneration. Nat Commun 5: 4561 Doi 10.1038/ncomms5561

Gonzalez-Billault C, Avila J, Caceres A (2001) Evidence for the role of MAP1B in axon formation. Mol Biol Cell 12: 2087-2098

Goold RG, Owen R, Gordon-Weeks PR (1999) Glycogen synthase kinase 3beta phosphorylation of microtubule-associated protein 1B regulates the stability of microtubules in growth cones. J Cell Sci 112 (Pt 19): 3373-3384

Grozdanov V, Muller A, Sengottuvel V, Leibinger M, Fischer D (2010) A method for preparing primary retinal cell cultures for evaluating the neuroprotective and neuritogenic effect of factors on axotomized mature CNS neurons. Curr Protoc Neurosci Chapter 3: Unit3 22 Doi 10.1002/0471142301.ns0322s53

Guo X, Snider WD, Chen B (2016) GSK3beta regulates AKT-induced central nervous system axon regeneration via an eIF2Bepsilon-dependent, mTORC1-independent pathway. Elife 5: e11903 Doi 10.7554/eLife.11903

Guzman ML, Rossi RM, Neelakantan S, Li X, Corbett CA, Hassane DC, Becker MW, Bennett JM, Sullivan E, Lachowicz JLet al (2007) An orally bioavailable parthenolide analog selectively eradicates acute myelogenous leukemia stem and progenitor cells. Blood 110: 4427-4435 Doi 10.1182/blood-2007-05-090621

Heinrich PC, Behrmann I, Haan S, Hermanns HM, Muller-Newen G, Schaper F (2003) Principles of interleukin (IL)-6-type cytokine signalling and its regulation. Biochem J 374: 1-20

Hexum J, CM B, AM K, JR O, DA L, DA H (2015) Parthenolide prodrug LC-1 slows growth of intracranial glioma. Bioorganic & medicinal chemistry letters 25:   Doi 10.1016/j.bmcl.2015.04.058

Kath C, Goni-Oliver P, Müller R, Schultz C, Haucke V, Eickholt B, Schmoranzer J (2018) PTEN suppresses axon outgrowth by down-regulating the level of detyrosinated microtubules. PLoS One 13: e0193257 Doi 10.1371/journal.pone.0193257

Kiehn O (2016) Decoding the organization of spinal circuits that control locomotion. Nat Rev Neurosci 17: 224-238 Doi 10.1038/nrn.2016.9

Kim JE, Liu BP, Park JH, Strittmatter SM (2004) Nogo-66 receptor prevents raphespinal and rubrospinal axon regeneration and limits functional recovery from spinal cord injury. Neuron 44: 439-451 Doi 10.1016/j.neuron.2004.10.015

Lang BT, Cregg JM, DePaul MA, Tran AP, Xu K, Dyck SM, Madalena KM, Brown BP, Weng YL, Li Set al (2015) Modulation of the proteoglycan receptor PTPσ promotes recovery after spinal cord injury. Nature 518: 404-408 Doi 10.1038/nature13974

Lee JK, Geoffroy CG, Chan AF, Tolentino KE, Crawford MJ, Leal MA, Kang B, Zheng B (2010) Assessing spinal axon regeneration and sprouting in Nogo-, MAG-, and OMgp-deficient mice. Neuron 66: 663-670 Doi 10.1016/j.neuron.2010.05.002

Leibinger M, Andreadaki A, Gobrecht P, Levin E, Diekmann H, Fischer D (2016) Boosting Central Nervous System Axon Regeneration by Circumventing Limitations of Natural Cytokine Signaling. Mol Ther 24: 1712-1725 Doi 10.1038/mt.2016.102

Leibinger M, Andreadaki A, Golla R, Levin E, Hilla AM, Diekmann H, Fischer D (2017) Boosting CNS axon regeneration by harnessing antagonistic effects of GSK3 activity. Proc Natl Acad Sci U S A 114: E5454-E5463 Doi 10.1073/pnas.1621225114

Leibinger M, Hilla AM, Andreadaki A, Fischer D (2019) GSK3-CRMP2 signaling mediates axonal regeneration induced by Pten knockout. Communications Biology 2: 1-13 Doi doi:10.1038/s42003-019-0524-1

Leibinger M, Muller A, Andreadaki A, Hauk TG, Kirsch M, Fischer D (2009) Neuroprotective and axon growth-promoting effects following inflammatory stimulation on mature retinal ganglion cells in mice depend on ciliary neurotrophic factor and leukemia inhibitory factor. J Neurosci 29: 14334-14341 Doi 10.1523/JNEUROSCI.2770-09.2009

Leibinger M, Muller A, Gobrecht P, Diekmann H, Andreadaki A, Fischer D (2013) Interleukin-6 contributes to CNS axon regeneration upon inflammatory stimulation. Cell Death Dis 4: e609 Doi 10.1038/cddis.2013.126

Leibinger M, Zeitler C, Gobrecht P, Andreadaki A, Gisselmann G, Fischer D (2021) Transneuronal delivery of hyper-interleukin-6 enables functional recovery after severe spinal cord injury in mice. Nat Commun 12: 391 Doi 10.1038/s41467-020-20112-4

Levin E, Diekmann H, Fischer D (2016) Highly efficient transduction of primary adult CNS and PNS neurons. Sci Rep 6: 38928 Doi 10.1038/srep38928

Lewandowski G, Steward O (2014) AAVshRNA-mediated suppression of PTEN in adult rats in combination with salmon fibrin administration enables regenerative growth of corticospinal axons and enhances recovery of voluntary motor function after cervical spinal cord injury. J Neurosci 34: 9951-9962 Doi 10.1523/JNEUROSCI.1996-14.2014

Liu K, Lu Y, Lee JK, Samara R, Willenberg R, Sears-Kraxberger I, Tedeschi A, Park KK, Jin D, Cai Bet al (2010) PTEN deletion enhances the regenerative ability of adult corticospinal neurons. Nat Neurosci 13: 1075-1081 Doi 10.1038/nn.2603

Liz MA, Mar FM, Santos TE, Pimentel HI, Marques AM, Morgado MM, Vieira S, Sousa VF, Pemble H, Wittmann Tet al (2014) Neuronal deletion of GSK3beta increases microtubule speed in the growth cone and enhances axon regeneration via CRMP-2 and independently of MAP1B and CLASP2. BMC Biol 12: 47 Doi 10.1186/1741-7007-12-47

Lu Y, Belin S, He Z (2014) Signaling regulations of neuronal regenerative ability. Curr Opin Neurobiol 27: 135-142 Doi 10.1016/j.conb.2014.03.007

Lucas FR, Goold RG, Gordon-Weeks PR, Salinas PC (1998) Inhibition of GSK-3beta leading to the loss of phosphorylated MAP-1B is an early event in axonal remodelling induced by WNT-7a or lithium. J Cell Sci 111 (Pt 10): 1351-1361

Moon LD, Asher RA, Rhodes KE, Fawcett JW (2001) Regeneration of CNS axons back to their target following treatment of adult rat brain with chondroitinase ABC. Nat Neurosci 4: 465-466 Doi 10.1038/87415

Müller A, Hauk TG, Fischer D (2007) Astrocyte-derived CNTF switches mature RGCs to a regenerative state following inflammatory stimulation. Brain 130: 3308-3320

Muller A, Hauk TG, Leibinger M, Marienfeld R, Fischer D (2009) Exogenous CNTF stimulates axon regeneration of retinal ganglion cells partially via endogenous CNTF. Mol Cell Neurosci 41: 233-246 Doi 10.1016/j.mcn.2009.03.002

Neelakantan S, Nasim S, Guzman ML, Jordan CT, Crooks PA (2009) Aminoparthenolides as novel anti-leukemic agents: Discovery of the NF-kappaB inhibitor, DMAPT (LC-1). Bioorg Med Chem Lett 19: 4346-4349 Doi 10.1016/j.bmcl.2009.05.092

Nieuwenhuis J, Adamopoulos A, Bleijerveld OB, Mazouzi A, Stickel E, Celie P, Altelaar M, Knipscheer P, Perrakis A, Blomen VAet al (2017) Vasohibins encode tubulin detyrosinating activity. Doi 10.1126/science.aao5676

Owen R, Gordon-Weeks PR (2003) Inhibition of glycogen synthase kinase 3beta in sensory neurons in culture alters filopodia dynamics and microtubule distribution in growth cones. Mol Cell Neurosci 23: 626-637

Park KK, Liu K, Hu Y, Smith PD, Wang C, Cai B, Xu B, Connolly L, Kramvis I, Sahin Met al (2008) Promoting axon regeneration in the adult CNS by modulation of the PTEN/mTOR pathway. Science 322: 963-966 Doi 10.1126/science.1161566

Silver J, Miller JH (2004) Regeneration beyond the glial scar. Nat Rev Neurosci 5: 146-156 Doi 10.1038/nrn1326

Skoumal R, Toth M, Serpi R, Rysa J, Leskinen H, Ulvila J, Saiho T, Aro J, Ruskoaho H, Szokodi Iet al (2011) Parthenolide inhibits STAT3 signaling and attenuates angiotensin II-induced left ventricular hypertrophy via modulation of fibroblast activity. J Mol Cell Cardiol 50: 634-641 Doi 10.1016/j.yjmcc.2011.01.001

Sobota R, Szwed M, Kasza A, Bugno M, Kordula T (2000) Parthenolide inhibits activation of signal transducers and activators of transcription (STATs) induced by cytokines of the IL-6 family. Biochem Biophys Res Commun 267: 329-333 Doi 10.1006/bbrc.1999.1948

Song JM, Qian X, Upadhyayya P, Hong KH, Kassie F (2014) Dimethylaminoparthenolide, a water soluble parthenolide, suppresses lung tumorigenesis through down-regulating the STAT3 signaling pathway. Curr Cancer Drug Targets 14: 59-69

Sutherland C, Cohen P (1994) The alpha-isoform of glycogen synthase kinase-3 from rabbit skeletal muscle is inactivated by p70 S6 kinase or MAP kinase-activated protein kinase-1 in vitro. FEBS letters 338: Doi 10.1016/0014-5793(94)80112-6

Sutherland C, Leighton IA, Cohen P (1993) Inactivation of glycogen synthase kinase-3 beta by phosphorylation: new kinase connections in insulin and growth-factor signalling. Biochem J 296 (Pt 1): 15-19

Terheyden-Keighley D, Leibinger M, Fischer D (2022) Transneuronal delivery of designer cytokines: perspectives for spinal cord injury. Neural Regeneration Research 17: 338-340 Doi 10.4103/1673-5374.317974

Terheyden-Keighley D, Leibinger M, Zeitler C, Fischer D (2023) Transneuronal Delivery of Cytokines to Stimulate Mammalian Spinal Cord Regeneration. Methods Mol Biol 2636: 85-99 Doi 10.1007/978-1-0716-3012-9_6

van den Brand R, Heutschi J, Barraud Q, DiGiovanna J, Bartholdi K, Huerlimann M, Friedli L, Vollenweider I, Moraud EM, Duis S et al (2012) Restoring voluntary control of locomotion after paralyzing spinal cord injury. Science 336: 1182-1185 Doi 10.1126/science.1217416

Yin Y, Cui Q, Li Y, Irwin N, Fischer D, Harvey AR, Benowitz LI (2003) Macrophage-derived factors stimulate optic nerve regeneration. J Neurosci 23: 2284-2293

Zheng B, Tuszynski MH (2023) Regulation of axonal regeneration after mammalian spinal cord injury. Nat Rev Mol Cell Biol: Doi 10.1038/s41580-022-00562-y

Zolotukhin S, Byrne BJ, Mason E, Zolotukhin I, Potter M, Chesnut K, Summerford C, Samulski RJ, Muzyczka N (1999) Recombinant adeno-associated virus purification using novel methods improves infectious titer and yield. Gene Ther 6: 973-985 Doi 10.1038/sj.gt.3300938

Zukor K, Belin S, Wang C, Keelan N, Wang X, He Z (2013) Short hairpin RNA against PTEN enhances regenerative growth of corticospinal tract axons after spinal cord injury. J Neurosci 33: 15350-15361 Doi 10.1523/jneurosci.2510-13.2013