Dynamic chromatin architecture identifies new autoimmune-associated

enhancers for *IL2* and novel genes regulating CD4+ T cell activation

Matthew C. Pahl*[1,2], Prabhat Sharma*[1,3], Rajan M. Thomas*[1,3], Zachary Thompson[1,3], Zachary

Mount[1,3], James Pippin[1,2], Peter A. Morawski[4], Peng Sun[1,3], Chun Su[1,2], Daniel J. Campbell[4,5],

Struan F.A. Grant[1,2,6,7,8], Andrew D. Wells[1,3,9,10]

[1]Center for Spatial and Functional Genomics, The Children's Hospital of Philadelphia

[2]Division of Human Genetics, The Children's Hospital of Philadelphia

[3]Department of Pathology, The Children's Hospital of Philadelphia

[4]Benaroya Research Institute at Virginia Mason, Seattle, WA

[5]Department of Immunology, University of Washington School of Medicine, Seattle, WA

[6]Department of Genetics, Perelman School of Medicine, University of Pennsylvania

[7]Department of Pediatrics, Perelman School of Medicine, University of Pennsylvania

[8]Division of Endocrinology and Diabetes, The Children's Hospital of Philadelphia

[9]Department of Pathology and Laboratory Medicine, Perelman School of Medicine, University of

Pennsylvania

[10]Institute for Immunology, Perelman School of Medicine, University of Pennsylvania

*Co-lead authors

Correspondence:

Andrew D. Wells, PhD

The Children's Hospital of Philadelphia Research Institute

916D Abramson Research Center

3615 Civic Center Boulevard

Philadelphia, PA 19104

Email: wellsa@chop.edu

Phone: 215-590-8710

ABSTRACT

Genome-wide association studies (GWAS) have identified hundreds of genetic signals associated with autoimmune disease. The majority of these signals are located in non-coding regions and likely impact *cis*-regulatory elements (cRE). Because cRE function is dynamic across cell types and states, profiling the epigenetic status of cRE across physiological processes is necessary to characterize the molecular mechanisms by which autoimmune variants contribute to disease risk. We localized risk variants from 15 autoimmune GWAS to cRE active during TCR-CD28 costimulation of naïve human CD4+ T cells. To characterize how dynamic changes in gene expression correlate with cRE activity, we measured transcript levels, chromatin accessibility, and promoter-cRE contacts across three phases of naive CD4+ T cell activation using RNA-seq, ATAC-seq, and HiC. We identified ~1,200 protein-coding genes physically connected to accessible disease-associated variants at 423 GWAS signals, at least one-third of which are dynamically regulated by activation. From these maps, we functionally validated a novel stretch of evolutionarily conserved intergenic enhancers whose activity is required for activation-induced *IL2* gene expression in human and mouse, and is influenced by autoimmune-associated genetic variation. The set of genes implicated by this approach are enriched for genes controlling CD4+ T cell function and genes involved in human inborn errors of immunity, and we pharmacologically validated eight implicated genes as novel regulators of T cell activation. These studies directly show how autoimmune variants and the genes they regulate influence processes involved in CD4+ T cell proliferation and activation.

INTRODUCTION

GWAS has linked hundreds of regions of the human genome to autoimmune disease susceptibility. The majority of GWAS variants are located in non-coding regions of the genome, and likely contribute to disease risk by modulating cis-regulatory element (cRE) activity to influence gene expression[1]. Identifying causal variants and effector genes molecularly responsible for increased disease risk is critical for identifying targets for downstream molecular study and therapeutic intervention. An understanding of how non-coding variants function is often limited by incomplete knowledge of the mechanism of action, *i.e.*, whether a variant is located in a cRE, in which cell types a cRE is active, and which genes are regulated by which cRE.

CD4+ T cells are key regulators of innate and adaptive immune responses that combat infection by orchestrating the activity of other immune cells. In their quiescent, naïve state, 'helper' T cells traffic between the blood and secondary lymphoid tissues as part of an immunosurveillance program maintained by transcription factors such as KLF2, TOB, and FOXO[2–4]. Upon encountering specific antigen, a cascade of signals activated through the TCR and costimulatory receptors results in large-scale changes in gene expression driven by factors like NFKB, NFAT, IRF4, and STATs, leading to proliferation and differentiation into specialized Th1, Th2, and Th17 subsets capable of participating in protective immunity[5,6]. CD4+ T cells are also key players in the induction and pathogenesis of autoimmunity. The cis-regulatory architecture of CD4+ T cells is enriched for autoimmune disease GWAS variants[7–14], and T cells from autoimmune patients harbor distinct epigenetic and transcriptomic signatures linking dysregulated gene expression with disease pathogenesis. Autoimmune variants may contribute to the breakdown of immune self-tolerance by shifting the balance between autoreactive conventional vs. regulatory CD4+ T cells, by altering cytokine production, or by promoting auto-antigen production[15,16].

Chromatin conformation assays allow for identification of putative target genes of

autoimmune variant-containing cRE in close spatial proximity to gene promoters. Previous work

using promoter-capture HiC and HiC in combination with other epigenetic marks have

implicated sets of autoimmune variants and effector genes that may participate in T cell

activation[17–19]. In addition to chromatin conformation approaches, multiple complementary

approaches have been developed to link disease associated SNPs to their downstream effector

genes. Expression quantitative trait mapping (eQTL) and chromatin co-accessibility data can be

used to implicate effector genes through statistical association of genotypes with readouts of

expression or other molecular markers[20]. However, these approaches do not account for the

relevant 3D structure of the genome in the nucleus, and are highly susceptible to *trans*-effects

and other confounding factors.

In this study, we measured the impact of TCR-CD28 activation on the autoimmune-

associated *cis*-regulatory architecture of CD4+ helper T cells, and by comparing our data to

those of several orthogonal GWAS effector gene nomination studies, identify hundreds of

effector genes not implicated previously. We find that 3D chromatin-based approaches exhibit

2- to 10-fold higher predictive sensitivity than eQTL and ABC statistical approaches when

benchmarked against a 'gold standard set' of genes underlying inborn errors in immunity and

tolerance. Our maps of autoimmune SNP-gene contacts also predicted a stretch of

evolutionarily conserved, intergenic enhancers that we show are required for normal expression

of the canonical T cell activation gene *IL2* in both human and mouse, whose activity is

influenced by autoimmune risk variants. The set of variant-connected effector genes defined by

3D physical proximity to autoimmune-associated cRE is enriched for genes that regulate T cell

activation, as validated pharmacologically in this study and by CRISPR-based screens in

orthogonal studies[21–23].

RESULTS

Gene expression dynamics as a function of naïve CD4+ T cell activation

To explore the universe of genes and cREs that are affected by and may contribute to CD4+

activation, we characterized the dynamics of gene expression, chromatin accessibility, and 3D

chromatin conformation in human CD4+CD62L^{hi}CD44^{lo} naïve T cells purified directly *ex vivo*

and in response to *in vitro* activation through the T cell receptor (TCR) and CD28 for 8 or 24

hours using RNA-seq (N=3 donors), ATAC-seq (N=3 donors), and HiC (N=2 donors, **Figure**

1A). As sequencing depth and restriction enzyme cutting frequency affects the resolution and

reliability of HiC experiments[24], we constructed libraries with two four-cutter restriction enzymes

and sequenced to a total of approximately 4 billion unique-valid reads per timepoint. We verified

the reproducibility of replicate ATAC-seq and RNA-seq libraries with principal component

analysis and HiC libraries using distance-controlled, stratum-adjusted correlation coefficient

(SCC). Replicate samples were highly correlated and clustered by activation stage (**Figure 1 -**

Figure Supplement 1A-C). As expected, quiescent naïve CD4+ T cells expressed high levels

of *SELL, TCF7, CCR7,* and *IL7R*, and rapidly upregulated *CD69, CD44, HLADR,* and *IL2RA*

upon stimulation (**Figure 1B, Supplementary File 1**). Genome-scale gene set variance

analysis based on MsigDB hallmark pathways showed that genes involved in KRAS and

Hedgehog signaling are actively enriched in quiescent naive CD4+ T cells, while stimulated cell

transcriptomes are enriched for genes involved in cell cycle, metabolism, and TNF-, IL-

2/STAT5-, IFNg-, Notch-, and MTORC1-mediated signaling pathways (**Supplementary File 2**).

To define global gene expression dynamics during the course of CD4+ T cell activation, we

performed pairwise comparisons and k-means clustering, identifying 4390 differentially

expressed genes after 8 hours of stimulation (3289 upregulated and 1101 downregulated) and

3611 differentially expressed genes between 8 hours and 24 hours (3015 upregulated and 596

downregulated, **Figure 1 - Figure Supplement 1D**) that could be further separated into five

clusters based on their distinct trajectories (**Figure 1C**, **Figure 1 - Figure Supplement 1E-G**,

Supplementary File 1). Genes upregulated early upon activation (cluster 1; n=1621 genes) are

enriched for pathways involved in the unfolded protein response, cytokine signaling, and

translation (*e.g.*, *IL2, IFNG, TNF, IL3, IL8, IL2RA, ICOS, CD40LG, FASLG, MYC, FOS, JUNB,*

REL, NFKB1, STAT5A, **Supplementary File 3**). Genes downregulated late (cluster 2; n=1600

genes) are moderately enriched for receptor tyrosine kinase signaling, cytokine signaling, and

extracellular matrix organization. Genes monotonically increasing (cluster 3; n=2676 genes) are

highly enriched for pathways involved in infectious disease and RNA stability, translation and

metabolism, and moderately enriched for pathways involved in the unfolded protein response,

cellular responses to stress, regulation of apoptosis, and DNA repair (*e.g.*, *TBX21, BHLHE40,*

IL12RB2, STAT1, CCND2, CDK4, PRMT1, ICAM1, EZH2, **Supplementary File 3**). Genes

downregulated early (cluster 4; n=1628 genes) are enriched for pathways involved in inositol

phosphate biosynthesis, neutrophil degranulation, and metabolism of nucleotides (*e.g.*, *KLF2,*

IL7R, RORA, IL10RA), and genes upregulated late (cluster 5; n=2154 genes) are highly

enriched for pathways involved in cell cycle, DNA unwinding, DNA repair, chromosomal

maintenance, beta-oxidation of octanoyl-CoA, and cellular response to stress (*e.g.*, *CDK2,*

E2F1, CDK1, CCNE1, CCNA2, PCNA, WEE1, CDC6, ORC1, MCM2). These patterns are

consistent with known changes in the cellular processes that operate during T cell activation,

confirming that *in vitro* stimulation of CD4+ T cells recapitulates gene expression programs

known to be engaged during a T cell immune response.

Dynamic changes in chromosomal architecture and genome accessibility during naïve

CD4+ T cell activation

To understand the *cis*-regulatory dynamics underlying the observed activation-induced

changes in gene expression, we examined CD4+ T cell nuclear chromosome conformation and

chromatin accessibility as a function of stimulation state using HiC and ATAC-seq. The human

genome consists of ~3 meters of DNA that is incorporated into chromatin and compacted into

the ~500 cubic micron nucleus of a cell in a hierarchically ordered manner. This degree of

compaction results in only ~1% of genomic DNA being accessible to the machinery that

regulates gene transcription[25], therefore a map of open chromatin regions (OCR) in a cell

represents its potential gene regulatory landscape. Open chromatin mapping of human CD4+ T

cells at all states identified a consensus *cis*-regulatory landscape of 181,093 reproducible OCR

(**Supplementary File 4**). Of these, 14% (25,291) exhibited differential accessibility following 8

hours of stimulation (FDR<0.05). Most differentially accessible regions (DAR) became more

open (18,887), but some DAR (6,629) showed reduced accessibility (**Figure 2A**,

Supplementary File 5). The change in accessibility over the next 16 hours of stimulation

showed the opposite dynamic, with 6,629 regions exhibiting reduced accessibility, and only

4,417 DAR becoming more open (total of 11,046 DAR, **Figure 2A**, **Supplementary File 5**).

These OCR represent the set of putative cRE with dynamic activity during T cell activation.

The vast majority of putative cRE are located in intergenic or intronic regions of the genome

far from gene promoters, meaning that the specific impact of a given cRE on gene expression

cannot be properly interpreted from a one-dimensional map of genomic or epigenomic features.

To predict which cRE may regulate which genes in CD4+ T cells across different states of

activation, we created three-dimensional maps of cRE-gene proximity in the context of genome

structure. The highest order of 3D nuclear genome structure is represented by A/B

compartments, which are large chromosomal domains that self-associate into transcriptionally

active (A) *vs*. inactive (B) regions[26]. In agreement with prior studies, we find that OCR located in

active A compartments exhibit higher average accessibility than those OCR located in less

active B compartments (**Figure 2 – Figure Supplement 1A, Supplementary File 6**), a trend

observed across all cell states. Likewise, genes located in A compartments show higher

average expression than those located in B compartments (**Figure 2 – Figure Supplement

1B**). A quantitative comparison across cell states showed that 94% of the CD4+ T cell genome

remained stably compartmentalized into A (42%) and B (52%), indicating that activation does

not cause a major shift in the large-scale organization of the genome within the nucleus (**Figure**

2B).

Within each A or B compartment, the genome is further organized into topologically

associating domains (TADs)[27]. These structures are defined by the fact that genomic regions

within them have the potential to interact with each other in 3D, but have low potential to interact

with regions outside the TAD. The location of TAD boundaries can influence gene expression by

limiting the access of cRE to specific, topologically associated genes. While ~80% of TAD

boundaries remained stable across all states, 20% of TAD boundaries (8925) changed as a

function of T cell activation (**Figure 2C**, **Figure 2 – Figure Supplement 1C-D**, **Supplemental**

File 7). TAD boundary dynamics were categorized and 2198 boundaries exhibited a change in

strength, 2030 boundaries shifted position, and 4697 boundaries exhibited more complex

changes such as loss of a boundary resulting in merger of two neighboring TAD, addition of a

boundary splitting one TAD into two, or a combination of any of these changes. Genes nearby

dynamic TAD boundaries are enriched for pathways involved in RNA metabolism, cellular

response to stress, and the activity of PTEN, p53, JAK/STAT, Runx and Hedgehog

(**Supplementary File 6**). We also detected chromatin stripes, which are TAD-like structures

that consist of a genomic anchor region that is brought into contact with a larger loop domain via

an active extrusion mechanism. Chromatin stripes are contained within and/or overlap TAD

regions, and are enriched for active enhancers and super-enhancers[28,29]. We identified 1526

chromatin stripes in quiescent naïve CD4+ T cells, 1676 stripes in 8 hour stimulated cells, and

2028 stripes at 24 hours post-stimulation (**Figure 2 - Figure Supplement 2A, Supplementary**

File 8). Consistent with prior studies in other cell types, chromatin stripes were preferentially

located in A compartments, and genes and OCR within stripe regions showed increased

expression and chromatin accessibility (**Figure 2 - Figure Supplement 2B-D**).

Within active topological structures, transcriptional enhancers can regulate the expression of

distant genes by looping to physically interact with gene promoters[30,31]. To identify potential

regulatory cRE-gene interactions, we identified high confidence loop contacts across all cell

states using Fit-HiC (merged 1kb, 2kb, and 4kb resolutions, **Figure 2D**). This approach detected

933,755 loop contacts in quiescent naïve CD4+ T cells, 900,267 loop contacts in 8-hour

stimulated cells, and 551,802 contacts in 24-hour stimulated cells (2,099,627 total unique

loops). Approximately 23% of these loops involved a gene promoter at one end and an OCR at

the other, and these promoter-interacting OCR were enriched for enhancer signatures based on

flanking histone marks from CD4+ T cells in the epigenome roadmap database[32] (**Figure 2 -**

Figure Supplement 1E). T cell activation resulted in significant reorganization of the open

chromatin-promoter interactome, as 907 promoter-OCR exhibited increased contact and 1333

showed decreased contact following 8 hours of stimulation (**Figure 2E**). Continued stimulation

over the next 16 hours was associated with an increase in the contact frequency of 41

promoter-OCR pairs, while only 4 pairs exhibited decreased contact (**Figure 2E**). Activation-

induced changes in chromatin architecture and gene expression were highly correlated, as

genomic regions exhibiting increased promoter connectivity became more accessible at early

stages of stimulation, which was associated with increased gene transcription from connected

promoters (**Figure 2F, Supplementary File 9**). The accessibility of promoter-connected OCR

and the expression of their associated genes decreased globally from 8 to 24 hours of

stimulation (**Figure 2F**). We compared these loop calls to a prior chromatin capture analysis in

CD4+ T cells by Burren et al.[18] and found that roughly 40% of stable loops in both stimulated

and unstimulated cells were identical in both studies, despite differing in approach (HiC vs.

PCHiC), analysis (HiC vs. CHiCAGO), sample (naïve CD4+ vs. total CD4+), timepoint (8 vs. 4

hour), and donor individuals (**Figure 2 - Figure Supplement 1F**). As expected, unstimulated

samples were more similar than activated samples.

We next focused on the 5 sets of genes with the dynamic expression patterns defined in

Figure 1D, and identified 57,609 OCR that contact dynamic gene promoters in at least one

stage. Most dynamic genes contacted between 1 and ~35 OCR, with a median of 10 OCR per

gene, but a handful of dynamic genes were observed in contact with over 100 distinct open

chromatin regions (**Figure 2 - Figure Supplement 1G**). Similarly, most OCR were connected to

a single dynamic gene, but many contacted more than one gene (median 2 genes per OCR),

suggesting that most dynamic genes have a complex regulatory architecture. Increased gene

expression upon activation correlated with an increase in the accessibility and promoter contact

frequency of distant cRE (**Figure 2 - Figure Supplement 1H**), as exemplified by *GEM* and *IRF4*

(**Figure 2 - Figure Supplement 3A**,**B**). Conversely, the 3D regulatory architecture of genes like

KLF2 and DPEP2, which were downregulated following stimulation, exhibited decreased contact

and accessibility (**Figure 2 - Figure Supplement 3C**,**D**).

Transcription factor footprints enriched in dynamic open chromatin identify regulators of T cell activation

To explore what factors drive dynamic changes in the regulatory architecture of the CD4+ T

cell genome during activation, we conducted a quantitative footprinting analysis at 1,173,159

transcription factor (TF) motifs located in the consensus CD4+ T cell open chromatin landscape

using the average accessibility of the region surrounding each motif as a measure of regulatory

activity (**Supplementary File 10**). Activation of naïve CD4+ T cells resulted in increased

chromatin accessibility around bZIP motifs at tens of thousands of genomic regions by 8 hours

post-stimulation (**Figure 3A** and **C**), which was associated with increased expression of Fos and

Jun family members constituting the AP-1 complex, as well as the bZIP factors BATF, BACH1,

and BACH2 (**Figure 3A**). The activity of NFE2 and SMAD was increased without increased

expression (**Figure 3A** and **C**), likely due to post-translational regulation of these factors by

phosphorylation[33]. Conversely, the motifs for a number of TF exhibited significantly reduced

accessibility early after stimulation, including those for EGR2 and FOXP3 that are known to

negatively regulate T cell activation[34,35] (**Figure 3A**). By 24 hours post-activation, bZIP activity

remained largely unchanged compared to 8 hours (**Figure 3B**), but a number of factors showed

decreased activity. These include several members of the Sp family, the Myc cofactor MAZ that

also cooperates with CTCF to regulate chromatin looping[36], KLF2, which controls genes

involved in naïve CD4+ T cell quiescence and homing[2,37], NRF1, a factor implicated in age-

associated T cell hypofunction[38], and EGR2 and 3, which are known to oppose T cell activation

and promote tolerance[34,39,40] (**Figure 3B** and **C**).

To explore how transcription factor activity may operate *via* the CD4+ T cell open chromatin

landscape to regulate distinct programs of dynamic gene expression during TCR/CD28-induced

activation, we focused on TF motifs enriched among those OCR specifically contacting

promoters of dynamic genes identified by our clustering analysis (**Figure 3 – Figure**

Supplement 1A). The set of OCR contacting dynamic gene promoters were enriched for the

motifs of 89 expressed (TPM>5) transcription factors, as compared to motifs present in the total

open chromatin landscape (**Figure 3D**). The majority of this TF activity was enriched in OCR

connected to genes highly upregulated at 24 hours post-activation (clusters 3 and 5, **Figure**

3D), with the exception of CREB3, ELF1, ESRRA, GABPA, RELA, XPB1, ZNF384, and the

transcriptional repressor IKZF1 known as a strong negative regulator of T cell activation and

differentiation[41–44]. Conversely, motifs for IKZF1, ZNF384, GABPA, ESRRA, and ELF1 were

highly enriched in the set of OCR contacting genes down-regulated early after activation (cluster

4, **Figure 3D**). Motifs for KLF2 and the metabolic gene regulator SREBF1 were likewise

enriched in OCR connected to down-regulated genes. Open chromatin regions interacting with

genes in cluster 2 are negatively enriched for this set of TF except for CTCF (**Figure 3D**).

Finally, we integrated TF footprint, promoter connectome, and gene co-expression data to

construct TF-gene regulatory networks likely operating at each timepoint. The connections

between regulatory nodes are based on physical promoter-TF footprint interactions with

confidence weighted by gene co-expression (**Figure 3E**, **Supplementary File 11**). Highly co-

expressed genes at the core of the unstimulated CD4+ T cell regulatory network encode

transcription factors such as KLF2, ETS1, IKZF1, and TCF7[45–47] that are known to be involved

in T cell gene silencing, quiescence, homeostasis, and homing. Genes connected to the top

factors in this network were enriched for pathways involved in immune signaling, DNA

replication and repair, protein secretion, and programmed cell death (**Figure 3 – Figure**

Supplement 1B). Costimulation through the TCR and CD28 induced a set of core network

genes active at both time points with known roles in T cell activation and differentiation (*NFKB1,*

JUNB, MYC, IRF4, STAT5, STAT1, LEF1, ATF4). Also part of this set is *PLAGL2*, an oncogene

in the Wnt pathway that regulates hypoxia-induced genes[48] with no prior defined role in T cell

activation. Additional nodes specifically implicated at 8 hours post-activation are HIF1A, the

major sensor of cellular hypoxia[49], and XBP1, a major transcriptional mediator of the unfolded

ER protein response with defined roles in T cell activation, differentiation, and exhaustion[50,51].

Factors specifically implicated at 24 hours post-activation include E2F1, a transcriptional

regulator of both cell cycle and apoptosis in T cells[52,53], BHLHE40, a factor known to control

multiple aspects of T cell metabolism and differentiation[54], and the Myc cofactor MAZ that has

not been previously studied in the context of T cell function. Genes connected to factors in the

activated T cell networks were enriched for pathways involved in cytokine signaling, the

interferon response, transcription, cell cycle, DNA replication and repair, and programmed cell

death (**Figure 3 – Figure Supplement 1B**). Together, these data indicate that concurrent but

separable stimulation-induced gene programs are the result of the activity of distinct sets of

DNA binding factors mobilized by antigen and costimulatory receptor signaling in naïve CD4+ T

cells.

Identification of autoimmune variants associated with CD4+ T cell cRE and predicted

effector genes

Following our established variant-to-gene (V2G) mapping approach to implicate functional

SNPs and their effector genes using the combination of GWAS and chromatin conformation

capture data[55–59], we intersected promoter-interacting OCR with autoimmune SNPs from the

95% credible set derived from 15 autoimmune diseases (**Figure 4A**, **Supplementary Files 12**

and **13**). Constraining the GWAS SNPs in this way reduced the credible set size from an

average of 14 variants per sentinel to 3 variants per sentinel. To determine whether open

chromatin in physical contact with dynamically regulated genes in CD4+ T cells is enriched for

autoimmune disease heritability, we performed a partitioned LD score regression analysis. This

landscape was enriched for variants associated with susceptibility to inflammatory bowel

disease (IBD), ulcerative colitis (UC), type I diabetes (T1D), lupus (SLE), celiac disease (CEL),

allergy (ALG), eczema (ECZ), and rheumatoid arthritis (RA), but not for variants associated with

psoriasis (PSO) or juvenile idiopathic arthritis (JIA) (**Figure 4B**, **Supplementary File 14**). The

OCR connected to genes upregulated early and/or progressively upon activation (clusters 1 and

3) were most strongly enriched for ALG, CEL, IBD/UC, RA and T1D heritability, while SLE and

ECZ heritability was most enriched in OCR connected to genes upregulated later post-activation

(clusters 3 and 5, **Figure 4B**). SLE was also the only disease (besides PSO and JIA) that was

not enriched in open chromatin connected to down-regulated genes (cluster 4, **Figure 4B**).

The promoter-connected open chromatin landscape for all CD4+ T cell states in this study

contains 2606 putatively causal variants linked to ~half of the sentinel signals (423) for the 15

diseases analyzed, and are in contact with a total of 1836 genes (**Supplementary File 13**). A

total of 1151 autoimmune variants localized to the promoters of 400 genes (-1500/+500bp from

the TSS, **Supplementary File 15**). These variants were on average 103 kb from the TSS of

their connected gene (**Figure 4C**), and each variant contacted an average of 5 genes (**Figure**

4D). The majority of linked SNPs interact with genes in addition to the nearest gene, and ~half

of linked SNPs 'skip' the nearest gene to target only distant genes (**Figure 4E**). Approximately

60% of connected genes were implicated across all timepoints (**Figure 4F**, **Supplementary File**

13), while ~40% (753) are dynamically regulated (clusters 1-5) in response to TCR/CD28 co-

stimulation. Examples of SNP-genes pairs that exhibit dynamic accessibility, chromosome

contact, and expression in response to T cell activation are the *SIK1*, *PARK7*, *DUSP5*,

CLEC2D, *TRIP10*, *GPR108* and *IL2* loci (**Figure 4G**, **Supplementary File 13**). The *TRIP10* and

GPR108 promoters were each captured in contact with a high confidence variant rs1077667

(PP>0.99), which is located in an intron of *TNFSF14* and is associated with multiple sclerosis

(**Figure 4H**). The accessibility of this SNP and it's contact with *TRIP10* and *GPR108* increased

following activation (**Figure 4H**). Conversely, the allergy-associated SNP rs7380290 is

accessible and contacts the *SIK1* promoter in resting cells, but shows reduced accessibility and

promoter connectivity upon activation (**Figure 4I**). *TRIP10* encodes a cytoskeletal binding

protein involved in endocytosis that suppresses glucose uptake in response to insulin

signaling[60], *GPR108* encodes an orphan G-protein coupled receptor, and *SIK1* encodes a salt-

inducible kinase with roles in cancer, epilepsy, and myeloid signaling[61]. None of these genes

have been previously studied in the context of T cell activation.

Comparative predictive power against orthogonal variant-to-gene mapping approaches

In an attempt to gauge the predictive power of our approach relative to other V2G

approaches, we compared our chromatin capture-based autoimmune GWAS effector gene

predictions to the predictions of four other chromosome capture-based studies in human CD4+

T cells[17–19,62], four single-cell eQTL studies in human CD4+ T cells[13,62–64], and a set of 449

genes that when mutated in humans cause inborn errors of immunity[65]. As part of the

comparison, we harmonized the chromatin-based datasets using our chromatin loop calling

approach and integrated with the same 95% credible set of autoimmune variants to create

comparable physical contact maps. A total of 6936 unique genes were implicated by all studies.

Our HiC-ATAC-seq approach in naïve and activated human CD4+ implicated 1947 effector

genes, 400 of which through proxies in open promoters and 1836 through contacts with distal

variants, of which 752 were dynamically regulated throughout activation. The HiC dataset from

Gate et al. implicated 110 genes, and the capture-HiC datasets from Burren et al., Yang et al.,

and Javierre et al. implicated 1408, 2572, and 2368 genes, respectively (**Supplementary File**

16). A total of 369 autoimmune GWAS effector genes were predicted in common by our HiC

approach and the pcHiC-based approaches. The eQTL studies by Soskic et al., Gate et al., Ye

et al., and Schmiedel et al. (DICE) implicated 171, 15, 15, and 221 genes, and the co-

accessibility data from Gate et al. implicated 45 genes. Concordance between our gene

predictions and the pcHiC predictions ranged from 15-24% (**Figure 5A**), while concordance

between our predictions and the eQTL predictions was low (0.6%~5%, **Figure 5A**), and

concordance among eQTL studies was also low (1.6%-13%, **Figure 5A**). However, when we

co-localized the Soskic variant-gene pairs with autoimmune GWAS SNPs, 41 of these eQTL

were also predicted by our study, representing a concordance 10-fold higher than obtained by a

random sampling of genes within 500 kb of autoimmune variants (**Figure 5B**). A total of 1519 of

the genes implicated in our study were not predicted by eQTL approaches, while 644 genes

from our CD4 T cell V2G data were not implicated by the other chromatin-based datasets, and

altogether 562 putative autoimmune effector genes were uniquely predicted by our study.

Potential sources of variation between the results of the studies are outlined in the Discussion.

A total of 17 genes were implicated by all CD4+ T cell-based approaches that nominated at

least 150 genes (APIP, BCL2L11, ERAP1, ERAP2, CD5, FIGNL1, IL18R1, METTL21B,

MRPL51, PPIF, PTGER4, PXK, RMI2, RNASET2, SERPINB1, TAPBPL, VAMP1).

To measure the predictive power of the sets of genes implicated by each approach above, a

precision-recall analysis was performed against the 'gold standard' human inborn errors in

immunity gene set, based on the hypothesis that genes under the control of autoimmune-

associated variants would cause monogenic immune disease phenotypes if subjected to

deleterious coding mutations. The set of genes from our study with disease-associated SNPs

located in their open promoters overlapped with only 5% of the genes from the gold standard

set (**Supplementary File 16**, **Figure 5C**). However, leveraging our HiC data to include distal

autoimmune variants captured interacting with open gene promoters increased the sensitivity of

the approach 3-fold to 16% (71 overlapping genes, **Figure 5C**), confirming the relevance of

long-range promoter-cRE/SNP contacts. This level of sensitivity is comparable to that of other

chromosome capture-based datasets for predicting HIEI genes in CD4+ T cells (**Figure 5C**).

Conversely, the eQTL-based approaches were 3- to ~12-fold less sensitive than the chromatin-

constrained approaches in predicting HIEI genes, with most recalling 0-2.5% and the DICE

eQTL dataset recalling 6% (**Figure 5C**). Restricting our 3D chromatin V2G to only those genes

dynamically regulated during T cell activation reduced predictive power from 16% to 10%, but

the fraction of these genes in the HIEI gold standard set (precision) increased from ~4% to 6.2%

(**Figure 5C**), indicating that focusing the V2G set on those SNP-gene pairs that are dynamically

regulated by T cell activation reduces potential false positive predictions. The precision of our

dataset was superior to that of the other chromosome capture-based datasets except for the

Burren pcHiC dataset (**Figure 5C**). The eQTL approaches were comparable to the 3D

chromatin-based datasets in precision (~3-6%, **Figure 5C**).

We also analyzed the relative predictive power of a recently proposed 'activity-by-contact'

(ABC) model[66] that uses a multi-tissue averaged HiC dataset instead of cell type-matched 3D

chromatin-based data to link variants to genes. This model constrains the input data by

removing bidirectional, antisense, and small RNAs, and eliminate genes ubiquitously expressed

across tissues based on the assumption that these genes are driven primarily by elements

adjacent to their promoters. Also, while most human genes have multiple alternative

transcriptional start sites (median = 3), ABC only annotates only one promoter per gene. To

apply the activity-by-contact gene nomination model to our CD4+ T cell chromatin-based V2G

data, we used our ATAC-seq data and publicly available CD4+ T cell H3K27ac ChIP-seq data

as input, and integrated this with GWAS and the averaged HiC dataset from the original ABC

study[66,67]. The ABC model nominated 650 genes compared to 1836 genes using our cell type-

matched HiC data and analysis pipeline. Only 357 genes were nominated by both approaches,

while 1479 genes nominated by our approach were not implicated by the ABC model, and 293

genes not implicated by our approach were newly implicated by ABC. To measure the predictive

power of the ABC approach, we re-ran the precision-recall re-analysis with all datasets

subjected to the ABC gene-promoter filter (**Fig. 5D**). We found that applying the restricted ABC

model promoter annotation to all datasets did not have a large effect on recall, however, the

precision of several of the datasets were affected. For example, using the restricted

promoter/gene set reduced the precision of our V2G approach and artificially inflated the

precision of the 'nearest gene to SNP' metric. The precision-recall analysis also shows that the

ABC score-based approach is only half as powerful at predicting HIEI genes as the chromatin-

based V2G approaches (**Fig. 5D**). This indicates that informing GWAS data with cell type- and

state-specific 3D chromatin-based data brings more target gene predictive power than

application of the multi-tissue-averaged HiC used by the ABC model. Together, these analyses

indicate that chromosome capture-based V2G is a more sensitive approach for identifying 'true'

effector genes than eQTL or ABC approaches, but comes with additional predicted genes that

represent either false positives or true effector genes for which monogenic LOF/GOF mutations

have not yet been characterized in humans.

Functional validation of novel autoimmune V2G-predicted enhancers at the *IL2* locus

Our 3D chromatin-based analysis specifically predicts dynamic, disease-associated

regulatory elements in intergenic space at the *IL2* locus. The *IL2* gene encodes a cytokine with

crucial, pleotropic roles in the immune system, and dysregulation of IL-2 and IL-2 receptor

signaling leads to immunodeficiency and autoimmune disorders in mice and humans[68–71].

Activation-induced transcription of *IL2* involves an upstream regulatory region (URR) ~375 bp

from the TSS that has served as a paradigm of tissue-specific, inducible gene transcription for

nearly four decades[72–74]. However, the presence of evolutionarily conserved non-coding

sequences (CNS) in the ~150 kb of intergenic space 46, 51, 80, 83, 85, 96, 122, and 128 kb

upstream of the TSS suggest that additional regulatory elements may have evolved to control

IL2[75] (**Figure 6A**). The -51 kb CNS contains a SNP linked to T1D, IBD, PSO, CEL and allergy

(rs72669153), while the -85 kb CNS contains a SNP linked to RA (rs6232750) and the -128 kb

CNS contains two SNPs linked to T1D, JIA, and SLE (rs1512973 and rs12504008, **Figure 6A**).

In TCR/CD28-stimulated naïve CD4+ T cells, these CNS are remodeled to become highly

accessible (**Figure 6A**), and they loop to interact physically with the *IL2* URR (**Figure 6B**) at

both time points. ChIP-seq analyses in human T cells[76] show that the URR and all distal CNS

except -85 are occupied by TF such as Jun/Fos (AP-1), NFAT, and NFkB that are known

regulators of *IL2* (**Figure 6 – Figure Supplement 1**), and the -85, -122, and -128 CNS are

occupied by additional TF not previously thought to be direct regulators of *IL2*, such as MYC,

BCL6, and STAT5 (**Figure 6 – Figure Supplement 1**). Recombinant reporter assays in primary

activated human CD4+ T cells showed that the -46, -51, -83, and -128 CNS/OCR can enhance

transcription from the URR (**Figure 6C**). To determine whether the native elements contribute to

the expression of *IL2*, we targeted each CNS/OCR individually in primary human CD4+ T cells

or Jurkat T cells using CRISPR/CAS9 (**Figure 6D**) and measured IL-2 secretion following

TCR/CD28 stimulation (**Figure 6E**). Deletion of the -46, -51, -83, -85, -122, and -128 kb

elements in primary human CD4+ T cells each resulted in a ~50% reduction in IL-2 production,

while deletion of the -80 kb element had little effect. A very similar pattern of impact was

observed when these elements were deleted individually in Jurkat T cells (**Figure 6E**). The URR

has a stronger contribution to IL-2 production than any individual intergenic element, as deletion

of the URR almost completely abrogated activation-induced IL-2 production by both primary

CD4+ or Jurkat T cells (**Figure 6E**). To determine whether these intergenic enhancers exist in

synergistic epistasis[77] necessary for *IL2* transactivation, we generated Jurkat T cell clones in

which the stretch of all 7 elements located 46 kb to 128 kb upstream of *IL2* was deleted using

CRISPR/CAS9 (**Figure 6F**). Despite the URR and 46 kb of upstream sequence being intact in

these clones, loss of the 81.3 kb stretch of intergenic enhancers renders these cells incapable

of expressing *IL2* at both the mRNA and protein level in response to stimulation (**Figure 6G**).

These results show that the URR is not sufficient for activation-induced expression of *IL2*, and

464 that *IL2* has a previously unappreciated, complex, and autoimmune disease-associated

465 regulatory architecture that was accurately predicted by our 3D epigenomic V2G approach.

466 Importantly, we find that the distal *IL2* cRE are highly accessible in quiescent memory T cell

467 subsets (Th1, Th2, Th17, Th1-17, Th22) isolated directly *ex vivo* from human blood, whereas

468 naïve CD4+ T cells and non-IL-2-producing Treg showed little accessibility at these elements

469 (**Figure 6 – Figure Supplement 1D**). This suggests that stable remodeling of distal *IL2* cRE

470 can persist *in vivo* after TCR signals cease, and that this epigenetic imprinting contributes to the

471 immediate, activation-induced production of IL-2 exhibited by memory, but not naïve or

472 regulatory, CD4+ T cells[78–80].

474 **Distal *IL2* enhancers are evolutionarily conserved and impact *in vivo* T cell-mediated**

475 **immunity in mice**

476  The intergenic *IL2* enhancers defined above are conserved at the sequence level between

477 human and mouse (**Figure 7A**). To test whether enhancer function is likewise evolutionarily

478 conserved, we used zygotic CRISPR/CAS9 targeting to generate mice with a 583 bp deletion of

479 the ortholog of the -128 kb human enhancer CNS/OCR situated 83 kb upstream from mouse *Il2*

480 (*Il2*-83-cRE-ko, C57BL6 background). Deletion of this genomic region did not discernably affect

481 T lymphocyte development in *Il2*-83-cRE-ko mice. However, peripheral CD4$^+$ T cells from *Il2*-83-

482 cRE-ko mice produced substantially less IL-2 at both the protein and mRNA level, and exhibited

483 reduced proliferation, in response to *in vitro* stimulation (**Figure 7B**), confirming that the

484 enhancer function of the orthologous -128 and -83 kb elements is conserved between human

485 and mouse. The *in vitro* induction of Foxp3+ regulatory T cells (Treg) from conventional CD4+ T

486 cell precursors in response to antigenic stimulation in the present of TGF-beta, a differentiation

487 process highly dependent upon IL-2, was reduced 4-fold in *Il2*-83-cRE-ko mice compared to

488 wild-type mice (**Figure 7C**), but could be rescued by addition of exogenous IL-2 to the culture

489 (**Figure 7C**).

To test whether the distal -83 kb *Il2* enhancer contributes to physiologic immune responses

in vivo, we immunized wild-type and *Il2*-83-cRE-ko mice with the model antigen chicken

ovalbumin. *Il2*-83-cRE-ko animals showed a nominal increase in the differentiation of CD4+ T

cells into CXCR5+PD-1[hi] follicular helper T cells (Tfh) in the spleen compared to wild-type

animals (**Figure 7D**), a process known to be antagonized by IL-2, and generated significantly

elevated levels of ovalbumin-specific IgG antibody following immunization (**Figure 7E**). To

determine whether the 83 kb *Il2* enhancer contributes to auto-inflammatory disease

susceptibility *in vivo*, we used an adoptive transfer model of IBD in which the development of

colitis is determined by the balance between conventional and regulatory CD4+ T cell

activities[81,82]. Upon transfer into lymphocyte-deficient Rag1-ko recipients, wild-type CD4+ T cells

respond to microbiota in the gut by inducing inflammatory colitis that leads to weight loss

(**Figure 7F**, WT Tconv), while co-transfer of wild-type Treg effectively controls inflammation

(**Figure 7F**, WT Tconv + WT Treg). Transfer of *Il2*-83-cRE-ko Tconv led to significantly less

colitis (**Figure 7F**, *Il2*-83-cRE-ko Tconv), and was associated with a nominal decrease in *Il2*-83-

cRE-ko Tconv in the spleen, and a significant decrease in the accumulation of *Il2*-83-cRE-ko

Tconv in the mesenteric lymph nodes (MLN) that drain the intestines (**Figure 7F** inset). Treg

require IL-2 from Tconv for their function and homeostasis[83], and despite their reduced

numbers, *Il2*-83-cRE-ko Tconv were able to support the homeostasis and function of co-

transferred wild-type Treg (**Figure 7F**, *Il2*-83-cRE-ko Tconv + WT Treg). That the -83 kb

ortholog of the human -128 kb *IL2* enhancer contributes to physiologic immune responses and

inflammatory disease susceptibility *in vivo*.

Impact of autoimmune risk-associated genetic variation on *cis*-regulatory element

activity

The chromatin conformation approach used here employs GWAS variants as 'signposts' to

identify disease-relevant regulatory elements and connect them to their target genes, but does

not *per se* determine the effect of disease-associated genetic variation on enhancer activity or

gene expression. We experimentally measured variant effects at the *IL2* locus, where the -128

enhancer defined above contains two SNPs linked to T1D, JIA, and SLE (rs1512973 and

rs12504008). Using a recombinant reporter assay in primary activated CD4+ T cells, we

confirmed that disease-associated genetic variation influences intergenic *IL2* enhancer activity

at the -128 kb element, in that the risk allele contributes significantly less transcriptional activity

than the reference allele (**Figure 8A**).

Autoimmune variants are likely to influence disease risk by altering the activity of *cis*-

regulatory elements in T cells. At genome scale, we identified over 1000 cRE likely impacted by

autoimmune disease-associated genetic variation, in that they contain autoimmune risk SNPs

that are predicted to decrease or increase transcription factor binding affinity. Overall, 1370

autoimmune risk variants in open chromatin were predicted to influence the activity of 495 DNA

binding factors (**Supplementary File 17**), including PLAG1, PRDM1, BACH2, MYC, TBX21,

BHLHE40, LEF1, TCF7, BCL6, IRF, p53, STAT (**Figure 8 – Figure Supplement 1A**), and

hundreds of NFkB, EGR, KLF, FKH/FOX, and FOS-JUN sites (**Figure 8B**). For example, the

T1D SNP rs3024505 (PP = 0.200) connected to the promoters of *IL9* and *FAIM3* (**Figure 8 –**

Figure Supplement 1B) and the celiac SNP rs13010713 (PP = 0.154) contacting the *ITGA4*

promoter (**Figure 8 – Figure Supplement 1C**) are predicted to disrupt binding sites for *MZF1*

(**Figure 8 – Figure Supplement 1D**) and SOX4 (**Figure 8 – Figure Supplement 1E**). Similarly,

the MS SNP rs1077667 contacting the promoters of *GPR108* and *TRIP10* (**Figure 4H**) is

predicted to reduce affinity for TP53, TP63, and OCT2/POU2F2 (**Figure 8 – Figure**

Supplement 1F).

Functional validation of chromosome capture-based V2G-implicated effector genes

To determine whether genes identified *via* their physical interaction with autoimmune

variants in CD4+ T cell chromatin contact maps tend to be directly involved in T cell activation

and function, we compared the set of autoimmune genes implicated by chromatin contacts in

this study to sets of genes identified in CRISPR-based screens that control aspects of CD4+ T

cell activation like proliferation and expression of the inflammatory genes *IFNG, CTLA4, IL2,*

IL2RA, and *TNF*[21–23]. The set of all V2G-implicated genes was highly enriched for genes shown

to regulate IL-2, IL-2 receptor, CTLA-4, and proliferation (**Figure 8C, Figure 8 – Figure**

Supplement 2A, Supplementary File 18). For example, 202 genes shown to regulate IL-2

production and 166 genes shown to regulate proliferation were also implicated in our

autoimmune V2G set (**Figure 8 – Figure Supplement 2A, Supplementary File 16**). Genes

implicated by V2G in activated CD4+ T cells were moderately enriched for genes known to

control the production of IFN-G, but at the individual disease level, only genes connected to

CRO- and ATD-associated variants were enriched for IFNG regulatory genes (**Figure 8C,**

Figure 8 – Figure Supplement 2A). Genes contacting CRO, PSO, RA, SLE, T1D and VIT

variants were moderately enriched for TNF regulatory genes, but the set of all V2G genes was

not enriched for TNF genes. We also queried the orthologs of our V2G-implicated genes from

the international mouse phenotype consortium (IMPC) database and identified 97 genes that

when knocked out give an immune phenotype and 126 V2G genes that result in a

hematopoietic phenotype (**Figure 8D**, **Figure 8 – Figure Supplement 2B**, **Supplementary File**

18). This frequency of observed immune/hematopoietic phenotypes represents a significant

(adjP<0.05) ~30% enrichment over expected. This gene set was also enriched for mortality,

homeostasis/metabolism, growth/body size, skeleton, and embryonic phenotypes (**Figure 8D**,

Supplementary File 18).

An important application of this V2G approach is the identification of novel regulators of T

cell activation and their potential as drug targets, as nearly 20% of implicated genes have at

least one chemical modulator currently available (**Figure 8E**, **Supplementary File 19**). As

shown above, this approach validates genes that are well-studied regulators of T cell function,

however, a significant portion of implicated genes are not well-studied and are not currently

known to regulate T cell activation (**Figure 8F**). We observed a trend that genes expressed

more highly in immune tissues (GTEX) have in general been better investigated, and identified

several less-studied genes that could be novel targets, including the kinases GRK6, PTK6,

SIK1, and MAP3K11, the G protein-coupled receptors OXER1, GPR183, GPR18, and KISS1R,

the acetylcholine receptor CHRNB1, and the *de novo* purine pathway enzyme GART. To

determine whether these V2G-implicated genes are novel regulators of T cell activation, we

used commercially available pharmacologic modulators in dose-response assays of activation-

induced T cell proliferation. Stimulation of T cells in the presence of ligands for CHRNB1,

KISS1R and OXER1 did not significantly affect T cell proliferation, however, small molecules

targeting GRK6, PTK6, MAP3K11, GPR183, GART, and SIK1 inhibited T cell activation in the

nanomolar to micromolar range (**Figure 8G**). Together, these data show that maps of dynamic,

3D variant-gene chromatin contacts in stimulated CD4+ T cells are able to identify genes with

bona fide roles in T cell activation.

DISCUSSION

By measuring dynamic changes in chromosome folding, chromatin accessibility, and gene

expression in naïve CD4+ T cells as a function of TCR-CD28 costimulation, we identified the

putative *cis*-regulatory landscape of autoimmune disease-associated genetic variation and

physically connected these elements to their putative effector genes. This and prior

chromosome capture-based studies show that most cRE and their variants interact with, and

therefore have the potential to regulate, more than one gene (median 5 in this study), supporting

a scenario in which multiple effector genes are operative at a GWAS locus. We validated a

stretch of novel distal elements predicted by our V2G approach as *bona fide* enhancers for the

canonical immune gene *IL2*, and showed that autoimmune-associated genetic variation at one

of these elements influences it's activity. We also conducted pharmacologic targeting

experiments and compared our results with CRISPR-based studies to validate sets of *bona fide*

effector genes with a confluence of multiple orthogonal lines of evidence supporting their role in
CD4+ T cell activation.

We also observe that the vast majority (87-95%) of GWAS effector genes predicted by
chromosome capture-based approaches are not the genes nearest to the sentinel SNPs
queried in this study, while roughly one-third of eGenes predicted by eQTL approaches are the
nearest to a sentinel. This and the low concordance between 3D chromatin vs. eQTL eGene
predictions is consistent with the view that eQTL-based and GWAS-based approaches
inherently implicate different types of genes. While eQTLs cluster strongly near transcription
start sites of genes with simple regulatory structures and low enrichment for functional
annotations, GWAS variants are generally far from the TSS of genes with complex regulatory
landscapes[84]. Moreover, eGene discovery by eQTL studies using scRNA-seq approaches are
significantly limited by the low gene detection power inherent to these methods, particularly in
rare cell types.

Our systematic comparison of 3D chromatin-based target gene nomination studies in human
CD4+ T cells revealed significant variability between datasets, with the highest concordance
exhibited by the Javierre and Burren datasets (37%) and the next highest exhibited by our V2G
and the Javierre dataset (24%). Although we harmonized the comparisons as much as possible
by subjecting each dataset to the same HiC loop calling and GWAS integration steps, there are
several potential sources for the observed discrepancy between the studies. The modes of
stimulation are largely comparable, but timepoints and donors varied, and ours was the only
study that sorted naïve CD4+ T cells prior to stimulation. The higher concordance among
promoter-capture datasets compared to our HiC dataset is likely due to their lower resolution
compared to our HiC and their greater sequencing depth focused at promoters compared to
HiC. The lower resolution of HindIII-based capture-HiC results in loops called to the wrong
promoters[24], and will miss distal SNP interactions at any promoters excluded from the capture
set. While HiC is unbiased in this regard, high resolution HiC will fail to call some SNP-promoter

loops because the sequencing depth is spread across the whole genome instead of focused at

promoters. However, despite variation between studies, the results show the clear value of

tissue-matched chromatin conformation maps compared to tissue-averaged HiC for

understanding the complex genetic and epigenetic mechanisms that regulate gene expression

and for predicting autoimmune effector genes.

Why do the V2G approaches not capture a larger proportion of genes from the human

inborn errors in immunity 'truth set'? Low recall/sensitivity could result from the fact that a

substantial portion of the mutations in the full HIEI gene set result in immunodeficiency but not

autoimmune phenotypes. However, restricting the HIEI truth set to only those disease gene

mutations that result in an autoimmune phenotype (143 genes) only increased recall from 16%

to 17.5%, suggesting that this is not a major factor reducing sensitivity. Also, our method uses

GWAS signals as an input, and unlike GWAS for height, BMI, blood traits, etc., most

autoimmune GWAS signals are likely not saturated, which limits discoverability. Another reason

for reduced sensitivity is the likelihood that many of the genes in the HIEI truth set operate in

cells other than CD4+ T cells, and consistent with this, when the Javierre pcHiC V2G is

extended to all immune cell types tested, recall increases from 17% to 27%. This observation

emphasizes how limited inclusion cell types and states in a study can significantly limit the

power to detect autoimmune effector genes. Why are so few of our 3D chromatin-based V2G

genes present in the HIEI truth set? Low precision could be due to false positives in the V2G

gene set; *i.e.*, contacts between disease-associated cRE and genes that are not in fact involved

in disease susceptibility. This does not argue *per se* against the biological relevance of the cRE

or the gene, only that the linkage to disease susceptibility is not relevant. Alternatively, the

absence of chromatin-GWAS-implicated genes in the truth set could be a false negative from

the point of view of the truth set; *i.e.*, monogenic, disease-causing mutations in these genes

may exist in the human population, but have not yet been discovered. For example, in the year

2000 approximately 100 monogenic mutations causing human inborn errors of immunity were

known, but this increased by ~11 disease mutations per year until 300 human inborn errors of

immunity had been identified by 2017. The advent of next-generation exome/genome

sequencing and the COVID-19 pandemic resulted in an increase in the rate of HIEI discovery

between 2018-2021 to ~40 per year to the current recognized HIEI of ~450. If we assume a

conservative continued rate of HIEI discovery of 25 disease genes per year, the next 10 years

will show us an additional ~250 disease genes that are not currently contained in the truth set.

Many of the genes identified in this 3D epigenomic V2G screen have known roles in T cell

activation and function. An example is *IL2*, and we used the resulting maps to identify and

validate a stretch of previously unknown distal enhancers whose activity is required for *IL2*

expression and is influenced by autoimmune genetic variation. Another example is the

phosphatase DUSP5 that regulates MAPK signaling during T cell activation[85,86]. However, roles

for many of the genes implicated here in T cell activation are not known. For example, one of

the top implicated genes, PARK7, is a deglycase studied in the context of Parkinson's disease,

but has a recently a been shown to modulate regulatory T cell function[87]. The orphan G protein-

coupled receptor GPR108 is another top gene uniquely implicated by our chromatin-based V2G

that has not been studied in T cells, but was identified in a recent CRISPR screen for genes

affecting IL-2 levels[23]. Also co-implicated by our study and recent CRISPR screens are the

cannabidiol receptor GPR18[88] and the purine biosynthetic enzyme GART[89]. The GPR18 agonist

arachidonyl glycine inhibited CD4+ T cell activation above 10 uM, while the GPR18 antagonist

O-1918 slightly enhanced T cell activation. This GPR was implicated by both chromatin- and

eQTL-based approaches. Antagonism of GART, an enzyme we previous identified as a V2G

effector gene in COVID19 severity[58], with the FDA-approved drug lometrexol inhibited T cell

activation in the 10 nM range. Antagonism of GRK6, a member of the G-coupled receptor

kinase family associated with insulin secretion and T2D susceptibility[90], and PTK6, an

oncogenic kinase studied in the context of cancer[91], led to inhibition of T cell activation in the nM

to uM range. These targets were implicated by the other chromatin-based approaches but not

by eQTL. Inhibition of MAP3K11, a kinase that facilitates signaling through JNK1 and IkappaB

kinase in cancer[92,93] inhibited stimulation-induced CD4+ T cell proliferation in the 100 nM range,

as did 25-hydroxycholesterol, a ligand of the G protein-coupled receptor GPR183 that was

implicated by both chromatin- and eQTL-based approaches. SIK1 is a member of the salt-

inducible kinase family uniquely implicated in our study that negatively regulates TORC

activity[61], and a small molecule SIK1 inhibitor potently antagonized stimulation-induced CD4+ T

cell activation in the pM range.

Our integration of high-resolution Hi-C, ATAC-seq, RNA-seq and GWAS data in a single

immune cell type across multiple activation states identified hundreds of autoimmune variant-

gene pairs at ~half of all GWAS loci studied, and application of this technique to additional

immune cell types will likely identify effector genes at many of the remaining loci. This study

highlights the value of chromosome conformation data as a powerful biological constraint for

focusing variant-to-gene mapping efforts[94], and shows that dynamic changes in the spatial

conformation of the genome that accompany cell state transitions alter gene expression by

exposing promoters to a varying array of *cis*-regulatory elements, transcription factors, and

genetic variants.

MATERIALS AND METHODS

<u>T cell isolation and *in vitro* stimulation</u>

Human primary CD4+ T cells were purified from the apheresis products obtained from healthy,

screened human donors through University of Pennsylvania Human Immunology Core (HIC).

Naïve CD4+ T cells were purified using EasySepTM human naïve CD4+ T cell isolation kit II

(STEM cells Technologies, cat#17555) by immunomagnetic negative selection as per

manufacturer's protocol. Isolated untouched, highly purified (93-98%) naïve human CD4 T cells

were activated using anti-CD3+anti-CD28 Dynabeads (1:1, Thermofisher scientific, cat #

11161D) for 8-24 hours. Cells were then used to prepare sequencing libraries. The human

leukemic T cell line Jurkat was obtained from ATCC, cloned by limiting dilution, and clones with

high activation-induced secretion of IL-2 were selected for further study. Single-cell mouse

lymphocyte suspensions were prepared from spleen and lymph nodes isolated from 6-week-old

female wild-type or *Il2*-83cRE-ko mice (C57BL6 background). Mouse CD4+ T cells were purified

by negative selection using a CD4+ T cell isolation kit (Miltenyi Biotec, cat. # 130-104-454). For

CD8 depletion, CD8+ T cells were positively stained using CD8a (Ly-2) microbeads (Miltenyi

Biotec, cat. # 130-117-044) and the negative flow-through fraction was collected. Tregs were

purified from the mouse lymphocytes using a CD4+CD25+ regulatory T cell isolation kit (Miltenyi

Biotec, cat. # 130-091-041). The purity of the isolated cells was checked by flow cytometry,

showing approximately 95% cell purity. For iTreg induction, CD4+ CD25- T cells (1x 10^6) were

activated in a 24-well plate pre-coated with anti-CD3 (1 µg/mL) and the cells were cultured for

72 hours in RPMI culture medium with soluble anti-CD3 (0.5 µg/mL), IL-2 (25 units/mL), TGF-

beta (3 ng/mL), anti-IFN-gamma (5 µg/mL), and anti-IL-4 (5 µg/mL). iTreg induction cultures

were also set up without adding IL-2. iTreg cultures were harvested after 72 hours, and cells

were stained for Foxp3 expression.

<u>*Il2*-83-cRE ko Mice</u>

The CRISPR/CAS method was employed to delete the intergenic -83 CNS sequence between

Il2 and *Il21*. CRISPR guide RNAs were designed (two guide RNAs upstream of -83CNS) and

one guide RNA (downstream of -83CNS) using guide RNA design tools

(http://crispr.mit.edu.guides). The pX335 plasmid (Addgene #42335) was used to generate a

DNA template for sgRNA *in vitro* transcription. To *in vitro* transcribe the sgRNA, a T7 sequence

was incorporated at the 5' end of the guide RNA sequence, and a part of the sgRNA scaffold

sequence from px335 was added at the 3' end of the guide RNA. Oligonucleotide sequences

containing the guide RNA (underlined) along with these added sequences were synthesized by

IDT. The oligonucleotide sequences were as follows:

T7_guideRNA#1_scaffold#1:

TTAATACGACTCACTATAGGTTTTCCACGGATCTGCTCGGGTTTTAGAGCTAGAAATAGC

T7_guideRNA#1_scaffold#2:

TTAATACGACTCACTATAGGTGCTTTCTAGGTGAAGCCCCGTTTTAGAGCTAGAAATAGC

T7_guideRNA#1_scaffold#3:

TTAATACGACTCACTATAGGTCATTTGAGCCTAACTACTCGTTTTAGAGCTAGAAATAGC

Along with the above-cited sequences, a reverse primer sequence

(AGCACCGACTCGGTGCCACT) from PX335 was used to amplify a PCR-generated template

for sgRNA *in vitro* transcription. The PCR product (~ 117 bp) was verified on a 2% agarose gel

and then gel-purified using the QIAQuick gel extraction kit. The T7 sgRNA PCR product (500

ng) was used as a template for *in vitro* transcription with the T7 High Yield RNA synthesis kit

(NEB, cat # E2040S). The reaction was incubated at 37°C for 4 hours, and then the sgRNA was

purified using the MEGAclear kit (Life Technologies, cat # AM1908) according to the kit

protocol. The sgRNA was eluted with elution buffer preheated to 95°C. The eluted sgRNA was

centrifuged at 13,000 rpm for 20 minutes at 4°C, and the suspension was transferred into a new

RNase-free tube. The sgRNA quality was checked by a bioanalyzer using an RNA nano Chip.

The sgRNA was diluted to 500 ng/µl in injection buffer (sterile 10 mM Tris/0.1 mM EDTA, pH

7.5) and stored in a -80°C freezer until use. An injection mixture (final volume 30 µl) was

prepared in injection buffer by mixing 500 ng/µl of each of the sgRNAs (left and right) with Cas9

mRNA (1 µg/µl, TriLink, cat # L-6125). Fertilized eggs collected from B6/129 mice were

microinjected at the CHOP transgenic core and transferred into pseudo-pregnant B6 females.

The pups were genotyped using primers flanking the targeted sequence (Forward primer:

TTAGGACCTCACCCATCACAA and reverse primer: CATGCCCAGCTACTCTGACAT). The

PCR product was cloned into the Promega PGEM T Easy TA cloning vector, and plasmid DNA

was Sanger sequenced to determine the size of the deletion. The targeting resulted in mice with

a 500 bp and 583 bp deletion at the targeted -83 CNS site and these mutant mice showed same

phenotype. The *Il2*-83-cRE mutant heterozygous male mice were backcrossed with B6 females

for 10 successive generations.

<u>ELISA</u>

Cell culture supernatants collected at various time intervals *from in vitro* stimulated T cells of

mice or humans were analyzed for IL-2 by ELISA using kits purchased from ThermoFisher

Scientific (mouse IL-2 ELISA kit: cat # 88-7024-88) and human IL-2 ELISA kit, cat # 88-7025-

76). IL-2 ELISA was performed following the protocol provided by the vendor.

<u>*In vivo* ovalbumin immunization</u>

Eight week-old female WT and Il2-CNS-83 KO mice were injected intraperitoneally with 50 µg of

ovalbumin (Sigma cat # A5503-5G) mixed with IFA (Sigma cat # F5506). Blood was collected

from the immunized mice after 10 days, and serum was separated. The level of ovalbumin-

specific IgG in the serum was determined by ELISA using an ovalbumin-specific IgG ELISA kit

purchased from MyBiosource (cat # MBS763696). The immunized mice were sacrificed on day

10 of immunization, and spleen and lymph nodes were collected. Splenocytes and lymph node

cells were stained with CD4 BV785, CD25 BV650, CD44 Percp Cy5.5, CD62L APC eFL780,

CXCR5 BV421, PD-1 APC, Bcl-6 PE antibodies. TFH frequency was determined by flow

cytometry analysis.

<u>*In vivo* inflammatory colitis model</u>

Conventional CD4+CD25-ve T cells and CD4+CD25+ Tregs were purified from spleens and

lymph nodes of male wild-type or Il2-CNS-83 ko mice (6-8 weeks of age). T cells were

transferred into Rag1-ko male mice (5 per group) by retro-orbital injection of 1 million wild-type

or Il2-CNS-83 ko Tconv cells alone or along with (0.25×10^6) wild-type Tregs. Experimental

Rag1-ko recipients were weighed 3 times per week and scored for IBD-induced clinical

symptoms. Mice were sacrificed 72 days post-transfer, and spleen, mesenteric lymph nodes,

and colon were collected. Single-cell suspensions were prepared, cells were stained for CD4,

CD8, CD25, CD44, and Foxp3, and analyzed by flow cytometry. The absolute T cell count was

estimated using cell count and cell frequency derived from the flow cytometry analysis.

<u>RNA-seq library generation and sequencing</u>

RNA was isolated from ~ 1 million of each cell stage using Trizol Re- agent (Invitrogen), purified

using the Directzol RNA Miniprep Kit (Zymo Research), and depleted of contaminating genomic

DNA using DNAse I. Purified RNA was checked for quality on a Bioanlayzer 2100 using the

Nano RNA Chip and samples with RIN>7 were used for RNA-seq library preparation. RNA

samples were depleted of rRNA using QIAseq Fastselect RNA removal kit (Qiagen). Samples

were then processed for the preparation of libraries using the SMARTer Stranded Total RNA

Sample Prep Kit (Takara Bio USA) according to the manufacturer's instructions. Briefly, the

purified first-strand cDNA is amplified into RNA-seq libraries using SeqAmp DNA Polymerase

and the Forward and the Reverse PCR Primers from the Illumina Indexing Primer Set HT for

Illumina. Quality and quantity of the libraries was assessed using the Agilent 2100 Bioanalyzer

system and Qubit fluorometer (Life Technologies). Sequencing was performed on the NovaSeq

6000 platform at the CHOP Center for Spatial and Functional Genomics.

<u>ATAC-seq library generation and sequencing</u>

A total of 50,000 to 100,000 sorted cells were centrifuged at 550 g for 5 min at 4 °C. The cell

pellet was washed with cold PBS and resuspended in 50 µL cold lysis buffer (10 mM Tris-HCl,

pH 7.4, 10 mM NaCl, 3 mM MgCl2, 0.1% NP-40/IGEPAL CA-630) and immediately centrifuged

at 550 g for 10 min at 4 °C. Nuclei were resuspended in the Nextera transposition reaction mix

(25 µL 2x TD Buffer, 2.5 µL Nextera Tn5 transposase (Illumina Cat #FC-121-1030), and 22.5 µL

nuclease free H2O) on ice, then incubated for 45 min at 37 °C. The tagmented DNA was then

purified using the Qiagen MinElute kit eluted with 10.5 µL Elution Buffer (EB). Ten microliters of

purified tagmented DNA was PCR amplified using Nextera primers for 12 cycles to generate

each library. PCR reaction was subsequently cleaned up using 1.5x AMPureXP beads

(Agencourt), and concentrations were measured by Qubit. Libraries were paired-end sequenced

on the Illumina HiSeq 4000 platform (100 bp read length).

<u>Hi-C library preparation</u>

Hi-C library preparation on FACS-sorted CD4+ T cells was performed using the Arima-HiC kit

(Arima Genomics Inc), according to the manufacturer's protocols. Briefly, cells were crosslinked

using formaldehyde. Crosslinked cells were then subject to the Arima-HiC protocol, which

utilizes multiple restriction enzymes to digest chromatin. Arima-HiC sequencing libraries were

prepared by first shearing purified proximally-ligated DNA and then size-selecting 200-600 bp

DNA fragments using AmpureXP beads (Beckman Coulter). The size-selected fragments were

then enriched using Enrichment Beads (provided in the Arima-HiC kit), and then converted into

Illumina-compatible sequencing libraries with the Swift Accel-NGS 2SPlus DNA Library Kit

(Swift, 21024) and Swift 2S Indexing Kit (Swift, 26148). The purified, PCR-amplified DNA

underwent standard QC (qPCR, Bioanalyzer, and KAPA Library Quantification [Roche,

KK4824]) and was sequenced with unique single indexes on the Illumina NovaSeq 6000

Sequencing System using 200 bp reads.

<u>ATAC-seq data analysis</u>

ATAC-seq peaks from libraries unstimulated and stimulated Naïve CD4+ T cells were called

using the ENCODE ATAC-seq pipeline (https://www.encodeproject.org/atac-seq/). Briefly, pair-

end reads from three biological replicates for each cell type were aligned to hg19 genome using

bowtie2[95], and duplicate reads were removed from the alignment. Narrow peaks were called

independently for each replicate using macs2[96] (-p 0.01 --nomodel --shift -75 --extsize 150 -B --

SPMR --keep-dup all --call-summits) and ENCODE blacklist regions (ENCSR636HFF) were

removed from peaks in individual replicates. Reproducible peaks, peaks called in at least two

replicates, were used to generate a consensus peakset. Signal peaks were normalized using

csaw[97] in 10kb bins background regions and low abundance peaks (CPM>1) were excluded

from the analysis. Tests for differential accessibility was conducted with the glmQLFit approach

implemented in edgeR[98] using the normalization factors calculated by csaw. OCRs with FDR <

0.05 and abs($\log_2$FC > 1) between stages were considered differentially accessible.

<u>Hi-C data analysis</u>

Paired-end reads from two replicates were pre-processed using the HICUP pipeline v0.7.4[99],

with bowtie as aligner and hg19 as the reference genome. The alignment .bam file were parsed

to .pairs format using pairtools v0.3.0 (https://github.com/open2c/pairtools) and pairix v0.3.7

(https://github.com/4dn-dcic/pairix), and eventually converted to pre-binned Hi-C matrix in .cool

format by cooler v0.8.10[100] with multiple resolutions (500bp, 1kbp, 2kbp, 2.5kbp, 4kbp, 5kbp,

10kbp, 25kbp, 40kbp, 50kbp, 100kbp, 250kbp, 500kbp, 1Mbp and 2.5Mbp) and normalized with

ICE method[101]. Replicate similarity was determined by HiCRep v1.12.2[100] at 10K resolution. For

each sample, eigenvectors were determined from an ICE balanced Hi-C matrix with 40kb

resolution using cooltools v0.3.2[102] and first principal components were used to determine A/B

compartments with GC% of genome region as reference track to determine the sign. Differential

TAD comparison was performed using TADcompare with the default settings for each

chromosome (v1.4.0)[103]. Finally, for each cell type, significant intra-chromosomal interaction

loops were determined under multiple resolutions (1kb, 2kb and 4kb) using the Hi-C loop caller

Fit-Hi-C2 v2.0.7[104] (FDR<1e-6) on merged replicates matrix. The consensus chromatin loops

within resolution were identified by combining all three stages. These sets of loops were used

as consensus for quantitative differential analysis explained below. The final consensus

interaction loops for visualization were collected by merging loops from all the resolutions with

preference to keep the highest resolution. Quantitative loop differential analysis across cell

types was performed on fast lasso normalized interaction frequency (IF) implemented in

multiCompareHiC v1.8.0[105] for each chromosome at resolution 1kb, 2kb and 4kb independently.

The contacts with zero interaction frequency (IF) among more than 80% of the samples and

average IF less than 5 were excluded from differential analysis. The QLF test based on a

generalized linear model was performed in cell type-pairwise comparisons, and p-values were

corrected with FDR. The final differential loops were identified by overlapping differential IF

contacts with consensus interaction loops.

<u>Bulk RNA-seq data analysis</u>

Bulk RNA-seq libraries were sequenced on an Illumina Novaseq 6000 instrument. The pair-end

fastq files were mapped to the genome assembly hg19 by STAR (v2.6.0c) independently for

each replicate. The GencodeV19 annotation was used for gene feature annotation and the raw

read count for gene feature was calculated by htseq-count (v0.6.1)[106] with parameter settings -f

bam -r pos -s yes -t exon -m union. The gene features localized on chrM or annotated as

rRNAs, small coding RNA, or pseudo genes were removed from the final sample-by-gene read

count matrix. Gene set variation analysis was performed using GSVA 1.42.0[107] with the MSigDB

hallmark geneset[108], with resulting scores analyzed using limma (limma_3.50.3)[109]. Low

abundance peaks (CPM>1) were excluded from the analysis. Testing for differential expression

was conducted with the glmQLFit approach implemented in edgeR[98]. Genes with FDR<0.05 and

abs($\log_2$FC>1) between stages were considered differentially expressed. Differential genes

were then clustered using k-means clustering. The number of clusters was determined using

the elbow method on the weighted sum of squares, where was set to k=5. Score for how similar

each gene followed the clusters expression pattern was determined by calculating pearson

correlation coefficients between each gene in the cluster and the cluster centroid.

<u>Transcription factor footprinting and motif analysis</u>

Transcription factor footprints were called using Regulatory Analysis Toolbox HINT-ATAC

(v0.13.0) with pooled ATAC-seq data for each stage and consensus peak calls[110]. The rgt-hint

footprinting was run with parameters –atac-seq, --paired-end, and organism=hg19 set. The

output footprint coordinates were subsequently matched using rgt-motifanalysis matching with

parameters --organism hg19 and –pseudocount 0.8 set. The JASPAR2020 position weight

matrix database was used to match footprints[111]. Differential analysis of TF binding across

conditions was performed using rgt-hint differential with parameters –organism hg19, --bc, --nc

24 using the motif matched transcription factor footprints. An activity score is then calculated

based on the accessibility surrounding the footprint.

Partitioned heritability LD score regression enrichment analysis

880
Partitioned heritability LD score regression[112] (v1.0.0) was used to identify heritability

enrichment with GWAS summary statistics and open chromatin regions annotated to genes.

The baseline analysis was performed using LDSCORE data

(https://data.broadinstitute.org/alkesgroup/LDSCORE) with LD scores, regression weights, and

allele frequencies from the 1000G phase 1 data. The summary statistics were obtained from

studies as described in **Supplementary File 14** and harmonized with the munge_sumstats.py

script. Annotations for Partitioned LD score regression were generated using the coordinates of

open chromatin regions that contact gene promoters through Hi-C loops for each cell type.

Finally, partitioned LD scores were compared to baseline LD scores to measure enrichment fold

change and enrichment p-values, which were adjusted with FDR across all comparisons.

Variant-to-gene mapping using HiC-derived promoter contacts

95% credible sets were determined as previously described[113]. Briefly, P values from GWAS

summary statistics were converted to Bayes factors. Posterior probabilities were then calculated

for each variant. The variants were ordered from highest to lowest posterior probabilities added

to the credible set until the cumulative sum of posterior probabilities reached 95%. Variants in

the 95% credible set were then intersected with the CD4+ T cell promoter interacting region

OCRs from the three timepoints using the R package GenomicRanges (v1.46.1)[114].

Genomic reference and visualizations

All analyses were performed using the hg19 reference genome using gencodeV19 as the gene

reference. Genomic tracks were visualized with pyGenomeTracks v3.5[115]. HiC matrices depict

the log1p(balanced count) from the cooler count matrix. ATAC-seq tracks were generated from

bigwig files that were normalized using deeptools[116].

ABC model predictions

We used the ABC model (PMID: 31784727) using the our Tcell ATAC-seq data (unstimulated or

24 hours stimulated) generated from Naive CD4+ T cells merged acrossed replicates. For

H3K27ac we retrieved paired fastq files from H3K27ac MINT-ChIP data from ENCODE for

resting (unstimated) and activated (36hr stimulated) CD4 T cells derived from thymus. The

deduplicated bam files were used as signal files for the ABC model, the consensus peak set

were used as the input set of enhancers. The accession numbers for unstimulated cells are:

ENCFF732NOS,   ENCFF658TLX,   ENCFF605OGN,   ENCFF012XYW,   ENCFF407QZP,

ENCFF556QWM,   ENCFF888IHV,   ENCFF260FUF,   ENCFF550EVW,   ENCFF747HGZ,

ENCFF067XWQ,   ENCFF939YBH,   ENCFF749LGW,   ENCFF788LWV,   ENCFF610FRB,

ENCFF538GQX,   ENCFF355PCA,   ENCFF040OMT,   ENCFF878SHZ,   ENCFF210WNQ,

ENCFF849ZJH,   ENCFF925ARM,   ENCFF112YNU,   ENCFF808JFV,   ENCFF448KHE,

ENCFF591BFV, ENCFF366UTF, ENCFF172EWR. The accession numbers for stimulated cells

are: ENCFF442DFF,   ENCFF870ZYC,   ENCFF189UPS,   ENCFF725KSA,   ENCFF618YAV,

ENCFF765AJK,   ENCFF854QUS,   ENCFF180TKV,   ENCFF761RCK,   ENCFF449EAE,

ENCFF915VPK,   ENCFF575XFH,   ENCFF227AGX,   ENCFF810XFH,   ENCFF367OMH,

ENCFF114BBZ,   ENCFF855XPP,   ENCFF132AKN,   ENCFF663MHJ,   ENCFF244TPB,

ENCFF230RIJ,   ENCFF648FPC,   ENCFF797CIK,   ENCFF096JPW,   ENCFF704KMT,

ENCFF385POD,   ENCFF395RIG,   ENCFF043RRJ,   ENCFF027RNR,   ENCFF333RGD,

ENCFF048PMV,   ENCFF887XSA,   ENCFF188CNA,   ENCFF614IZY,   ENCFF984TTZ,

ENCFF977RDC,   ENCFF393EWO,   ENCFF781VQT,   ENCFF638UCZ,   ENCFF873QZH,

ENCFF553JPM,   ENCFF589HRL,   ENCFF721ZCN,   ENCFF496HYV,   ENCFF455WBD,

ENCFF679UTJ, ENCFF200MYN, ENCFF459ARH.

The retrieved files were trimmed/preprocessed using fastp with the default settings. Following

this, matched R1 and R2 files were concatenated to a single R1/R2 file pair. The reads were

aligned to hg19 and duplicates were removed as described for the ATAC-seq data analysis.

The deduplicated bam files were used as input to the ABC model. We ran the ABC model with

the default settings for hg19 and provided reference files for the blacklist, promoter annotation,

chromosome sizes reference, and set of ubiquitiously expressed genes. The HiC data used

was the KR normalized averaged HiC dataset that was derived from a set of cell lines

(GM12878, NHEK, HMEC, RPE1, THP1, IMR90, HU- VEC, HCT116, K562, KBM7;

average_hic.v2.191020.tar.gz) at 5kb resolution then scaled by powerlaw distribution as

described in the original manuscript. The default threshold of 0.2 was used to link enhancers to

genes. The resulting elements were intersected with the credible set list used in this study for

precision-recall analyses.

<u>Precision-recall analyses against HIEI genes</u>

To benchmark our autoimmune effector gene predictions against prior eQTL and chromatin

capture studies, we curated a list of 449 expert curated genes from Human Errors in Inborn

Immunity 2022, where mutations have been reported to cause immune phenotypes in

humans[65]. Most of these genes were identified through rare loss of function mutations in the

coding region. In addition to this we also took a subset of 145 genes reported to specifically

result in autoimmune phenotypes. These two sets of genes were treated as "gold standard" sets

of genes to compare different approaches to predict GWAS effector genes. We curated the

results of several other chromatin conformation or eQTL based methods of assigning variants to

genes to compare with our study. For the chromatin confirmation studies, we obtained loop calls

from the associated publications and intersected with the list variants in the 95% credible set

from 15 GWAS used to predict effector genes using GenomicRanges (v1.46.1)[114]. For eQTL

studies, we obtained the summary stats files and identified eGenes linked to any member of the

credible set reported as an eQTL in the transcriptome wide association analysis (TWAS; p-value

< 1e-4). Precision was calculated as the ratio of the number predicted genes in the truth set to

the total number of predicted genes, while recall was calculated as the ratio of the number

predicted genes in the truth set to the total number of genes in truth set.

<u>Colocalized eQTL comparisons</u>

In addition to comparisons with the eGenes identified by TWAS in the Soskic et al. study, we

also compared overlap of gene nominations in our study with the eQTLs that colocalized with an

autoimmune GWAS[13]. Enrichment of sentinel-gene assignments was conducted similarly as

described previously[56]. Briefly, a null distribution was constructed by randomly selecting genes

within 1 Mb of the sentinel compared to the set of colocalized cis-eQTL[13] found in through

10,000 iterations. The observed overlap reports the set of gene identified by both our HiC based

approach with the set of colocalized eQTLs. We report the empirical P value of the observed

value relative to null distribution.

<u>International mouse phenotyping consortium comparisons</u>

The set of HiC implicated genes were compared to the mouse international phenotyping

consortium set of genes with reported phenotypes[117]. We converted the list of V2G implicated

genes to mouse homologs using homologene. We tested for enrichment for each phenotype

using a one-sided proportion test implicated in R prop.test with type set to "upper".

<u>Identification of pharmacological agents</u>

We queried the Drug-Gene Interaction Database with the set of V2G implicated genes for

chemicals using rDGIdb (v1.20.0)[118]. To identify the number of papers for each gene with at

least one drug annotated to target it, we queried pubmed titles and abstracts using the R

package RISmed (v2.3.0) with each gene's name and either "autoimmune" and the list of

autoimmune diseases (**Supplementary File 17**). A score to approximate expression specificity

was computed using the sum GTEX median expression values (v8) for whole blood or spleen

divided by other tissues[119].

<u>Lentiviral-based CRISPR/CAS9 targeting in Jurkat cells</u>

LentiCRISPRv2-mCherry vectors encoding gRNA-CAS9 and the fluorescent reporter mCherry

were used for Jurkat targeting. CRISPR guide RNAs (sgRNA) targeting human IL-2-21

intergenic -46, -51, -80, -83, -85, -122, -128 CNS regions were designed using

http://crispr.tefor.net and cloned into lentiCRISPRv2-mCherry. Empty vector without gRNA

insert was used as a control. Below is the list of CRISPR gRNA for causing deletion of human

IL2-21 intergenic regions:

CNS region   CRISPR gRNA

-46 CNS   gRNA #1 AGGATGCCTACCTCCAAATG

986   gRNA # 2 AGGTGACAACATTTAGTCAG

987   -51 CNS   gRNA #1 GGCAACGAAATTCACTGTGA

988   gRNA # 2 ATTCTAACAGGAATCATTCG

989   -80 CNS   gRNA # 1 GTTCTACCTATGCCGCATTG

990   gRNA # 2 GAGATTTACTCAGTCCAATG

991   -83 CNS   gRNA # 1 GTGACAAGCATGACTCTACA

992   gRNA # 2 GTGATGGTGAATTAAGCTGA

993   -85 CNS   gRNA #1 AGGGTTTTCTAGTTACGAGA

994   gRNA # 2 ATGGTTAGTTAGCTCCCAAG

995   -96 CNS   gRNA#1   TGGGAAAAACATCTTACCTG

996   gRNA # 2 TGGCCCATGAACCATCAAAG

997   -122 CNS   gRNA# 1 GTTATTAATCTAAGCGGAGA

998   gRNA# 2 GGAAGTTAGGCAGTCAATCG

999   -128 CNS   gRNA#1   CTTCAATCATTGCATTCCAC

1000   gRNA# 2 TGACACCACCCCTGCTTGAG

HEK 293T cells were grown in RPMI1640 complete medium (RPMI1640 + 1X P/S, 1x L-Glu,

10% FBS), 37C, 7%CO2. 293T cells were transfected with 10 ug of lenti-CRISPR-V2-CRE

construct along with packaging plasmid 6 ug of PsPAX2 (Addgene, Cat #12260) and 3.5 ug of

PmD2.G (Addgene, Cat #12259) using Lipofectamine 2000 transfection reagent (Invitrogen cat

11668019). After 6 hours, the transfection medium was replaced with complete culture

medium. Transfected cells were incubated at 37C for 48-72 hours in a cell culture incubator.

and then the Lentiviral supernatants were harvested and spun at 300g for 5 minutes to remove

cellular debris. Lentiviral supernatants were concentrated using Lenti-X™ Concentrator (Takara

Bio, Cat # 631232) and then centrifuged at 1500g for 30 minutes at 4°C and supernatant was

discarded. The lentiviral pellet was resuspended at a ratio of 1:20 of the original volume using

RPMI media and concentrated virus supernatant aliquots were prepared and stored until use at

-80°C. To achieve high transduction efficiency, the viral supernatant was titrated in Jurkat cells

through transduction using various dilutions of the viral supernatants and transduction efficiency

was determined by mcherry expression analyzed through flow cytometry. Jurkat cells were

seeded in a 24 well plate at 0.5 x10⁶/well in culture media, viral supernatant with 8 ug/mL of

polybrene was added to each well. Spinfection was performed for 90 min. at 2500 rpm, and

transduced cells were equilibrated at 37C for ~6 hrs, followed by incubation at 37C 5%CO2 for

~72 hours culture. Transduced Jurkat Cells were then harvested and stimulated using PMA (15

ng/mL) + Ionomycin (1 uM) + human anti-CD28 (2 ug/ml), BioXcell cat # BE0248 in 96 well

culture plates (TPP, cat # 92097) in triplicates. Cell culture supernatants were collected at the

end of culture and analyzed for IL-2 by ELISA using a kit (cat # 88-7025-76) purchased from

Thermofisher Scientific.

<u>gRNA-CAS9 RNP-based targeting in primary human CD4+ T cells</u>

Primary Human CD4 T cells derived from 5 normal healthy donors were obtained from Human

Immunology core (University of Pennsylvania). Alt-R S.p. HiFi Cas9 Nuclease V3 cat # 1081061

CAS9 and following list of Alt-R® CRISPR-Cas9 sgRNA targeting IL-2-21 CRE were purchased

from Integrated DNA Technologies, USA.

Name	20nt gRNA Seq
46CNS-1	GACGTATATAGTCATCTGAT
46CNS-2	TCTGTGGAGCTGCTGCGTTA
46CNS-3	AGGATGCCTACCTCCAAATG
51CNS-1	ATTCTAACAGGAATCATTCG

51CNS-2	GAGTAAAAAGACGTGTTACC
51CNS-3	GGCAACGAAATTCACTGTGA
80CNS-1	AATGCGGCATAGGTAGAACT
80CNS-2	GAGATTTACTCAGTCCAATG
80CNS-3	TGGTTGTCACAGTAACTAGG
83CNS-1	GTGACAAGCATGACTCTACA
83CNS-2	GTGATGGTGAATTAAGCTGA
83CNS-3	TGGGCTCTGACTCACTTAGA
85CNS-1	GCTAACATTGACTTCTCTAC
85CNS-2	AATCTATGCAAGGGGTGAAT
85CNS-3	GCATCATGATGAAGCTTATC
122CNS-1	GTTATTAATCTAAGCGGAGA
122CNS-2	GGAAGTTAGGCAGTCAATCG
122CNS-3	CACTTTGTGCTCGGATGCTC
128CNS-1	CTTCAATCATTGCATTCCAC
128CNS-2	TCAAGCAGGGGTGGTGTCAA
128CNS-3	TGGTGATTCATCTTTAGCAT
IL-2URR-1	TCCATTCAGTCAGTCTTTGG
IL-2URR-2	TAGTGTCCCAGGTGATTTAG
IL-2URR-3	TAGAGCTATCACCTAAGTGT
B2m-1	GAGTAGCGCGAGCACAGCTA
B2m-2	CACGCGTTTAATATAAGTGG

1028

Primary Human CD4 T (5-10e6) were incubated with sgRNA and CAS9 protein complex and

electroporation was done using P3 Primary Cell 4D-Nucleofector™ X Kit L (Lonza, cat # V4XP-

3024) and Lonza 4D nucleofector system. B2M gene was CRISPR targeted as a positive

control. As per manufacturer's protocol cells were electroporated pulse code FI-115 in 100ul

cuvette format. After nucleofection, cells were allowed to rest in the complete media for ~2

days. Cells were then harvested, washed with PBS and aliquots of cells were used for further

experimentation such as flow staining and cell activation. Primary human CD4 T Cells

(0.1e6/well) were seeded in triplicates (for each experimental condition) in 96 well plate format

in RPMI complete medium and stimulated using ImmunoCult™ Human CD3/CD28 T Cell

Activator (STEM cells Technologies, cat # 10971). Cell culture supernatants were collected at

4h of stimulation and stored at -80C until assayed. IL-2 ELISA was performed using

Thermofisher Scientific IL-2 Human Uncoated ELISA Kit with Plates, cat # 88-7025-76.

Recombinant reporter assays

The *IL2* URR was cloned into Xho I and Hind III sites of PGL4 Luc vector (Promega) and then

~500 bp individual sequence representing *IL2-IL21* intergenic CNS at -46, -51, -80, -83, -85, -

128 were cloned upstream to IL-2 URR at the Xho I site of the pGL4 vector. Primary human

CD4 T cells obtained from 5 normal healthy donors were transfected with PGL4-cRE-URR

constructs. Briefly, primary human CD4 T cells were activated with anti-CD3+ anti-CD28

Dynabeads (1:1) (Thermofisher scientific, cat # 11161D) + IL-2 overnight and then 1 million

aliquots of cells (triplicates) were electroporated using Nucleofector 2b, human T cell

Nuclefactor kit (Lonza VPA # 1002, program # T-020) with 2ug of PGL4 firefly vector constructs

along with 0.2 ug of PGL4 Renilla vector; cells were allowed to rest ON in RPMI + IL-2 and then

re-stimulated with plate-bound anti-CD3+ anti-CD28 (2 ug/ml each) for 5 hrs. Dual luciferase

assay was performed with cell lysate prepared using Promega dual luciferase assay kit. Cell

lysate was prepared in PLB as per manufacturer's protocols and then firefly and renilla

luciferase activities were analyzed by spectramax ID5 (Molecular Devices). Firefly luciferase

activity was normalized against the internal control renilla luciferase activity.

Pharmacologic validation of novel T cell activation regulatory genes

The drugs lometrexol (GART antagonist), 25-hydroxycholesterol (GPR183 ligand), epinephrine

& norepinephrine (CHRNB agonists), and arachidonoyl glycine (GPR18 agonist) were

purchased from Sigma. The GPR18 antagonist O-1918 was purchased from ChemCruz.

CEP1347 (MAP3K11 antagonist), kisspeptin 234 (KISSIR ligand) and the PTK6 antagonist

tilfrinib were purchased from Tocris. The GRK6 antagonist GRK-IN-1 was purchased from DC

Chemicals. The SIK1 antagonist HG-91-9-1 was purchased from Selleckchem. The OXER1

agonist 5-Oxo-ete was purchased from Cayman chemicals. The drugs were dissolved in DMSO

or ethanol as suggested by the vendor and working stocking concentrations were prepared in

RPMI1640 medium or PBS. The drug effect on T cell proliferation was assayed using murine

lymphocytes cultured under TCR and CD28 activation conditions. Spleen and lymph nodes

were collected from 6-8 weeks old female C57BL/7 mice and single cell lymphocyte cell

suspensions were prepared in RPMI 1640 complete medium. 20 million lymphocytes were

labeled with CFSE and re-suspended in RPMI 1640 medium. Cells (0.5 million labeled

cells/well) were loaded into 48 well culture plate and activated with mouse anti-CD3 and anti-

CD28 agonistic antibodies (1μg mL each). Drugs at the indicated concentrations were added in

the culture medium and both untreated and drug treated cell cultures were incubated at 37°C for

72 hours in a cell culture incubator. Cells were harvested after 3 days of culture, washed with

PBS and then stained with live-dead aqua dye. After washing with FACS buffer (PBS containing

2% FBS), cells were stained with fluorochrome conjugated antibodies CD4-APC, CD8-PB,

CD44-Percp Cy5.5 and CD25-BV650. Stained cells were analyzed on a Beckman Coulter

Cytoflex S flow cytometer. The division profile of CD4+ CFSE+ T cells were gated on live

populations. The flow data was analyzed using Flowjo10 software and the number of divided

CD4+ T cells were determined as described previously[120].

1080

FIGURE LEGENDS

Figure 1: Defined gene expression dynamics throughout activation of naïve CD4+ T

cells. (**A**) Diagram of study design: RNA-seq, ATAC-seq, and HiC libraries were prepared from

sorted CD4+CD62L^{hi}CD44^{lo} naïve T cells (donor N=3) prior to or following 8hr or 24hr

stimulation with anti-CD3/28 beads. cREs are classified to genes allowing for interrogation of

the pattern of expression, accessibility, and chromatin structure changes of autoimmune-

associated variants. (**B**) Heatmap showing normalized expression of known markers of T cell

activation. (**C**) K-means clustering of differentially expressed genes into five groups using the

elbow and within-cluster-sum of squares methods to select the number of clusters. The centroid

of the cluster is depicted as a black line, and members of the cluster are depicted as colored

lines. Color indicates the Pearson's correlation coefficient between the gene and the cluster

centroid, with red indicating higher correlation, and blue lower correlation.

Figure 1 – Figure Supplement 1: Library sequencing reproducibility and expression

clustering. (A) PCA of RNA-seq libraries. (B) PCA of ATAC-seq libraries. (C) Stratified

correlation (SCC) of HiC libraries from two donors by stage of activation. (D) Volcano plot of

RNA-seq data for CD4+ T cells unstimulated vs. 8 hr stimulation (top) and 8 hr vs. 24 hr

stimulation (number of DEG indicated). (E) Elbow plot of the within group sum of squares used

to determine optimal cluster number. (F) Genes per expression cluster. (G) Centroids +/-

standard deviation for each cluster.

Figure 2: Chromatin accessibility and architecture changes across three stages of T cell

activation. (**A**) Volcano plots depicting the log$_2$ change in accessibility in all reproducible OCRs

(present in 2 of the 3 replicates) compared to the -log$_{10}$(FDR), significant points are indicated in

red (FDR < 0.05; abs(log$_2$FC) > 1). (**B**) Genomic regions were classified by membership in A vs.

B compartments at each time point. The pie chart depicts regions with differential A/B

compartment assignment in response to activation. (**C**) TAD structure was determined at each

time point, and regions exhibiting a shift, split/merge, change in strength, or more complex

rearrangements are depicted in the pie chart. (**D**) Loop calls identified from HiC data for each

timepoint called at three resolutions (1kb, 2kb, 4kb bins). (**E**) Differential loop-calls called across

all resolutions. (**F**) Density plots of cRE accessibility and gene expression separated by whether

contact frequency increased, decreased, or remained stable from the transition from

unstimulated to 8 hour activation and 8 hour to 24 hour activation.

Figure 2 – Figure Supplement 1: Characterization of CD4+ T cell epigenomic data. (A)

Accessibility of OCR and (B) expression of genes located in

A vs. B compartments (log2 fkpm). Differential accessibility (C) and expression (D) of cRE-gene

pairs in regions exhibiting differential TAD structure. (E) Enrichment of promoter interacting

region OCR for annotated regulatory elements in CD4+ T cells from the epigenome roadmap

project (E038_15). (F) Overlap of cRE-promoter loops defined in this study to loops defined by

Burren et al. in a prior promoter capture Hi-C study. (G) Distribution of OCR contacted per gene

(top, median = 8 OCRs per gene). Distribution of genes contacted per OCR (bottom, median =

3). (H) Differential promoter-OCR connectivity by HiC as a function of differential OCR

accessibility (central bar = median, boxplot edges = 25-75 quantiles, whiskers = 1.5x inter-

quantile range).

Figure 2 – Figure Supplement 2: Characterization of chromatin stripes during CD4+ T cell

activation. (A) Stripes called per timepoint. (B) Proportion of SNPs annotated in A vs B

compartments. Accessibility of OCR (C) and expression of genes (D) located in anchor (small

more interactive region of stripe), stripe excluding the anchor (stripe), or outside stripes.

Figure 2 – Figure Supplement 3: Large-scale epigenetic changes at dynamically

expressed genes during CD4+ T cell activation. HiC contact frequency matrix (heatmap),

chromatin accessibility (grey), and loop calls (red) for (A) GEM, (B) IRF4, (C) KLF2, and (D)

DPEP2. The IRF4 matrix is truncated at by end of the chromosome annotation, and inset shows

the expression (TPM) at each time point for each gene.

Figure 3: Transcription factors prediction potential regulators of chromatin status and

expression changes. Footprints were annotated to each TF motif by sequence matching to the

JASPAR motif database, and the average accessibility of the region surrounding each motif was

used as a measure of activity at each time point. Scatterplots depict the change in accessibility

for each TF motif (activity score) and $\log_2$FC of TF gene expression between unstimulated and 8

hour activation (**A**) or 8 hour and 24 hour activation (**B**). TF were classified as differentially

expressed (orange), differentially active (brown), both (red), or neither (grey). Dot size indicates

the number of predicted footprints occupied by each motif. (**C**) Average accessibility (normalized

by depth and motif number) in a 200bp window surrounding motif footprints for FOSL2::JUNB,

Smad2::Smad3, MAZ, and KLF2 from three timepoints (unstim: green; 8hr: blue; 24hr: red). (**D**)

Z-score of TF motif enrichment for cRE connected to the five clusters compared to all OCR.

Lighter color indicates higher specificity of enrichment for that TF cluster. (**E**) Connections

between different TFs based on physical interactions and predicted regulation-based co-

expression determined by GENIE3 for the three timepoints. Node color indicates expression

(TPM; darker = higher expression, lighter = lower expression), edge color reflects confidence in

the interaction called by GENIE3 (darker higher confidence).

Figure 3 – Figure Supplement 1: Enrichment of transcription family members and gene

regulatory network construction. (A) -log10(P) enrichment of transcription factors annotated

by family for each expression cluster. (B) Top five pathways enriched for a subset of TF-

regulated gene expression.

Figure 4: Variant-to-gene mapping of autoimmune-associated loci implicates genetic

variants in the control of T cell activation. (**A**) Identification of genes in contact with open

chromatin regions harboring autoimmune-associated SNPs. Sentinel SNPs from 15

autoimmune GWAS (**Table S11**) were used to identify the 95% credible set of proxy SNPs for

each lead SNP. SNP locations were integrated with ATAC-seq and HiC data to identify the 95%

credible set of accessible SNPs in physical contact with a gene promoter. Genes are further

refined based on expression dynamics over the time course of T cell activation. (**B**) Partitioned

LD score regression for autoimmune GWAS studies using the OCR in contact with the genes in

the five clusters defined by RNA-seq. Circle size = enrichment, color = significance,

FDR<0.01, ***FDR<0.001. (C**) Log distribution of the 1D distance between each proxy SNP

and its interacting gene based on 3D chromatin conformation (median=103 kbp). (**D**)

Distribution of the number of genes contacted by each accessible variant (median=5). (**E**)

Fraction of open disease-associated variants that interact with no gene promoters (grey), only

with the nearest gene promoter (purple), with the nearest gene and a distant gene(s) (orange),

and with only a distant gene(s) (blue). (**F**) Number of genes identified by variant-to-gene

mapping at each time point. Black = shared in all stages, blue = shared in two stages, red =

specific to one stage. (**G**) Set of implicated proxy-genes pairs that are both differentially

expressed, display differential accessibility, and chromatin contact across T cell activation. (**H**)

Example MS variant rs1077667 (PP=1.0) exhibits increased accessibility and contact with the

promoters of the *GPR108* and *TRIP10* at 8 hours post activation, which are increased in

expression at this time point. (**I**) Example allergy variant rs7380290 interacts with the *SIK1* gene

that is upregulated after activation.

Figure 5: **Comparative predictive power of orthogonal V2G approaches**. (**A**) Percent

overlap in predicted genes among chromatin-based and eQTL-based V2G approaches. (**B**)

Physical variant-gene associations (HiC) are enriched for statistical variant-gene associations

(eQTL) in activated CD4+ T cells. The histogram depicts the null distribution of shared variant-

gene pairs expected at random (~5) while the red line indicates the observed number of variant-

gene pairs (41) shared with the 127 eQTL identified by Soskic et al. in a similar CD4+ T cell

activation system. (**C**) Expression of the shared genes. (**D**) Precision-Recall analysis of V2G

gene predictions against the set of monogenic human inborn errors in immunity.

Figure 6: Functional validation of autoimmune V2G-implicated cRE at the *IL2* locus. (**A**)

The combination of evolutionary conservation (blue), open chromatin (red), and autoimmune

disease-associated SNPs at the *IL2* locus identify putative cRE in quiescent vs. 8-hour activated

naïve CD4+ T cells. (**B**) Activation-dependent TAD/sub-TAD structure (heatmaps), chromatin

remodeling (grey) and promoter-OCR interactions (red) at the *IL2* locus. (**C**) Recombinant

reporter assay showing transcriptional activity of the *IL2* URR (+35 to -500 from the TSS) in

activated primary CD4+ T cells (N= 7 donors) compared to a basal promoter (pGL4, left panel).

The right panel depicts transcriptional activity of the CNS regions indicated in **A** cloned

upstream of the URR. All regions except the -80 CNS show statistically significant activity

relative to the URR alone (N=7, P<0.05, line = median, box = 95/5% range). (**D**) Scheme of

CRISPR/CAS9-based deletion of individual *IL2* CNS using flanking gRNAs. (**E**) Activation-

induced secretion of IL-2 protein by CRISPR-targeted primary CD4+ T cells (N=5 donors, left

panel) or Jurkat cells (N=3 replicates, right panel) relative to untargeted control (CAS9, no

gRNA) cells (****P<0.0001, ***P<0.001, **P<0.01, *P<0.05, line = median, box = 95/5% range).

In primary cells, *B2M* gRNAs served as an irrelevant targeted control. (**F**) Scheme of

CRISPR/CAS9-based deletion of the 81.3 kb region containing all distal *IL2* cRE using flanking

gRNAs in Jurkat cells. (**G**) Activation-induced IL-2 protein (left panel) and mRNA (right panel) by

control (black) vs. 81.3 kb deleted (red) Jurkat cells (N=3 separate clones).

Figure 6 – Figure Supplement 1: TF occupancy and stability of distal IL2 cRE in CD4+ T

cell subsets. Evolutionary conservation (blue), open chromatin (red), and autoimmune disease-

associated SNPs at the IL2 locus in quiescent (A) and 8-hour activated (B) naïve CD4+ T cells.

(C) Curated TF occupancy at the IL2 URR and distal cRE from the ReMap atlas of regulatory

regions.

Figure 7: Distal *IL2* enhancers are evolutionarily conserved and impact *in vivo* T cell-

mediated immunity in mice. (**A**) Map of the human and mouse *IL2* locus depicting mammalian

conservation (dark blue), ATAC-seq from activated human and mouse CD4+ T cells (red), and

orthologous non-coding sequences conserved between mouse and human (light blue). (**B**) IL-2

protein secretion (top left panel) and *Il2* mRNA expression (top right panel) by CD4+ T cells

from wild-type or *Il2*-83 cRE ko mice in response to stimulation with plate-bound anti-CD3 and

anti-CD28 Ab *in vitro*. Bottom panels show soluble anti-CD3-induced *in vitro* clonal expansion of

wild-type or *Il2*-83 cRE ko CD4+ (left) and CD8+ (right) T cells measured by dye dilution[120]. (**C**)

Foxp3 expression by murine CD4+CD25- Tconv stimulated with anti-CD3 Ab and TGF-beta in

the absence (top panels) vs. presence (bottom panels) of exogenous IL-2. Wild-type or *Il2*-83

cRE ko mice (N=6) were immunized intraperitoneally with chicken ovalbumin in incomplete

Freund's adjuvant. The frequency of CD4+PD1+CXCR5^{hi} follicular helper T cells in the spleens

of 3 animals was measured by flow cytometry at day 5 post-immunization (**D**), and levels of

ovalbumin-specific IgG in the serum of 3 animals were measured at day 10 post-immunization

(**E**). (**F**) In vivo IBD model. CD4+CD25- Tconv from wild-type (pink, red) or *Il2*-83 cRE ko

(purple, blue) mice were transferred alone (purple, pink) or together with wild-type CD4+CD25+

Treg (red, blue) into RAG1-ko mice (N=5 per group). Animal weight was monitored for 40 days,

and the number (in millions) of activated CD4+ Tconv in the spleens and mesenteric lymph

nodes was measured by flow cytometry (inset). Statistical significance (*p<0.05) was measured

by ANOVA or t-test.

Figure 8: Functional validation of autoimmune V2G-implicated genes. (**A**) Recombinant

reporter assay in primary activated CD4+ T cells (N= 7 donors) showing transcriptional activity

of the reference vs. risk alleles of the *IL2* -128 cRE relative to the URR alone (P<0.0001). (**B**)

Prominent TF motifs predicted to be disrupted (blue) or stabilized (red) by promoter-connected

autoimmune SNPs. (**C**) 3D chromatin-based V2G genes are enriched for CRISPR-implicated

genes that regulate CD4+ T cell activation. Observed enrichment of genes regulating multiple

aspects of CD4+ T cell activation (IL-2, IL-2 receptor, CTLA-4, IFNg, TNFa or proliferation) from

CRISPR screens by Freimer, Schmidt, and Shifrut among sets of 3D chromatin-implicated

genes among individual diseases (green, FDR<0.05) or all diseases (purple, FDR<0.05). (**D**)

Enrichment for 3D chromatin-based autoimmune V2G genes among genes with germline

knock-out mouse immune (red) and other (black) phenotypes (adjP<0.05, IMPC database). (**E**)

Autoimmune V2G-implicated genes with at least one pharmacologic modulator (rDGIdb). (**F**)

Comparison of the number of manuscripts retrieved from PubMed related to autoimmune

disease for each V2G gene with pharmaceutical agents available (x-axis) with an immune-

specific expression score computed using the sum GTEX median expression values (v8) for

whole blood or spleen divided by other tissues (y-axis). Genes highlighted in red were selected

for functional validation in **G**. (**G**) Dose-dependent impact of the indicated pharmacologic agents

targeting the V2G-implicated genes *KISS1R, CHRNB, OXER1, GPR18, GRK6, PTK6,*

MAP3K11, GPR183, GART, and *SIK1* on proliferation of anti-CD3/28 activated murine (left

panel) or human (right panel) CD4+ T cells *in vitro* (N=4).

Figure 8 – Figure Supplement 1: Predicted impact of autoimmune disease-associated genetic

variation at V2G-implicated loci. (A) Top 50 TF motifs impacted by autoimmune SNPs (see

Table S15). HiC contact frequency matrix (heatmap), chromatin accessibility (grey), and loop

calls (blue-IL19/ITGA4, red-FAIM3) at the T1D-rs3024505 locus (B) and the celiac- rs13010713

locus (C). Dashed box indicates the SNP affected by allelic variation. Genetic variation at

rs3024505 is predicted to disrupt a MZF1 motif (D), and variation at rs13010713 is predicted to

disrupt a SOX4 binding motif (E). (F) Variation at MS SNP rs1077667 is predicted to disrupt

TP53, TP63, and POU2F2 (OCT2) binding sites.

Figure 8 – Figure Supplement 2: Orthogonal validation of 3D chromatin-based V2G

genes. Expression (scaled TPM) of genes implicated by chromatin contacts in (A) prior

CRISPRi/a screens and (C) genes with known immune phenotypes in the international mouse

phenotyping consortium. Red = increased, blue = decreased.

1257

Supplemental Files

Supplementary File 1: Expression and clustering of differentially expressed genes.

Supplementary File 2: Pathway enrichment of differentially expressed RNA-seq genes

Supplementary File 3: Pathway enrichment of differentially expressed RNA-seq genes by

cluster

Supplementary File 4: Accessible chromatin regions

Supplementary File 5: Differential accessible open chromatin regions

Supplementary File 6: A/B compartment calls

Supplementary File 7: Differential TAD boundaries

Supplementary File 8: Stripe calls

Supplementary File 9: Differential contact frequency

Supplementary File 10: Summary of TF footprinting

Supplementary File 11: TF target gene pathway enrichment

Supplementary File 12: List of all GWAS studies included

Supplementary File 13: Variant to gene mapping across all timepoints

Supplementary File 14: Partitioned LD score regression output

Supplementary File 15: Autoimmune variants in open gene promoters

Supplementary File 16: Comparison of V2G approaches

Supplementary File 17: Motifs predicted to disrupt transcription factor binding sites.

Supplementary File 18: V2G genes implicated by orthogonal data

Supplementary File 19: V2G target gene drug repurposing results

1279

1280

1281

REFERENCES

1. Zhang, F. & Lupski, J. R. Non-coding genetic variants in human disease. *Hum Mol Genet* **24**, R102-10 (2015).
2. Weinreich, M. A. *et al.* KLF2 Transcription-Factor Deficiency in T Cells Results in Unrestrained Cytokine Production and Upregulation of Bystander Chemokine Receptors. *Immunity* **31**, 122–130 (2009).
3. Tzachanis, D. *et al.* Tob is a negative regulator of activation that is expressed in anergic and quiescent T cells. *Nat Immunol* **2**, 1174–82 (2001).
4. Yusuf, I. & Fruman, D. A. Regulation of quiescence in lymphocytes. *Trends Immunol* **24**, 380–386 (2003).
5. Zhu, J., Yamane, H. & Paul, W. E. Differentiation of Effector CD4 T Cell Populations. *Annu Rev Immunol* **28**, 445–489 (2010).
6. Saravia, J., Chapman, N. M. & Chi, H. Helper T cell differentiation. *Cell Mol Immunol* **16**, 634–643 (2019).
7. Farh, K. K.-H. *et al.* Genetic and epigenetic fine mapping of causal autoimmune disease variants. *Nature* **518**, 337 343 (2014).
8. Chen, L. *et al.* Genetic Drivers of Epigenetic and Transcriptional Variation in Human Immune Cells. *Cell* **167**, 1398 1414.e24 (2016).
9. Soskic, B. *et al.* Chromatin activity at GWAS loci identifies T cell states driving complex immune diseases. *Nat Genet* 1–8 (2019) doi:10.1038/s41588-019-0493-9.
10. Robertson, C. C. *et al.* Fine-mapping, trans-ancestral and genomic analyses identify causal variants, cells, genes and drug targets for type 1 diabetes. *Nat Genet* **53**, 962–971 (2021).
11. Lu, X. *et al.* Global discovery of lupus genetic risk variant allelic enhancer activity. *Nat Commun* **12**, 1611 (2021).
12. Chandra, V. *et al.* Promoter-interacting expression quantitative trait loci are enriched for functional genetic variants. *Nat Genet* **53**, 110–119 (2021).
13. Soskic, B. *et al.* Immune disease risk variants regulate gene expression dynamics during CD4+ T cell activation. *Nat Genet* 1–10 (2022) doi:10.1038/s41588-022-01066-3.
14. Mouri, K. *et al.* Prioritization of autoimmune disease-associated genetic variants that perturb regulatory element activity in T cells. *Nat Genet* 1–10 (2022) doi:10.1038/s41588-022-01056-5.
15. Rioux, J. D. & Abbas, A. K. Paths to understanding the genetic basis of autoimmune disease. *Nature* **435**, 584–589 (2005).
16. Buckner, J. H. Mechanisms of impaired regulation by CD4(+)CD25(+)FOXP3(+) regulatory T cells in human autoimmune diseases. *Nat Rev Immunol* **10**, 849 859 (2010).
17. Javierre, B. M. *et al.* Lineage-Specific Genome Architecture Links Enhancers and Non-coding Disease Variants to Target Gene Promoters. *Cell* **167**, 1369 1384.e19 (2016).
18. Burren, O. S. *et al.* Chromosome contacts in activated T cells identify autoimmune disease candidate genes. *Genome Biol* **18**, 165 (2017).
19. Yang, J. *et al.* Analysis of chromatin organization and gene expression in T cells identifies functional genes for rheumatoid arthritis. *Nat Commun* **11**, 4402 (2020).
20. Bykova, M., Hou, Y., Eng, C. & Cheng, F. Quantitative trait locus (xQTL) approaches identify risk genes and drug targets from human non-coding genomes. *Hum Mol Genet* **31**, R105–R113 (2022).
21. Shifrut, E. *et al.* Genome-wide CRISPR Screens in Primary Human T Cells Reveal Key Regulators of Immune Function. *Cell* **175**, 1958-1971.e15 (2018).
22. Schmidt, R. *et al.* CRISPR activation and interference screens decode stimulation responses in primary human T cells. *Science* **375**, eabj4008 (2022).

23. Freimer, J. W. *et al.* Systematic discovery and perturbation of regulatory genes in human T cells reveals the architecture of immune networks. *Nat Genet* **54**, 1133–1144 (2022).

24. Su, C., Pahl, M. C., Grant, S. F. A. & Wells, A. D. Restriction enzyme selection dictates detection range sensitivity in chromatin conformation capture-based variant-to-gene mapping approaches. *Hum Genet* (2021) doi:10.1007/s00439-021-02326-8.

25. Thurman, R. E. *et al.* The accessible chromatin landscape of the human genome. *Nature* **489**, 75 (2012).

26. Lieberman-Aiden, E. *et al.* Comprehensive Mapping of Long-Range Interactions Reveals Folding Principles of the Human Genome. *Science* **326**, 289–293 (2009).

27. Dixon, J. R. *et al.* Topological domains in mammalian genomes identified by analysis of chromatin interactions. *Nature* **485**, 376 380 (2012).

28. Mirny, L. A., Imakaev, M. & Abdennur, N. Two major mechanisms of chromosome organization. *Curr Opin Cell Biol* **58**, 142–152 (2019).

29. Yoon, S., Chandra, A. & Vahedi, G. Stripenn detects architectural stripes from chromatin conformation data using computer vision. *Nat Commun* **13**, 1602 (2022).

30. Kim, T.-K. & Shiekhattar, R. Architectural and Functional Commonalities between Enhancers and Promoters. *Cell* **162**, 948 959 (2015).

31. Hamamoto, K. & Fukaya, T. Molecular architecture of enhancer–promoter interaction. *Curr Opin Cell Biol* **74**, 62–70 (2022).

32. Consortium, R. E. *et al.* Integrative analysis of 111 reference human epigenomes. *Nature* **518**, 317 330 (2015).

33. Yoon, J.-H. *et al.* Phosphorylation status determines the opposing functions of Smad2/Smad3 as STAT3 cofactors in TH17 differentiation. *Nat Commun* **6**, 7600 (2015).

34. Safford, M. *et al.* Egr-2 and Egr-3 are negative regulators of T cell activation. *Nat Immunol* **6**, 472–480 (2005).

35. Schubert, L. A., Jeffery, E., Zhang, Y., Ramsdell, F. & Ziegler, S. F. Scurfin (FOXP3) Acts as a Repressor of Transcription and Regulates T Cell Activation. *J Biol Chem* **276**, 37672–37679 (2001).

36. Xiao, T., Li, X. & Felsenfeld, G. The Myc-associated zinc finger protein (MAZ) works together with CTCF to control cohesin positioning and genome organization. *Proc National Acad Sci* **118**, e2023127118 (2021).

37. Sebzda, E., Zou, Z., Lee, J. S., Wang, T. & Kahn, M. L. Transcription factor KLF2 regulates the migration of naive T cells by restricting chemokine receptor expression patterns. *Nat Immunol* **9**, 292–300 (2008).

38. Moskowitz, D. M. *et al.* Epigenomics of human CD8 T cell differentiation and aging. *Sci Immunol* **2**, eaag0192 (2017).

39. Harris, J. E. *et al.* Early Growth Response Gene-2, a Zinc-Finger Transcription Factor, Is Required for Full Induction of Clonal Anergy in CD4+ T Cells. *J Immunol* **173**, 7331–7338 (2004).

40. Williams, J. B. *et al.* The EGR2 targets LAG-3 and 4-1BB describe and regulate dysfunctional antigen-specific CD8+ T cells in the tumor microenvironment. *J Exp Med* **214**, 381–400 (2017).

41. Avitahl, N. *et al.* Ikaros Sets Thresholds for T Cell Activation and Regulates Chromosome Propagation. *Immunity* **10**, 333–343 (1999).

42. Bandyopadhyay, S. *et al.* Interleukin 2 gene transcription is regulated by Ikaros-induced changes in histone acetylation in anergic T cells. *Blood* **109**, 2878–86 (2007).

43. Thomas, R. M. *et al.* Ikaros Enforces the Costimulatory Requirement for IL2 Gene Expression and Is Required for Anergy Induction in CD4+ T Lymphocytes. *J Immunol* **179**, 7305–7315 (2007).

44. Thomas, R. M. *et al.* Ikaros Silences T-bet Expression and Interferon-γ Production during T Helper 2 Differentiation. *J Biol Chem* **285**, 2545–2553 (2010).

45. Weber, B. N. *et al.* A critical role for TCF-1 in T-lineage specification and differentiation.
Nature **476**, 63–68 (2011).
46. Johnson, J. L. *et al.* Lineage-Determining Transcription Factor TCF-1 Initiates the Epigenetic
Identity of T Cells. *Immunity* **48**, 243-257.e10 (2018).
47. Wang, W. *et al.* TCF-1 promotes chromatin interactions across topologically associating
domains in T cell progenitors. *Nat Immunol* 1–11 (2022) doi:10.1038/s41590-022-01232-
z.
48. Furukawa, T. *et al.* Involvement of PLAGL2 in activation of iron deficient- and hypoxia-
induced gene expression in mouse cell lines. *Oncogene* **20**, 4718–4727 (2001).
49. Wu, Q. *et al.* Hypoxia-inducible factors: master regulators of hypoxic tumor immune escape.
J Hematol Oncol **15**, 77 (2022).
50. Fu, W. *et al.* A multiply redundant genetic switch "locks in" the transcriptional signature of
regulatory T cells. *Nat Immunol* **13**, 972–980 (2012).
51. Pramanik, J. *et al.* Genome-wide analyses reveal the IRE1a-XBP1 pathway promotes T
helper cell differentiation by resolving secretory stress and accelerating proliferation.
Genome Med **10**, 76 (2018).
52. Gao, X., Tewari, K., Svaren, J. & Suresh, M. Role of cell cycle regulator E2F1 in regulating
CD8 T cell responses during acute and chronic viral infection. *Virology* **324**, 567–576
(2004).
53. DeRyckere, D. & DeGregori, J. E2F1 and E2F2 Are Differentially Required for Homeostasis-
Driven and Antigen-Induced T Cell Proliferation In Vivo. *J Immunol* **175**, 647–655 (2005).
54. Cook, M. E., Jarjour, N. N., Lin, C.-C. & Edelson, B. T. Transcription Factor Bhlhe40 in
Immunity and Autoimmunity. *Trends Immunol* **41**, 1023–1036 (2020).
55. Chesi, A. *et al.* Genome-scale Capture C promoter interactions implicate effector genes at
GWAS loci for bone mineral density. *Nat Commun* **10**, 1260 (2019).
56. Su, C. *et al.* Mapping effector genes at lupus GWAS loci using promoter Capture-C in
follicular helper T cells. *Nat Commun* **11**, 3294 (2020).
57. Pahl, M. C. *et al.* Cis-regulatory architecture of human ESC-derived hypothalamic neuron
differentiation aids in variant-to-gene mapping of relevant complex traits. *Nat Commun*
12, 6749 (2021).
58. Pahl, M. C. *et al.* Implicating effector genes at COVID-19 GWAS loci using promoter-
focused Capture-C in disease-relevant immune cell types. *Genome Biol* **23**, 125 (2022).
59. Su, C. *et al.* 3D chromatin maps of the human pancreas reveal lineage-specific regulatory
architecture of T2D risk. *Cell Metab* **34**, 1394-1409.e4 (2022).
60. Feng, Y. *et al.* The Cdc42-interacting Protein-4 (CIP4) Gene Knock-out Mouse Reveals
Delayed and Decreased Endocytosis*. *J Biol Chem* **285**, 4348–4354 (2010).
61. Darling, N. J. & Cohen, P. Nuts and bolts of the salt-inducible kinases (SIKs). *Biochem J*
478, 1377–1397 (2021).
62. Gate, R. E. *et al.* Genetic determinants of co-accessible chromatin regions in activated T
cells across humans. *Nat Genet* **9**, 1 (2018).
63. Schmiedel, B. J. *et al.* Single-cell eQTL analysis of activated T cell subsets reveals
activation and cell type–dependent effects of disease-risk variants. *Sci Immunol* **7**,
eabm2508 (2022).
64. Ye, C. J. *et al.* Intersection of population variation and autoimmunity genetics in human T
cell activation. *Science* **345**, 1254665 1254665 (2014).
65. Tangye, S. G. *et al.* Human Inborn Errors of Immunity: 2022 Update on the Classification
from the International Union of Immunological Societies Expert Committee. *J. Clin.
Immunol.* **42**, 1473–1507 (2022).
66. Nasser, J. *et al.* Genome-wide enhancer maps link risk variants to disease genes. *Nature* 1–
6 (2021) doi:10.1038/s41586-021-03446-x.

67. Fulco, C. P. *et al.* Activity-by-contact model of enhancer–promoter regulation from
thousands of CRISPR perturbations. *Nat. Genet.* **51**, 1664–1669 (2019).
68. Cerosaletti, K. *et al.* Multiple autoimmune-associated variants confer decreased IL-2R
signaling in CD4+ CD25(hi) T cells of type 1 diabetic and multiple sclerosis patients. *Plos*
One **8**, e83811 (2013).
69. Spolski, R., Li, P. & Leonard, W. J. Biology and regulation of IL-2: from molecular
mechanisms to human therapy. *Nat Rev Immunol* **18**, 648 659 (2018).
70. Abbas, A. K., Trotta, E., Simeonov, D. R., Marson, A. & Bluestone, J. A. Revisiting IL-2:
Biology and therapeutic prospects. *Sci Immunol* **3**, eaat1482 (2018).
71. Joosse, M. E. *et al.* Duplication of the IL2RA locus causes excessive IL-2 signaling and may
predispose to very early onset colitis. *Mucosal Immunol* 1–11 (2021)
doi:10.1038/s41385-021-00423-5.
72. Fujita, T., Shibuya, H., Ohashi, T., Yamanishi, K. & Taniguchi, T. Regulation of human
interleukin-2 gene: Functional DNA sequences in the 5′ flanking region for the gene
expression in activated T lymphocytes. *Cell* **46**, 401–407 (1986).
73. Durand, D. B., Bush, M. R., Morgan, J. G., Weiss, A. & Crabtree, G. R. A 275 basepair
fragment at the 5' end of the interleukin 2 gene enhances expression from a
heterologous promoter in response to signals from the T cell antigen receptor. *J Exp*
Medicine **165**, 395 407 (1987).
74. Novak, T. J., White, P. M. & Rothenberg, E. V. Regulatory anatomy of the murine
interleukin-2 gene. *Nucleic Acids Res* **18**, 4523–4533 (1990).
75. Mehra, P. & Wells, A. D. Long-Range Transcriptional Control of the Il2 Gene by an
Intergenic Enhancer. *Mol Cell Biol* **35**, 3880–3891 (2015).
76. Consortium, T. E. P. The ENCODE (ENCyclopedia Of DNA Elements) Project. *Science* **306**,
636–640 (2004).
77. Lin, X. *et al.* Nested epistasis enhancer networks for robust genome regulation. *Science*
377, 1077–1085 (2022).
78. Hedlund, G., Dohlsten, M., Ericsson, P. O. & Sjögren, H. O. Rapid response to Con A by
CD4+CD45R− rat memory lymphocytes as compared to CD4+CD45R+ lymphocytes.
Cell Immunol **119**, 317–326 (1989).
79. Picker, L. *et al.* Direct demonstration of cytokine synthesis heterogeneity among human
memory/effector T cells by flow cytometry. *Blood* **86**, 1408–1419 (1995).
80. Sojka, D. K., Bruniquel, D., Schwartz, R. H. & Singh, N. J. IL-2 Secretion by CD4+ T Cells In
Vivo Is Rapid, Transient, and Influenced by TCR-Specific Competition. *J Immunol* **172**,
6136–6143 (2004).
81. Powrie, F., Correa-Oliveira, R., Mauze, S. & Coffman, R. L. Regulatory interactions between
CD45RBhigh and CD45RBlow CD4+ T cells are important for the balance between
protective and pathogenic cell-mediated immunity. *J Exp Medicine* **179**, 589–600 (1994).
82. Morawski, P. A., Mehra, P., Chen, C., Bhatti, T. & Wells, A. D. Foxp3 Protein Stability Is
Regulated by Cyclin-dependent Kinase 2. *J Biol Chem* **288**, 24494–24502 (2013).
83. Smigiel, K. S. *et al.* CCR7 provides localized access to IL-2 and defines homeostatically
distinct regulatory T cell subsets. *J Exp Medicine* **211**, 121–136 (2014).
84. Mostafavi, H., Spence, J. P., Naqvi, S. & Pritchard, J. K. Systematic differences in discovery
of genetic effects on gene expression and complex traits. *Nat. Genet.* 1–10 (2023)
doi:10.1038/s41588-023-01529-1.
85. Kovanen, P. E. *et al.* T-cell development and function are modulated by dual specificity
phosphatase DUSP5. *J Biological Chem* **283**, 17362–9 (2008).
86. Chuang, H.-C. & Tan, T.-H. MAP4K Family Kinases and DUSP Family Phosphatases in T-
Cell Signaling and Systemic Lupus Erythematosus. *Cells* **8**, 1433 (2019).
87. Danileviciute, E. *et al.* PARK7/DJ-1 promotes pyruvate dehydrogenase activity and
maintains Treg homeostasis during ageing. *Nat Metabolism* **4**, 589–607 (2022).

88. McHugh, D. *et al.* N-arachidonoyl glycine, an abundant endogenous lipid, potently drives directed cellular migration through GPR18, the putative abnormal cannabidiol receptor. *Bmc Neurosci* **11**, 44 (2010).

89. Kan, J. L., Jannatipour, M., Taylor, S. M. & Moran, R. G. Mouse cDNAs encoding a trifunctional protein of de novo purine synthesis and a related single-domain glycinamide ribonucleotide synthetase. *Gene* **137**, 195–202 (1993).

90. Varney, M. J. *et al.* G protein–coupled receptor kinase 6 (GRK6) regulates insulin processing and secretion via effects on proinsulin conversion to insulin. *J Biol Chem* **298**, 102421 (2022).

91. Gilic, M. B. & Tyner, A. L. Targeting protein tyrosine kinase 6 in cancer. *Biochimica Et Biophysica Acta Bba - Rev Cancer* **1874**, 188432 (2020).

92. Slattery, M. L., Lundgreen, A. & Wolff, R. K. MAP kinase genes and colon and rectal cancer. *Carcinogenesis* **33**, 2398–2408 (2012).

93. Knackmuss, U. *et al.* MAP3K11 is a tumor suppressor targeted by the oncomiR miR-125b in early B cells. *Cell Death Differ* **23**, 242–252 (2016).

94. Hammond, R. K. *et al.* Biological constraints on GWAS SNPs at suggestive significance thresholds reveal additional BMI loci. *Elife* **10**, e62206 (2021).

95. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat Methods* **9**, 357–359 (2012).

96. Zhang, Y. *et al.* Model-based Analysis of ChIP-Seq (MACS). *Genome Biol* **9**, R137 (2008).

97. Lun, A. T. L. & Smyth, G. K. csaw: a Bioconductor package for differential binding analysis of ChIP-seq data using sliding windows. *Nucleic Acids Res* **44**, e45–e45 (2016).

98. Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinform Oxf Engl* **26**, 139–40 (2009).

99. Wingett, S. *et al.* HiCUP: pipeline for mapping and processing Hi-C data. *F1000research* **4**, 1310 (2015).

100. Abdennur, N. & Mirny, L. A. Cooler: scalable storage for Hi-C data and other genomically labeled arrays. *Bioinform Oxf Engl* **36**, 311–316 (2019).

101. Imakaev, M. *et al.* Iterative correction of Hi-C data reveals hallmarks of chromosome organization. *Nat Methods* **9**, 999 1003 (2012).

102. Nora, E. P. *et al.* Molecular basis of CTCF binding polarity in genome folding. *Nat Commun* **11**, 5612 (2020).

103. Cresswell, K. G. & Dozmorov, M. G. TADCompare: An R Package for Differential and Temporal Analysis of Topologically Associated Domains. *Frontiers Genetics* **11**, 158 (2020).

104. Kaul, A., Bhattacharyya, S. & Ay, F. Identifying statistically significant chromatin contacts from Hi-C data with FitHiC2. *Nat Protoc* **15**, 991–1012 (2020).

105. Stansfield, J. C., Cresswell, K. G., Vladimirov, V. I. & Dozmorov, M. G. HiCcompare: an R-package for joint normalization and comparison of HI-C datasets. *Bmc Bioinformatics* **19**, 279 (2018).

106. Anders, S., Pyl, P. T. & Huber, W. HTSeq--a Python framework to work with high-throughput sequencing data. *Bioinform Oxf Engl* **31**, 166–9 (2014).

107. Hänzelmann, S., Castelo, R. & Guinney, J. GSVA: gene set variation analysis for microarray and RNA-Seq data. *Bmc Bioinformatics* **14**, 7–7 (2013).

108. Liberzon, A. *et al.* The Molecular Signatures Database Hallmark Gene Set Collection. *Cell Syst* **1**, 417–425 (2015).

109. Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Res* **43**, e47–e47 (2015).

110. Li, Z. *et al.* Identification of transcription factor binding sites using ATAC-seq. *Genome Biol* **20**, 45 (2019).

111. Fornes, O. *et al.* JASPAR 2020: update of the open-access database of transcription factor
binding profiles. *Nucleic Acids Res* **48**, D87–D92 (2020).
112. Finucane, H. K. *et al.* Partitioning heritability by functional annotation using genome-wide
association summary statistics. *Nat Genet* **47**, 1228–35 (2015).
113. Su, C. *et al.* 3D promoter architecture re-organization during iPSC-derived neuronal cell
differentiation implicates target genes for neurodevelopmental disorders. *Prog Neurobiol*
102000 (2021) doi:10.1016/j.pneurobio.2021.102000.
114. Lawrence, M. *et al.* Software for Computing and Annotating Genomic Ranges. *Plos*
Comput Biol **9**, e1003118 (2013).
115. Lopez-Delisle, L. *et al.* pyGenomeTracks: reproducible plots for multivariate genomic
datasets. *Bioinform Oxf Engl* **37**, 422–423 (2020).
116. Ramírez, F., Dündar, F., Diehl, S., Grüning, B. A. & Manke, T. deepTools: a flexible
platform for exploring deep-sequencing data. *Nucleic Acids Res* **42**, W187–W191
(2014).
117. Muñoz-Fuentes, V. *et al.* The International Mouse Phenotyping Consortium (IMPC): a
functional catalogue of the mammalian genome that informs conservation. *Conservation*
Genetics Print **19**, 995–1005 (2017).
118. Wagner, A. H. *et al.* DGIdb 2.0: mining clinically relevant drug–gene interactions. *Nucleic*
Acids Res **44**, D1036–D1044 (2016).
119. Consortium, T. Gte. *et al.* The Genotype-Tissue Expression (GTEx) pilot analysis:
Multitissue gene regulation in humans. *Science* **348**, 648–660 (2015).
120. Wells, A. D., Gudmundsdottir, H. & Turka, L. A. Following the fate of individual T cells
throughout activation and clonal expansion. Signals from T cell receptor and CD28
differentially regulate the induction and duration of a proliferative response. *J Clin Invest*
100, 3173 3183 (1997).
1560
1561

Figure 1

Figure 2

A Chromatin accessibility dynamics (ATAC-seq)

B AB compartment

C Topologically associating domains

D

E

F Integrated accessibility, connectivity, and expression dynamics

Figure 3

Figure 4

Figure 5

Figure 6

Figure 8

A

Luciferase activity (fold over URR)

p<0.0001
***

-128 OCR NON-RISK (blue)   -128 OCR RISK SNP (gray)

B

TF affinity

# of SNPs

higher
lower

TF motif: NFKB   KLF   FOS::JUN   EGR   FKH

C

V2G geneset enrichment for CRISPR-validated targets

IL2   CTLA4   IL2RA   IFNG   proliferation   TNF

Any GWAS, ALG, AS, ATD, CEL, CRO, ECZ, IBD, JIA, MS, PSO, RA, SLE, T1D, UC, VIT

Enriched
not significant

Enrichment −log10(P value)

D

Reported Mouse Phenotypes

V2G implicated genes (−log10 Pvalue)

Reported phenotype: muscle, pigmentation, respiratory system, adipose tissue, liver/biliary system, renal/urinary system, hearing/vestibular/ear, edocrine/exocrine, cardiovascular system, reproductive system, nervous system, digestive/alimentary, vision/eye, craniofacial, limbs/digits/tail, integument, immune system, hematopoietic system, embryo, skeleton, behvior/neurological, growth/size/body region, homeostasis/metabolism, mortality/aging

E

chemical available
19.7%
80.3%
chemical not available

F

immune specificity score

LTB
IL1R2
LST1
CD244
RAC2   PTPRC   CCR2
ELANE   CD247   FASLG
TNFRSF9
IKZF1 CD6   CD5
ICOS
O XER1
MAP3K11
GPR183
GRK6   GPR18
RCSD1
TXK   INPP5D   CD27   CD28
IL2RB   CCR6 CCR5
CD84   PDCD1 CTLA4   CD10
ICOS   IL10
TNF
C3

log2(pubmed_count + 1)

G

mouse   human

% max. CD4+ T cell activation

log [drug] uM

nM   uM   mM

- GPR18 (O-1918)
- KISS1R (kisspeptin)
- CHRNB (epinephrine)
- CHRNB (norepinephrine)
- OXER1 (5-oxo-ETE)
- GPR18 (Arachidonylglycine)
- GRK6 (GRK-IN-1)
- PTK6 (Tilfrinib)
- MAP3K11 (CEP-1347)
- GPR183 (25-HC)
- GART (lometrexol)
- SIK1 (HG-91-9-1)

Figure 1 - Figure Supplement 1

Figure 2 - Figure Supplement 1

Figure 2 - Figure Supplement 2

Figure 2 - Figure Supplement 3

Figure 3 - Figure Supplement 1

Figure 6 - Figure Supplement 1

Figure 8 - Figure Supplement 1

A Top 50 SNP-affected TF binding sites

number of sites affected

affinity
decreased
increased

B T1D locus
rs3024505

C CEL locus
rs13010713

Unstim

8hr

24hr

IL19 *FAIM3*

ITGA4

206,750 206,850 206,950 207,050 207,150 Kb
chr1

181.0 181.5 182.0 182.5 183.0 Mbp
chr2

D MZF1
rs3024505 (*FAIM3/IL19*)

E SOX4
rs13010713 (*ITGA4*)

F rs1077667 (*GPR108/TRIP10*)
TP53
TP63
POU2F2

Figure 8 - Figure Supplement 2