The bloodstream form of *Trypanosoma brucei* displays non-canonical gluconeogenesis

Julie Kovářová[1,*], Martin Moos[2], Michael P. Barrett[3], David Horn[4], Alena Zíková[1]

[1] Institute of Parasitology, Biology Centre of the Czech Academy of Sciences, České

Budějovice, Czech Republic

[2] Institute of Entomology, Biology Centre of the Czech Academy of Sciences, České

Budějovice, Czech Republic

[3] Wellcome Centre for Integrative Parasitology, University of Glasgow, Glasgow, United

Kingdom

[4] Wellcome Centre for Anti-Infectives Research, University of Dundee, Dundee, United

Kingdom

* Corresponding author: julie.kovarova@paru.cas.cz

<u>Abstract</u>

Trypanosoma brucei is a causative agent of the Human and Animal African Trypanosomiases.

The mammalian stage parasites infect various tissues and organs including the bloodstream,

central nervous system, skin, adipose tissue and lungs. They rely on ATP produced in

glycolysis, consuming large amounts of glucose, which is readily available in the mammalian

host. In addition to glucose, glycerol can also be used as a source of carbon and ATP and as a

substrate for gluconeogenesis. However, the physiological relevance of glycerol-fed

gluconeogenesis for the mammalian-infective life cycle forms remains elusive. To demonstrate

its (in)dispensability, first we must identify the enzyme(s) of the pathway. Loss of the

canonical gluconeogenic enzyme, fructose-1,6-bisphosphatase, does not abolish the process

hence at least one other enzyme must participate in gluconeogenesis in trypanosomes. Using a

combination of CRISPR/Cas9 gene editing and RNA interference, we generated mutants for four enzymes potentially capable of contributing to gluconeogenesis: fructose-1,6-bisphoshatase, sedoheptulose-1,7-bisphosphatase, phosphofructokinase and transaldolase, alone or in various combinations. Metabolomic analyses revealed that flux through gluconeogenesis was maintained irrespective of which of these genes were lost. Our data render unlikely a previously hypothesised role of a reverse phosphofructokinase reaction in gluconeogenesis and preclude the participation of a novel biochemical pathway involving transaldolase in the process. The sustained metabolic flux in gluconeogenesis in our mutants, including a triple-null strain, indicates the presence of a unique enzyme participating in gluconeogenesis. Additionally, the data provide new insights into gluconeogenesis and the pentose phosphate pathway, and improve the current understanding of carbon metabolism of the mammalian-infective stages of *T. brucei*.

Author Summary

Trypanosoma brucei is a unicellular parasite causing sleeping sickness in humans and nagana disease in cattle. The parasite invades the bloodstream and cerebrospinal fluid and only recently, it has been shown to infect additional tissues such as skin, adipose tissue, or lungs. While the glucose-based metabolism of the bloodstream form is well understood, the parasite's metabolism in these secondary tissues has not been sufficiently explored, despite its importance for drug development. One possibility is the use of gluconeogenesis since the mammalian-infective stages can use glycerol as a carbon and ATP source. First, enzymes involved in gluconeogenesis have to be identified, then it can be tested if the pathway is advantageous for the survival of the parasite. We generated mutants in four different enzymes potentially involved in this metabolic pathway. Surprisingly, the flux in gluconeogenesis was maintained

in all cell lines tested, implying that another non-canonical enzyme participates in the

production of glucose from glycerol in these parasites.

Introduction

Trypanosoma brucei brucei is the causative agent of Human African Trypanosomiasis, also

termed sleeping sickness [1]. The mammalian-infective stage of the parasite is called the

bloodstream form (BSF), and it is transmitted between hosts by blood-feeding tsetse flies. In

the first stage of the disease, these extracellular parasites divide in the bloodstream of the

mammalian host. If left untreated, the trypanosomes invade the central nervous system,

manifesting as the second stage of the disease. BSF parasites can also inhabit skin, adipose

tissue, lungs and other tissues [1-5]. Previous dogma stated that BSF trypanosomes are

absolutely glucose-dependent. However, we [6], and others [7] have shown that they also

employ gluconeogenesis (GNG) and can use glycerol as a carbon source. Glycerol utilisation

is expected to be most physiologically relevant to parasites that inhabit adipose tissue or skin,

but the significance of glycerol as a carbon source remains elusive. Nevertheless, it is clear that

the parasite's metabolism is highly flexible and adaptable, and may differ between the various

mammalian forms, which should be considered when developing drugs inhibiting metabolic

enzymes. Notably, the adipose tissue forms are less responsive to several trypanocidal drugs

[8].

Under standard culture conditions in medium containing glucose, glycolysis provides the

majority of cellular ATP and is indispensable to BSF trypanosomes [9]. However, in culture

medium containing glycerol, BSF trypanosomes use GNG to produce sugars from non-sugar

carbon sources by converting glycerol to glucose 6-phosphate (G6P) [6, 7]. GNG primarily

uses the same enzymes as glycolysis, but operating in the opposite direction. The key difference

between the two pathways is the enzymatic step between fructose 6-phosphate (F6P), and fructose 1,6-bisphosphate (F1,6bP) when phosphofructokinase (PFK) phosphorylates F6P to F1,6bP in glycolysis, while fructose-1,6-bisphosphatase (FBPase) dephosphorylates F1,6bP to form F6P in GNG [10] (Fig 1A). In addition, since glycerol is used as the non-sugar substrate, glycerol kinase (GK) becomes a key GNG enzyme [7].

Previously, we demonstrated the presence of GNG in the BSF parasites by RNA interference (RNAi) silencing of the glucose transporters, where the observed lethal phenotype was rescued by the addition of glycerol to the culture media [6]. As verified by targeted LC-MS metabolomics, glycerol was incorporated into fructose 6-phosphate (F6P) and other metabolites via GNG. The same conclusion was reached by Pineda and colleagues after adapting BSF parasites to glucose-free, glycerol-containing medium [7]. Surprisingly, however, GNG was not abolished after deletion of the *FBPase* gene indicating an involvement of another, so far unknown, enzyme [6, 7].

The role of FBPase in GNG was also studied in the insect procyclic form (PCF) of *T. brucei* [11]. This stage has a more elaborate mitochondrion in terms of both morphology and metabolism, which can provide additional substrates for GNG. Hence, proline is utilised by PCF as a carbon source, via proline degradation in the mitochondrion, and fed into GNG via phosphoenolpyruvate carboxykinase and pyruvate phosphate dikinase [11]. FBPase knock-out (KO) PCF cells are viable when grown in medium containing proline and they incorporate proline-derived metabolites into G6P by GNG. This suggests that, similarly to BSF cells, the activity of FBPase can be compensated by another unknown enzyme. Interestingly, the FBPase KO PCF cells display a mild growth defect, defects in metacyclogenesis and transmission through the tsetse fly [11].

The presence of another enzyme catalyzing the F1,6bP to F6P conversion is intriguing, as in the majority of eukaryotic cells studied to date, this metabolic reaction is performed solely by the activity of FBPase. Therefore, the balance between FBPase and PFK activity is strictly and tightly regulated [10]. FBPase is positively regulated by ATP, and negatively by AMP and fructose 2,6-bisphosphate (F2,6bP). PFK is controlled by the same metabolites in a reciprocal manner, i.e. it is positively regulated by AMP, ADP, and F2,6bP. F2,6bP is an activator of most eukaryotic PFK enzymes, normally produced from F1,6bP by the bifunctional enzyme, 6-phosphofructo-2-kinase/fructose-2,6-bisphosphatase [12]. The finely tuned regulatory mechanism between PFK and FBPase ensures mutual exclusivity of these two reactions, and prevents futile cycling (a situation when both reactions would be running in a cycle, only consuming ATP). Interestingly, *T. brucei* PFK is much less sensitive to inhibition by ATP than many eukaryotic PFKs, and its activity is not affected by the presence of F2,6bP [13]. Therefore, the well-established PFK regulatory mechanism appears to be missing from *T. brucei* [14]. Regulation of *T. brucei* FBPase has not yet been elucidated.

Although *T. brucei* PFK is an ATP-dependent enzyme, its amino acid sequence is a 'chimera' between representative pyrophosphate- and ATP-dependent enzymes [15]. PFK isolated from *T. brucei* is regulated by AMP, which serves as the only allosteric activator, although 10-fold less potent than for the leishmania enzyme [13, 15], and by phosphoenolpyruvate acting as an allosteric inhibitor. F1,6bP or F2,6bP did not significantly influence enzyme activity [13]. As predicted by a mathematical model, and experimentally validated, PFK is present in excess in *T. brucei* and therefore, glycolytic flux is reduced only after depletion of PFK beyond 60 % of WT activity. It is unclear, however, whether the effect is direct, or a secondary consequence of decreased activity of other glycolytic enzymes (hexokinase, enolase, pyruvate kinase) [9].

Highly specific allosteric inhibitors of *T. brucei* PFK have been developed, and shown to be

effective in killing BSF parasites *in vitro* and in infected mice [16]. Fernandes and colleagues

[17] reported reversed activity of PFK *in vitro*, but the physiological relevance is unclear as

reversed activity of three mammalian PFK isoforms *in vitro* is not thought to occur *in vivo* [18].

In trypanosomatids, both glycolytic and gluconeogenic enzymes are localised within

specialized organelles called glycosomes. Glycosomes are highly adapted peroxisomes, and

they share related PEX-dependent protein import machineries [19]. These specialized

organelles also harbour parts of the pentose phosphate pathway (PPP), nucleotide metabolism,

and other associated pathways. The arrangement and function of the PPP is not fully resolved

in *T.brucei* BSF because transketolase is not expressed in this stage, but transaldolase is,

resulting in an incomplete non-oxidative branch of the PPP (Fig 1A) [20, 21]. The glycosomal

membrane is 'semi-permeable', thought to allow free passage of small molecules (up to 340

Da based on a mathematical model of trypanosome metabolism) through open channels, but

to be impermeable to larger molecules [22, 23]. Hence, cofactors such as ADP/ATP or

NAD^+/NADH are balanced inside glycosomes, and glycosomal localisation of the glycolytic

enzymes has important implications for their regulation [24].

We and others have shown previously that deletion of the *FBPase* gene does not eliminate the

enzymatic activity responsible for dephosphorylation of F1,6bP [6, 7], indicating that other

enzymes capable of this activity are also present. Notably, a gene encoding sedoheptulose-1,7-

bisphosphatase (SBPase), an enzyme typically involved in the Calvin cycle of plants, is present

in the genome of *T. brucei*. Due to a relatively high sequence identity (26 %) to FBPase, we

considered it a possible candidate for possessing FBPase activity. We also explored two other

scenarios in which the reverse PFK activity [17] or a pathway involving transaldolase (TAL)

[25] could bypass FBPase in GNG. To our surprise, double knock-out cell lines for FBPase

and SBPase (*Δfbp.sbp*), triple knock out cell lines for FBPas, SBPase and TAL (*Δfbp.sbp.tal*)

and PFK RNAi cell lines in a background of *Δfbp.sbp,* retained GNG flux, suggesting an

unknown enzyme performing the FBPase reaction. Nevertheless, we present several unique

metabolomic datasets that provide new insights into non-canonical GNG and PPP in BSF

trypanosomes.

Materials and Methods

T. brucei cell culture and cell line construction

Bloodstream form *Trypanosoma brucei brucei* Lister 427 cells were cultured in HMI-11

medium [26] supplemented with 10 % FCS (BioSera) at 37° C and 5 % CO_2. The fructose-1,6-

bisphosphatase and sedoheptulose-1,7-bisphosphatase knock-out cell line (*Δfbp.sbp*) was

generated by CRISPR/Cas9 gene editing in the 2T1$^{T7-Cas9}$ cell line [27]. In order to delete *FBP*

(Tb927.9.8720), a template for sgRNA transcription was synthesised by end-filling PCR with

primers FBPgRNA_F and sgRNA_R, and a repair template for DNA integration was

synthesised with primers FBP.NPT50_5 and FBP.NPT50_3. Cas9 expression was induced

with tetracycline at 1 µg/ml (InvivoGen) 24 h prior to transfection with an Amaxa nucleofector

(Lonza) using programme Z-001. First, the *Δfbp* cell line was generated, and after validation

of successful replacement of both *FBP* alleles with a G418 resistance cassette, a subsequent

transfection was performed to replace the *SBP* (Tb927.2.5800) alleles with a puromycin

resistance cassette (primer pairs SBPgRNA_F and sgRNA_R for sgRNA, and SBP.PAC50_5

and SBP.PAC50_3 for the repair template with puromycin resistance). Likewise, Δ*tal* (TAL

is encoded by Tb927.8.5600) was generated in the parental and *Δfbp.sbp* cell lines, using primer

pairs AZ1550 and sgRNA_R for sgRNA, and AZ1551 and AZ1552 for the repair template with

phleomycin resistance. The primers used are listed in Table 1. Hygromycin was used at 5 µg/ml

(InvivoGen), blasticidin at 5 µg/ml (InvivoGen), G418 at 2.5 µg/ml (Invivogen), and

puromycin at 0.1 µg/ml (InvivoGen).

For depletion of PFK (Tb927.3.3270), the RNAi plasmid p2T7-177 [28] was used, digested

with *Bam*HI and *Hind*III. A *PFK* PCR product was amplified using primer pair AZ1339 and

AZ1340, digested with the same restriction enzymes and ligated with the plasmid. The resulting

p2T7-177-PFK construct was linearized with *Not*I prior to electroporation of 2T1$^{\text{T7-Cas9}}$ and

Δfbp.sbp cell lines in parallel. Clones obtained after phleomycin selection (at 2.5 μg/ml) were

tested for a growth defect after tetracycline induction at 1 μg/ml, and validated by qRT-PCR.

The $^{\text{Ty}}$PFK cell line was also created by Cas9 editing in the 2T1$^{\text{T7-Cas9}}$ cell line. Following 24 h

of Cas9 induction, electroporation was performed with a sgRNA template (primer pair AZ1388

and sgRNA_R), and PURO.$^{\text{Ty}}$PFK cassette (primer pair AZ1281 and AZ1282). Clones were

selected with puromycin at 0.1 μg/ml and integration validated by PCR (primer pair AZ1403

and AZ1404) and western blot.

Table 1: Sequences of DNA primers used in the study.

Primer	Sequence	Description
FBPgRNA_F	TAATACGACTCACTATAGGGAGGGTGTGGTGCTTTTCGTGGGGCGTTTTAGAGCTAGAAATAGCAAG	FBP F sgRNA
sgRNA_R	GCACCGACTCGGTGCCACTTTTTCAAGTTGATAACGGACTAGCCTTATTTTAACTTGCTATTTCTAGCTCTAAAAC	R sgRNA universal
FBP.NPT 50_5	ttttgcgactgtttttcaatttcattaacgacaccactcttcccagatttATGATTGAACAAGATGGATTGC	FBP.G418 cassette F
FBP.NPT 50_3	TTTTAAAAAAAAACCTGTACCCTTTCCACACGCATCGAAGCAACCATTGGCTCAGAAGAACTCGTCAAGAAGG	FBP.G418 cassette R
SBPgRNA_F	TAATACGACTCACTATAGGGTGTGGTGCTTTTCGTGGGGCGTTTTAGAGCTAGAAATAGCAAG	SBP F sgRNA
SBP.PAC 50_5	gcccgttacaggggttccctttcactcaacctttcgtggaaagggagaatATGACCGAGTACAAGCCC	SBP.puro cassette F
SBP.PAC 50_3	gatcggaccatcaccttgtgcggcacaaaaataaaaacaagaagaaaaaaGGTACCGAGCTCGAATTCTC	SBP.puro cassette R
AZ1211	GATGGCGATTCAAACGTCCG	*FBP* ORF 177 – 981 bp
AZ1212	TTATCCCCCGGGTAGCAGAA	*FBP* ORF 177 – 981 bp
AZ1213	ATTGGCTTCTACCGCGCATA	*SBP* ORF 438 – 1,050 bp
AZ1214	GTTCGTCAACCCCGTTGTTG	*SBP* ORF 438 – 1,050 bp
AZ1339	ATGGATCCTTCACTTCAACCCGACGGAG	PFK RNAi F, *PFK* 338 – 894 bp
AZ1340	ATAAGCTTCTGTCACGACCCATGAGCTT	PFK RNAi R, *PFK* 338 – 894 bp
AZ1388	TAATACGACTCACTATAGGGGTGACGAGCTTGCTCGTAACGGTTTTAGAGCTAGAAATAGCAAG	Ty.PFK sgRNA
AZ1281	GTGAACTGGAGGCAATCAACAGGAATCGCGACCGCCTCCACGAGGAACTGGAGGTCCATACTAACC	PURO.$^{\text{Ty}}$PFK cassette F

	AAGATCCACTTGACGCCAAGCTCTAATCTCCGCT CTTATTTAGTTTTGC	
AZ1282	GTGCCCTCCTCATTCTCCCTTTCCCAGAACT CCGTTACCAGAGGGTCATTCAATCATGTCG ACACACCAAG	PURO.^{Ty}PFK cassette R
AZ1403	GGACGTATTGCATGTGCTGTC	validation of ^{Ty}PFK F
AZ1404	TGAGTCACGCTGTTCAGCAT	validation of ^{Ty}PFK R
AZ1550	TAATACGACTCACTATAGGGCGCTACACTT TTGTAAGGGAGTTTTAGAGCTAGAAATAGC AAG	TAL sgRNA
AZ1551	CTTAGAAGGGGGAAGGCAACAAGACATGG CCAAGTTGACCAGTG	TAL phleo cassette F
AZ1552	GTTACTTGATGGTGGGTACCCTCCCTCAGTC CTGCTCCTCGG	Tal phleo cassette R
AZ1609	ATGGATCCGTCTCTCGGCTACGTGATGG	validation of Δ*tal*
AZ1610	ATAAGCTTTCGTTACAGCAATGACGCCT	validation of Δ*tal*

qRT-PCR

Total RNA was extracted from $1 - 2 \times 10^8$ cells using the RNeasy kit (Qiagen). DNA was

removed using Turbo DNase (Applichem) at 37° C for 30 min, which was subsequently treated

with DNase inactivation reagent (Ambion) for 5 min at RT. Following ethanol precipitation,

cDNA was synthesised from 2 μg of RNA using TaqMan Reverse Transcription Reagent

(Applied Biosystems) and random hexamer primers. Real-time PCR amplification was

performed using LightCycler 480 SYBR Green I Master (Roche) and LightCycler 480

thermocycler (Roche). Primers used for the PFK target and reference genes are listed in

Table 2.

Table 2: Sequences of DNA primers used for qRT-PCR.

Target gene	Primer name	Sequence
PFK F	AZ1489	CCTCACGGAGAAAGTGAAGG
PFK R	AZ1490	GGGTAGCGAGACTTGTTTGC
18S F	AZ50	GCGAAACGCCAAGCTAATAC
18S R	AZ51	AGCCGCGACATAGAAAAAGA
Tubulin F	AZ52	GCAGAGTCCAACATGAACGA
Tubulin R	AZ53	CGTCCGCGTCTAGTATTGCT

<u>Subcellular fractionation and western blotting</u>

In order to separate organellar fractions containing glycosomes from the cytosol, cells were subjected to digitonin-based subcellular fractionation and the obtained samples used for western blots. Briefly, 1×10^8 cells were harvested, washed in 1 x PBS and resuspended in 500 μl of SoTE buffer (0.6 M sorbitol, 2 mM EDTA, 20 mM Tris-HCl, pH 7.5). A further 500 μl of SoTE buffer containing 0.03 % digitonin (Sigma-Aldrich) was added, samples were incubated on ice for 5 min and subsequently centrifuged at 4,500 g, 4° C for 3 min. The obtained supernatant was used as cytosolic fraction, and pellets resuspended in an equivalent volume of 1 x PBS and used as organellar fractions. 40 μl of samples were loaded for western blots.

For whole cell lysates, the equivalent of 1×10^7 cells was used. For western blots (WB) 4-12 % NuPAGE polyacrylamide gels (Invitrogen) and 1 x SDS running buffer (25 mM Tris, 192 mM glycine, 1 % SDS) were used. Subsequently, proteins were transferred to a PVDF membrane (Pierce) in transfer buffer (39 mM glycine, 48 mM Tris, 20 % methanol) at 90 V for 90 min at 4 °C. Following 30 min blocking in 5 % milk (Serva) in PBS-Tween (0.05 %), primary antibody was incubated in milk solution overnight at 4 °C. Following 3 x 10 min wash in PBS-Tween, secondary antibody was incubated in milk solution for 1 h at RT. Following 3 x wash in PBS-T, signal was visualised using Western ECL Substrate (BioRad). The following antibodies were used: α-FBP at 1:500 , α-SBP at 1:500 (both kind gifts from Frederick Bringaud), anti-Ty 1:1,000 (ThermoFisher), α-APRT at 1:500 was used as a marker for cytosolic fraction, and α-hexokinase at 1:2,000 as an organellar marker. Secondary α-mouse (BioRad) and α-rabbit (BioRad) antibodies conjugated to HRP were used at 1:2,000.

 Metabolomics

For the experiment with *Δfbp.sbp* in [13]C-glycerol, cells were grown in the standard HMI-11

medium supplemented with 5 mM [13]C-U-glycerol (Cambridge Isotope Laboratories). For the

experiment with *Δfbp.sbp*/[RNAi]PFK, Δ*tal,* and Δ*fbp.sbp.tal*, HMI-11 medium was prepared from

components according to the recipe (Hirumi), but glucose was omitted and instead 5 mM [13]C_3-

U-glycerol (Cambridge Isotope Laboratories) was supplied. Cells were grown in this medium

for 24 h (PFK cell lines) or 48 h (TAL cell lines) prior to sample extraction.

Samples for the metabolomic experiments were prepared by the same extraction protocol as

reported previously [29]. Briefly, 5 x 10^7 cells were used per 100 μl sample, which were first

rapidly cooled to 4° C in a dry ice – ethanol bath. Following a wash with 1 x PBS, cell pellets

were resuspended in 100 μl of chloroform:methanol:water (1:3:1) suspension and incubated

with shaking at 4° C for 1 h in order to achieve full extraction into the solvent. Subsequently,

samples were centrifuged (12,000 g, 10 min, 4° C), supernatants were collected and stored at -

80° C until analysis.

The metabolomic methods used were described in detail elswere [30]. Briefly, an Orbitrap Q

Exactive Plus mass spectrometer coupled to an LC Dionex Ultimate 3000 (Thermo Fisher

Scientific, San Jose, CA, USA) was used for metabolite profiling. LC condition: column

SeQuant ZIC-pHILIC 150 mm x 4.6 mm i.d., 5 μm, (Merck KGaA, Darmstadt, Germany);

flow rate of 450 μl/min; injection volume of 5 μl; column temperature of 35°C; mobile phase

A = acetonitrile and B = 20 mmol/l aqueous ammonium carbonate (pH = 9.2; adjusted with

NH4OH); gradient: 0 min, 20 % B; 20 min, 80 % B; 20.1 min, 95 % B; 23.3 min, 95 % B; 23.4

min, 20 % B; 30.0 min 20 % B. The Q-Exactive settings were: mass range 70-1050 Daltons;

70 000 resolution; electrospray ion source operated in the positive and negative modes.

The analysis of the Δ*fbp.sbp* cell line in HMI-11 medium was performed at Glasgow

Polyomics, using separation on 150 x 4.6 mm ZIC-pHILIC (Merck) on a Dionex UltiMate

3000 RSLC system (Thermo Scientific) followed by mass detection on Orbitrap QExactive

(Thermo Fisher Scientific) mass spectrometer (Thermo Fisher). Analysis was operated in

polarity switching mode, using 10 μl injection volume and a flow rate of 300 μl/min.

The analyses was performed in four replicates, and a set of standards was run in parallel.

Metabolite identification was based on matches with standards where possible or otherwise

predicted based on mass and retention time. Data were analysed using mzMatch [31] and

mzMatch.ISO [32], Xcalibur™ software, version 4.0 (Thermo Fisher Scientific, San Jose, CA,

USA), and an in-house developed Metabolite Mapper® platform.

FBPase assay

2×10^7 cells were used per sample. They were centrifuged (1,300 g, 10 min), washed with 1 x

PBS and resuspended in 100 μl of SoTE buffer (10 mM Tris-HCl pH 8, 1 mM EDTA, 0.15 %

Triton X-100, protease inhibitor cocktail (Roche)). After 20 min incubation at RT, samples

were centrifuged (14,000 g, 10 min, 16° C) and supernatant collected. The reaction mixture

containing 20 mM Tris pH 7.8, 10 mM $MgCl_2$, 1 mM NADP, 1 U PGI (Sigma-Aldrich), 1 U

G6PDH, 100 μl cell extract was incubated at 30° C for 5 min prior to activity measurement.

The reaction was triggered by addition of 5 mM F1,6bP (Sigma-Aldrich) immediately prior to

measurement of NADPH production at 340 nm for 5 min at 30° C using a UV-1601

spectrophotometer (Shimadzu).

Immunofluorescence assay (IFA)

For the immunofluorescence assay, cells were fixed in 7.4 % formaldehyde in 1 x PBS for 15

min, and subsequently washed three times with 1 x PBS. For permeabilisation, 0.1 % Triton

X-100 (AppliChem) in 1 x PBS was applied for 10 min, and subsequently washed 3 x with 1 x

PBS. Following blocking in 5.5 % FBS in 1 x PBS-Tween (0.05 % Tween) and 2 x wash with

1 x PBS, the primary antibodies were applied (α-Ty at 1:100 (ThermoFisher) and α-FBP at

1:1,000) for 1 h at RT. Following three washes with 1 x PBS-T and two washes with 1 x PBS,

the secondary antibodies were applied (Alexa Fluor 647 α-mouse (Life Technologies) at

1:2,000 and Alexa Fluor 488 α-rabbit (Life Technologies) at 1:2,000) for 1 h at RT. After an

additional three washes with 1 x PBS-T and two washes with PBS, ProLong Gold Antifade

mounting solution (Invitrogen) was applied. Imaging was performed using an Axioplan

microscope (Zeiss).

Results

<u>Generation of double knock-out *Δfbp.sbp* by two step Cas9 editing</u>

We took advantage of CRISPR/Cas9-based gene editing [27] to generate a double knock-out

of FBPase and SBPase (*Δfbp.sbp*) in *T. brucei* BSF. First, an sgRNA template and a repair

template were simultaneously electroporated in a transient transfection, resulting in

replacement of both alleles of the *FBPase* gene with a G418 resistance cassette. Subsequently,

applying the same approach, both alleles of *SBPase* were replaced with a puromycin cassette

(Fig. 1B). Full replacement of all alleles was validated by PCR and western blotting (Fig 1C,

D). Such a straightforward replacement of both target genes indicated that FBPase and SBPase

most likely do not play essential roles under the growth conditions used. Accordingly, the

Δfbp.sbp cell line showed no growth defect in the standard nutrient-rich HMI-11 medium (Fig

1E). When Creek's Minimal Medium (CMM), which better approximates the composition of

blood serum, was used for cell culture, *Δfbp.sbp* cells suffered a mild growth defect with higher

variability, possibly due to different metabolic adaptations required for growth in CMM (Fig

1E) [33]. Because these experiments showed a growth defect under nutrient-restrictive

conditions, we decided to assess infectivity *in vivo.* Infection of mice showed that *Δfbp.sbp*

cells were infectious, although they reached significantly lower parasitaemia on day 2

compared to the parental cell line (0.6-fold, $p = 0.04$ (S1 Fig)).

Figure 1. Generation and growth analysis of Δ*fbp.sbp* strains.

A – The scheme shows glycolysis, gluconeogenesis, and the pentose phosphate pathway in BSF *T. brucei* with highlighted FBPase, PFK, and TAL. Missing reactions of transketolase are depicted by grey arrows.

B – A scheme for the Cas9 editing method by transient transfection. The templates for an sgRNA and for an antibiotic resistance cassette were transfected simultaneously, which resulted in replacement of both alleles for *FBPase* in the first step, and for *SBPase* in the second step.

C – Validation of the Δ*fbp.sbp* cell line by PCR, shows parts of ORFs for *FBPase* and *SBPase*

amplified in the parental 2T1$^{T7.Cas9}$ cell line, but absent from Δ*fbp.sbp*.

D - Validation of the Δ*fbp.sbp* cell line by western blot. Only the parental cell line shows signal

for FBPase (whole cell lysates) and SBPase (organellar fractions), * - cross-reacting protein,

gel loading – fluorescent protein detection on a TGX gel.

E – Growth curves of two independent clones of Δ*fbp.sbp* show no defect in the standard HMI-

11 medium. Growth curves in the CMM medium show a mild growth defect of the Δ*fbp.sbp*

clones and higher variability.

<u>Metabolomic analysis of *Δfbp.sbp* shows sugar phosphate substrate accumulation and</u>

<u>maintained gluconeogenesis</u>

We next examined the effects of the *fbp* and *sbp* double gene deletion on central carbon

metabolism. First, a metabolomic analysis was performed on *Δfbp.sbp* cells cultured in CMM.

The most striking change was approx. 30-fold accumulation of S1,7bP (p = 3 x 10^{-6}), the

substrate of SBPase (Fig 2A, B), confirming disruption of the SBPase reaction. Sedoheptulose

7-phosphate (S7P) was 4-fold increased (p = 10^{-5}), probably due to non-enzymatic conversion

of highly accumulated S1,7bP. The metabolites of the FBPase reaction were also affected,

although to a much lesser extent than for SBPase, with F6P increased 1.3-fold (p = 0.03), and

F1,6bP not changed significantly (Fig 2B). Since F6P and F1,6bP are also involved in other

metabolic pathways, especially glycolysis, absence of FBPase does not have such a substantial

impact on their abundance.

Figure 2. Gluconeogenic activity is maintained in Δ*fbp.sbp* strains.

A – Changes in Δ*fbp.sbp* cells grown in CMM and subjected to non-targeted LC-MS

metabolomics when compared to the parental cell line. The most changed metabolites are

S7P, S1,7bP, and phosphothreonine. The plot includes 87 metabolites from glycolysis, PPP,

TCA cycle or amino acid metabolism.

B – Sedoheptulose 1,7-bisphosphate (S1,7bP), sedoheptulose 7-phosphate (S7P), fructose

1,6-bisphosphate (F1,6bP), and fructose 6-phosphate (F6P) as detected in the parental and

Δ*fbp.sbp* cells grown in CMM and subjected to non-targeted LC-MS metabolomics.

C - Δ*fbp.sbp* cells were grown in HMI-11 supplemented with U-^{13}C$_3$-glycerol and analysed by

LC-MS metabolomics, showing that ^{13}C from glycerol is incorporated into GNG products.

H6P – hexose 6-phosphates.

To test whether GNG flux is present, we grew the parental and *Δfbp.sbp* cells in HMI-11

medium supplemented with 5 mM U-^{13}C$_3$-glycerol and subjected the samples to metabolomic

analysis. Overall, few changes were observed in the *Δfbp.sbp* metabolome compared to the

parental cell line, with a 2-fold accumulation of S7P being most evident in *Δfbp.sbp* cells grown

with glycerol (Fig 2C). Since glucose was also present in the medium, the predominant portion

of the hexose phosphate isotopomer pool was unlabelled in both cell lines, however, $^{13}C_3$

labelling represented 10 % of the hexose phosphate pool. 12 % of F1,6bP contained the $^{13}C_3$

label, providing clear evidence that these labelled carbons originate from U-$^{13}C_3$-glycerol fed

through GNG, in both parental and *Δfbp.sbp* cell lines (Fig 2C).

Changes in the metabolome of *Δfbp.sbp* cells grown in CMM (Fig 2B) correspond with the

growth defect observed in that medium and reflect higher sensitivity of cells to perturbations

in nutrient-restricted conditions, in contrast to the HMI-11 medium, where minimal changes in

growth and the metabolome of *Δfbp.sbp* cells were observed (Fig 1E, 2C). Overall, the

differences detected in sugar phosphates correlate with expectations for FBPase and SBPase

deletion, except for maintainance of gluconeogenic flux. This maintained flux was also

confirmed by conversion of F1,6bP to F6P in cell extracts; comparable levels of enzymatic

activity were measured using a specific enzyme activity assay from the parental cell line (15

µmol/min/mg, n = 3) and *Δfbp.sbp* (16 µmol/min/mg, n = 3).

<u>Depletion of PFK reduces glycolysis but not GNG</u>

Since we did not observe changes in GNG flux in *Δfbp.sbp* cell lines, we next determined

whether reversed activity of PFK (dephosphorylation of F1,6bP into F6P) can contribute to

GNG. PFK was depleted by RNAi in the parental cell line (PFK^{RNAi}) and in the *Δfbp.sbp*

background (*Δfbp.sbp*/PFK^{RNAi}). Induction of knockdown using tetracycline caused a severe

growth defect in both cell lines, as expected due to the essential role of PFK in glycolysis (Fig

3A). RT-PCR confirmed a drop in target gene expression to 20 % in the parental background

and 60 % in the *Δfbp.sbp* background (insets, Fig 3A). We performed metabolomic analysis of

the PFK^{RNAi} and *Δfbp.sbp*/PFK^{RNAi} cells fed with U-$^{13}C_3$-glycerol as the sole carbon source (no

glucose was present in the HMI-11 medium, except for ~0.5 mM glucose from 10 % FBS).

Samples were harvested 24 h after tetracycline induction in the glycerol medium and subjected

to LC-MS metabolomics. The profile of glycerol 3-phosphate (G3P) shows that glycerol was

taken up and incorporated, since the $^{13}C_3$-labelled part represents around 80 % (78 - 85 %) of

the total in all sample groups (Fig 3B). Crucially, RNAi induced cells with decreased PFK

expression still utilised glycerol via GNG and incorporated it into hexose phosphate sugars.

The profile of F6P shows that the majority (44 – 66 %) is $^{13}C_6$-fructose 6-phosphate, made

from $^{13}C_3$-glycerol, in all samples (Fig 3B). The major difference after PFK depletion is in the

unlabelled part that originates from glucose via glycolysis, and relies on the forward reaction

of PFK. This fraction decreased from 18 to 8 % of the total amount in PFK[RNAi] induced

samples, confirming a reduction in PFK glycolytic activity. On the other hand, the proportion

of the $^{13}C_6$-labelled part increased from 44 to 66 %, demonstrating undiminished glycerol

incorporation. A substantial part of F6P is $^{13}C_3$-labelled (26 – 38 %), indicating that both

reactions, phosphorylation of F6P into F1,6P and the reverse dephosphorylation, take place.

Analysis of the F1,6bP profile shows an even higher proportion of $^{13}C_6$-labelled, representing

64 – 74 % across the samples. Smaller changes occur after induction of PFK knockdown, but

the same trend as in F6P is observed. The unlabelled part dropped from 6.5 to 2.5 %, $^{13}C_3$

decreased from 29 to 22 %, and $^{13}C_6$ increased from 65 to 75 % (Fig 3B). In the

Δfbp.sbp/[RNAi]PFK cell line, no significant changes were detected after induction of PFK

silencing, suggesting that any further perturbation of metabolic flux is lethal for these cells.

Figure 3. PFK knockdown causes a severe growth defect, but only mild changes in the metabolome.

A – Growth curves of PFK^RNAi and in *Δfbp.sbp*/^RNAiPFK show a severe growth defect after RNAi. Levels of PFK mRNA are shown as detected by qRT-PCR, compared to non-induced cells, and normalised to 18S, 24 h tetracycline induction (insets).

B – Targeted LC-MS metabolomics in HMI-11 medium depleted of glucose, and supplemented with $^{13}C_3$-glycerol. The proportion of the C_{13} label incorporation relative to the total amount is indicated. G3P – glycerol 3-phosphate, F6P – fructose 6-phosphate, F1,6bP – fructose 1,6-bisphosphate. 2T1 ^{13}C – parental cell line in medium with ^{13}C-glycerol,

PFK^RNAi – non-induced PFK RNAi cell lines in medium with ^{13}C-glycerol,

PFK^RNAi + Tet – PFK RNAi induced for 24 h in medium with ^{13}C-glycerol,

Δfbp.sbp/PFK – non-induced *Δfbp.sbp*/^RNAiPFK cell line in medium with ^{13}C-glycerol,

Δfbp.sbp/PFK + Tet - *Δfbp.sbp*/^RNAiPFK cell line induced for 24 h in medium with ^{13}C-glycerol

C – Same as in B), but relative changes in the total levels of metabolites are depicted. P5P – pentose 5-phosphates, O8P – octulose 8-phosphate, S7P – sedoheptulose 7-phosphate.

Pentose phosphate pathway (PPP) intermediates are labelled to a lesser extent than the hexose

phosphates. Around 70 % of the total pentose phosphate pool is unlabelled, while $^{13}C_5$

represents between 11 to 18 %, and $^{13}C_3$ about 5 % (Fig 3C). Octulose 8-phosphate (O8P) has

a similar labelling pattern, suggesting its synthesis from pentose phosphates condensed with a

three carbon moiety ($^{13}C_3$ represents ~ 65 %, Fig 3C). S7P, another intermediate of the non-

oxidative PPP, is substantially labelled. The $^{13}C_3$ part comprised around 20 % of total S7P in

the WT control and PFK^{RNAi}, and 28 % in $\Delta fbp.sbp/^{RNAi}$PFK cells, regardless of whether

knockdown was induced or not, suggesting that PFK depletion has little impact on the

incorporation of glycerol. However, depletion of FBPase and SBPase caused an increase in

$^{13}C_3$, in addition to the increase in total amount of S7P more than 4-fold (compared to WT)

(Fig 3C), consistent with the previous measurement (Fig 2B). Moreover, the $^{13}C_7$ part

represented about 30 % of total S7P in $\Delta fbp.sbp$ cells, confirming glycerol as a substrate for

the production of S7P. These larger sugar phosphates have been observed in *T. brucei* before

[29], and although their role in metabolism remains unclear, the current targeted metabolomic

data offers new insights into their biosynthesis.

Metabolomic profiles of other metabolites, such as intermediates of glycolysis, the succinate

shunt, or mitochondrial carbon metabolism, provide additional valuable information.

Substantial $^{13}C_3$ labelling was observed in phosphoenolpyruvate (PEP), pyruvate, fumarate,

malate, aspartate, and alanine (Fig 3C, S2 Fig). 72 – 78 % of PEP is $^{13}C_3$, demonstrating a high

level of utilisation of labelled glycerol in glycolysis. The unlabelled fraction dropped from 22

% to 13 % after PFK knockdown, again confirming a drop in glucose utilisation in glycolysis,

relative to glycerol. Pyruvate comprised 28 – 36 % of unlabelled, and 60 – 68 % of $^{13}C_3$-

pyruvate. Malate comprised 43 % to 57 % of unlabelled, 10 – 16 % of $^{13}C_2$ (synthesized in the

mitochondrion via acetyl-CoA), and 28 % - 36 % of $^{13}C_3$ (synthesized in the glycosomal

succinate shunt), without significant differences between the sample groups (S2 Fig). For

aspartate, the $^{13}C_3$ content ranges from 42 % to 50 % in all cell samples. These data indicated

that glycerol was used in glycolysis for pyruvate and acetyl-CoA production and further fed

into mitochondrial enzyme reactions classically associated with the TCA cycle (S2 Fig). The

data are consistent with our previous work, where we observed similar labelling patterns with

glycerol, indicating an increase in mitochondrial metabolism when BSF cells rely on GNG [6].

Overall, few changes were observed after PFK knockdown, despite the severe growth defect.

However, we saw clear evidence for diminished glycolysis, but not GNG. The data

demonstrated that glycerol was utilised as the main carbon source under these conditions, since

the majority of glycolytic intermediates were labelled from $^{13}C_3$-glycerol. In addition to use as

a substrate for GNG, glycerol was used for synthesis of S7P, which may reflect a novel

variation of the non-oxidative PPP.

PFK and FBPase co-localise in glycosomes

We next sought to establish the localisation of the key glycolytic and GNG enzymes, PFK and

FBPase. Both of these enzymes are glycosomal, but their mutual co-localisation has never been

reported. For this purpose, one of the endogenous PFK alleles was tagged in the 2T1[T7-Cas9] cell

line with the Ty-tag, generating [Ty]PFK, which was used for microscopy in a combination with

a specific α-FBPase antibody. We cultured the cells in medium supplemented with glucose or

glycerol only, or in a combination of both carbon sources, and then subjected these cells to

super-resolution confocal microscopy (Fig 4). PFK and FBPase always co-localised in

glycosomes and no differences were observed under the various conditions tested. The two

enzymes are not separated by compartmentalisation. Further work will be required to reveal

potential regulation, although futile cycling does occur to some extent as seen in the

metabolomic data (Fig 3B).

Figure 4. PFK and FBPase co-localise in glycosomes.

A – PCR validation of ^{Ty}PFK clones.

B – Western blot validation of the ^{Ty}PFK cell line.

C – Immunofluorescence assay with ^{Ty}PFK cell line, α-Ty, and α-FBPase staining. Cells were

cultured in CMM with different carbon sources (glucose, glycerol, or both), but it had no effect

on co-localisation of PFK and FBPase.

<u>TAL does not contribute to gluconeogenesis</u>

One possible explanation for the observed GNG activity (i.e. generation of F6P from glycerol)

may be a novel metabolic pathway involving three enzymes (SBPase, TAL, and fructose-1,6-

bisphosphate aldolase) that operate in a cycle, using glycolytic intermediates as substrates to

produce F6P (Fig 5A) [25]. Considering the unprecedented presence of SBPase, expression of

TAL in the absence of transketolase, and the persistence of GNG flux, such a novel pathway

could offer an explanation. Hence, we decided to test the role of TAL in GNG, and generated

a knock-out cell line lacking *FBPase, SBPase*, and *TAL (Δfbp.sbp.tal)*. *TAL* was deleted using

Cas9 and replaced with a phleomycin resistance cassette in the parental 2T1^{T7-Cas9} and in the

Δfbp.sbp cell lines (Fig 5B). The *Δfbp.sbp.tal* cells displayed no growth defect in HMI-11

medium supplemented either with glucose or glycerol as the main carbon source (Fig 5C). The

cells were also still infective to mice at levels comparable to the parental cell line (S3 Fig).

A targeted LC-MS metabolomic analysis was performed on *Δfbp.sbp.tal* cells grown in CMM

supplemented with 5 mM $^{13}C_3$-glycerol and no glucose (~0.5 mM glucose was present due to

supplementation with 10 % FCS). The results showed substantial uptake of, and reliance upon,

glycerol as the major carbon source. The profile of G6P is surprisingly similar between the

parental and *Δfbp.sbp.tal* cells when fed with $^{13}C_3$-glycerol, with the majority of the G6P

metabolite being $^{13}C_6$-labelled (62 % in WT and 49 % in *Δfbp.sbp.tal*) (Fig 5D). Even more

substantial is the fully labelled part in F1,6bP, representing 84 % of total in WT and 78 % in

Δfbp.sbp.tal. Additionally, the total amount of the G6P metabolite is 2-fold increased in the

triple knock-out (Fig 5D). S7P is a sugar phosphate formed by an unknown mechanism,

however, our data show that S7P is synthesized from glycerol. The total amount of S7P is

increased 3-fold in WT when fed on glycerol relative to glucose, indicating that glycerol

increases S7P synthesis. In addition, S7P is increased 2-fold in *Δtal* and 6-fold in the

Δfbp.sbp.tal cell line. The majority is $^{13}C_3$- and $^{13}C_7$-labelled (28 % and 49 % respectively, in

Δfbp.sbp.tal, Fig 5D). In summary, our metabolomic data show persistence of GNG flux from

glycerol even in the *Δfbp.sbp.tal* cell line. We did not observe other major changes in the

metabolome associated with deletion of TAL, thus the function of TAL in BSF trypanosomes,

where its usual PPP partner enzyme transketolase is absent, remains unknown.

Figure 5. Deletion of TAL does not decrease GNG flux.

A - The novel pathway as suggested by Hannaert [25]: Erythose 4-phosphate (E4P) is condensated with dihydroxyacetone phosphate (DHAP) by fructose-1,6-bisphosphate aldolase into S1,7bP, which is dephosphorylated by SBPase into S7P. That is used as a substrate by transaldolase, together with glyceraldehyde 3-phosphate (GA3P), and converted into F6P and E4P, which can enter another cycle.

B – PCR detection of a 500-bp product from the *TAL* ORF, proving deletion of the gene.

C – Growth curves of the parental (2T1[T7-Cas9]), *Δtal,* and *Δfbp.sbp.tal* cell lines in HMI-11 medium supplemented with glucose, or glycerol only (and 10 % FBS).

D - Targeted LC-MS metabolomics in CMM supplemented with 5 mM $^{13}C_3$-glycerol. G6P – glucose 6-phosphate, F1,6bP - fructose 1,6-bisphosphate, S7P - sedoheptulose 7-phosphate. 2T1 – parental cell line in medium with glucose, 2T1 ^{13}C – parental cell line in medium with ^{13}C-glycerol, *Δtal*^{13}C – *Δtal* cell line in medium with ^{13}C-glycerol, *Δfbp.sbp.tal* - *Δfbp.sbp.tal* cell line in medium with glucose, *Δfbp.sbp.tal*^{13}C - *Δfbp.sbp.tal* cell line in medium with ^{13}C-glycerol.

Discussion

We took advantage of Cas9-based gene editing to delete multiple genes and in an attempt to abolish GNG in BSF *T. brucei*. However, our data show that the combined deletion of FBPase, SBPase, and TAL had little impact on gluconeogenic activity. To the best of our knowledge, this is the first case of deletion of three diploid gene loci simultaneously in BSF *T. brucei*. Depletion of PFK was possible only to a limited extent, but our data support the view that this enzyme is not involved in GNG, as suggested previously [17]. Current understanding of trypanosome metabolism does not provide an alternative explanation for how GNG operates, and which enzymes contribute to GNG flux.

In our previous work and that of others, it was demonstrated that deletion of the canonical *FBPase* gene in *T. brucei* does not deplete FBPase activity, i.e. conversion of F1,6bP to F6P [6, 7]. This is in contrast to closely related leishmania parasites, where *FBPase* deletion disrupted GNG and caused a severe phenotype in mammalian infective amastigotes [34]. Since SBPase is not present in the genome of leishmania, this enzyme was a promising candidate for the observed FBPase activity. The enzyme has 26 % sequence identity to FBPase, and its catalytic activity is predicted to be very similar, using a sugar phosphate backbone extended by one carbon. Additionally, SBPase from yeast was demonstrated to posess 'FBPase' activity [35]. The most probable origin of SBPase in trypanosomes is acquisition by horizontal gene transfer [36].

The streamlined protocol for Cas9-based gene editing allowed us to generate a double gene knock-out combined with a knock-down. Knock-out for both *FBPase* and *SBPase* genes were readily obtained by sequential transfections. The *Δfbp.sbp* cell line had no growth defect under nutrient-rich culture conditions. However, there was a mild and variable growth defect in

CMM. The high variability in growth rate could be caused by different mechanisms of

adaptation to nutrient-restricted conditions. The metabolomic experiments with *Δfbp.sbp*

revealed that SBPase bears its canonical enzymatic activity (converting S1,7bP to S7P), since

its substrate, S1,7bP was highly accumulated in the knockout cell line. A concomitant increase

in S7P is most likely explained by non-enzymatic loss of a single phosphate from the

accumulated S1,7bP. Similarly, accumulation of S1P and S1,7bP was observed previously in

an SBPase yeast deletion mutant [37]. SBPase was also identified in *Toxoplasma gondii*, and

its deletion resulted in a similar phenotype to what we observed here, both in metabolomics

and in decreased infectivity [38].

Fernandes and colleagues [17] showed reverse PFK activity (equal to canonical FBPase) *in*

vitro. Our data do not support reverse activity of trypanosome PFK *in vivo*, however, since we

observed a decrease in glycolytic flux but not GNG flux following PFK knockdown. We

detected a severe growth defect when the PFK mRNA was depleted to 20 % in parental and to

60 % in the *Δfbp.sbp* background. More efficient PFK knockdown by RNAi is unlikely to be

achievable in the *Δfbp.sbp* double knock-out mutant, because metabolic flux is flexible and

enzymes can compensate for each other to some extent, which is impossible in the double

knock-out background. Metabolomics showed surprisingly few changes in the levels of

detected metabolites in the PFK knockdown cell lines. Nevertheless, we noted a decrease in

glycolytic flux, but not in GNG flux. The directionality of the PFK activity is dependent on

concentrations of substrates and products. Fernandes *et al.* calculated that the reversal should

occur if the concentration of substrates is 500-fold higher than the concentration of products

[17]. The changes in ADP, ATP, F6P, and F1,6bP in our metabolomic datasets were much less

prominent, for instance one of the largest changes is the 2-fold accumulation of F1,6bP in

Δfbp.sbp/[RNAi]PFK compared with WT (S3 Fig). However, we measured metabolites extracted

from whole cells, whereas the ADP/ATP ratio within the glycosomal sub-compartment may be

substantially shifted relative to that of the whole cell. Altogether, although it is challenging to

test involvement of an essential glycolytic enzyme, our data do not support participation of

PFK on GNG.

The PPP intermediates detected by metabolomics can be separated into two groups. The first

represents the oxidative PPP and metabolites derived originally from glucose, where less

labelling from glycerol was detected (pentose phosphates, Fig 3C). The second group,

representing the non-oxidative PPP, contains S7P or O8P, which have significant proportions

labelled (Fig 3C, 5D). Since BSF *T. brucei* does not express transketolase [20, 21], the

metabolites cannot be produced in the canonical non-oxidative PPP. This suggests that in a low

concentration of glucose, as used here, glucose is used preferentially to feed the oxidative PPP,

whereas glycerol is used to feed glycolysis and to synthesize large sugar phosphates such as

S7P. Although O8P has previously been shown to be synthesized *in vitro* from F6P and R5P

by TAL [29], there appears to be a separate route for synthesis in the parasites, as evidenced

by the presence of this metabolite in *TAL*-depleted cell lines. In both WT and *Δfbp.sbp.tal*,

much higher quantities of O8P were detected in the presence of glucose than in the same cell

lines grown in glycerol-based medium (S4 Fig). Our datasets provide new insights into the role

of TAL and the whole PPP in BSF, however, further studies will be required to resolve the

arrangement of the pathway in this life stage.

A novel metabolic pathway that includes the activities of SBPase, TAL, and fructose-1,6-

bisphosphate aldolase was proposed [25]. If active, this pathway conforms to the labelling

patterns detected in S7P (and O8P), and also with S1,7bP production catalysed by aldolase, as

reported previously [37]. Most importantly, this would present an alternative pathway for the

production of F6P from glycerol. Nevertheless, our experiments with the Δ*tal* and Δ*fbp.sbp.tal*

mutants show that carbons from glycerol are still incorporated into F1,6bP, F6P, and other

metabolites, demonstrating continued flux though GNG, and precluding the existence of the

alternative pathway.

It is also unclear how the flux between glycolysis and GNG, i.e. between PFK and FBPase

activity, is regulated. Potentially, the forward or reverse flux could be controlled by upstream

kinases, i.e. hexokinase in glycolysis and glycerol kinase in GNG, due to competition for ATP,

as reported recently [39]. Another possibility would be an exclusive compartmentalisation of

these enzymes, but our immunofluorescence analysis shows that PFK and FBPase co-localise

under all conditions tested. These results suggest that some futile cycling does take place, as

substantial $^{13}C_3$-labelling of hexose phosphates (Fig 3B) occurs. However, since FBPase is not

a key GNG enzyme, we suggest that regulation is achieved by a distinct enzyme that possesses

FBPase activity but whose identity is yet to be established.

When glucose is limiting, bloodstream form trypanosomes adapt by activating alternative

metabolic pathways. Labelling patterns in succinate and citrate indicate flux in mitochondrial

metabolism. Since only a small proportion of labelling in citrate was detected (72 – 89 % of

citrate was unlabelled), incorporation of additional substrates, such as glutamine or other amino

acids likely occurs, as reported previously [40]. If glycerol is available to trypanosomes, it is

used as a carbon source, being incorporated into sugar phosphates under all of conditions tested

here. Deletion or depletion of four different enzymes failed to substantially diminish GNG flux

in BSF trypanosomes. Taken together, our results indicate that (an)other enzyme(s), currently

unrecognized by *in silico* searchers, is/are responsible for the activity. The question as to

whether GNG is essential for BSF *T. brucei* cannot be readily addressed until these key

enzyme(s) are identified.

<u>Acknowledgements</u>

We thank Prof. Frederic Bringaud (University of Bordeaux) and Prof. Paul Michels (University

of Edinburgh) for the kind gift of antibodies α-FBPase, α-SBPase, and α-PFK. This work was

supported by CZ.02.2.69/0.0/0.0/19_074/0016248 LeishWeb to J.K.; by the European

Regional Development Fund (ERDF) and Ministry of Education, Youth and Sport (MEYS)

project CZ.02.1.01/0.0/0.0/16_019/0000759, by the European Research Council (ERC,

MitoSignal, grant agreement no. 101044951), and GACR 20-14409S to A.Z.; D. H. was funded

by an Investigator Award from Wellcome [217105/Z/19/Z]. M.P.B was funded by an MRC

Newton grant 'Bridging epigenetics, metabolism and cell cycle in pathogenic

trypanosomatids' MR/S019650/1.

Supporting Figures

S1 Figure: Mice infection with *Δfbp.sbp* cell line.

S2 Figure. Targeted LC-MS metabolomics in HMI-11 medium depleted of glucose, but

supplemented with $^{13}C_3$-glycerol. 2T1 ^{13}C – parental cell line in medium with ^{13}C-glycerol,

PFK^{RNAi} – non-induced PFK RNAi cell lines in medium with ^{13}C-glycerol,

PFK^{RNAi} + Tet – PFK RNAi induced for 24 h in medium with ^{13}C-glycerol,

Δfbp.sbp/PFK – non-induced *Δfbp.sbp*/^{RNAi}PFK cell line in medium with ^{13}C-glycerol,

Δfbp.sbp/PFK + Tet - *Δfbp.sbp*/^{RNAi}PFK cell line induced for 24 h in medium with ^{13}C-glycerol.

664   S3 Figure. Mice infection with *Δfbp.sbp.tal* cell line.

675   S4 Figure. Octulose 8-phosphate (O8P) as detected in the *Δtal* cell lines by LC-MS

676   metabolomics. [13]C indicates medium supplemented with [13]C-glycerol as the only carbon

677   source.

References

1.   Crilly NP, Mugnier MR. Thinking outside the blood: Perspectives on tissue-resident

Trypanosoma   brucei.   PLoS   Pathog.   2021;17(9):e1009866.   Epub   2021/09/17.   doi:

10.1371/journal.ppat.1009866.   PubMed   PMID: 34529724;   PubMed   Central   PMCID:

PMCPMC8445408.

2.   Caljon G, Van Reet N, De Trez C, Vermeersch M, Pérez-Morga D, Van Den Abbeele J. The

Dermis as a Delivery Site of Trypanosoma brucei for Tsetse Flies. PLoS Pathog. 2016;12(7):e1005744.

Epub 2016/07/22. doi: 10.1371/journal.ppat.1005744. PubMed PMID: 27441553; PubMed Central

PMCID: PMCPMC4956260.

3.   Capewell P, Cren-Travaillé C, Marchesi F, Johnston P, Clucas C, Benson RA, et al. The skin is a

significant but overlooked anatomical reservoir for vector-borne African trypanosomes. Elife. 2016;5.

Epub 2016/09/23. doi: 10.7554/eLife.17716. PubMed PMID: 27653219; PubMed Central PMCID:

PMCPMC5065312.

4.   Trindade S, Rijo-Ferreira F, Carvalho T, Pinto-Neves D, Guegan F, Aresta-Branco F, et al.

Trypanosoma brucei Parasites Occupy and Functionally Adapt to the Adipose Tissue in Mice. Cell Host

Microbe. 2016;19(6):837-48. Epub 2016/05/31. doi: 10.1016/j.chom.2016.05.002. PubMed PMID:

27237364; PubMed Central PMCID: PMCPMC4906371.

5.   Mabille D, Dirkx L, Thys S, Vermeersch M, Montenye D, Govaerts M, et al. Impact of pulmonary

African trypanosomes on the immunology and function of the lung. Nat Commun. 2022;13(1):7083.

Epub 2022/11/19. doi: 10.1038/s41467-022-34757-w. PubMed PMID: 36400767; PubMed Central

PMCID: PMCPMC9674601.

6.   Kovářová J, Nagar R, Faria J, Ferguson MAJ, Barrett MP, Horn D. Gluconeogenesis using

glycerol   as   a   substrate   in   bloodstream-form   Trypanosoma   brucei.   PLoS   Pathog.

2018;14(12):e1007475. Epub 2018/12/28. doi: 10.1371/journal.ppat.1007475. PubMed PMID:

30589893; PubMed Central PMCID: PMCPMC6307712.

7. Pineda E, Thonnus M, Mazet M, Mourier A, Cahoreau E, Kulyk H, et al. Glycerol supports

growth of the Trypanosoma brucei bloodstream forms in the absence of glucose: Analysis of metabolic

adaptations on glycerol-rich conditions. PLoS Pathog. 2018;14(11):e1007412. Epub 2018/11/02. doi:

10.1371/journal.ppat.1007412. PubMed PMID: 30383867; PubMed Central PMCID:

PMCPMC6245841.

8. Trindade S, De Niz M, Costa-Sequeira M, Bizarra-Rebelo T, Bento F, Dejung M, et al. Slow

growing behavior in African trypanosomes during adipose tissue colonization. Nat Commun.

2022;13(1):7548. Epub 2022/12/09. doi: 10.1038/s41467-022-34622-w. PubMed PMID: 36481558;

PubMed Central PMCID: PMCPMC9732351.

9. Albert MA, Haanstra JR, Hannaert V, Van Roy J, Opperdoes FR, Bakker BM, et al. Experimental

and in silico analyses of glycolytic flux control in bloodstream form Trypanosoma brucei. J Biol Chem.

2005;280(31):28306-15. Epub 2005/06/16. doi: 10.1074/jbc.M502403200. PubMed PMID: 15955817.

10. M C Scrutton a, Utter MF. The Regulation of Glycolysis and Gluconeogenesis in Animal Tissues.

Annual Review of Biochemistry. 1968;37(1):249-302. doi: 10.1146/annurev.bi.37.070168.001341.

11. Wargnies M, Bertiaux E, Cahoreau E, Ziebart N, Crouzols A, Morand P, et al. Gluconeogenesis

is essential for trypanosome development in the tsetse fly vector. PLoS Pathog.

2018;14(12):e1007502. Epub 2018/12/18. doi: 10.1371/journal.ppat.1007502. PubMed PMID:

30557412; PubMed Central PMCID: PMCPMC6312356.

12. Rider MH, Bertrand L, Vertommen D, Michels PA, Rousseau GG, Hue L. 6-phosphofructo-2-

kinase/fructose-2,6-bisphosphatase: head-to-head with a bifunctional enzyme that controls

glycolysis. Biochem J. 2004;381(Pt 3):561-79. Epub 2004/06/02. doi: 10.1042/bj20040752. PubMed

PMID: 15170386; PubMed Central PMCID: PMCPMC1133864.

13. Cronin CN, Tipton KF. Purification and regulatory properties of phosphofructokinase from

Trypanosoma (Trypanozoon) brucei brucei. Biochem J. 1985;227(1):113-24. Epub 1985/04/01. doi:

10.1042/bj2270113. PubMed PMID: 3158309; PubMed Central PMCID: PMCPMC1144815.

14.   Chevalier N, Bertrand L, Rider MH, Opperdoes FR, Rigden DJ, Michels PA. 6-Phosphofructo-2-

kinase and fructose-2,6-bisphosphatase in Trypanosomatidae. Molecular characterization, database

searches, modelling studies and evolutionary analysis. Febs j. 2005;272(14):3542-60. Epub

2005/07/13. doi: 10.1111/j.1742-4658.2005.04774.x. PubMed PMID: 16008555.

15.   McNae IW, Martinez-Oyanedel J, Keillor JW, Michels PA, Fothergill-Gilmore LA, Walkinshaw

MD. The crystal structure of ATP-bound phosphofructokinase from Trypanosoma brucei reveals

conformational transitions different from those of other phosphofructokinases. J Mol Biol.

2009;385(5):1519-33. Epub 2008/12/17. doi: 10.1016/j.jmb.2008.11.047. PubMed PMID: 19084537.

16.   McNae IW, Kinkead J, Malik D, Yen LH, Walker MK, Swain C, et al. Fast acting allosteric

phosphofructokinase inhibitors block trypanosome glycolysis and cure acute African trypanosomiasis

in mice. Nat Commun. 2021;12(1):1052. Epub 2021/02/18. doi: 10.1038/s41467-021-21273-6.

PubMed PMID: 33594070; PubMed Central PMCID: PMCPMC7887271.

17.   Fernandes PM, Kinkead J, McNae IW, Bringaud F, Michels PAM, Walkinshaw MD. The kinetic

characteristics of human and trypanosomatid phosphofructokinases for the reverse reaction. Biochem

J. 2019;476(2):179-91. Epub 2018/11/09. doi: 10.1042/bcj20180635. PubMed PMID: 30404924;

PubMed Central PMCID: PMCPMC6340114.

18.   Hofmann E. The significance of phosphofructokinase to the regulation of carbohydrate

metabolism. Rev Physiol Biochem Pharmacol. 1976;75:1-68. Epub 1976/01/01. doi:

10.1007/BFb0030484. PubMed PMID: 181810.

19.   Michels PAM, Gualdrón-López M. Biogenesis and metabolic homeostasis of trypanosomatid

glycosomes: New insights and new questions. J Eukaryot Microbiol. 2022;69(6):e12897. Epub

2022/02/18. doi: 10.1111/jeu.12897. PubMed PMID: 35175680.

20.   Stoffel SA, Alibu VP, Hubert J, Ebikeme C, Portais JC, Bringaud F, et al. Transketolase in

Trypanosoma brucei. Mol Biochem Parasitol. 2011;179(1):1-7. Epub 2011/05/17. doi:

10.1016/j.molbiopara.2011.04.006. PubMed PMID: 21570429.

21.   Cronín CN, Nolan DP, Voorheis HP. The enzymes of the classical pentose phosphate pathway

display differential activities in procyclic and bloodstream forms of Trypanosoma brucei. FEBS Lett.

1989;244(1):26-30. Epub 1989/02/13. doi: 10.1016/0014-5793(89)81154-8. PubMed PMID: 2924907.

22.   Gualdrón-López M, Brennand A, Avilán L, Michels PA. Translocation of solutes and proteins

across the glycosomal membrane of trypanosomes; possibilities and limitations for targeting with

trypanocidal   drugs.   Parasitology.   2013;140(1):1-20.   Epub   2012/08/24.   doi:

10.1017/s0031182012001278. PubMed PMID: 22914253.

23.   Achcar F, Barrett MP, Breitling R. Explicit consideration of topological and parameter

uncertainty gives new insights into a well-established model of glycolysis. Febs j. 2013;280(18):4640-

51. Epub 2013/07/20. doi: 10.1111/febs.12436. PubMed PMID: 23865459; PubMed Central PMCID:

PMCPMC4768353.

24.   Bakker BM, Westerhoff HV, Michels PA. Regulation and control of compartmentalized

glycolysis in bloodstream form Trypanosoma brucei. J Bioenerg Biomembr. 1995;27(5):513-25. Epub

1995/10/01. doi: 10.1007/bf02110191. PubMed PMID: 8718456.

25.   Hannaert V, Bringaud F, Opperdoes FR, Michels PA. Evolution of energy metabolism and its

compartmentation in Kinetoplastida. Kinetoplastid Biol Dis. 2003;2(1):11. Epub 2003/11/14. doi:

10.1186/1475-9292-2-11. PubMed PMID: 14613499; PubMed Central PMCID: PMCPMC317351.

26.   Hirumi H, Doyle JJ, Hirumi K. Cultivation of bloodstream Trypanosoma brucei. Bull World

Health Organ. 1977;55(2-3):405-9. Epub 1977/01/01. PubMed PMID: 303956; PubMed Central

PMCID: PMCPMC2366732.

27.   Rico E, Jeacock L, Kovářová J, Horn D. Inducible high-efficiency CRISPR-Cas9-targeted gene

editing and precision base editing in African trypanosomes. Sci Rep. 2018;8(1):7960. Epub 2018/05/23.

doi: 10.1038/s41598-018-26303-w. PubMed PMID: 29785042; PubMed Central PMCID:

PMCPMC5962531.

28.   Wickstead B, Ersfeld K, Gull K. Targeting of a tetracycline-inducible expression system to the

transcriptionally silent minichromosomes of Trypanosoma brucei. Mol Biochem Parasitol. 2002;125(1-

2):211-6. Epub 2002/12/07. doi: 10.1016/s0166-6851(02)00238-4. PubMed PMID: 12467990.

29.   Creek DJ, Mazet M, Achcar F, Anderson J, Kim DH, Kamour R, et al. Probing the metabolic

network in bloodstream-form Trypanosoma brucei using untargeted metabolomics with stable

isotope labelled glucose. PLoS Pathog. 2015;11(3):e1004689. Epub 2015/03/17. doi:

10.1371/journal.ppat.1004689. PubMed PMID: 25775470; PubMed Central PMCID:

PMCPMC4361558.

30.   Moos M, Korbelová J, Štětina T, Opekar S, Šimek P, Grgac R, et al. Cryoprotective Metabolites

Are Sourced from Both External Diet and Internal Macromolecular Reserves during Metabolic

Reprogramming for Freeze Tolerance in Drosophilid Fly, Chymomyza costata. Metabolites. 2022;12(2).

Epub 2022/02/26. doi: 10.3390/metabo12020163. PubMed PMID: 35208237; PubMed Central

PMCID: PMCPMC8877510.

31.   Scheltema RA, Jankevics A, Jansen RC, Swertz MA, Breitling R. PeakML/mzMatch: a file format,

Java library, R library, and tool-chain for mass spectrometry data analysis. Anal Chem.

2011;83(7):2786-93. Epub 2011/03/16. doi: 10.1021/ac2000994. PubMed PMID: 21401061.

32.   Chokkathukalam A, Jankevics A, Creek DJ, Achcar F, Barrett MP, Breitling R. mzMatch-ISO: an

R tool for the annotation and relative quantification of isotope-labelled mass spectrometry data.

Bioinformatics. 2013;29(2):281-3. Epub 2012/11/20. doi: 10.1093/bioinformatics/bts674. PubMed

PMID: 23162054; PubMed Central PMCID: PMCPMC3546800.

33.   Creek DJ, Nijagal B, Kim DH, Rojas F, Matthews KR, Barrett MP. Metabolomics guides rational

development of a simplified cell culture medium for drug screening against Trypanosoma brucei.

Antimicrob Agents Chemother. 2013;57(6):2768-79. Epub 2013/04/11. doi: 10.1128/aac.00044-13.

PubMed PMID: 23571546; PubMed Central PMCID: PMCPMC3716122.

34.   Naderer T, Ellis MA, Sernee MF, De Souza DP, Curtis J, Handman E, et al. Virulence of

Leishmania major in macrophages and mice requires the gluconeogenic enzyme fructose-1,6-

bisphosphatase. Proc Natl Acad Sci U S A. 2006;103(14):5502-7. Epub 2006/03/30. doi:

10.1073/pnas.0509196103. PubMed PMID: 16569701; PubMed Central PMCID: PMCPMC1459384.

35.   Kuznetsova E, Xu L, Singer A, Brown G, Dong A, Flick R, et al. Structure and activity of the metal-

independent fructose-1,6-bisphosphatase YK23 from Saccharomyces cerevisiae. J Biol Chem.

2010;285(27):21049-59. Epub 2010/04/30. doi: 10.1074/jbc.M110.118315. PubMed PMID: 20427268;

PubMed Central PMCID: PMCPMC2898295.

36.   Teich R, Zauner S, Baurain D, Brinkmann H, Petersen J. Origin and distribution of Calvin cycle

fructose and sedoheptulose bisphosphatases in plantae and complex algae: a single secondary origin

of complex red plastids and subsequent propagation via tertiary endosymbioses. Protist.

2007;158(3):263-76. Epub 2007/03/21. doi: 10.1016/j.protis.2006.12.004. PubMed PMID: 17368985.

37.   Clasquin MF, Melamud E, Singer A, Gooding JR, Xu X, Dong A, et al. Riboneogenesis in yeast.

Cell. 2011;145(6):969-80. Epub 2011/06/15. doi: 10.1016/j.cell.2011.05.022. PubMed PMID:

21663798; PubMed Central PMCID: PMCPMC3163394.

38.   Olson WJ, Martorelli Di Genova B, Gallego-Lopez G, Dawson AR, Stevenson D, Amador-Noguez

D, et al. Dual metabolomic profiling uncovers Toxoplasma manipulation of the host metabolome and

the discovery of a novel parasite metabolic capability. PLoS Pathog. 2020;16(4):e1008432. Epub

2020/04/08. doi: 10.1371/journal.ppat.1008432. PubMed PMID: 32255806; PubMed Central PMCID:

PMCPMC7164669.

39.   Allmann S, Wargnies M, Plazolles N, Cahoreau E, Biran M, Morand P, et al. Glycerol suppresses

glucose consumption in trypanosomes through metabolic contest. PLoS Biol. 2021;19(8):e3001359.

Epub 2021/08/14. doi: 10.1371/journal.pbio.3001359. PubMed PMID: 34388147; PubMed Central

PMCID: PMCPMC8386887.

40.   Johnston K, Kim DH, Kerkhoven EJ, Burchmore R, Barrett MP, Achcar F. Mapping the

metabolism of five amino acids in bloodstream form Trypanosoma brucei using U-(13)C-labelled

substrates and LC-MS. Biosci Rep. 2019;39(5). Epub 2019/04/28. doi: 10.1042/bsr20181601. PubMed

PMID: 31028136; PubMed Central PMCID: PMCPMC6522824.