Parallelized screening of virus accessory genes reveals diverse defense and counter-defense mechanisms

Sukrit Silas[1,10], Héloïse Carion[1], Kira S. Makarova[2], Eric Laderman[1], David Sanchez Godinez[3,4], Matthew Johnson[1], Andrea Fossati[5,6,7], Danielle Swaney[5,6,7], Michael Bocek[8], Eugene V. Koonin[2], Joseph Bondy-Denomy[1,6,9,10]

[1] Department of Microbiology and Immunology, University of California, San Francisco, San Francisco, CA 94158, USA

[2] National Center for Biotechnology Information, National Library of Medicine, National Institutes of Health, Bethesda, MD 20894, USA

[3] Cardiovascular Research Institute, University of California, San Francisco, San Francisco, CA 94158, USA

[4] Department of Biochemistry and Biophysics, University of California, San Francisco, San Francisco, CA 94158, USA

[5] J. David Gladstone Institutes, San Francisco, 94158, California, USA

[6] Quantitative Biosciences Institute, University of California, San Francisco, San Francisco, CA 94158, USA

[7] Department of Cellular and Molecular Pharmacology, University of California, San Francisco, San Francisco, CA 94158, USA

[8] Twist Biosciences, South San Francisco, CA 94080, USA

[9] Innovative Genomics Institute, Berkeley, CA 94720, USA

[10] Corresponding authors: Sukrit.Silas@ucsf.edu, Joseph.Bondy-Denomy@ucsf.edu

Abstract

Genomes of bacterial and archaeal viruses are replete with fast-evolving, uncharacterized accessory genes (AGs), most of which likely antagonize host defenses or other viruses. We developed a computational pipeline to find AGs in bacteriophage genomes and built a high-throughput screening platform to assay their functions. This approach is targeted towards identifying the most salient antiviral mechanisms in any bacterial niche. We tested 200 Enterobacteriophage AGs in 20 wild *Escherichia coli* strains challenged with 8 phages. The most prominent AG functions we observed were antagonism of O-antigen-based barrier defense and restriction-modification (R-M) systems. In response to phage-encoded anti-R-M strategies, some Type I and III R-M systems surprisingly act as programmed-cell-death modules. Moreover, several hyper-variable AGs trigger other abortive defense systems, demonstrating that phage-associated molecular patterns that activate immunity need not be well-conserved. Our approach yields insights into the multiple levels of virus-host competition and can be rapidly deployed in various bacteria.

Introduction

To a large extent, the evolution of life is a story of the virus-host arms race[1], during which the hosts evolved diverse, elaborate antiviral defense mechanisms, and viruses retorted by evolving equally versatile counter-defenses. Analysis of the rapidly growing genomic and especially metagenomic sequence databases uncovered a vast pool of proteins with mostly unknown functions encoded by viruses infecting bacteria and archaea[2]. The characterized examples of such proteins comprise only a small fraction of that pool, but possess diverse, exquisitely evolved capacities. These functionalities include creating novel interfaces in protein complexes[3], mimicking biomolecules to competitively inhibit enzymes[4], allosterically modifying cellular enzymes[5], interfering with host processes such as transcription[6], translation[7], adaptive immunity[8], motility[9], quorum sensing[10] and DNA transfer[11], inhibiting the reproduction of other viruses[12], and even participating in inter-kingdom competition[13]. The study of model phage and phage-defense systems over the past few decades has yielded key molecular biology tools, notably, recombinant DNA methodology that relies on restriction-modification (R-M) systems[14] and CRISPR genome editing[15]. A comprehensive investigation of the genes encoded by viruses can reveal an enormous variety of biological mechanisms involved in the virus-host arms race.

Bacterial and archaeal defense systems often cluster together in genomic loci known as defense islands[16]. Recently, a broad variety of distinct defense mechanisms have been discovered by investigating genes located in the same islands with known defense systems[17,18]. Counter-defense genes also often cluster together in virus genomes[19], but these "counter-defense islands" have not been studied as systematically as the defense islands in prokaryotic genomes. Because experiments with individual virus proteins can be slow and laborious, systematic approaches are essential to prioritize genes for study.

We developed an approach to identify bacteriophage genes that are likely to interfere with host biology. Core genes are present in (nearly) all members of a group of related phages and typically encode essential structural components of the virions as well as components of the replication and transcription machineries. In contrast, Accessory Genes (AGs) are present in some, but not all, closely-related genomes[20,21] and most are of unknown function, whereas those that have been characterized (such as numerous anti-CRISPR proteins) are often involved in counter-defense[22-24] or superinfection exclusion[11,12]. Our algorithm identifies "accessory regions" (ARs) that interrupt blocks of conserved core genes or are located at genome ends in closely related virus genomes (Figure 1a). AGs are typically non-essential, undergo frequent HGT, and are subject to enhanced evolutionary pressure due to their involvement in the virus-host arms race, resulting in extensive diversification.

To test AG functions, we developed the Phage Accessory Gene Exploration aNd Testing (PhAGENT) platform, which can experimentally evaluate thousands of phage genes at once. The platform utilizes genetic screening to rapidly identify AGs that affect host sensitivity to the phage or trigger programmed cell death (PCD). This yields host-phage-AG combinations that can then be interrogated using further screens, proteomics, and classical microbiology. AG libraries can be synthesized from any phage genomic dataset, and the PhAGENT platform can be deployed in any bacterial or archaeal strain collection where conjugation and transposon mutagenesis are supported. To validate the approach, we synthesized 200 phage AGs from diverse Enterobacteriophages and tested them in 20 wild strains of the *E. coli* Reference (ECOR) collection. We identified layered bacterial defenses and extensive co-evolution of defense and counter-defense mechanisms: a distinct mode of bacterial seroconversion by phage proteins that alter host metabolism, phage-encoded inhibitors of antiviral restriction mechanisms, an R-M system that can trigger PCD in addition to its canonical defense functions,

98 and decoy immunity in bacteria functionally similar to that found in plants[25] which can alert the
99 host to viral attempts to interfere with cellular pathways.
100
101 **Combined Results + Discussion**
102
A high throughput screening platform for identifying and assaying accessory genes
AGs are usually defined within a cluster of closely-related viruses[20-23] and AG
identification is highly sensitive to the method used to cluster virus genomes. We therefore
sought to identify ARs without *a priori* sequence comparison or clustering of virus genomes
(Supplementary File 1). First, all virus genes from the entire dataset were collected, and
pairwise combinations of genes (the query pair) were evaluated sequentially (Figure 1b). Hence,
every virus genome that contained both genes of the query pair was grouped into a possible
virus cluster, and genes encoded between the genes of the query pair were considered
members of a possible AR. Second, a series of filters were applied. Clusters containing few
viruses (<6) or highly diverged ones were discarded. The longest ARs (>10 genes) or those that
consisted of the same genes in every genome were also discarded. Third, all retained ARs were
scored and ranked by diversity (see Methods). Crucially, we did not rely on any gene ontology
to identify AGs (eg. genes similar to or encoded next to known counter-defense factors). A
sample AR is shown in Figure 1c. Redundant, low scoring ARs that comprised mostly of genes
which had already been found in higher scoring ARs were then discarded. Using an initial set of
1706 Enterobacteriophage genomes, 2014 non-redundant ARs containing 10888 putative AGs
were identified in 1217 phages. A typical AR contained 11-30 unique AGs ~50-100 aa in length.
A typical phage genome contained <5 ARs and <5 AGs (Figure 1d).
We manually inspected the highest-scoring 200 ARs and chose multiple (2-5) AGs from
>60 of these (see Supplementary File 2 for detailed notes on each AR). A barcoded library of
200 AGs was constructed (with 65 AGs from ARs containing known counter-defense factors,
and 135 AGs from ARs with all genes of unknown function) to enable multiplexed identification
of AGs that target bacterial defenses. In the subset of 65 AGs, 11 phage and mobile genetic
element (MGE)-encoded proteins with previously elucidated counter-defense or superinfection-
exclusion functions were included as controls (Abc2, ArdA, Arn, Cor, Dmd, Imm, Ip1, Ocr, Ral,
SieA, Stp), with the remaining 54 of unknown function (Supplementary File 3).
To select phage-host combinations, we used a subset of the *E. coli* Reference (ECOR)
strains[26] and assayed their ability to resist infection by 8 model phages. We reasoned that we
may find new antiviral mechanisms if the phage receptors were present on the host cell surface,
but some host factor(s) were blocking infection. Hence, 45 phage-host combinations with
attenuated infection were chosen, that is, those where a phage was able to produce clearings
on the bacterial lawn, but only with significantly reduced titer (Supplementary Figure 1a).
To identify AG function, we developed a scalable, high-throughput screen to
simultaneously assess how all AGs affect host immunity to infection (See Methods). Barcoded,
chromosomally integrated, IPTG-inducible AGs were expressed in each host (one AG per cell)
in pools, and fitness effects were measured in triplicate upon AG-induction and phage infection
by next-generation sequencing (Figure 1e) (Supplementary File 4). Importantly, although a set
of 8 model phages was used in our experiments, the AGs can be taken from any available
phage genome. AG induction was limited to 1-1.5 hrs, and subsequent infections were only
allowed to proceed for ~1 phage replication cycle, to prevent prolonged induction of various
AGs from biasing the results. This setup also avoids biases resulting from epigenetic
modification of phages by widespread restriction-modification (R-M) defenses in *E. coli*.
AGs that drastically reduced the fitness of some (but not all) hosts either upon
expression, or upon phage infection, were selected for further study.
147

AGs target conserved host pathways and barrier defenses

AGs target conserved host pathways and barrier defenses
To identify AGs with counter-defense phenotypes, we compared infected samples with
uninfected controls and found that several host-phage-AG combinations were depleted from the
infected pools, presumably because the respective AGs facilitate phage replication (Figure 2a).
Each AG that sensitized any wild strain to infection was then individually validated in that strain
against the entire phage set. In these validation experiments, some phage-host combinations
that were omitted in the PhAGENT screen also exhibited counter-defense phenotypes of AGs
(Supplementary Figure 1b). In all, increased plaquing of several phages was observed upon
expression of 3 known type I R-M inhibitors (Ocr, Ral, ArdA), 6 novel AGs, and the T4 internal
protein 2 (IpII) in a total of 28 phage-host combinations. We hypothesized that these AGs
antagonize host factors to help the phage(s) replicate.
Six novel AGs (orfs 48, 63, 92, 116, 126, 148), which have not been previously
implicated in phage-host interactions (Supplementary File 5), all exhibited counter-defense
phenotypes in multiple host strains. Because the broad-spectrum effects of these AGs did not
correlate with the defense systems in the hosts (Supplementary Figure 1c), we sought to
determine whether they could be targeting conserved host pathways that have antiviral
functions.
To identify the antiviral mechanisms inhibited by AGs, we employed two unbiased
approaches. (1) Transposon-knockout screens in the hosts helped pinpoint gene disruptions
that phenocopy AG expression (i.e. increased phage infection) (Supplementary File 6). (2)
Affinity-purification and mass spectrometry (AP-MS) of AGs helped identify their binding
partners in the wild strains (Supplementary File 7). We performed these assays for every host
found to be sensitized to infection by any AG.
Genome-wide knockout screens indicated that disrupting the host O-antigen or capsule
genes phenocopied the phage-sensitization effects of counter-defense AGs with almost all
tested phages (Table 1a). In particular, transposon-mediated O-antigen disruption (Figure 2b) or
expression of AGs 48, 63, and 92 (Figure 2c) both converged on the same phenotype, namely,
enhanced infection by phage T5 (which does not natively encode these AGs). We hypothesized
that T5 adsorption was blocked by the wild-type O-antigen structures in these hosts, but
heterologous AG expression enabled infection by modifying or removing the O-antigen. We
therefore examined LPS preparations from every host where any AG exhibited a counter-
defense phenotype. Hosts expressing AGs 48, 63, and 92 exhibited subtle downshifts in the O-
antigen banding pattern (Figure 2d), as observed previously in some seroconverting lysogens
as a means to exclude superinfecting phages[27]. Specifically, the following host-AG
combinations showed visible downshifts in O-antigen bands: orfs 63 and 92 in ECOR1, orfs 48
and 92 in ECOR3, and orfs 48 and 63 in ECOR22. These downshifts correlate with the host-AG
pairs where T5 infection was especially (10^{5-6}-fold) enhanced (Figure 2c). We reasoned that the
AGs might interfere with O-antigen biosynthesis, and tentatively named orfs 48, 63 and 92,
gnarl1, 2, and *3* respectively, after the skin-eating demon in the popular TV series *Buffy the*
Vampire Slayer.
Next, we examined the distribution of *gnarl* genes in the context of O-antigen
seroconversion. All known seroconverting prophages modify O-antigens using enzymes with
distinct biochemical activities (such as glycosyltransferases and acetyltransferases)[27-31]. *Gnarl-*
encoding phages in our genomic dataset do not encode any such seroconverting enzymes.
Gnarl proteins (40aa – 90aa) are also much smaller than typical enzymes, suggesting that these
AGs do not possess O-antigen-modifying enzymatic activity but instead affect O-antigen
biosynthesis. This potentially constitutes a distinct mode of host seroconversion. *Gnarl* genes
are not restricted to either lytic or lysogenic phages; *gnarl1* and *gnarl3* are present in lytic
phages (Supplementary File 5), whereas *gnarl2* is present in the classical temperate phage Mu
(gene E6/Mup07). Seroconversion could occur in lysogenic infections to exclude superinfecting

phages, or in lytic infections as receptor-masking to prevent newly synthesized virions from
unproductively binding lysed cell fragments[32].
To identify the cellular targets of gnarl proteins, we analyzed AP-MS data for all counter-
defense AGs. Each AG-encoded protein bound distinct host proteins although the profiles of
AGs 116 and 126 were somewhat correlated (Figure 2e). We performed Gene Ontology
pathway analysis for each co-IP to and found that UDP-glucose biosynthesis factors were
significantly enriched in the *gnarl3* set. Among these, GalU catalyzes the formation of UDP-
glucose[33] to build the LPS outer core[34]; GalF possibly regulates the levels of UDP-glucose[35];
and GalE reversibly converts UDP-glucose to UDP-galactose (also an LPS outer core
component)[36]. If *gnarl3* inhibits UDP-glucose biosynthesis, we reasoned that overexpressing the
target of *gnarl3* in the wild hosts might overcome its phage-sensitization phenotype. As a
control, we also overexpressed *wecB*, which was enriched in *gnarl3* AP-MS but is not involved
in UDP-glucose synthesis.
Overexpression of *galU* completely reversed the *gnarl3* T5-sensitization phenotype in all
tested hosts, namely, ECOR1, ECOR3, ECOR21 (not shown), and ECOR22 (Figure 2f),
whereas overexpression of *galE, galF,* and *wecB* had no effect. The same effect was observed
with unrelated phages T4 (Supplementary Figure 2) and P1 (not shown), indicating the *gnarl3-*
galU interaction was phage-agnostic. Overexpressing *galU* without *gnarl3* did not affect phage
sensitivity except in ECOR22, where it partially potentiated infection for unknown reasons. No
other genes we tested (*galE, galF,* wecB) affected phage-sensitization by *gnarl3*, suggesting a
specific interaction between *galU* and *gnarl3*. Coupled with LPS electrophoresis and AP-MS
data, this observation suggests a hypothetical model where *gnarl3* inhibits GalU, reducing the
availability of UDP-glucose and resulting in a modification of the cell envelope that potentiates
phage absorption. Although the possibility remains that the *gnarl3* phenotype is reversed
indirectly by *galU* overexpression (eg. through its role in galactose metabolism), the inability of
certain phages to adsorb to UDP-glucose pyrophosphorylase mutants of both *E. coli*[37] and *B.*
subtilis[38] suggests that glycosylation of cell-surface biopolymers is an ancient phage-blocking
strategy.
226
Barrier defenses and restriction systems provide layered immunity
The existence of broad-spectrum counter-defense AGs underscores the importance of
barrier defenses. Consistent with this, our genome-wide transposon screens identified an O121-
specific O-antigen[39] as a principal barrier to infection in ECOR21 (Figure 3a). While disrupting
O-antigen allowed other phages to infect this strain, we could not obtain corresponding
transposon insertions that sensitized ECOR21 to T2 (Figure 4a). Because the same ECOR21-
knockout library was used for transposon screens with every phage, we hypothesized that an
additional defense mechanism in ECOR21 restricted T2 infection (Figure 3b). In the PhAGENT
screen, we recovered a phenotype for a T4 internal-protein, IpII (orf143), that partially sensitized
ECOR21 to T2 (Figure 2a), suggesting that a T2-restricting defense system was disabled by
IpII. If this hypothetical defense system indeed complements barrier defenses, this would
explain why no single transposon insertion in ECOR21 seems to enable T2 infection.
To test this hypothesis, we eliminated O-antigen by deleting *wecA*[40] from ECOR21
(Figure 3c) and attempted the genome-wide knockout screen again. The screen now yielded
several insertions in an atypical BREX system (Figure 3d) in which a fused type IV restriction-
modification enzyme GmrSD (also known as BrxU) is embedded[41]. Thus, the O-antigen barrier
indeed obscured this defense locus.
Type IV R-M systems have co-evolved extensively with T-even phages. T-even phage
DNA contains hydroxymethylcytosines (hmC) that are further modified to glucosyl-hmC (g-
hmC), presumably to escape cleavage by type I R-M systems. Type IV R-M systems
(comprising two factors, GmrS and GmrD) can restrict g-hmC modified DNA, but T4 encodes an

accessory gene product, internal protein 1 (IpI), that is packaged into the phage capsid,
delivered to the host with the virus DNA, and disables type IV R-M. However, IpI cannot block
restriction by a modified type IV R-M system that combines both GmrS and GmrD activities in a
single effector GmrSD[32].
We wondered whether BREX or GmrSD was the target of IpII in ECOR21. We cloned
the entire operon along with its natural promoter and recapitulated its phenotype in *E. coli* K12.
The operon robustly protected against all tested T-even phages (T2, T4, T6). To isolate the
defense activity, we first deleted *pglX*, an essential component of BREX, and found that the
phages were still blocked. Next, we tested an active-site mutant of GmrSD (GmrSD*) and found
that infection was almost completely restored. Coupled with previous observations that type IV
R-M systems digest glycosylated T-even phage DNA whereas BREX systems do not[42], our
results indicate that GmrSD, and not BREX, restricted T-even phages (Figure 3e). Moreover,
overexpressing GmrSD alone was sufficient to block all T-even phages whereas a catalytic
mutant (GmrSD*) had no effect (Figure 3f).
Next, we tested whether various AGs (IpII, Ocr, IpI) might modify BREX/GmrSD function.
Ocr (a canonical R-M inhibitor) was recently demonstrated to have anti-BREX activity[43],
whereas IpI blocks the ancestral two-component type IV R-M system but not GmrSD. We found
that Ocr and IpI did not affect any configurations of the BREX/GmrSD system we tested.
However, expressing IpII restored infection by T-even phages despite the presence of
BREX/GmrSD, or the Δ*pglX* construct (Figure 3e), indicating that IpII inhibited GmrSD.
Of the many T-even phage internal-proteins, IpII is now only the second (after IpI) with
an identified target. T4 naturally encodes IpII, and was previously known to infect GmrSD-
containing hosts via an undiscovered mechanism[32]. Our results suggest that IpII provides this
activity. Consistent with this, over-expressed GmrSD targeted T2, T4, and T6, but the
uninduced leaky GmrSD construct protected against only T2 and T6 (Figure 3f). When IpI or IpII
were supplied, IpII restored T2 and T6 plaquing but did not further enhance T4 infection (Figure
3g). The observed redundancy between barrier defenses and restriction systems might obscure
phenotypes of other internal-proteins of T-even phages.
276
Restriction-modification systems can provide abortive defense in response to their
inhibitors
A second group of host-interacting AGs produced "conditional lethal" phenotypes, that is,
several AGs imposed a severe infection-independent fitness burden in a minority of strains
(Figure 4a). We hypothesized that these strains might harbor abortive defenses that detect
specific phage proteins and induce PCD in response to a perceived infection. Notably, several
of these trigger AGs were known anti-restriction factors (Supplementary Figure 3a). Putative
anti-restriction-induced abortive (ARIA) mechanisms might serve as a backup to thwart
phage/MGE-encoded counter-defense strategies.
We transformed the respective hosts with trigger AGs and confirmed their abortive
phenotypes (Figure 4b). We then performed genome-wide transposon suppressor screens to
find the genetic locus responsible for PCD. Transposon insertions in the abortive system would
be expected to inactivate PCD and allow survival of the bacteria despite expression of the
trigger AG. We identified putative ARIAs in all hosts except in ECOR66, where no suppressor
mutants (other than disruptions of the trigger AG) could be obtained (Table 1b, Supplementary
Figure 4, Supplementary File 6).
Wild strains encoding these ARIAs respond variously to R-M inhibitors, even
discriminating between the two DNA mimics, Ocr and ArdA, suggesting a diversity of
mechanisms (Supplementary Figure 3a). Notably, Ral, which operates by allosterically
hyperactivating the methyltransferase (unlike Ocr and ArdA)[5] did not trigger PCD in any strains.
Conversely, a novel AG (orf7) that triggers PCD by activating the same ARIAs triggered by Ocr

and ArdA (Supplementary Figure 3a) does not inhibit the *EcoKI* type I R-M system
(Supplementary Figure 3b). Further, the predicted structure of orf7 – hereafter, *aoa1* for
Activator Of ARIA – does not resemble a DNA mimic (Supplementary File 8), and it is also not
as negatively charged (pI ~4 for Ocr/ArdA, ~8 for *aoa1*). It thus seems possible for a canonical
R-M inhibitor to not trigger ARIAs, and for ARIAs to be triggered by R-M inhibitors as well as
other phage factors.
We were surprised to find that a type III R-M system in ECOR22 (hereafter, EcoR22III)
was required for cell-death in response to Ocr (Table 1b, Figure 4c), which blocks both type I
and type III R-M[44]. While R-M defenses are ubiquitous in bacteria (in particular, *E. coli*), so are
anti-restriction factors in phages. In response, bacteria evolved so-called anti-anti-restriction
mechanisms (such as retrons[45] and PARIS[46]) to serve as backup for their frontline defenses.
Our results suggest that in ECOR22, the frontline (R-M) and backup (PCD) defense functions
have coalesced into the same system. To test this explicitly, we reconstituted the PCD and R-M
activities of EcoR22III in a model *E. coli* strain.
In its native host, the *mod* and *res* genes of EcoR22III are associated with an SNF2-like
helicase found in DISARM systems, *drmD*[47], and all 3 genes are encoded in a P4 satellite
prophage hotspot[46]. However, transposon-suppressors of Ocr-toxicity in ECOR22 were found
only in *mod* and *res* (Figure 4c, Supplementary Figure 4). When overexpressed in a lab strain,
the 3-gene system did not produce a PCD response to Ocr (not shown). We then
overexpressed EcoR22III alone (only the *mod* and *res* genes) and found that EcoR22III was
now toxic upon Ocr induction but did not react to ArdA or Aoa1, consistent with the behavior of
the wild host (Figure 4d). Ocr was not toxic in the lab strain on its own (see Figure 5a). An F54V
Ocr mutant, previously reported to evade detection by an ARIA known as PARIS[46], also lost the
ability to trigger EcoR22III PCD (Supplementary Figure 3c). Since there were no other R-M loci
in the lab strain nor its native host ECOR22, EcoR22III likely detects Ocr directly rather than its
effect on another R-M system.
Next, we tested whether EcoR22III possessed the canonical R-M activity. We
constructed an active-site mutant of the PD-(D/E)xK restriction nuclease (res*: D1024A,
D1043A double mutant) and compared immunity conferred by wildtype and res* EcoR22III
against a panel of phages (not shown). EcoR22III could block λvir, but infection was not
restricted by the putative restriction-dead mutant (Figure 4e). Furthermore, λvir passaged
through the res* host (and thus modified by the presumably intact EcoR22III methyltransferase)
regained the ability to infect cells carrying wildtype EcoR22III, indicating that EcoR22III indeed
restricts only unmodified phage DNA.
We wondered if the targeted (R-M) and abortive (PCD) defense functions of EcoR22III
could be uncoupled. EcoR22III genes *mod* and *res* individually did not induce cell death,
indicating that PCD is a property of the complex (Figure 4f). The res* nuclease-dead mutation in
EcoR22III that had completely abrogated its ability to target phage λvir (Figure 4e), also
decreased PCD by 100-fold but did not fully abolish Ocr-induced toxicity (Figure 4f). Therefore,
the EcoR22III PD-(D/E)xK nuclease motif appears largely responsible for its PCD and R-M
activities.
Overall, EcoR22III is capable of mounting targeted antiviral defenses as well as an
abortive response in the presence of certain phage proteins, possibly through different
mechanisms mediated by the same effector (Figure 4g). Coupling restriction and PCD could
allow cells to respond dynamically depending on the progress of the infection[48].
343
A distinct Type I R-M derived system deploys a small toxin to provide decoy immunity
In contrast, a second R-M-like ARIA we identified in ECOR55 from our transposon
screens (Table 1b, Supplementary Figure 4) has acquired a similar PCD function but lost the
restriction capability. This locus includes *hsdS* and *hsdM* genes typical of type I R-M but lacks

the nuclease (*hsdR*), and instead encodes a tightly linked 61aa protein. We named this system
ronin, for the master-less samurai in the 1962 Japanese film, *Harakiri*.
Ronin induces cell suicide when Ocr, ArdA, or *aoa1* are expressed in ECOR55 (Figure
4b), but only responds to ArdA and *aoa1* (and not Ocr) when cloned into a lab strain (Figure 5a).
The same results were obtained with a nearly identical system found bioinformatically in
ECOR66 (not shown).
Several lines of evidence suggest that the 61aa protein (hereafter, *ronA*) in Ronin is the
PCD effector. A highly hydrophobic transmembrane helix suggests that RonA might interfere
with membrane integrity (Figure 5b). Over-expression of *ronA* alone is lethal to the host, but this
effect is suppressed by HsdMS co-expression (Figure 5c). Ronin activity is independent of the
methyltransferase function of HsdMS as PCD was not abrogated by the inactivating F287G
mutation in motif IV (NPPF) of the *hsdM* catalytic domain[49] (Figure 5d). However, removal of
ronA inactivates Ronin, such that anti-restriction proteins no longer elicit a PCD response
(Figure 5d). Live cell imaging revealed that Ronin activation arrested growth but did not lyse
cells, consistent with RonA-mediated inner-membrane disruption as a hypothetical PCD
mechanism (Figure 5e). As *ronA* is not lethal when co-expressed with other Ronin components,
we suggest that its activity is sequestered by the putative complex formed by HsdM and HsdS
(hereafter, HsdMS). This hypothetical model is reminiscent of PrrC activation upon blocking of
EcoPrrI restriction by T4-encoded peptide Stp[50]: similarly to how Stp binds the EcoPrrI HsdRMS
complex to release the abortive anticodon nuclease PrrC, an anti-restriction protein might bind
Ronin HsdMS and release the lethal effector.
Next, we examined the gene neighborhoods of *ronA* homologs in all complete bacterial
genomes. We found that they were always linked to *hsdS* and *hsdM* genes, and the vast
majority were arranged in one of two genetic architectures (Figure 5f). The more prevalent
architecture (~70%) contained an uninterrupted (>400 aa) integrase upstream, and an atypical
R-M system downstream (with two DUF262 proteins often found in type IV R-M in addition to
the type I R-M *hsdRMS* genes) transcribed from the opposite strand. This arrangement was
never found in *E. coli* genomes. The less prevalent architecture (~30%) was exclusive to *E. coli*
and often contained a downstream BREX system, also encoded in the opposite strand.
Importantly, most (77%) *ronA*-encoding *E. coli* genomes did not contain any type I R-M systems
anywhere in the genome, and therefore lacked *hsdR* genes (Supplementary File 9). Although it
is certainly possible that an unlinked HsdR is recruited *in trans* in other species, the
predominantly nuclease-lacking configuration in *E. coli* suggests that ARIA activity is Ronin's
primary function. This is reminiscent of decoy immunity (especially common in plants), where a
host factor mimics the target of a pathogen effector. When the pathogen effector binds a decoy
instead of its actual target, it alerts the host to the presence of the pathogen[25].
Thus, Ronin appears to be a type I R-M derivative that has gained the ability to scout for
anti-restriction effectors but lost its prototypical defense function. Ronin encodes the canonical
target of Ocr/ArdA (the type I R-M HsdMS complex), and its activation possibly depends on the
ability of these anti-restriction proteins to bind and inactivate HsdMS (Figure 5g). Consistent
with this, the Ocr F54V mutant, which evades PARIS and EcoR22III but does not lose anti-
restriction activity[46], could not evade Ronin (Supplementary Figure 3c). Notably, Ronin almost
always appears linked to a type I R-M or BREX system, both of which can be blocked by Ocr[43].
Hence, Ronin might drive evolution of Ocr-encoding phages to a BREX/R-M susceptible state,
suggesting that decoy immunity could be a powerful host strategy to constrict the evolution of
virus escape mutants.
394
Prophage-encoded abortive infection mechanisms can be triggered by specific AGs
Three additional abortive loci triggered by counter-defense-associated AGs were
identified through whole-genome suppressor screens (Table 1b, Figure 6). These were all

prophage-linked, implying that prophages can activate superinfection exclusion mechanisms in
response to individual phage products.
First, we found that transposon-insertions in the P4 satellite prophage (especially the
~100-bp intergenic region marked by a star in Figure 6a) alleviated the lethality due to orf116
expression in ECOR15 (Table 1b, Supplementary Figure 4). We cloned the satellite prophage
onto a plasmid and reconstituted this effect in a lab strain of *E. coli*. While P4 often encodes a
hotspot typically occupied by superinfection exclusion systems[46], the P4 construct we cloned
does not contain this hotspot. Therefore, the P4 prophage itself kills the host in response to
expression of orf116 (Figure 6a). Orf116 is the *abc1* gene from phage P22 and is adjacent to a
known recBCD inhibitor *abc2* (Supplementary File 5), but its own function is less clear[51]. Abc1
contains an HTH domain suggesting that it could be a DNA-binding protein (Supplementary
Files 3, 9). *Abc1* expression triggers the PCD function of the P4 prophage, but this toxicity is
blocked when expression of P4 is also simultaneously driven with the strong pBAD promoter
(Figure 6a). Because the toxicity of Abc1 in this strain can be alleviated by interfering with P4
prophage regulation, this result confirms the specificity of the P4-Abc1 interaction. The P4
satellite prophage cannot be induced without a co-resident P2 prophage (which was absent in
our lab strain), ruling out a prophage-induction-based mechanism of action. We speculate that
abc1 interferes with some aspect of transcriptional regulation of the prophage, perhaps, with the
repression of the P4 *kil* gene known to have a cytotoxic effect[52]. This hypothesis seems to be
supported by the observation that the intergenic region that likely contains the *kil* promoter was
heavily targeted by transposon insertions in our suppressor screens (Supplementary Figure 4).
The gop-beta-cII abortive system has been previously identified as a toxin-antitoxin-like
locus[53], and suggested to exert anti-phage activity against coliphages λ and P1[46]. Our
suppressor screens showed that gop-beta-cII toxicity is activated by orf184 (Table 1b,
Supplementary Figure 4), which is widely distributed among coliphages and often present in the
same AR with the anti-restriction gene *ral* (Supplementary File 5). We cloned gop-beta-cII into a
lab strain of *E. coli* and confirmed that orf184 triggered abortive defense (Figure 6b). However,
orf184 is not present in λ or P1.
Finally, a novel prophage-encoded system in ECOR61 inhibits growth upon expression
of orf24 (Table 1b, Supplementary Figure 4), which is extremely narrowly distributed and is
associated with the anti-restriction gene *ocr* in phage genomes (Supplementary File 5). We
reconstituted this effect in a lab strain of *E. coli* and found that the effect is more muted (Figure
6c) than in the native host (Figure 4b). An Icd-like cell division inhibitor protein is the likely
effector, as the system loses function when it is removed (Figure 6c). Our results highlight
parallels between guardianship of defense systems by ARIAs and prophage-mediated
superinfection exclusion (Figure 6d).
434
435 **Conclusion**
436
The PhAGENT platform harnesses the prediction that AGs manipulate host processes to
provide a snapshot of the phage-host interface. This approach can systematically measure the
fitness effects of hundreds of AGs in parallel, as well as their ability to trigger or block bacterial
responses to infection. Here, we focused PhAGENT towards the discovery of counter-defense
AGs.
In addition to higher throughput, the PhAGENT screen offers several advantages over
inferential experiments with individual strains and phages. First, wild bacterial isolates are often
not sensitive to many phages. Even when extensive host-phage collections can be curated,
infection experiments have binary outcomes (i,e. survival or death) which can present
challenges for deconvoluting the multiple mechanisms at play. A gene-delivery approach

bypasses this barrier and allows for the discovery of both the interacting defense and counter-
defense factors at once.
Second, phages that might encode modulators of specific defense systems frequently
prove impossible to procure and grow. Gene-synthesis allows AGs to be derived from any
sequenced phage. In particular, assaying AGs individually in many wild bacterial strains can
directly identify thus-far elusive triggers of defense systems. Abortive defense systems are
generally thought to detect conserved phage factors (nucleic acids, capsid, terminase, portal, or
a general effect such as host transcriptional arrest)[54-56], but our results show that poorly
conserved phage genes can also trigger suicide mechanisms in the host and its prophages.
Broadly understanding the principles of protein detection in antiviral immunity could yield a
generalizable protein-targeting toolkit.
Third, we identified exaptations of well-studied R-M systems for mounting PCD
responses to anti-restriction proteins that were not identified by sequence comparisons with
known defense/counter-defense factors, which can miss subtle differences or miss divergent
genes entirely. For instance, EcoR22III does not contain known domains that might suggest its
PCD activity. Similarly, *ronA*-like toxins have not previously been linked with type I R-M genes
(missed likely due to their small size).
Fourth, the PhAGENT screen offers more specificity than overexpressing putative
defense systems in model strains to determine whether they are involved in antiviral defense in
nature because the phage-host interaction is (a) revealed principally by a phage
countermeasure to thwart the defense system, and (b) discovered in wild host strains where the
defense systems are natively regulated. While redundant defense pathways may lead to missed
interactions, co-expressing colinear configurations of multiple AGs in the same hosts could
further increase the sensitivity of the approach.
Finally, any genetically tractable bacteria or archaea can be queried by PhAGENT. Our
approach is especially suitable for discovering rare interactions that may be present only in a
particular evolutionary niche. Deploying this platform more broadly in various pathogenic
bacteria will help determine how phage-therapy approaches will need to be tailored to overcome
species-specific antiviral mechanisms.
While we synthesized a library of 200 AGs to test the PhAGENT platform, the same
workflow can scale to substantially larger libraries, since we routinely perform transposon
mutagenesis screens with ~10^5 elements. The throughput could further be expanded by a
double-barcoding strategy that allows a single read to identify both the host strain and AG. The
platform is therefore only limited by our ability to synthesize phage genes. Stepwise screens
could help explore the phage pangenome more efficiently; for instance, we sampled >60
accessory regions here, but could focus subsequent libraries on ARs with confirmed host-
interacting AGs. Expanding the scope beyond AGs could also highlight secondary host-
modifying functions of conserved phage genes such as capsids, polymerases, tail proteins, etc.
Overall, PhAGENT uncovers the fundamentally layered nature of anti-phage immunity in
bacteria, and can tease the layers apart, providing a holistic view of bacterial defenses. We
discovered phage effectors that interact with this multilayered defense architecture at every
step: virus proteins that inhibit metabolism to modify the cell surface and block viral entry,
counter-defense effectors that block the activity of antiviral restriction mechanisms inside the
cell, cellular abortive immunity that guards against counter-defense strategies deployed by
viruses, and phage-encoded triggers of superinfection exclusion mechanisms in prophages and
other mobile elements in the host genome. With more detailed maps of phage-host interactions
in many species, we hope to generalize these themes in phage-host co-evolution.

Figure 1. Phage Accessory Gene Exploration aNd Testing (PhAGENT) platform. (A)
Schematic of a genomic region showing core (shown in grey) and accessory (in various colors)
genes in a hypothetical phage genome cluster. **(B)** Schematic of algorithm to exhaustively
identify accessory regions using pairwise combinations of genes, without *a priori* genome
clustering. **(C)** A sample high-scoring AR in a family of *Salmonella* phages. Purple shading
denotes >80% and green denotes >90% nucleotide identity. **(D)** Statistical features of ARs and
AGs. 2D chart depicts saturation curves (cumulative distribution functions) of unique AGs (left
axis) and phage genomes (right axis) represented in ~2000 non-redundant ARs ranked by
score on the X-axis. 3D bar graphs show various distributions (length of AGs, number of AGs
per AR, number of AGs per phage, and number of ARs per phage). Color-coded histograms in
each bar graph are cumulative i.e., the first histogram (dark blue) shows the top 20% highest-
scoring ARs, the next (light blue) represents top 40% and so on. **(E)** Schematic of the
PhAGENT screen. A barcoded library of IPTG-inducible AGs is transformed into each host in
triplicate via Tn7 transposition resulting in single-copy chromosomal insertions. Each replicate
host library is then challenged with each phage, and samples are collected at time 0, and 30-60
min post infection. Uninfected but induced controls are maintained for every experiment.
Another round of plating and scraping depletes barcodes from cells killed by phage infection or
AG expression, and high-throughput sequencing is used to quantify remaining barcodes. >>10X
as many colonies as the number of AGs are scraped at each step to avoid bottlenecking.

Figure 2. AGs that sensitize hosts to infection by modifying barrier defenses. (A)
Heatmap of log2-transformed fitness scores comparing infected and uninfected samples. Of 200
AGs tested, only AGs with a host-sensitizing phenotype are shown on the X-axis. Red
boxes/bold names highlight counter-defense AGs selected for further study. Named AGs are
orf1:*ocr*, orf7:*aoa1*, orf48:*gnarl1*, orf63:*gnarl2*, orf87:*ral*, orf92:*gnarl3*, orf116:*abc1*, orf143:*lpII*
and orf169:*ardA*. **(B)** Schematic of follow-up transposon knockout screens, performed in the
native host without AGs. Graphs show fitness of transposon-mediated knockouts in infected (Y-
axis) and uninfected (X-axis) Tn5-libraries from three representative hosts challenged by T5.
Red dots are gene-disruptions that lower bacterial fitness upon infection (e.g., disruptions in O-
antigen biosynthesis). **(C)** Plaque assays with phage T5 and hosts from (D) with *gnarl* genes
repressed (top) or induced (bottom). **(D)** Lipopolysaccharide (LPS) visualized by electrophoresis
from host-AG combinations in (E) where AGs produced a phage-sensitizing effect. **(E)** Log2-
transformed protein representation scores (MaxLFQ) for the top 30 enriched host proteins upon
expression of AGs. **(F)** log10-transformed T5 infection scores in wild hosts ECOR1 (green),
ECOR3 (orange), ECOR22 (blue) with or without UDP-glucose biosynthesis pathway genes
galU, galE, galF, and ECA precursor *wecB* cloned onto a plasmid and overexpressed. (See also
Figure S2)

Figure 3. AGs that block canonical antiviral defenses. (A) Fitness of transposon-mediated
knockouts in infected and uninfected Tn5-libraries from ECOR21 challenged by various phages.
(B) Schematic of ECOR21 defense against T2 through both barrier defenses and a defense
system which is putatively blocked by orf143/lpII. **(C)** LPS visualized from ECOR21 with the O-
antigen biosynthesis initiator *wecA* deleted by allelic exchange. **(D)** Whole-genome transposon-
knockout screens as in (A) but with the ECOR21::Δ*wecA*. **(E)** Variants of the BREX/GmrSD
operon cloned onto a plasmid along with their native promoters and tested in a lab strain of *E.
coli* (TOP10) against T-even phages (T2, T4, T6). Removal of the PglX gene inactivates BREX.
Active-site mutations that disable GmrSD are shown as two parallel red bars (GmrSD* double
mutant; D474A, H475A). Ocr, lpI and lpII are expressed from single-copy chromosomal
insertions. Bar graphs show log10 infection scores. **(F)** GmrSD and its inactive variant GmrSD*
cloned onto a plasmid and tested in TOP10 for restriction activity against T-even phages.
"pBAD" indicates the defense construct was overexpressed from the plasmid. **(G)** Plaque

545   assays with TOP10 cells with uninduced GmrSD (i.e., not overexpressed) challenged by T-even
546   phages.
547
Figure 4. AGs that trigger abortive defense in the host. (A) Heatmap of log2-transformed
fitness scores comparing induced and uninduced samples without phage infection. Of 200 AGs
tested, only AGs that exhibited lethality in some (but not all) hosts are shown. Red boxes/bold
names highlight conditional-lethal AGs selected for further study. Named AGs are orf7:*aoa1*,
orf87:*ral*, orf63:*gnarl2*, orf1:*ocr*, orf92:*gnarl3*, orf116:*abc1*, orf169:*ardA* and orf2:*arn*. **(B)** Host-
AG combinations in red boxes from (A) tested individually for AG-induced lethality (serial tenfold
dilutions of saturated culture; AGs repressed: left, induced: right). **(C)** Schematic depicts
hypothetical strain-specific abortive defense system being triggered by AGs, and inactivated in
follow-up transposon knockout screens (inset). Data from a representative screen with a Tn5
library constructed from ECOR22 with chromosomally encoded *ocr*. Proportion of transposon
insertions per gene in surviving cells with *ocr* repressed is plotted on the X-axis, and with *ocr*
induced on the Y-axis. Y-axis interrupted to resolve genes other than type III R-M *mod*, *res*.
(See also Figure S4) **(D)** Abortive phenotype of EcoR22III in its native host, and reconstituted
when cloned onto a plasmid and expressed in TOP10 (AGs repressed: left, induced: right)**. (E)**
Plaque assay showing R-M phenotype of EcoR22III against methylated and unmethylated λvir
in TOP10. λvir* was methylated by passaging in TOP10 strain with nuclease-dead EcoR22III
(res* double mutant; D1024A, D1043A) and tested for its ability to evade restriction by wildtype
or res* EcoR22III. **(F)** As in (D), PCD activity of EcoR22III variants in TOP10 (wildtype, nuclease
dead (res*), and *mod*, *res* genes expressed individually). **(G)** Hypothetical model of EcoR22III
R-M and PCD activities. EcoR22III kills its host in the presence of Ocr, partially relying on its
nuclease activity. "pBAD" indicates the construct was overexpressed from the plasmid in (F-H).
569
Figure 5. Decoy immunity in bacteria. (A) Schematic of Ronin system from ECOR55.
Abortive phenotype reconstituted in TOP10 upon expression of Ronin from a plasmid (serial
tenfold dilutions of saturated culture; AGs repressed: left, induced: right). **(B)** Prediction of
transmembrane character along the 61 aa RonA sequence. **(C)** Lethality of Ronin components
tested when overexpressed in TOP10 (indicated as "pBAD") without AGs. **(D)** As in (A), genetic
requirements for the abortive phenotype of Ronin. Red bar in *hsdM* (second row) denotes the
F287G active-site mutation. (See also Figure S5) **(E)** Time-lapse imaging of TOP10 cells
expressing Ronin with (bottom) or without (top) chromosomally-integrated ArdA. **(F)** Two
prevalent architectures of *ronA*-linked type I R-M-like abortive system (ronin), based on the
presence of an upstream DUF4102 integrase. Number of genomes with a full type I R-M system
(with *hsdR, hsdM, hsdS* genes) anywhere in the genome is indicated. **(G)** Schematic depicting
hypothetical RonA "uncaging" when ArdA is present.
582
Figure 6. AGs that trigger prophage-encoded superinfection-exclusion mechanisms. (A)
Abortive phenotype of P4 satellite prophage from ECOR15 cloned onto a plasmid and
expressed in TOP10. Orf116 is expressed from a single-copy chromosomal insertion (serial
tenfold dilutions of saturated culture; AGs repressed: left, induced: right). Star marks an
intergenic region that was strongly targeted in transposon suppressor screens. (See also Figure
S4) **(B)** As in (A), abortive phenotype of gop-beta-cII system from ECOR59 upon expression of
orf184. **(C)** As in (A), abortive phenotype of novel prophage-encoded superinfection exclusion
system from ECOR61 upon expression of orf24. **(D)** Schematic representation of similar
defense architecture of ARIAs and superinfection exclusion systems. "pBAD" indicates the ARIA
was overexpressed from the plasmid in (A-C).
593
Table 1. Summary of transposon-mediated whole-genome knockout screens. Follow-up
screens to find **(A)** phage exclusion mechanisms that could be the targets of counter-defense

AGs identified in PhAGENT screen and **(B)** abortive defense mechanisms that might trigger
PCD in response to conditional-lethal AGs. A total of 26 whole-genome knockout screens were
performed in (A), with 7 Tn5 libraries (one in each host) challenged by the indicated phages. In
contrast, each host-AG combination in (B) was screened as a separate Tn5 library.

Acknowledgements

S.S. was supported by the Damon Runyon Fellowship Award (DRG 2352-19). Experiments were funded by awards to J.B.D by the Vallee, Searle, and Kleberg Foundations. We thank Jason Peters for technical discussions, Andy Millard for the Enterobacteriophage genome dataset, and Ry Young, Andrew Fire, Carol Gross and Jonathan Weissman for guidance.

Author Contributions

Conceptualization, S.S., J.B.D.; Methodology, S.S., E.L.; Resources, A.F., D.S.; Software, S.S.; Investigation, S.S., H.C., D.S.G.; Formal Analysis, S.S., H.C., M.J., K.S.M., E.V.K.; Visualization, S.S., H.C., M.J., K.S.M.; Writing – Original Draft, S.S., H.C.; Writing – Reviewing & Editing, S.S., H.C., M.B., E.V.K., J.B.D.; Supervision, S.S., J.B.D.; Funding Acquisition, S.S., J.B.D.

Declaration of Interests

J.B.D. is a scientific advisory board member of SNIPR Biome, Excision Biotherapeutics, and LeapFrog Bio, consults for BiomX, and is a scientific advisory board member and co-founder of Acrigen Biosciences. The Bondy-Denomy lab received research support from Felix Biotechnology.

Inclusion and Diversity

We support inclusive, diverse, and equitable conduct of research.

Supplementary Figure 1. Phage-host combinations included in PhAGENT screen (A)
Phage susceptibility of select wild ECOR strains. Numbers in boxes are log10-transformed
infection scores (7 = 10^7-fold higher plaquing than 0). Borders indicate that individual plaques
were visible in that phage-host combination. Combinations with attenuated phage infection that
were selected for the PhAGENT screen are in yellow. Combinations with attenuated infection
that were attempted but yielded no survivors in liquid infection experiments (perhaps due to
abortive defenses) are in red. **(B)** Phage-host combinations where counter-defense phenotypes
were confirmed for any AGs. Red denotes combinations where the screen could not be
completed (as in (A)). Yellow: combinations that were tested where none of the 200 AGs
produced counter-defense effects. Green: combinations where phage-sensitizing effects could
not be verified by plaque assays. Blue: combinations where counter-defense phenotypes were
confirmed by plaque assay (at least 100X higher EOP upon AG expression). **(C)** Defense
repertoire of selected ECOR strains. Computational prediction of the presence or absence of
various defense systems in ECOR strains used in the PhAGENT screen. Predictions were
generated using the ISLAND software suite[57].

Supplementary Figure 2. Over-expression of *galU* reverses T4 sensitization by *gnarl3*.
log10-transformed T4 infection scores in wild hosts ECOR1 (green), ECOR3 (orange), ECOR22
(blue) with or without UDP-glucose biosynthesis pathway genes *galU, galE, galF,* and ECA
precursor *wecB* cloned onto a plasmid and overexpressed.

Supplementary Figure 3. Abortive effects of known R-M inhibitors in wild *E. coli.* (A)
Schematic representation of conditional-lethality data from Figure 5b, showing only known R-M
inhibitors and *aoa1*. **(B)** AGs from (A) tested for their ability to block EcoKI restriction of phage
λvir in plaque assays with restriction-competent *E. coli* MG1655. **(C)** F54V mutant of Ocr (known
to evade a previously described ARIA, PARIS) still triggers ARIA activity in ECOR55 but not
ECOR22.

**Supplementary Figure 4. Transposon insertions that suppress AG-induced conditional-
lethal phenotypes.** Number of unique transposon insertions recovered from survivors from Tn5
libraries, mapped to putative abortive loci. Tn5 libraries were prepared in the indicated wild
hosts and survivors obtained upon AG-induction. E.g., in ECOR15, >500 insertions were
recovered from the ~100 bp intergenic region marked by a star, and at the very 3'end of the
DNA primase CDS. Scale is indicated for each locus.

Supplementary Figure 5. Upstream non-coding region is required for Ronin PCD activity.
(A) The first 135-bp of the 235-bp upstream non-coding region (marked by a thick red line) are
required for activity as truncation down to 100-bp causes inactivation. ArdA is chromosomally
integrated under pLlacO-1 control and strains in the left (ArdA repressed) and right (ArdA
induced) panels are isogenic. **(B)** Over-expression of 235-bp upstream region is not toxic in
TOP10 indicating that it is not the PCD effector. **(C)** As in (A), abortive phenotype of truncated
Ronin system when overexpressed (indicated as "pBAD"). Over-expression of truncated Ronin
did not rescue the PCD defect indicating that the upstream region contributes something in
addition to the promoter.

Supplementary File 1. Worked example of PhAGENT algorithm. Readme.txt describes how
to run the code. Sample database of Salmonella phage genomes is included. All ARs from this
phage dataset are reported in the output file SPFMphages_accRegions_nr_all.txt. Any AR can
be extracted and visualized using PhamLite[58]. The AR shown in Figure 1d is extracted by
default.

Supplementary File 2. Manually inspected accessory regions from Enterobacteriophage
genomes. Visualizations of all ~200 accessory regions (ARs) that were manually inspected.
The first slide in the file shows the layout for the rest, with the AR score calculated according to
the formula in Methods, followed by a unique identifier for each hypothetical phage group
("phamily"). Each AR was also given a short, memorable name (top right). Slides with green
backgrounds indicate ARs from which genes were selected for PhAGENT screen. Authors'
impressions upon AR inspection are included in the Notes section.

Supplementary File 3. Phage accessory genes selected for PhAGENT screen. 200
accessory genes (AGs) selected for screening. Genomic locations, protein and phage genome
identifiers, and protein descriptions are indicated. IDs denoted by "UDP" correspond to the
orf/AG numbers used throughout the manuscript.

Supplementary File 4. Experimental record of PhAGENT screen. A detailed record of all
PhAGENT experiments is presented. The first sheet contains phage-host combinations, dates
of the experiment, and preliminary results. The next three sheets each correspond to one
replicate of PhAGENT screening. The timeline of each experiment indicates library growth
characteristics. Empirically measured multiplicity at time of infection, and dilutions used before
plating are also recorded. Screens that had to be repeated for technical reasons are color-
coded.

Supplementary File 5. Accessory regions of AGs selected for further study. Visualizations
of ARs where AGs that produced counter-defense or conditional-lethal phenotypes in PhAGENT
screen (see Table 1) are located.

Supplementary File 6. Experimental record of follow-up transposon whole-genome
knockout screens. A detailed record of all follow-up transposon screens is presented. The first
sheet contains details of screens designed to phenocopy counter-defense phenotypes of AGs
identified in PhAGENT screen. These screens are performed with RB-TnSeq libraries prepared
with naïve ECOR hosts (no AGs present). Each library is challenged by various phages one at a
time. The timeline of each experiment indicates library growth characteristics. Empirically
measured multiplicity at time of infection, and dilutions used before plating are also recorded.
Screens were performed in two stages: first, a pilot with one phage for each RB-TnSeq library,
and then a larger experiment with all other phages that showed enhanced infection upon
expression of a counter-defense AG. The second sheet contains details of screens designed to
inactivate putative abortive defense mechanisms responsible for AG-triggered conditionally-
lethal phenotypes. Tn5 libraries are constructed in ECOR strains already transformed with Tn7
transposons carrying trigger AGs (maintained under strict glucose suppression). No phages are
involved. Survivors are selected and sequenced after trigger AG induction upon plating on
IPTG. Screens are color coded to indicate whether they were ultimately attempted.

Supplementary File 7. Host-AG combinations for affinity-purification/mass-spectrometry.
Matrix of all host-AG combinations where potential binding partners of AGs were affinity-purified
from the native host expressing counter-defense AGs, and identified by mass-spectrometry.
Experiments were performed in triplicate. Orf74 was inert in PhAGENT screening and was used
as a control for all experiments in the relevant hosts.

Supplementary File 8. AlphaFold predictions for all AGs that produced phenotypes in
PhAGENT screens. PDB files containing predicted structures of orfs 7 (aoa1; triggers PCD by
Ronin), 24 (triggers PCD in ECOR61), 48, 63, 92, (*gnarl*1-3; putative O-antigen modifiers), 116
(*abc1*; triggers PCD by P4 prophage), 126 (broad-spectrum counter-defense), 143 (IpII; inhibitor

of GmrSD), 145 (triggers PCD in MG1655), 148 (broad-spectrum counter-defense), and 184
(triggers PCD by gop-beta-cII).
725
Supplementary File 9. Genomic architectures of all ECOR55-like Ronin systems.
Spreadsheet contains accession numbers and genomic locations of all *ronA* homologs.
Grouping variables indicate whether a type I R-M system (containing *hsdR, hsdM,* and *hsdS*)
was found anywhere in the genome, and also whether a full (>400 aa) DUF4102 integrase was
present immediately upstream of Ronin. HTML files split Ronin systems into two groups based
on the presence of the integrase, and gene annotations can be viewed by mouse-over.

Methods
733
Algorithm to identify phage accessory regions
1706 non-redundant Enterobacteriophage genomes were obtained (with help from Dr.
Andrew Millard, Leicester) from public databases. These phages were both lytic and lysogenic,
but were sequenced and assembled from cultured isolates (i.e. prophages and metagenomic
samples were not included). All annotated protein records from all the phages were assigned
unique numeric IDs and clustered by MMseqs2[59]. ORF IDs in phage genomes were then
replaced by their MMseqs2 cluster representative IDs in each genome. Thus, each phage
genome can be represented as an ordered list of IDs of clustered proteins. The gene-
neighborhood of every ORF (up to 10 genes upstream and downstream) was extracted from
every virus. Given that phage genomic termini can be difficult to identify, every genome was
considered potentially circularly permuted. Allowing for circularization also ensures all gene-
neighborhoods were the same size and putative ARs towards ends of genomes would not be
missed. A 2D matrix of all possible pairwise combinations of genes was constructed, with every
matrix position containing lists of (a) the phages that each gene appears in, (b) the 10 upstream,
and (c) 10 downstream neighboring ORFs of each gene in every genome. We then traversed
the matrix and assessed every gene pair to identify all possible ARs, defined as the region
between a pair of genes that are both present in at least 6 different phage genomes but are no
more than 10 genes apart in any genome. Regions trivially containing the same genes in each
participating genome were discarded. Genomes involved in a putative AR were then compared
pairwise at the nucleotide level using FastANI[60], and ARs with divergent genomes were
removed. In practice, this was done by pre-computing an all-vs-all matrix of genome ANI and
then traversing subsets of the matrix containing all the genomes participating in the putative AR
to ensure no cells were empty (FastANI produces no output when very divergent genomes are
compared). Various parameters were extracted from surviving ARs. N is the number of unique
genes in the AR. The vector c contains conservation numbers (the number of different genomes
that each gene appears in; also indicated on top of each gene) for all the unique genes in this
AR. The vector l contains the lengths (in numbers of genes) of the AR in each genome. CV
denotes coefficient of variation. $GCdev$ is a vector containing one minus the absolute deviation
in GC content of each accessory gene from its resident genome. A scaling factor is also
included to normalize the score by the number of phage genomes in the group that have at least
one gene in the AR. ARs were scored according to the formula below.
765

$$ARscore = \frac{N * median(GCdev) * CV(c)}{CV(l) * mean(c)} * \frac{\#non\_empty}{\#genomes}$$

766
767   To filter out ARs that were subsets of other ARs, we successively removed redundant,
768 low scoring ARs with at least 90% of genes that had already been encountered in higher scoring
769 ARs. The top 200 ARs were examined manually and 1-15 AGs were selected from 62 ARs
770 (Supplementary Files 2,3). A worked example of the code is provided in Supplementary File 1.
771
Strains and Vectors
The *E. Coli* Reference (ECOR) collection of wild *E. coli* strains was obtained from the
STEC Center (Michigan State University). Lab strains were obtained as follows: MG1655 (from
Dr. Carol Gross, UCSF), C990 (from Dr. Ry Young, Texas A&M), DH5α (NEB), DH10ß/TOP10
(Life Technologies), BW25113 and BW25141 (from Dr. Vivek Mutalik, LBNL), and WM6026
(from Dr. Jason Peters, UW-Madison). All strains were cultured at 37°C with shaking (180-225
rpm) in lysogeny broth (LB) (10 g/l tryptone, 5 g/l yeast extract, 10 g/l NaCl with 15 g/l agar for
plates) supplemented with 100 µg/ml of ampicillin/carbenicillin, 10-20 µg/ml gentamicin (except

WM6026, which yielded more consistent colony sizes with 50 µg/ml), 50 µg/ml kanamycin, and
0.5% w/v glucose, 0.1% w/v L-arabinose, and 1 mM IPTG (isopropyl-b-D-thiogalactopyranoside)
as needed. The WM6026 strain required diaminopimelic acid (DAP) for growth, which was
supplied at 300 µM. All strains were stored in 20% glycerol at -80°C for long term storage.
R6K-origin pTn7 plasmids (from Dr. Jason Peters, UW-Madison) – the
pJMP1039/pTn7C1 helper plasmid (carbenicillin) and the pJMP1360/pTn7C185 transposon
vector (gentamicin, carbenicillin) – were used for integration of phage AGs into the *E. coli*
genome. All host defense systems identified in the various screens were placed under the
araBAD promoter (arabinose inducible) in the pBAD Myc/His A vector (Invitrogen). Conjugative
allelic exchange was carried out using R6K-origin pKEK2201 (kanamycin) (from Karl E. Klose,
UT San Antonio). All plasmids were constructed by Gibson Assembly.
R6K-origin plasmids were maintained in BW25141 (routine cloning) and WM6026
(conjugative donor strain). All other plasmids were maintained in DH5α or TOP10. All plasmid
sequences were verified by Sanger or whole-plasmid sequencing (Plasmidsaurus, Primordium)
and are available upon request. Frequent plasmid dimerization during the Gibson Assembly
procedure (especially when using the NEBuilder reagent) was ameliorated by using 10-fold less
of one of the insert DNA fragments in a reaction involving at least 2 insert fragments (not
including the backbone).
798
AG conjugation into host strains
AGs were delivered to *E. coli* hosts by Tn7 transposition via conjugation performed by
triparental mating as described previously[61]. Briefly, the transposon and helper plasmids were
delivered to the recipient from WM6026 donor strains by overnight incubation on LB agar plates
supplemented with 300 µM DAP and 0.5% glucose to repress AG expression. The next day,
cells were scraped and resuspended in 1 ml LB, and 1000X-20000X dilutions of this conjugation
mixture were plated on LB plates (lacking DAP) supplemented with the appropriate antibiotics
and 0.5% glucose. All strains were maintained under strict glucose repression of the pLlacO-1
promoter at every step until screens were performed. Thus, all AGs were maintained as single-
copy chromosomal inserts at the same attTn7 site in the genomic region between *glmS* and
pstS genes.
Efficient Tn7 transposition is necessary to construct a AG library in any given *E. coli*
host. Tn7 conjugation-transposition efficiencies were determined in 44 ECOR stains containing
an intact *glmS-pstS* region (we note that other strains can support Tn7 integration as well). Tn7
transposons carried only a barcode sequence and the gentamicin-resistance gene to minimize
payload-dependent variation in conjugation efficiency between strains. Efficiency of conjugation
was determined by streaking the conjugation mixture on two different gentamicin concentrations
(25 µg/ml and 10 µg/ml) to account for possible differences in natural antibiotic resistance
between the wild strains. The barcode sequence was verified by Sanger sequencing and both
flanks were checked by PCR to ensure proper integration of the Tn7 transposon at the expected
attTn7 site. Strains that supported Tn7 transposition at high levels and were amenable to agar-
overlay plaque assays were selected for further experiments.
821
DNA extraction + Nextera Genomic Sequencing + Analysis
Genomic DNA was extracted using a modified SDS–proteinase K method: Briefly, cells
pelleted from 50 µl of saturated culture (or 5 µl of the cell mixture scraped from selective plates
during AG library construction) were re-suspended in 200 µl of lysis buffer (10 mM tris, 10 mM
EDTA, 400 mg/ml proteinase K, 0.5% SDS) and incubated at 50-55°C for 1 hour. Subsequently,
the temperature was lowered to 37°C and RNase A (Thermo Scientific) was added to a
concentration of 1 mg/ml. After 30-60 min of incubation, the digests subsequently purified using
the Genomic DNA Clean & Concentrator Kit (Zymo Research). DNA was prepared for whole-
genome sequencing using the Nextera Flex Library Prep Kit (now, the Illumina DNA Prep Kit)

according to manufacturer's instructions, except that we lowered the volumes of all
tagmentation reactions 5-fold, and used a custom dual-indexed primer set JSW-SS-22:33
(CAAGCAGAAGACGGCATACGAGAT NNNNNNNN GTCTCGTGGGCTCGG) and JSW-SS-
34:41 (AATGATACGGCGACCACCGAGATCTACAC NNNNNNNN TCGTCGGCAGCGTC) to
amplify libraries for 11-13 cycles using Phusion High Fidelity PCR Master Mix (NEB) instead of
the Illumina-supplied PCR reagents. The N_8 sequences correspond to reverse-complemented
Nextera DNA indexes N701 to N712 and N501 to N508, respectively (Illumina). Libraries were
resolved by agarose gel electrophoresis and DNA was excised in the 300-400 bp range. Gel
slices were purified using the Zymo Gel DNA Recovery Kit and sequenced on the Illumina
MiSeq using 150-cycle v3 or 600-cycle v2 reagent kits.
Sequencing adaptors were trimmed from reads using cutadapt, mapped to reference
genomes using bowtie2, and visualized using IGV. Reference genomic contigs of ECOR strains
were obtained from the USDA[62]. SNPs, indels, and large structural variants were analysed
using a mapping-free variant finder[63].
845
Phage propagation and plaque assays
High-titer phage stocks were generated by growth in liquid culture. All phage infections
were performed at 37°C in LB supplemented with 10 mM $MgSO_4$ and 5 mM $CaCl_2$. *E. coli*
BW25113 or C990 were grown to an optical density (at 600 nm; OD_{600}) ~ 0.5 (1-5 X 10^8 cfu/ml)
and infected with the desired phage at MOI ~ 5. Infections were allowed to proceed at 37°C with
agitation for 5-6 hours until the cultures were clear. Lysates were clarified by centrifugation
(8000 X g for 2 min) and sterile-filtered using a 0.45 µm SFCA syringe filter (Millipore). Phage
lysates were stored at 4°C.
Bacteriophages T2, T3, T4, T5, T6, T7, P1vir were obtained from Dr. Vivek Mutalik
(LBNL), and λvir was obtained from Dr. Ry Young (Texas A&M). To the extent possible, all
phages were grown on *E. coli* C990 which lacks restriction-modification. T4 and P1vir could not
be grown on C990 and were instead grown on *E. coli* BW25113. To ensure that our phage
lysates were free from ancestral type I R-M methylation marks or other host-passage derived
DNA modifications, we serially passaged all phages in the desired hosts 3 times. All phage
genomes were subsequently verified by Nextera high-throughput sequencing.
Phage titers were determined against various hosts by the agar-overlay method with LB
top agar containing 0.7% agar, 10 mM $MgSO_4$ and 5 mM $CaCl_2$. Phage spot-titration assays
were performed with 10-fold serial dilutions of phage lysates (3-5 µl of each dilution pipetted
onto bacteria immobilized in top-agar overlay using a multichannel pipette).
865
AG Library Preparation + NGS verification + testing post-conjugation evenness
200 AGs with unique 10 bp DNA barcodes were synthesized (Twist and IDT) and cloned
individually under pLlacO-1 control (IPTG inducible, repressed by glucose) into a modified
pTn7C185 vector (with mobile CRISPRi components sgRNA and dCas9 removed). Plasmid
assembly reactions were performed in 96-well format using the 2X Gibson assembly master mix
(NEB) and transformed into chemically-competent BW25141 cells in 96-well format.
Transformations were individually plated on LB agar supplemented with 20 µg/ml gentamicin
and 0.5% glucose and one colony was initially checked by Sanger sequencing from each plate.
Where a successful transformant could not be obtained on the first attempt, more colonies were
sequenced in successive rounds until a plasmid with no mutations in the region spanning the
pLlacO-1 promoter, the AG ORF, and the barcode sequence was obtained. Verified vectors
were miniprepped using the ZR Plasmid miniprep – Classic kit (Zymo) and transformed into
chemically-competent WM6026 cells in 96-well format, and each transformation was plated
individually on LB agar supplemented with 50 µg/ml gentamicin, 300 µM DAP, and 0.5%
glucose.

One colony from each of the 200 WM6026 transformants was grown in 2 ml square-well
96-well plates with shaking overnight, and 100 ul of each culture was then pooled together to
yield ~20 ml of the donor mixture. This mixture was concentrated down to 10 ml by
centrifugation at 6000 g for 10 min, and 5 ml of 60% glycerol (20% final) was added to yield the
donor library. The 15 ml library was divided into single-use 50 ul aliquots (each containing >50M
cells, i.e. >300K of each donor strain) and stored at -80°C.
Genomic DNA was extracted from one of the library aliquots and all constructs in the
final pool were verified by Nextera high-throughput sequencing. This allowed us to assess
library evenness as well as verify the fidelity of our cloning, colony picking, and strain pooling
process. Overall, 195 AGs were cloned successfully. One AG (orf88) never yielded any colonies
in multiple cloning attempts, possibly due to its cytotoxicity. Three AGs (orfs 36, 52, 94) were
dropouts during the WM6026 growth step, and another two (orfs 25, 84) were discovered to
contain mutations upon high-throughput sequencing that had been missed by Sanger
sequencing. Orf25 had previously presented difficulties in cloning, and did not yield a mutation-
free construct even after checking >12 colonies (whereas most transformations yielded a
correct construct on the first attempt). The orf84 construct contained a mixed base at the last
position of the second repeat of the Lac-operator sequence upstream of the ORF, which would
not be expected to affect any downstream assays significantly.
The donor library was conjugated into MG1655 *E. coli* according to the Tn7 transposition
procedure (above) and a 1000X dilution of the conjugation mixture was plated on LB agar
supplemented with 20 µg/ml gentamicin and 0.5% glucose. We routinely use multiple 150 mm X
15 mm bacteriological petri dishes (Corning) to obtain sufficient numbers of transconjugant
colonies to assemble libraries. The next day, >>2,000-20,000 transconjugant colonies (10-100X
the number of AGs in the library) were scraped and combined thoroughly, and the thick cell
mixture was directly frozen at -80°C in 20% glycerol. Genomic DNA was extracted from this
mixture and the distribution of DNA barcodes corresponding to individual AGs was measured in
both this post-conjugation strain mixture as well as the genomic DNA extracted from the
WM6026 donor library aliquot.
To amplify barcodes for high-throughput sequencing, genomic DNA was subjected to
two successive rounds of PCR. The DNA barcodes were amplified from genomic DNA using
universal primers JSW-SS-170 (CGACGCTCTTCCGATCTNNNNN
<u>TGATGTCGTTGTTGCCATCG</u>) and JSW-SS-171 (ACTGACGCTAGTGCATCA
<u>CTTTCTGAGCCAGTGTTGCT</u>) and the Q5 Hot Start High-Fidelity 2X PCR master mix (NEB)
according to manufacturer's instructions. Sequencing adaptors were attached in a second round
of PCR using amplicons from the first PCR round as the templates, with dual-indexed primer
sets JSW-SS-42:53 (CAAGCAGAAGACGGCATACGAGAT NNNNNNNN
GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCT <u>ACTGACGCTAGTGCATCA</u>) and JSW-
SS-54:61 (AATGATACGGCGACCACCGAGATCTACAC NNNNNNNN
ACACTCTTTCCCTACA<u>CGACGCTCTTCCGATCT</u>), where the N_8 sequences correspond to
reverse-complemented TruSeq HT indexes D701 to D712 and D501 to D508, respectively
(Illumina). Template-matching regions in the primers are underlined. Cycling conditions for
round 1 were as follows: one cycle at 98°C for 1 min; two cycles at 98°C for 10s, 66°C for 30s,
and 72°C for 10s; 22 cycles at 98°C for 10s, and 72°C for 20s; and one cycle at 72°C for 2 min.
Conditions for round 2 were one cycle at 98°C for 1 min; two cycles at 98°C for 10 s, 64°C for
30 s, and 72°C for 10 s; 4 cycles at 98°C for 10 s, and 72°C for 20 s; and one cycle at 72°C for
2 min. 0.5-1 ul of unpurified 1[st] round reaction product was used as template for the 2[nd] round of
PCR. Amplicons from the 2[nd] round were gel-purified by electrophoresis (3% agarose gel, 4.2
V/cm, 2 hours) and quantified using the 1X dsDNA HS kit with the Qubit 4.0 Fluorometer
(Invitrogen). Amplicons were sequenced on a MiSeq using the 150-cycle v3 reagent kit
(Illumina) in single-end format (100 bp) with two (8 bp) index reads. DNA barcodes were

trimmed from the reads and the proportions of various AGs in the mixtures were determined
using the prevalence of the corresponding barcode sequence.
Concordance between the Nextera whole-genome sequencing and the barcode
sequencing of the WM6026 donor mixture showed that barcode representation was a suitable
proxy for the true distribution of AGs in the mixture and that amplification biases during library
preparation were not a significant source of noise. Concordance between barcode sequencing
in the WM6026 donor and MG1655 post-conjugation recipient mixtures indicated that the Tn7
transposition and colony scraping process does not meaningfully bottleneck AG distribution.
The AG library was then conjugated (in triplicate) into ECOR strains that were previously
confirmed to support robust Tn7 transposition. Strain mixtures were prepared in a procedure
analogous to library construction in MG1655, except higher dilutions of the ECOR+AG
conjugation mixtures (10,000-20,000X) were plated and the gentamicin concentration on
selective plates was lowered to 10 µg/ml to account for higher steady state cell densities and
higher gentamicin sensitivity of the wild *E. coli* strains. As with the MG1655 library, thick cell
mixtures comprised of >>2,000-20,000 colonies scraped from each replicate conjugation were
directly stored at -80°C in 20% glycerol and used to seed cultures for the PhAGENT screens
(below). Genomic DNA was extracted from each mixture and barcode sequencing was
performed to measure the AG distribution in every replicate of all ECOR strain derived libraries.
The replicates were very highly correlated, and AG distributions in all libraries were confirmed to
be free of significant bottlenecking. These datasets are the reference libraries that serve as the
baseline for AG induction and phage infection experiments.
952
MOI calculation for liquid killing assays
Phage adsorption to host cells may be inefficient in very dilute cultures (10^{6-7} cfu/ml).
Therefore, the apparent multiplicity of infection (MOI-a) might be lower than the expected MOI
(MOI-e). We added 2x serial dilutions of a λvir lysate to a 1:100 dilution of a saturated culture of
E. coli BW25113 that had been allowed to acclimate at 37°C with shaking for 15 min. The
proportion of surviving cells was determined (relative to uninfected controls) after a 60-75 min
incubation (one phage replication cycle) by plating on LB agar. An apparent MOI value was
calculated using the Poisson distribution function with zero occurrences $MOIa = -ln\,(MOIe)$ for
each phage dilution, assuming that all infected cells would be killed by the virus (i.e. the
proportion of surviving cells would be the zero class in the Poisson distribution). The apparent
and expected MOI values were plotted against each other and the inverse of the slope (1/m) of
the line (MOI-a = m*MOI-e + 0) fit through the data was determined as the fold-change
reduction in MOI-a as compared to MOI-e for these experimental conditions. This fold-change
reduction was determined to be ~26x. Hence, all phage infection screens were targeted to an
MOI-a value of ~100 to ensure efficient phage adsorption and infection within the timeframe of a
single phage replication cycle.
969
AG Screen + Plaque assay confirmation + Data analysis
Thick cell mixtures from each of the three replicates of the various ECOR+AG libraries
were thawed on ice and diluted into LB supplemented with Mg^{2+}, Ca^{2+}, and 10 µg/ml gentamicin.
Typically, 1-5 µl of the library mixture was diluted in 2 ml broth to yield an initial OD_{600} of 0.1-
0.15. The cultures were immediately split x2, with one tube used to monitor OD_{600} values as the
experiments progressed, and the other left untouched for the screen. AG expression was
induced by the addition of 1 mM IPTG after acclimatization for 15 min at 37°C in a shaking
incubator and OD_{600} values were measured again for each ECOR+AG library at time of
induction. Libraries were then grown for ~80-120 min depending on the relative growth rates of
the various ECOR strains, to OD_{600} ~ 1.2-1.5. A 10^{-5} dilution of each late-log phase culture was
then plated on LB agar (with 10 µg/ml gentamicin and 0.5% glucose to repress AGs and avoid
fitness costs) to obtain a count of viable colony forming units (cfu). Cultures were immediately

diluted ~1:20 in fresh LB (with IPTG, Mg^{2+}, Ca^{2+}, and 10 µg/ml gentamicin) and 1 ml aliquots
were gently mixed with phage lysates at an MOI-e of ~30-100 (the actual MOI-e values were
recorded the next day when cfu counts were obtained). Parallel uninfected controls were
maintained for each experiment, even when the same ECOR+AG library was being tested
against different phages and the uninfected controls would be interchangeable. Infections were
allowed to proceed for 30-35 min for T-phages and 60-70 min for P1vir and λvir, i.e. one
replication cycle. We note that allowing sufficient time for the phages to complete cell lysis was
necessary as fast-growing cells were sometimes able to form colonies on LB agar despite
having been infected. These colonies had an uneven size distribution and a "splotchy"
appearance, perhaps due to ongoing infection on the plate that was nevertheless not able to
outpace colony formation. After the infection period was complete, the cultures were vortexed,
diluted 1:100, vortexed again, and 50 µl of the dilution was plated on 2 large (150 mm X 15 mm)
LB agar plates (with 10 µg/ml gentamicin and 0.5% glucose to repress AGs and avoid fitness
costs). The next day >>2,000-20,000 colonies were scraped from each infected and uninfected
set of plates. 5 µl of the thick resulting cell mixtures was reserved for genomic DNA extraction,
and the rest was frozen at -80°C in 20% glycerol as a backup. A full technical record of our
screens is provided in Supplementary File 4.
AG barcodes were amplified and sequenced as with the WM6026 donor library, with a
slight modification. The three biological replicate sets of ECOR+AG experiments were amplified
using JSW-SS-171 and different versions of primer JSW-SS-170 containing extra N's at the
start of the sequencing read. The first replicate was amplified with JSW-SS-170 (containing 5
N's), the second replicate with JBD-SS-254 (CGACGCTCTTCCGATCTNNNNNN
TGATGTCGTTGTTGCCATCG) with 6 N's, and the third replicate with JBD-SS-255
(CGACGCTCTTCCGATCTNNNNNNN TGATGTCGTTGTTGCCATCG) with 7 N's. This ensured
that reads from the three sets would not cross contaminate analyses of replicate experiments.
Amplicons were sequenced to the same depth as the baseline libraries on Illumina MiSeq and
NextSeq instruments in single-end format (100 bp) with two (8 bp) index reads.
We required exact matches for the 20 bp primer-derived invariant sequences flanking
the barcodes. These sequences were trimmed and the proportions of various AG-encoding
strains in the mixtures were determined using the prevalence of the corresponding 10 bp
barcode sequences. Barcode counts per AG were normalized by dividing each count by the
median of each experiment, which would represent neutral AG fitness. A log2 transformation
was subsequently applied to obtain fitness scores. Care was taken to sequence each library to
similar depth such that fitness scores across experiments were distributed in the same range.
Poorly represented AGs were identified in the baseline libraries as well as the infected and
uninfected experimental samples using a low read cutoff of 100, or if the log2 normalized
average read counts across the three replicates for any AG were >3x the median absolute
deviation below the median for that experiment. Reproducibility was determined using T-tests
across the three replicates, and we required a P-value of at least 0.2. The magnitude of the AG
fitness effect was determined by (a) subtracting the AG fitness scores in the uninfected library
from the infected libraries to assess whether AGs changed host sensitivity to phage infection,
and (b) subtracting the AG fitness scores in the baseline library from the uninfected libraries to
assess whether AGs change host fitness directly. AGs were selected for further study if they
were not poorly represented to begin with, produced reproducible effects across replicates, and
changed host fitness by at least 4-fold. Although we had targeted our experiments to phage-
host combinations where the host was mostly resistant to the phage in order to find counter-
defense AGs, a few hosts turned out to be more sensitive to certain phages in liquid infection
than suggested by agar-overlay plaque assays (perhaps due to abortive antiviral defenses). In
these cases, our platform was also able to robustly identify AGs that conferred a protective
advantage to the host (e.g. Imm and Cor, which both block phage DNA entry [11,12]). In practice

however, these effects can be very variable and the reproducibility filters typically needed to be
relaxed in order to find such superinfection exclusion AGs.
All counter-defense AGs were subsequently conjugated into the wild ECOR hosts where
they demonstrated an effect, and tested against all 8 phages (not just the ones that were used
in the screen with that host). In several hosts, the AG was found to additionally sensitize the
host to phages that were omitted in the screen. Not all the screen phenotypes could be
confirmed in agar-overlay plaque assays, possibly due to variations in cell growth between liquid
and solid media, or due to the requirement for successive infection cycles for plaque formation
that would render plaque assays less sensitive than single-infection cycle dynamics in our
screen. Nevertheless, further experiments were performed only in phage-host-AG combinations
that could be confirmed by plaque assay due to ease of follow up testing.
Conditional lethal AGs were identified directly from the heatmap of fitness scores, with a
subset selected for further experimentation using an arbitrary fitness cutoff of ~6-7 (~100-fold).
1045
Tn5 Dropout Screen + Data analysis to find phage exclusion systems
Tn5 whole-genome knockout strain libraries were constructed using the RB-TnSeq
donor library (pKMW7-derived strain APA766 from Dr. Adam Deutschbauer) conjugated with the
recipient ECOR strain in a biparental mating on LB plates supplements with 300 µM DAP. The
next day, cell mixtures were scraped and resuspended in 1 ml LB and 1000X dilutions of this
conjugation mixture were plated on 10 large (150 mm X 15 mm) LB plates (lacking DAP)
supplemented with 50 µg/ml kanamycin. Approximately 0.5-1 million colonies were scraped
taking care to break colonies apart on the surface of the plates, mixed very thoroughly,
aliquoted and stored at -80°C with 20% glycerol, with ~5-10 µl reserved for genomic DNA
extraction and reference library preparation.
RB-TnSeq allows the same mutant library to be challenged under various conditions
such as challenge by various phages, and simplifies the downstream sequencing-library
preparation by substituting barcode sequencing for all experimental conditions after initial
characterization of the reference library. To characterize the Tn5-knockout reference libraries in
the various ECOR strains, genomic DNA was extracted from 5-10 µl of the cell scrape and
prepared for sequencing using the NEBNext Ultra II FS DNA Library Prep Kit (NEB) with the
following modifications. 500 ng DNA was fragmented for 25 min according to manufacturer's
instructions. The adaptor ligation step was performed as directed albeit with a custom pre-
annealed oligo adaptor comprising of YL001 (/5Phos/GATCGGAAGAG/3ddC/) annealed to one
of YL002:005
(AGCGGCAATTTCACACAGGACAAGCAGAAGACGGCATACGAGAT NNNNNNNNN<u>AGTG</u>GTG
ACTGGAGTTCAGACGTGTGCTCTTCCGATC*T) where the 4-base underlined sequence
serves as a variable barcode unique to each oligo. The USER Enzyme steps post-ligation were
skipped and the bead-cleanup was performed using 40 µl and 20 µl of Sample Purification
Beads successively as instructed for library size selection in the 150-250 bp range. The PCR
amplification of transposon-genome junctions was performed using the cycling parameters
described in the kit for 11 cycles with Q5 Ultra II FS Master Mix (NEB) using primers YL006
(AGCGGCAATTTCACACAGGA) and JBD-SS-272/oAD493
(ACACTGGCAGAGCATTACGCCCT) instead of the kit-supplied primers. A second nested-PCR
enrichment step was performed for 11 cycles using the same cycling parameters as before,
using the entire bead-purified reaction product from the first step as a template with primers
YL009 (CAAGCAGAAGACGGCATACGAG) and JBD-SS-273:280
(AATGATACGGCGACCACCGAGATCTACAC TATAGCCT
ACACTCTTTCCCTACACGACGCTCTTCCGATCT NNNNNNN
GCAGGGATGTCCACGAGGTCTCT) where N_8 barcodes correspond to reverse-complemented
Illumina TruSeq HT D501-508 indexes. Bead-purified amplicons were quantified using the 1X

dsDNA HS kit with the Qubit 4.0 Fluorometer (Invitrogen) and sequenced on Illumina MiSeq and
NovaSeq instruments in single-end format (150 bp) with two (8 bp) index reads.
The first 100 bp of these reads contain transposon derived sequences and a DNA
barcode that uniquely identifies a particular Tn5 insertion. The last 50 bp of these reads contain
host-genome derived sequence that help determine the location of the transposon insertion and
hence construct a map between the Tn5 barcode and the insertion site (IS) in the genome.
Briefly, 20 bp barcodes and 16-50 bp IS sequences were trimmed from the sequencing reads.
To isolate the barcode and IS, we allowed 3 mismatches in the 23 bp invariant sequence
upstream and up to 6 mismatches in the 50 bp downstream of the barcode using a Hamming
distance function to compare the expected and actual sequences. Any reads failing these
sequence-fidelity requirements were discarded. Reads containing an IS that matched the Tn5
transposon vector in the donor strain were also discarded. The 4 Ns adjacent to the underlined
index sequences in YL002:005 were also sequenced as part of one 8 bp index read and served
as a tag to remove PCR duplicates. This allowed us to distinguish between sequencing "reads"
and DNA "fragments" present in the sample at the adaptor ligation step. Thus, reads containing
unique combinations of barcode, PCR tag, and IS were entered into the barcode/IS map for
each Tn5 library, each representing a distinct DNA fragment (but also machine errors in
sequencing, chimeric PCR amplicons, 3' truncations etc.). The filtered ISs were then mapped to
genes within their respective host genomes. As described previously[64], it is possible for
transposon barcodes to map to the genomes ambiguously. Barcodes associated with multiple
distinct ISs were retained only if their most frequently represented IS accounted for over 90% of
the fragments containing that barcode. We then counted the number of DNA fragments
recovered per gene and used this as a reference library for subsequent BarSeq experiments to
assay depletion of strains with mutations in specific genes upon phage infection.
Tn5 mutant strain libraries of the various ECOR strains were then challenged by various
phages in a procedure analogous to the ECOR+AG screens. Cultures of Tn5 libraries were
initiated with 100 µl of thick cell scrape mixture in 100 ml LB supplemented with 50 µg/ml
kanamycin, Mg^{2+}, and Ca^{2+}, to an initial OD_{600} value of 0.1-0.15. Cultures were grown at 37°C
with shaking for 80-90 min to an OD_{600} of 0.7-1. A 10^{-5} dilution of each late-log phase culture
was then plated on LB agar (with 50 µg/ml kanamycin) to obtain a count of viable colony forming
units (cfu). Cultures were immediately diluted ~1:10 in fresh LB (with Mg^{2+}, Ca^{2+}, and 50 µg/ml
kanamycin) and 1 ml aliquots were gently mixed with lysates of the appropriate phages at an
MOI-e of ~30-100 (the actual MOI-e values were recorded the next day when cfu counts were
obtained). Since the parental ECOR strains were typically quite resistant to phage infection to
begin with, only mutants with transposon-inactivated defense systems would be expected to
drop out upon infection. In practice however, some strains exhibited more pronounced mortality
upon prolonged phage infection in the Tn5 screens. In particular, in experiments with T2 and the
Tn5 library constructed in the Δ*wecA* strain of ECOR21 (which was generally more susceptible
to infection by T2 and other phages), MOI was titrated in three parallel infection screens (with
varying amounts of phage lysate added: 350 µl, 100 µl, and 25 µl) to empirically ensure specific
depletion of only the phage-sensitive mutants. Parallel uninfected controls were maintained for
each experiment, and each set of experiments was performed in biological duplicates. Infections
were allowed to proceed for 5 min at the benchtop without shaking, and 75 min at 37°C with
shaking. Since an interaction between the host and phage had already been identified in the
earlier AG screen, infections were not restriction to a single phage replication cycle, but allowed
to proceed further to achieve full depletion of the phage-sensitive Tn5-knockout mutants in the
pool. After the infection period was complete, the cultures were vortexed, diluted 1:10, vortexed
again, and 100 µl of the dilution was plated on 2 large (150 mm X 15 mm) LB agar plates (with
50 µg/ml kanamycin). We note that this dilution yields bacterial lawns and in our experience this
preserves diversity of surviving colonies without degrading the signal of dropouts by phage-
killing in the experiment. The next day, bacterial lawns were scraped from each infected and

uninfected set of plates. 5 µl of the thick resulting cell mixtures was reserved for genomic DNA
extraction, and the rest was frozen at -80°C in 20% glycerol as a backup. A full technical record
of our screens is provided in Supplementary File 6.
To amplify barcodes from the post-selection infected and uninfected Tn5 libraries, we
followed a modified BarSeq98 method[64], with two successive rounds of PCR. The transposon
barcode sequence was amplified by PCR using Q5 Ultra II FS Master Mix (NEB) with primers
JBD-SS-310 (CGACGCTCTTCCGATCTNNNNNNNN GATGTCCACGAGGTCTCT) and JBD-SS-
311 (ACTGACGCTAGTGCATCA GTCGACCTGCAGCGTACG). Template-matching regions in
the primers are underlined. Sequencing adaptors were attached in a second round of PCR
using amplicons from the first PCR round as the templates, with dual-indexed primer sets JSW-
SS-42:53 and JSW-SS-54:61 (sequences already provided in section describing AG screening).
Cycling conditions for round 1 were as follows: one cycle at 98°C for 4 min; two cycles at 98°C
for 30s, 63°C for 30s, and 72°C for 30s; 22 cycles at 98°C for 30s, and 72°C for 30s; and one
cycle at 72°C for 4 min. Conditions for round 2 were one cycle at 98°C for 1 min; two cycles at
98°C for 10 s, 64°C for 30 s, and 72°C for 10 s; 4 cycles at 98°C for 10 s, and 72°C for 20 s;
and one cycle at 72°C for 2 min. 0.5-1 ul of unpurified 1[st] round reaction product was used as
template for the 2[nd] round of PCR. Amplicons from the 2[nd] round were gel-purified by
electrophoresis (3% agarose gel, 4.2 V/cm, 2 hours) and quantified using the 1X dsDNA HS kit
with the Qubit 4.0 Fluorometer (Invitrogen). Amplicons were sequenced on Illumina MiSeq and
NovaSeq platforms in single-end format (150 bp) with two (8 bp) index reads.
DNA barcodes were trimmed from the reads and the proportions of various Tn5-
knockout mutants in the mixtures were determined using the prevalence of the corresponding
barcode sequence. Briefly, we allowed 3 mismatches in the 18 bp invariant sequence upstream,
and 4 mismatches in the 18 bp downstream of the transposon barcodes using a Hamming
distance function to compare the expected and actual sequences. The frequency of occurrence
of each barcode (representation number) was counted. Transposon barcodes were matched to
those found in the reference libraries, inferred to represent knockouts in the genes containing
the corresponding ISs, and the various barcode representation numbers were added together
for each gene. Cumulative transposon insertions per gene were then compared between the
infected and uninfected samples (with the reference library as a baseline to filter out poorly
represented, presumably essential genes) using the same data analysis procedure as for the
PhAGENT screen, albeit with two biological replicates instead of three.
1165
Tn5 Suppressor Screen + Data analysis to find abortive loci
Tn5 whole-genome knockout strain libraries were constructed using the RB-TnSeq
donor conjugated with the recipient ECOR strain as before, with a slight modification. The
recipient ECOR strain was equipped with the AG (as a single copy chromosomal Tn7 insertion
as usual) which triggers PCD, presumably by activating an abortive defense system. The trigger
AG was inserted prior to Tn5 library construction in order to minimize passages of the library
prior to screening and avoid potential bottlenecking due to fitness effects of the various Tn5
knockouts. Libraries were constructed as above, except that conjugation was carried out under
strict glucose repression to prevent PCD activation at this stage, and the conjugation mixture
scraped from the LB+DAP+glucose plates was then plated directly on LB plates containing 10
µg/ml gentamicin and 1 mM IPTG in addition to 50 µg/ml kanamycin. We typically needed to
plate the entire conjugation mixture scraped from five 100 mm LB+DAP+glucose plate onto five
150 mm LB+IPTG+Kan+Gm plate in order to obtain sufficient numbers of suppressor colonies
for characterization by high-throughput sequencing. Control samples were obtained by plating
1000-2000X dilutions on LB+glucose+Kan+Gm plates. Suitable dilutions were empirically
determined for the various ECOR hosts. The next day, suppressor colonies were scraped from
the large plates and stored at -80°C with 20% glycerol, with ~5-10 µl reserved for genomic DNA
extraction and sequencing. Genomic DNA was extracted and libraries were prepared and

sequenced akin to the treatment of the RB-TnSeq/NEBNext reference libraries detailed above.
There was no need for barcode sequencing since the screen identifies survivors instead of
dropouts, and each library was only characterized once after IPTG induction/selection for
suppressor mutants. Tn5 reference libraries were constructed from the data using the same
procedures for the baseline libraries constructed in the various naïve (without AGs) ECOR
strains above. The number of transposon insertions (i.e. the number of uniquely mapping ISs
from distinct DNA fragments) recovered per gene were used directly as a measure of survival of
mutants despite expression of PCD-triggering AGs. In practice, no statistical analysis was
necessary to identify the genetic loci responsible for PCD, as the signal-to-noise ratio was
sufficiently high to allow visual identification of hits when transposon insertion frequency was
plotted in control vs experimental samples (i.e. when the trigger AG was repressed by plating on
glucose-containing media, vs induced with IPTG).
1196
LPS gel electrophoresis
Lipopolysaccharide was prepared from liquid overnight cultures of wild *E. coli* strains
expressing the relevant AGs (LB + IPTG). Briefly, 50-100 μl of saturated culture was spun down
and cells were thoroughly resuspended in 100-200 μl of lysing buffer (62.5 mM Tris-HCl, pH6.8,
10% glycerol, 2% SDS, 4% ß-mercaptoethanol). We routinely supplement 1X Laemmli Buffer
(BioRad) with SDS and ßME to make lysing buffer. Samples were boiled at 100°C for 15 min,
and allowed to rest at room temperature for 15 min. 100 μg of Protease K were added to each
sample and protein was digested at 60°C for at least 1 hr. 15 μl of each sample were run onto a
12% polyacrylamide TGX Mini-Protean Precast gel (BioRad) at 200V for 35 min, and visualized
using the Pro-Q Emerald 300 LPS stain kit (Thermo Fisher) precisely according to
manufacturer's instructions.
1208
Allelic Exchange for *wecA* deletion
The R6K-origin pKEK2201 vector (from Prof. Karl E. Klose) carrying the desired genetic
modification (eg. a *wecA* deletion) flanked by 1000 bp homology arms was prepared by Gibson
assembly, and delivered as a suicide vector into wild *E. coli* strains via conjugation from the
WM6026 auxotrophic donor. Conjugation was performed using the same method used to deliver
AGs, albeit with biparental instead of triparental mating. Successful transconjugants (containing
a chromosomally integrated plasmid after the first homologous recombination event HR1) were
selected by incubation on LB plates supplemented with kanamycin. The second recombination
event HR2 to complete the allelic exchange was selected via *sacB*-mediated counterselection
on LB plates supplemented with 300 mM sucrose after overnight growth of HR1
transconjugants. Successful allele exchange was verified by PCR, and by LPS electrophoresis
to assay for the absence of O-antigen in the case of the *wecA* deletion.
1221
1222 **Live cell Microscopy**
1223 Cells were prepared for microscopy, immobilized under an LB agarose pad, and imaged
1224 as described previously[65]. Where necessary, AGs were pre-induced with 1 mM IPTG for 30 min
1225 immediately before harvesting for microscopy.
1226
Verification of abortive loci activity
Abortive/PCD systems identified by transposon screening of conditionally lethal AGs
were cloned by Gibson assembly using the NEBuilder HiFi DNA assembly master mix (NEB)
onto pBAD Myc/His A vectors along with their native promoters wherever possible. AGs that
trigger PCD systems (or appropriate controls) were conjugated into DH10ß/TOP10 strains
carrying pBAD-PCD plasmids using the Tn7 transposition method (above). Where the
restriction/PCD systems prevented conjugation of AGs, the order of transformations was
reversed – the AG was conjugated into naïve DH10ß/TOP10 cells first, and subsequently these

strains were transformed with the pBAD-PCD vectors. Successful transconjugants were grown
overnight with glucose to repress trigger AGs, and viability was tested the next day by spotting
10-fold serial dilutions of saturated cultures on LB agar plates with four additives: 1) Glucose
repression to assess background toxicity, (2) Arabinose induction of PCD systems alone to
assess fitness costs of over-expression of abortive components without the trigger AGs, (3)
IPTG to assess ability of AGs to trigger PCD without over-expression of abortive components
(i.e. under leaky expression conditions, or when expressed from the native promoters where
present), and (4) IPTG and Arabinose to assess ability of AGs to trigger PCD when abortive
components were also over-expressed.
1244
Affinity purification of AGs from wild *E. coli* hosts
FLAG-tagged AGs were conjugated into ECOR strains by Tn7-transposition (above)
according to Supplementary File 7. Each strain was grown overnight at 37°C in 2 mL of LB
supplemented with .5% glucose and 10 mg/mL gentamicin, then diluted 1:100 into 100 mL LB
supplemented with 10 mg/mL gentamicin and grown at 37°C for 30 minutes. The media was
then supplemented with 1 mM IPTG and grown for another 2.5 hours. "Popcorn" pellets were
collected by centrifugation for 30 minutes at 4,000 X g, resuspension in 100 µL of ice-cooled
lysis buffer [50 mM Tris pH 7.4, 150 mM NaCl, 1 mM EDTA, 1 mM $MgCl_2$, 0.5% NP40
(Research Products International), 125 U/mL Benzonase (Millipore), 1x protease inhibitor
cocktail (Roche, cOmplete ULTRA tablets, mini, EDTA-free), 0.5 mg/ml lysozyme (Fisher
Scientific)], followed by dropwise addition into liquid nitrogen and stored at -80°C. Popcorn
pellets were lysed by 10 cycles of cryomilling for 2 minutes at 12 counts per second in a SPEX
SamplePrep 6870D Freezer/Mill and stored at -80°C.
For each phage-host combination, 3 replicate lysates were reconstituted using 200 mg
of the cryomilled powder resuspended in 750 µL of lysis buffer. Lysates were cleared by
centrifugation at 16,000 X g at 4°C for 15 minutes. Anti-FLAG magnetic agarose beads (Pierce)
were washed on a magnet with 1 mL of ice-cooled IP buffer (50 mM Tris pH 7.4, 150 mM NaCl,
1 mM EDTA, 1mM $MgCl_2$) twice and resuspended to their original volume in IP buffer. Cleared
lysates were incubated with 30 µL of FLAG beads on an end-over-end tube rotator at 10 rpm at
4°C overnight. The supernatant was separated from the beads on a magnet. Beads were
washed once with 700 µL of .05% NP-40 and 1x protease inhibitor cocktail in IP, once with 700
µL of .05% NP-40 in IP, and 3 times with 700 µL of IP. Proteins were eluted by incubation with
25 µL of 100 µg/ml 3xFLAG peptide (Sigma) in 0.05% RapiGest (Waters Corp) in IP buffer on
an end-over-end tube rotator at 10 rpm at 4°C overnight. The eluate was separated from the
beads on a magnet. The elution step was then repeated with the same beads and fresh elution
buffer at room temperature for 1 hour. The eluates were combined and stored at -20°C.
1271
Mass spectrometry sample preparation and data acquisition
Samples were prepared for Mass Spectrometry (MS) as described previously in the
Filter-Aided Sample Preparation Protocol[66] with the following modifications. Before transferring
to an AcroPrep Advance 10k Omega plate, each eluate was resuspended in 100 µL of 8M Urea
(Thermo Scientific) dissolved in 50 mM ammonium bicarbonate (MP Biomedicals) and 5 mM
TCEP (Aldrich), followed by centrifugation to dryness. During alkylation of cysteine residues, 10
mM chloroacetamide (Thermo Scientific) was used instead of 10 mM iodoacetamide. Wash
steps prior to proteolysis were carried out with 200 µL of 20 mM ammonium bicarbonate
(instead of 100 µL). Proteolysis was performed without the addition of lysyl endopeptidase.
Proteolysis was quenched by addition of 10% trifluoroacetic acid (Thermo Scientific) to reach ≈
pH 2. Peptides were then purified over a BioPureSPE Mini 96-Well Plate (Nest Group). Briefly,
C18 columns were activated by washing once with 400 µL of acetonitrile (Fisher Scientific) and
twice with 400 µL of 0.1% formic acid (Thermo Scientific) in water solution, LC–MS grade
(Thermo Scientific). Samples were loaded and centrifuged at 1500 X g for 2 minutes. The bound

peptides were washed 3 times with 400 μL of 1% formic acid before elution with 150 μL of 50%
acetonitrile in 0.1% trifluoroacetic acid. Peptides were vacuum dried and stored at -80°C.
Samples were resuspended in 15 μl of 0.1% formic acid prior to loading for MS. The MS
acquisition was performed as described[67] with same MS parameters and LC configuration.
1290
Mass Spectrometry data analysis and statistical analysis
Raw files were processed in FragPipe[68] using the LFQ-MBR workflow with minor
modifications. The files were searched against a combined database of protein sequences from
ECOR1, ECOR3, ECOR15, ECOR21, ECOR22, ECOR66[62], and MG1655 with duplicate entries
removed and common contaminants added. Decoys were generated by pseudo-inversion.
Protein and peptide FDR was fixed to 1% and MBR was disabled. Cysteine carbamylation was
set as fixed modification while N-term acetylation, methionine oxidation and pyro-glu formation
were set as variable modification with a maximum of 3 modifications per peptide.
MaxLFQ intensities were used for statistical analysis[69]. Each triplicate set of host-AG
samples was quantile normalized against a triplicate set of samples from the same host
expressing orf74. For quantile normalization, all proteins in each sample were ranked according
to their MaxLFQ intensity. Next, the mean value of the intensities of proteins occupying a given
rank was calculated, then each protein's value was replaced with the mean value according to
its rank. Fold changes were calculated by taking the log2 ratio of average normalized values
between the test set and the control set. For each host-AG sample, all positively enriched
proteins were ranked according to their fold change. For each AG, proteins were assigned
"summed rank" values by calculating the sum of the ranks of each protein across all hosts
where that AG was tested. P-values were calculated by student's t-test on the normalized sets
of replicates from the test and control samples. Top enriched candidates for each AG were
selected based on their summed ranks and p-values.
1311
Ronin systems phylogenetic analysis
The *ronA* nucleotide sequence in the ECOR55 Ronin system was used to identify similar
genes by BLASTN. *RonA*-encoding bacterial genomes were downloaded using the entrez direct
e-utilities command line tools. Of 100 complete genomes retrieved, *ronA* homologs had been
annotated in 84. *hsdR* genes and other type I R-M components were identified using phmmer
and Defense Finder[70] respectively. Ronin-adjacent genetic architectures were visualized using
the PhamLite software suite[58]. All ronin-adjacent architectures are shown in HTML files with
mouse-over annotations for genes.

References

1. Koonin, E.V., and Wolf, Y.I. (2012). Evolution of microbes and viruses: a paradigm shift in evolutionary biology? Front Cell Infect Microbiol *2*, 119. 10.3389/fcimb.2012.00119.

2. Kristensen, D.M., Mushegian, A.R., Dolja, V.V., and Koonin, E.V. (2010). New dimensions of the virus world discovered through metagenomics. Trends Microbiol *18*, 11-19. 10.1016/j.tim.2009.11.003.

3. Bingham, R., Ekunwe, S.I., Falk, S., Snyder, L., and Kleanthous, C. (2000). The major head protein of bacteriophage T4 binds specifically to elongation factor Tu. J Biol Chem *275*, 23219-23226. 10.1074/jbc.M002546200.

4. Wang, H.C., Ho, C.H., Hsu, K.C., Yang, J.M., and Wang, A.H. (2014). DNA mimic proteins: functions, structures, and bioinformatic analysis. Biochemistry *53*, 2865-2874. 10.1021/bi5002689.

5. Loenen, W.A., and Murray, N.E. (1986). Modification enhancement by the restriction alleviation protein (Ral) of bacteriophage lambda. Journal of molecular biology *190*, 11-22. 10.1016/0022-2836(86)90071-9.

6. Nechaev, S., and Severinov, K. (2003). Bacteriophage-induced modifications of host RNA polymerase. Annual review of microbiology *57*, 301-322. 10.1146/annurev.micro.57.030502.090942.

7. Mizuno, C.M., Guyomar, C., Roux, S., Lavigne, R., Rodriguez-Valera, F., Sullivan, M.B., Gillet, R., Forterre, P., and Krupovic, M. (2019). Numerous cultivated and uncultivated viruses encode ribosomal proteins. Nat Commun *10*, 752. 10.1038/s41467-019-08672-6.

8. Bondy-Denomy, J., Pawluk, A., Maxwell, K.L., and Davidson, A.R. (2013). Bacteriophage genes that inactivate the CRISPR/Cas bacterial immune system. Nature *493*, 429-432. 10.1038/nature11723.

9. Chung, I.Y., Jang, H.J., Bae, H.W., and Cho, Y.H. (2014). A phage protein that inhibits the bacterial ATPase required for type IV pilus assembly. Proc Natl Acad Sci U S A *111*, 11503-11508. 10.1073/pnas.1403537111.

10. Shah, M., Taylor, V.L., Bona, D., Tsao, Y., Stanley, S.Y., Pimentel-Elardo, S.M., McCallum, M., Bondy-Denomy, J., Howell, P.L., Nodwell, J.R., et al. (2021). A phage-encoded anti-activator inhibits quorum sensing in Pseudomonas aeruginosa. Molecular cell *81*, 571-583 e576. 10.1016/j.molcel.2020.12.011.

11. Lu, M.J., and Henning, U. (1994). Superinfection exclusion by T-even-type coliphages. Trends Microbiol *2*, 137-139. 10.1016/0966-842x(94)90601-7.

12. Uc-Mass, A., Loeza, E.J., de la Garza, M., Guarneros, G., Hernandez-Sanchez, J., and Kameyama, L. (2004). An orthologue of the cor gene is involved in the exclusion of

temperate lambdoid phages. Evidence that Cor inactivates FhuA receptor functions.
Virology *329*, 425-433. 10.1016/j.virol.2004.09.005.

13.   Brussow, H., Canchaya, C., and Hardt, W.D. (2004). Phages and the evolution of
bacterial pathogens: from genomic rearrangements to lysogenic conversion. Microbiol
Mol Biol Rev *68*, 560-602, table of contents. 10.1128/MMBR.68.3.560-602.2004.

14.   Berg, P., and Mertz, J.E. (2010). Personal reflections on the origins and emergence of
recombinant DNA technology. Genetics *184*, 9-17. 10.1534/genetics.109.112144.

15.   Doudna, J.A., and Charpentier, E. (2014). Genome editing. The new frontier of genome
engineering with CRISPR-Cas9. Science *346*, 1258096. 10.1126/science.1258096.

16.   Makarova, K.S., Wolf, Y.I., Snir, S., and Koonin, E.V. (2011). Defense islands in
bacterial and archaeal genomes and prediction of novel defense systems. Journal of
bacteriology *193*, 6039-6056. 10.1128/JB.05535-11.

17.   Doron, S., Melamed, S., Ofir, G., Leavitt, A., Lopatina, A., Keren, M., Amitai, G., and
Sorek, R. (2018). Systematic discovery of antiphage defense systems in the microbial
pangenome. Science *359*. 10.1126/science.aar4120.

18.   Gao, L., Altae-Tran, H., Bohning, F., Makarova, K.S., Segel, M., Schmid-Burgk, J.L.,
Koob, J., Wolf, Y.I., Koonin, E.V., and Zhang, F. (2020). Diverse enzymatic activities
mediate antiviral immunity in prokaryotes. Science *369*, 1077-1084.
10.1126/science.aba0372.

19.   Pinilla-Redondo, R., Shehreen, S., Marino, N.D., Fagerlund, R.D., Brown, C.M.,
Sorensen, S.J., Fineran, P.C., and Bondy-Denomy, J. (2020). Discovery of multiple anti-
CRISPRs highlights anti-defense gene clustering in mobile genetic elements. Nat
Commun *11*, 5652. 10.1038/s41467-020-19415-3.

20.   Juhala, R.J., Ford, M.E., Duda, R.L., Youlton, A., Hatfull, G.F., and Hendrix, R.W.
(2000). Genomic sequences of bacteriophages HK97 and HK022: pervasive genetic
mosaicism in the lambdoid bacteriophages. Journal of molecular biology *299*, 27-51.
10.1006/jmbi.2000.3729.

21.   Comeau, A.M., Bertrand, C., Letarov, A., Tetart, F., and Krisch, H.M. (2007). Modular
architecture of the T4 phage superfamily: a conserved core genome and a plastic
periphery. Virology *362*, 384-396. 10.1016/j.virol.2006.12.031.

22.   Bondy-Denomy, J., Qian, J., Westra, E.R., Buckling, A., Guttman, D.S., Davidson, A.R.,
and Maxwell, K.L. (2016). Prophages mediate defense against phage infection through
diverse mechanisms. The ISME journal *10*, 2854-2866. 10.1038/ismej.2016.79.

23.   Tsao, Y.F., Taylor, V.L., Kala, S., Bondy-Denomy, J., Khan, A.N., Bona, D., Cattoir, V.,
Lory, S., Davidson, A.R., and Maxwell, K.L. (2018). Phage Morons Play an Important
Role in Pseudomonas aeruginosa Phenotypes. Journal of bacteriology *200*.
10.1128/JB.00189-18.

24.   Cumby, N., Davidson, A.R., and Maxwell, K.L. (2012). The moron comes of age.
1395   Bacteriophage *2*, 225-228. 10.4161/bact.23146.

1396   25.   Moffett, P. (2016). Using Decoys to Detect Pathogens: An Integrated Approach. Trends
1397   Plant Sci *21*, 369-370. 10.1016/j.tplants.2016.04.002.

1398   26.   Ochman, H., and Selander, R.K. (1984). Standard reference strains of Escherichia coli
1399   from natural populations. Journal of bacteriology *157*, 690-693. 10.1128/JB.157.2.690-
1400   693.1984.

1401   27.   Kulikov, E.E., Golomidova, A.K., Efimov, A.D., Belalov, I.S., Letarova, M.A.,
1402   Zdorovenko, E.L., Knirel, Y.A., Dmitrenok, A.S., and Letarov, A.V. (2021). Equine
1403   Intestinal O-Seroconverting Temperate Coliphage Hf4s: Genomic and Biological
1404   Characterization. Applied and environmental microbiology *87*, e0112421.
1405   10.1128/AEM.01124-21.

1406   28.   Allison, G.E., and Verma, N.K. (2000). Serotype-converting bacteriophages and O-
1407   antigen modification in Shigella flexneri. Trends Microbiol *8*, 17-23. 10.1016/s0966-
1408   842x(99)01646-7.

1409   29.   Vander Byl, C., and Kropinski, A.M. (2000). Sequence of the genome of Salmonella
1410   bacteriophage P22. Journal of bacteriology *182*, 6472-6481. 10.1128/JB.182.22.6472-
1411   6481.2000.

1412   30.   Newton, G.J., Daniels, C., Burrows, L.L., Kropinski, A.M., Clarke, A.J., and Lam, J.S.
1413   (2001). Three-component-mediated serotype conversion in Pseudomonas aeruginosa by
1414   bacteriophage D3. Mol Microbiol *39*, 1237-1247. 10.1111/j.1365-2958.2001.02311.x.

1415   31.   Losick, R. (1969). Isolation of a trypsin-sensitive inhibitor of O-antigen synthesis
1416   involved in lysogenic conversion by bacteriophage epsilon-15. Journal of molecular
1417   biology *42*, 237-246. 10.1016/0022-2836(69)90040-0.

1418   32.   Labrie, S.J., Samson, J.E., and Moineau, S. (2010). Bacteriophage resistance
1419   mechanisms. Nature reviews. Microbiology *8*, 317-327. 10.1038/nrmicro2315.

1420   33.   Weissborn, A.C., Liu, Q., Rumley, M.K., and Kennedy, E.P. (1994). UTP: alpha-D-
1421   glucose-1-phosphate uridylyltransferase of Escherichia coli: isolation and DNA sequence
1422   of the galU gene and purification of the enzyme. Journal of bacteriology *176*, 2611-2618.
1423   10.1128/jb.176.9.2611-2618.1994.

1424   34.   Genevaux, P., Bauda, P., DuBow, M.S., and Oudega, B. (1999). Identification of Tn10
1425   insertions in the rfaG, rfaP, and galU genes involved in lipopolysaccharide core
1426   biosynthesis that affect Escherichia coli adhesion. Arch Microbiol *172*, 1-8.
1427   10.1007/s002030050732.

1428   35.   Marolda, C.L., and Valvano, M.A. (1996). The GalF protein of Escherichia coli is not a
1429   UDP-glucose pyrophosphorylase but interacts with the GalU protein possibly to regulate

cellular levels of UDP-glucose. Mol Microbiol *22*, 827-840. 10.1046/j.1365-
2958.1996.01531.x.

36. Pierson, D.E., and Carlson, S. (1996). Identification of the galE gene and a galE homolog
and characterization of their roles in the biosynthesis of lipopolysaccharide in a serotype
O:8 strain of Yersinia enterocolitica. Journal of bacteriology *178*, 5916-5924.
10.1128/jb.178.20.5916-5924.1996.

37. Franklin, N.C. (1969). Mutation in gal U gene of E. coli blocks phage P1 infection.
Virology *38*, 189-191. 10.1016/0042-6822(69)90144-5.

38. Young, F.E. (1967). Requirement of glucosylated teichoic acid for adsorption of phage in
Bacillus subtilis 168. Proc Natl Acad Sci U S A *58*, 2377-2384. 10.1073/pnas.58.6.2377.

39. Fratamico, P.M., Briggs, C.E., Needle, D., Chen, C.Y., and DebRoy, C. (2003). Sequence
of the Escherichia coli O121 O-antigen gene cluster and detection of enterohemorrhagic
E. coli O121 by PCR amplification of the wzx and wzy genes. J Clin Microbiol *41*, 3379-
3383. 10.1128/JCM.41.7.3379-3383.2003.

40. Lehrer, J., Vigeant, K.A., Tatar, L.D., and Valvano, M.A. (2007). Functional
characterization and membrane topology of Escherichia coli WecA, a sugar-phosphate
transferase initiating the biosynthesis of enterobacterial common antigen and O-antigen
lipopolysaccharide. Journal of bacteriology *189*, 2618-2628. 10.1128/JB.01905-06.

41. Picton, D.M., Luyten, Y.A., Morgan, R.D., Nelson, A., Smith, D.L., Dryden, D.T.F.,
Hinton, J.C.D., and Blower, T.R. (2021). The phage defence island of a multidrug
resistant plasmid uses both BREX and type IV restriction for complementary protection
from viruses. Nucleic acids research *49*, 11257-11273. 10.1093/nar/gkab906.

42. Gordeeva, J., Morozova, N., Sierro, N., Isaev, A., Sinkunas, T., Tsvetkova, K.,
Matlashov, M., Truncaite, L., Morgan, R.D., Ivanov, N.V., et al. (2019). BREX system of
Escherichia coli distinguishes self from non-self by methylation of a specific DNA site.
Nucleic acids research *47*, 253-265. 10.1093/nar/gky1125.

43. Isaev, A., Drobiazko, A., Sierro, N., Gordeeva, J., Yosef, I., Qimron, U., Ivanov, N.V.,
and Severinov, K. (2020). Phage T7 DNA mimic protein Ocr is a potent inhibitor of
BREX defence. Nucleic acids research *48*, 5397-5406. 10.1093/nar/gkaa290.

44. Kruger, D.H., Reuter, M., Hansen, S., and Schroeder, C. (1982). Influence of phage T3
and T7 gene functions on a type III(EcoP1) DNA restriction-modification system in vivo.
Mol Gen Genet *185*, 457-461. 10.1007/BF00334140.

45. Millman, A., Bernheim, A., Stokar-Avihail, A., Fedorenko, T., Voichek, M., Leavitt, A.,
Oppenheimer-Shaanan, Y., and Sorek, R. (2020). Bacterial Retrons Function In Anti-
Phage Defense. Cell *183*, 1551-1561 e1512. 10.1016/j.cell.2020.09.065.

46. Rousset, F., Depardieu, F., Miele, S., Dowding, J., Laval, A.L., Lieberman, E., Garry, D.,
Rocha, E.P.C., Bernheim, A., and Bikard, D. (2022). Phages and their satellites encode

hotspots of antiviral systems. Cell host & microbe *30*, 740-753 e745.
10.1016/j.chom.2022.02.018.

47.   Ofir, G., Melamed, S., Sberro, H., Mukamel, Z., Silverman, S., Yaakov, G., Doron, S.,
and Sorek, R. (2018). DISARM is a widespread bacterial defence system with broad anti-
phage activities. Nat Microbiol *3*, 90-98. 10.1038/s41564-017-0051-0.

48.   Koonin, E.V., and Zhang, F. (2017). Coupling immunity and programmed cell suicide in
prokaryotes: Life-or-death choices. Bioessays *39*, 1-9. 10.1002/bies.201600186.

49.   Murray, N.E. (2000). Type I restriction systems: sophisticated molecular machines (a
legacy of Bertani and Weigle). Microbiol Mol Biol Rev *64*, 412-434.
10.1128/MMBR.64.2.412-434.2000.

50.   Penner, M., Morad, I., Snyder, L., and Kaufmann, G. (1995). Phage T4-coded Stp:
double-edged effector of coupled DNA and tRNA-restriction systems. Journal of
molecular biology *249*, 857-868. 10.1006/jmbi.1995.0343.

51.   Murphy, K.C. (2000). Bacteriophage P22 Abc2 protein binds to RecC increases the 5'
strand nicking activity of RecBCD and together with lambda bet, promotes Chi-
independent recombination. Journal of molecular biology *296*, 385-401.
10.1006/jmbi.1999.3486.

52.   Forti, F., Polo, S., Lane, K.B., Six, E.W., Sironi, G., Deho, G., and Ghisotti, D. (1999).
Translation of two nested genes in bacteriophage P4 controls immunity-specific
transcription termination. Journal of bacteriology *181*, 5225-5233.
10.1128/JB.181.17.5225-5233.1999.

53.   Ghisotti, D., Finkel, S., Halling, C., Deho, G., Sironi, G., and Calendar, R. (1990).
Nonessential region of bacteriophage P4: DNA sequence, transcription, gene products,
and functions. Journal of virology *64*, 24-36. 10.1128/JVI.64.1.24-36.1990.

54.   Lopatina, A., Tal, N., and Sorek, R. (2020). Abortive Infection: Bacterial Suicide as an
Antiviral Immune Strategy. Annu Rev Virol *7*, 371-384. 10.1146/annurev-virology-
011620-040628.

55.   Gao, L.A., Wilkinson, M.E., Strecker, J., Makarova, K.S., Macrae, R.K., Koonin, E.V.,
and Zhang, F. (2022). Prokaryotic innate immunity through pattern recognition of
conserved viral proteins. Science *377*, eabm4096. 10.1126/science.abm4096.

56.   Zhang, T., Tamman, H., Coppieters 't Wallant, K., Kurata, T., LeRoux, M., Srikant, S.,
Brodiazhenko, T., Cepauskas, A., Talavera, A., Martens, C., et al. (2022). Direct
activation of a bacterial innate immune system by a viral capsid protein. Nature *612*, 132-
140. 10.1038/s41586-022-05444-z.

57.   Johnson, M.C., Laderman, E., Huiting, E., Zhang, C., Davidson, A., and Bondy-Denomy,
J. (2022). Core Defense Hotspots within *Pseudomonas aeruginosa* are a

consistent and rich source of anti-phage defense systems. bioRxiv,
2022.2011.2011.516204. 10.1101/2022.11.11.516204.

58. Johnson, M.C. (2023). PhamLite. https://console.latch.bio/explore/61249/info.

59. Steinegger, M., and Soding, J. (2017). MMseqs2 enables sensitive protein sequence
searching for the analysis of massive data sets. Nat Biotechnol *35*, 1026-1028.
10.1038/nbt.3988.

60. Jain, C., Rodriguez, R.L., Phillippy, A.M., Konstantinidis, K.T., and Aluru, S. (2018).
High throughput ANI analysis of 90K prokaryotic genomes reveals clear species
boundaries. Nat Commun *9*, 5114. 10.1038/s41467-018-07641-9.

61. Peters, J.M., Koo, B.M., Patino, R., Heussler, G.E., Hearne, C.C., Qu, J., Inclan, Y.F.,
Hawkins, J.S., Lu, C.H.S., Silvis, M.R., et al. (2019). Enabling genetic analysis of diverse
bacteria with Mobile-CRISPRi. Nat Microbiol *4*, 244-250. 10.1038/s41564-018-0327-z.

62. Patel, I.R., Gangiredla, J., Mammel, M.K., Lampel, K.A., Elkins, C.A., and Lacher, D.W.
(2018). Draft Genome Sequences of the Escherichia coli Reference (ECOR) Collection.
Microbiol Resour Announc *7*. 10.1128/MRA.01133-18.

63. Hansen, L. (2021). SeqDiff. https://github.com/hansenlo/SeqDiff.

64. Wetmore, K.M., Price, M.N., Waters, R.J., Lamson, J.S., He, J., Hoover, C.A., Blow,
M.J., Bristow, J., Butland, G., Arkin, A.P., and Deutschbauer, A. (2015). Rapid
quantification of mutant fitness in diverse bacteria by sequencing randomly bar-coded
transposons. mBio *6*, e00306-00315. 10.1128/mBio.00306-15.

65. Li, Y., Guan, J., Hareendranath, S., Crawford, E., Agard, D.A., Makarova, K.S., Koonin,
E.V., and Bondy-Denomy, J. (2022). A family of novel immune systems targets early
infection of nucleus-forming jumbo phages. bioRxiv, 2022.2009.2017.508391.
10.1101/2022.09.17.508391.

66. Fossati, A., Frommelt, F., Uliana, F., Martelli, C., Vizovisek, M., Gillet, L., Collins, B.,
Gstaiger, M., and Aebersold, R. (2021). System-Wide Profiling of Protein Complexes
Via Size Exclusion Chromatography-Mass Spectrometry (SEC-MS). Methods in
molecular biology *2259*, 269-294. 10.1007/978-1-0716-1178-4_18.

67. Gordon, D.E., Jang, G.M., Bouhaddou, M., Xu, J., Obernier, K., White, K.M., O'Meara,
M.J., Rezelj, V.V., Guo, J.Z., Swaney, D.L., et al. (2020). A SARS-CoV-2 protein
interaction map reveals targets for drug repurposing. Nature *583*, 459-468.
10.1038/s41586-020-2286-9.

68. Kong, A.T., Leprevost, F.V., Avtonomov, D.M., Mellacheruvu, D., and Nesvizhskii, A.I.
(2017). MSFragger: ultrafast and comprehensive peptide identification in mass
spectrometry-based proteomics. Nature methods *14*, 513-520. 10.1038/nmeth.4256.

69.   Cox, J., Hein, M.Y., Luber, C.A., Paron, I., Nagaraj, N., and Mann, M. (2014). Accurate
1539   proteome-wide label-free quantification by delayed normalization and maximal peptide
1540   ratio extraction, termed MaxLFQ. Mol Cell Proteomics *13*, 2513-2526.
1541   10.1074/mcp.M113.031591.

1542   70.   Tesson, F., Herve, A., Mordret, E., Touchon, M., d'Humieres, C., Cury, J., and Bernheim,
1543   A. (2022). Systematic and quantitative view of the antiviral arsenal of prokaryotes. Nat
1544   Commun *13*, 2561. 10.1038/s41467-022-30269-9.
1545

A. Core vs. Accessory genes in a hypothetical phage cluster

phage 1
phage 2
phage 3
phage 4
phage 5
phage 6

Accessory Genes

B. Algorithm to find accessory regions without a priori genome clustering

choose any two genes

3 genomes contain both genes from chosen pair and become a cluster

A candidate accessory region between and

all gene families from all phage genomes

try every possible pairwise combination of genes

All possible accessory regions

... etc.

filter and rank accessory regions

high scoring

low scoring

C. Sample accessory region found by the algorithm

1 kb   2 kb   3 kb   4 kb

SPFM7
SPFM6
SPFM5
SPFM20
SPFM15
SPFM14
SPFM12
SPFM10

putative virion structural protein

accessory region

conserved hypothetical protein

D. Characteristics of ARs and AGs

AGs   # phages

10000
8000
6000
4000
2000

1200
1000
800
600

ARs ranked by score

AGs

4000
3000
2000
1000

51-100
101-150
<50
151-200
201-300
>300

Protein length (aa)

AR quintiles by score

ARs

1000
500
0

11-20
<10
21-30
31-50 >50

AGs

phages

400
200
0

1-5
6-10
11-15
16-20 >20

AGs

phages

600
400
200
0

1-5
6-10
11-15
16-20 >20

ARs

AR quintiles by score

E. PhAGENT screen to assay AG phenotypes

Conjugate barcoded AG library into host

N = number of genes

Scrape >>10N colonies from each conjugation

x3 replicates per host strain

Challenge each replicate with phages one-by-one

phage 1
phage 2
phage 8

Three parallel measurements for each infection

AG only   AG   phage

t = 0   t = 30-60 min

Scrape >>10N colonies for each measurement

High-throughput sequencing

to quantify barcodes

A. Accessory genes with counter-defense phenotypes in any host

Phage - Host combination (y-axis):
MG1655-λvir, C990-T4, ECOR1-P1vir, ECOR3-P1vir, ECOR3-T2, ECOR3-T6, ECOR10-T2, ECOR10-T4, ECOR10-T6, ECOR13-T5, ECOR15-T2, ECOR15-T6, ECOR18-P1vir, ECOR18-T6, ECOR19-P1vir, ECOR19-T6, ECOR21-P1vir, ECOR21-T2, ECOR21-T4, ECOR22-P1vir, ECOR22-T2, ECOR22-T4, ECOR22-T6, ECOR24-P1vir, ECOR24-T6, ECOR51-P1vir, ECOR51-T4, ECOR53-P1vir, ECOR53-T4, ECOR53-T6, ECOR55-P1vir, ECOR55-T4, ECOR59-P1vir, ECOR59-T4, ECOR59-T6, ECOR61-P1vir, ECOR61-T2, ECOR61-T3, ECOR61-T4, ECOR61-T5, ECOR61-T6, ECOR61-T7, ECOR66-P1vir, ECOR66-T4, ECOR66-T6K

log(2) fitness score (Infected vs Uninfected) — color scale from ~ -10 to 2

AGs (x-axis): ocr, aoa1, 0009, gnarl1, 0050, gnarl2, 0068, 0074, ral, gnarl3, 0101, 0107, abc1, 0126, lplI, 0148, 0151, 0153, ardA

AGs that had a counter-defense effect in at least one host strain

B. Whole genome Tn5-knockout screening to find phage exclusion mechanisms

transposon insertion inactivates defense system

virus kills host

ECOR1 Tn5 library infected by T5 — Fitness (Infected) vs Fitness (Uninfected); LPS O-antigen pathway

ECOR3 Tn5 library infected by T5 — LPS O-antigen pathway

ECOR22 Tn5 library infected by T5 — LPS O-antigen pathway

C. Plaque assays show very strong sensitization to phage T5 by AGs 48, 63, 92 (gnarl 1, 2, 3)

Accessory gene:

gnarl1 — repressed / induced
gnarl2 — repressed / induced
gnarl3 — repressed / induced

ECOR 1   ECOR 3   ECOR 22

D. Gel electrophoresis of LPS from wild E. coli expressing AGs 48, 63, 92 (gnarl 1, 2, 3)

WT 48 63 92 (ECOR 1)   WT 48 92 (ECOR 3)   WT 48 63 92 (ECOR 22)

ECOR 1   ECOR 3   ECOR 22

E. Heatmap of enriched host-proteins upon AP-MS of broad-spectrum phage AGs

proteins (y-axis)

color scale: 0 to 25

gnarl1, gnarl2, gnarl3, abc1, orf 126, lplI, orf 148

F. Over-expression of galU reverses T5-sensitization by gnarl3

no Tn7 insert $P_{L lacO-1}$ — gnarl3

Empty vector — ECOR1, ECOR3, ECOR22
P_{BAD} — galU — ECOR1, ECOR3, ECOR22
P_{BAD} — galE — ECOR1, ECOR3, ECOR22
P_{BAD} — galF — ECOR1, ECOR3, ECOR22
P_{BAD} — wecB — ECOR1, ECOR3, ECOR22

A. Whole genome Tn5-knockout screening in ECOR21 with various phages

B. ECOR21 might block T2 at two levels simultaneously

C. O-antigen removed in ECOR21::ΔwecA

D. Tn5 screens after removing O-antigen layer

E. BREX-operon configurations tested against T-even phages with various inhibitors

F. GmrSD is sufficient to confer resistance against T-even phages

G. T4 naturally encodes IpII and does not require additional IpII to overcome basal levels of GmrSD restriction

A. Accessory genes with conditional lethal effects in any host

Host (rows): MG1655, C990, ECOR1, ECOR3, ECOR10, ECOR13, ECOR15, ECOR18, ECOR19, ECOR21, ECOR22, ECOR24, ECOR51, ECOR53, ECOR55, ECOR59, ECOR61, ECOR66

AGs that were conditionally lethal in at least one host strain (columns): 0008, 0145, aoa1, ral, 0151, gnarl2, 0024, ocr, gnarl3, abc1, ardA, 0148, 0184, 0062, 0076, arn, 0013, 0066

log(2) fitness score (Uninfected vs Baseline): 0, -2, -4, -6, -8, -10, -12

B. Fitness cost of conditional lethal interactions in native hosts

- MG1655 + orf145
- ECOR15 + Abc1
- ECOR22 + Ocr
- ECOR55 + Ocr
- ECOR55 + Aoa1
- ECOR55 + ArdA
- ECOR59 + orf184
- ECOR61 + orf24
- ECOR66 + Ocr
- ECOR66 + Aoa1
- ECOR66 + ArdA

Accessory gene: repressed — induced

C. Whole genome Tn5 suppressor screens

same AG expressed in multiple strains of same species → strain with abortive system triggered by AG dies

transposon insertion inactivates abortive system → conditional lethal phenotype is abrogated

Proportion of Tn5 insertions per gene in surviving cells

ECOR22 + Ocr

Trigger (Ocr) induced vs Trigger (Ocr) repressed

mod, Type III R-M, res

D. Type III R-M from ECOR22 kills cells upon Ocr expression

ECOR22 wild host: Ocr, Aoa1, ArdA

P_BAD — type III mod — type III res

TOP10: Ocr, Aoa1, ArdA

repressed — induced

E. λvir grown on nuclease-dead (res*) mutant can evade EcoR22III restriction

P_BAD — type III mod — type III res

TOP10: λvir, λvir*

P_BAD — type III mod — type III res*

TOP10: λvir, λvir*

F. EcoR22III restriction endonuclease site is partially responsible for Ocr-induced PCD

- P_BAD — type III mod — type III res
- P_BAD — type III mod — type III res*
- P_BAD — type III mod
- P_BAD — type III res

Ocr repressed — Ocr induced

G. Hypothetical model of the same R-M system mounting both targeted and abortive defenses

type III R-M — type III R-M blocks infection

Ocr-encoding phage or MGE — Ocr — Ocr-modified type III R-M kills host

A. Ronin system from ECOR55 kills cells upon aoa1 or ardA induction

repressed   induced

Ocr
Aoa1
ArdA

B. RonA contains a predicted transmembrane region

TMHMM posterior probability

Protein sequence (61 aa)

transmembrane   inside   outside

C. RonA is toxic upon overexpression but not with hsdMS present

no ArdA

D. RonA is necessary for ArdA-induced PCD activity

ArdA repressed   ArdA induced

E. Ronin activation causes growth arrest but not cell lysis

no Tn7 insert

$P_{L\ lacO-1}$ ArdA

0   30   60   90   120   min

F. RonA-associated type I R-M systems (ronin) lack the restriction endonuclease

genomes with *ronA* (84)

ronA linked to *hsdMS* (84)

DUF4102 integrase upstream (58)

lacking full integrase upstream (26)

	present in *E. coli*	full type I R-M anywhere
	0	55
	26	6

DUF4102 integrase

ronin

hsdR   DUF262   hsdS   hsdM

Type I R-M

ronA hsdS hsdM

ronin

BREX operon

G. Hypothetical model of RonA "uncaging"

RonA is "caged" by type I M-S

ArdA-encoding phage or MGE

ArdA

"uncaged" RonA kills host

A. P4 satellite prophage from ECOR15 arrests TOP10 upon abc1 induction

$P_{L\ lacO-1}$ Abc1

repressed   induced

B. gop-beta-cII system from ECOR59 arrests TOP10 upon orf184 induction

$P_{L\ lacO-1}$ orf 184

gop beta cII

P_{BAD} gop beta cII

repressed   induced

C. Novel prophage-encoded system from ECOR61 arrests TOP10 upon orf24 induction

$P_{L\ lacO-1}$ orf 24

lcd DNA primase

P_{BAD} lcd DNA primase

DNA primase

repressed   induced

D. Parallels between Immune Guardianship by ARIAs and Superinfection Exclusion

phage factor

superinfection exclusion system

prophage/MGE

anti-immune factor

guardian immunity

frontline defenses

genome

A. Phage susceptibility of ECOR strains

B. Phage-host combinations with confirmed counter-defense phenotypes

counter-defense phenotype

...not detected

...detected by screen

...confirmed by plaque assay (>10²-fold)

n   EOP = n

n   EOP = n, and single plaques visible

selected for PhAGENT screen

attempted but screen could not be performed

C. Defense systems in ECOR strains

A. R-M inhibitors trigger abortive defenses in wild E. coli

B. R-M inhibition by AGs in MG1655

C. Effect of Ocr F54V mutation on ARIA activity

A. Ronin upstream region is required for PCD activity

235 bp | hsdS | hsdM

100 bp | hsdS | hsdM

ArdA repressed   ArdA induced

B. Ronin upstream region is not toxic when overexpressed on its own

P_BAD | 235 bp

no ArdA

C. Over-expression of Ronin with truncated upstream region

P_BAD | 100 bp | hsdS | hsdM   P_L lacO-1 | ArdA

repressed

induced