Neural Geometry from Mixed Sensorimotor Selectivity for Predictive Sensorimotor Control

Running Title: Effect of Target Motion on Reach Directionality in Motor Cortex

Yiheng Zhang[1,2,3,4] Yun Chen[1,2,3,4] Tianwei Wang[1,2] and He Cui[1,2]*

1 Center for Excellence in Brain Science and Intelligence Technology, Institute of Neuroscience,
Chinese Academy of Sciences, Shanghai 200031, China

2 Chinese Institute for Brain Research, Beijing, 102206, China

3 University of Chinese Academy of Sciences, Beijing 100049, China

4 equal contributions

* Corresponding author: He Cui

Email: hecui@cibr.ac.cn

Abstract

Although recent studies suggest that activity in the motor cortex, in addition to generating motor outputs, receives substantial information regarding sensory inputs, it is still unclear how sensory context adjusts the motor commands. Here, we recorded population neural activity in the motor cortex via microelectrode arrays while monkeys performed flexible manual interceptions of moving targets. During this task, which requires predictive sensorimotor control, the activity of most neurons in the motor cortex encoding upcoming movements, was influenced by ongoing target motion. Single-trial neural states at the movement onset formed staggered orbital geometries, suggesting that target motion modulates peri-movement activity in an orthogonal manner. This neural geometry was further evaluated with a representational model and recurrent neural networks (RNNs) with task-specific input-output mapping. We propose that the sensorimotor dynamics can be derived from neuronal mixed sensorimotor selectivity and dynamic interaction between modulations.

Introduction

The motor cortex, a central brain region generating motor commands, is widely known for its relation to movement kinetics (Evarts, 1968; Kalaska et al., 1989) and kinematics(Georgopoulos et al., 1982; Wang et al., 2022). More than this, the motor cortex also carries substantial sensory information and is significantly involved in a variety of sensorimotor processes. Many studies have reported that the motor cortex reveals early visuomotor responses for action selection (Cisek and Kalaska, 2005; Wang et al., 2019) and motor planning (Hatsopoulos and Amit, 2012; Pesaran et al., 2006; Rao and Donoghue, 2014), anticipative activity for complex rules and sequences (Lu and Ashe, 2005; Wang et al., 2024; Zimnik and Churchland, 2021), as well as fast closed-loop modulation for somatosensory and visual feedback control (Omrani et al., 2016; Sobinov and Bensmaia, 2021; Stavisky et al., 2017; Suway and Schwartz, 2019; Tkach et al., 2007). However, it is still unclear how moving targets requiring forward control affect peri-movement activity in the motor cortex.

Nowadays, the dynamical systems perspective provides a population view to interpret mixed coding for task variables and rules (Mante et al., 2013; Rigotti et al., 2013). With respect to motor control, the preparatory population activity not only sets initial states to seed the motor generation (Churchland et al., 2012; Churchland and Shenoy, 2024; Vyas et al., 2020), but also presumably contains redundant information – that should be 'held' rather than 'released' to trigger muscles – in an output-null space (Kaufman et al., 2014). Therefore, we hypothesize that the potential

sensory modulation, which results from target motion and does not intervene the motor output,

functions in orthogonal dimensions of output-potent neural states.

To investigate the mixed selectivity and neural dynamics shaped by concurrent sensorimotor

signals, we recorded population activity in the primary motor cortex (M1) from monkeys

performing a flexible manual interception task. Unlike previous studies constraining interception at

a fixed location (Merchant et al., 2004a, 2004b), our task demands predictive spatiotemporal

mappings to displace a body effector to a trial-varying location. We found that the activity of most

neurons was jointly tuned to both reach direction and target motion via directional selectivity

shifts, gain modulations, offset adjustments, or their combinations. Strikingly, such mixed

sensorimotor selectivity exists throughout the entire trial, in contrast to the gradient of sensory-to-

motor tuning from cue to movement epochs in posterior parietal cortex (PPC) that we recently

reported (Li et al., 2022, 2018). Principal component analysis (PCA) on the neural population

revealed a clear orbital neural geometry in low-dimensional space at the movement onset: The

neural states were distributed in reach-direction order and formed ring-like structures whose

slopes were determined by target-motion conditions. This target-motion effect is maintained

independently of hand speed. Neuronal simulation indicates that these characteristics of neural

population dynamics could be derived from the mixed sensorimotor selectivity of single neurons.

Recurrent neural networks (RNNs) trained with proper input-output mappings offer insights into

the relationship between neuronal modulation and neural geometry in a dynamical system. We

propose that this sensory modulation occurs at both single-neuron and population levels as a

general element of neural computations for predictive sensorimotor control.

Results

Mixed tuning of M1 single neurons during flexible manual interception task

Three monkeys (*Macaca mulatta,* C, G, and D, male, weight 7 to 10 kg) were trained to perform a delayed manual interception task (Figure 1A), which was modified from the task utilized in our recent studies (Li et al., 2022, 2018). To initiate a trial, the monkey needed to hold the center dot of a standing touch screen for 600 ms. Then, a peripheral target appeared, either stationary or rotating around the center dot at a constant angular velocity. The monkey was required to wait during a randomized delay (400-800 ms) until the central dot turned dark (GO signal) and then to immediately reach to the target. Once the monkey touched the screen (Touch) again, the target stopped and another dot showed the touched location, in red for a successful interception or in blue for a failure. The error tolerance between the target and the touched location (reach endpoint) was 3 cm. There were five target-motion conditions, consisting of clockwise (CW) conditions - 240 °/s and -120 °/s, counterclockwise (CCW) conditions 120 °/s and 240 °/s, along with 0 °/s (static) condition. We define the magnitude and direction of the target's velocity as TV_{mag} (for example, 0 °/s, 120 °/s, and 240 °/s) and TV_{dir} (namely CCW and CW), respectively. These target-motion conditions were interleaved trial by trial, and the initial location of the target was random. The reach endpoints of a well-trained monkey distributed uniformly around the circle (Figure 1B, Rayleigh's test: p=0.36; data from monkey C, 772 correct trials in one session). Because the reach direction was defined as the angle of the reach endpoint, for simplicity, we divided the circular space equally into eight sectors (45° per each) and grouped trials according to the eight reach directions and five target-motion conditions.

We recorded neural data with Utah arrays from monkeys C, G, and D (implanted sites are shown in Figure 1C, and all datasets are listed in Table S1) and hand trajectories from monkeys C and G (Figure S1), during the interception task. The hand trajectory was launched to the final interception location. The temporal profiles of hand speed were unimodal bell-like curves and similar across target-motion conditions (correlation coefficients is 0.97 ± 0.05 for monkey C and 0.99 ± 0.02 for monkey G, mean ± sd.).

Notably, we found that neuronal directional tuning during peri-movement period (MO ± 100 ms, MO for movement onset) was modulated by target motion, according to our statistical criteria, mainly in three ways: preferred direction (PD) shift, gain modulation, and offset addition (Table 1, Figure S2-S4; see Methods). We determined these modulations on the basis of the classical cosine tuning model (Georgopoulos et al., 1982) and several previous studies (Bremner and Andersen, 2012; Pesaran et al., 2010; Sergio et al., 2005). Specifically, PD-shift neurons had their PDs shifted in moving-target conditions compared to the static-target condition. As illustrated by the example neuron (Figure 1D, *PD shift*), whose PDs corresponding to CCW conditions (red and yellow) and CW conditions (blue and purple) exhibited obvious differences, TV_{dir} rather than

TV_{mag} dominated this modulation. Gain-modulation neurons exhibited reach-direction tuning

multiplied by target velocity: while their directionality remained invariant, the neuronal responses

at PD differed across target-motion conditions. This modulation was dominated by TV_{dir} as well.

The turning curves of the example neuron (Figure 1D, *Gain*) had higher responses at PD in CW

conditions (blue and purple) than in the others (green, yellow, and red), indicating a varying

tuning depth for reach direction. Neurons with addition modulation underwent changes of offset

activity induced by target velocity (both TV_{dir} and TV_{mag}). As shown by the example neuron

(Figure 1D, *Addition*), this effect was roughly the same in all reach directions.

The activity of these example neurons could be well described by PD shift, gain, and additive

models (Figure 1E; see Methods), respectively. Nevertheless, it was difficult to classify all

neurons with mixed sensorimotor selectivity into one of these three groups exclusively, because

many of them experienced a mixture of two or three of above modulations. We found that the

adjusted R^2 of a full model (0.55 ± 0.24, mean ± sd.) can be higher than that of the PD shift (0.47

± 0.24), gain (0.46 ± 0.22), additive (0.41 ± 0.26), and simple models (only reach direction, 0.34 ±

0.25) for three monkeys (1162 neurons, rank-sum test, one-tailed, p<0.01, Figure S5). These

target-motion modulations on neuronal directionality suggested the participation of sensory

signals in shaping neural dynamics during interception execution.

Figure 1. The flexible manual interception task and example neurons

A Task diagram. In each trial, the monkey first holds the center dot for 600 ms (Hold), then a target appears (Target on), after 400-800 ms delay the center dot turns dark (GO), immediately the monkey moves its hand to reach to the target (Touch). The movement time (from GO to Touch) is required to be within 800 ms, otherwise the trial will be aborted. The feedback dot, which is presented at the touched location of the screen, will be in red for success or in blue for failure.

B Distribution of touch endpoints. Left panel shows fifteen reaching-up example trials in five target-motion conditions, three trials in each condition. The squares mark the touch endpoints, while the circles and triangles are the target onset and stop location. The five target-motion conditions (-240°/s, - 120°/s, 0°/s, 120°/s, and 240°/s) are indicated in five colors (purple, blue, green, yellow, and red). Target onset location is randomly distributed. Right panel shows the

touch endpoints of all trials, each point represents a trial, colored according to target-motion

conditions. The distribution was uniform around the circle (monkey C 772 trials, Rayleigh's test,

p=0.36).

C Implanted locations of microelectrode array in the motor cortex of the three well-trained

monkeys. Neural data were recorded from the cortical regions contralateral to the used hand. AS,

arcuate sulcus; CS, central sulcus.

D Three example neurons with PD shift, gain modulation, and offset addition. The peri-stimulus

time histograms (PSTH) show the activity of example neurons when monkeys reached to upper

areas in five target-motion conditions. The solid lines represent the trial-averaged firing rates, the

colored shadow represents the standard error. The gray shadow indicates the time window

between MO-100 ms and MO+100 ms.

E The directional tuning curves of the three example neurons with PD shift, gain, and addition

modulation around movement onset (MO ± 100 ms, adjusted R^2: 0.70, 0.84, and 0.60). Dots and

bars denote the average and standard error of firing rates, colored according to target-motion

conditions.

Table 1 Ratio of mixed selectivity neurons around movement onset

Units across sessions	Gain (G)	PD shift (S)	Addition (A)	None
Monkey C, 7 sessions, N=84.9±15.9	46.9±15.9%	79.0±8.9%	61.9±14.5%	6.5±6.2%
Monkey G, 4 sessions, N=97.5±17.7	29.3±4.9%	51.6±29.1%	25.9±8.3%	25.5±15.8%
Monkey D, 4 sessions, N=44.5±7.5,	45.6±13.8%	74.4±8.8%	49.1±9.1%	10.8±5.4%
RNN, N=200, 100 models	46.5±4.6%	40.7±4.2%	57.9±8.9%	14.2±4.2%

Single neurons were classified by modulation patterns (more details in **Methods**). The first

column shows the subject, the number of recording sessions, and the number of isolated units

(mean±sd.). Other columns show the ratio of units with any target-velocity modulations in all

active units. 'None' means the units were not specifically tuned by either reach direction nor

target velocity.

Coding of sensory and motor information in neural populations

To quantify target motion information embodied in neural response of the motor cortex, we

performed a series of decoding analyses on the neural data from monkey C (n=95, 772 correct

trials). To begin with, we trained a support vector machine (SVM) classifier for target-motion

conditions (chance level: one in five) and another for reach directions (chance level: one in eight;
see Methods) on neural data. As Figure 2A shows, the decoding accuracy of target-motion
condition increased quickly and peaked at over 70% around GO, while the decoding accuracy of
reach direction climbed in an approximately linear manner and reached a plateau of about 80%
before MO. These results were stable on the data of the other two monkeys and the pseudo-
population of all three monkeys (Figure S6) and reconfirmed by the continuous decoding results
with support vector regressions (Figure S7A), suggesting that target motion information existed in
M1 throughout almost the entire trial. According to the demixed PCA (dPCA) (Kobak et al., 2016)
results, the reach-direction components occupied high ranks; although the target-velocity
components explained few variance, the interaction components were non-negligible (Figure S8).
This implies that the target motion information was intertwined with reach-direction information,
rather than being processed independently and in parallel.
Then, we performed another decoding analysis to probe the potential interaction between
reach direction and target velocity during execution period. We trained a reach-direction decoder
(chance level: one in eight) to check if the decoder of one certain target-motion condition could be
transferred to other conditions (Figure 2B left). It turned out that the performance of the
transferred decoder deteriorated more significantly for $TV_{dir.}$ (CCW *vs.* CW, mean ± sd. of
accuracy 0.26±0.06), compared with that for TV_{mag} (120 *vs.* 240, 0.50±0.06, paired t-test, p<0.01)
and for target's state (static *vs.* motion 0.55±0.06, paired t-test, p<0.01). This result suggests that
the coding of reach direction was rather sensitive to $TV_{dir,}$ but contained similarities for static and
moving targets. We also compared the neural coding rules across different reach-direction
conditions. We trained a target-velocity decoder (chance level: one in five), and similarly checked
the transferred decoding accuracy (Figure 2B right). We observed that the target-velocity decoder
was locked with reach direction, as the transferred decoding accuracy diminished with increasing
difference of reach direction. These results qualitatively imply the interaction as that target
velocities affected the reach-directional tuning, especially by $TV_{dir.}$ This target-motion effect was
most obvious at the MO (Figure S7C).
To explore how sensory information influences neural dynamics while preserving motor output,
we performed PCA on the normalized population activity. We obtained the trial-averaged neural
trajectories (five target velocities by eight reach directions, totally 40 conditions) after TO or after
GO (Figure S9A). The distance between neural trajectories grouped by reach-direction conditions
was larger than the distance grouped by target-motion conditions, especially after GO (Figure
S9B), consistent with the dPCA results that M1 primarily encoded reach direction rather than
target velocity. In addition, the distance of neural trajectories between CCW and CW was much

larger than the distance between 120°/s and 240 °/s conditions, indicating that TV$_{dir}$ dominated

the target-motion effect, agreeing with the decoding results (Figure S9C).

However, these neural trajectories were not yet the ideal description, because they were

shaped mostly by time. Therefore, to highlight the proposed target-motion effect on reach

direction, we focused on four key time windows and snapshotted the neural trajectory as neural

state to extract the coding rule at single-trial level and from a geometric view. Here, we define the

"neural states" as the projection of single-trial data during a specific time bin on principal

components (Mante et al., 2013; Parthasarathy et al., 2017; Sun et al., 2022). At the MO, the first

two principal components (PCs) of the neural states explained the most variance ([24.8%,

13.8%]) and were most related to reach direction (the goodness of fitting reach direction, [R^2_{pc1},

R^2_{pc2}] = [0.82, 0.81], Figure 2C). While reach direction was represented by the first two PCs at GO

and during movement execution, target velocity influenced the tuning pattern of the first three

PCs in various ways (Figure 2D).

Figure 2. Features of the encoding pattern at population level

A The decoding accuracy (SVM with 10-fold cross-validation) of reach direction (black line) and target velocity (blue line) by population activity (monkey C, n=95, 772 trials), is aligned to target on (TO), GO, and movement onset (MO). The dash-dotted lines are chance level of decoding

reach direction (black, one in eight) and target velocity (blue, one in five). The shaded area is the

standard deviation of the decoding accuracy for 10 permutations.

B The left panel shows the performance of reach-direction decoder (chance level: one in eight)

transferred between different target-motion conditions. The SVM decoder was built on randomly

selected 100 trials in training dataset and tested in another 100 trials from a dataset of different

conditions (CCW *vs.* CW, 120 *vs.* 240, static *vs.* motion). The distributions of decoding accuracy

were from 1000 repetitions and compared with one tailed t-test (p<0.01, with three stars). The

right panel shows the performance of target-velocity decoder (chance level: one in five) in

different reach-direction conditions. The accuracy distribution was also obtained from 1000

repetitions.

C The explained variance and representation of the principal components. The first row shows

the explained variance of each PC (cumulatively over 70%). The second row shows the PCs'

fitting goodness (R^2_{PCs}) of reach direction and target velocity in four epochs.

D Directional tuning curves of the principal components. Each row shows the directional tuning of

one PC (the first three PCs in **C**) in four epochs. Each dot represents a trial, tuning curves are

averaged in eight reach directions, and PDs of PCs are indicated by the short lines in the top of

subplot by a weighted sum of response. The colors of the lines and dots mean the target-motion

conditions, as the same as the legend on the left. The goodness of fitting reach direction (R^2_{DirVel})

for the single-trial PCs under single target-motion conditions is shown by mean ± sd. across

conditions.

Orbital neural geometry in latent space

We visualized the neural states in the low-dimensional space spanned by the above three PCs

(Figure S7D). At the MO, the projections of the single-trial neural states onto the PC1-PC2

subspace distributed in reach-direction order, as reach direction was divided into eight sectors,

there appeared to be eight reach-direction clusters (Figure 3A top and Figure 3B left).

Interestingly, the neural states under each single target-motion condition formed ring-like

structures and the fitted ellipses exhibited concentric shapes (the fitting goodness of ellipses, R^2 =

0.92 ± 0.01, mean ± sd. across target-motion conditions; see Methods). Moreover, these ellipses

tilted in condition-dependent angles, which is particularly evident in the PC2-PC3 subspace

(Figure 3A bottom and Figure 3B right). Note that here we performed a linear transformation on

all resulting neural state points to make the ellipse of the static condition orthogonal to the z-axis

for better visualization.

Next, we quantified the spatial features of these ellipses by calculating the tilting angles, which

were defined as the angles between the normal vectors of the moving-target and static-target

conditions. Strikingly, these tilting angles were linearly correlated with target velocity (both TV_{mag}

and TV$_{dir}$), and the relationship was robust in nine datasets from three monkeys (Figure 3C and

Figure S10).

To eliminate hand-speed effect, we resampled trials to construct a new dataset with similar

distributions of hand speed in each target-motion condition and found similar orbital neural

geometry. Moreover, the target-motion gain model provided a better explanation compared to the

hand-speed gain model (Figure S11).

Given these results, we propose that this orbital neural geometry epitomizes the sensorimotor

dynamics of M1 at the population level. The sensory input can regularly modulate neural states in

an orthogonal dimension (PC3), without interfering with motor generation (in PC1 and PC2).

Figure 3. The orbital neural geometry in latent dynamics

A Three-dimensional neural state of M1 population activity obtained by PCA. Each point

represents a single trial. The upper subplot is colored according to five target-motion conditions,

while the bottom is in colors corresponding to eight reach directions. The explained variances of

the first three PCs were 17.7%, 11.3%, and 5.9%. Neural data were from monkey C (N=480,

merged six sessions), for each of the total 40 conditions, 15 trials were randomly sampled.

B Fitted ellipses of neural states. The ellipses fitted in **A** are projected onto three two-dimensional

subspaces, colored by target velocities (left column) or reach directions (right column).

C The relation between the tiling angle and target velocity. The tilting angle is calculated between

ellipses of the moving-target conditions and the static-target condition (0 °/s) in the range from -

90° to 90°, CCW is defined as positive. Circles, squares, and triangles correspond to monkeys C

(7 sessions), G (4 sessions), and D (4 sessions), respectively. The lines indicate the linear fitting

between the tilting angle (θ) and target velocity (vel.), with solid line for monkey C (θ =

0.23*vel.+4.2, R^2=0.91), dashed line for monkey D (θ = 0.26*vel.+4.3, R^2=0.81), and dotted line

for monkey G (θ = 0.15*vel.-1.4, R^2=0.89).

Population neural geometry relies on neuronal tuning

To test whether a group of single neurons with a certain type of mixed sensorimotor selectivity

could exhibit the orbit neural geometry, three neuronal models were constructed based on the

three fitting models described above (PD shift, gain, and addition, see Figure 1D and Figure S2-

S4). We ran a simulation with these representational neuronal models (Figure 4A). Here, each

group consisted of 300 model neurons with their PDs uniformly distributed, and was solely

modulated as PD-shift, gain, or addition (see Methods). The resulting neural geometry of the

three simulation groups showed distinct features (Figure 4B): The single-condition ellipses were

inclined with target-motion-dependent angles in the PD-shift and gain groups, similar to the real

neural data, but the ellipses in the addition group were layered in parallel. The reach-direction

clusters in the first two PCs were conservative in the gain and addition group, but not in the PD-

shift group. These results indicate that the neural states of the real data mainly resembled the

geometry feature of the gain modulation group. Nonetheless, we found that a population with a

uniform mixture of all three modulations was able to simulate the neural geometry as well (Figure

4C).

Comparing with the static-target condition, we calculated relative rotation angle and tilting angle

between ellipses along with the vertical shift of neural states, in order to quantify the simulated

structure (Figure 4D). The results show that the real data yielded a smaller rotation angle than the

PD-shift group, a smaller vertical shift than the additive group, but larger tilting angles than all

models. The mixed group had the most similar tilting angle, though with moderate performance in

rotation angle and state shift.

These simulations suggest that the existence of PD-shift and additive modulation would not

disrupt the neural geometry which is primarily driven by gain modulation; rather, it is possible that

these three modulations support each other in a mixed population.

Figure 4. The shape of neural dynamics relies on neuronal mixed selectivity

A The tuning curves of three ideal neurons in five target-motion conditions. From up to down, they are PD shift, gain modulation, and addition.

B The simulated neural states from three groups of ideal neurons, colored according to target-motion conditions (first row) and reach directions (second row). For each simulation, the neural states were obtained from 300 model neurons by PCA. The neural state in 180° reach direction is marked with a red dot. The first two principal components can explain more than 95% of the variance in the data (the explained variance of the first three PCs, Gain: 49.5%, 46.6%, and 2.0%; PD shift: 50.1%, 47.1%, and 1.6%; Addition:50.8%, 47.9%, and 1.4%).

C The neural states of a mixed group of 100*3 model neurons, as in **B.** The explained variance of the first three principal components were 48.4%, 44.2%, and 3.1%.

D Quantification of the difference between neural-state ellipses in four simulated groups and a real dataset (monkey C, n=95). Rotational angle is the angular differences in the first two neural

320   state. Tilting angle is the relative angle of the normal vector of ellipses. State shift is the root of

321   mean squared distance between two ellipses.

The recurrent neural network provides dynamic insights

To infer how such modulated subpopulations would interact with each other in a dynamical

system, we trained 100 RNN models with random weight initialization (Figure 5A; see Methods).

The inputs included motor intention, target location, and a GO signal. Motor intention was defined

as an abstract motor command predicted to compensate for sensorimotor delays (Cui, 2016), and

could be provided by the PPC (Andersen and Buneo, 2002; Andersen and Cui, 2009), here

simplified as the interception location. The network was to generate hand velocity after MO. For a

fixed validation set of 500 trials, these trained network models performed well (distance between

the reach endpoint and the target was 0.0046 ± 0.0027, a. u., mean ± sd., while the radius of

target motion was 0.15).

In these network models, we found three comparable features. First, from the decoding result,

target motion information existed in nodes' population dynamics shortly after TO (Figure S12A).

Second, most of the activated nodes (for example, Figure 5B) could be classified into the above-

mentioned three modulations by the same statistical standard as for the real neural data (Table

1). Third, the states reduced from node population activity were arranged in a way resembling the

actual neural geometry at the MO (the fitting goodness of ellipses, R^2 = 0.98 ± 0.05, mean ± sd.;

Figure 5C). The tilting angles followed the same pattern as suggested by the actual results

(Figure 5D). We then performed canonical component analysis (CCA) and Procrustes analysis

(Table S2; see Methods), the results also indicated the similarity between network dynamics and

neural dynamics.

With these RNN models, we tried perturbations to investigate the function of certain modulation

groups and their connection. The ablation experiments showed that losing any kind of modulation

nodes would largely deteriorate the performance, and those nodes merely with PD-shift

modulation could mostly impact the neural state structure (Table S3). The connections within and

between gain modulation nodes and additive modulation nodes were stronger than the others

(p<0.01, K-W Test) but comparable to each other (p=1.00, K-W Test; Figure 5E). However,

further strengthening the connection from gain to additive nodes or within additive nodes would

impair the orbital structure more significantly (Table S3). We speculate that the influence, and

perhaps the formation, of modulations both relate to the robust input and output weight pattern

across models (Figure S12B-C). From these results, all the three modulations are necessary for

anticipative performance and their interactions likely mold the ring-like structure, though their
specific roles are not yet elucidative.
We also tested three alternative network models: (1) only receives motor intention and a GO-
signal; (2) only receives target location and a GO-signal; (3) initialized with sparse connection
(sparsity=0.1); the unmentioned settings and training strategies were as the same as those for
original models (Table S4; see Methods). The results showed that the three modulations could
emerge in these models as well, but with obviously distinctive distributions. In (1), the ring-like
structure became overlapped rings parallel to the PC1-PC2 plane or barrel-like structure instead;
in (2), the target-motion related tilting tendency of the neural states remained, but the projection
of the neural states on the PC1-PC2 plane was distorted and the reach-direction clusters
dispersed. These implies that both motor intention and target location seem to be needed for the
proposed ring-like structure. The initialization of connection weights of the hidden layer can
influence the network's performance and neural state structure, even so, the ring-like structure
and especially the target-velocity dependent tilting remained in a degree. However, we did not
find specific connection patterns in these sparsely-initialized networks (p=0.73, K-W Test).

Figure 5. The neural geometry in RNNs

A Network architecture. The network inputs consist of motor intention, target location, and GO signal. The motor intention is the two-dimensional Cartesian coordinate of the interception location, and exists fixed during MO-50ms to MO; the target location is the two-dimensional Cartesian coordinate of the moving target, and appears time-varying during the whole trial; the GO-signal is a step function, jumping from 0 to 1 at GO. The RNN with 200 hidden units is expected to output hand velocity in two-dimensional Cartesian coordinates for accurate interception.

B Three example nodes with PD-shift modulation, gain modulation, and additive modulation. Similar to Figure 1D.

C Three-dimensional neural state of node activity obtained by PCA, colored according to target-motion conditions (top) and reach-direction conditions (Bottom). Similar to Figure 3A.

D The tilting angle of ellipses. Similar to Figure 3D. The fitted line is θ = 0.15*vel.+0.11, R^2=0.96, across five target velocities.

383 **E** The connectivity between different types of modulations. On the left is a boxplot representing

384 the averaged absolute connection weight, across 100 models. S for PD-shift nodes, G for gain

385 nodes, and A for additive nodes. On the right is a diagram of the connectivity, with linewidth

386 representing the relative connection strength.

Discussion

Mixed selectivity in M1 has been widely reported; but to our knowledge, sensory modulation on anticipatory motor control, as examined in this study, has been scarcely studied. To reveal how sensory context influence neural dynamics during movement, we recorded M1 population activity from monkeys performing a flexible manual interception that is highly dependent on predictive sensorimotor transformations. Single-neuron activity showed that the movement tuning of M1 neurons varied with target-motion conditions in complicated ways, including PD shift, gain modulation, addition, or their mixtures. Unsupervised dimensionality reduction (PCA) on population activity revealed an orbital neural geometry, where neural states of single trials spontaneously formed ellipses and were tilted according to target-motion conditions. Such a neural geometry, which could be simulated with a group of representational neurons for gain modulation alone, also emerged in the RNNs with appropriate input-output mappings. As suggested by the dynamic and connected population of the RNNs, the interactions between modulations were sophisticated in preserving the encoding of motor output. These results reveal sensory modulation of peri-movement neural dynamics at the single-neuron and population levels, and bridge the neuronal mixed sensorimotor selectivity and a low-dimensional neural geometry.

A main concern about our finding is that the observed modulation would actually be attributed to interactions between kinematic variables, such as reach direction and hand speed (Inoue et al., 2018; Moran and Schwartz, 1999; Paninski et al., 2004). However, we can eliminate the hand-speed effect for at least two reasons: First, the hand speed pattern and neural activity pattern were not consistent. We found hand speed vary with TV_{mag}, but the largest gap in neuronal activity related to TV_{dir} (Figure 1 & S1); Second, the neural geometry was not contingent upon hand speed distribution, as similar hand-speed distribution across target-motion conditions did not change the phenomenon (Figure S11). It is a pity that we did not record sufficient muscle data during interception to rule out the motor cortical representation for kinetic variables such as force (Evarts, 1968; Scott and Kalaska, 1997; Sun et al., 2022), but we believe that this would not invalidate our main conclusions.

Another concern is whether the neural geometry was representing sensory information. Previous studies have showed that M1 has transient responses to behaviorally relevant visual cues (Alexander and Crutcher, 1990; Lamarre et al., 1983; Lurito et al., 1991; Port et al., 2001; Rao and Donoghue, 2014; Rizzolatti et al., 1981), but it is seldom evoked by purely visual stimuli, only weakly tuned (Kruse, 2002; Kwan et al., 1985; Merchant et al., 2001). Here, the modulated neural dynamics, though distinguished according to target-motion conditions, are ultimately shaped for motor generation rather than sensory representation. To further clarify, the discussing target-motion effect is different from the sensory modulation in action selection (Cisek and

Kalaska, 2005), motor planning (Pesaran et al., 2006), visual replay and somatosensory feedback

(Pruszynski et al., 2011; Stavisky et al., 2017; Suway and Schwartz, 2019; Tkach et al., 2007),

because it occurred around movement onset and in predictive control trial-by-trial.

In the present study, we focused on the orbital neural geometry by taking 'snapshots' of neural

activities with PCA. In addition to this unsupervised dimensionality reduction method, we also

tried another supervised one – dPCA (Kobak et al., 2016). However, the dPCA results (Figure

S8) showed that the condition-independent temporal component explained the largest variance,

followed by the component of reach direction, whereas the interactive and target-velocity

components only covered very small proportions. The dPCA did not distinguish an independent

subspace coding target-velocity (Figure S8C), suggesting the target-velocity information persisted

in the shared subspace with reach direction.

It would be interesting to explore whether other motor areas also allow sensory modulation

during flexible interception. The functional differences between M1 and other areas lead to

uncertain speculations. Although M1 has pre-movement activity, it is more related to task

variables and motor outputs. Recently, a cycling task sets a good example that the

supplementary motor area (SMA) encodes context information and the entire movement (Russo

et al., 2020), while M1 preferably relates to cycling velocity (Saxena et al., 2022). The dorsal

premotor area (PMd) has been reported to capture potential action selection and task probability,

while M1 not (Cisek and Kalaska, 2005; Glaser et al., 2018; Wang et al., 2019). If the neural

dynamics of other frontal motor areas are revealed, we might be able to tell whether the orbital

neural geometry of mixed selectivity is unique in M1, or it is just inherited from upstream areas

like PMd. Either outcome would provide us some insights into understanding the interaction

between M1 and other frontal motor areas in motor planning.

From a bigger view, how could the motor cortex cooperate with other brain regions during

flexible interception? It could be true that the communication and collaboration of PPC-motor

cortex circuity translate abstract motor intention to executable motor command: PPC transforms

sensory modulation to the motor cortex, while the motor cortex sends potential movement

direction to PPC, and then their recurrent interaction generates mature motor intention and

command. Our simulations with neuronal models and RNNs demonstrate that differences in

modulations and the interaction between them can be essential. In fact, modulations in the form

of PD shift and gain have been widely found in the motor cortex and PPC. For example, motor

cortex neurons experience gain and PD shift modulation by arm posture(Kakei et al., 2001, 1999;

Scott and Kalaska, 1997). Neurons in PPC have eye and hand gain fields for a visually-guided

reach plan (Batista et al., 1999; Buneo et al., 2002; Chang et al., 2009), integrating target position

related to gaze and hand position to form motor intention (Andersen et al., 1997; Andersen and

Buneo, 2002; Cohen and Andersen, 2002). Therefore, we posit that the three types of

modulations should all be involved in sensorimotor computation in predictive motor control.

Furthermore, recent studies show that interactions of PPC-motor cortex circuity are involved in

motor planning, spatial transformation and motor selection (Dann et al., 2016; Martínez-Vázquez

and Gail, 2018; Schaffelhofer and Scherberger, 2016). The PPC-M1 circuit, as a key part of

cortico-subcortical networks for the predictive sensorimotor control (Brozović et al., 2007), will be

a topic for future studies.

Methods

Experimental model and subject details

Three adult male rhesus macaques (monkey C, D, and G, *Macaca mulatta*, 7-10 kg) were recruited in this study. The monkeys sat in primate chairs to perform the task. The stimuli were back-projected onto a vertical touch screen (Elo Touch system; sampling at 100 Hz, spatial resolution <0.1 mm) about 30 cm in front of the monkeys. Hand trajectory was tracked by an optical camera (VICON Inc.) with an infrared marker on the fingertip. All animal maintenance and procedures were in accordance with NIH guidelines and approved by the Institutional Animal Care and Use Committee (IACUC) of Institute of Neuroscience, CAS.

Task and behavior

The monkeys were trained to perform a flexible manual interception task in a dark room. The task paradigm was modified based on the visually-guided reaching interception task in a previous study (Li et al., 2018). In the beginning, the monkey held the green central dot of a touch screen for 600 ms to initiate a trial (Figure 1A). Then, a green target dot appeared at a random location, moving along a circular path with a radius of 15 cm around the central dot. The central dot turned dark as a GO cue after a random delay (400-800 ms); then the monkey could intercept the target at any moment within 150-800 ms after the GO cue. Once any peripheral location was touched, the target stopped. The tolerance range of the touch endpoint for correct trials was within 3 cm of the target. The monkey would be rewarded with juice after each correct trial. Conditions where targets moved clockwise (CW; -240 °/s, -120 °/s) or counterclockwise (CCW; 120 °/s, 240 °/s), as well as targets stayed stationary (0°/s), were pseudo-randomly interleaved trial by trial. Additional target velocities (-360 °/s, -180 °/s, 180 °/s, 360 °/s added) were introduced in subsequent sessions (Table S1). The single-trial MO is defined as the moment when hand velocity first rose to 5% of the maximum.

Data collection

After the monkeys were adequately trained for the task (successful rate > 90%), head-posts were implanted stereotaxically under anesthesia (induced by 10 mg/kg ketamine, then sustained by 2% Isoflurane). After a few weeks of recovery and adaptation, the monkeys were implanted with Utah microelectrode arrays (Blackrock Microsystems, Salt Lake City, UT) in M1 of the hemisphere contralateral to the handedness (Figure 1C, a 128-channel array for monkey C, 96-channel arrays for monkey G and D). The recording areas were identified by Magnetic Resonance Imaging (MRI) and cortical sulcus features. Neuronal activity was recorded via a Blackrock Microsystem 256-channel recording system, sampled at 30 kHz. We collected 85 ± 16, 45 ± 8, 98 ± 18 well-isolated units from monkey C, G, and D across sessions, respectively (mean ± sd., more details in Table S1). Hand trajectory was collected by optical camera (VICON Inc.) with an

infrared marker on the fingertip from GO to Touch, and touch endpoint was collected by

touchscreen.

Peri-stimulus time histograms (PSTHs)

The PSTHs and spike rasters of single neurons are shown in Figure 1D and Figure S2-S4. All

trials were classified into 40 conditions, eight reach-endpoint sectors by five target-motion

conditions. Condition-averaged firing rates were calculated with 50-ms bins and smoothed with a

Gaussian kernel (standard deviation = 20 ms). The standard error of firing rates was estimated

from the 10 bootstrap samples in the trials of corresponding condition.

Classification of neuronal tuning properties

To depict target-motion modulation for single-neuron reach-direction tuning, we applied a series

of statistical analyses for classification.

We first calculated three indices of each neuron for each target-motion conditions, using the

trial-averaged firing rate around movement onset (MO ± 100ms). These indices were preferred

direction (PD), tuning depth, and offset activity. The PD of each neuron for a certain target-motion

condition was calculated by the weighted sum of neuronal firing rates averaged by eight reach-

direction sectors. The tuning depth of each neuron was determined by the range (max - min) of

firing rates in corresponding target-motion conditions. The offset activity of each neuron was

calculated by the mean firing rate (MO ± 100 ms) of each target-motion condition.

Then, we classified PD shift, gain, and addition groups (in Table 1). A neuron was classified as

'PD shift', if its PDs were significantly different between the moving-target conditions and the

static-target condition (Watson-Williams test in circular data, CircStat by Berens, 2009); as 'gain',

if its tuning depths were significantly different between the moving-target conditions and the

static-target condition (two-tailed Wilcoxon signed-rank test, p<0.05); as 'offset', if its offset

activities were significantly different between the moving-target conditions and the static-target

condition (two-tailed Wilcoxon signed-rank test, p<0.05).

Population decoding

The population activity of the motor cortex was used to decode target motion and reach direction

by support vector machine (SVM). Neuronal firing rate was soft-normalized as

$$FR_{norm.} = \frac{FR_{raw}}{FR_{max} - FR_{min} + 5}$$

where raw firing rates were divided by the range of firing rates plus five (Churchland et al., 2012).

We trained two SVM classifiers (MATLAB function 'fitcecoc', 10-fold cross-validation) to decode

reach direction (chance level: one in eight) and target motion (chance level: one in five) of single

trials in 100-ms window sliding with 50-ms step (Figure 2A). The temporal decoding was repeated

ten times to obtain the mean and standard deviation of decoding accuracy.

We tested the generalization of reach-direction and target-motion decoders (SVM, MATLAB

function 'fitcecoc') in different conditions during execution period (MO ± 100 ms, Figure 2B). The

decoder predicted single-trial reach direction (one of the eight 45° sectors) or target motion (one

of the five ones) across conditions based on normalized population activity. The reach-direction

decoder (Figure 2B left), which was trained by trials in a certain set of target-motion conditions,

was tested with trials from another set of target-motion conditions (CCW vs. CW, or 120 vs. 240,

or static vs. motion, 100 trials randomly selected without replacement from corresponding

training-test datasets), and the training-test decoding was repeated 1000 times to compare the

distribution of accuracy with paired t-test. The target-velocity decoder (Figure 2B right), which was

trained by trials in a given reach-direction condition within 45°, was tested with trials from other

reach-direction conditions.

Unsupervised dimensionality reduction

The population activity was stored in NKT datasets, where N, K, and T denote the neurons, trials,

and time bins, respectively. For neural state, we averaged T dimension of neural activity in a

100-ms bin (for example, the two 50-ms bins around MO), and normalized neuronal firing rates by

Z-score (MATLAB function 'zscore') to get a K x N dataset. After preprocessing, we used PCA

(MATLAB function 'pca') to reduce the dimension from K x N to K x P (P is the number of

principal components), and we fitted the PCs with reach direction ($PC = a_1 * cos(\theta) + a_2 *$

$sin(\theta) + c$) and target velocity ($PC = \frac{a_1}{1+e^{-a_2(vel.)}} + c$) in Figure 2. We also tried independent

component analysis (ICA, MATLAB function 'rica'), with similar results in Figure S7B. Neural

states of single trials were colored according to target velocity or reach direction and fitted as

three-dimensional (first three PCs, Figure 3) ellipses by MATLAB package 'MatGeom' (Legland,

n.d.). To show the relative position of ellipses in the best viewing angle, we used an isometric

affine transformation to globally map all neural state points on new axes, while preserving the

proportional relationships between points. After this linear transformation, the azimuth and

elevation of ellipses changed slightly, but the tilting angle between ellipses remained constant

(Figure 3). The tilting angles, rotation angles, and state shift were calculated between ellipses of

the moving-target conditions and the static-target condition in each session, with CW defined as

negative angles and CCW as positive angles. A set of tilting angles were obtained from

corresponding conditions in one dataset, and a linear regression model was used to fit all ellipses

angles (θ) and target velocities (vel.) in nine sessions. For neural trajectory, we averaged NKT to

NCT (40 condition, five target velocities by eight reach directions) and reduced neural dimension

N to explain concatenated C x T (40 conditions * 10 bins) with PCA. The neural trajectories of

preparatory and peri-movement period (TO~ +500ms, GO~ +500ms) were shown in Figure S9

with Euclidian distance across conditions.

Fitting and simulating single-neuron activity

We used PD shift, gain, offset, and full models to fit neuronal activity. Neurons are fitted by single-

trial data. We introduced a special sigmoid function to fit the nonlinear target-motion effects

because target-velocity direction (CCW vs. CW) has a stronger effect than target-velocity

magnitude (120 vs. 240) (Churchland and Lisberger, 2001; Pouget and Sejnowski, 1997). The

gain and additive models refer to hand-velocity gain studies (Amirikian and Georgopulos, 2000;

Inoue et al., 2018; Moran and Schwartz, 1999).

In the gain model, the target-motion effect on the amplitude of cosine tuning is denoted as:

$$FR = \left(\frac{a_1}{1 + e^{-a_2(vel.)}} + c_2\right) * cos(\theta - \theta_{pd}) + c_1$$

where FR is the firing rate at the movement onset (MO ± 100 ms). θ and *vel*. are the reach

direction and target velocity, respectively; θ_{pd} is the fitted preferred direction of the neuron;

a_1, a_2, c_1 are constants to be fitted.

In the additive model, the target velocity adjusts the offset activity, as:

$$FR = a_1 * cos(\theta - \theta_{pd}) + \frac{a_3}{1 + e^{-a_2(vel.)}} + c_1$$

with similar symbols to the gain models, plus the new constant a_3.

In the PD shift model, the target-motion effect on PDs is represented as:

$$FR = a_1 * cos\left(\theta - \theta_{pd} + \frac{a_3}{1 + e^{-a_2(vel.)}}\right) + c_1$$

with the similar symbols to the above models.

The full model integrates all the three of the above effects:

$$FR = a_1 * cos(\theta - pd_{vel}) + \frac{a_2}{1 + e^{-a_2(vel.)}} + \frac{a_3}{1 + e^{-a_2(vel.)}} * cos(\theta - pd_{vel}) + a_4$$

$$* cos\left(\theta - pd_{vel} - \frac{a_5}{1 + e^{-a_2(vel.)}}\right) + c_1$$

with constants a_1, a_2, a_3, a_4, a_5.

We fitted neuronal activity with these four models (MATLAB 'fit' function), and compared the

fitting goodness with adjusted R-squares ($R^2_{adj.} = \frac{(1-r^2)(n-1)}{n-p-1}$, where r is the original R-square, n

is the trial number, and p is the degree of the polynomial).

Simulations with model neurons were based on three models to investigate the relationship

between neuronal tuning (Figure 4A) and population neural geometry (Figure 4B). We first built

three model neuron groups (each n=300) and performed PCA to obtain the neural state. Then,

we repeated this in a mixed group, including 100 neurons in each of PD shift, gain and additive

model (Figure 4C). Model neurons had three components: cosine-tuning for reach direction

($\theta_{pd,n}$, PD uniformly distributed around the circle for each neuron n), Gaussian temporal profiles

(t=1:200, σ=30, peak time $t_{\mu,n} \in N(100, 10)$, the 100-th bin is the MO, random distribution for $t_{\mu,n}$

of neurons), and distinct target-motion modulation for each group. We designed five target

velocity values ($vel. = [1,2,3,4,5]$) and 64 reach directions ($\theta = 360 * \frac{[1:64]}{64}$), for 320 trials in total

(Michaels et al., 2016). The concrete expression of model neurons follows as below:

The gain-model neuron:

610
$$FR_{t,n,vel.,\theta} = e^{\frac{-(t-t_{\mu,n})^2}{2*30^2}} * \left(\frac{1}{1 + e^{-g_n(vel.-3)}} * cos(\theta - \theta_{pd,n}) + 1 \right)$$

611   here, g_n is randomized within [0, 1] for different neurons, modulating the target-motion gain in a

612   sigmoidal function.

613   The PD-shift model neuron:

614
$$FR_{t,n,vel.,\theta} = e^{\frac{-(t-t_{\mu,n})^2}{2*30^2}} * \left(cos\left(\theta - \theta_{pd,n} - \frac{90}{1 + e^{-s_n(vel.-3)}}\right) + 1 \right)$$

615   here, s_n is a random value within [0, 1.5] for different neurons, and contributes to a sigmoidal

616   function with target-motion shift on neuronal PD ($\theta_{pd,n}$).

617   The additive model:

618
$$FR_{t,n,vel.,\theta} = e^{\frac{-(t-t_{\mu,n})^2}{2*30^2}} * \left(cos(\theta - \theta_{pd,n}) + \frac{1}{1 + e^{-g_n(vel.-3)}} + 1 \right)$$

here, g_n is also a random value within [0, 1] for different neurons, and adjusts the offset activity in

a sigmoidal function with target-motion.

The firing rates of three groups of model neurons (K x N x T, 320 x 300 x 200) were averaged

at the MO (mean T = [50:150]) to get the K x N (320 x 200) dataset for PCA. As with the real

neural data, we selected the first three PCs (K x C, 320 x 3) to derive the simulated neural state

shown in Figure 4.

RNNs training

For the inputs of RNNs, motor intention appears from MO-50 ms to MO, and is represented as

constant variables in the form of two-dimensional Cartesian coordinates; target location is

designed as time-varying two-dimensional Cartesian coordinates of the target throughout the

entire trial; the GO-signal is a step function jumping from 0 to 1 at GO. Each RNN consists of 200

hidden units, and outputs hand velocity for accurate interception after the MO.

The RNN nodes evolve according to a standard dynamic differential equation:

633
$$\tau \dot{x} = -x + Jr + Bu$$

634   where τ is a time constant (here 50 ms), x is the activity, r is the firing rate, and u denotes the

635   combined inputs. r can be calculated from x following:

636
$$r = \begin{cases} \tanh(x), x > 0 \\ 0, \qquad x \le 0 \end{cases}$$

637 The connection matrix J of hidden layer was randomly initialized with a normal distribution (mean

638 = 0, sigma = $g/\sqrt{N}$, g = 1.5, N = 200), and the connection between inputs and hidden units, the

639 matrix B, was initialized as all zero. The initial states were zero vectors. The output z is obtained

640 by

641
$$z = Wr$$

642 where W is the read-out weight, and is expected to reproduce the desired hand velocity

643 generated by bell-shaped physical equation (Kao et al., 2021). W was also initiated as zero

644 matrix. The loss function is:

645
$$E = e + \alpha r_1$$

where e is the mean squared error of z and training target. r_1 is a regularity (Sussillo et al., 2015),

denoting the magnitude of the nodes' activity and calculated as activity squared summed across

time. $\alpha = 1e - 7$. The optimization was realized with optim.Adam() based on PyTorch, the

learning rate was 0.001.

RNNs analyses

The classification of modulation type on network nodes was finished with the same procedure for

real data as above. The firing rates of nodes ([0, 1]) were used for all analyses.

For decoding, similar to **Population decoding**, two SVM classifiers (Python, SVC from

sklearn.svm, 10-fold cross-validation) were trained to decode reach direction and target motion of

single trials in 100-ms window sliding with 50-ms step. It should be noted that, instead of original

firing rates, the first 100 PCs were used for decoding, in order to decrease the influence of

inactivated nodes. The decoding results were first averaged cross 10 cross-validation repetitions

and then the resulting means were averaged across 100 models to obtain mean ± sd.

We performed Canonical Component Analysis (CCA) and Procrustes analysis to validate the

similarity between data and network dynamics. The data were around the TO or around the MO

each in 2 seconds (±1s, 50 ms x 40 bins). The RNN activity were collected between MO-200 ms

and MO+120 ms, considering the shortest movement time, and then averaged into 40 bins for

comparison. First, we performed PCA (Python, PCA from sklearn.decomposition) on actual and

model data to get the first 30 PCs (KT x N → KT x 30), respectively. Then, we used these two-

dimensional matrices to compute their first ten canonical components with CCA (Python, CCA

from sklearn.cross_decomposition). The Pearson correlation coefficients (Python, pearsonr from

scipy.stats) were calculated between paired canonical components. The Procrustes disparity

(Python, procrustes from scipy.spatial) were computed between pairs of data and network first 30

PCs.

RNNs experiments

We performed ablation of nodes and manipulation of connection weights. In former, we selected the nodes from a certain type of modulation, and set all its connection weights to zero to simulate 'ablation'. When manipulating the connection between certain types of modulations, the selected connection weight was adjusted to be 1.5 times the original. Both above perturbations were retained for the entire trial. The perturbed network was tested with the same validation set of 500 trials as for the intact network.

We also constructed three alternative models to check the influence of input and weight initialization. The first one only receives motor intention and a GO-signal (GM); the second only receives target location and a Go-signal (GT); these two were initialized and trained as the main model. The third one was initialized with sparse connection (sparsity=0.1); except for this, the input and the training were as the same as the main model. All involved modes shared the same validation set.

Software

We used the Python 3.11 (python.org) and MATLAB 2018b (The MathWorks, Inc.).

Acknowledgments

We thank C. Li, J. Malpeli, C. Zheng, and R. Zheng for helpful comments and discussions; C. Guan for veterinary assistance; and P. Ding, L. Du, and Z. Xiao for administrative support. This work was supported by the National Key R&D Program (Grants 2020YFB1313400 and 2017YFA0701102), National Science Foundation of China (Grants 31871047 and 31671075), and CAS Strategic Priority Research Program (Grant XDB32040103).

Additional information

Competing interests

The authors declare no conflict of interests.

Funding

Funder	Grant reference number	Author
National Key Research and Development Program of China	2020YFB1313400 2017YFA0701102	He Cui
National Science Foundation of China	31871047 31671075	He Cui
Strategic Priority Research Program of Chinese Academy of Sciences	XDB32040103	He Cui

The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

Author contributions

Y. Zhang and H. Cui designed the experiment, Y. Zhang and T. Wang collected the data, Y. Zhang and Y. Chen analyzed the data, Y. Chen built RNNs, Y. Zhang performed the simulation, Y. Zhang, Y. Chen, T Wang, and H. Cui prepared the manuscript.

Author ORCIDs

Yiheng Zhang https://orcid.org/0000-0002-5370-1316

Yun Chen https://orcid.org/0000-0002-0817-2160

Tianwei Wang https://orcid.org/0000-0002-5192-5594

712   He Cui https://orcid.org/0000-0001-6277-9804

714   **Peer review material**

715   https://doi.org/10.7554/eLife.100064.1.sa1

Additional files

Supplementary files

719   MDAR checklist

Data availability

722   The example experimental datasets and relevant analysis code have been deposited in Mendeley
723   Data at https://data.mendeley.com/datasets/8gngr6tphf. The RNN relevant code and example
724   model datasets are available at https://github.com/yunchenyc/RNN_ringlike_structure.
725

References

Alexander GE, Crutcher MD. 1990. Neural representations of the target (goal) of visually guided arm movements in three motor areas of the monkey. *J Neurophysiol* **64**:164–178. doi:10.1152/jn.1990.64.1.164

Amirikian B, Georgopulos AP. 2000. Directional tuning profiles of motor cortical cells. *Neurosci Res* **36**:73–79. doi:10.1016/S0168-0102(99)00112-1

Andersen RA, Buneo CA. 2002. Intentional maps in posterior parietal cortex. *Annu Rev Neurosci* **25**:189–220. doi:10.1146/annurev.neuro.25.112701.142922

Andersen RA, Cui H. 2009. Intention, Action Planning, and Decision Making in Parietal-Frontal Circuits. *Neuron* **63**:568–583. doi:10.1016/j.neuron.2009.08.028

Andersen RA, Snyder L, Bradley D, Xing J. 1997. Multimodal Representation of Space in the Posterior Parietal Cortex and Its Use in Planning Movements. *Annu Rev Neurosci* **20**:303–330. doi:10.1146/annurev.neuro.20.1.303

Batista AP, Buneo C a, Snyder LH, Andersen R a. 1999. Reach Plans in Eye-Centered Coordinates. *Science (80-)* **285**:257–260. doi:10.1126/science.285.5425.257

Berens P. 2009. CircStat : A MATLAB Toolbox for Circular Statistics. *J Stat Softw* **31**:128–129. doi:10.18637/jss.v031.i10

Bremner LR, Andersen RA. 2012. Coding of the Reach Vector in Parietal Area 5d. *Neuron* **75**:342–351. doi:10.1016/j.neuron.2012.03.041

Brozović M, Gail A, Andersen RA. 2007. Gain mechanisms for contextually guided visuomotor transformations. *J Neurosci* **27**:10588–10596. doi:10.1523/JNEUROSCI.2685-07.2007

Buneo C a, Jarvis MR, Batista AP, Andersen R a. 2002. Direct visuomotor transformations for reaching. *Nature* **416**:632–636. doi:10.1038/416632a

Chang SWC, Papadimitriou C, Snyder LH. 2009. Using a Compound Gain Field to Compute a Reach Plan. *Neuron* **64**:744–755. doi:10.1016/j.neuron.2009.11.005

Churchland MM, Cunningham JP, Kaufman MT, Foster JD, Nuyujukian P, Ryu SI, Shenoy K V. 2012. Neural population dynamics during reaching. doi:10.1038/nature11129

Churchland MM, Lisberger SG. 2001. Shifts in the population response in the middle temporal visual area parallel perceptual and motor illusions produced by apparent motion. *J Neurosci* **21**:9387–9402. doi:10.1523/jneurosci.21-23-09387.2001

Churchland MM, Shenoy K V. 2024. Preparatory activity and the expansive null-space. *Nat Rev Neurosci* **25**:213–236. doi:10.1038/s41583-024-00796-z

Cisek P, Kalaska JF. 2005. Neural Correlates of Reaching Decisions in Dorsal Premotor Cortex: Specification of Multiple Direction Choices and Final Selection of Action. *Neuron* **45**:801–814. doi:10.1016/j.neuron.2005.01.027

Cohen YE, Andersen RA. 2002. A common reference frame for movement plans in the posterior
parietal cortex. *Nat Rev Neurosci* **3**:553–562. doi:10.1038/nrn873
Cui H. 2016. Forward Prediction in the Posterior Parietal Cortex and Dynamic Brain-Machine
Interface. *Front Integr Neurosci* **10**:1–6. doi:10.3389/fnint.2016.00035
Dann B, Michaels JA, Schaffelhofer S, Scherberger H. 2016. Uniting functional network topology
and oscillations in the fronto-parietal single unit network of behaving primates. *Elife* **5**:1–27.
doi:10.7554/eLife.15719
Evarts E V. 1968. Relation of pyramidal tract activity to force exerted during voluntary movement.
J Neurophysiol **31**:14–27. doi:10.1152/jn.1968.31.1.14
Georgopoulos A, Kalaska J, Caminiti R, Massey J. 1982. On the relations between the direction
of two-dimensional arm movements and cell discharge in primate motor cortex. *J Neurosci*
2:1527–1537. doi:10.1523/JNEUROSCI.02-11-01527.1982
Glaser JI, Perich MG, Ramkumar P, Miller LE, Kording KP. 2018. Population coding of conditional
probability distributions in dorsal premotor cortex. *Nat Commun* **9**:1788.
doi:10.1038/s41467-018-04062-6
Hatsopoulos NG, Amit Y. 2012. Synthesizing complex movement fragment representations from
motor cortical ensembles. *J Physiol Paris* **106**:112–119.
doi:10.1016/j.jphysparis.2011.09.003
Inoue Y, Mao H, Suway SB, Orellana J, Schwartz AB. 2018. Decoding arm speed during
reaching. *Nat Commun* **9**:1–14. doi:10.1038/s41467-018-07647-3
Kakei S, Hoffman DS, Strick PL. 2001. Direction of action is represented in the ventral premotor
cortex. *Nat Neurosci* **4**:1020–1025. doi:10.1038/nn726
Kakei S, Hoffman DS, Strick PL. 1999. Muscle and Movement Representations in the Primary
Motor Cortex. *Science (80-)* **285**:2136–2139. doi:10.1126/science.285.5436.2136
Kalaska JF, Cohen D a, Hyde ML, Prud'homme M. 1989. A comparison of movement direction-
related versus load direction-related activity in primate motor cortex, using a two-
dimensional reaching task. *J Neurosci* **9**:2080–2102. doi:10.1523/JNEUROSCI.09-06-
02080.1989
Kao TC, Sadabadi MS, Hennequin G. 2021. Optimal anticipatory control as a theory of motor
preparation: A thalamo-cortical circuit model. *Neuron* **109**:1567-1581.e12.
doi:10.1016/j.neuron.2021.03.009
Kaufman MT, Churchland MM, Ryu SI, Shenoy K V. 2014. Cortical activity in the null space:
permitting preparation without movement. *Nat Neurosci* **17**:440–448. doi:10.1038/nn.3643
Kobak D, Brendel W, Constantinidis C, Feierstein CE, Kepecs A, Mainen ZF, Qi X-L, Romo R,
Uchida N, Machens CK. 2016. Demixed principal component analysis of neural population
data. *Elife* **5**:1–36. doi:10.7554/eLife.10989

Kruse W. 2002. Temporal Relation of Population Activity in Visual Areas MT/MST and in Primary

Motor Cortex during Visually Guided Tracking Movements. *Cereb Cortex* **12**:466–476.

doi:10.1093/cercor/12.5.466

Kwan HC, Mackay WA, Murphy JT, Wong YC. 1985. Properties of visual cue responses in

primate precentral cortex. *Brain Res* **343**:24–35. doi:10.1016/0006-8993(85)91154-0

Lamarre Y, Busby L, Spidalieri G. 1983. Fast ballistic arm movements triggered by visual,

auditory, and somesthetic stimuli in the monkey. I. Activity of precentral cortical neurons. *J*

Neurophysiol **50**:1343–1358. doi:10.1152/jn.1983.50.6.1343

Legland D. n.d. MatGeom library for geometric computing with MATLAB.

doi:10.5281/zenodo.5768192

Li Y, Wang Y, Cui H. 2022. Posterior parietal cortex predicts upcoming movement in dynamic

sensorimotor control. *Proc Natl Acad Sci* **119**:e2118903119. doi:10.1073/pnas.2118903119

Li Y, Wang Y, Cui H. 2018. Eye-hand coordination during flexible manual interception of an

abruptly appearing, moving target. *J Neurophysiol* **119**:221–234. doi:10.1152/jn.00476.2017

Lu X, Ashe J. 2005. Anticipatory Activity in Primary Motor Cortex Codes Memorized Movement

Sequences. *Neuron* **45**:967–973. doi:10.1016/j.neuron.2005.01.036

Lurito JT, Georgakopoulos T, Georgopoulos AP. 1991. Cognitive spatial-motor processes. 7. The

making of movements at an angle from a stimulus direction: studies of motor cortical activity

at the single cell and population levels. *Exp Brain Res* **87**:562–580.

doi:10.1007/BF00227082

Mante V, Sussillo D, Shenoy K V., Newsome WT. 2013. Context-dependent computation by

recurrent dynamics in prefrontal cortex. *Nature* **503**:78–84. doi:10.1038/nature12742

Martínez-Vázquez P, Gail A. 2018. Directed interaction between monkey premotor and posterior

parietal cortex during motor-goal retrieval from working memory. *Cereb Cortex* **28**:1866–

1881. doi:10.1093/cercor/bhy035

Merchant H, Battaglia-Mayer A, Georgopoulos AP. 2004a. Neural Responses during Interception

of Real and Apparent Circularly Moving Stimuli in Motor Cortex and Area 7a. *Cereb Cortex*

14:314–331. doi:10.1093/cercor/bhg130

Merchant H, Battaglia-Mayer A, Georgopoulos AP. 2004b. Neural responses in motor cortex and

area 7a to real and apparent motion. *Exp Brain Res* **154**:291–307. doi:10.1007/s00221-003-

1664-5

Merchant H, Battaglia-Mayer A, Georgopoulos AP. 2001. Effects of Optic Flow in Motor Cortex

and Area 7a. *J Neurophysiol* **86**:1937–1954. doi:10.1152/jn.2001.86.4.1937

Michaels JA, Dann B, Scherberger H. 2016. Neural Population Dynamics during Reaching Are

Better Explained by a Dynamical System than Representational Tuning. *PLoS Comput Biol*

12:1–22. doi:10.1371/journal.pcbi.1005175

Moran DW, Schwartz AB. 1999. Motor Cortical Representation of Speed and Direction During

Reaching. *J Neurophysiol* **82**:2676–2692. doi:10.1152/jn.1999.82.5.2676

Omrani M, Murnaghan CD, Pruszynski JA, Scott SH. 2016. Distributed task-specific processing of

somatosensory feedback for voluntary motor control. *Elife* **5**:1–17. doi:10.7554/eLife.13141

Paninski L, Fellows MR, Hatsopoulos NG, Donoghue JP. 2004. Spatiotemporal Tuning of Motor

Cortical Neurons for Hand Position and Velocity. *J Neurophysiol* **91**:515–532.

doi:10.1152/jn.00587.2002

Parthasarathy A, Herikstad R, Bong JH, Medina FS, Libedinsky C, Yen SC. 2017. Mixed

selectivity morphs population codes in prefrontal cortex. *Nat Neurosci* **20**:1770–1779.

doi:10.1038/s41593-017-0003-2

Pesaran B, Nelson MJ, Andersen RA. 2010. A Relative Position Code for Saccades in Dorsal

Premotor Cortex. *J Neurosci* **30**:6527–6537. doi:10.1523/jneurosci.1625-09.2010

Pesaran B, Nelson MJ, Andersen RA. 2006. Dorsal Premotor Neurons Encode the Relative

Position of the Hand, Eye, and Goal during Reach Planning. *Neuron* **51**:125–134.

doi:10.1016/j.neuron.2006.05.025

Port NL, Kruse W, Lee D, Georgopoulos AP. 2001. Motor Cortical Activity during Interception of

Moving Targets. *J Cogn Neurosci* **13**:306–318. doi:10.1162/08989290151137368

Pouget A, Sejnowski TJ. 1997. Spatial transformations in the parietal cortex using basis

functions. *J Cogn Neurosci* **9**:222–37. doi:10.1162/jocn.1997.9.2.222

Pruszynski JA, Kurtzer I, Nashed JY, Omrani M, Brouwer B, Scott SH. 2011. Primary motor

cortex underlies multi-joint integration for fast feedback control. *Nature* **478**:387–390.

doi:10.1038/nature10436

Rao NG, Donoghue JP. 2014. Cue to action processing in motor cortex populations. *J*

Neurophysiol **111**:441–453. doi:10.1152/jn.00274.2013

Rigotti M, Barak O, Warden MR, Wang X-J, Daw ND, Miller EK, Fusi S. 2013. The importance of

mixed selectivity in complex cognitive tasks. *Nature* **497**:585–90. doi:10.1038/nature12160

Rizzolatti G, Scandolara C, Matelli M, Gentilucci M. 1981. Afferent properties of periarcuate

neurons in macaque monkeys. II. Visual responses. *Behav Brain Res* **2**:147–163.

doi:10.1016/0166-4328(81)90053-X

Russo AA, Khajeh R, Bittner SR, Perkins SM, Cunningham JP, Abbott LF, Churchland MM. 2020.

Neural Trajectories in the Supplementary Motor Area and Motor Cortex Exhibit Distinct

Geometries, Compatible with Different Classes of Computation. *Neuron* **107**:745-758.e6.

doi:10.1016/j.neuron.2020.05.020

Saxena S, Russo AA, Cunningham JP, Churchland MM. 2022. Motor cortex activity across

movement speeds is predicted by network-level strategies for generating muscle activity.

Elife **11**:1–31. doi:10.7554/eLife.67620

Schaffelhofer S, Scherberger H. 2016. Object vision to hand action in macaque parietal,
premotor, and motor cortices. *Elife* **5**:1–24. doi:10.7554/eLife.15278

Scott SH, Kalaska JF. 1997. Reaching Movements With Similar Hand Paths But Different Arm
Orientations. I. Activity of Individual Cells in Motor Cortex. *J Neurophysiol* **77**:826–852.
doi:10.1152/jn.1997.77.2.826

Sergio LE, Hamel-pa C, Kalaska JF, Lauren E, Hamel-pa C, John F. 2005. Motor Cortex Neural
Correlates of Output Kinematics and Kinetics During Isometric-Force and Arm-Reaching
Tasks 2353–2378. doi:10.1152/jn.00989.2004.

Sobinov AR, Bensmaia SJ. 2021. The neural mechanisms of manual dexterity. *Nat Rev Neurosci*
22:741–757. doi:10.1038/s41583-021-00528-7

Stavisky SD, Kao JC, Ryu SI, Shenoy K V. 2017. Motor Cortical Visuomotor Feedback Activity Is
Initially Isolated from Downstream Targets in Output-Null Neural State Space Dimensions.
Neuron **95**:195-208.e9. doi:10.1016/j.neuron.2017.05.023

Sun X, O'Shea DJ, Golub MD, Trautmann EM, Vyas S, Ryu SI, Shenoy K V. 2022. Cortical
preparatory activity indexes learned motor memories. *Nature* **602**:274–279.
doi:10.1038/s41586-021-04329-x

Sussillo D, Churchland MM, Kaufman MT, Shenoy K V. 2015. A neural network that finds a
naturalistic solution for the production of muscle activity. *Nat Neurosci* **18**:1025–1033.
doi:10.1038/nn.4042

Suway SB, Schwartz AB. 2019. Activity in Primary Motor Cortex Related to Visual Feedback. *Cell
Rep* **29**:3872-3884.e4. doi:10.1016/j.celrep.2019.11.069

Tkach D, Reimer J, Hatsopoulos NG. 2007. Congruent Activity during Action and Action
Observation in Motor Cortex. *J Neurosci* **27**:13241–13250. doi:10.1523/JNEUROSCI.2895-
07.2007

Vyas S, O'Shea DJ, Ryu SI, Shenoy K. 2020. Causal role of motor preparation during error-driven
learning_Suppl. *Neuron* **106**:1–11. doi:10.1016/j.neuron.2020.01.019

Wang M, Montanède C, Chandrasekaran C, Peixoto D, Shenoy K V., Kalaska JF. 2019. Macaque
dorsal premotor cortex exhibits decision-related activity only when specific stimulus–
response associations are known. *Nat Commun* **10**. doi:10.1038/s41467-019-09460-y

Wang T, Chen Y, Cui H. 2022. From Parametric Representation to Dynamical System: Shifting
Views of the Motor Cortex in Motor Control. *Neurosci Bull*. doi:10.1007/s12264-022-00832-x

Wang T, Chen Y, Zhang Y, Cui H. 2024. Multiplicative joint coding in preparatory activity for
reaching sequence in macaque motor cortex. *Nat Commun* **15**:3153. doi:10.1038/s41467-
024-47511-1

Zimnik AJ, Churchland MM. 2021. Independent generation of sequence elements by motor
cortex. *Nat Neurosci* **24**:412–424. doi:10.1038/s41593-021-00798-5