Boreal moss-microbe interactions are revealed through metagenome assembly of novel

bacterial species

Sarah Ishak[13*+], Jonathan Rondeau-Leclaire[1+], Maria Faticov[123], Sébastien Roy[12], and Isabelle

Laforest-Lapointe[123*]

[1]Département de biologie, Université de Sherbrooke, Sherbrooke, Québec, Canada

[2]Centre SÈVE, Université de Sherbrooke, Sherbrooke, Québec, Canada

[3]Centre d'Étude de la Forêt, Université du Québec à Montréal, Montréal, Québec, Canada

[+]These authors contributed equally

*Co-corresponding authors: isabelle.laforest.lapointe@gmail.com ;

s.isaac1905@protonmail.com

ORCID NUMBERS

Sarah Ishak   (0000-0003-3412-3642)

Jonathan Rondeau-Leclaire   (0000-0002-4433-3566)

Maria Faticov   (0000-0001-8206-9332)

Sébastien Roy   (0000-0001-7183-0691)

Isabelle Laforest-Lapointe   (0000-0003-3630-9475)

Abstract (max 200 words)

Moss-microbe interactions contribute to ecosystem processes in boreal forests. Yet, how host-specific characteristics and the environment drive the composition and metabolic potential of moss microbiomes is still poorly understood. In this study, we use shotgun metagenomics to identify the taxonomy and metabolic potential of the bacteria of four moss species of the boreal forests of Northern Québec, Canada. To characterize moss bacterial community composition and diversity, we assembled the genomes of 110 potentially novel bacterial species. Our results highlight that moss genus, species, gametophyte section, and to a lesser extent soil pH and soil temperature, drive moss-associated bacterial community composition and diversity. In the brown gametophyte section, two *Stigonema* spp. showed partial pathway completeness for photosynthesis and nitrogen fixation, while all brown-associated Hyphomicrobiales had complete assimilatory nitrate reduction pathways and many nearly complete carbon fixation pathways. Several brown-associated species showed partial to complete pathways for coenzyme M and F420 biosynthesis, important for methane metabolism. In addition, green-associated Hyphomicrobiales (*Methylobacteria* spp.) displayed potential for the anoxygenic photosystem II pathway. Overall, our findings demonstrate how host-specific characteristics and environmental factors shape the composition and metabolic potential of moss bacteria, highlighting their roles in carbon fixation, nitrogen cycling, and methane metabolism in boreal forests.

Introduction

Boreal forests are one of Earth's largest biomes, covering 29% of land surfaces [1] and storing at least 30% of global carbon [2]. Within this ecosystem, a diverse understory vegetation thrives, consisting of vascular plants, lichens, and bryophytes (small non-vascular plants including liverworts, hornworts, and mosses). Ubiquitous in boreal forests, mosses are crucial contributors to key ecosystem processes, notably to nitrogen (N) fixation and carbon (C) sequestration [3], which are also facilitated by moss symbiotic microorganisms [4, 5]. Moss microbes contribute around 50% of N inputs to boreal forests, play crucial roles in facilitating C sequestration in peatlands, and facilitate host adaptation in degraded landscapes [6]. Moss-associated microbial communities are notably dominated by bacteria from the Proteobacteria, Acidobacteria, Actinobacteria, and Cyanobacteria phyla [7–9]. Driven by the pressures of accelerating climate change, boreal ecosystems face a temperature increase of ~2.5ºC before the end of the century, which could alter moss-microbe interactions with major implications for global N [10] and C cycles [11]. In addition, boreal forests are often harvested intensively for forestry or disturbed by mining activities [12, 13]. Considering the strong pressures of climate change on boreal forests combined with the ever-increasing anthropogenic impacts on land use, improving our understanding of the drivers and functions of moss-microbe interactions is imperative to mitigate the risks of altering this precious habitat [14, 15].

Moss microbial communities can be affected by many local abiotic and biotic factors [8, 16]. For example, soil pH and soil temperature have been shown to structure bacterial communities associated with *Sphagnum* moss species [9, 17, 18]. Phylogenetic relationships and host species are also important drivers of moss bacterial community diversity and community composition [5, 9, 20]. While different moss species have been shown to associate with contrasting bacterial communities, closely related moss species are likely to

harbor more similar microbial taxa than more distantly related ones [18, 21, 22]. This

convergence can be attributed to shared host physiological traits and ecological niches.

For example, Holland-Moritz *et al.* [23], showed that phylogenetic relationship between

Sphagnum species explained up to 65% of the variation in bacterial community composition.

Interestingly, moss species identity also influences the effects of environmental factors on

bacterial diversity and community composition [18]. While previous studies have often

focused on characterizing the microbial taxonomy and functions of dominant boreal mosses

such as *Pleurozium schreberi* [24] and peat mosses of the genus *Sphagnum* [25], scant

research is available on other moss genera such as *Polytrichum* and *Dicranum* (but see [26]),

which are common in Canadian boreal forest [27]. Importantly, these taxa are key candidates

for remediation and restoration efforts as they often thrive in degraded landscapes [28]. Thus,

there is an important knowledge gap on moss-microbe interactions beyond common moss

genera. Considering the urgency to develop conservation strategies for bryophytes, it is

crucial to deepen our understanding of moss-microbe interactions and their functional roles in

unaltered boreal ecosystems by investigating understudied moss species, their phylogenetic

relationships, and local environmental factors.

Beyond moss species identity, physiological differences within a single host, such as

disparities in tissue types along the moss gametophyte, could also influence moss-microbe

interactions. Gametophytes are the dominant life stage of mosses, separated along their

vertical gradient in two distinct sections: a living "green" section and an underlying, decaying

"brown" section [29]. These sections can differ in microbial diversity, as well as in the

composition and functional potential of their microbial communities. [30, 31]. The green

section can harbor photosynthetically active microbes [32, 33] and is associated with N-

fixing cyanobacteria [34]; it is often the focus of studies on moss-bacteria associations.

However, the brown section is much less studied even though it is thought to be an important

source of nutrients for specific groups of microbes, such as bacteria [33, 35]. Notably, green and brown sections of the moss gametophyte are exposed to distinct microenvironmental conditions [36], whereby variations in light, soil pH, moisture, and temperature can impact moss microbial diversity and functions. Overall, to gain a detailed overview of moss-microbe interactions, it is necessary to measure how gametophyte sections assemble divergent microbial communities and functional profiles, and whether these differences are consistent among moss species as well as along environmental gradients.

In recent years, metagenome-resolved community profiling became a widespread approach to estimate environmental microbial composition and functions [37]. The advent of whole metagenome sequencing (WMS) revolutionized the field of microbial ecology, making it possible to assemble metagenome-assembled genomes (MAGs) and thus delve into the so-called "dark matter" of microbiomes [38]. By leveraging the entire genomic material of a sample, WMS not only enables the discovery of novel or uncultured microbial species; it also provides a more complete picture of the metabolic potential of community members [39]. WMS is therefore particularly useful in characterizing understudied microbiomes, such as that of boreal mosses. The discovery of novel bacterial species, along with an improved understanding of their metabolic potential in different gametophyte sections and environmental conditions, is important for predicting how these interactions will shift as climate change and human activities alter boreal ecosystems.

Here, we measure the influence of moss identity, gametophyte section, and abiotic environment on moss bacterial communities and their metabolic potential. We focus on four boreal moss species in the Eeyou Istchee (James Bay) region of Québec, Canada: *Polytrichum juniperinum*, *P. commune*, *P. piliferum*, and *Dicranum undulatum* (Fig. 1ab). We use WMS and a combination of cutting-edge bioinformatic tools to (i) characterize moss-microbe interactions, (ii) assemble metagenomes, and (iii) identify novel bacterial species

116   associated with boreal mosses. Our objectives are to (1) characterize the diversity,

117   community composition, and metabolic potential of moss-associated bacteria; and

118   (2) quantify the relative importance of moss species, gametophyte section, soil pH, and soil

119   temperature as drivers of moss-bacteria interactions.

Results and Discussion

Bacterial community composition and diversity are driven by moss identity, gametophyte section, soil pH, and soil temperature

First, we investigated the influence of moss genus, moss species (nested within moss genus), gametophyte section, soil pH, and soil temperature on bacterial community composition and diversity (Table 1a). We then narrowed our focus on the three *Polytrichum* spp. (excluding *Dicranum undulatum*; Table 1b). This sequential approach allowed us to disentangle whether closely related moss species are likely to harbor more similar microbial taxa than more distantly related ones. Both moss genus and species (nested within genera) were strong drivers of bacterial community composition (Genera: $R^2 = 13.8\%$, p-value = 0.001; Species: $R^2 = 12.3\%$, p-value = 0.001; Fig. 2a, Table 1a). Notably, when restricting the analysis to the three *Polytrichum* spp., the effect of moss species on bacterial community composition remained significant ($R^2 = 19.7\%$, p-value = 0.001; Table 1b). Thus, our results highlight that differences in bacterial community composition are driven by both moss genus and, within these genera, by moss species. These results are in line with previous studies in which moss phylogeny was shown to be an important driver of bacterial community composition [4, 8, 25]. This finding illustrates a potential co-evolutionary relationship between mosses and some of the bacteria members within the community [17, 40]. Yet, more studies are needed to fully understand the nature and mechanisms driving these genus- and species-specific interactions. We also found that bacterial community composition differed between green and brown gametophyte sections, regardless of whether *D. undulatum* samples were included in the model or not ($R^2 = 12.7\%$, p-value = 0.001; $R^2 = 17.2\%$, p-value = 0.001, respectively; Fig. 2b, Table 1ab). This variation between sections can be explained by differences in physiology [41] and micro-environmental conditions (e.g., soil pH, moisture, temperature) [36]. In addition, we observed a significant, albeit small, interaction between gametophyte

section and moss genus ($R^2 = 2.6\%$, p-value = 0.001; Table 1a), as well as moss species ($R^2 =$

1.3%, p-value = 0.042; Table 1b), highlighting that the effect of gametophyte section on

bacterial community composition differs among moss species. In both models, whether

D. undulatum was included or not, we found that soil temperature (with *Dicranum:*

$R^2 = 7.4\%$, p-value = 0.001; without *Dicranum*: $R^2 = 6.9$, p-value = 0.001) and pH

($R^2 = 5.1\%$, p-value = 0.001 and $R^2 = 4.1\%$, p-value = 0.001 respectively) were significant

drivers of bacterial community composition. Interestingly, the effect of abiotic factors on

community composition was weaker than that of moss genus and species (Table 1). Finally,

the interaction of soil pH and temperature affected moss bacterial community composition

(Table 1), in coherence with the important roles of abiotic factors in structuring moss-

microbe interactions across moss species [9, 17, 18].

Bacterial alpha diversity differed between moss genera, with alpha diversity (bacterial

Shannon diversity) being higher for *Polytrichum* than for *D. undulatum* ($\chi^2 = 8.6$, p-value =

0.003; Table 2a, Fig. 2c). However, we did not detect a significant association between moss

species and bacterial alpha diversity, neither when nesting species within moss genera

($\chi^2 = 0.07$, p-value = 0.97; Table 2a, Fig. 2c) nor when narrowing the model to *Polytrichum*

spp. ($\chi^2 = 0.08$, p-value = 0.96; Table 2b, Fig. 2c). Of note, alpha diversity in the green

section was consistently lower than in the brown section of the gametophyte. This pattern

was consistent between moss genera ($\chi^2 = 29.2$, p-value < 0.001; Fig. 2c, Table 2a) and

among *Polytrichum* spp. ($\chi^2 = 35.7$, p-value < 0.001; Fig. 2c, Table 2b). This could be

explained by (i) a phenomenon of niche dominance of a few taxa in the green section or by

(ii) the proximity of the brown section to the soil, which is known to be extremely rich in

microorganisms. As such, the brown section could behave as an interface in which both soil

and moss-associated bacteria can thrive. This section is also mostly shaded from UV and may

contain less chlorophyll than the above active, green section [42]. Finally, the underlying

brown section tends to have higher moisture contents due to the boundary layer effect (i.e., the physical phenomenon in which wind velocity is reduced); an effect that minimizes moisture loss [43]. Among abiotic factors, soil temperature was the only factor showing a significant association with moss bacterial alpha diversity, evident in both models, with and without *D. undulatum* ($\chi^2 = 6.3$, p-value = 0.01 and $\chi^2 = 6.6$, p-value = 0.01; Fig. 2c, Table 2ab). Furthermore, we did not detect a significant effect of the interaction between gametophyte section and moss genus or species on bacterial alpha diversity (Table 2).

In summary, these results show that moss genus, gametophyte section, soil pH, and soil temperature are important drivers of moss bacterial community composition and diversity, but that moss species only affects the former. Our work further demonstrates that green and brown sections within a moss gametophyte differ in bacterial community composition (Fig. 2d) and alpha diversity, emphasizing the relationship between moss characteristics and the associated bacterial communities.

Metagenome assembly reveals that novel bacteria constitute most of moss bacterial community

By assembling metagenomes, we discovered 110 potentially novel bacterial species whose genomes were not represented in reference databases. Specifically, we performed metagenome co-assembly, binning, and dereplication, which produced 157 non-redundant species-level MAGs. We filtered out MAGs with quality scores (QS) ≤ 0.50 (QS = completeness - 5 $\times$ contamination), yielding a subset of 118 MAGs of medium to high quality (82.46 ± 11.68 % completeness, 1.97 ± 1.26 % contamination). In order to estimate which MAGs belong to undescribed species, we compared them to every known representative bacterial genome using a combination of tools based on mash distance and average nucleotide identity (see *Methods*). This resulted in a set of draft genomes from 110

putative novel bacterial species spanning 23 orders across nine phyla. Hereafter, we refer to

this set as the novel species-level MAGs (nMAGs), in that they constitute genomes that are

non-redundant with any published reference genome at the species level. These nMAGs have

a mean length of 4.0 ± 1.2 Mbp, QS of 0.73 ± 0.13, and L50 of 11.9 kb $\pm$ 10.5. Full quality

statistics of the nMAGs, including the number of contigs and proportion of GC content, are

reported in Supplementary Data 1 and summarized in Fig. 3.

To gain a more detailed understanding of the microbial communities of boreal

mosses, we added the 110 nMAGs to the 85,205-reference genome GTDB used for

community profiling, resulting in an average 4.3-fold increase in sample containment (a

measure of proportion of sample k-mers included in the minimal reference genomes set that

encompasses all k-mers in a sample; see Methods; Supplementary Fig. S1). This increase

demonstrates the effectiveness of using *de novo* metagenome assembly to identify and

describe previously unknown microbial species, which is crucial for environmental datasets,

and to increase the proportion of sample information to perform ecological analyses. As

detailed in Methods, all results presented here (including the above section on diversity) rely

on community composition estimation performed using the improved reference database.

Of note, we modified the GTDB taxonomy used to annotate our genomes according to

the latest nomenclature in the *List of Prokaryotic names with Standing in Nomenclature*

(LPSN) [44]. Specifically, GTDB classifies Hyphomicrobiales (LPSN nomenclature) as

Rhizobiales, so we corrected our annotations to use Hyphomicrobiales throughout. GTDB

also uses the order Cyanobacteriales, which encompasses species of many different orders

(e.g., Oscillatoriales, Synechococcales, Chroococcales, etc.) according to LPSN. Since every

member of the class Cyanobacteria found in our samples belonged to the Nostocaceae family,

we renamed the Cyanobacteriales order to Nostocales, following LPSN.

We recovered seven MAGs belonging to the order Baltobacterales [45] from the

phylum Eremiobacterota [46] (Fig. 3). Despite the presence of 71 Baltobacterales species-

representative genomes in the GTDB R214 database (28th April 2023), all seven recovered

genomes were identified as nMAGs. Many Eremiobacterota species were recently described

by metagenome assembly from a number of boreal ecosystems such as Alaskan mosses [18],

Canadian acidic springs [47], and Swedish peatlands [48]. Baltobacterales were highly

prevalent in our samples, detected in 109/131 (~83%) samples, and two *Nyctobacter* MAGs

were of high quality (QS = 0.95 and 0.91 respectively; Supplementary Data 1). While there is

little information on the ecological traits of Eremiobacterota, several genera within this group

were shown to have the capacity for C fixation [39]. Our study supports this finding, as each

Eremiobacterota we detected showed partial completeness in most prokaryotic carbon

fixation pathway considered (Supplementary Fig. S2). Pessi *et al.* [50] detected *nif*

homologues in the Eremiobacterota species Candidatus *Lamibacter sapmiensis*. However, we

only found partial completeness of an assimilatory nitrate reduction module (M00531) in

Candidatus *Velthaea versatilis*, whose nitrogen reduction capacity has already been well

described [49]. Interestingly, all Eremiobacterota nMAGs except *Tumulicola* BrAA6 can

encode either partial or complete pathways for anoxygenic photosynthesis, supporting

findings by Ward *et al.* [45]. Additionally, we detected partial completeness of the methane

oxidation pathway across all detected species within this phylum, supporting the findings of

Holland-Moritz *et al.* [51], who identified only two cases of the methane oxidation marker

gene *pmoA*. Moreover, Ray *et al*. [52] suggested Eremiobacterota might use atmospheric

chemosynthesis by assimilating formaldehyde through the serine pathway. Our results

support this hypothesis, as all Eremiobacterota we detected exhibited partial completeness of

this pathway (see Supplementary Fig. S2). Our study thus contributes novel Eremiobacterota

genomes of potentially undiscovered species and information about their metabolic potential,

and further supports existing knowledge about moss-associated members of this clade.

Bacterial species associated with moss species

In order to identify differentially abundant taxa across groups of samples, we used ANCOM-

BC2 [53], which estimates $\log_2$-fold changes in absolute abundances between sample groups

through pairwise comparisons. To assess differences in abundance across moss species, we

used the Dunnett's test, with *D. undulatum* as a reference group. *D. undulatum* was selected

because of its contrasting bacterial community composition, which served as a baseline for

evaluating differential abundant bacteria among moss species (Fig. 4a). A total of 28 species

were found to be differentially abundant across the three *Polytrichum* species compared to

D. undulatum (Fig. 4a; Supplementary Fig. S3). Of these 28 bacterial species, 24 were

nMAGs, underlining the contribution of the assembly process to the depth of our analyses.

We also performed a differential abundance test at the order level in a pairwise

manner (i.e., without a reference host). Interestingly, we found that the order Baltobacterales

was more abundant in *P. commune* by 2.7 to 3.3 $\log_2$-folds compared to all other moss species

(Supplementary Fig. S3). Previous metagenomic studies showed that Baltobacterales encode

genes necessary for hopanoid synthesis (a biochemical process in which hopanoid, steroid-

like compounds, are produced) [49]. Hopanoid molecules were shown to contribute to

bacterial tolerance to low pH soils [54], temperature stress [49], microbial defense [56], and

are linked to N-fixation [57, 58]. Notably, *P. commune* is known for its antibacterial

properties [60] and specimens from this moss species were sampled from acidic soils

(Supplementary Data S2) in our study. The stress-tolerance properties of hopanoid structures

in Baltobacterales could provide an adaptive advantage for colonization and persistence of

bacteria in acidic and anti-microbial habitats as found in *P. commune* mosses.

268   Two *Stigonema* spp. (Nostocales) were differentially abundant in *P. juniperinum* and

269 *P. piliferum* (relative to *D. undulatum*), but not in *P. commune* (Fig. 4a). One major

270 difference between *Polytrichum* species is the habitat in which *P. commune* was growing in.

271 We found that *P. commune* grew in a significantly more shaded environment than the other

272 *Polytrichum* species (Supplementary Data S2), and *Stigonema* may be more sensitive to these

273 differences in shading. Indeed, Jean *et al.* (2020) found that Nostocaceae abundance was

274 negatively associated with canopy cover [8]. Furthermore, the aforementioned anti-microbial

275 properties of *P. commune* could select against *Stigonema* spp. growth. Finally, *Stigonema*,

276 known for having an alternative nitrogenase, is also common in boreal lichens (*Stereocaulon*)

277 growing near *Polytrichum* and *D. undulatum*. [57]. The proximity of *Stigonema* to these moss

278 species, combined with its unique functional adaptation, such as its alternative nitrogenase,

279 may influence the abundance or composition of the bacterial communities associated with

280 these mosses. In general, all Acetobacterales and Terriglobales species were positively

281 associated with *Polytrichum* species (Fig. 4a). Of note, Terriglobales, one of the most overall

282 relatively abundant species in our dataset (mean $20.8 \pm 10.8\%$ of attributed sequences across

283 all samples, Fig. 2d), were also found to be highly present in sub-arctic and boreal peatland

284 moss microbiomes [62].

285   Three species from the orders Solirubrobacterales, Steroidobacterales, and

286 Streptosporangiales were found to be much more abundant in *D. undulatum* compared to

287 *Polytrichum* hosts (3.20 to 4.63 $\log_2$-folds, Fig. 4a). The most closely related species from

288 these orders were incidentally all species identified through metagenome assembly and were

289 described in soil and roots in non-boreal ecosystems [52, 63]. A potential reason for the

290 differences in bacterial communities between *Dicranum* and *Polytrichum* genera could be

291 due to host environmental preferences (*D. undulatum* preferring moist soils), as well as moss

292 N and phosphorus (P) content [64]. Although our results show that moss-associated bacterial

communities are influenced by moss identity, experimental studies are needed to disentangle

the simultaneous effects of biotic and abiotic conditions on moss-associated bacteria and on

their contribution to ecosystem functions.

Bacterial species associated with brown and green gametophyte sections

We identified 44 species as differentially abundant between green and brown gametophyte

sections, of which 18 are nMAGs (Fig. 4b). These species are hereafter described as

"associated" to the section in which they were more abundant. Our work identified 27 species

as brown-associated, including all nine *Bradyrhizobia*, thus suggesting a strong specificity for

brown gametophytes for members of this genus (Fig. 4b).

303   Bacterial species with the highest log_2-fold change in brown gametophyte sections

included three nMAGs from the genus *Edaphobacter*, uncultured species from the order

Solirubrobacterales (*VAYN01*) and Terriglobales (*CAIQPK01*), as well as three uncultured

species from the order Streptosporangiales (*Bog-532 sp019240215* and two *Trebonia*

nMAGs; Fig. 4b). Various *Edaphobacter* species, such as *E. lichenicola sp. nov.,*

Edaphobacter modestus gen. nov., and *Edaphobacter aggregans sp. nov.*, were previously

described from alpine forest soils [65, 66]. Although Solirubrobacterales are poorly studied,

researchers have detected members of this order in diverse ecosystems including agricultural

lands [67] and Antarctic soils [68]. Several other brown-associated taxa, such as *Rudaea* spp.,

CAIMXF01, *Trebonia*, and *Acidisarcina*, are also mainly associated with soils [69–71].

Finally, two *Stigonema* species (cyanobacteria of the order Nostocales) were differentially

abundant in brown gametophytes (Fig. 4b). In line with our findings, Renaudin *et al*. [10]

found that cyanobacteria biomass increased towards the bottom of moss gametophyte,

peaking at approximately 2-3 cm from the apex. In their study on the bacterial communities

of two *Racomitrium* moss species, Klarenberg *et al.* [72] demonstrated that there is little

transmission of bacteria from soil to moss gametophytes [69]. Meanwhile, Escolástico-Ortiz *et al.* [73] showed overlap between soil bacterial communities and the moss *Racomitrium lanuginosum*. However, the authors also revealed the presence of a core microbiome for moss gametophytes and soil, with each harbouring distinct, non-overlapping core bacterial communities. While we did not characterize the bacterial community of the surrounding soil, our finding of a number of soil species in the brown gametophyte section suggests this habitat may be colonized by both soil and moss-associated microbes.

In the green gametophyte section, nMAGs from the genera *CAHJXF01*, *CAHJXG01*, *Lichenicola*, and *Palsa-465* had the highest change in absolute abundance (log$_2$-fold > 1.29; Fig. 4b) compared to the brown gametophyte section. Hay *et al*. [75] described MAGs of *CAHJXF01* and *CAHJXG01* from high Arctic glaciers, containing the gene form I RubisCo – an important gene for C fixation. We also found certain species of Hyphomicrobiales (Rhizobiales) to be green-associated, specifically from the genera *Methylobacteria*, *Lichenihabitans*, and *RH-AL1*, as well as all species of the genus *Bradyrhizobium* (Fig 4b). *Methylobacteria*, for example, were found in the core microbiome of boreal shrubs [76] and were shown to colonize non-boreal mosses (e.g., *Didymodon tectorum* and *Funaria hygrometrica* [77, 78]). Similarly, *Lichenihabitans minor* and *RH-AL1* spp. were shown to harbor genes associated with methanol consumption [74]. Although *Lichenihabitans minor* tends to be associated with lichens, the detection of this species in our samples is not surprising, given the prevalence of lichens in the habitat of boreal mosses [79]. *Lichenicola* and *Lichenicoccus* species, eponymously, can associate with lichens. For example, *Lichenicola cladoniae* was previously found in Antarctic lichens [80]. Overall, through our analysis, we observed the association of several species from genera such as *CAHJXF01*, *CAHJXG01*, *Methylobacteria*, *Lichenihabitans*, and *RH-AL1* spp. with the green gametophyte

section. Future studies would be warranted to quantify their impact on C fixation and

methanol consumption in boreal forests.

Bacterial metabolic potential differs in green and brown gametophyte sections

Next, we investigated the metabolic potential of the gametophyte section-associated bacteria

by analyzing which pathways their genomes have the potential to express. We focused on

four pathway groups relevant to boreal ecosystem functioning: photosynthesis, nitrogen

metabolism, methane metabolism, and carbon fixation.

First, we found that two brown-associated *Stigonema* species, also known as true-

branched filamentous cyanobacteria, had partial pathway completeness for photosystems I

and II (*Stigonema sp903903925* = 83.3% and *Stigonema* BrZ1.MAG = 66.7%) (Fig. 5;

Supplementary Data S3). These were the only two differentially abundant species capable of

at least partially encoding these pathways. Moreover, two *Stigonema* spp. stood out because

of their partially complete N fixation (nifHDK + vnfDKGH) pathway (*Stigonema*

sp903903925 = 75% and *Stigonema* BrZ1.MAG = 50% completeness) (Fig. 5;

Supplementary Data S3). This supports a previous finding in feathermosses, which showed

Stigonema spp. to be important contributors to N fixation [81, 82]. *Stigonema* spp. also

showed partial pathway completeness in all methane modules relevant to methanogenesis,

and not methane oxidation. Although methanogenesis has not been shown to occur in any

organisms except archaea [83], partial completeness of several methanogenesis pathways

alludes to the potential for certain cyanobacteria to contribute to non-methanogenesis

methane production (Fig. 5; Supplementary Data S3). Overall, our findings suggest that

Nostocales species associated with the brown gametophyte section are photosynthetically

active and have the potential for nitrogen fixation [85, 86].

Second, we detected complete assimilatory nitrate reduction pathways in both green-

and brown-associated Hyphomicrobiales (Rhizobiales), in line with earlier findings on

nitrogen metabolic capacities of this clade (Fig. 5; Supplementary Data S3).

Hyphomicrobiales taxa were previously shown to be N-fixers for nodule-producing

leguminous plants [87] and Sphagnum mosses [17]. Interestingly, we found that nitrogen

metabolism pathways were not significantly more complete in bacteria associated with either

green or brown gametophyte sections. This suggests that the potential of the bacterial

community to encode these pathways is not influenced by section type (Supplementary Table

S1). Furthermore, most methane metabolic pathways were also partially complete in both

green and brown-associated Hyphomicrobiales genomes (Fig. 5; Supplementary Data S3).

For example, the methanogenesis acetate pathway was significantly more complete in brown-

associated species of *Bradyrhizobium* (p-value = 0.021; Fig. 5; Supplementary Table S1).

This aligns with the results of a previous study on *Sphagnum* mosses, which found that

methanotrophy is primarily carried out by Hyphomicrobiales species [88]. Additionally, two

of the three green-associated Methylobacteria showed potential to fully encode the

anoxygenic photosystem II pathway, whereas the third one (one of the nMAGs) can only

encodes 50% of the required genes (Fig. 5; Supplementary Data S3). Of note, the only

photosynthesis pathway whose completeness was significantly associated with the green-

associated species was also the anoxygenic photosystem II (p-value = 0.008 Supplementary

Table S1). This is in line with recent findings that aerobic anoxygenic photosynthetic bacteria

are present among plant phyllosphere microbial communities [89], including certain species

of Methylobacteria [90]. Next, not a single green-associated Hyphomicrobiales encodes any

gene of the phosphate acetyltransferase-acetate kinase pathway (carbon fixation), whereas

among brown-associated Hyphomicrobiales, six *Bradyrhizobium* spp. encode 100% of the

required genes (Fig. 5; Supplementary Data S3). In line with this finding, the phosphate

acetyltransferase-acetate kinase pathway was also the only pathway significantly more

complete in differentially brown-associated species (p-value = 0.012; Supplementary

Table S1). This pathway is part of a central metabolism of bacteria [91]; its high

completeness in *Bradyrhizobia* indicates the metabolic potential for C fixation [92]. To sum

up, our findings suggest that green- and brown-associated Hyphomicrobiales provide a range

of metabolic contributions such as C fixation, methane metabolism, and nitrate reduction to

their ecosystem.

Third, brown-associated Burkholderiales (*Caballeronia* spp.) showed potential for

denitrification activity, dissimilatory and assimilatory nitrate reduction, and a complete

nitrification pathway (Fig. 5; Supplementary Data S3). However, the only green-associated

Burkholderiales species (CAHJXF01 nMAG) lacked all three nitrification pathways

(denitrification, nitrification, and complete nitrification, Fig. 5). While we did not find

methane oxidation pathways in these species, they still showed completeness in several

methane metabolism pathways, such as formaldehyde assimilation (Fig. 5; Supplementary

Data S3). These findings suggest the distinct metabolic capabilities of Burkholderiales in

different sections of the gametophyte [93, 94]. Curiously, pathways for coenzyme M (p-value

= 0.015) and F420 biosynthesis (p-value = 0.169) were partially to fully complete in several

brown-associated species (Fig. 5; Supplementary Table S1 and Supplementary Data S3).

F420 is a costly co-enzyme to synthesize; nevertheless, Ney *et al*. [95] demonstrated it was

commonly synthesized by aerobic bacteria. Meanwhile, coenzyme M is found in

methanotrophic archaea and used by bacteria in metabolizing of different substrates, for

example alkenes [96], which are present in boreal forest soils [97]. More interestingly,

alkenes are known to be produced by bacteria associated with lichens [98]. In our sampling

sites, lichens frequently grew alongside mosses, indicating that some of the bacterial taxa may

be shared between the groups. Our results suggest that the presence of coenzyme M in

bacteria associated with brown gametophyte sections may be attributed to their utilization of

alkenes commonly found in lichens.

Overall, our findings demonstrate the diverse metabolic potential and adaptations of

bacterial species associated with different sections of moss gametophyte, shedding light on

their roles in N cycling, methane metabolism and C fixation in boreal forests.

Conclusion

In this study, using a whole metagenome sequencing approach, we demonstrated that moss

genus, moss species, gametophyte section, and, to a lesser extent, soil pH and soil

temperature are significant drivers of bacterial community composition and diversity

(Tables 1-2; Fig. 2). Through metagenomic assembly, we identified 110 potentially novel

bacterial species (Fig. 3) and characterized distinct bacterial communities associated with

four different moss species (Fig. 4). We found that bacterial communities in green and brown

gametophyte sections occupy unique functional niches (Fig. 5), with partial partial overlap.

Our results show that the brown gametophyte section, despite their senescent nature, can

support bacterial communities that contribute to ecological functions such as nitrogen

fixation, carbon fixation, and methane metabolism, while green-associated bacteria exhibit

potential for anoxygenic photosynthesis.

To build on these findings, several key areas for future research emerge. First,

validating the specific metabolic pathways of the 110 potentially novel bacterial species and

their contributions to processes like nitrogen fixation and carbon sequestration is important.

For example, experimental studies and transcriptomic studies could help to identify which

genes are actively transcribed under different environmental conditions. Second, exploring

the interactions between moss-associated bacteria, fungi, and other microbial members could

provide more insights into their joint roles in ecosystem functioning. Gaining a clear

understanding of moss metabolites could further elucidate the potential co-evolutionary or

symbiotic relationships between non-cyanobacterial microbial communities and mosses,

revealing how metabolites influence microbial interactions and ecosystem processes. To

summarize, our results contribute to our understanding of how moss-bacteria interactions

contribute to ecosystem functions, such as nutrient cycling and carbon sequestration,

emphasizing the importance of both green and brown gametophyte section-associated

bacteria in these processes.

Methods

<u>Study area</u>

This study was conducted within the Eeyou Istchee (James Bay) region of Northern Québec

(Fig. 1a; N51.7040721°, W76.0521982°). The sampling area is part of Québec's boreal forest

biome and is dominated by closed-crown forest stands of Jack pine (*Pinus banksiana*) and

black spruce (*Picea mariana*) [99]. The monthly average annual temperature of this region is

0.3□, while mean annual precipitation is 741 mm [100]. The soils are predominantly acidic

and nutrient poor [101].

<u>Study design and moss species</u>

Samples were taken from four acrocarpous (stems harbor archegonia at the top) moss species:

Polytrichum juniperinum, *Polytrichum commune*, *Polytrichum piliferum*, and *Dicranum*

undulatum (Fig. 1b), which are native to the Eeyou-Istchee region of Québec [28]. These

were sampled in September 2021, at five different locations (Fig. 1a). Three microsites, each

5 to 10 meters apart, were randomly selected and then, at each microsite, 3-by-3-inch moss

patches were sampled directly (Fig. 1a). At each microsite/species, we measured soil

temperature, by inserting the probe of the Fisherbrand^{TM} Envirometer^{TM} directly in the soil of

where the moss was sampled after it had been removed. Soil pH was measured in the

laboratory using the method described by Hendershot, Lalande, and Duquette [102]. Mosses

were first visually identified to species level in the field and then confirmed by microscopy.

For each moss species, samples were taken from both the brown and green sections of the

gametophyte, which were determined visually (Fig. 1c). To maintain the integrity of the

samples, the green and brown sections were separated while on site, using ethanol and DNA-

away treated tweezers and scissors. Samples were placed in sterile 50 mL tubes and

immediately stored at -20□ for the return trip to the laboratory, where they were stored at -

80□. A total of 131 samples were collected for brown and green gametophyte sections across

P. juniperinum ($N_{brown} = 24$, $N_{green} = 22$), *P. commune* ($N_{brown} = 9$, $N_{green} = 9$), *P. piliferum*

($N_{brown} = 26$, $N_{green} = 26$), and *D. undulatum* ($N_{brown} = 8$, $N_{green} = 7$) from five sites (see

Supplementary Table S2 for sample breakdown).

<u>DNA extraction and whole metagenome sequencing (WMS)</u>

Green and brown gametophyte sections were processed separately, but in the same manner.

DNA extraction of moss samples was performed as modified versions of the protocols used

by Holland-Moritz *et al.* [4] and Klarenberg *et al.* [16]. Using tweezers and scissors sterilized

with 80% ethanol solution and 'DNA Away' solution (Thermo Fisher Scientific, MA, USA),

0.25 g of each moss sample were weighed out and divided into two sterile 2 mL bead beating

tubes that also contained three 2.3 mM BioSpec sterilized stainless-steel beads. These tubes

were then frozen in liquid N_2 for approximately 30s and mosses were broken down using the

FisherbrandTM Bead Mill 24 Homogenizer for 30s at 1.75 m/s. DNA was subsequently

extracted using the QIAGEN DNEasy PowerSoil DNA Extraction Kit with some minor

changes. Each moss sample was divided into two 2 mL PowerBead extraction tubes. The kit

buffer and beads were added to each tube, along with 60 uL of C1 Solution from the kit.

All tubes with buffer were homogenized at 3.25 m/s for 30s and then at 2.40 m/s for 10

minutes using the FisherbrandTM Bead Mill 24 Homogenizer. After centrifugation as per the

extraction kit protocol, supernatant from both tubes of one sample were combined in one tube

with 250 uL of C2 Solution from the extraction kit. The rest of the extraction followed the

manufacturer's instructions. All extractions were performed under a Type A2 Bio Safety

Cabinet. Extracted DNA were subsequently quantified using the NanoDropTM One

Spectrophotometer (Thermo Fisher Scientific, MA, USA). Samples under 20 ng/uL or with

A260/230 ratios less than 1.8 were tested for presence of bacteria by PCR amplification of

the V4-V5 region of bacterial 16S rRNA using 515F (GTGCCAGCMGCCGCGGTAA) as

forward primer and 806R (GGACTACHVGGGTWTCTAAT) as reverse primer. The 25 uL

PCR reaction mix for these tests consisted of 5 uL of 1X GC Buffer, 0.5 uL of 10uM dNTPs,

0.5 uL each of 10uM forward and reverse primers, 0.25 uL HotStart DNA polymerase and

17.25 uL RNAse free MilliQ water. PCR thermocycling conditions were an initial

denaturation at 94□ for 3 minutes, followed by 35 cycles of: denaturation at 95□ for 30s,

annealing at 50□ for 30s, elongation at 72□ for 1-minute 30s; lastly there was a final

elongation step at 72□ for 10 minutes. All tested samples showed amplification at the

predicted length of 290 base pairs.

In total, 131 metagenomic libraries were generated. Samples were sequenced on a

single lane of Illumina NovaSeq6000 with $2 \times 150bp$ chemistry at Génome Québec

(Montréal, Canada).

Bioinformatic analyses

Samples preprocessing (quality control), assembly, MAG generation and annotation, as well

as novelty analyses, were performed on high-performance computer with the aid of SLURM

workload management for parallel processing. The pipeline for these steps is written in bash.

In the next steps, the R programming language was used for diversity and differential

abundance analyses, as well as production of figures and tables. R packages are listed in

Supplementary Table S4, along with all bioinformatic software (version and non-default

options, when applicable) outlined in the following methods. The complete workflow is

provided in Supplementary Fig. S4.

Sample quality control

The raw sample library size was $14.8 \times 10^6 \pm 2.2 \times 10^6$ reads. Raw reads were trimmed and

quality-controlled using Kneaddata 0.12.0, keeping reads with a minimum length of 50 bp

and an average quality score of 30 over a 4-nucleotide sliding window. At the time of the

experiment, all four moss species sampled in this study lacked reference genomes. Thus, host

DNA was partially removed by aligning reads against the closely related genome of

Physcomitrella patens, as well as the human genome GRCh38 to remove human

contamination, using Bowtie 2.4.4 (from within the Kneaddata pipeline). On average, clean

samples contained $10 \times 10^6 \pm 1.6 \times 10^6$ paired-end reads.

Choice of a co-assembly strategy

Metagenome assembly quality depends on depth of coverage of each genome within samples

[103]. A common strategy is to co-assemble samples based on biological similarities, by

pooling sequences submitted to the assembler, increasing the coverage of genomes, and

therefore retrieving more good quality genomes [104]. In this study, we pooled samples

(i.e., concatenated the *.fastq* metagenome files) taken from the same moss species,

gametophyte section, and microsite into 46 sample pools. We applied the pipeline described

in the following section identically to both assembly and co-assembly strategies.

Co-assembly generated 157 MAGs with completeness $> 50\%$ and contamination $< 10\%$,

compared to 139 through regular sample-level assembly. Given this, we focused on the

MAGs generated by co-assembly, which had a slightly higher average completeness,

increasing from 73.6% ± 16.5 to 74.2% ± 15.4%.

<u>Metagenome co-assembly and MAG generation</u>

Samples were co-assembled using MetaWRAP modules [105]. First, paired-end sequences

were assembled using metaSPAdes 3.15.4 and re-assembled unassembled reads using

MEGAHIT 1.2.9. Contigs were then binned with three binners (MetaBAT 2.12.1,

CONCOCT 1.1.0, and MaxBin 2.2.4). Afterwards, these three sets of bins were hybridized

with the MetaWRAP 1.3 *bin_refinement* module to produce a final set of refined bins for

each co-assembly. Bin quality was then assessed using CheckM 1.0.18 and bins with

completeness $< 50\%$ and contamination $> 10\%$ were excluded [106], yielding a set of

260 bins (5.7 ± 3.3 bins per co-assembly).

To obtain a unique set of species-level MAGs for the whole study, bins were pooled

from all co-assemblies and dereplicated them using dRep 3.4.0 [107]. dRep uses a two-step

clustering strategy, whereby bins are first clustered using Mash 2.3 [108]; then, within each

cluster, bins are dereplicated at a 95% average nucleotide identity (ANI) using

MUMmer 3.23 [109]. GUNC 1.0.5 [110] was used to identify chimeric MAGs, which were

then removed from the final set of species-level metagenome-assembled genomes (MAGs).

<u>MAG novelty analysis</u>

MAGs which potentially belonged to species with no published reference genome, and

therefore were absent from the reference database, were identified by applying three

bioinformatic validations. First, we taxonomically annotated MAGs using GTDB-Tk [111].

Second, we calculated the Mash distance between these 156 MAGs and the 31,332 non-

redundant bacterial and archaeal genomes from EnsemblBacteria. Third, since Mash distance

could be underestimated when comparing incomplete MAGs, we used skani 0.2.1 [112] to

compute ANI between these MAGs and the 85,205 GTDB species representative genomes.

MAGs were labelled as novel MAGs (nMAGs) if they had an ANI < 0.95 having a Mash

distance > 0.05 and no species-level taxonomy assignment. Additionally, after adding the

nMAGs to the reference genome index as described below, two were dropped because they

were not detected by Sourmash 4.7.0 [113]. nMAGs were only kept if they exhibited an

assembly quality score (defined as Completeness – 5 × Contamination) > 0.50, yielding a

final set of 110 nMAGs.

<u>nMAG annotations</u>

nMAGs were further annotated by predicting their coding sequences using Prodigal 2.6.3

[114] and passing these to the MicrobeAnnotator 2.0.5 [115] pipeline, generating an

estimation of their metabolic potential through pathway completeness reconstruction (see

Pathway completeness analysis below for further details). Finally, phylogeny was estimated

using PhyloPhlAn 3.0.3 [116] and phylogenetic trees built using ggtree 3.10.0 [117] and

ggtreeExtra 1.12.0, allowing a cross-validation with their attributed GTDB taxonomy.

nMAGs statistics (length, L50, number of contigs and GC contents) were assessed using

bbmap 38.86.

<u>Taxonomy characterization and abundance estimation of bacterial community</u>

Sample community composition and relative abundance at the species level was estimated

using Sourmash gather [113] with a *k*-mer size of 31, using the GTDB representative species

genome index [111] to which nMAGs were added. This approach also measures a sample's

containment, which represents the proportion of sample *k*-mers found in the smallest set of

reference genomes required to cover all sample *k*-mers with an exact match. Of note, *k*-mer-

based community profiling tools generate a *sequence* abundance table; these abundance

values are not normalized for genome length (as opposed to marker-gene-based tools),

making them distinct from taxonomic abundance [118]. However, the necessity to introduce

nMAGs in the reference database precludes the use of marker-gene referencing for taxonomic

assignment.

Containment was used to estimate the improvement in sample coverage caused by the

addition of nMAGs to the GTDB R214 reference index. The GenBank 2022-03 reference

index (comprising 1.22 M microbial genomes) was also compared, and the containment

increase is reported for both scenarios (Supplementary Fig. S1). Finally, the sequence

abundance matrix was built into a phyloseq (version 1.46.0) object using the raw counts; this

object was then used as a starting point for all subsequent analyses.

<u>Diversity analyses</u>

All statistical analyses and plots were made using the open-access software R 4.2.1 [119].

For community composition (multivariate) analyses, uneven sequencing depth and

heteroskedasticity were addressed by applying variance stabilizing transformations (VST,

DESeq2 1.42 [120]) to the abundance matrix. This matrix was then used to estimate pairwise

Bray-Curtis dissimilarity between samples [121], and to compute principal coordinates using

the *capscale* function in the vegan package (version 2.6-4) [122]. Finally, the *adonis2*

function from the vegan package [122] was used to run PERMANOVAs, assessing the

sensitivity of model outputs to the order of its parameters.

In brief, we explored the effect of moss genus, moss species, gametophyte section,

and soil abiotic factors on bacterial community composition. We ran models with and without

Dicranum samples, thus testing the differences in bacterial community composition between

moss genera and among *Polytrichum* spp. (*P. juniperinum*, *P. commune*, and *P. piliferum*),

respectively. This approach allowed us to disentangle whether closely related moss species

are likely to harbor more similar microbial taxa than more distantly related ones, because of

shared physiological traits and similar ecological niches. In the models with *Dicranum*

samples, we modelled multivariate bacterial community composition as a function of

categorical variables '*moss genus*', '*moss species*' nested within '*moss genus*', '*gametophyte*

section', and continuous variables '*soil pH*' and '*soil temperature*', and their two-way

interactions. We nested '*moss species'* within the genera to account for the hierarchical

taxonomic structure and to explore species-specific effects on bacterial community

composition and alpha diversity (see below). In the models without *Dicranum* samples, we

modelled multivariate bacterial community composition as a function of categorical variables

'*moss species*', '*gametophyte section*', and continuous variables '*soil pH*' and '*soil*

temperature', and their two-way interactions. A fixed effect '*moss species*' was included in

the models, to quantify the portion of variance in bacterial community composition explained

by *Polytrichum* spp. In all multivariate models, permutations were restricted within blocks

(i.e., '*Location'*), using the function *how* of the permute package [123].

For alpha diversity analyses, reads were rarefied to 24,900 (slightly less than the

lowest sample sum of reads) using *rarefy_even_depth* (phyloseq) to account for uneven

sequencing depth [40, 124]. Linear models were run on Shannon index using function *lmer* in

the lme4 1.1-35.1 [125] package, then the *anova* function in the car 3.1-2 package was used

to assess model significance. Final models were further simplified by removing insignificant

interactions. For each of the alpha diversity models, the functions *simulateResiduals* and *plot*

from the DHARMa 0.4.6 package were used to assess model fit [126].

To explore the differences in bacterial diversity among moss identities, gametophyte

section and abiotic factors, bacterial diversity (Shannon index) was modelled as a function of

categorical variables 'moss *genus*', '*moss species*' (nested within '*moss genus*'),

'*gametophyte section*', and continuous variables '*soil pH*' and '*soil temperature*', and their

two-way interactions. Finally, models excluding *Dicranum* samples were run to confirm

effect persistence intra-genera, for example by including fixed factor '*moss species*'. In all

models, to account for variation among sampling locations, the random effect '*Location'* was

included, and to account for nested design, we nested '*Microsite'* under '*Location'*. See

Supplementary Table S3 for more details on model structures, fixed and random effects

tested, and link functions used.

<u>Differential abundance analyses (DAA)</u>

To identify species associated with moss species and gametophyte section, DAA were

performed using ANCOMBC 2.4.0 package [53]. This approach was chosen because it takes

into account the compositionality of next-generation sequencing data (i.e., sequencing data is

commonly expressed in terms of relative abundances, meaning they sum up to an arbitrary

constant and reside in a simplex mathematical space) [127], controls for false discovery rate

by accounting for multiple testing and multiple group comparisons, and corrects for both

taxon- and sample-specific biases (i.e., differences in extraction efficiency across taxa and

sampling fraction across samples). In brief, differentially abundant taxa were identified

between (1) gametophyte section (green vs. brown), controlling for host moss species, soil

pH, and soil temperature; and (2) host moss species, controlling for gametophyte section,

using Dunnett's test with *D. undulatum* as a reference group. In the latter, environmental

covariates were omitted because they were correlated with location (SoilTemp ~ Location,

$R^2 = 0.58$, p < 1e-10) and moss species (SoilpH ~ Host, $R^2 = 0.62$, p-value < 1e-10), and

location is significantly associated with moss species ($\chi^2 = 179.1$, p-value < 1e-10). For both

approaches, we only considered species present in at least 10% (n = 13) sample [128]. The

Holm procedure was used to account for multiple comparisons. We only reported results that

passed ANCOMBC's sensitivity test to pseudo-count addition and had an adjusted p-value <

0.01.

<u>Pathway completeness analysis</u>

Pathway completeness is a value from 0 to 100% representing the proportion of proteins

making up a given pathway (or module) whose coding sequence could be found in a genome.

The relationship between pathway completeness and gametophyte green and brown sections

was investigated for the 44 differentially abundant species. The tests were performed on

pathways from four groups of interest: nitrogen metabolism, photosynthesis, methane

metabolism, and carbon fixation (Supplementary Table S1), as described in Kyoto

Encyclopedia of Genes and Genomes (KEGG) [129]. Pathways for which completeness was

$\leq$ 20% in more than 80% of the 44 DA species considered were excluded, yielding a subset of

23 pathways. Since pathway completeness correlates with genome completeness [130], we

controlled for genome completeness (attributing 100% to non-MAGs). Specifically, the

model *Module proteins (%) ~ Gametophyte sections (green or brown) + MAG completeness*

was tested, using a generalized linear model with a quasibinomial distribution and a logit link

function to fit the response variable, which is a proportion that can include both boundaries

(0% and 100%). P-values were corrected using the Holm method and only pathways with

adjusted p-values < 0.05 were reported as significant. The heatmap in Fig. 5 was generated

using the ComplexHeatmap 2.18.0package and shows a slightly different set of pathways

compared to Supplementary Table S1, as certain pathways that were excluded for the

statistical test on the basis of completeness-prevalence (as mentioned above) were still of

interest for the discussion.

<u>Environmental DNA contamination assessment</u>

The quasi absence of non-bacterial genomes in our assemblies prompted us to investigate the

high-level taxonomy of sample reads. 100,000 reads per sample were randomly selected and

used to perform a BLASTn search against the entire *ncbi_nt* database (across all Kingdoms)

using BLAST+ 2.12.0. Although this approach cannot be used to estimate the relative

abundance of non-microbial reads, our data showed that numerous reads belonged to

eukaryotic, non-microbial species (see Supplementary Fig. S5-6). Therefore, the effective

sequencing depth of bacterial species, even after adding our nMAGs, is lower than the sample

sequencing depth. Of note, as the focus of this work being on bacteria, we designed the

metagenome assembly pipeline to focus on metagenomic bins of bacterial origin, implicitly

excluding eukaryotes that might have been assembled.

Furthermore, since no reference genome for the studied mosses was available at the

time the assembly and analyses were performed, we assessed *a posteriori* the potential of

host DNA to have contaminated our MAGs. To this end, reads from pooled sample were

mapped to every MAG separately using Bowtie2 *(--very-sensitive)*. Reads aligning with

MAGs were then mapped to the recently published genome of *P. commune*

(GCA_950295325.1). With one exception, the proportion of sample reads mapping to both,

any MAG and *P. commune*'s genome, was extremely small, at 0.02 ± 0.07%. The exceptional

MAG, *Green_N.bin.3*, had 3.7% of reads mapping to it also mapping to *P. commune*'s

genome. This MAG was annotated as a *Buchnera* species (Supplementary Data 1); a

phylogenetically close species, *Buchnera aphidicola*, has a particularly small genome (618

kbp), which could explain why the proportion of its genome conserved with a plant could be

higher than for a bacteria with a more common genome size. Nevertheless, we tested our

differential abundance analyses anew by excluding this MAG from the dataset, and the

results presented in this work remained completely unaltered, as this MAG had not been

found to be differentially abundant across gametophyte section or host mosses.

Acknowledgements

First and foremost, the authors would like to thank the Cree Community of Nemaska for

welcoming our team and allowing us to sample on their land. We thank Nemaska Lithium for

their collaboration and support of the project. We would also like to acknowledge J. Chamard

and members of S. Roy's Lab (L. Garneau, J. Beaudin, O. Alix-Paré, and S. Malo) for the

much-appreciated help with sampling and moss dissections, E. Lussier for extensive support

with laboratory work, J.-F. Lucier for assistance with metagenomic analysis, and

M. Desrochers for map generation. We are grateful to Génome Québec and to the Digital

Research Alliance of Canada for providing us with access to super computers to perform our

bioinformatic analyses. We also thank J. Beaudin, P. Moffett, and P. B. Beauregard for

critical reading of the manuscript. We are grateful to Isabel Ramirez for the beautiful

depictions of mosses presented in Fig. 1b. This work was supported by funding from

(1) Fonds de recherche du Québec – Nature et technologies (FRQNT) Programme partenariat

sur le développement durable du secteur minier (Grant 285683 lead PI Pr. Roy and co-PI Pr.

Laforest-Lapointe); and (2) Pr. Laforest-Lapointe's Canada Research Chair on Applied

Microbial Ecology.

Data availability statement

734 Scripts for all bioinformatic and statistical analyses, along with abundance tables and

735 metadata used for analysis, were made public in an online repository

736 (https://github.com/jorondo1/borealMoss). Raw sequencing reads, assemblies, and MAGs

737 were deposited in the European Nucleotide Archive (ENA) at EMBL-EBI under accession

738 number PRJEB76464.

References

1. Kayes I, Mallik A. Boreal Forests: Distributions, Biodiversity, and Management. In: Leal Filho W, Azul AM, Brandli L, Lange Salvia A, Wall T (eds). *Life on Land*. 2020. Springer International Publishing, Cham, pp 1–12.

2. Walker XJ, Baltzer JL, Cumming SG, Day NJ, Ebert C, Goetz S, et al. Increasing wildfires threaten historic carbon sink of boreal forest soils. *Nature* 2019; **572**: 520–523.

3. Turetsky MR. The role of bryophytes in carbon and nitrogen cycling. *The Bryologist* 2003; **106**: 395–409.

4. Holland-Moritz H, Stuart J, Lewis LR, Miller S, Mack MC, McDaniel SF, et al. Novel bacterial lineages associated with boreal moss species. *Environmental Microbiology* 2018; **20**: 2625–2638.

5. Ininbergs K, Bay G, Rasmussen U, Wardle DA, Nilsson M-C. Composition and diversity of nifH genes of nitrogen-fixing cyanobacteria associated with boreal forest feather mosses. *New Phytologist* 2011; **192**: 507–517.

6. Rousk K, Michelsen A. Ecosystem nitrogen fixation throughout the snow-free period in subarctic tundra: effects of willow and birch litter addition and warming. *Global Change Biology* 2017; **23**: 1552–1563.

7. Klarenberg IJ, Keuschnig C, Russi Colmenares AJ, Warshan D, Jungblut AD, Jónsdóttir IS, et al. Long-term warming effects on the microbiome and nifH gene abundance of a common moss species in sub-Arctic tundra. *New Phytologist* 2022; **234**: 2044–2056.

8. Jean M, Holland-Moritz H, Melvin AM, Johnstone JF, Mack MC. Experimental assessment of tree canopy and leaf litter controls on the microbiome and nitrogen fixation rates of two boreal mosses. *New Phytologist* 2020; **227**: 1335–1349.

9. Tveit AT, Kiss A, Winkel M, Horn F, Hájek T, Svenning MM, et al. Environmental patterns of brown moss- and Sphagnum-associated microbial communities. *Sci Rep* 2020; **10**: 22412.

10. Renaudin M, Blasi C, Bradley RL, Bellenger J-P. New insights into the drivers of moss-

associated nitrogen fixation and cyanobacterial biomass in the eastern Canadian boreal forest.

Journal of Ecology 2022; **110**: 1403–1418.

11. Masson-Delmotte V, Zhai P, Pirani A, Connors SL, Pean C, Chen Y, et al. Climate Change

2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment

Report of the Intergovernmental Panel on Climate Change. 2021. Intergovernmental Panel on

Climate Change, Switzerland.

12. Martin M, Grondin P, Lambert M-C, Bergeron Y, Morin H. Compared to Wildfire, Management

Practices Reduced Old-Growth Forest Diversity and Functionality in Primary Boreal

Landscapes of Eastern Canada. *Front For Glob Change* 2021; **4**.

13. Yin X, Martineau C, Fenton NJ. How big is the footprint? Quantifying offsite effects of mines

on boreal plant communities. *Global Ecology and Conservation* 2023; **41**: e02372.

14. Jonsson M, Kardol P, Gundale MJ, Bansal S, Nilsson M-C, Metcalfe DB, et al. Direct and

Indirect Drivers of Moss Community Structure, Function, and Associated Microfauna Across a

Successional Gradient. *Ecosystems* 2015; **18**: 154–169.

15. Royles J, Griffiths H. Invited review: climate change impacts in polar regions: lessons from

Antarctic moss bank archives. *Global Change Biology* 2015; **21**: 1041–1057.

16. Klarenberg IJ, Keuschnig C, Colmenares AJR, Warshan D, Jungblut AD, Jónsdóttir IS, et al.

Long-term warming effects on the microbiome and nifH gene abundance of a common moss

species in sub-Arctic tundra. *New Phytologist* 2021; **234**: 2044–2056.

17. Bragina A, Berg C, Cardinale M, Shcherbakov A, Chebotar V, Berg G. Sphagnum mosses

harbour highly specific bacterial diversity during their whole lifecycle. *ISME J* 2012; **6**: 802–

813.

18. Holland-Moritz H, Stuart JEM, Lewis LR, Miller SN, Mack MC, Ponciano JM, et al. The

bacterial communities of Alaskan mosses and their contributions to N2-fixation. *Microbiome*

2021; **9**: 53.

19. McLaren JR, Buckeridge KM, van de Weg MJ, Shaver GR, Schimel JP, Gough L. Shrub

encroachment in Arctic tundra: Betula nana effects on above- and belowground litter

decomposition. *Ecology* 2017; **98**: 1361–1376.

20. Cao W, Xiong Y, Zhao D, Tan H, Qu J. Bryophytes and the symbiotic microorganisms, the

pioneers of vegetation restoration in karst rocky desertification areas in southwestern China.

Appl Microbiol Biotechnol 2020; **104**: 873–891.

21. Tian Y, Li YH. Comparative analysis of bacteria associated with different mosses by 16S rRNA

and 16S rDNA sequencing. *J Basic Microbiol* 2017; **57**: 57–67.

22. Wang Y, Xue D, Chen X, Qiu Q, Chen H. Structure and Functions of Endophytic Bacterial

Communities Associated with Sphagnum Mosses and Their Drivers in Two Different Nutrient

Types of Peatlands. *Microb Ecol* 2024; **87**: 47.

23. Holland-Moritz H, Stuart JEM, Lewis LR, Miller SN, Mack MC, Ponciano JM, et al. The

bacterial communities of Alaskan mosses and their contributions to N2-fixation. *Microbiome*

2021; **9**: 1–14.

24. Elzen E van den, Bengtsson F, Fritz C, Rydin H, Lamers LPM. Variation in symbiotic N2

fixation rates among Sphagnum mosses. *PLOS ONE* 2020; **15**: e0228383.

25. Kostka JE, Weston DJ, Glass JB, Lilleskov EA, Shaw AJ, Turetsky MR. The Sphagnum

microbiome: new insights from an ancient plant lineage. *New Phytologist* 2016; **211**: 57–64.

26. Stuart JEM, Holland-Moritz H, Jean M, Miller SN, Ponciano JM, McDaniel SF, et al. The

relationship of C and N stable isotopes to high-latitude moss-associated N2 fixation. *Oecologia*

2021; **197**: 283–295.

27. Macdonald SE, Fenniak TE. Understory plant communities of boreal mixedwood forests in

western Canada: Natural patterns and response to variable-retention harvesting. *Forest Ecology*

and Management 2007; **242**: 34–48.

28. Liu C, Pouliot K, Roy S, Rochefort L. Moss regeneration for lithium mine waste rock

revegetation in Québec, Canada. *Botany* 2024; **102**: 26–40.

29. Lindo Z, Gonzalez A. The bryosphere: an integral and influential component of the Earth's

biosphere. *Ecosystems* 2010; **13**: 612–627.

30.   Arróniz-Crespo M, Bougoure J, Murphy DV, Cutler NA, Souza-Egipsy V, Chaput DL, et al.

819   Revealing the transfer pathways of cyanobacterial-fixed N into the boreal forest through the

820   feather-moss microbiome. *Front Plant Sci* 2022; **13**: 1036258.

821   31.   Chen K-H, Liao H-L, Bellenger J-P, Lutzoni F. Differential gene expression associated with

822   fungal trophic shifts along the senescence gradient of the moss Dicranum scoparium.

823   *Environmental Microbiology* 2019; **21**: 2273–2289.

824   32.   Solheim B, Zielke M. Associations Between Cyanobacteria and Mosses. In: Rai AN, Bergman

825   B, Rasmussen U (eds). *Cyanobacteria in Symbiosis*. 2002. Springer Netherlands, Dordrecht, pp

826   137–152.

827   33.   Liu X, Rousk K. The moss traits that rule cyanobacterial colonization. *Annals of Botany* 2022;

828   **129**: 147–160.

829   34.   Renaudin M, Blasi C, Bradley RL, Bellenger J-P. New insights into the drivers of moss-

830   associated nitrogen fixation and cyanobacterial biomass in the eastern Canadian boreal forest.

831   *Journal of Ecology* 2022; **110**: 1403–1418.

832   35.   Rousk K. Biotic and abiotic controls of nitrogen fixation in cyanobacteria–moss associations.

833   *New Phytologist* 2022; **235**: 1330–1335.

834   36.   Mohanasundaram B, Pandey S. Effect of environmental signals on growth and development in

835   mosses. *Journal of Experimental Botany* 2022; **73**: 4514–4527.

836   37.   Quince C, Walker AW, Simpson JT, Loman NJ, Segata N. Shotgun metagenomics, from

837   sampling to analysis. *Nat Biotechnol* 2017; **35**: 833–844.

838   38.   Thomas AM, Segata N. Multiple levels of the unknown in microbiome research. *BMC Biology*

839   2019; **17**: 48.

840   39.   Carter MM, Olm MR, Merrill BD, Dahan D, Tripathi S, Spencer SP, et al. Ultra-deep

841   sequencing of Hadza hunter-gatherers recovers vanishing gut microbes. *Cell* 2023; **186**: 3111-

842   3124.e13.

843   40.   Wicaksono WA, Cernava T, Berg C, Berg G. Bog ecosystems as a playground for plant–

844   microbe coevolution: bryophytes and vascular plants harbour functionally adapted bacteria.

845   *Microbiome* 2021; **9**: 170.

41.   Budke JM, Bernard EC, Gray DJ, Huttunen S, Piechulla B, Trigiano RN. Introduction to the

Special Issue on Bryophytes. *Critical Reviews in Plant Sciences* 2018; **37**: 102–112.

42.   Martin CE, Churchill SP. Chlorophyll concentrations and a/b ratios in mosses collected from

exposed and shaded habitats in Kansas. *Journal of Bryology* 1982.

43.   Halpern CB, Dovčiak M, Urgenson LS, Evans SA. Substrates mediate responses of forest

bryophytes to a gradient in overstory retention. *Canadian Journal of Forest Research* 2014; **44**:

855–866.

44.   Parte AC, Sardà Carbasse J, Meier-Kolthoff JP, Reimer LC, Göker M. List of Prokaryotic

names with Standing in Nomenclature (LPSN) moves to the DSMZ. *Int J Syst Evol Microbiol*

2020; **70**: 5607–5612.

45.   Ward LM, Cardona T, Holland-Moritz H. Evolutionary Implications of Anoxygenic

Phototrophy in the Bacterial Phylum Candidatus Eremiobacterota (WPS-2). *Front Microbiol*

2019; **10**.

46.   Ji M, Greening C, Vanwonterghem I, Carere CR, Bay SK, Steen JA, et al. Atmospheric trace

gases support primary production in Antarctic desert surface soil. *Nature* 2017; **552**: 400–403.

47.   Sheremet A, Jones GM, Jarett J, Bowers RM, Bedard I, Culham C, et al. Ecological and

genomic analyses of candidate phylum WPS-2 bacteria in an unvegetated soil. *Environmental*

Microbiology 2020; **22**: 3143–3157.

48.   Woodcroft BJ, Singleton CM, Boyd JA, Evans PN, Emerson JB, Zayed AAF, et al. Genome-

centric view of carbon processing in thawing permafrost. *Nature* 2018; **560**: 49–54.

49.   Ji M, Williams TJ, Montgomery K, Wong HL, Zaugg J, Berengut JF, et al. Candidatus

Eremiobacterota, a metabolically and phylogenetically diverse terrestrial phylum with acid-

tolerant adaptations. *ISME J* 2021; **15**: 2692–2707.

50.   Pessi IS, Delmont TO, Zehr JP, Hultman J. Discovery of Eremiobacterota with nifH

homologues in tundra soil. *Environ Microbiol Rep* 2024; **16**: e13277.

51.   Holland-Moritz H, Stuart J, Lewis LR, Miller S, Mack MC, McDaniel SF, et al. Novel bacterial

lineages associated with boreal moss species. *Environmental Microbiology* 2018; **20**: 2625–

2638.

52.   Ray AE, Zaugg J, Benaud N, Chelliah DS, Bay S, Wong HL, et al. Atmospheric chemosynthesis

875   is phylogenetically and geographically widespread and contributes significantly to carbon

876   fixation throughout cold deserts. *ISME J* 2022; **16**: 2547–2560.

877   53.   Lin H, Peddada SD. Multigroup analysis of compositions of microbiomes with covariate

878   adjustments and repeated measures. *Nat Methods* 2024; **21**: 83–91.

879   54.   Brenac L, Baidoo EEK, Keasling JD, Budin I. Distinct functional roles for hopanoid

880   composition in the chemical tolerance of Zymomonas mobilis. *Molecular Microbiology* 2019;

881   **112**: 1564–1575.

882   55.   Kulkarni G, Busset N, Molinaro A, Gargani D, Chaintreuil C, Silipo A, et al. Specific Hopanoid

883   Classes Differentially Affect Free-Living and Symbiotic States of Bradyrhizobium

884   diazoefficiens. *mBio* 2015; **6**: 10.1128/mbio.01251-15.

885   56.   Schmerk CL, Bernards MA, Valvano MA. Hopanoid Production Is Required for Low-pH

886   Tolerance, Antimicrobial Resistance, and Motility in Burkholderia cenocepacia. *Journal of

887   Bacteriology* 2011; **193**: 6712–6723.

888   57.   Dobritsa SV, Potter D, Gookin TE, Berry AM. Hopanoid lipids in Frankia: identification of

889   squalene-hopene cyclase gene sequences. *Can J Microbiol* 2001; **47**: 535–540.

890   58.   Cornejo-Castillo FM, Zehr JP. Hopanoid lipids may facilitate aerobic nitrogen fixation in the

891   ocean. *Proc Natl Acad Sci U S A* 2019; **116**: 18269–18271.

892   59.   Kiss L. Chapter 3 Intracellular mycoparasites in action: Interactions between powdery mildew

893   fungi and *Ampelomyces*. *British Mycological Society Symposia Series*. 2008. Elsevier, pp 37–52.

894   60.   Klavina L, Springe G, Nikolajeva V, Martsinkevich I, Nakurte I, Dzabijeva D, et al. Chemical

895   Composition Analysis, Antimicrobial Activity and Cytotoxicity Screening of Moss Extracts

896   (Moss Phytochemistry). *Molecules* 2015; **20**: 17221–17243.

897   61.   Villarreal JC, Renaudin M, Beaulieu-Laliberté A, Bellenger JP. Stigonema associated with

898   boreal Stereocaulon possesses the alternative vanadium nitrogenase. *The Lichenologist* 2021;

899   **53**: 215–220.

62.   Ortiz-Rivero J, Garrido-Benavent I, Heiðmarsson S, de los Ríos A. Moss and Liverwort Covers

Structure Soil Bacterial and Fungal Communities Differently in the Icelandic Highlands. *Microb*

Ecol 2023; **86**: 1893–1908.

63.   Taş N, Conejo N, Salmon VG. Metagenomes and Metagenome-Assembled Genome Sequences

from Nitrogen-Fixing Alder Nodules. *Microbiol Resour Announc* 2023; **12**: e01266-22.

64.   Gordon C, Wynn JM, Woodin SJ. Impacts of increased nitrogen supply on high Arctic heath:

the importance of bryophytes and phosphorus availability. *New Phytologist* 2001; **149**: 461–471.

65.   Belova SE, Suzina NE, Rijpstra WIC, Sinninghe Damsté JS, Dedysh SN. Edaphobacter

lichenicola sp. nov., a member of the family Acidobacteriaceae from lichen-dominated forested

tundra. *International Journal of Systematic and Evolutionary Microbiology* 2018; **68**: 1265–

1270.

66.   Koch IH, Gich F, Dunfield PF, Overmann J. Edaphobacter modestus gen. nov., sp. nov., and

Edaphobacter aggregans sp. nov., acidobacteria isolated from alpine and forest soils.

International Journal of Systematic and Evolutionary Microbiology 2008; **58**: 1114–1122.

67.   Shange RS, Ankumah RO, Ibekwe AM, Zabawa R, Dowd SE. Distinct Soil Bacterial

Communities Revealed under a Diversely Managed Agroecosystem. *PLOS ONE* 2012; **7**:

e40338.

68.   Chong CW, Convey P, Pearce DA, Tan IKP. Assessment of soil bacterial communities on

Alexander Island (in the maritime and continental Antarctic transitional zone). *Polar Biol* 2012;

35: 387–399.

69.   Rapoport D, Sagova-Mareckova M, Sedláček I, Provaznik J, Králová S, Pavlinic D, et al.

Trebonia kvetii gen. nov., sp. nov., an acidophilic actinobacterium, and proposal of the new

actinobacterial family Treboniaceae fam. nov. *Int J Syst Evol Microbiol* 2020; **70**: 5106–5114.

70.   Weon H-Y, Yoo S-H, Kim Y-J, Lee C-M, Kim B-Y, Jeon Y-A, et al. Rudaea cellulosilytica gen.

nov., sp. nov., isolated from soil. *International Journal of Systematic and Evolutionary*

Microbiology 2009; **59**: 2308–2312.

71.   Mieszkin S, Trouche B, Ancousture J, Raouf Y, Uroz S, Alain K. Scleromatobacter humisilvae

gen. nov., sp. nov., a novel bacterium isolated from oak forest soil. *International Journal of*

Systematic and Evolutionary Microbiology 2023; **73**: 005793.

72.   Klarenberg IJ, Keuschnig C, Salazar A, Benning LG, Vilhelmsson O. Moss and underlying soil

bacterial community structures are linked to moss functional traits. *Ecosphere* 2023; **14**: e4447.

73.   Escolástico-Ortiz DA, Blasi C, Bellenger J-P, Derome N, Villarreal-A JC. Differentially

abundant bacteria drive the N2-fixation of a widespread moss in the forest-tundra transition

zone. *Symbiosis* 2023; **90**: 193–211.

74.   Tagirdzhanova G, Saary P, Cameron E, Garber A, Díaz Escandón D, Goyette S, et al. Evidence

for a core set of microbial lichen symbionts from a global survey of metagenome. 2023.

75.   Hay MC, Mitchell AC, Soares AR, Debbonaire AR, Mogrovejo DC, Els N, et al. Metagenome-

assembled genomes from High Arctic glaciers highlight the vulnerability of glacier-associated

microbiota and their activities to habitat loss. *Microb Genom* 2023; **9**: 001131.

76.   Bragina A, Berg C, Berg G. The core microbiome bonds the Alpine bog vegetation to a

transkingdom metacommunity. *Molecular Ecology* 2015; **24**: 4795–4807.

77.   Schauer S, Kutschera U. A novel growth-promoting microbe, Methylobacterium funariae sp.

nov., isolated from the leaf surface of a common moss. *Plant Signaling & Behavior* 2011; **6**:

510–515.

78.   Gao Y-Q, Yu J-L, Ren D-M, Zhao D-P, Zhang D-J. Endophytic Methylobacterium in Tissue

Culture of the Moss Didymodon tectorum. *anbf* 2022; **59**: 117–122.

79.   Noh H-J, Park Y, Hong SG, Lee YM. Diversity and Physiological Characteristics of Antarctic

Lichens-Associated Bacteria. *Microorganisms* 2021; **9**: 607.

80.   Noh H-J, Shin SC, Park Y, Choi A, Baek K, Hong SG, et al. Lichenicola cladoniae gen. nov.,

sp. nov., a member of the family Acetobacteraceae isolated from an Antarctic lichen.

International Journal of Systematic and Evolutionary Microbiology 2020; **70**: 5918–5925.

81.   Warshan D, Bay G, Nahar N, Wardle DA, Nilsson M-C, Rasmussen U. Seasonal variation in

nifH abundance and expression of cyanobacterial communities associated with boreal feather

mosses. *ISME J* 2016; **10**: 2198–2208.

82.   Pederson ERA, Warshan D, Rasmussen U. Genome Sequencing of Pleurozium schreberi: The

955   Assembled and Annotated Draft Genome of a Pleurocarpous Feather Moss. *G3*

956   *Genes|Genomes|Genetics* 2019; **9**: 2791–2797.

957   83.   Liu L-Y, Xie G-J, Ding J, Liu B-F, Xing D-F, Ren N-Q, et al. Microbial methane emissions

958   from the non-methanogenesis processes: A critical review. *Science of The Total Environment*

959   2022; **806**: 151362.

960   84.   Bižić M, Klintzsch T, Ionescu D, Hindiyeh MY, Günthel M, Muro-Pastor AM, et al. Aquatic

961   and terrestrial cyanobacteria produce methane. *Science Advances* 2020; **6**: eaax5343.

962   85.   Renaudin M, Laforest-Lapointe I, Bellenger J-P. Unraveling global and diazotrophic

963   bacteriomes of boreal forest floor feather mosses and their environmental drivers at the

964   ecosystem and at the plant scale in North America. *Science of The Total Environment* 2022;

965   **837**: 155761.

966   86.   Gavazov KS, Soudzilovskaia NA, van Logtestijn RSP, Braster M, Cornelissen JHC. Isotopic

967   analysis of cyanobacterial nitrogen fixation associated with subarctic lichen and bryophyte

968   species. *Plant Soil* 2010; **333**: 507–517.

969   87.   Garrido-Oter R, Nakano RT, Dombrowski N, Ma K-W, AgBiome Team, McHardy AC, et al.

970   Modular Traits of the Rhizobiales Root Microbiota and Their Evolutionary Relationship with

971   Symbiotic Rhizobia. *Cell Host Microbe* 2018; **24**: 155-167.e5.

972   88.   Kolton M, Weston DJ, Mayali X, Weber PK, McFarlane KJ, Pett-Ridge J, et al. Defining the

973   Sphagnum Core Microbiome across the North American Continent Reveals a Central Role for

974   Diazotrophic Methanotrophs in the Nitrogen and Carbon Cycles of Boreal Peatland Ecosystems.

975   *mBio* 2022; **13**: e03714-21.

976   89.   Nissinen R, Franz O, Kovanen S, Mäkelä M, Kraft V, Ketola K, et al. Aerobic anoxygenic

977   phototrophic bacteria are ubiquitous in phyllo- and endosphere microbiomes of boreal and

978   subarctic plants. 2023. bioRxiv. , 2023.02.19.529139

979   90.   Zervas A, Zeng Y, Madsen AM, Hansen LH. Genomics of Aerobic Photoheterotrophs in Wheat

980   Phyllosphere Reveals Divergent Evolutionary Patterns of Photosynthetic Genes in

981   Methylobacterium spp. *Genome Biology and Evolution* 2019; **11**: 2895–2908.

982   91.   Ingram-Smith C, Martin SR, Smith KS. Acetate kinase: not just a bacterial enzyme. *Trends in*

983   *Microbiology* 2006; **14**: 249–253.

984   92.   Smith AR, Kieft B, Mueller R, Fisk MR, Mason OU, Popa R, et al. Carbon fixation and energy

985   metabolisms of a subseafloor olivine biofilm. *ISME J* 2019; **13**: 1737–1749.

986   93.   Estrada-De Los Santos P, Bustillos-Cristales R, Caballero-Mellado J. Burkholderia, a Genus

987   Rich in Plant-Associated Nitrogen Fixers with Wide Environmental and Geographic

988   Distribution. *Applied and Environmental Microbiology* 2001; **67**: 2790–2798.

989   94.   Li J, Yu C, Liu Z, Wang Y, Wang F. Microplastic accelerate the phosphorus-related metabolism

990   of bacteria to promote the decomposition of methylphosphonate to methane. *Sci Total Environ*

991   2023; **858**: 160020.

992   95.   Ney B, Ahmed FH, Carere CR, Biswas A, Warden AC, Morales SE, et al. The methanogenic

993   redox cofactor F420 is widely synthesized by aerobic soil bacteria. *ISME J* 2017; **11**: 125–137.

994   96.   Wu H-H, Pun MD, Wise CE, Streit BR, Mus F, Berim A, et al. The pathway for coenzyme M

995   biosynthesis in bacteria. *Proceedings of the National Academy of Sciences* 2022; **119**:

996   e2207190119.

997   97.   Yang Y, Cheng S, Fang H, Guo Y, Li Y, Zhou Y, et al. Linkages between the molecular

998   composition of dissolved organic matter and soil microbial community in a boreal forest during

999   freeze–thaw cycles. *Front Microbiol* 2023; **13**.

1000   98.   Ikeda MA, Nakamura H, Sawada K. Aliphatic hydrocarbons in the lichen class

1001   Lecanoromycetes and their potential use as chemotaxonomic indicators and biomarkers.

1002   *Organic Geochemistry* 2023; **179**: 104588.

1003   99.   Moussaoui L, Leduc A, Girona MM, Bélisle AC, Lafleur B, Fenton NJ, et al. Success Factors

1004   for Experimental Partial Harvesting in Unmanaged Boreal Forest: 10-Year Stand Yield Results.

1005   *Forests* 2020; **11**: 1199.

1006   100.   Environment and Climate Change Canada Historical Climate Data. Monthly Climate

1007   Summaries. 2021.

1008   101.   Collin A, Messier C, Kembel SW, Bélanger N. Can sugar maple establish into the boreal forest?

1009   Insights from seedlings under various canopies in southern Quebec. *Ecosphere* 2018; **9**: e02022.

102. Carter MR, Gregorich EG (eds). Soil Sampling and Methods of Analysis, 2nd ed. 2007. CRC

Press, Boca Raton.

103. Sims D, Sudbery I, Ilott NE, Heger A, Ponting CP. Sequencing depth and coverage: key

considerations in genomic analyses. *Nat Rev Genet* 2014; **15**: 121–132.

104. Parks DH, Rinke C, Chuvochina M, Chaumeil P-A, Woodcroft BJ, Evans PN, et al. Recovery of

nearly 8,000 metagenome-assembled genomes substantially expands the tree of life. *Nat*

Microbiol 2017; **2**: 1533–1542.

105. Uritskiy GV, DiRuggiero J, Taylor J. MetaWRAP—a flexible pipeline for genome-resolved

metagenomic data analysis. *Microbiome* 2018; **6**: 158.

106. Bowers RM, Kyrpides NC, Stepanauskas R, Harmon-Smith M, Doud D, Reddy TBK, et al.

Minimum information about a single amplified genome (MISAG) and a metagenome-assembled

genome (MIMAG) of bacteria and archaea. *Nat Biotechnol* 2017; **35**: 725–731.

107. Olm MR, Brown CT, Brooks B, Banfield JF. dRep: a tool for fast and accurate genomic

comparisons that enables improved genome recovery from metagenomes through de-replication.

ISME J 2017; **11**: 2864–2868.

108. Ondov BD, Treangen TJ, Melsted P, Mallonee AB, Bergman NH, Koren S, et al. Mash: fast

genome and metagenome distance estimation using MinHash. *Genome Biology* 2016; **17**: 132.

109. Cirkoska E, Kishore C, Bose SC, Mubarakali A, Reshmi TR, Marina N. A Study on Suffix

Trees and Their Applications in Genome Sequences Using MUMmer. In: Smys S, Palanisamy

R, Rocha Á, Beligiannis GN (eds). *Computer Networks and Inventive Communication*

Technologies. 2021. Springer Nature, Singapore, pp 15–28.

110. Orakov A, Fullam A, Coelho LP, Khedkar S, Szklarczyk D, Mende DR, et al. GUNC: detection

of chimerism and contamination in prokaryotic genomes. *Genome Biology* 2021; **22**: 178.

111. Chaumeil P-A, Mussig AJ, Hugenholtz P, Parks DH. GTDB-Tk: a toolkit to classify genomes

with the Genome Taxonomy Database. *Bioinformatics* 2020; **36**: 1925–1927.

112. Shaw J, Yu YW. Fast and robust metagenomic sequence comparison through sparse chaining

with skani. *Nat Methods* 2023; **20**: 1661–1665.

113. Brown CT, Irber L. sourmash: a library for MinHash sketching of DNA. *Journal of Open*

Source Software 2016; **1**: 27.

114. Hyatt D, Chen G-L, LoCascio PF, Land ML, Larimer FW, Hauser LJ. Prodigal: prokaryotic

gene recognition and translation initiation site identification. *BMC Bioinformatics* 2010; **11**:

119.

115. Ruiz-Perez CA, Conrad RE, Konstantinidis KT. MicrobeAnnotator: a user-friendly,

comprehensive functional annotation pipeline for microbial genomes. *BMC Bioinformatics*

2021; **22**: 11.

116. Asnicar F, Thomas AM, Beghini F, Mengoni C, Manara S, Manghi P, et al. Precise

phylogenetic analysis of microbial isolates and genomes from metagenomes using PhyloPhlAn

3.0. *Nat Commun* 2020; **11**: 2500.

117. Yu G, Smith DK, Zhu H, Guan Y, Lam TT-Y. ggtree: an r package for visualization and

annotation of phylogenetic trees with their covariates and other associated data. *Methods in*

Ecology and Evolution 2017; **8**: 28–36.

118. Sun Z, Huang S, Zhang M, Zhu Q, Haiminen N, Carrieri AP, et al. Challenges in benchmarking

metagenomic profilers. *Nat Methods* 2021; **18**: 618–626.

119. R Core Team. R: A language and environment for statistical computing. 2022.

120. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq

data with DESeq2. *Genome Biology* 2014; **15**: 550.

121. Tian W, Wang H, Xiang X, Wang R, Xu Y. Structural Variations of Bacterial Community

Driven by Sphagnum Microhabitat Differentiation in a Subalpine Peatland. *Frontiers in*

Microbiology 2019; **10**: 1–11.

122. Oksanen J, Simpson GL, Blanchet FG, Kindt R, Legendre P, Minchin PR, et al. vegan:

Community Ecology Package. 2022.

123. Simpson GL. permute: Functions for Generating Restricted Permutations of Data. R package

version 0.9-5. 2019.

124. McMurdie PJ, Holmes S. phyloseq: An R Package for Reproducible Interactive Analysis and

Graphics of Microbiome Census Data. *PLOS ONE* 2013; **8**: e61217.

125. Bates D, Mächler M, Bolker B, Walker S. Fitting linear mixed-effects models using lme4.

Journal of Statistical Software 2015; **67**: 1–48.

126. Hartig F. DHARMa: residual diagnostics for hierarchical (multi-level/mixed) regression models.

2020.

127. Gloor GB, Macklaim JM, Pawlowsky-Glahn V, Egozcue JJ. Microbiome Datasets Are

Compositional: And This Is Not Optional. *Front Microbiol* 2017; **8**.

128. Nearing JT, Douglas GM, Hayes MG, MacDonald J, Desai DK, Allward N, et al. Microbiome

differential abundance methods produce different results across 38 datasets. *Nat Commun* 2022;

13: 342.

129. Kanehisa M, Goto S. KEGG: Kyoto Encyclopedia of Genes and Genomes. *Nucleic Acids Res*

2000; **28**: 27–30.

130. Eisenhofer R, Odriozola I, Alberdi A. Impact of microbial genome completeness on

metagenomic functional inference. *ISME Communications* 2023; **3**: 12.

1078

Author contributions

SI, SR, and ILL designed the experiment. SI conducted the sample collection, processing, and

molecular work. JRL developed the bioinformatic workflow and conducted the bioinformatic

analyses with the contribution from SI, MF, and ILL. SI and JRL analyzed the data with

support from MF and ILL. SI wrote the first draft with the contribution from JRL, MF, ILL,

and SR. All authors reviewed the final manuscript.

Additional Information

Authors declare no competing interests.

 Table legends

1088 **Table 1**. Moss genus, moss species nested within moss genus, gametophyte section, soil pH,

1089 soil temperature, and their interactions drive moss bacterial community composition. Shown

1090 are the PERMANOVA on Bray-Curtis dissimilarity (n = 131 and n = 121 samples

1091 respectively) degrees of freedom (df), determination coefficients (R^2), and p-values

1092 (significant values are shown in bold).

	Variables	a) with *D. undulatum*			b) without *D. undulatum*		
		df	R^2(%)	p-value	df	R^2(%)	p-value
1st level	Moss Genus	1	13.8	**0.001**	-	-	-
	Moss Species	-	-	-	2	19.7	**0.001**
	Gametophyte Section	1	12.7	**0.001**	1	17.2	**0.001**
	Soil Temperature	1	7.4	**0.001**	1	6.9	**0.001**
	Soil pH	1	5.1	**0.001**	1	4.1	**0.001**
2nd level	Genus × Species	2	12.3	**0.001**	-	-	-
	Genus × Gametophyte Section	1	2.6	**0.001**	-	-	-
	Genus × Soil Temperature	1	0.6	0.232	-	-	-
	Genus × Soil pH	1	0.2	0.789	-	-	-
	Species × Gametophyte Section	-	-	-	2	1.3	**0.042**
	Species × Soil Temperature	-	-	-	2	3.0	**0.003**
	Species × Soil pH	-	-	-	2	3.0	**0.001**
	Gametophyte Section × Soil Temp.	1	0.9	**0.003**	1	1.1	**0.011**
	Gametophyte Section × Soil pH	1	0.3	0.447	1	0.2	0.749
	Total R^2 (%)		**57.3**			59	

1093

1094

Table 2. The effects of moss genus, moss species nested within moss genus, gametophyte section, soil pH, and temperature on moss bacterial alpha-diversity (Shannon index). Shown are the linear mixed-effects model (n = 131 and n = 121 samples respectively) test statistics (χ^2), degrees of freedom (df), and p-values (significant values are shown in bold).

	Variables	df	χ2	p-value	df	χ2	p-value
			a) with *D. undulatum*			b) without *D. undulatum.*	
	Moss Genus	1	8.6	**0.003**	-	-	-
	Moss Species	-	-	-	2	0.08	0.960
1st level	Gametophyte Section	1	29.3	<0.001	1	35.7	**<0.001**
	Soil Temperature	1	6.3	**0.012**	1	6.6	**0.010**
	Soil pH	1	1.2	0.279	1	1.7	0.192
	Genus × Species	2	0.07	0.966	-	-	-
	Genus × Gametophyte Section	1	2.1	0.148			
	Genus × Soil Temperature	1	0.04	0.840	-	-	-
2nd level	Genus × Soil pH	1	0.08	0.784	-	-	-
	Species × Gametophyte Section	-	-	-	2	2.8	0.246
	Species × Soil Temperature	-	-	-	2	2.3	0.320
	Gametophyte Section × Soil Temp.	1	1.7	0.194	1	3.0	0.084
	Gametophyte Section × Soil pH	1	12.6	**<0.001**	1	6.8	**0.009**
	Total χ² (%)		**62**			**59**	

Figure legends

Figure 1. Sampling design, location, and collected moss species.

Panel (a) shows the sampling layout: at each location, three microsites were chosen at random (indicated by yellow circles). Within each microsite, three replicate samples were collected (shown as green circles). Inset in the top right of Panel (a) shows a map of Québec with the sampling sites located within the continuous boreal forest region. Panel (b) shows the illustrations of each of four moss species (drawings were created by Isabel Ramirez). Panel (c) shows the images of green and brown sections of the moss gametophyte.

Figure 2. (a, b) Two-dimensional principal coordinates analysis (PCoA) of bacterial communities based on Bray-Curtis dissimilarity index, calculated using variance-stabilizing-

transformed abundance table. Each point represents one moss sample, and ellipses indicate a

95% bivariate confidence interval. Each ellipse colour represents a moss species (a) or a moss

gametophyte section (b). (c) Differences in alpha diversity (Shannon diversity) of moss

bacterial communities. Showed are four moss species: *P. juniperinum* (N_{brown} = 24,

N_{green} = 22), *P. commune* (N_{brown} = 9, N_{green} = 9), *P. piliferum* (N_{brown} = 26, N_{green} = 26), and

D. undulatum (N_{brown} = 8, N_{green} = 7). (d) Mean sequence relative abundance by bacterial

orders between green (top panel) and brown (bottom panel) gametophyte sections across four

moss species.

1119

Figure 3. Novel species-level metagenome-assembled genomes obtained with co-assembly.

From inner to outer layers: proportion of guanine and cytidine in assembly nucleotides

(%□GC), genome size (length, Mbp), number of contigs in each MAG, length of the shortest

of the largest contigs required to cover 50% of the total assembly length (N50) and quality

score (QS). The inner section shows the phylogenetic tree of 110 metagenome-assembled

genomes constructed using PhyloPhlAn and built phylogenetic trees using the ggtree

package, allowing a cross-validation with their assigned taxonomy in the GTDB database.

Two clades were renamed using NCBI taxonomy, as the GTDB nomenclature used

illegitimate names: Nostocales (Cyanobacteriales) and Hyphomicrobiales (Rhizobiales). The

different colored dots on the branches of the tree correspond to the colors of the bacterial

orders listed in the legend.

1131

1132 **Figure 4**. Differential abundance of bacterial species between moss species and between

1133 green and brown gametophyte sections.

1134 LFC: $\log_2$-fold change in absolute abundances, with all values showing significant

1135 differential abundance (p < 0.01). (a) Differentially abundant bacterial species in moss

species relative to *D. undulatum*. LFC are relative to *D. undulatum*. Positive LFC (blue) indicates enrichment in specified moss species relative to *D. undulatum*, whereas negative LFC (red) indicates enrichment in *D. undulatum* relative to specified moss species. (b) Bacterial species differentially abundant in either green or brown sections of moss gametophyte. Bar colours indicate in which section (green or brown) the species was differentially enriched, with LFC calculated in relation to the reference group. Blank cells correspond to species for which no significant differences in abundance were detected, and therefore no LFC values reported.

Figure 5. Metabolic abilities of the differentially abundant species in brown and green gametophyte sections.

The pathway completeness of selected metabolic pathway groups, such as photosynthesis, nitrogen and methane metabolism, and prokaryotic carbon fixation, is indicated for each genome (blue scale). This figure also shows the taxonomic rank (e.g., bacterial order) for each of 44 differentially abundant genomes.

(a)

Sampling Sites

Continuous Boreal Forest (Rowe, 1972)

Sherbrooke

Montréal

0   300 km

419 m

Collines Lescar

Eastmain Site 1

Eastmain Site 2

449 m

Pine Forest

51°40'N

Nemaska Community Intersection

Loc des Montagnes

448 m

Chemin PK

76°W

75°40'W

51°40'N

10m

0   5 km

Projection: NAD 1983 MTM 9 | Author: CEF, 2023

(b)

Polytrichum piliferum

0.3 cm

Polytrichum juniperinum

0.8 cm

Dicranum undulatum

0.7 cm

Polytrichum commune

0.8 cm

(c)