**Title:** Synthetically mannosylated antigens induce antigen-specific humoral tolerance and reduce

anti-drug antibody responses to immunogenic biologics

**Authors**: Rachel P. Wallace[1], Kirsten C. Refvik[1], Jennifer T. Antane[1], Kym Brünggel[1], Andrew

C. Tremain[2], Michal R. Raczy[1], Aaron T. Alpar[1], Mindy Nguyen[3], Ani Solanki[3], Anna J.

Slezak[1], Elyse A. Watkins[1], Abigail L. Lauterbach[1], Shijie Cao[1], D. Scott Wilson[1,4*], Jeffrey A.

Hubbell[1,2,5*]

**Affiliations:**

[1]Pritzker School for Molecular Engineering, University of Chicago; Chicago, IL 60637, USA.

[2]Committee on Immunology, University of Chicago; Chicago, IL 60637, USA.

[3]Animal Resources Center, University of Chicago; Chicago, IL 60637, USA

[4]Biomedical Engineering Department, Johns Hopkins University; Baltimore, MD 21211, USA.

[5]Committee on Cancer Biology, University of Chicago; Chicago, IL 60637, USA.

*Corresponding authors. Email: jhubbell@uchicago.edu, scott.wilson@jhmi.edu
15

**Summary:** Immunogenic biologics trigger an anti-drug antibody (ADA) response in patients,

which reduces efficacy and increases adverse reactions. Our laboratory has previously shown

that targeting protein antigen to the liver microenvironment can reduce antigen-specific T cell

responses; herein, we present a strategy to increase delivery of otherwise immunogenic biologics

to the liver via conjugation to a synthetic mannose polymer (p(Man)). This delivery leads to

reduced antigen-specific T follicular helper cell and B cell responses resulting in diminished

ADA production, which is maintained throughout subsequent administrations of the native

biologic. We found that p(Man)-antigen treatment impairs the ADA response against

recombinant uricase, a highly immunogenic biologic, without a dependence on hapten

immunodominance or control by Tregs. We identify increased TCR signaling and increased

apoptosis and exhaustion in T cells as effects of p(Man)-antigen treatment via transcriptomic

analyses. This modular platform may enhance tolerance to biologics, enabling long-term

solutions for an ever-increasing healthcare problem.

**Introduction:**

Since the approval of recombinant insulin in 1982, the development of therapeutic

proteins, or biologics, has exploded, such that in 2020 biologics accounted for 20% of FDA

approvals, a number that is expected to increase in the years to come[1]. Novel therapeutic classes

such as monoclonal antibodies and enzyme replacement therapies enabled increasingly precise

ways to treat or cure debilitating genetic diseases, cancer and autoimmunity. Compared to

traditional small-molecule therapeutics, the protein composition of biologics enable them to

directly interact with native pathways and mimic endogenous molecules to improve efficacy, but

it also makes them vulnerable to recognition by the immune system. The immune response is

primarily due to a T cell-dependent antibody response against the protein drug[2,3]. At sufficient

levels, these anti-drug antibodies (ADAs) generated in response to the therapeutic can cause a

large number of complications including neutralization of the drug's therapeutic function,

premature drug clearance, hypersensitivity and life-threatening anaphylaxis reactions upon

repeated administration[4–8].

A biologic's immunogenicity is determined by a variety of factors including

characteristics of the protein such as its homology to native human proteins expressed in that

individual, the number of subunits, and post-translational modifications, as well as genetic

factors and dose route, quantity, and frequency[3]. The pharmaceutical industry has employed a

number of strategies to try to reduce immunogenicity throughout the development of the biologic

including selecting sequences with maximum homology to human proteins, however

immunogenicity remains a hurdle for regulatory approval[9,10]. Strategies to design less

immunogenic proteins using MHC associated peptide proteomics, predictive algorithms and

machine learning[11–16] are under development, but the only treatments currently available to

patients who require immunogenic drugs is to co-administer broad immunosuppressants such as

methotrexate and thiopurines, which leave patients at increased risk for infection and

malignancy[17–21]. There is a critical need for an antigen-specific approach to reduce

immunogenicity to biologics. Here we present a modular platform that targets protein

therapeutics to the liver's innate tolerance inducing environment, and thus reduces the

immunogenicity of protein therapeutics, allowing post-tolerization administration of the native

therapeutic.

Our lab has previously developed a liver-targeted amine-reactive glycoconjugate system

and demonstrated that antigen-glycopolymer conjugates elicit tolerogenic antigen-specific T cell

signatures after both prophylactic and therapeutic administration[22,23]. The polymeric N-acetyl

galactosamine-decorated glycopolymer (p(GalNAc)) used previously were designed to target

hepatic antigen presenting cells expressing C-type lectins containing the QPD binding motif (i.e.

hepatocytes, Kupffer cells, and liver sinusoidal endothelial cells). We demonstrated that

intravenous administration of out antigen-p(GalNAc) resulted in antigen presentation to CD4$^+$

and CD8$^+$ T cells in the liver's innate tolerogenic environment, resulting in CD4$^+$ and CD8$^+$ T

cell deletion and anergy. Due to the ubiquitous presence of mannose on the surface of

endogenous and foreign antigens from yeast, bacteria, and viruses, many immune and

parenchymal cells have evolved to recognize mannosylated antigens for uptake and presentation

upon binding to EPN motif-containing lectins[24,25]. We previously developed a polymer

containing mannose residues along with TLR7 agonists as a vaccine platform and demonstrated

that mannose polymer binding to C-type lectins was critical for antigen uptake and presentation

to a wide variety of professional APCs such as DCs and macrophages[26–28]. Here, we sought to

utilize a pure mannose glycopolymer to act as a ligand for a broader range of mannose-binding

lectins and investigate its potential as a tolerogenic therapy to reduce both T cell and B cell-

mediated ADA responses to immunogenic therapeutics. We show that after intravenous injection

our p(Man) polymer construct localizes to the liver as other macromolecules do when injected

intravenously. We find this localization to be significantly more efficient than unconjugated

antigen likely due to the polymer's interactions with liver-resident APCs through their mannose-

binding C-type lectin receptors. This increased antigen localization in the liver appears to result

in reduced antigen-specific T cell, B cell and antibody responses to a variety of immunogenic

proteins. In addition, through depletion studies and transcriptional analysis, we demonstrate that

the reduced ADA response generated by our platform likely occurs due to a reduced follicular

helper T cell response and is independent of regulatory T cells or hapten immunodominance, i.e.

any immune dominance of the conjugated polymer.

**Results:**

**Conjugation to p(Man) increases antigen uptake and prolongs presentation in the liver**

We sought to utilize a novel glycopolymer as a ligand for a broader range of mannose-

binding type II C-type lectins and investigate its potential as a tolerogenic therapy (**Figure 1A,**

**Supplemental Figures 1-3**). Mannose-decorated polymers (p(Man)) were synthesized via

reversible addition fragmentation chain-transfer polymerization (RAFT) from a mannose-

decorated monomer, N-(*2-hydroxyethyl*)*methacrylamide* —a biologically inert comonomer—and

an azide-terminated RAFT agent.   The model antigen ovalbumin (OVA) was first modified with

an azide- and amine-reactive di-functional polyethylene glycol linker that upon conjugation to

the protein forms a self-immolative linkage that is relatively stable in the extracellular space, but

degrades upon internalization, releasing the protein in its unmodified form[26] (**Supplemental**

**Figure 4A**). To measure the liver-targeting effects of p(Man) conjugation compared to the

glycopolymers previously explored by our group, we performed a biodistribution study. Mice

were treated with saline, dye-labeled OVA ($OVA_{647}$), or $OVA_{647}$ conjugated to p(Man),

p(GalNAc), or a synthetic glycopolymer decorated with N-acetyl glucosamine (p(GlcNAc)) via

intravenous injection (i.v.). Organs were collected 3 hr after injection and measured via IVIS. At

the current dose, only the mice treated with p(Man)-$OVA_{647}$ experienced a significant increase in

liver fluorescence signal compared to the $OVA_{647}$ control (**Figure 1B-C**). In addition, there was

no off-target $OVA_{647}$ signal detected in other organs (**Supplemental Figure 4B**).

**Pre-treatment with p(Man)-OVA results in decreased antigen-specific Tfh cells and**

**prevents αOVA antibody response**

We next investigated the effects of p(Man)-antigen therapy on antigen-specific T cell

phenotypes using an OT-I/ OT-II vaccination model, in which adoptive transfer of OVA-specific

$CD8^+$ T cells (OT-I) or $CD4^+$ T cells (OT-II) is followed by tolerization with p(Man)-antigen and

then vaccination to challenge the tolerization regime (**Figure 1D**). When mice were given

prophylactic treatment with p(Man)-OVA, we saw an impaired expansion of OVA-specific

$CD4^+$ T cells upon vaccination compared to the untreated mice (**Figure 1E**). Specifically, we

found the greatest reduction in $CD4^+PD-1^{hi}CD44^+CXCR5^+Bcl-6^+$ T follicular helper (Tfh) cells

within the OT-II compartment after p(Man)-antigen treatment (**Figure 1F**). Given the essential

role played by Tfh cells in germinal center formation, antibody affinity maturation, and memory

B cell formation[29], the reduction in Tfh cells prompted us to investigate the effects of p(Man)-

antigen therapy on humoral immune responses. We treated mice weekly via i.v. injection with

p(Man)-OVA, followed by weekly i.v. challenges of unmodified OVA (**Figure 1G**). We found

that p(Man)-OVA antigen therapy prevented an antibody response to OVA, while treatment with

p(Man) co-administered with unconjugated OVA expedited and increased the antibody response

to OVA (**Figure 1H**).

**Pre-treatment with p(Man)-conjugated antigen prevents ADA responses to immunogenic**

**therapeutic uricase**

128           Once we established that p(Man)-OVA treatment can prevent ADA responses to OVA,

we sought to investigate whether these tolerogenic effects would prevent a strong antibody

response against a more immunogenic antigen like uricase. Uricase is an enzyme approved for

the short-term treatment of hyperuricemia, however its long-term use is often limited by

immunogenicity. Upon exposure to *Candida* uricase, mice generated a robust antigen-specific

antibody response. This antibody response was shown to be T cell dependent and was

completely abrogated when αCD4 and αCD8 blocking antibodies were administered

(**Supplemental Figure 5A,B**).

136           We produced p(Man)-uricase conjugates and verified the reaction and the self-

immolative characteristics of the linker via SDS-PAGE (**Supplemental Figure 6A**). While the

enzymatic byproduct of uricase, hydrogen peroxide, is known to have an inflammatory influence

on the immune system[30], the enzymatic activity of uricase was measured and found to be

significantly decreased upon conjugation to p(Man) and is unlikely to affect APCs upon uptake

(**Supplemental Figure 6B,C**). We administered p(Man)-uricase, unmodified uricase or saline

i.v. to mice weekly during a 3 wk tolerization window followed by weekly challenge doses of

unmodified uricase (**Figure 2A**). Serum was collected weekly, and uricase-specific antibodies

were measured via ELISA. The uricase-specific antibody response was represented as the area

under the curve of the absorbance over background across a 10-fold log-transformed dilution

series. Pre-treatment with p(Man)-uricase was found to significantly decrease the uricase-

specific IgG response after high-dose antigen challenge compared to treatment with unmodified

uricase and saline-treated mice (**Figure 2B,C**).  This reduction in antibody response at endpoint

was consistent across all subclasses of IgG (**Figure 2D**). To ensure this reduction in antibody

response was consistent across immunogenic antigens, we repeated the experiment using *E. coli*

asparaginase, an immunogenic protein used in oncology, and found consistent suppression of the

antigen-specific IgG response after p(Man)-antigen treatment (**Supplemental Figure 7**).

We next quantified the number of uricase-specific antibody-secreting cells using

ELISpot. Only the untreated group was found to have significantly more uricase-specific

antibody-secreting cells in the spleen on day 26 after initial uricase challenge, indicative of a

smaller plasmablast and plasma cell response after both antigen and p(Man)-antigen pre-

treatment (**Figure 2E**).

We next looked at the effect of p(Man)-uricase treatment on the B cell and T cell

compartments (**Supplemental Figure 8A,B**). Fluorescent uricase-streptavidin multimers were

used to identify antigen-specific B cells via flow cytometry (**Figure 3A**). Antigen-specificity was

verified by incubating splenocytes from uricase-experienced mice with 5 mM uricase or OVA

prior to multimer staining. Pre-incubation with uricase, but not OVA, resulted in the abrogation

of uricase multimer binding (**Supplemental Figure 9**). Pre-treatment with p(Man)-uricase but

not unmodified uricase prevented a significant rise in uricase-specific B cells in the spleen

compared to the naïve controls as well as a noteworthy reduction in CD38$^+$ IgD$^-$ memory B cells

and CD38$^+$IgD$^-$IgM$^-$ class-switched memory B cells compared to the saline control treatment,

which may indicate the potential for long-term B cell tolerance after p(Man)-antigen therapy

(**Figure 3B**). We also saw trends towards reduction in plasmablasts and germinal center B cells,

indicating a broadly reduced uricase-specific B cell response in the p(Man)-treated groups. The

phenotype of the corresponding bulk B cell compartment, in contrast, showed no difference

between treatment groups, indicating that p(Man)-uricase treatment is an antigen-specific

treatment without broad non-specific effects on B cells (**Figure 3C**). Similarly, the cell counts of

the bulk T cell compartment were consistent across groups, with the exception that all groups

exposed to foreign uricase antigen had increased frequency of Tfh cells within the T cell

compartment compared to the naïve control (**Supplemental Figure 10A,B**). The p(Man)-uricase

treated group was the only group without a significant increase in overall cell numbers of Tfh

cells and ICOS$^{hi}$ Tfh cells compared to the naïve control (**Supplemental Figure 10B,C**).

**Reduction in antigen-specific antibody response is not due to shift in the response to**

**αp(Man) antibodies**

After we established that pre-treatment with p(Man)-uricase leads to a reduced uricase-

specific antibody response, we wanted to probe the potential mechanisms contributing to this

effect. One possible mechanism leading to a reduced antigen-specific response is that the B cells

recognize the p(Man) polymer instead of uricase and out-compete the uricase-specific B cells for

T cell help in the germinal center, leading to a strong p(Man)-specific response instead of a

uricase-specific response. To test the p(Man)-specific antibody response, we coated ELISA

plates with uricase, p(Man)-uricase, p(Man)-OVA or p(Man) alone and measured antibodies

from mice pre-treated with p(Man)-uricase, uricase or saline as in Figure 2A (**Figure 4A**). All of

these mice were naïve to the OVA protein. We found the p(Man)-uricase treated mice generated

no antibodies that bound p(Man)-OVA or p(Man) and the average antibody response to p(Man)-

uricase was not higher than unmodified uricase in p(Man)-uricase treated mice (**Figure 4B**).

These data indicate there was no antibody response to the p(Man) polymer regardless of

treatment and thus that the possibility of immunodominance can be excluded.

**Tregs are not required to maintain humoral tolerance generated by p(Man)-antigen**

**treatment in an ADA context**

We next decided to investigate if the antigen-specific T regulatory (Treg) cells induced

by p(Man)-antigen treatment were required during high-dose challenge to maintain the reduced

ADA response. In the OT-II vaccination study, we found that treatment with p(Man)-OVA

resulted in an increased percentage of $FoxP3^+CD25^+$ Treg cells (**Supplemental Figure 11**).

These cells could be responsible for the reduced antibody response through either Tfh

suppression or direct B cell suppression. In order to probe the role of Tregs in the reduced

uricase-specific antibody response, we administered αCD25 depleting antibody to the mice after

tolerization and waited 2 wk for the antibody to be cleared before challenging with uricase

(**Figure 4C**). We found that administration of the depleting antibody was not sufficient to

prevent the reduction in antibody response seen after p(Man)-uricase treatment (**Figure 4D,E**).

Therefore, active suppression of Tfh or B cells by Tregs during subsequent high dose-antigen

challenge is likely not the mechanism by which p(Man)-antigen therapy imparts humoral

immune tolerance in the ADA context.

**p(Man)-OVA treatment results in transcriptionally distinct profiles within OT-II cells**

To further investigate the antigen-specific T cell-intrinsic mechanisms by which p(Man)-

antigen therapy could result in reduced antigen-specific antibody responses, we performed bulk

RNA sequencing on OT-II cells 5 days after antigen treatment (**Figure 5A**). We found the

p(Man)-OVA treatment group resulted in 1,994 differentially expressed genes (DEGs) compared

to the saline control, while treatment with unmodified OVA resulted in 132 DEGs compared to

saline. Of the 1,994 DEGs altered by p(Man)-treatment, only 114 were found to also be

differentially expressed in the OVA treatment comparison (**Figure 5B**). This indicates that

within the antigen-specific CD4$^+$ T cells, p(Man)-OVA treatment is driving transcriptional

changes that are distinct from those up-regulated by native antigen presentation. Examination of

the nature of the transcriptional distinctions between these two treatments provided evidence

that, even at this early stage, pro-apoptotic and immunoregulatory markers distinguish the

transcriptome after p(Man)-OVA treatment (**Figure 5C**). While both treatments exhibit the

anticipated activation and proliferative profiles of naïve T cells in the early stages after antigen

recognition, overexpression of caspases and downregulation of heat shock proteins in the p(Man-

OVA)-treated cells depicts a stronger pro-apoptotic profile, suggestive of a pro-tolerogenic

deletion of these cells from the repertoire. Simultaneously, a stronger tolerogenic signature

among the OT-II cells in the p(Man)-OVA group likely indicates the development of a persisting

population of regulatory cells. Significant upregulation of co-inhibitory receptors, anergy-

associated markers, and Treg-associated chemokine receptors like CCR10 all indicate that

p(Man)-OVA treatment imprints a unique tolerogenic phenotype on these cells[31] **(Figure 5D).**

To understand which pathways these DEGs may be affecting in the p(Man)-OVA treatment

compared to OVA treatment, we performed ingenuity pathway analysis (IPA). We identified 36

T cell-relevant immune pathways significantly altered in p(Man)-OVA treated groups compared

to unmodified OVA (**Figure 5E**). The majority of these pathways were found to be associated

with T cell activation and TCR signaling. Gene signatures were found for Th1, Th2, Tfh and

Th17 pathways, although these pathways share many activation signatures such as increases in

CD3ζ, NFAT, JAK and STAT signaling, which may explain why all are identified as significant.

Some pathways associated with T cell activation such as Nur77 and TWEAK signaling are also

associated with apoptosis and controlling T cell proliferation. The upregulation of signaling

genes after p(Man)-OVA treatment is also significantly associated with an increase in cytokine

signaling pathways such as IL-2, interferons and IL-23. Along with increased T cell signaling

and activation, IPA identified pathways associated with exhaustion and apoptosis. The T cell

exhaustion pathway is significantly upregulated due to increases in expression of co-inhibitory

receptors CTLA4, EOMES, and PD-1. Upregulation of caspases and BAD are associated with

apoptosis pathways, which are significantly upregulated in OT-II cells after p(Man)-OVA

treatment. The combination of T cell activation along with upregulated apoptosis and exhaustion

pathways suggest that p(Man)-OVA may lead to a deficient antigen-specific T follicular helper

response through the activation of T cells towards different pathways such as Th1, Th2 or Th17,

or alternatively that the increased antigen experience and activation signaling may lead to broad

apoptosis and exhaustion of OVA-specific T cells. Alongside responses leading to lack of T cell

help, we observe at this early time point of 5 days post p(Man)-antigen administration indicators

of development of regulatory immunity, including upregulation of *ikzf2*, which encodes Helios,

although *foxp3* was not upregulated at this time point. Together, these T cell fates result in

insufficient T cell help available to drive strong anti-OVA antibody responses.

**Discussion:**

In this study, we synthesized a novel mannosylated polymer, p(Man), that can be easily

conjugated to any protein antigen (such as a biologic drug) to reduce the T follicular helper and

resulting antibody responses against the unmodified antigen upon future administration. We

showed that pre-treatment with p(Man)-antigen was able to significantly reduce the antigen-

specific antibody response not only to OVA as a model protein but also to therapeutic doses of

uricase and asparaginase, both highly immunogenic protein drugs[32,33]. We first demonstrated that

p(Man) conjugation significantly increased the delivery of the conjugated antigen to the liver to a

greater extent than even our previously published liver-targeted glycopolymers[22,23]. This

difference may be due to the more promiscuous binding of mannose residues to a wider variety

of C-type lectins compared to GalNAc or GlcNAc, resulting in increased endocytosis and

antigen presentation[25,34–36], or it may be mediated by signaling through a specific mannose-

binding lectin in the liver we have yet to identify[37,38]. Our lab has previously shown that

polymeric mannose can increase antigen uptake and presentation by a variety of APCs and this

increased antigen presentation in the absence of co-stimulation may be a driving mechanism of

tolerance in this system[26–28,39]. Like the previously published glycopolymers, pre-treatment with

p(Man)-OVA resulted in an impaired CD4$^+$ antigen-specific T cell response in the OT

vaccination model, with an exceptional decrease in the number of Tfh cells within the OT-II

population.  Once we saw this lack of Tfh cell help, we confirmed that p(Man)-OVA

pretreatment was sufficient to prevent an OVA-specific antibody response. p(Man)-OVA

treatment fully prevented a detectable OVA-specific antibody response, while treatment with

unmodified OVA, which contains native mannosylation[40], did not. This provides evidence for

the unique capacity of our linear p(Man) polymer to reduce humoral immune responses to an

antigen over other types of mannosylation, at least native mannosylation in the chicken. To

investigate the capacity for p(Man)-antigen treatment to prevent antibody responses to a highly

immunogenic therapeutic protein, we conjugated it to uricase. Uricase is a tetrameric enzyme

that degrades uric acid into the more easily excreted allantoin and is used as a therapeutic to treat

hyperuricemia[41,42]. However, even when PEGylated, it elicits an immune response in the

majority of patients, limiting therapeutic efficacy[43–45]. When p(Man)-uricase was given as a pre-

treatment prior to multiple doses of unconjugated therapeutic uricase doses, we found a

significant reduction in the uricase-specific immune response via a reduction in uricase-specific

antibodies, fewer uricase-specific antibody-secreting cells, and fewer uricase-specific B cells in

the spleen, indicating a broad reduction in the humoral response to this immunogenic antigen.

The effect of p(Man)-uricase treatment was different than p(Man)-OVA or p(Man)-

asparaginase, as the antibody response to uricase was not fully eliminated. Despite the presence

of detectible antibodies after p(Man)-uricase treatment, they were significantly reduced

compared to untreated mice. This reduction may be clinically relevant, as the magnitude of the

ADA response has been shown to affect the extent of clinical consequences[46]. Furthermore

ADA-reducing therapeutics have demonstrated in both preclinical and clinical studies that a

reduction in ADAs to uricase is correlated with reduced symptoms of gout such as serum uric

acid control and reduction in tophi[47,48]. It is not yet clear how antigenic differences alter the

efficacy of p(Man)-antigen treatment. The increased immunogenicity of uricase may be the

result of its enzymatic function, which produces hydrogen peroxide, a reactive oxygen species,

as a by-product, which may increase inflammation[49,50,51]. Alternatively, there may be differences

in post-translational glycosylation of the different antigens, which has been shown to alter

immune response[52,53]. The efficacy of p(Man)-antigen treatment may also differ across antigens

due to the differences between the natural affinity of an individual's T cell or B cell compartment

to each antigen's epitopes[12,54,55].

To probe the immune mechanisms responsible for the effects of p(Man)-antigen delivery

we performed an αCD25 depletion after p(Man) treatment. We found that pretreatment with

p(Man)-uricase lead to a significant reduction in the uricase-specific antibody response despite

the depletion. αCD25 depletion has been shown to deplete Tregs in addition to other recently

activated CD25 expressing cells[56]. We interpret these data to suggest that Tregs generated during

p(Man)-antigen treatment likely do not play a predominant role actively suppressing Tfh or B

cells to cause the reduction of antibodies we see after p(Man)-antigen treatment. It is possible

that the reduction of other CD25-expressing cells such as NK cells, B cells and activated T cells

could confound this readout, but we do not expect the activated T cells or NK cells to have a role

in maintaining the antibody response once germinal center activity has been initiated, and if

critical antibody-generating B cells were depleted we would have expected antibody responses to

decrease after treatment with αCD25. We also recognized the possibility that the presence of

p(Man), a non-protein component and potential antigen, associated with the protein antigen

during initial priming of B cells could affect downstream antibody responses though hapten

immunodominance, resulting in a p(Man)-dominated antibody response. This immunodominance

has been shown with 4-hydroxy-3-nitrophenylacetyl (NP)-OVA when more copies of NP are

conjugated to an OVA protein, the B cells that are positively selected by T cells in the germinal

center predominantly bind NP while expressing the OVA epitopes on their MHC after they

endocytose the entire construct[57]. To test for this effect, we looked for antibody responses against

the p(Man) polymer, but we saw no detectable response. Therefore, hapten immunodominance is

unlikely to account for the reduction in antibody response we see to uricase after p(Man)-uricase

treatment.

To further investigate mechanism, we performed transcriptional analysis on the OT-II T

cells early (5 days) after p(Man)-OVA treatment and found upregulation of signatures associated

with TCR-signaling, proliferation, apoptosis, and alternative mechanisms of T cell activation.

These data suggest that p(Man)-antigen treatment increases presentation of the antigen to the T

cells and resultant TCR signaling; however these T cells undergo non-Tfh fates of apoptosis,

regulatory responses, exhaustion, or other non-germinal center responses limiting the Tfh help

available to the B cells, ultimately preventing strong antibody responses against the antigen.

Elevation in *nrp1/ikzf2*, genes that have canonically been used to differentiate thymic-derived

Tregs from the FoxP3$^+$ Tregs that are induced later in the periphery, show evidence of some of

the earliest regulatory response to the antigen on our brief experimental timeline, even as the

traditional CD25$^+$FoxP3$^+$ Tregs demonstrated to mediate effects in longer timeline depletion

experiments may not have developed[58]. Of additional interest might be the elevation in

chemokine receptor CCR10, which was found to be expressed by a subset of FoxP3$^+$ Tregs

homing via CCL28 secreted by cholangiocytes in the inflamed liver, where we hypothesize many

of the lectin receptors capable of binding p(Man) for antigen uptake are located[31]. Lacking in

costimulation from APCs that interacted with p(Man)-OVA, markers of anergy like *izumo1r*

(FR4), as well as co-inhibitory receptors like *cd200, lag3,* and *ctla4* were also upregulated on

OT-IIs in the pMan treated groups and likely represent key checkpoints that prevent ADA

formation. Within the pathway analysis, we also saw upregulation of pathways associated with

uptake such as endocytosis. This may be due to receptor endocytosis such as TCR and other

surface receptor down-regulation associated with tolerance[59–61] which could also account for

some reduced T cell responses we see in the OT model.

Here we investigated the capacity of p(Man)-antigen treatment to prevent T cell

dependent IgG antibody responses to subsequent antigen exposure in an antigen-specific manner.

This technology has the potential to be applied in combination as a pretreatment to a number of

immunogenic biologics in which ADA responses can limit therapeutic efficacy. Examples of

potential use cases include preventing ADAs to chronically administered monoclonal antibody

therapies such as adalimumab[62], enzyme replacement therapies such as phenylalanine ammonia

lyase for the treatment of phenylketonuria[63] and factor VIII replacement for hemophilia A[64].

With further validation, it may be used in combination with adeno-associated viral vectors to

prevent ADAs and allow re-dosing[65]. In other studies, Cao et al. found that p(Man)-antigen

pretreatment can be used to prevent food allergic responses and reduce antigen-specific IgE as

well as IgG[66]. Future work can inform how p(Man) conjugation may impact IgE responses

relevant in a minority of ADA responses[67,68] and other diseases such as asthma[69]. Since we

hypothesize p(Man)-antigen acts through the abrogation of T cell help, it may not be able to

reduce T cell independent antibody responses, and further work should be done to investigate

this application.

The experiments conducted in this study were all done in mouse models, however recent

success of conceptually similar tolerance strategies have progressed to clinical trials. A

glycosylation-modified antigen therapy has recently demonstrated safety and tolerability in a

phase 1 clinical trial for celiac disease[70] and is currently undergoing trials for multiple sclerosis

(NCT04602390) as well. It is important to consider that while mannose-binding c-type lectins

such as the mannose receptor have been identified as one-to-one orthologs between mice and

humans, others have not[24]. Subsequent studies of p(Man)-antigen's effects on antibody responses

in humanized mice and non-human primates would further inform its translational potential.

In conclusion we have demonstrated that p(Man)-antigen conjugate therapy can

significantly diminish the resulting antibody response to subsequent administration of an

immunogenic antigen. This technology has the potential to be used with any immunogenic

protein drug to prevent a patient from eliciting a strong ADA response to the drug and reduce the

risk of adverse effects associated with ADAs, including loss of therapeutic efficacy, infusion

reactions and anaphylaxis. The ability the reduce antibodies in an antigen-specific manner may

have other broad applications in the treatment of antibody-mediated diseases such as asthma,

allergy and antibody-mediated autoimmune diseases where specific antigens can be identified

such as pemphigus vulgaris and myasthenia gravis [71,72]. Further work is necessary to investigate

how p(Man)-antigen therapy would function in the therapeutic setting after an antibody response

has already developed. It could potentially be combined with antibody reducing treatments such

as anti-CD20 treatments, plasma cell depletion or IVIG to reduce an antibody response and use

p(Man)-antigen therapy to keep the antigen-specific response of interest low once broad

immunosuppressants are discontinued. Ultimately, this modular platform may enhance tolerance

to current and future biologics, enabling long-term therapeutic solutions for an ever-increasing

healthcare problem.

**Acknowledgements:** We thank the Cytometry and Antibody Core Facility at the University of

Chicago. We thank The University of Chicago HIM Facility (RRID:SCR_017916).  We thank

the Center for Research Informatics, which is funded by the Biological Sciences Division at the

University of Chicago with addition funding from the Institute for Translational Medicine,

CTSA grant number UL1 TR000430 from the NIH, and the University of Chicago

Comprehensive Cancer Center Support Grant (NIH P30CA014599). We thank Suzana Gomes

for her laboratory support throughout this project.

**Funding:** This work was supported by the University of Chicago's Chicago Immunoengineering

Innovation Center, the Alper Family Foundation, and Anokion S.A.

**Author Contributions:** Conceptualization, R.P.W, K.B., D.S.W., and J.A.H.; Methodology,

R.P.W., A.C.T., K.B., and D.S.W.; Investigation, R.P.W., A.C.T., K.B., J.T.A., A.T.A., M.N.,

A.S., E.A.W., A.J.S., A.L.L., K.C.R., and D.S.W.; Visualization, R.P.W., and K.C.R.;

Resources, M.R.R., A.J.S., and D.S.W.; Writing- Original Draft, R.P.W. and J.A.H.; Writing-

Reviewing & Editing, R.P.W., K.C.R., A.T.A., D.S.W., and J.A.H.,; Funding Acquisition,

J.A.H.; Supervision, D.S.W., and J.A.H.

**Competing Interests:** This work was funded in part by Anokion SA. D.S.W., and J.A.H., are

inventors on patents related to synthetically glycosylated antigens, licensed to Anokion SA.

J.A.H. consults for, is on the Board of Directors of, is on the Scientific Advisory Board of, and

holds equity in Anokion. All other authors declare that they have no competing interests.

**Data and materials availability:** Gene expression data obtained by RNA-seq will be deposited

in the NCBI Gene Expression Omnibus. All other data needed to evaluate the conclusions in the

paper are present in the paper or the Supplementary Materials. The datasets generated and

analyzed during the study are available from the corresponding author upon request.

**Methods:**

**Study Design**

The purposes of this study were to test the feasibility of using p(Man)-antigen therapy to induce

humoral tolerance and lessen an antigen-specific antibody response when administered before

treatment with an immunogenic protein. We used OVA as a model protein along with TCR

transgenic OT-I and OT-II to study antigen-specific T cell phenotypes and responses using flow

cytometry and RNA-seq readouts. For RNA-seq experiments, $n$=4-5 samples were analyzed per

treatment group, 3 mice were pooled per sample. For other OVA experiments, $n = 5$ mice were

used per group. We next tested p(Man)-antigen pre-treatment to reduce antibodies against

immunogenic therapeutic proteins uricase from *Candida* and asparaginase from *E. coli.* Pre-

immune mice with an initial antigen-specific antibody titer >2 prior to antigen experience were

excluded from the study.  $N = 8$ mice were used per group in immunogenic challenge

experiments. Each experiment was repeated at least twice.

**Mice**

Mice were maintained in a specific pathogen-free facility at the University of Chicago. The

experiments in this study were performed in accordance with the Institutional Animal Care and

Use Committee. For OT-I and OT-II adoptive transfer experiments, 9 week old C57BL/6 mice

were purchased from The Jackson Laboratory. OT-I (stock no: 003831) and OT-II (stock no:

004194) were crossed to CD45.1 mice (stock no: 002014) to yield congenically labeled OT-I and

OT-II. For the immunogenic protein challenge experiments 9 week old female Balb/c mice were

purchased from Charles River Laboratory.

**Transgenic T cell adoptive transfer**

Spleens and lymph nodes (axillary, brachial, inguinal, and popliteal) were isolated from TCR

transgenic mice. OT-I and OT-II T cells were isolated by negative magnetic bead selection using

a CD8 and CD4 (Stemcell) negative selection kit, respectively. Cell purity was assessed by flow

cytometry. OT-I and OT-II cells in DMEM were injected through the tail vein.

**Immunogenic antigens**

All OVA injections were performed with 15 µg Endofit OVA (Sigma). All asparaginase

injections were performed with 15 µg clinical wild-type asparaginase from *E. coli* (Asparaginase

5000, Medac GmbH). Asparaginase was purified via HIS-tag affinity purification and size

exclusion chromatography. All uricase injections were performed with 0.5U *S. candida* uricase

expressed in *E. coli* (Sigma) measured via Amplex red uricase activity assay (Thermo Fisher). *S.*

*candida* uricase was purchased from Sigma and purified via FPLC on the Äkta system (Cytiva).

Uricase was reconstituted in 50 mM sodium phosphate pH 6.4 and purified via anion exchange

chromatography on a Sepharose Q FF column (Cytiva) followed by buffer exchange into 50 mM

sodium carbonate bicarbonate pH 9.2 and size exclusion chromatography via HiLoad$^{TM}$ 16/600

Superdex$^{TM}$ 200pg (Cytiva). Activity of purified uricases and p(Man)-uricase constructs were

measured using Amplex$^{TM}$ Red Uric Acid/Uricase Assay kit (Thermo Fisher). Injections were

administered once per week intravenously in the tail vein.

**p(Man) polymer synthesis**

Unless otherwise stated, chemicals were reagent grade and purchased from Sigma-Aldrich. All

NMR spectra were collected on a Bruker Avance-II 400 MHz NMR and analyzed with

MnovaNMR (Mestrelab). GPC characterization was performed on Tosoh EcoSEC size exclusion

chromatography System using Tosoh SuperAW3000 column at 50°C, with the eluent DMF with

0.01 M LiBr. A refractive index (RI) detector was used to detect polymers, and mass values were

determined by column calibration with PMMA standards.  Mannose monomer and azide-

terminated RAFT chain transfer agent were synthesized as previously described[26,28]. Full

synthesis schema with structures are provided in supplementary information.

For RAFT polymerization, mannosylated monomer (300 mg, 1.027 mmol) and HEMA

(Combi Blocks, 300 mg, 2.326 mmol) were dissolved in dry DMF in a Schlenk tube. To that

solution,  azide CTA (17 mg, 0.029 mmol) and free radical initiator AIBN (1 mg, 0.0061 mmol)

were added. The reaction vessel was degassed via four freeze-pump-thaw cycles and then heated

to 70°C to initiate polymerization. After 14 hr, the reaction vessel was immersed in liquid

nitrogen to stop the reaction. The polymer was the precipitated in cold diethyl ether three times.

The final product was dried under reduced pressure. The product was white powder (550 mg,

yield 89%). The polymer was characterized by $^{1}$H-NMR and GPC (**Supplemental Figures 1-3**)

Resulting polymer had a number average molecular weight of 18 kDa.

**p(Man) antigen constructs**

Azido-terminated p(Man) polymer was conjugated to lysine residues on the antigen via click

chemistry using a self-immolative heterobifunctional linker, as previously described[22].

Conjugation was verified through an increase in molecular weight visible on SDS-PAGE

separation (BioRad) which was reduced upon addition of 0.05% β-mercaptoethanol in the

loading buffer. Mice were treated weekly for 2-3 weeks with 2.5-5μg p(Man)-antigen

intravenously in the tail vein.

**Antigen biodistribution**

C57BL/6 mice were treated by tail-vein injection with fluorescently labelled OVA in the form of

free OVA (OVA$_{647}$),OVA$_{647}$–p(GalNAc), OVA$_{647}$–p(GlcNAc); the subscripts indicate the

fluorophores Dy-647 from Dyomics. After 3 hr, the livers of mice treated with OVA$_{647}$, OVA$_{647}$–

p(GalNAc)or OVA$_{647}$–p(GlcNAc) were perfused via the hepatic portal vein with 5–10 ml of

42°C Krebs–Ringer modified buffer (KRB) supplemented with 0.5 mM EDTA. The perfused

livers and spleens were collected, placed in 4°C PBS, then analyzed for total fluorescence using

an IVIS Spectrum in vivo imaging system (PerkinElmer).

**Preparation of single-cell suspensions**

Draining lymph nodes (inguinal, popliteal, inguinal, axillary) and/or spleens were collected and

filtered through a 70 μm filter. Lymph nodes were mechanically disrupted and digested at 37°C

for 45 min in collagenase D. Digested lymph nodes and spleens were processed into single-cell

suspensions via mechanical disruption and passage through a sterile 70 μm strainer. Red blood

cells in splenocyte suspensions were lysed in ACK lysing buffer (Gibco) for 5 min before

quenching with 15 mL DMEM. The single cell suspensions were then resuspended in IMDM +

10% FBS + 1% pen/strep.

**Flow cytometry**

For phenotypic analysis of cells, $1.5 \times 10^6$ cells were stained in PBS with 1:200 CD16/CD32 Fc

Block (Biolegend) and 1:500 Live/Dead fixable dye (ThermoFisher) at 4$^o$C for 15 min. Cells

were washed then stained with 1:200 surface antibodies and multimers at 4$^o$C for 30 min. Cells

were washed then fixed in 2% paraformaldehyde. For intracellular staining, Cytofix/Cytoperm

(BD Biosciences) was used at 4$^\circ$C for 20 min, and cells stained intracellularly in Perm Wash

Buffer (Biolegend) with 1:200 antibodies at 4$^\circ$C for 30 min. For transcription factor stain, FoxP3

Transcription Factor Kit (eBioscience) was used per manufactures protocol. For complete

antibody panel see **Supplemental Table 1**. Representative gating found in **Supplemental**

**Figure 5**.

**Antigen-specific multimer production**

Uricase was biotinylated using EZ-link NHS-biotin (Thermo Scientific). Unreacted NHS-biotin

was removed using Zeba spin desalting columns, 7kDa MWCO (Thermo Scientific). The extent

of biotinylation was measured using the QuantTag Biotin Quantification kit (Vector

Laboratories) to ensure 1:1 molar ratio of uricase and biotin. Biotinylated uricase was reacted for

20 min on ice with 4:1 molar ratio of biotin to streptavidin-conjugated PE or streptavidin-

conjugated APC (Biolegend). Streptavidin-conjugated FITC (BioLegend) was reacted with

excess free biotin to form a non-antigen-specific streptavidin probe as a control. Multimer

formation was confirmed using SDS-PAGE gel. Cells were stained for flow cytometry with all

three streptavidin probes at the same time as other fluorescent surface markers at a volumetric

ratio of 1:200. The antigen-specific staining of B cells with multimer was verified by staining

with and without 5M uricase on splenocytes from vaccinated and antigen-naive mice.

**Antigen-binding ELISA**

For all experiments, blood was collected from mice at least 4 days after last treatment or

challenge dose and the plasma was isolated. Plasma was assessed for antigen-specific IgG by

ELISA. 96-well ELISA plates (Costar Highbind flat-bottom plates, Corning) were coated with

10 µg/mL antigen in carbonate buffer (50 mM sodium carbonate/sodium bicarbonate, pH 9.6)

overnight at 4°C. The following day, plates were washed three times in PBS with 0.05% Tween

20 (PBS-T) and then blocked with 1x casein (Sigma) for 2 h at RT. Following blocking, wells

were washed three times with PBS-T and further incubated with six 10-fold dilutions of plasma

in 1x casein for 2 hr at RT. Wells were then washed five times with PBS-T and incubated for an

additional 1 hr at RT with horseradish peroxide (HRP)- conjugated antibodies against mouse

IgG, IgG1, IgG2b, IgG2c, or IgG3 (Southern Biotech). After five washes with PBS-T, bound

antigen-specific Ig was detected with tetramethylbenzidine (TMB) substrate. Stop solution (3%

$H_2SO_4$ + 1% HCl) was added after 18 min of TMB incubation at RT, and the OD was measured

at 450 and 570 nm on an Epoch Microplate Spectrophotometer (BioTek). Background signal at

570 nm was subtracted from the OD at 450 nm. The area under the curve (AUC) of background-

subtracted absorbance versus log-transformed dilution was then calculated. Titer was determined

as the log-transformed dilution at which the background-subtracted absorbance was greater than

0.01.

**Uricase-binding IgG ELISpot assay**

Uricase was inactivated by heating at 98°C for 5 min. ELISpot plates (Millipore IP Filter plate)

were coated with 20 µg/mL heat-inactivated uricase in sterile PBS overnight at 4°C. Plates were

then blocked using ELISpot Media (RPMI 1640, 1% glutamine, 10% fetal bovine serum, 1%

penicillin-streptomycin) for 2 hr at 37°C. Splenocytes from vaccinated mice were seeded in

triplicate at a starting concentration of $6.75 \times 10^5$ cell/well and diluted seven times in 3-fold serial

dilutions. Plates were incubated for 18 hr at 37°C and 5% $CO_2$ after which the cells were washed

five times in PBS. Wells were incubated with 100 µL IgG-biotin HU adsorbed (Southern

Biotech) for 2 hr at RT. Next, plates were washed four times in PBS followed by the addition of

100 µL HRP-conjugated streptavidin/well for 1 hr at RT. Plates were washed again and

incubated with 100 µL TMB/well for 5 min. Finally, plates were washed three times with

distilled water and left to dry completely in a laminar flow hood. A CTL ImmunoSpot Analyzer

was used to image plates, count spots, and perform quality control.

**RNAseq Data Collection**

Mice were adoptively transferred with 750,000 OT-I and OT-II cells at day 0. At day 1, mice

received i.v. saline, 5 µg p(Man)-OVA, or equimolar OVA. Four days later, mice were sacrificed

and spleens and liver and peripheral lymph nodes were pooled. OT-II cells were sorted into

Trizol using fluorescence activated cell sorting (FACS) and frozen at -80$^{o}$C. RNA was extracted

(Qiagen, RNeasy Micro), and three mice were pooled per replicate. RNA integrity was measured

by the Agilent 2100 Bioanalyzer. RNA was converted to cDNA using SmartSeq-v4 (Takara, cat

no. R400752). The library was prepared using Nextera XT DNA library preparation kit

(Illumina, cat no. FC-131-1096). For each setting, all samples were pooled and sequenced using

Illumina HiSeq 4000 (2x100 paired end).

**RNAseq analysis**

The RNA-seq analysis was performed under the R programming and software

environment for statistical computing and graphics version 3.6 (R Core Team, 2019). We

assessed the quality of FastQ files using the FastQC tool (version 0.11.5). Raw reads were

aligned to the GRCh38 primary genome assembly using Spliced Transcripts Alignment to a

Reference (STAR) aligner (version 2.7.2a) 1-pass algorithm[73]. Bam files were sorted in

lexicographical order with the sambamba program (v0.5.4). Reads were assigned to exon

features annotated in Ensembl Mus musculus GRCm38 annotation (release 97) using the

FeatureCounts tool from the subread package (version 1.5.2) and the read counts were

summarized by genes[74,75]. Picard tools (version 2.18.7) was used for post alignment quality

control. Alignment-free transcriptome quantification method kallisto (v0.46.1) was used to

estimate the transcript abundance of each sample[76]. Transcript-level estimates were summarized

for gene-level analysis using R package tximport. Genes with zero read counts across all samples

were removed. Raw library sizes were scaled using normalization factors calculated with the

calcNormFactors function in the edgeR R package with trimmed mean of M-values (TMM)

option enabled[77]. The normalized count per million (CPM) values were $\log_2$-transformed. We

used the voomWithQualityWeights function from the limma package to remove heteroscedascity

from the count data. A linear model for each gene was fit with the limma algorithm, adjusted for

any batch effects, and the genes were ranked by differential expression using the empirical Bayes

method. Differentially expressed genes were identified with the Benjamini-Hochberg procedure

for multiple testing correction. Significance was set at an adjusted P-value threshold of 0.05. We

used the Ingenuity Pathway Analysis Fall 2020 release (Qiagen) to identify canonical pathways,

causal networks, and upstream regulators that related to significant differentially expressed genes

in the p(Man)-OVA to OVA comparisons.

**CD25 depletion study**

Mice were treated weekly for 3 weeks with either p(Man)-uricase or saline. Two weeks

following last treatment, mice were intravenously injected with 500 µg of either anti-CD25

depletion antibody of the IgG2a isotype control (BioXCell). Two weeks following depletion

mice were challenged weekly intravenously with 0.5U uricase for 4 wk.

**T cell depletion study**

Mice were treated weekly for 3 wk with 500 µg αCD4 and αCD8 depleting antibodies or IgG2b

isotype control antibodies (BioXCell). 4 days after initial depletion mice were administered 1U

of uricase or saline i.v. weekly for 3 wk. Serum was collected for antibody measurement.

**Statistical Analysis**

Statistically significant differences between experimental groups were determined using Prism

software (v9, GraphPad). All statistical analyses are stated specifically in the figure legends for

all experiments. For most experiments, unless otherwise specified in figure legend, one-way

ANOVA was performed with a Tukey's test to correct for multiple comparisons. For showing

statistical significance ***$P \leq 0.001$; **$P \leq 0.01$; *$P \leq 0.05$, unless otherwise stated. For the RNA-

seq experiment, a detailed description of the statistical analysis can be found in Methods.

Analyses were performed using R and GraphPad Prism V.7 software. The threshold for

statistical significance was fixed at $p < 0.05$.

**Figure 1: p(Man) conjugation increases delivery of antigen to the liver and impairs the activation of CD4$^+$ antigen-specific T follicular helper cells** (A) Structure of the p(Man) polymer conjugated to protein antigen. (B) *N*=4 C57Bl/6 mice were administered saline, free OVA-AF647 or OVA-AF647 conjugated to glycopolymers p(Man), p(GlcNAc), and p(GalNAc). 3 hr after injection organs were isolated to measure radiant efficiency of the fluorescent antigen in the livers. (C) Representative IVIS images of the liver, treated as in (B). (D) 500,000 CD45.1 congenically labeled OT-I and OT-II T cells were adoptively transferred into *n*=5 C57Bl/6 mice. Mice were then treated with saline, free OVA, or OVA conjugated to p(Man), p(GalNAc) or p(GlcNAc) before vaccination with OVA+ LPS. OT-II cells were isolated on day 19 for phenotypic analysis. (E) Frequency of OT-II in the dLN after vaccination. (F) Frequency of PD-1$^+$ Bcl6$^+$ CXCR5$^+$ T follicular helper (Tfh) cells within the OT-II population after vaccination. (G) Balb/c mice were treated weekly with p(Man)-OVA, a mixture of unconjugated p(Man) and OVA or saline for three weeks followed by a weekly challenge of high dose OVA. (H) Time course of OVA-specific IgG response represented as $\log_{10}$ titer. Symbols represent mean across *n*=8 mice. Data are shown as means ± SEM. Unless otherwise stated symbols represent individual mice and statistical differences in all graphs were determined by one-way ANOVA with Tukey's *p<0.05, **p<0.01 and ***p<0.001.

**Figure 2: Prophylactic administration of p(Man)-antigen, impairs anti-drug antibody**

**responses to immunogenic therapeutic uricase** (A) *N*=8 Balb/c mice were treated 3 times with

p(Man)-uricase, free uricase or saline followed by a therapeutic dose of uricase weekly for 4 wk.

(B) Time course of uricase-specific IgG response represented as the area under the curve of

absorbance vs log-transformed dilution (AUC). Symbols represent mean across *n*=8 mice. (C)

Uricase-specific IgG response at day 21 represented by AUC. (D) Comparison of uricase-

642    specific IgG subclasses at day 21 represented by AUC. (E) Quantification and representative

643    wells of the uricase-specific IgG-secreting splenocytes by ELISpot. Data are shown as means $\pm$

644    SEM.  Unless otherwise stated symbols represent individual mice and statistical differences in all

645    graphs were determined by one-way ANOVA with Tukey's *p<0.05, **p<0.01 and ***p<0.001

**Figure 3: Antigen-specific memory B cell responses are impaired after prophylactic**

**p(Man)-uricase treatment** (A) Representative flow plots of antigen-specific uricase-multimer

double-positive B cells of naïve mice and uricase-challenged mice on day 25 after treatment with

p(Man)-uricase, uricase, or saline. (B) Quantification of uricase-specific B220$^+$ B cells,

B220$^+$CD138$^+$ plasmablasts, B220$^+$CD38$^-$GL7$^+$ germinal center B cells, CD38$^+$IgD$^-$ memory B

cells, CD38$^+$IgM$^-$IgD$^-$ class-switched memory B cells and CD38$^+$IgD$^+$ naïve B cells in the spleen

on day 25 after treatment. (C) Quantification of bulk B220$^+$ B cells as in (B). (D) Quantification

of bulk CD3$^+$ T cells, CD4$^+$CXCR5$^+$PD-1$^+$Bcl6$^+$ Tfh cells and ICOS$^{hi}$ Tfh cells in the spleen on

659    day 25 after treatment. Data are shown as means ± SEM.  Unless otherwise stated symbols

660    represent individual mice and statistical differences in all graphs were determined by one-way

661    ANOVA with Tukey's *p<0.05, **p<0.01 and ***p<0.001.

**Figure 4: p(Man)-antigen treatment leads to an impaired antigen-specific antibody**

**response independently of hapten immunodominance or regulatory T cell control** (A)

Quantification of the antigen-specific antibody response to uricase, p(Man)-uricase, p(Man)-

OVA or free p(Man) after treatment as in 2D represented as $\log_{10}$ titer. (B) Heatmap

representation of the average $\log_{10}$ titer from each treatment group against each antigen as in (A).

(C) N=8 Balb/c mice were treated three times with saline or p(Man)-uricase followed by

administration of an isotype control or αCD25. High dose uricase injections were administered 2

wk after depletion for 4 wk. (D) Time course of the uricase-specific IgG response represented as

AUC. Symbols represent mean across *n*=8 mice. (E) Uricase-specific IgG at day 45. Data are

shown as means ± SEM. Unless otherwise stated symbols represent individual mice and

statistical differences in all graphs were determined by one-way ANOVA with Tukey's *p<0.05,

**p<0.01 and ***p<0.001.

**Figure 5: p(Man)-antigen treatment increases T cell intrinsic molecular signatures of TCR signaling along with apoptosis and exhaustion** (A) Strategy for RNA-seq. OT-I and OT-II T cells were adoptively transferred into C57Bl/6 mice on day 0. On day 1, mice were treated with p(Man)-OVA, free OVA or saline and OT-II cells were isolated from the spleens and lymph nodes for RNA-seq on day 5. *N*=4-5 samples per treatment group, 3 mice were pooled per sample. (B) Number of DEGs (FDR < 0.05; no FC cutoff) in p(Man)-OVA vs saline treated groups compared to OVA vs saline treated groups in OT-II. (C) Volcano plot of p(Man)-OVA vs OVA treatment in O-II. (D) Heatmap showing selected significant DEGs for the three comparisons (OVA vs saline; p(Man)-OVA vs saline; p(Man)-OVA vs OVA) of OT-II cells. Significant DEGs (FDR < 0.05; no FC cutoff) are indicated with an asterisk. (E) Significantly up- or down-regulated canonical immune pathways expressed in p(Man)-OVA groups compared with OVA groups as identified by ingenuity pathway analysis (IPA). Bars represent calculated z score. The BH-adjusted P value and associated genes are displayed to the right. Up/down arrows indicate gene expression.

**References**

1. de la Torre, B. G. & Albericio, F. The pharmaceutical industry in 2020. An analysis of fda drug approvals from the perspective of molecules. *Molecules* **26**, 1–14 (2021).

2. De Groot, A. S. & Scott, D. W. Immunogenicity of protein therapeutics. *Trends Immunol.* **28**, 482–490 (2007).

3. Jawa, V. *et al.* T-cell dependent immunogenicity of protein therapeutics pre-clinical assessment and mitigation–updated consensus and review 2020. *Front. Immunol.* **11**, (2020).

4. Hansel, T. T., Kropshofer, H., Singer, T., Mitchell, J. A. & George, A. J. T. The safety and side effects of monoclonal antibodies. *Nat. Rev. Drug Discov.* **9**, 325–338 (2010).

5. Mayer, L. & Young, Y. Infusion reactions and their management. *Gastroenterol. Clin. North Am.* **35**, 857–866 (2006).

6. Schellekens, H. & Casadevall, N. Immunogenicity of recombinant human proteins: Causes and consequences. *J. Neurol. Suppl.* **251**, 4–9 (2004).

7. Bartelds, G. M. *et al.* Clinical response to adalimumab: Relationship to anti-adalimumab antibodies and serum adalimumab concentrations in rheumatoid arthritis. *Ann. Rheum. Dis.* **66**, 921–926 (2007).

8. Wolbink, G. J., Aarden, L. A. & Dijkmans, B. A. C. Dealing with immunogenicity of biologicals: Assessment and clinical relevance. *Curr. Opin. Rheumatol.* **21**, 211–215 (2009).

9. Büttel, I. C. *et al.* Taking immunogenicity assessment of therapeutic proteins to the next level. *Biol. J. Int. Assoc. Biol. Stand.* **39**, 100–109 (2011).

10. FDA, CDER, & CBER. *Immunogenicity Assessment for Therapeutic Protein Products.* https://www.fda.gov/regulatory-information/search-fda-guidance-documents/immunogenicity-assessment-therapeutic-protein-products (2014).

11. Lamberth, K. *et al. Post hoc assessment of the immunogenicity of bioengineered factor VIIa*

*demonstrates the use of preclinical tools*. https://www.science.org.

12. Hamze, M. *et al.* Characterization of CD4 T cell epitopes of infliximab and rituximab

identified from healthy donors. *Front. Immunol.* **8**, (2017).

13. Karle, A., Spindeldreher, S. & Kolbinger, F. Secukinumab, a novel anti–IL-17A antibody,

shows low immunogenicity potential in human in vitro assays comparable to other marketed

biotherapeutics with low clinical immunogenicity. *mAbs* **8**, 536–550 (2016).

14. Abelin, J. G. *et al.* Defining HLA-II Ligand Processing and Binding Rules with Mass

Spectrometry Enhances Cancer Epitope Prediction. *Immunity* **51**, 766-779.e17 (2019).

15. Chen, B. *et al.* Predicting HLA class II antigen presentation through integrated deep learning.

*Nat. Biotechnol.* **37**, 1332–1343 (2019).

16. Racle, J. *et al.* Robust prediction of HLA class II epitopes by deep motif deconvolution of

immunopeptidomes. *Nat. Biotechnol.* **37**, 1283–1286 (2019).

17. Kazi, Z. B. *et al.* An immune tolerance approach using transient low-dose methotrexate in

the ERT-naïve setting of patients treated with a therapeutic protein: experience in infantile-

onset Pompe disease. *Genet. Med.* **21**, 887–895 (2019).

18. Smedby, K. E., Baecklund, E. & Askling, J. Malignant lymphomas in autoimmunity and

inflammation: A review of risks, risk factors, and lymphoma characteristics. *Cancer*

*Epidemiol. Biomarkers Prev.* **15**, 2069–2077 (2006).

19. Kandiel, A., Fraser, A. G., Korelitz, B. I., Brensinger, C. & Lewis, J. D. Increased risk of

lymphoma among inflammatory bowel disease patients treated with azathioprine and 6-

mercaptopurine. *Gut* **54**, 1121–1125 (2005).

20. Kotlyar, D. S. *et al.* Risk of lymphoma in patients with inflammatory bowel disease treated

with azathioprine and 6-mercaptopurine: A meta-analysis. *Clin. Gastroenterol. Hepatol.* **13**,

847-858.e4 (2015).

21. Engels, E. A. *et al.* Immune-related conditions and immune-modulating medications as risk

factors for non-Hodgkin's lymphoma: a case-control study. *Am. J. Epidemiol.* **162**, 1153–

1161 (2005).

22. Wilson, D. S. *et al.* Synthetically glycosylated antigens induce antigen-specific tolerance and

prevent the onset of diabetes. *Nat. Biomed. Eng.* (2019) doi:10.1038/s41551-019-0424-1.

23. Damo, M., Wilson, D. S., Watkins, E. A. & Hubbell, J. A. Soluble N-Acetylgalactosamine-

Modified Antigens Enhance Hepatocyte-Dependent Antigen Cross-Presentation and Result

in Antigen-Specific CD8+ T Cell Tolerance Development. *Front. Immunol.* **12**, 1–15 (2021).

24. Drickamer, K. & Taylor, M. E. Recent insights into structures and functions of C-type lectins

in the immune system. *Curr. Opin. Struct. Biol.* **34**, 26–34 (2015).

25. Brown, G. D., Willment, J. A. & Whitehead, L. C-type lectins in immunity and homeostasis.

*Nat. Rev. Immunol.* **18**, 374–389 (2018).

26. Wilson, D. S. *et al.* Antigens reversibly conjugated to a polymeric glyco-adjuvant induce

protective humoral and cellular immunity. *Nat. Mater.* **18**, (2019).

27. Gray, L. T. *et al.* Generation of potent cellular and humoral immunity against SARS-CoV-2

antigens via conjugation to a polymeric glyco-adjuvant. *Biomaterials* **278**, 121159 (2021).

28. Slezak, A. J. *et al.* Tumor Cell-Surface Binding of Immune Stimulating Polymeric Glyco-

Adjuvant via Cysteine-Reactive Pyridyl Disulfide Promotes Antitumor Immunity. *ACS Cent.*

*Sci.* **8**, 1435–1446 (2022).

29. Crotty, S. T Follicular Helper Cell Biology: A Decade of Discovery and Diseases. *Immunity*

**50**, 1132–1148 (2019).

30. Nathan, C. & Cunningham-Bussel, A. Beyond oxidative stress: an immunologist's guide to

reactive oxygen species. *Nat. Rev. Immunol.* **13**, 349–361 (2013).

31. Eksteen, B. *et al.* Epithelial Inflammation Is Associated with CCL28 Production and the

Recruitment of Regulatory T Cells Expressing CCR101. *J. Immunol.* **177**, 593–603 (2006).

32. Pokrovsky, V. S., Kazanov, M. D., Dyakov, I. N., Pokrovskaya, M. V. & Aleksandrova, S. S.

Comparative immunogenicity and structural analysis of epitopes of different bacterial L-

asparaginases. *BMC Cancer* **16**, 89 (2016).

33. Allen, K. C. *et al.* Risk of Anaphylaxis with Repeated Courses of Rasburicase: A Research

on Adverse Drug Events and Reports (RADAR) Project. *Drug Saf.* (2015)

doi:10.1007/s40264-014-0255-7.

34. Mi, Y. *et al.* Functional Consequences of Mannose and Asialoglycoprotein Receptor

Ablation*. *J. Biol. Chem.* **291**, 18700–18717 (2016).

35. Burgdorf, S., Kautz, A., Böhnert, V., Knolle, P. A. & Kurts, C. Distinct Pathways of Antigen

Uptake and Intracellular Routing in CD4 and CD8 T Cell Activation. *Science* **316**, 612–616

(2007).

36. Tan, M. C. A. A. *et al.* Mannose receptor-mediated uptake of antigens strongly enhances

HLA class II-restricted antigen presentation by cultured dendritic cells. *Eur. J. Immunol.* **27**,

2426–2435 (1997).

37. Schuette, V. *et al.* Mannose receptor induces T-cell tolerance via inhibition of CD45 and up-

regulation of CTLA-4. *Proc. Natl. Acad. Sci. U. S. A.* (2016) doi:10.1073/pnas.1605885113.

38. van Kooyk, Y. & Geijtenbeek, T. B. H. DC-SIGN: escape mechanism for pathogens. *Nat.*

*Rev. Immunol.* **3**, 697–709 (2003).

39. ElTanbouly, M. A. & Noelle, R. J. Rethinking peripheral T cell tolerance: checkpoints across

a T cell's journey. *Nat. Rev. Immunol.* **21**, 257–267 (2021).

40. Yang, Y., Barendregt, A., Kamerling, J. P. & Heck, A. J. R. Analyzing Protein Micro-

Heterogeneity in Chicken Ovalbumin by High-Resolution Native Mass Spectrometry

Exposes Qualitatively and Semi-Quantitatively 59 Proteoforms. *Anal. Chem.* **85**, 12037–

12045 (2013).

41. Vadhan-Raj, S. *et al.* A randomized trial of a single-dose rasburicase versus five-daily doses

in patients at risk for tumor lysis syndrome. *Ann. Oncol.* **23**, 1640–1645 (2012).

42. Sundy, J. S. *et al.* Efficacy and Tolerability of Pegloticase for the Treatment of Chronic Gout

in Patients Refractory to Conventional Treatment: Two Randomized Controlled Trials.

*JAMA* **306**, 711–720 (2011).

43. Kelly, S. J. *et al.* Diabetes Insipidus in Uricase-Deficient Mice⬜: A Model for Evaluating

Therapy with Poly ( Ethylene Glycol ) -Modified Uricase. *J. Am. Soc. Nephrol.* **12**, 1001–

1009 (2001).

44. Lipsky, P. E. *et al.* Pegloticase immunogenicity: the relationship between efficacy and

antibody development in patients treated for refractory chronic gout. *Arthritis Res. Ther.* **16**,

R60 (2014).

45. Albert, J. A., Hosey, T. & LaMoreaux, B. Increased Efficacy and Tolerability of Pegloticase

in Patients With Uncontrolled Gout Co-Treated With Methotrexate: A Retrospective Study.

*Rheumatol. Ther.* **7**, 639–648 (2020).

46. Shankar, G. *et al.* Assessment and Reporting of the Clinical Immunogenicity of Therapeutic

Proteins and Peptides—Harmonized Terminology and Tactical Recommendations. *AAPS J.*

**16**, 658–673 (2014).

47. Kishimoto, T. K. *et al.* Improving the efficacy and safety of biologic drugs with tolerogenic

nanoparticles. *Nat. Nanotechnol.* (2016) doi:10.1038/nnano.2016.135.

48. Kivitz, A. *et al.* Phase 2 Dose-Finding Study in Patients with Gout Using SEL-212, a Novel

PEGylated Uricase (SEL-037) Combined with Tolerogenic Nanoparticles (SEL-110).

*Rheumatol. Ther.* **10**, 825–847 (2023).

49. Bentley, R. & Neuberger, A. The mechanism of the action of uricase. *Biochem. J.* **52**, 694–

699 (1952).

50. Niethammer, P., Grabher, C., Look, A. T. & Mitchison, T. J. A tissue-scale gradient of

hydrogen peroxide mediates rapid wound detection in zebrafish. *Nature* **459**, 996–999

(2009).

51. El-Kenawi, A. & Ruffell, B. Inflammation, ROS, and Mutagenesis. *Cancer Cell* **32**, 727–729

(2017).

52. Fernandes, A. I. & Gregoriadis, G. The effect of polysialylation on the immunogenicity and

antigenicity of asparaginase: implication in its pharmacokinetics. *Int. J. Pharm.* **217**, 215–

224 (2001).

53. Jefferis, R. Posttranslational Modifications and the Immunogenicity of Biotherapeutics. *J.*

*Immunol. Res.* **2016**, 5358272 (2016).

54. Mazor, R. *et al.* Dual B- and T-cell de-immunization of recombinant immunotoxin targeting

mesothelin with high cytotoxic activity. *Oncotarget* **7**, 29916–29926 (2016).

55. Cassotta, A. *et al.* A single T cell epitope drives the neutralizing anti-drug antibody response

to natalizumab in multiple sclerosis patients. *Nat. Med.* **25**, 1402–1407 (2019).

56. Sakaguchi, S., Sakaguchi, N., Asano, M., Itoh, M. & Toda, M. Immunologic self-tolerance

maintained by activated T cells expressing IL-2 receptor alpha-chains (CD25). Breakdown of

a single mechanism of self-tolerance causes various autoimmune diseases. *J. Immunol.* **155**,

1151–1164 (1995).

57. Brooks, J. F. *et al.* Negative feedback by NUR77/Nr4a1 restrains B cell clonal dominance

during early T-dependent immune responses. *Cell Rep.* **36**, 109645 (2021).

58. Tan, B. J. Y., Ono, M. & Satou, Y. Single-Cell Transcriptome Analysis of Treg. in

*Regulatory T-Cells: Methods and Protocols* (ed. Ono, M.) 259–278 (Springer US, 2023).

doi:10.1007/978-1-0716-2647-4_17.

59. Boyer, C. *et al.* T cell receptor/CD3 complex internalization following activation of a

cytolytic T cell clone: Evidence for a protein kinase C-independent staurosporine-sensitive

step. *Eur. J. Immunol.* **21**, 1623–1634 (1991).

60. Balagopalan, L., Barr, V. A. & Samelson, L. E. Endocytic events in TCR signaling: focus on

adapters in microclusters. *Immunol. Rev.* **232**, 84–98 (2009).

61. Subtil, A., Hemar, A. & Dautry-Varsat, A. Rapid endocytosis of interleukin 2 receptors when

clathrin-coated pit endocytosis is inhibited. *J. Cell Sci.* **107**, 3461–3468 (1994).

62. Hoshitsuki, K. *et al.* Adalimumab Immunogenicity Is Negatively Correlated with Anti-Hinge

Antibody Levels in Patients with Rheumatoid Arthritis. *J. Pharmacol. Exp. Ther.* **375**, 488–

497 (2020).

63. Qi, Y. *et al.* Pharmacokinetic, pharmacodynamic, and immunogenic rationale for optimal

dosing of pegvaliase, a PEGylated bacterial enzyme, in adult patients with phenylketonuria.

*Clin. Transl. Sci.* **14**, 1894–1905 (2021).

64. Prezotti, A. N. L., Frade-Guanaes, J. O., Yamaguti-Hayakawa, G. G. & Ozelo, M. C.

Immunogenicity of Current and New Therapies for Hemophilia A. *Pharmaceuticals* **15**, 911

(2022).

65. Verdera, H. C., Kuranda, K. & Mingozzi, F. AAV Vector Immunogenicity in Humans: A

Long Journey to Successful Gene Transfer. *Mol. Ther.* **28**, 723–746 (2020).

66. Cao, S. *et al.* Glycopolymer-conjugated antigens as an inverse vaccine platform prevent

anaphylaxis in a pre-clinical model of food allergy. 2023.03.23.534004 Preprint at

https://doi.org/10.1101/2023.03.23.534004 (2023).

67. Vultaggio, A. *et al.* Anti-infliximab IgE and non-IgE antibodies and induction of infusion-

related severe anaphylactic reactions. *Allergy* **65**, 657–661 (2010).

68. Chung, C. H. *et al.* Cetuximab-Induced Anaphylaxis and IgE Specific for Galactose-α-1,3-

Galactose. *N. Engl. J. Med.* **358**, 1109–1117 (2008).

69. Gould, H. J. & Sutton, B. J. IgE in allergy and asthma today. *Nat. Rev. Immunol.* **8**, 205–217

(2008).

70. Murray, J. A. *et al.* Safety and tolerability of KAN-101, a liver-targeted immune tolerance

therapy, in patients with coeliac disease (ACeD): a phase 1 trial. *Lancet Gastroenterol.*

*Hepatol.* S2468125323001073 (2023) doi:10.1016/S2468-1253(23)00107-3.

71. Sajda, T. & Sinha, A. A. Autoantibody Signaling in Pemphigus Vulgaris: Development of an

Integrated Model. *Front. Immunol.* **9**, 692 (2018).

72. Gilhus, N. E. *et al.* Myasthenia gravis — autoantibody characteristics and their implications

for therapy. *Nat. Rev. Neurol.* **12**, 259–268 (2016).

73. Dobin, A. *et al.* STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, (2013).

74. Tarasov, A., Vilella, A. J., Cuppen, E., Nijman, I. J. & Prins, P. Sambamba: Fast processing

of NGS alignment formats. *Bioinformatics* **31**, (2015).

75. Liao, Y., Smyth, G. K. & Shi, W. The Subread aligner: Fast, accurate and scalable read

mapping by seed-and-vote. *Nucleic Acids Res.* **41**, (2013).

76. Bray, N. L., Pimentel, H., Melsted, P. & Pachter, L. Near-optimal probabilistic RNA-seq

quantification. *Nat. Biotechnol.* **34**, (2016).

77. Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: A Bioconductor package for

differential expression analysis of digital gene expression data. *Bioinformatics* **26**, (2009).

902