Spatial-linked alignment tool (SLAT) for aligning heterogenous slices properly

Chen-Rui Xia[1,2,†], Zhi-Jie Cao[1,2,†,*], Xin-Ming Tu[1,#], and Ge Gao[1,2,*]

[1] State Key Laboratory of Protein and Plant Gene Research, School of Life Sciences, Biomedical Pioneering Innovative Center (BIOPIC) and Beijing Advanced Innovation Center for Genomics (ICG), Center for Bioinformatics (CBI), Peking University, Beijing 100871, China

[2] Changping Laboratory, Beijing 102206, China.

[†] These authors contributed equally to this work.

[*] Authors to whom correspondence should be addressed: gaog@mail.cbi.pku.edu.cn (for G.G.), or caozj@mail.cbi.pku.edu.cn (for Z.J.C)

[#] Current address: Paul Allen School of Computer Science and Engineering, University of Washington, Seattle, WA 98195, United States

Abstract

Spatially resolved omics technologies reveal the spatial organization of cells in various biological systems. Integrative and comparative analyses of spatial omics data depend on proper slice alignment, which should take both omics profiles and spatial context into account. Here we propose SLAT (Spatially-Linked Alignment Tool), a graph-based algorithm for efficient and effective alignment of spatial omics data. Adopting a graph adversarial matching strategy, SLAT is the first algorithm capable of aligning heterogenous spatial data across distinct technologies and modalities. Systematic benchmarks demonstrate SLAT's superior precision, robustness, and speed vis à vis existing methods. Applications to multiple real-world datasets further show SLAT's utility in enhancing cell-typing resolution, integrating multiple modalities for regulatory inference, and mapping fine-scale spatial-temporal changes during development. The full SLAT package is available at https://github.com/gao-lab/SLAT.

Introduction

Recently emerging spatial omics technologies enable profiling the location, intercommunication, and functional cooperation of native cells through fluorescence *in situ* hybridization (seqFISH[1], MERFISH[2], seqFISH+[3] and Xenium[4]) and spatial barcoding (10x Visium[5], HDST[6], Slide-seqV2[7], Stereo-seq[8] and spatial-ATAC-seq[9]) from multiple tissue "slices", revealing tissue structure heterogeneity and shedding light on the underlying physiological and pathological mechanisms[8,10].

Properly aligning cells that share molecular features (e.g., cell type) and spatial context across multiple slices, especially these generated from distinct sources, is critical for their follow-up analysis. E.g., inter-technology alignment effectively bridges technologies complementary in spatial resolution and omics coverage[4], while aligning various time points during spatially dynamic processes like embryogenesis helps identify key spatial temporal changes and their molecular underpinnings. However, current spatial alignment algorithms[11–13] are mostly designed for homogeneous alignments (e.g., three-dimensional reconstruction from consecutive slices[14,15]), and can hardly handle heterogeneous slices which often involve complex non-rigid deformations, uneven spatial resolutions as well as complex batch effects.

Here, we introduce SLAT (Spatially-Linked Alignment Tool), a unified framework for aligning both homogenous and heterogeneous single-cell spatial datasets. By modeling the intercellular relationship as a spatial graph, SLAT adopts graph neural networks and adversarial matching for robustly aligning spatial slices. In addition to its superior performance revealed by systematic benchmarks, as the first algorithm capable of aligning heterogenous spatial data, SLAT introduces a wide range of application scenarios including alignment across distinct technologies and experimental conditions. SLAT is publicly accessible at https://github.com/gao-lab/SLAT and will be continuously updated as spatial omics technologies evolve.

Results

Align heterogenous spatial omics data via graph adversarial matching

By modeling the spatial topology per slice as a spatial graph where each cell is connected to its nearest neighbors by edges, we reformulate the slice-alignment task as a graph-matching problem. In efforts to correct potential cross-dataset batch effect, SLAT employed an SVD-based matrix decomposition strategy to project omics profiles of the cells into a shared low-dimensional space (**Methods**), which in turn serves as node features of the spatial graphs.

Multilayer lightweight graph convolutional networks are incorporated to propagate and aggregate information between cells and their neighbors via stepwise concatenations, generating a holistic representation with information at multiple scales from individual cells to local niches as well as global positions. Then, SLAT solves a minimum-cost bipartite matching problem between the spatial graphs through a dedicated adversarial component[16] to align cells from different slices (Fig. 1, **Methods**).

SLAT: Spatially-Linked Alignment Tool

Fig. 1 Architecture of SLAT framework.
The SLAT algorithm can be divided into three main steps: model, represent and align. We model omics data from
two single-cell spatial slices as low dimensional representations $\mathbf{X}_1 \in \mathbb{R}^{N_1 \times M}, \mathbf{X}_2 \in \mathbb{R}^{N_2 \times M}$ obtained via
SVD-based cross-dataset decomposition, where N_1, N_2 are cell numbers of each slice and M is the SVD
dimensionality. We use spatial coordinates of the cells to build K-neighbor spatial graphs $\mathcal{G}_1 = (\mathcal{V}_1, \mathcal{E}_1, \mathbf{X}_1)$, $\mathcal{G}_2 =$
$(\mathcal{V}_2, \mathcal{E}_2, \mathbf{X}_2)$ in each slice, respectively. The neighbor size can be set dynamically to adjust for uneven spatial
resolution across technologies. To encode local and global spatial information, SLAT uses L-layer lightweight
graph convolution networks to propagate and aggregate information across the spatial graph into cell embeddings
$\widehat{\mathbf{X}}_1, \widehat{\mathbf{X}}_2$, which are then projected into alignment space $\mathbf{Z}_1, \mathbf{Z}_2$ with a multi-layer perceptron trained adversarially
against a discriminator $f_D(\mathbf{Z})$ to align the two graphs by minimizing their Wasserstein distance (see **Methods**).
Node labels with the same letters (e.g., A_1, A_2) represent corresponding cells in different slices.

Notably, apart from congruent regions that can be well aligned across slices, heterogeneous

alignment often involves distinct regions that reflect biologically relevant spatial alterations. To

avoid artificially aligning such regions (i.e., over-alignment), SLAT adopts an adaptive clipping

strategy during adversarial matching to retain only the closest pairs between slices in terms of their

cosine similarity in the embedding space, taken as reliable anchors for guiding the whole alignment

procedure (Extended Data Fig. 1a). Furthermore, SLAT calculates a similarity score for each aligned

pair, making it possible to pinpoint spatially discrepant regions (see below and **Methods** for more

details).

Systematic benchmarks suggest SLAT is accurate robust and fast

To evaluate SLAT's performance over existing algorithms which are designed for homogeneous

alignment[12,14], we first ran systematic benchmarks based on consecutive, homogeneous slices from

the same tissue generated by three representative technologies with a wide range of throughput,

resolution, and technological routes: 10x Visium[5], MERFISH[2], and Stereo-seq[8] (Fig. 2a,

Supplementary Data Table 1, Supplementary Fig. 1). All slices were randomly rotated before feeding

to the alignment methods, which in turn attempt to recover the correct cell matching in expert

curated cell types and spatial regions (as visualized by vertical lines connecting the two slices in Fig.

2b and as heatmaps in Fig. 2c).

	Visium	MERFISH	Stereo-seq
Resolution	50μm	subcellular	0.2μm
Cells/Spots	~3,000	~6,500	~100,000
Features	>20,000 genes	151 genes	>20,000 genes
Tissue	Human DLPFC	Mouse hypothalamic preoptic	Whole E15.5 mouse embryo

Fig. 2 Evaluation on homogeneous spatial alignment.

a, Summary of the three benchmark datasets. For each dataset, two slices were used for alignment, see
Supplementary Fig. 1 for more details of every slice. **b**, Visualization of alignment results of different methods on
the benchmark datasets in **a**. Slices are colored by spatial region (left) and cell types (right), respectively. Vertical
lines connect aligned cell pairs (subsampled to 300 alignment pairs for clear visualization). Blue lines indicate
accurate alignments (with matching cell types and spatial regions) while red lines indicate inaccurate alignments.
PASTE failed to run on the Stereo-seq dataset due to GPU memory overflow (capping at 80 GB). **c**, Heatmaps
quantifying the region matching accuracy and cell type matching accuracy respectively in **b**, in the form of
confusion matrices. The number in each cell is the average proportion across eight repeats with different random
seeds. **d**, Running time of each method on subsampled datasets of varying sizes. $n = 8$ repeats with different
subsampling random seeds. Error bars indicate mean ± s.d.

Overall, SLAT outperformed PASTE[14] and STAGATE[12] in all three datasets (Fig. 2b and c). Of note,

PASTE exhibited particularly low cell type accuracy in the MERFISH dataset where different cell

types are spatially interlaced (Fig. 2c), probably due to its excessive reliance on the spatial distance

between cells over molecular features. Meanwhile, we found that STAGATE produced suboptimal

alignments albeit employing graph neural networks for modeling spatial omics data too, while its

performance improved for split slices which are free of batch effects (**Methods**, Supplementary Fig.

2).

By combining spatial context with transcriptome profiles, SLAT is better equipped to distinguish

transcriptionally similar but spatially distinct cell groups. Consistently, we found that SLAT also

achieved significantly higher alignment accuracy than the conventional spatially unaware algorithms

Seurat[17] and Harmony[18] (Supplementary Fig. 2-5). In particular, although these algorithms matched

cell types reasonably well, their alignment largely disarranged spatial regions (Supplementary Fig.

2-5).

SLAT's lightweight graph convolutional component effectively improves model robustness.

Evaluation with subsampled data of various sizes showed that SLAT consistently provides the best

results, even with as few as 200 cells (Supplementary Fig. 5, **Methods**). Further inspection

confirmed that the performance of SLAT remains highly robust to a wide range of hyperparameter

settings and against random corruption of the spatial graph (Extended Data Fig. 2, **Methods**).

As technologies continue to evolve, the throughput of spatial single-cell experiments is constantly

increasing[19]. Implemented as a neural network and optimized for parallelism, SLAT is highly

scalable: benchmarks showed that SLAT is consistently the fastest method across all data sizes and

aligns slices each with over 100,000 cells in just 3 minutes (Fig. 2d). We also noticed that PASTE

fails to run as cell number exceeds 25,600, partly due to its memory intensive implementation for

optimal transport.

Matching heterogeneous datasets across distinct technologies and modalities

Spatial alignment across multiple technologies and modalities amalgamates complementary

information towards a comprehensive *in situ* view of cellular states[20]. Benefiting from its novel

design, SLAT provides reliable alignment across heterogeneous datasets, which is largely beyond the

capability of current alignment algorithms.

We first used SLAT to align spatial transcriptomic datasets generated by two distinct technologies

with varying scale and detectability[8,21]. Specifically, the seqFISH dataset contains over 10,000 cells

with only 350 genes detected in total[3,21], whereas the Stereo-seq dataset contains about 5,000 cells

covering over 20,000 genes[8]. Meanwhile, the cell types in the two datasets were annotated at

different resolutions: the seqFISH dataset has finer-grained annotation (21 cell types, Extended Data

Fig. 3a) than the Stereo-seq dataset (11 cell types, Extended Data Fig. 3b). The high-quality

alignment produced by SLAT (Fig. 3a) enables an accurate label transfer for cell typing improvement

(Extended Data Fig. 3c). For example, cells labeled as "Neural crest" in the Stereo-seq dataset were

aligned to seqFISH regions with four major cell types (Fig. 3b, Extended Data Fig. 3d), refining cell

typing which was further validated by known marker genes (Fig. 3c and Extended Data Fig. 3e).

Notably, slices in the two datasets also differ in spatial structure with substantial non-rigid

deformations (Extended Data Fig. 3a and 3b). Such challenging scenario effectively failed methods

other than SLAT (Extended Data Fig. 4). Similar cross-scale alignment can also be achieved on

Visium and Xenium slices[4], where SLAT accurately pinpoints a rare group of triple positive breast

tumor cells, while all other methods failed (Extended Data Fig. 5).

Fig. 3 Spatial alignment across distinct technologies and modalities.

a, Visualization of the alignment of E8.75 seqFISH mouse embryo and E9.5 Stereo-seq mouse embryo datasets,
colored by cell types (subsampled to 300 alignment pairs for clear visualization). **b**, Highlighting the alignment of
cells labeled as "Neural crest" in Stereo-seq (top) to seqFISH (bottom), alignment lines are colored by cell type of
aligned cells in seqFISH. **c**, UMAP visualization of cells labeled as "Neural crest" in Stereo-seq, colored by cell
type of aligned cells in seqFISH, cell types with fractions less than 5% are collectively labeled as "Other". Dashed
circles indicate manual annotation by marker genes (corresponding to Extended Data Fig. 3c and 3d). **d**,
Visualization of the alignment of E11.5 spatial-ATAC-seq mouse embryo and E11.5 Stereo-seq mouse embryo
datasets, with spatial-ATAC-seq dataset colored by clusters (see Extended Data Fig. 6e) and Stereo-seq colored by
cell types, respectively (subsampled to 300 alignment pairs for clear visualization). **e**, Cell type labels of
spatial-ATAC-seq transferred from Stereo-seq via SLAT alignment. **f**, Showing chromatin accessibility score and
gene expression pattern of heart marker *Tnnt2* on SLAT alignment.

More challenging is the cross-modality spatial alignment, which is partly due to the disjoint feature

spaces that invalidate the use of our SVD-based cross dataset matrix decomposition strategy as well

as other canonical batch correction methods. Benefitting from the modular design of SLAT (Fig. 1),

we employed the graph-linked multi-modality embedding strategy we proposed before[22] to project

cells of different modalities into a shared embedding space before feeding them into the LGCNs

(**Methods**). With this extension, SLAT successfully produced a spatial alignment across RNA

(Stereo-seq) and ATAC (spatial-ATAC-seq) slices. While the different modalities featured drastically

different spatial resolutions (0.2 μm for Stereo-seq but 20 μm for spatial-ATAC-seq), SLAT managed

to align them well (Fig. 3d, Extended Data Fig. 6a). Cell-type labels transferred from Stereo-seq to

spatial-ATAC-seq based on the alignment were consistent with anatomical features and with the

accessibility of tissue-specific genes (Fig. 3e and 3f, Extended Data Fig. 6b). Joint regulatory

analysis using the aligned cell pairs and transferred labels effectively identified various key

regulators in the heart, such as *Jund* and *Ctnnb1*[23,24] (**Methods**, Extended Data Fig. 6c). We also

experimented with other spatial alignment methods by feeding the same multi-modality embeddings

as input, but found suboptimal results (Supplementary Fig. 7).

Mapping fine-scale spatial-temporal transitions by developmental alignment

Embryonic development is a highly dynamic process with extensive spatial-temporal transitions

involving the generation, maturation, and functional alteration of organs and tissues at specific

timepoints. To probe spatial-temporal dynamics during early development, we used SLAT to align

two spatial atlases of mouse embryonic development at E11.5 and E12.5[8] (Fig. 4a).

Most cells in the brain, heart, and liver were well aligned with high alignment credibility scores,

consistent with their spatial-temporal conservation at the two timepoints (Fig. 4b, Extended Data Fig.

7a and 7b). In addition, several regions were enriched with less-aligned cells of lower alignment

credibility scores, which may be attributed to both biological development (e.g., the newly emerged

organs kidney and ovary at region I[8,25–27], the rapid enlargement of lung primordium and its

displacement from the upper part of the heart to the lower part at region II[28] as shown in Extended

Data Fig. 7c, and the disappearance of branchial arch at corresponding position of region III in

E11.5[8]) and technical variation (e.g., tissue loss at corresponding position of region IV in E11.5).

Fig. 4 Developmental alignment of mouse embryo.

a, Visualization of the alignment of E11.5 and E12.5 mouse embryo Stereo-seq datasets, colored by cell types (subsampled to 300 alignment pairs for clear visualization). **b**, Similarity score of the alignment. Higher scores indicate higher alignment confidence. Dashed circles highlight five regions with low similarity scores. **c**, Sanky plot showing cell type correspondence of SLAT alignment between the two datasets. The green box highlights cells labeled as "Kidney" and "Ovary" in E12.5. **d**, Alignment visualization highlighting cells labeled as "Ovary" (left) and "Kidney" (right) in E12.5 and their aligned cells in E11.5, respectively. **e**, UMAP visualization of cell type annotations for cells labeled as "Kidney" in E12.5 and their aligned cells in E11.5. **f**, Dot plot showing marker gene expression of cell types in **e**.

We next followed up on kidney and ovary, two newly emerging organs at around E11.5–12.5[8,25–27]. SLAT accurately identified them both as developing from the "Urogenital ridge" at E11.5 (highlighted by the green box in Fig. 4c). Consistent with previous reports, SLAT alignment showed precisely that the ovary develops directly from a single area[29] (left panel of Fig. 4d) while the kidney develops from two separate areas corresponding to the mesonephros and metanephros structures in

early kidney development[25,30,31] (right panel of Fig. 4d, Extended Data Fig. 7d). In addition, based on

the clustering analysis of aligned cells, we further identified a group of rare nephron progenitor cells

located in the mesonephros, likely corresponding to an ephemeral mesonephric tubule during

mesonephros development[32] (Fig. 4e and 4f). Interestingly, the results show that the mesonephros is

spatially adjacent to the ovary, which is consistent with the well-documented developmental cascade

whereby a part of the mesonephros develops into the fallopian tube after degeneration (Fig. 4d). Our

findings revealed the unique value of heterogeneous spatial alignment for interrogating the

spatial-temporal process of organogenesis.

We also attempted the same alignment task with other methods (Extended Data Fig. 8). The spatially

unaware algorithms (Seurat, Harmony) simply missed mesonephros, while the spatially aware

algorithms (PASTE, STAGATE) align them to incorrect cells, further demonstrating the unique

capability of SLAT on heterogeneous spatial alignment.

Discussion

One of the essential challenges in spatial omics alignment is to appropriately model spatial context. Early methods such as Splotch[13] and Eggplant[33] model slices as rigid bodies and require manual annotation of landmark spots to guide the alignment. PASTE[14] eliminates the need for landmark annotation by considering gene expression of all spatial spots, but its reliance on exact spatial distance impedes application to heterogeneous alignments involving complex non-rigid structural alterations. By combining spatial graph convolution and adversarial matching, SLAT achieves reliable spatial alignment for both homogeneous and heterogeneous slices in an unsupervised, data-oriented manner.

3D reconstruction from consecutive slices is a common application of spatial alignment. SLAT supposes reconstruction from multiple slices via progressive pairwise alignment (e.g., Extended Data Fig. 9) for a consecutive stack of four slices from the same E15.5 mouse embryo). While similar 3D stacking can also be achieved with PASTE[14], SLAT is better equipped to account for non-rigid structural shifting and alteration among slices, enabling adaptive correction for potential deformation artifacts (see Supplementary Fig. 9 for quantitative assessment).

SLAT's unique ability to conduct heterogeneous spatial alignments promises a wide range of biological applications. In particular, such spatial alignment enables identifying spatially-resolved changes such as key alterations of spatial patterns during development. Meanwhile, proper cross-modal alignment further sheds lights on key regulators and corresponding regulatory circuits. However, it should be noted that the alignments produced by SLAT do not necessarily presuppose one-to-one matchings: the best match returned by SLAT should be regarded as the cell with the highest similarity in terms of cell type and spatial context, and should not be taken as definite one-to-one matches as in single-cell multi-omics techniques where different modalities were generated from the same cell simultaneously.

SLAT is fast. In fact, generating the input cell embeddings is the most time-consuming step, while

training and inferring with the SLAT core model takes only about 10 seconds for 10^6 cells. Once the

input embeddings are ready, aligning millions of cells can be completed in near real-time, enabling

an efficient search of a massive database within an affordable timeframe. SLAT's blazing speed sets

the stage for some exciting applications such as inferring spatially dependent causal mechanisms by

systematically comparing multiple Perturb-map slices[34], or constructing whole organ 3D atlases

involving thousands of slices[35].

Last but not least, designed as a flexible framework, SLAT can be readily adapted and extended. For

instance, additional information such as expert curation may be incorporated into the coordinate

matching module to help distinguish symmetric structures and polish the final alignment (**Methods**).

Meanwhile, the spatial graph modeling technique employed in SLAT may also be adapted to address

other problems, such as comparative alignment across species.

Overall, SLAT provides a unified framework for various spatial integration scenarios. To promote its

application by the research community, the SLAT package, along with detailed tutorials and demo

cases, is available online at https://github.com/gao-lab/SLAT.

Acknowledgments

We thank Mr. H. Yang for his helpful guidance on mouse embryonic development and anatomy, as well as Drs. Z. Zhang, L. Tao, F. Tang, X.S. Xie, C. Li, and J. Lu at Peking University for their helpful discussions and comments during the study. This work was supported by funds from the National Key Research and Development Program of China (2022ZD0115004), as well as the State Key Laboratory of Protein and Plant Gene Research, the Beijing Advanced Innovation Center for Genomics (ICG) at Peking University, the Changping Laboratory, and the Shaw Foundation Hong Kong Limited. The research of G.G. was supported in part by the National Program for Support of Top-Notch Young Professionals.

Author contributions

G.G. and Z.J.C conceived the study and supervised the research. C.R.X and Z.J.C. designed and implemented the computational framework and conducted benchmarks and case studies with guidance from G.G. X.M.T proposed and implemented the matrix decomposition-based batch correction strategy. C.R.X, Z.J.C., X.M.T and G.G. wrote the manuscript.

Competing interests

The authors declare no competing interests.

Extended Data Figures

Extended Data Fig. 1 Illustration of dynamic clipping and alternative metrics.

a, Illustration of the dynamic clipping criteria used to select cells for adversarial training. Only a subset of cells from the two datasets with minimal Wasserstein distance are selected (see **Methods**). **b**, Illustration where the "edge score" metric is ineffective. "Graph 1" and "Graph 2" are two graphs with five nodes, where nodes with the same letters (e.g., a_1, a_2) represent ground truth node pairs. "Alignment 1" and "Alignment 2" are two graph-matchings: "Alignment 1" correctly matches all nodes between two graphs, while only one node is correctly matched in "Alignment 2". Nevertheless, both alignments give the same "edge score".

Extended Data Fig. 2 Effect of hyperparameter settings on alignment performance.

a, Joint accuracy, which is defined as the proportion of aligned cells with both cell type and spatial region matched correctly (**Methods**) for SLAT with different hyperparameter settings including: graph neighbor size (K), input embedding methods, number of training epochs, number of LGCN layers (L), MLP hidden layer dimension, embedding hidden size (P), weight of the adversarial loss (α), dynamic clipping ratio (l), and learning rate. $n = 8$ repeats with different model random seeds. We scaled the y-axis to [0.65, 0.75] to show subtle changes. See **Methods** for detailed explanations of the hyperparameters. **b**, Changes in joint accuracy at different spatial graph corruption rates. $n = 8$ repeats with different model random seeds.

Extended Data Fig. 3 Refining cell type annotation in Stereo-seq via label transfer.
a-b, Original annotation of **a**, the seqFISH dataset and **b**, the Stereo-seq dataset. **c**, Sanky plot
showing cell type correspondence of SLAT alignment between seqFISH and Stereo-seq dataset. **d**,
Proportions seqFISH transferred cell types in Stereo-seq cells originally labeled as "Neural crest".
All other cell types with proportions of less than 5% are collectively shown as "Other". **e**, Expression
of ectoderm and mesoderm marker genes as partitioned in Fig. 5c.

Extended Data Fig. 4 Alignment results of current methods on the seqFISH and Stereo-seq slices.

Visualization of the alignment results of seqFISH and Stereo-seq mouse embryo slices via current methods. The left panels show the complete alignment results (subsampled to 300 alignment pairs for clear visualization), while the right panels highlight cells labeled as "Neural crest" in Stereo-seq and their aligned cells in seqFISH.

Extended Data Fig. 5 Alignment results of current methods on the Xenium and Visium slices.
Xenium and Visium are spatial technologies designed by 10x with complementary capabilities[4]. Xenium achieves subcellular resolution but only detects a limited set of genes, while Visium profiles the complete transcriptome with coarse spatial resolution. Specifically, the Xenium slice covers more than 100,000 cells with 313 genes detected in total, and the Visium slice contains 3,841 spots with over 20,000 genes[4]. **a, b,** Visualization of the alignment results of two consecutive Xenium and Visium human breast cancer slices via different methods (subsampled to 300 alignment pairs for clear visualization), **b** Highlighting cells labeled as "Triple positive breast tumor cells" (ERBB2+/ESR1+/PGR+, Supplementary Fig. 6a and 6b) in Xenium and their aligned cells in Visium. PASTE failed due to GPU memory overflow (capping at 80 G). Of note, SLAT is the only algorithm that successfully aligned Xenium-resolved rare triple positive cells to Visium slice, even though they span less than 10 spots (Supplementary Fig. 6c and 6d). **c,** Running time of different methods. **d,** Overlapping of spots between the SLAT aligned and manually curated triple positive spots. **e,** Log$_2$ fold change of shared top 25 marker genes in **e** for SLAT aligned compared to manually curated triple positive spots (Supplementary Fig. 6e).

Extended Data Fig. 6 Alignment-based cross-modality annotation and regulatory reference.

a, Similarity score distribution in spatial-ATAC-seq and Stereo-seq alignment. **b**, Visualization of chromatin accessibility score of cell type marker genes in the spatial-ATAC-seq slice. **c**, Key heart regulators identified by SCENIC+ based on the SLAT alignment.

Extended Data Fig. 7 Spatial-temporal dynamics in mouse embryonic development.

a, Similarity score distribution in E11.5 and E12.5 mouse embryo Stereo-seq slice alignment. **b**, Original annotation of E11.5 (top) and E12.5 (bottom) mouse embryo datasets. **c**, Spatial location of "Heart" and "Lung primordium" cells in the E11.5 (left) and E12.5 (right) slice, respectively. **d**, Spatial location of "Kidney" aligned cells in E11.5 in the right panel of **d**, colored by cell type in Fig. 4**e**.

Extended Data Fig. 8 Alignment results of current methods in mouse embryonic development.
Alignment of E11.5 and E12.5 mouse embryo slices using current methods. The left panels show the complete alignments (subsampled to 300 alignment pairs for clear visualization), while the right panels highlight cells labeled as "Kidney" in E12.5 and their aligned cells in E11.5.

Extended Data Fig. 9 Multi-slice SLAT alignment for 3D reconstruction.
Simultaneous SLAT alignment across four slices from the same E15.5 mouse embryo, ordered by vertical position
(subsampled to 300 alignment pairs for clear visualization).

References

1. Lubeck, E., Coskun, A. F., Zhiyentayev, T., Ahmad, M. & Cai, L. Single-cell in situ RNA profiling by sequential hybridization. *Nat Methods* **11**, 360–361 (2014).

2. Chen, K. H., Boettiger, A. N., Moffitt, J. R., Wang, S. & Zhuang, X. Spatially resolved, highly multiplexed RNA profiling in single cells. *Science* **348**, aaa6090 (2015).

3. Eng, C.-H. L. *et al.* Transcriptome-scale super-resolved imaging in tissues by RNA seqFISH+. *Nature* **568**, 235–239 (2019).

4. Janesick, A. *et al.* High resolution mapping of the breast cancer tumor microenvironment using integrated single cell, spatial and in situ analysis of FFPE tissue. *bioRxiv* 2022.10.06.510405 (2022) doi:10.1101/2022.10.06.510405.

5. Ståhl, P. L. *et al.* Visualization and analysis of gene expression in tissue sections by spatial transcriptomics. *Science* **353**, 78–82 (2016).

6. Vickovic, S. *et al.* High-definition spatial transcriptomics for in situ tissue profiling. *Nat Methods* **16**, 987–990 (2019).

7. Stickels, R. R. *et al.* Highly sensitive spatial transcriptomics at near-cellular resolution with Slide-seqV2. *Nat Biotechnol* **39**, 313–319 (2021).

8. Chen, A. *et al.* Spatiotemporal transcriptomic atlas of mouse organogenesis using DNA nanoball-patterned arrays. *Cell* **185**, 1777-1792.e21 (2022).

9. Deng, Y. *et al.* Spatial profiling of chromatin accessibility in mouse and human tissues. *Nature* 1–9 (2022) doi:10.1038/s41586-022-05094-1.

10. Kuppe, C. *et al.* Spatial multi-omic map of human myocardial infarction. *Nature* **608**, 766–777

(2022).

11. Long, Y. *et al.* DeepST: A versatile graph contrastive learning framework for spatially informed

clustering, integration, and deconvolution of spatial transcriptomics. 2022.08.02.502407 Preprint

at https://doi.org/10.1101/2022.08.02.502407 (2022).

12. Dong, K. & Zhang, S. Deciphering spatial domains from spatially resolved transcriptomics with

an adaptive graph attention auto-encoder. *Nat Commun* **13**, 1739 (2022).

13. Äijö, T. *et al.* Splotch: Robust estimation of aligned spatial temporal gene expression data.

757096 Preprint at https://doi.org/10.1101/757096 (2019).

14. Zeira, R., Land, M., Strzalkowski, A. & Raphael, B. J. Alignment and integration of spatial

transcriptomics data. *Nat Methods* **19**, 567–575 (2022).

15. Bergenstråhle, J., Larsson, L. & Lundeberg, J. Seamless integration of image and molecular

analysis for spatial transcriptomics workflows. *BMC Genomics* **21**, 482 (2020).

16. Gao, J., Huang, X. & Li, J. Unsupervised Graph Alignment with Wasserstein Distance

Discriminator. in *Proceedings of the 27th ACM SIGKDD Conference on Knowledge Discovery &*

Data Mining 426–435 (ACM, 2021). doi:10.1145/3447548.3467332.

17. Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-1902.e21 (2019).

18. Korsunsky, I. *et al.* Fast, sensitive and accurate integration of single-cell data with Harmony.

Nature Methods **16**, 1289–1296 (2019).

19. Moses, L. & Pachter, L. Museum of spatial transcriptomics. *Nat Methods* **19**, 534–546 (2022).

20. Eisenstein, M. Seven technologies to watch in 2022. *Nature* **601**, 658–661 (2022).

21. Lohoff, T. *et al.* Integration of spatial and single-cell transcriptomic data elucidates mouse

organogenesis. *Nat Biotechnol* **40**, 74–85 (2022).

22. Cao, Z.-J. & Gao, G. Multi-omics single-cell data integration and regulatory inference with

graph-linked embedding. *Nat Biotechnol* 1–9 (2022) doi:10.1038/s41587-022-01284-4.

23. Ricci, R. *et al.* Distinct functions of junD in cardiac hypertrophy and heart failure. *Genes Dev* **19**,

208–213 (2005).

24. Ombrato, L. *et al.* Generation of neighbor-labeling cells to study intercellular interactions in vivo.

Nat Protoc **16**, 872–892 (2021).

25. Development of Murine Urinary System. *GUDMAP*

https://www.gudmap.org/tutorials/urogenital-dev/urinary-system/.

26. Liu, C.-F., Liu, C. & Yao, H. H.-C. Building Pathways for Ovary Organogenesis in the Mouse

Embryo. *Curr Top Dev Biol* **90**, 263–290 (2010).

27. Cao, J. *et al.* The single-cell transcriptional landscape of mammalian organogenesis. *Nature* **566**,

496–502 (2019).

28. Warburton, D. *et al.* Lung Organogenesis. *Curr Top Dev Biol* **90**, 73–158 (2010).

29. Wear, H. M., McPike, M. J. & Watanabe, K. H. From primordial germ cells to primordial

follicles: a review and visual representation of early ovarian development in mice. *Journal of*

Ovarian Research **9**, 36 (2016).

30. Bouchard, M. Transcriptional control of kidney development. *Differentiation* **72**, 295–306

(2004).

31. Reidy, K. J. & Rosenblum, N. D. Cell and molecular biology of kidney development. *Semin*

Nephrol **29**, 321–337 (2009).

32. Miao, Z. *et al.* Single cell regulatory landscape of the mouse kidney highlights cellular

differentiation programs and disease targets. *Nat Commun* **12**, 2277 (2021).

33. Andersson, A. *et al.* A Landmark-based Common Coordinate Framework for Spatial

Transcriptomics Data. 2021.11.11.468178 Preprint at https://doi.org/10.1101/2021.11.11.468178

(2021).

34. Dhainaut, M. *et al.* Spatial CRISPR genomics identifies regulators of the tumor

microenvironment. *Cell* **185**, 1223-1239.e20 (2022).

35. Peng, H. *et al.* Morphological diversity of single neurons in molecularly defined cell types.

Nature **598**, 174–181 (2021).

36. Wu, F. *et al.* Simplifying Graph Convolutional Networks. Preprint at

http://arxiv.org/abs/1902.07153 (2019).

37. Wei, R., Yin, H., Jia, J., Benson, A. R. & Li, P. Understanding Non-linearity in Graph Neural

Networks from the Bayesian-Inference Perspective. Preprint at

https://doi.org/10.48550/arXiv.2207.11311 (2022).

38. Arjovsky, M., Chintala, S. & Bottou, L. Wasserstein Generative Adversarial Networks. in

Proceedings of the 34th International Conference on Machine Learning 214–223 (PMLR, 2017).

39. Marstal, K., Berendsen, F., Staring, M. & Klein, S. SimpleElastix: A User-Friendly, Multi-lingual

Library for Medical Image Registration. in *2016 IEEE Conference on Computer Vision and*

Pattern Recognition Workshops (CVPRW) 574–582 (2016). doi:10.1109/CVPRW.2016.78.

40. Besl, P. J. & McKay, N. D. A method for registration of 3-D shapes. *IEEE Transactions on*

Pattern Analysis and Machine Intelligence **14**, 239–256 (1992).

41. Maynard, K. R. *et al.* Transcriptome-scale spatial gene expression in the human dorsolateral

prefrontal cortex. *Nat Neurosci* **24**, 425–436 (2021).

42. Douglas, J. *et al.* Metrics for Evaluating Network Alignment. 6.

43. Conte, D., Foggia, P., Sansone, C. & Vento, M. THIRTY YEARS OF GRAPH MATCHING IN

PATTERN RECOGNITION. *Int J Pattern Recogn* **18**, 265–298 (2004).

44. Wolf, F. A., Angerer, P. & Theis, F. J. SCANPY: large-scale single-cell gene expression data

analysis. *Genome Biology* **19**, 15 (2018).

45. González-Blas, C. B. *et al.* SCENIC+: single-cell multiomic inference of enhancers and gene

regulatory networks. 2022.08.19.504505 Preprint at https://doi.org/10.1101/2022.08.19.504505

(2022).

Methods

SLAT framework

Modeling spatial coordinates and omics feature of single cell

We denote a spatial omics dataset as $\mathcal{D} = \{(\mathbf{g}^{(i)}, \mathbf{s}^{(i)}), i = 1,2, \ldots, N\}$, where N is the number of spots or cells, $\mathbf{g}^{(i)} \in \mathbb{R}^G$, and $\mathbf{s}^{(i)} \in \mathbb{R}^2$ are the raw omics features (e.g., genes) and spatial coordinates of cell i, respectively, where G is the number of omics features. For datasets containing non-identical omics features, we use their overlapping features. For ease of notation, we denote the combination of omics features and spatial coordinates across all cells in a dataset as matrices $\mathbf{G} \in \mathbb{R}^{N \times G}$, $\mathbf{S} \in \mathbb{R}^{N \times 2}$, respectively. Subscripts such as $\mathbf{G}_1, \mathbf{G}_2$ and $\mathbf{S}_1, \mathbf{S}_2$ are added to distinguish two datasets being aligned.

In attempts to correct inter-sample batch effects, we employ an SVD-based cross-dataset matrix decomposition strategy as a preprocessing step. To begin with, we denote the log-normalized and scaled omics matrices of two spatial datasets as $\widetilde{\mathbf{G}}_1 \in \mathbb{R}^{N_1 \times G}$ and $\widetilde{\mathbf{G}}_2 \in \mathbb{R}^{N_2 \times G}$. We then apply SVD on their dot product as follows:

$$\widetilde{\mathbf{G}}_1 \widetilde{\mathbf{G}}_2^\top = \mathbf{U}\mathbf{\Sigma}\mathbf{V}^\top \#(1)$$

Using the decomposed matrices, we obtain the batch-corrected embeddings of the two datasets as:

$$\mathbf{X}_1 = \mathbf{U}_{1:M}\mathbf{\Sigma}_{1:M}^{\frac{1}{2}} \#(2)$$
$$\mathbf{X}_2 = \mathbf{V}_{1:M}\mathbf{\Sigma}_{1:M}^{\frac{1}{2}}$$

where $\mathbf{U}_{1:M} \in \mathbb{R}^{N_1 \times M}$, $\mathbf{V}_{1:M} \in \mathbb{R}^{N_2 \times M}$, $\mathbf{\Sigma}_{1:M} \in \mathbb{R}^{M \times M}$ are truncated decompositions corresponding to the M-largest singular values. Notably, this preprocessing step is flexible and allows for modular input (e.g., embedding produced by dedicated algorithms like GLUE[22] for cross-modality integration).

We model the spatial information of cells in $\mathcal{D}$ as a spatial graph $\mathcal{G} = (\mathcal{V}, \mathcal{E}, \mathbf{X})$, where each node $v_i \in \mathcal{V}$ corresponds to a cell with the batch corrected embedding $\mathbf{x}^{(i)} \in \mathbb{R}^M$ as its node attribute, and

the edges connect the *K*-nearest neighbors in the spatial space. We also denote the adjacency matrix

of the graph as $\mathbf{A} \in \{0,1\}^{N \times N}$. $\mathbf{A}_{i,j} = 1$ if the edge $(v_i, v_j) \in \mathcal{E}$, otherwise $\mathbf{A}_{i,j} = 0$. In particular,

for cross-technology alignment, different technologies may have distinct spatial resolutions, in which

case we can select different *K*'s for each slice according to its spatial resolution. SLAT also supports

building the spatial graph by radius, where all cells located within a specific radius are taken as

"neighbors".

SLAT formulates the alignment of two spatial datasets $\mathcal{D}_1 = \left\{ \left(\mathbf{x}_1^{(i)}, \mathbf{s}_1^{(i)} \right), i = 1,2,\ldots,N_1 \right\}$ and

$\mathcal{D}_2 = \left\{ \left(\mathbf{x}_2^{(i)}, \mathbf{s}_2^{(i)} \right), i = 1,2,\ldots,N_2 \right\}$ as a minimum-cost bipartite matching problem of their

corresponding spatial graphs $\mathcal{G}_1 = (\mathcal{V}_1, \mathcal{E}_1, \mathbf{X}_1)$ and $\mathcal{G}_2 = (\mathcal{V}_2, \mathcal{E}_2, \mathbf{X}_2)$:

$$\min_{\mathcal{M}} \sum_{(v_i, v_j) \in \mathcal{M}} \left\| \mathbf{z}_1^{(i)} - \mathbf{z}_2^{(j)} \right\|, \quad \text{s.t.} \quad \mathcal{M} \subset \mathcal{V}_1 \times \mathcal{V}_2, |\mathcal{M}| = m, \#(3)$$

where $\mathbf{z}_1^{(i)}, \mathbf{z}_2^{(j)} \in \mathbb{R}^P$ are node embeddings of v_i and v_j in $\mathcal{G}_1$ and $\mathcal{G}_2$, respectively, $\mathcal{M}$ is a set

of matches of fixed size, and P is the dimensionality of node embeddings. Similarly, we use

matrices $\mathbf{Z}_1 \in \mathbb{R}^{N_1 \times P}$ and $\mathbf{Z}_2 \in \mathbb{R}^{N_2 \times P}$ to denote the combination of cell embeddings of all cells in

the two datasets.

It has been demonstrated in a previous work, that the above matching problem is equivalent to

minimizing the Wasserstein distance between node embeddings from different graphs[16]. SLAT

follows the same approach with adaptations for spatial omics data. Below, we explain how node or

cell embeddings $\mathbf{Z}_1, \mathbf{Z}_2$ can be obtained and optimized for spatial graph alignment.

Construction of holistic cell representations

An accurate alignment of spatial omics datasets should align cells that are similar in both the

molecular modality and the spatial context. In particular, the spatial context can involve various

resolutions, ranging from microenvironments to global positions within the tissue. Inspired by

previous work[16,36,37], we first employ the lightweight graph-convolutional network (LGCN) to derive

a holistic cell representation with all such information integrated for each dataset. A LGCN

propagates and aggregates information along the spatial graph through stepwise concatenations:

$$\widetilde{\mathbf{X}} = f_{\text{LGCN}}(\mathbf{A}, \mathbf{X}) = \text{Concat}(\mathbf{X}, \widehat{\mathbf{A}}\mathbf{X}, \widehat{\mathbf{A}}^2\mathbf{X}, \dots, \widehat{\mathbf{A}}^L\mathbf{X}), \#(4)$$

where $\widehat{\mathbf{A}} = \widetilde{\mathbf{D}}^{-\frac{1}{2}}\widetilde{\mathbf{A}}\widetilde{\mathbf{D}}^{-\frac{1}{2}}$, $\widetilde{\mathbf{A}} = \mathbf{A} + \mathbf{I}$, $\widetilde{\mathbf{D}}$ is the diagonal degree matrix of $\widetilde{\mathbf{A}}$, and L is the maximal

number of steps. The resulting cell representation $\widetilde{\mathbf{X}} \in \mathbb{R}^{N \times (L+1)M}$ is a concatenation of multi-level

information. The first M dimensions correspond to no graph propagation, which is simply a copy of

the omics data $\mathbf{X}$. The second M dimensions correspond to one-step graph propagation, reflecting

the composition of a cell's immediate neighbors, which form its microenvironment. The information

coarsens as the number of steps increases, gradually becoming a representation of rough locations

within the tissue. Thus, $\widetilde{\mathbf{X}}$ contains informative features for spatial alignment at multiple levels of

spatial context.

The cell representation $\widetilde{\mathbf{X}}$ is constructed separately for each dataset by using dataset-specific omics

profiles and adjacency matrices:

$$\begin{aligned}\widetilde{\mathbf{X}}_1 &= f_{\text{LGCN}}(\mathbf{A}_1, \mathbf{X}_1), \\ \widetilde{\mathbf{X}}_2 &= f_{\text{LGCN}}(\mathbf{A}_2, \mathbf{X}_2).\end{aligned} \#(5)$$

521   **Adversarial graph alignment**

522   Based on the holistic cell representations $\widetilde{\mathbf{X}}_1, \widetilde{\mathbf{X}}_2$ described above, we use adversarial alignment to

523   learn cell embeddings $\mathbf{Z}_1, \mathbf{Z}_2$ that minimize the Wasserstein distance for graph matching[16].

524   Specifically, we apply a multilayer perceptron denoted as f_{Z} to mitigate systematic bias in $\widetilde{\mathbf{X}}_1$ and

525    $\widetilde{\mathbf{X}}_2$ that may arise from differences in omics distribution or spatial topology across datasets:

$$\begin{aligned}\mathbf{Z}_1 &= f_{\text{Z}}(\widetilde{\mathbf{X}}_1), \\ \mathbf{Z}_2 &= f_{\text{Z}}(\widetilde{\mathbf{X}}_2).\end{aligned} \#(6)$$

527   We then introduce the Wasserstein discriminator f_{D}, which uses $\mathbf{Z}_1, \mathbf{Z}_2$ as input and tries to

528   maximize the following Wasserstein loss L_W to estimate the Wasserstein distance:

$$L_W = \frac{1}{|\mathcal{V}_1|} \sum_{v_i \in \mathcal{V}_1} f_D\left(\mathbf{z}_1^{(i)}\right) - \frac{1}{|\mathcal{V}_2|} \sum_{v_j \in \mathcal{V}_2} f_D\left(\mathbf{z}_2^{(j)}\right), \#(7)$$

where $\mathbf{z}_1^{(i)}$ is a row in $\mathbf{Z}_1$ corresponding to cell v_i, and $\mathbf{z}_2^{(j)}$ is a row in $\mathbf{Z}_2$ corresponding to cell

v_j. The transformation f_Z can then be adversarially trained to minimize (7), for aligning the

distribution of cell embeddings in the two datasets properly. However, different single-cell spatial

datasets may contain different cell-type proportions or distinct spatial regions. It is thus unreasonable

to assume identical distribution of their cell embeddings as assumed in the standard scheme

described above[38]. Inspired by a previous study[16], we use the output of the Wasserstein discriminator

f_D as a dynamic clipping criterium to select $l \times N_1$ and $l \times N_2$ cells from the two datasets with

minimum Wasserstein distance for adversarial training (Extended Data Fig. 1a):

$$\begin{aligned}
\mathcal{V}_1' &= \arg k\min_{v_i \in \mathcal{V}_1' \subseteq \mathcal{V}_1} f_D\left(\mathbf{z}_1^{(i)}\right), \quad |\mathcal{V}_1'| = l \times N_1, \\
\mathcal{V}_2' &= \arg k\max_{v_j \in \mathcal{V}_2' \subseteq \mathcal{V}_2} f_D\left(\mathbf{z}_2^{(j)}\right), \quad |\mathcal{V}_2'| = l \times N_2,
\end{aligned} \#(8)$$

537   where l is a hyperparameter between 0 and 1. These cells correspond to the most reliable anchors to

538   guide the alignment. The Wasserstein discriminator loss L_W is then modified accordingly as follows:

$$L_W = \frac{1}{|\mathcal{V}_1'|} \sum_{v_i \in \mathcal{V}_1'} f_D\left(\mathbf{z}_1^{(i)}\right) - \frac{1}{|\mathcal{V}_2'|} \sum_{v_j \in \mathcal{V}_2'} f_D\left(\mathbf{z}_2^{(j)}\right). \#(9)$$

This approach ensures that distinct regions across two spatial datasets will not be forcibly aligned.

The results show that SLAT performs best when $l = 0.6$, although the performance depends only

weakly on l (Extended Data Fig. 2a).

To avoid a degenerate solution where all embeddings collapse to a singular point, we adopt an

additional reconstruction term to ensure that the embeddings have sufficient information to

reconstruct input, essentially enhancing model stability. We use a simple multilayer perceptron

network denoted as f_R for data reconstruction, making the following reconstruction loss:

$$L_R = \frac{1}{|\mathcal{V}_1|} \sum_{v_i \in \mathcal{V}_1} \left\| f_R\left(\mathbf{z}_1^{(i)}\right) - \mathbf{x}_1^{(i)} \right\| + \frac{1}{|\mathcal{V}_2|} \sum_{v_j \in \mathcal{V}_2} \left\| f_R\left(\mathbf{z}_2^{(j)}\right) - \mathbf{x}_2^{(j)} \right\|. \#(10)$$

549   **Overall objective**

550   Finally, the overall training objective of SLAT can be summarized as follows:

$$\max_{f_D} \alpha \cdot L_W, \#(11)$$

$$\min_{f_Z, f_R} \alpha \cdot L_W + (1 - \alpha) \cdot L_R, \#(12)$$

where L_W and L_R are defined by Eqs. (9) and (10), respectively, α is a hyperparameter

balancing the contribution of adversarial alignment and data reconstruction. We use stochastic

gradient descent (SGD) with the Adam optimizer to train the SLAT model.

Coordinate matching

Apart from the core model described above, we also provide options to use additional information to

match the spatial coordinates $\mathbf{S}$ of two slices which can help distinguish symmetric structures (e.g.,

left and right hemispheres of the brain) and improve the final matching quality. The goal of

coordinate matching is to roughly align different slices in terms of their overall direction by

estimating an affine transformation matrix $\mathbf{M}$. The exact strategy depends on the type of information

available.

First, guided by expert knowledge of tissue structures, $\mathbf{M}$ can be computed by combining the

scaling, rotation, and translation operations required to obtain a rough alignment between the two

slices. Second, if imaging data like H-E staining images are available, $\mathbf{M}$ can be estimated

following our tutorial based on SimpleElastix[39], which is a state-of-the-art medical image registration

tool. Finally, if no other information is available, we also provide a default solution based on iterative

closet point (ICP)[40], a point-cloud registration algorithm, where we treat the spatial datasets as point

clouds on a two-dimensional plane and uses geometric features for registration. With the obtained **M**

matrix, we consort the coordinates of the two datasets by the following transformation:

$$\mathbf{S}_2' = \mathbf{S}_2 \cdot \mathbf{M}. \#(13)$$

Cell matching and quality assessment

573   With the cell embeddings $\mathbf{Z}_1, \mathbf{Z}_2$ learned by the SLAT core model and the matched spatial

574   coordinates $\mathbf{S}_1$ and $\mathbf{S}_2'$, we match dataset $\mathcal{D}_2$ with $\mathcal{D}_1$ using the following strategy: For cell j in

575   dataset $\mathcal{D}_2$, SLAT first selects the top K most similar $\mathcal{D}_1$ cells in the embedding space $\mathbf{Z}$ as a

576   candidate set $\mathcal{C}_j$, then chooses the one with minimal distance in matched spatial coordinates $\mathbf{S}$ as

577   the final match.

$$\mathcal{C}_j = \arg\, k\max_{v_i \in \mathcal{C}_j \subseteq \mathcal{V}_1} \cos\left(\mathbf{z}_1^{(i)}, \mathbf{z}_2^{(j)}\right), |\mathcal{C}_j| = K, \#(14)$$

$$\text{match}(v_j) = \underset{v_i \in \mathcal{C}_j}{\arg\min} \left\| \mathbf{s}_1^{(i)} - \mathbf{s}_2'^{(j)} \right\|, \#(15)$$

where $\mathbf{s}_1^{(i)}$ and $\mathbf{s}_2'^{(j)}$ are rows in the coordinate matrices $\mathbf{S}_1$ and $\mathbf{S}_2'$, respectively. K is a

hyperparameter that can be adjusted according to the confidence in the coordinate matching result.

The cosine similarity $\cos\left(\mathbf{z}_1^{(i)}, \mathbf{z}_2^{(j)}\right)$ is used to quantify the matching quality. We filter out

unreliable matches by defining a cutoff for cosine similarity. Cell pairs with low matching quality

suggest either biologically distinct changes or technological artifacts and should be carefully checked

(see Results). Empirically, we introduce a default cutoff at 0.8 which can also be adjusted in a

case-by-case manner.

Systematic benchmarks

Benchmark datasets

We selected 10x Visium, MERFISH and Stereo-seq as representative spatial technologies for

benchmarking alignment methods. The 10x Visium dataset comes from consecutive slices of human

dorsolateral prefrontal cortex, containing about 3,000 spots per slice, each spanning 50 μm with over

20,000 genes detected[41]. The MERFISH dataset comes from consecutive slices of mouse

hypothalamic preoptic, with subcellular resolution but only 151 genes detected in total[21]. The

Stereo-seq dataset comes from consecutive slices of an E15.5 mouse embryo, containing over

100,000 single cells per slice, divided into 25 cell types with complex spatial organization, and

detects over 20,000 genes in total[8]. For each technology, we chose the first two slices if the dataset

contains more than two slices. Cell type annotation and tissue region segmentation were obtained

from the original authors whenever possible. Since the Stereo-seq dataset does not provide tissue

segmentation, we segmented the most prominent regions in the embryo, including Brain, Jaw and

face, Spinal cord, Heart, Lung, Liver, and Belly under the guidance of an expert in mouse anatomy.

The spatial segmentation of the MERFISH dataset is provided in publication figures but not in raw

data, so we re-segmented the data as guided by the figures.

Slice processing

For each technology, we first removed the cells/spots that are unannotated in both slices, then rotated

the second slice with a random angle before feeding to the alignment methods.

Benchmarked methods

The benchmarked methods Harmony, PASTE, STAGATE, and SLAT were executed using the

Python packages "harmonypy" (v0.0.6), "paste-bio" (v1.3.0), "STAGATE_pyG" (latest commit

8b9c8ef), and "scSLAT" (v0.2.0), respectively, in Python (v3.8). Seurat was executed using the R

package "Seurat" (v4.1.1) in R (v4.1.3). For each method, we used the default hyperparameter

settings and data preprocessing steps recommended by the original authors.

Evaluation metrics

Intuitively, a proper spatial alignment should align cells matched in both molecular profile and

spatial context. Thus, we reported the cell type matching accuracy and spatial region matching

accuracy simultaneously via the form of confusion matrices. For convenience, we also defined a

"joint accuracy" as the proportion of cells with both cell type and spatial region matched correctly,

which corresponds to the upper right corner of the confusion matrix.

Meanwhile, we also noticed that there exists other metrics commonly used for evaluating graph

alignment, e.g., the edge score[42,43]. However, we did not use it as we found that it can be deceived in

certain situations. For example, Extended Data Fig. 1b shows two graphs with known ground truth

node pairing information. "Alignment 1" is the correct matching (grey lines) and "Alignment 2" has

four mismatched pairs (highlighted by red lines), but the two alignments get the same edge score.

Benchmark tasks

We benchmarked the alignment methods based on the following four tasks: (1) real world alignment,

(2) split slice alignment, (3) scalability test:

In real world alignment, the slices to be aligned are different slices produced from the same position

of the same tissue[5,8,41] (Supplementary Fig. 1). Each algorithm was run eight times with different

model random seeds. Performance of the alignment methods is presented as the confusion matrix as

described above.

In split slice alignment, we split the first slice of each technology used in real world alignment into

two pseudo-slices of equal size by randomly sampling the cells without replacement (Supplementary

Fig. 3). Each algorithm was run eight times with different model random seeds. Performance

quantification is the same as in real world alignment.

For scalability benchmark, we randomly subsampled the Stereo-seq dataset used in real world

benchmark to a range of cell numbers (3200, 6400, 12800, 25600, 51200, 102400). The subsampling

process was repeated eight times with different random seeds (Supplementary Fig. 5).

Benchmark workflow

646 We used Snakemake (v7.12.0) to manage the whole benchmark workflow. All benchmarked methods

647 were allocated 16 cores of Intel Xeon Platinum 8358 CPU, 128 GB of RAM, and a NVIDIA A100

648 GPU with 80 GB VRAM by the Slurm workload management system.

Hyperparameter robustness

We tested SLAT's robustness to key hyperparameters including: (1) neighbor size K for the spatial

graph, (2) input embedding methods, (3) number of training epochs, (4) number of LGCN layers L,

(5) MLP hidden layer dimension, (6) dimension P of SLAT embedding, (7) weight α of the

adversarial loss, (8) dynamic clipping ratio l, (9) learning rate of the SLAT model.

We run SLAT in two homogeneous Stereo-seq datasets from the same E15.5 mouse embryo, each

containing more than 100,000 cells so we randomly subsampled 8,000 cells from each dataset to

save time. Every experiment was run 8 times with different model random seeds (Extended Data Fig.

2).

Robustness to noise in the spatial graph

662 In practice, the spatial graph may be imperfect due to technical limitations. Therefore, we tested

663 SLAT's robustness to graph corruption in the same dataset used in hyperparameter robustness

664 evaluation. Specifically, we randomly masked the edges in the graph by increasing ratios (from 0.1

665 and 0.9). Every experiment was run 8 times with different masking random seeds.

Heterogeneous alignment across distinct technologies and modalities

Visium and Xenium data alignment

The Visium and Xenium datasets were generated from consecutive slices of the human breast cancer

tissue sample[4]. In order to maintain the comparability with the original paper, we chose the exact

same slices used in their analysis (see Fig. 4c of ref[4]). We further selected the shared region between

the two slices as the original authors reported[4] for follow-up analysis.

Considering that the Xenium slice contains more than 100,642 cells while Visium only contains less

than 3,841 spots in same physical region, we used different neighbor sizes proportional to cell

density ($K = 5$ for Visium, and $K = 130$ for Xenium) when constructing the spatial graphs, in

order to ensure that the GCNs have similar spatial receptive fields. SLAT is then run with otherwise

default parameters. We selected Visium triple positive spots based on the number of aligned Xenium

triple positive cells (Supplementary Fig. 6b). 7 spots with more than 2 aligned cells were chosen.

Following the original paper, we did differential gene expression analysis of the SLAT identified and

manually curated triple positive spots against all other spots, respectively, using the Scanpy[44]

function "scanpy.tl.rank_genes_groups" with parameter "methods='wilcoxon'". For comparison, all

methods included in the benchmarks (PASTE, Harmony, Seurat, and Harmony) were applied to the

same Visium and Xenium datasets with their default parameters.

For comparison, all methods benchmarked (PASTE, Harmony, Seurat, and Harmony) were applied

to the same datasets with their default parameters (Extended Data Fig. 5).

SeqFISH data and Stereo-seq data alignment

For Stereo-seq and seqFISH, we chose the E9.5 slice with the most complete cell type annotation by

the original authors (i.e., the slices with the least proportion of unannotated cells), respectively. To

align the chosen seqFISH and Stereo-seq slices, we run SLAT using the default parameters and

filtered out alignment pairs with similarity scores lower than 0.8. We next refined cell type

annotations in the Stereo-seq dataset based on the higher resolution annotation in seqFISH through

label transfer for "Neural crest" cells (Fig. 3b). We also manually annotated "Neural crest" cells for

independent validation. Scanpy[44] was used for the analysis following its official tutorial: the data

were log-normalized using the functions "sc.pp.normalize_total" and "sc.pp.log1p". Highly variable

genes were identified with "sc.pp.highly_variable_genes". The first 50 principal components after

PCA ("sc.tl.pca") were used to generate neighborhood graphs ("sc.pp.neighbors") for computing

UMAP embeddings ("sc.tl.umap") and Leiden clustering ("sc.tl.leiden"). The "Neural crest" cells

were annotated via marker genes of different germ layers (*Foxc2* and *Vcan* for mesoderm, *Msx1*, *Mif*,

and *Dik1*for ectoderm, see Extended Data Fig. 3d).

For comparison, all methods benchmarked (PASTE, STAGATE, Seurat and Harmony) were applied

to the same datasets with their default parameters (Extended Data Fig. 4).

Spatial-ATAC-seq and Stereo-seq data alignment

For cross-modality alignment, the slices to align were two 11.5-day mouse embryo datasets from

Stereo-seq (RNA) and spatial-ATAC-seq (ATAC), respectively. For Stereo-seq, we chose the E11.5

slice with the most complete cell type annotation by the original authors (i.e., the slice with the least

proportion of unannotated cells). For spatial-ATAC-seq, we chose the E11.5 slice with the highest

spatial resolution (20 μm). Given that the current spatial-ATAC-seq data did not cover the entire

embryo due to technical limitations, we extracted the anatomically corresponding regions from the

Stereo-seq dataset under expert guidance.

To project cells from different modalities into a shared latent space, we employed the graph-linked

multi-modality embedding strategy we proposed before [22], then ran SLAT with the default

hyperparameters. Based on the outputted alignment, we transferred cell type labels from Stereo-seq

to spatial-ATAC-seq which was not annotated, and then applied SCENIC+[45] for joint regulatory

inference.

To compare with benchmarked methods (PASTE, Harmony, Seurat and Harmony), we also used the

same multi-modality embedding as their input followed by their default pipeline. Exceptions are

Seurat and STAGATE, which do not support low dimensional embeddings as input, thus cannot be

compared. In addition, we also compared with the original multi-modality embeddings[22] directly

(Supplementary Fig. 7).

Spatial-temporal alignment

We applied SLAT to align E11.5 and E12.5 heterogeneous mouse embryo Stereo-seq datasets with

the default hyperparameters. In order to maintain the comparability with the original paper, we chose

the exact same slices used in their analysis (see Fig. 3a of ref[8]). Regions with lower SLAT similarity

scores were marked in Fig. 5b. To focus on kidney development, we extracted cells labeled as

"Kidney" in E12.5 and their aligned cells in E11.5, then clustered these cells by using the standard

Scanpy clustering pipeline mentioned above and annotated them via well-defined kidney markers[31,32]

(*Osr1*, *Foxc1* and *Podxl* for nephron progenitors; *Uncx*, *Nr2f2*, *Dach*, *Wt1*, *Nphs1*, and *Cd44* for

kidney; see Fig. 4f). To further demonstrate the robustness of SLAT, we rerun the analysis against

two additional slices randomly chosen from E11.5 and E12.5, and obtained similar results

(Supplementary Fig. 8).

For comparison, all methods benchmarked (PASTE, STAGATE, Seurat and Harmony) were applied

to the same datasets with their default parameters (Extended Data Fig. 8).

Data availability

All datasets used in this study were already published and were obtained from public data

repositories. See Supplementary Table 1 for detailed information on the entire datasets used in this

study, including publication and downloading URLs.

Code availability

The SLAT framework was implemented in the "scSLAT" Python package, which is available at

https://github.com/gao-lab/SLAT. For reproducibility, the scripts for all benchmarks were assembled

using Snakemake (v7.12.0), which is also available in the above repository.