The BTB-ZF gene *Bm-mamo* regulates pigmentation in silkworm caterpillars

Songyuan Wu, Xiaoling Tong, Chenxing Peng, Jiangwen Luo, Chenghao Zhang, Kunpeng Lu,

Chunlin Li, Xin Ding, Xiaohui Duan, Yaru Lu, Hai Hu, DuanTan, Fangyin Dai[*]

Affiliations

State Key Laboratory of Resource Insects, Key Laboratory of Sericultural Biology and Genetic

Breeding, Ministry of Agriculture and Rural Affairs, College of Sericulture, Textile and Biomass

Sciences, Southwest University, Chongqing 400715, China

Songyuan Wu fly_jungle@163.com

Xiaoling Tong xltong@swu.edu.cn

Chenxing Peng mxtaste@sina.com

Jiangwen Luo luo0510@email.swu.edu.cn

Chenghao Zhang hangsdo0452@163.com

Kunpeng Lu lukunpeng@swu.edu.cn

Chunlin Li lclin0522@swu.edu.cn

Xin Ding dingxin2019@swu.edu.cn

Xiaohui Duan duanxiaohui@email.swu.edu.cn

Yaru Lu 17723940349@163.com

Hai Hu huhaiswu@163.com

DuanTan tanzeduan@163.com

Fangyin Dai fydai@swu.edu.cn

*Corresponding author

Fangyin Dai

fydai@swu.edu.cn

86-023-68250793

Southwest University, No. 2 Tiansheng Road, Beibei District, Chongqing 400715, China

Abstract

The color pattern of insects is one of the most diverse adaptive evolutionary phenotypes.

However, the molecular regulation of this color pattern is not fully understood. In this study, we

found that the transcription factor Bm-mamo is responsible for *black dilute* (*bd*) allele mutations

in the silkworm. Bm-mamo belongs to the BTB zinc finger family and is orthologous to mamo in

Drosophila melanogaster. This gene has a conserved function in gamete production in

Drosophila and silkworms and has evolved a pleiotropic function in the regulation of color

patterns in caterpillars. Using RNAi and clustered regularly interspaced short palindromic

repeats (CRISPR) technology, we showed that Bm-mamo is a repressor or has dark melanin

patterns in the larval epidermis. Using in vitro binding assays and gene expression profiling in

wild-type and mutant larvae, we also showed that *Bm-mamo* likely regulates the expression of

related pigment synthesis and cuticular protein genes in a coordinated manner to mediate its role

in color pattern formation. This mechanism is consistent with the dual role of this transcription

factor in regulating both the structure and shape of the cuticle and the pigments that are

embedded within it. This study provides new insight into the regulation of color patterns as well

as into the construction of more complex epidermis features in some insects.

43
Introduction

Insects often display stunning colors, and the patterns of these colors have been shown to be

involved in behavior(*1*), immunity(*2*), thermoregulation(*3*), and UV protection(*4*); in particular,

visual antagonism of predators via aposematism(*5*), mimicry(*6*), cryptic color patterns(*7*) or

some combination of the above (*8*). In addition, the color patterns are divergent and convergent

among populations(*9*). Due to these striking visual features and highly active adaptive

evolutionary phenotypes, the genetic basis and evolutionary mechanism of color patterns have

long been a topic of interest.

Insect coloration can be pigmentary, structural, or bioluminescent. Pigments are synthesized by

insects themselves and form solid particles that are deposited within the cuticle of the body

surface and the scales of the wings(*10, 11*). Interestingly, recent studies have shown that bile

pigments and carotenoid pigments synthesized through biological synthesis are incorporated into

body fluids and fill in the wing membranes of two butterflies (*Siproeta stelenes* and *Philaethria*

diatonica) via hemolymph circulation, providing color in the form of liquid pigments(*12*). These

pigments form colors by selective absorption and/or scattering of light depending on their

physical properties(*13*). However, structural color refers to colors, such as metallic colors and

iridescence, generated by optical interference and grating diffraction of the

microstructure/nanostructure of the body surface or appendages (such as scales)(*14, 15*). Pigment

color and structural color are widely distributed in insects and can only be observed by the naked

eye in illuminated environments. However, some insects, such as fireflies, exhibit colors (green

to orange) in the dark due to bioluminescence(*16*). Bioluminescence occurs when luciferase

catalyzes the oxidation of small molecules of luciferin(*17*). In conclusion, the color patterns of

insects have evolved to become highly sophisticated and are closely related to their living

environment. For example, cryptic color can deceive animals via high similarity to the

surrounding environment. However, the molecular mechanism by which insects form precise

color patterns to match their living environment has not been determined.

Recent research has identified the metabolic pathways associated with related pigments, such as

melanins, pterins, and ommochromes, in Lepidoptera(*18*). A deficiency of enzymes in the

pigment metabolism pathway can lead to changes in color patterns(*19, 20*). In addition to

pigment synthesis, the microstructure/nanostructure of the body surface and wing scales are

important factors that influence body color patterns. The body surface and wing scales of

Lepidoptera are composed mainly of cuticular proteins (CPs)(*21, 22*). There are multiple genes

encoding CPs in Lepidopteran genomes. For example, in *Bombyx mori,* more than 220 genes

encode CPs(*23*). However, the functions of CPs and the molecular mechanisms underlying their

fine-scale localization in the cuticle are still unclear.

In addition, several pleiotropic factors, such as *wnt1(24)*, *Apontic-like(25)*, *clawless(26)*,

abdominal-A(27), *abdominal-B(28)*, *engrailed(29)*, *antennapedia(30)*, *optix(31)*, *bric* à *brac*

(*bab*)(*32*), and *Distal-less* (*Dll*)(*33*), play important roles in the regulation of color patterns.

The molecular mechanism by which these factors participate in the regulation of body color

patterns needs further study. In addition, there are many undiscovered factors in the gene

regulatory network (GRN) involved in the regulation of insect color patterns. The identification

of new color pattern regulatory genes and the study of their molecular mechanisms would be

helpful for further understanding color patterns.

Silkworms (*B. mori*) have been completely domesticated for more than 5000 years and are

famous for their silk fiber production(*34*). Because of this long history of domestication,

artificial selection and genetic research, its genetic background has been well studied, and

approximately 200 strains of mutants with variable color patterns have been preserved(*35*),

which provides a good resource for color pattern research. The *black dilute* (*bd*) mutant, which

exhibits recessive Mendelian inheritance, has a dark gray larval body color, and the female is

sterile. *Black dilute fertile* (*bd^f*) is an allele that leads to a lighter body color than *bd* and fertile

females (Fig. 1). The two mutations were mapped to a single locus at 22.9 centimorgans (cM) of

linkage group 9(*36*). No pigmentation-related genes have been reported at this locus. Thus, this

research may reveal a new color pattern-related gene, which stimulated our interest.

Results

Candidate gene of the *bd* allele

To identify the genomic region responsible for the *bd* alleles, positional cloning of the *bd* locus

was performed. Due to the female infertility of the *bd* mutant and the fertility of females with the

bd^f allele, we used *bd^f* and the wild-type Dazao strain as parents for mapping analysis.

The 1162 back-crossed filial 1st (BC1M) generation individuals from *bd^f* and Dazao were

subjected to fine mapping with molecular markers (Fig. 2). A genomic region of approximately

390 kilobases (kb) was responsible for the *bd* phenotype (Table S1). According to the SilkDB

database(*37*), this region included five predicted genes (*BGIBMGA012516*, *BGIBMGA012517*,

BGIBMGA012518, *BGIBMGA012519* and *BGIBMGA014089*). In addition, we analyzed the

predictive genes for this genomic region from the GenBank(*38*) and SilkBase(*39*) databases (Fig.

S1). The number of predicted genes varied among the different databases. We performed

sequence alignment analysis of the predicted genes in the three databases to determine their

correspondence. Real-time quantitative polymerase chain reaction (qPCR) was subsequently

performed, which revealed that *BGIBMGA012517* and *BGIBMGA012518* were significantly

downregulated on Day 3 of the fifth instar of the larvae in the *bd* phenotype individuals, while

there was no difference in the expression levels of the other genes (Fig. S1). These two genes

were predicted to be associated with a single locus (*LOC101738295*) in GenBank. To determine

the gene structures of *BGIBMGA012517* and *BGIBMGA012518*, we used forward primers for

the *BGIBMGA012517* gene and reverse primers for the *BGIBMGA012518* gene to amplify

cDNA from the wild-type Dazao strain. By gene cloning, the predicted genes *BGIBMGA012517*

and *BGIBMGA012518* were proven to be one such gene. For convenience, we temporarily called

this the *12517/8* gene.

The *12517/8* gene produces two transcripts; the open reading frame (ORF) of the long transcript

is 2397 bp, and the ORF of the short transcript is 1824 bp, which is the same 5'-terminus as that

of the wild-type Dazao strain (Fig. S2). The *12517/8* gene showed significantly lower expression

in the *bd*^{f} mutant (Fig. S3), and multiple variations were found in the region nearby this gene by

comparative genomic analysis between *bd*^{f} and Dazao (Table S2). In addition, *12517/8* was

completely silenced due to the deletion of DNA fragments (approximately 168 kb) from the first

upstream intron and an insertion of 3560 bp in the *bd* mutant (Fig. S4).

To predict the function of the *12517*/8 gene, we performed a BLAST search using its full-length

sequence and found a transcription factor, the *maternal gene required for meiosis* (*mamo*) in *D.*

melanogaster, that had high sequence similarity to that of the *12517/8* gene (Fig. S5). Therefore,

we named the *12517/18* gene *Bm-mamo*; the long transcript was designated *Bm-mamo-L*, and the

short transcript was designated *Bm-mamo-S*.

The *mamo* gene belongs to the Broad-complex, Tramtrack and bric à brac/poxvirus zinc finger

protein (BTB-ZF) family. In the BTB-ZF family, the zinc finger serves as a recognition motif for

DNA-specific sequences, and the BTB domain promotes oligomerization and the recruitment of

other regulatory factors(*40*). Most of these factors are transcription repressors, such as nuclear

receptor corepressor (N-CoR) and silencing mediator for retinoid and thyroid hormone receptor

(SMRT)(*41*), but some are activators, such as p300(*42*). Therefore, these features commonly

serve as regulators of gene expression. *mamo* is enriched in embryonic primordial germ cells

(PGCs) in *D. melanogaster*. Individuals deficient in *mamo* are able to undergo oogenesis but fail

to execute meiosis properly, leading to female infertility in *D. melanogaster(43)*. *Bm-mamo* was

identified as an important candidate gene for further analysis.

Expression pattern analysis of *Bm-mamo*

To analyze the expression profiles of *Bm-mamo*, we performed quantitative PCR. The expression

levels of the *Bm-mamo* gene in the Dazao strain were investigated throughout the body at

different developmental stages, from the embryonic stage to the adult stage. The gene was highly

expressed in the molting stage of caterpillars, and its expression was upregulated in the later

pupal and adult stages (Fig. 3A). This finding suggested that the *Bm-mamo* gene responds to

ecdysone and participates in the processes of molting and metamorphosis in silkworms.

According to the investigation of tissue-specific Dazao strain expression in 5th-instar 3rd-day

larvae, the midgut, head, and epidermis exhibited high expression levels; the trachea, nerves, silk

glands, testis, ovary, muscle, wing disc and Malpighian tubules exhibited moderate expression

levels; and the blood and fat bodies exhibited low expression levels (Fig. 3B). These findings

suggested that *Bm-mamo* is involved in the developmental regulation of multiple silkworm

tissues. Due to the melanism of the epidermis of the *bd* mutant and the high expression level of

the *Bm-mamo* gene in the epidermis, we measured the expression level of this gene in the

epidermis of the 4th to 5th instars of the Dazao strain. In the epidermis, the *Bm-mamo* gene was

upregulated during the molting period, and the highest expression was observed at the beginning

of molting (Fig. 3C).

Functional analyses of *Bm-mamo*

To study the function of the *Bm-mamo* gene, we carried out an RNA interference (RNAi)

experiment. Short interfering RNA (siRNA) was injected into the hemolymph of silkworms, and

an electroporation experiment(*44*) was immediately conducted. We found significant melanin

pigmentation in the epidermis of the newly molted 5th-instar larvae. These findings indicate that

Bm-mamo deficiency can cause melanin pigmentation (Fig. 4). The melanistic phenotype of the

RNAi individuals was similar to that of the *bd* mutants.

In addition, gene knockout was performed. Ribonucleoproteins (RNPs) generated from guide

RNA (gRNA) and recombinant Cas9 protein were injected into 450 silkworm embryos. In the

G0 generation, individuals with a mosaic melanization phenotype were found. These melanistic

individuals were raised to moths and subsequently crossed. A homozygous line with a gene

knockout was obtained through generations of screening. The gene-edited individuals had a

significantly melanistic body color, and the female moths were sterile (Fig. 5).

These results indicated that the *Bm-mamo* gene negatively regulates melanin pigmentation in

caterpillars and participates in the reproductive regulation of female moths.

Downstream target gene analysis

Bm-mamo belongs to the zinc finger protein family, which specifically recognizes downstream

DNA sequences according to their zinc fingers. We identified homologous genes of *mamo* in

multiple species and conducted a phylogenetic analysis (Fig. S6). Sequence alignment revealed

that the amino acid residues of the zinc finger motif of the mamo-S protein were highly

conserved among 57 species. The insufficient ability of the different species in GenBank to

predict alternative transcripts of mamo may have resulted in the mamo-L protein being found in

only 30 species (Fig. S7). As the mamo protein contains a tandem Cys_2His_2 zinc finger (C2H2-

ZF) motif, it can directly bind to DNA sequences. Previous research has suggested that the ZF-

DNA binding interface can be understood as a "canonical binding model", in which each finger

contacts DNA in an antiparallel manner. The binding sequence of the C2H2-ZF motif is

determined by the amino acid residue sequence of its α-helical component. The first amino acid

residue in the α-helical region of the C2H2-ZF domain is at position 1, and positions -1, 2, 3, and

6 are key amino acids involved in the recognition and binding of DNA. The residues at positions

-1, 3, and 6 specifically interact with base 3, base 2, and base 1 of the DNA sense sequence,

respectively, while the residue at position 2 interacts with the complementary DNA strand(*45,*

46). To analyze the downstream target genes of mamo, we first predicted the DNA-binding

motifs of these genes using online software (http://zf.princeton.edu) based on the canonical

binding model(*47*). In addition, the DNA-binding sequence of mamo (<u>TGCGT</u>) in *Drosophila*

was confirmed by electrophoretic mobility shift assay (EMSA)(*48*), which has a consensus

sequence with the predicted binding site sequence of Bm-mamo-S (G<u>TGCGT</u>GGC), and the

predicted sequence was longer. This indicates that the predicted results for the DNA-binding site

have good reliability. Furthermore, the predicted DNA-binding sites of Bm-mamo-L and Bm-

mamo-S were highly consistent with those of mamo orthologs in different species (Fig. S8). This

finding suggested that the protein may regulate similar target genes between species.

C2H2-ZF transcription factors function by recognizing and binding to *cis*-regulatory sequences

in the genome, which harbor *cis*-regulatory elements (CREs)(*49*). CREs are broadly classified as

promoters, enhancers, silencers or insulators(*50*). CREs are often near their target genes, such as

enhancers, which are typically located upstream (5') or downstream (3') of the gene they regulate

or in introns, but approximately 12% of CREs are located far from their target gene(*51*).

Therefore, we first investigated the 2 kb upstream and downstream regions of the predicted

genes in silkworms.

The predicted position weight matrices (PWMs) of the recognized sequences of Bm-mamo

protein and the Find Individual Motif Occurrences (FIMO) software of MEME were used to

perform silkworm whole-genome scanning for possible downstream target genes. The 2 kb

upstream and downstream regions of the 14,623 predicted genes in silkworms were investigated.

A total of 10,622 genes contained recognition sites within 2 kb of the upstream/downstream

region of the Bm-mamo protein in the silkworm genome (Fig. S9, Table S3 & Table S4).

Moreover, we compared the transcriptome data of integument tissue between homozygotes and

heterozygotes of the *bd* mutant at the 4th instar/beginning molting stage[52]. In the integument

tissue, 10,072 genes (~69% of the total predicted genes of silkworm) were expressed in

heterozygotes, and 9,853 genes (~67% of the total predicted genes) were expressed in

homozygotes of the *bd* mutant. In addition, there were 191 genes whose expression significantly

differed between homozygotes (*bd*/*bd*) and heterozygotes (+/*bd*) according to comparative

transcriptome analysis (Table S5)[52]. Protein functional annotation was performed, and 19 CP

genes were significantly differentially expressed between heterozygotes and homozygotes of *bd*.

In addition, the orthologs of these CPs were analyzed in *Danaus plexippus*, *Papilio xuthu* and *D.*

melanogaster (Table S6). Furthermore, we identified 53 enzyme-encoding genes, 17

antimicrobial peptide genes, 6 transporter genes, 5 transcription factor genes, 5 cytochrome

genes, and others. Among the differentially expressed genes (DEGs), CP was significantly

enriched, and previous studies have shown that CPs can affect pigmentation[53]. Therefore, we

first investigated the expression of the CP genes. Among them, 18 CP genes had Bm-mamo

binding sites in the upstream and downstream 2 kb genomic regions. In addition, we investigated

the expression levels of the 18 CP genes in the integument from the 4th instar (Day 1) to the

beginning of the 5th instar in the *Bm-mamo* knockout lines (Fig. 7& Fig. S10). The CP genes

were significantly upregulated at one or several time points in homozygous (*mamo*$^{-}$/*mamo*$^{-}$)

individuals compared with heterozygous (*mamo*$^{-}$/+) *Bm-mamo* knockout individuals.

Interestingly, the expression of the CP gene *BmorCPH24* was significantly upregulated at the

feeding stage in *Bm-mamo* knockout homozygotes. Previous studies have shown that

BmorCPH24 deficiency can lead to a marked decrease in pigmentation in silkworm larvae[53].

Therefore, the expression of some CP genes may be necessary for determining the color patterns

of caterpillars.

In addition, the synthesis of pigment substances is an important driver of color patterns(*54*).

Eight key genes (*TH*, *DDC*, *aaNAT*, *ebony*, *black*, *tan*, *yellow* and *laccase2*) involved in melanin

synthesis(*11*) were investigated in heterozygous and homozygous *Bm-mamo* gene knockout lines.

Among them, *yellow*, *tan*, and *DDC* were significantly upregulated during the molting period in

the homozygous *Bm-mamo* knockout individuals (Fig. 6). The upstream and/or downstream 2 kb

genomic regions of *yellow*, *tan* and *DDC* contain the binding site of the Bm-mamo protein. In

addition, the expression of *yellow*, *DDC* and *tan* can promote the generation of melanin(*55*).

To explore the interaction between the Bm-mamo protein and its binding sequence, EMSA was

conducted. The binding site of Bm-mamo-S (C<u>TGCGT</u>GGT) was located approximately 70 bp

upstream of the transcription initiation site of the *Bm-yellow* gene. The EMSA results showed

that the Bm-mamo-S protein expressed in prokaryotes can bind to the C<u>TGCGT</u>GGT sequence

in vitro (Fig. S11). This finding suggested that the Bm-mamo-S protein can bind to the upstream

region of the *Bm-yellow* gene and regulate its transcription.

Therefore, the *Bm-mamo* gene may control the color pattern of caterpillars by regulating key

melanin synthesis genes and 18 CP genes.

Discussion

Insects have evolved many important phenotypes during the process of adapting to the

environment. Among these traits, the color pattern is one of the most interesting. The

biochemical metabolic pathways of pigments, such as melanin, ommochromes, and pteridines,

have been identified in insects(*18*). However, the regulation of pigment metabolism-related

genes and the processes involved in the transport and deposition of pigment substances are

unclear. In this study, we discovered that the *Bm-mamo* gene negatively regulates melanin

pigmentation in caterpillars. When this gene is deficient, the body color of silkworms exhibits

substantial melanism, and changes in the expression of some melanin synthesis genes and some

CP genes are also significant.

Structural genes of melanin synthesis

Deficiencies of some genes in the melanin synthesis pathway, such as *TH*, *yellow*, *ebony*, *tan*,

and *aaNAT,* can lead to variations in color patterns *(11)*. However, they often lead to localized

pigmentation changes only in later-instar caterpillars. For example, in *yellow*-mutant silkworms,

the eye spot, lunar spot, star spot, spiracle plate, head, and some sclerotized areas appear reddish

brown, and the other cuticle is consistent with that in the wild type in later instars of larvae (third

instar to fifth instar)(*56*). This situation is highly similar to that of the *tan* mutant in

silkworms(*57*). Why do the *yellow* and *tan* phenotypes appear only in a limited cuticle region in

later instars of silkworm larvae? One possible reason is that the expression of *yellow* and *tan* is

limited to a certain region by transcription regulators. Alternatively, other factors, such as CPs in

the cuticle, may limit pigmentation by interacting with pigment substances in the cuticle.

On the one hand, we investigated the expression levels of *yellow* and *tan* in the pigmented region

(lunar spot) and nonpigmented region of the epidermis during the 4th molting of the wild-type

Dazao strain. The expression level of *yellow* was significantly upregulated in the lunar spot (the

epidermis on the dorsum of the fifth body segment) compared with the nonpigmented region (the

epidermis on the dorsum of the sixth body segment) at 6 hours and 12 hours after molting.

Moreover, *tan* was significantly upregulated 18 hours after molting in the lunar spot region (Fig.

S12). This finding suggested that the upregulated expression of pigment synthesis genes at key

time points may be important for pigmentation. However, *yellow* and *tan* were still moderately

expressed in the nonpigmented epidermis, although they did not cause significant melanin

pigmentation. This finding indicates that pigment synthesis alone cannot determine the

predominant color pattern of the cuticle in caterpillars.

Cuticular proteins participate in pigmentation

On the other hand, synthesized pigment substances need to be transported from epidermal cells

and embedded into the cuticle to allow pigmentation. Therefore, the correct cuticle structure and

location of cuticular proteins in the cuticle may be important factors affecting pigmentation.

Previous studies have shown that a lack of expression of *BmorCPH24*, which encodes an

important component of the endocuticle, can lead to dramatic changes in body shape and a

significant reduction in the pigmentation of caterpillars(*53*). We crossed *Bo* (*BmorCPH24* null

mutation) and *bd* to obtain F_1 ($Bo/+^{Bo}$, *bd*/+) and then self-crossed F_1 to observe the phenotype of

F_2. The area of lunar spots and star spots decreased, and light-colored stripes appeared on the

body segments; however, the other areas still exhibited significant melanin pigmentation in

double-mutation (*Bo*, *bd*) individuals (Fig. S13). However, in previous studies, the introduction

of *Bo* into *L* (ectopic expression of *wnt1* results in lunar stripes generated on each body

segment)(*24*) and *U* (overexpression of *SoxD* results in excessive melanin pigmentation of the

epidermis)(*58*) strains by genetic crosses markedly reduced the pigmentation of *L* and *U*(*53*).

Interestingly, there was a more significant decrease in pigmentation in the double mutants (*Bo*, *L*)

and (*Bo*, *U*) than in (*Bo*, *bd*). These findings suggested that *Bm-mamo* has a greater ability than

does *wnt1* and *SoxD* to regulate pigmentation. On the one hand, mamo may be a stronger

regulator of the melanin metabolic pathway, and on the other hand, mamo may regulate other CP

genes to reduce the impact of BmorCPH24 deficiency.

How do CPs affect pigmentation? One study showed that some CPs can form "pore canals" to

transport macromolecules(*59*). In addition, some CPs can be crosslinked with catecholamines,

which are synthesized in the melanin metabolism pathway(*60*). Because there are no live cells in

the cuticle, melanin precursor substances may be transported by the pore canals formed by some

CPs and fixed to specific positions through cross-linking with CPs. The cuticular protein

TcCPR4 is needed for the formation of pore canals in the cuticle of *Tribolium castaneum(61)*. In

contrast, the vertical pore canal is lacking in the less pigmented cuticles of *T. castaneum(62)*.

This finding suggested that the pore canals constructed by TcCPR4 may transport pigments and

contribute to cuticle pigmentation in *T. castaneum*. Moreover, the melanin metabolites N-

acetyldopamine (NADA) and N-β-alanyldopamine (NBAD) can target and sclerotize the cuticle

by cross-linking with specific cuticular proteins (*57*). This finding suggested that pigments

interact with specific CPs, thereby affecting pigmentation, hardening properties and the structure

of the cuticle. Interestingly, a study showed that in addition to absorbing specific wavelengths,

pigments can affect cuticle polymerization, density, and the refractive index, which in turn

affects the reflected wavelengths that produce structural color in butterfly wing scales(*63*). This

implies that the interaction between pigments and CPs can be very subtle, resulting in the

formation of unique nanostructures, such as those on wing scales, that produce brilliant structural

colors.

Consequently, to maintain the accuracy of the color pattern, the localization of CPs in the cuticle

may be very important. In a previous study employing microarray analysis, different CPs were

found in differently colored areas of the epidermis in *Papilio xuthus* larvae(*64*). We investigated

whether the CP genes were highly expressed in the black region of *Papilio xuthus* caterpillars.

Thirteen orthologous genes were found in silkworms (Table S7). Among them, 11 genes

(*BmorCPR67*, *BmorCPR71*, *BmorCPR76*, *BmorCPR79*, *BmorCPR99*, *BmorCPR107*,

BmorCPT4, *BmorCPH5*, *BmorCPFL4*, *BmorCPG27* and *BmorCPG4*) were significantly

upregulated in homozygous (*mamo⁻/mamo⁻*) knockout individuals at some time points from the

3rd day of the 4th instar to the 5th instar; *BmorCPG4* was also among the 18 previously detected

CP genes, and two genes (*BmorCPG38* and *BmorCPR129*) were not differentially expressed

between homozygous (*mamo⁻/mamo⁻*) and heterozygous (*mamo⁻/+*) individuals (Fig. S14).

The expression of the 28 CP genes mentioned above was significantly upregulated in

homozygous (*mamo⁻/mamo⁻*) gene knockout individuals at several stages. These CPs may be

involved in the transportation or cross-linking of melanin in the cuticle. However, there were no

differences in the expression levels of these genes during some periods compared with those in

the control group, and the expression of some genes was significantly downregulated at some

time points in the melanic individuals (Fig. 7). This finding suggested that the regulation of CP

genes is complex and may involve other transcription factors and feedback effects. CPs are

essential components of the insect cuticle and are involved in cuticular microstructure

construction (*65*), body shape development(*66*), wing morphogenesis(*67*), and pigmentation(*53*).

CP genes usually account for more than 1% of the total genes in an insect genome and can be

categorized into several families, including CPR, CPG, CPH, CPAP1, CPAP3, CPT, CPF and

CPFL(*68*). The CPR family is the largest group of CPs and contains a chitin-binding domain

called the Rebers and Riddiford motif (R&R)(*69*). The variation in the R&R consensus sequence

can be divided into three subfamilies (RR-1, RR-2, and RR-3)(*70*). Among the 28 CPs, 11 RR-1

genes, 6 RR-2 genes, 4 hypothetical cuticular protein (CPH) genes, 3 glycine-rich cuticular

protein (CPG) genes, 3 cuticular protein Tweedle motif (CPT) genes, and 1 CPFL (like the CPFs

in a conserved C-terminal region) gene were identified. The RR-1 consensus among species is

usually more variable than the RR-2 consensus, which suggests that RR-1 may have a species-

specific function. RR-2 often clustered into several branches, which may be due to gene

duplication events in co-orthologous groups and may result in conserved functions between

species (*71*). The classification of CPH was based on the lack of known motifs. In the epidermis

of Lepidoptera, CPH genes often exhibit high expression levels. For example, *BmorCPH24* has

the highest expression level in the silkworm larval epidermis(*72*). The CPG protein is rich in

glycine. The CPH and CPG genes are less commonly found in insects outside the order

Lepidoptera (*73*). This finding suggested that these genes may provide species-specific functions

for Lepidoptera. CPT contains a Tweedle motif, and the *TweedleD1* mutation has a dramatic

effect on body shape in *D. melanogaster(74)*. The CPFL family members are relatively

conserved among species and may be involved in the synthesis of larval cuticles(*75*). CPT and

CPFL may have relatively conserved functions among insects. The CP genes are a group of

rapidly evolving genes, and their copy numbers may undergo significant changes in different

species. In addition, RNAi experiments on 135 CP genes in the brown planthopper (*Nilaparvata*

lugens) revealed that deficiency of 32 CP genes leads to significant defects in phenotypes, such

as lethality and developmental retardation. These findings suggested that the 32 CP genes are

indispensable and that other CP genes may have redundant and complementary functions (*76*). In

previous studies, it was found that the construction of the larval cuticle of silkworms requires the

precise expression of more than two hundred CP genes(*22*). The production, interaction, and

deposition of CPs and pigments are complex and precise processes, and our research showed that

Bm-mamo plays an important regulatory role in this process in silkworm caterpillars. To further

understand the role of CPs, future work should aim to identify the functions of important

cuticular protein genes and the deposition mechanism in the cuticle.

Maturation of pigment granules

In addition, among the 191 DEGs found in the comparative transcriptome data, we also

discovered several interesting genes. For example, *BGIBMGA013242*, which encodes a major

facilitator superfamily protein (MFS) named *BmMFS* and is responsible for the *cheek and tail*

spot (cts) mutant, was significantly upregulated in the *bd* mutant. Deficiency of this gene

(*BmMFS*) results in chocolate-colored head and anal plates on the silkworm caterpillar(*77*). In

the ommochrome metabolic pathway, *Bm-re*, which encodes an MFS protein, may function in the

transportation of several amino acids, such as cysteine or methionine, into pigment granules(*78,*

79). Therefore, the encoded product of *BmMFS* may participate in pigmentation by promoting

the maturation of pigment granules. Moreover, *BGIBMGA013576* and *BGIBMGA013656*, which

encode MFS domain-containing proteins belonging to the solute carrier family 22 (SLC22) and

solute carrier family 2 (SLC2) families, respectively, were significantly upregulated in *bd*. These

two genes may participate in pigmentation in a similar manner to that of the *BmMFS* gene. In

addition, the *red Malpighian tubule* (*red*) gene encodes a LysM domain-containing protein. *Red*

deficiency results in a significant decrease in orange pterin in the wing of *Colias* butterflies.

Research suggests that the product of *red* may interact with V-ATPase to modulate vacuolar pH

across a variety of endosomal organelles, thereby affecting the maturation of pigment granules in

cells(*80*). These findings indicate that the maturation of pigment granules plays an important role

in the coloring of Lepidoptera and that *Bm-mamo* may be involved in regulating the maturation

of pigment granules in the epidermal cells of silkworms.

Bm-mamo may affect the synthesis of melanin in epidermal cells by regulating *yellow*, *DDC*, and

tan, regulate the maturation of melanin granules in epidermal cells through *BmMFS*, and affect

the deposition of melanin granules in the cuticle by regulating CP genes, thereby

comprehensively regulating the color pattern of caterpillars.

The upstream sequence of *Bm-mamo*

Moreover, in *D. melanogaster*, *mamo* is needed for functional gamete production. In the

silkworm, *Bm-mamo* has a conserved function in female reproduction. The DNA-binding motifs

of the mamo proteins in silkworms and fruit flies are highly conserved, suggesting that they may

regulate the same downstream target genes. However, *mamo* has developed new functions in

color pattern regulation in silkworm caterpillars. We found that in *D. melanogaster*, the *mamo*

gene is expressed mainly during the adult stage; it is not expressed during the 1st instar or 2nd

instar larval stage and has very low expression at the 3rd instar (Table S8). In addition, several

binding sites of mamo were found near the TSS of *yellow* in *D. melanogaster* (Fig. S15). The

yellow gene is a key melanin metabolic gene with upstream and intron sequences that have been

identified as multiple CREs, and it has been considered a research model for CREs (*81, 82*). This

difference may be due to a change in the expression pattern of *mamo,* which led to the

development of a new function for this gene in regulating coloration in silkworm caterpillars.

Changes in gene expression patterns are generally believed to be the result of the evolution of

CREs. Because CREs play important roles in the spatiotemporal expression pattern regulation of

genes, many CREs are found in noncoding regions, which are relatively prone to sequence

variation compared with the sequences of coding genes(*49*). However, the molecular mechanism

underlying the sequence changes in CREs is unclear.

TFs, because they recognize relatively short sequences, generally between 4 and 20 bp in length,

can have many binding sites in the genome(*83*). Therefore, one member of the TF family has the

potential to regulate many genes in one genome. Single regulation of specific TFs can lead to

patterned changes in the expression of multiple downstream genes, enabling organisms to adapt

to the environment by altering a given type of trait. For example, the marine form of the three-

spined stickleback (*Gasterosteus aculeatus*) has thick armor, whereas the lake population (which

was recently derived from the marine form) does not. Research has shown that pelvic loss in

different natural populations of three-spined stickleback fishes occurs via regulatory mutations

resulting from the deletion of a tissue-specific enhancer (*Pel*) of the *pituitary homeobox*

transcription factor 1 (*Pitx1*) gene. The researchers genotyped 13 pelvic-reduced populations of

three-spined sticklebacks from disparate geographic locations. Nine of the 13 pelvic-reduced

stickleback populations had sequence deletions of varying lengths, all of which were located at

the *Pel* enhancer (*84*).

We investigated nucleotide diversity in 51 wild silkworms and 171 domesticated silkworms (*34,*

85). The nucleotide diversity of introns and upstream sequences of *Bm-mamo* in wild silkworms

was significantly greater than that in domestic silkworms. In addition, the approximately 1 kb

genomic region upstream had high fixation indices (F_{ST}) (Fig. S16). This indicates a significant

degree of differentiation between the wild strains in this genomic region. Multiple sequence

alignment analysis of this region was performed between 12 wild silkworms and 12 domestic

silkworms (Fig. S17 & Table S9). The results showed that the wild silkworm group had a greater

degree of nucleotide variation, while the sequences in domestic silkworms were highly

conserved. The domestic silkworm group had two different forms: one had a long interspersed

nuclear element (LINE) inserted at approximately 4.5 kb, and the other did not have this

transposon sequence. It is suggested that this genomic region is prone to variation in wild

silkworm and is fixed in domestic silkworms. The larvae of domesticated strains often have a

lighter or even pale body color, but the wild-type strains have a darker color. The *Bm-mamo* gene

may be involved in the domestication of silkworms.

The function of TFs is to precisely regulate the temporal and spatial expression of target genes.

Therefore, the expression of TFs requires strict regulation. There are long intergenic regions

upstream of many important TFs, dozens of kilobase pairs (Kb) to hundreds of Kb, which may

contain many CREs for better control its expression pattern. It has often been believed that

changes in CREs are caused by random mutations. However, recent studies have shown that

there is a mutation bias in the genome; compared with that in the intergenic region, the mutation

frequency is reduced by half inside gene bodies and by two-thirds in essential genes. In addition,

they compared the mutation rates of genes with different functions. The mutation rate in the

coding region of essential genes (such as translation) is the lowest, and the mutation rates in the

coding region of specialized functional genes (such as environmental response) are the highest.

These patterns are mainly affected by the features of the epigenome(*86*). Due to the plasticity of

epigenomic features, mutations bias associated with epigenomes may even lead to environmental

influences on mutations (*87*). This finding suggested that different functional regions in the

genome are subject to distinct controls and that some sequences can undergo procedural changes

under environmental changes. Therefore, some variations in the CRE of TFs in response to

environmental changes may be mutation bias.

Materials and Methods

Silkworm strains

The *bd* and *bd^f* mutant strains and the wild-type Dazao and N4 strains were obtained from the

bank of genetic resources of Southwest University. Silkworms were reared on mulberry leaves or

artificial diets at 25°C and 73% relative humidity in the dark for 12 hours and light for 12 hours.

Positional cloning of the *bd* locus

For mapping of the *bd* locus, F1 heterozygous individuals were obtained from a cross between a

bd^f strain and a Dazao strain. Then, an F1 female was crossed with a *bd^f* male (BC1F), and an F1

male was backcrossed with a *bd^f* female (BC1M). A total of 1162 BC1M individuals were used

for recombination analysis. Genomic DNA was extracted from the parents (Dazao, *bd^f* and F1)

and each BC1 individual using the phenol chloroform extraction method. Available DNA

molecular markers were identified through polymorphism screening and linkage analysis. The

primers used for mapping are listed in Table S10.

Phylogenetic analysis

To determine whether *Bm-mamo* orthologs were present in other species, the BlastX program of

the National Center for Biotechnology Information (NCBI)

(http://www.ncbi.nlm.nih.gov/BLAST/) was used. The *Bm-mamo-L* and *Bm-mamo-S* sequences

were subjected to BLAST searches against the nonredundant protein sequence (nr) database.

Sequences with a maximum score and an E-value$\leq 10^{-4}$ were downloaded. The sequences of

multiple species were subjected to multiple sequence alignment of the predicted amino acid

sequences by MUSCLE(*88*). A phylogenetic tree was subsequently constructed using the

neighbor☐joining method with the MEGA7 program (Pearson model). The confidence levels for

various phylogenetic lineages were estimated by bootstrap analysis (2,000 replicates).

Quantitative PCR

Total RNA was isolated from the whole body and integument of the silkworms using TRIzol

reagent (Invitrogen, California, USA), purified by phenol chloroform extraction and then reverse

transcribed with a PrimeScript™ RT Reagent Kit (TAKARA, Dalian, China) according to the

manufacturers' protocol.

qPCR was performed using a CFX96™ Real-Time PCR Detection System (Bio-Rad, Hercules,

CA) with a qPCR system and an iTaq Universal SYBR Green Supermix System (Bio-Rad). The

cycling parameters were as follows: 95°C for 3 min, followed by 40 cycles of 95°C for 10 s and

annealing for 30 s. The primers used for the target genes are listed in Table S10. The expression

levels of the genes in each sample were determined with biological replicates, and each sample

was analyzed in triplicate. The gene expression levels were normalized against the expression

levels of the ribosomal protein L3 (RpL3). The relative expression levels were analyzed using

the classical $R=2^{-\Delta\Delta Ct}$ method.

siRNA for gene knockdown

siRNAs for *Bm-mamo* were designed with the siDirect program (http://sidirect2.rnai.jp). The

target siRNAs and negative controls were synthesized by Tsingke Biotechnology Company

Limited. The siRNA (5 μl, 1 μl/μg) was injected from the abdominal spiracle into the

hemolymph at the fourth instar (Day 3) larval stage. Immediately after injection, phosphate-

buffered saline (pH 7.3) droplets were placed nearby, and a 20-voltage pulse for one second and

pause for one second were applied 3 times. The phenotype was observed for fifth-instar larvae.

The left and right epidermis were separately dissected from the injected larvae, after which RNA

was extracted. Then, cDNA was synthesized, and the expression level of the gene was detected

via qPCR.

sgRNA synthesis and RNP complex assembly

CRISPRdirect (http://crispr.dbcls.jp/doc/) online software was used to screen appropriate single

guide RNA (sgRNA) target sequences. The gRNAs were synthetized by the Beijing Genomics

Institute. The sgRNA templates were transcribed using T7 polymerase with RiboMAX™ Large-

Scale RNA Production Systems (Promega, Beijing, China) according to the manufacturer's

instructions. The RNA transcripts were purified using 3 M sodium acetate (pH 5.2) and

anhydrous ethanol (1:30) precipitation, washed with 75% ethanol, and eluted in nuclease-free

water. All injection mixes contained 300 ng/µL Cas9 nuclease (Invitrogen, California, USA) and

300 ng/µL purified sgRNA. Before injection, mixtures of Cas9 nuclease and gRNA were

incubated for 15 min at 37°C to reconstitute active ribonucleoproteins (RNPs)(*89*).

Microinjection of embryos

For embryo microinjection, microcapillary needles were manufactured using a PC-10 model

micropipette puller (Narishige, Tokyo, Japan). Microinjection was performed using an Eppendorf

TransferMan NK 2 and a FemtoJet 4i system (Eppendorf, Hamburg, Germany). The eggs used

for microinjection were generated by mating female and male wild-type Dazao moths. Within 4

hours of culture, the eggs were allowed to adhere to a clean glass slide. CRISPR/Cas9-messenger

RNP mixtures with volumes of approximately 1 nL were injected into the middle of the eggs,

and the wound was sealed with glue. All the injected embryos were allowed to develop in a

climate chamber at 25°C and 80% humidity(*90*).

Comparative genomics

The reference genome of Dazao was downloaded from Silkbase (https://silkbase.ab.a.u-

tokyo.ac.jp/cgi-bin/index.cgi). The bd^f genome was obtained from the silkworm pangenome

project(*34*).

The short reads of the bd^f strains were mapped to the silkworm reference genome by BWA55

v0.7.17 mem with default parameters. The SAMtools56 v1.11 and Picard v2.23.5

(https://broadinstitute.github.io/picard/) programs were used to filter the unmapped and

duplicated reads. A GVCF file of the samples was obtained using GATK57 v4.1.8.1

HaplotypeCaller with the parameter -ERC = GVCF. The VCF files of insertions/deletions (indels)

and single-nucleotide polymorphisms (SNPs) were used for further analysis via eGPS software.

Downstream target gene screening

Online software (http://zf.princeton.edu) was used to predict the DNA-binding site for Cys2His2

zinc finger proteins. The confident ZF domains with scores higher than 17.7 were chosen. RF

regression on the B1H model was used to predict the DNA-binding sites. Then, the sequence

logo and position weight matrices (PWMs) of the DNA-binding sites were obtained. The

sequences 2 kb upstream and downstream of the predicted genes were extracted by a Perl script.

The FIMO package of the MEME suite was used to search for binding sites in the silkworm

genome.

Analysis of EMSA

Primers with binding sites and flanking sequences were designed according to the FIMO results.

A biotin label was added to one end of the upstream primer (probe), and the downstream primer

was used as a reference. Primers with the same sequence as the labeled probes were used as

competitive probes. The EMSA was conducted with an EMSA reagent kit (Beyotime, Shanghai,

China) according to the manufacturers' protocol.

References

1.   G. A. Sword, S. J. Simpson, O. T. M. El Hadi, H. Wilps, Density-dependent aposematism

in the desert locust. *P Roy Soc B-Biol Sci* **267**, 63-68 (2000).

2.   A. I. Barnes, M. T. Siva-Jothy, Density-dependent prophylaxis in the mealworm beetle

Tenebrio molitor L. (Coleoptera: Tenebrionidae): cuticular melanization is an indicator of

investment in immunity. *P Roy Soc B-Biol Sci* **267**, 177-182 (2000).

3.   N. F. Hadley, A. Savill, T. D. Schultz, Coloration and its thermal consequences in the

New Zealand tiger beetle *Neocicindela perhispida*. *J Therm Biol* **17**, 55-61 (1992).

4.   Y. G. Hu, Y. H. Shen, Z. Zhang, G. Q. Shi, Melanin and urate act to prevent ultraviolet

damage in the integument of the silkworm, *Bombyx mori*. *Arch Insect Biochem* **83**, 41-55

(2013).

5.   M. Stevens, G. D. Ruxton, Linking the evolution and form of warning coloration in

nature. *P Roy Soc B-Biol Sci* **279**, 417-426 (2012).

6.   K. K. Dasmahapatra *et al.*, Butterfly genome reveals promiscuous exchange of mimicry

adaptations among species. *Nature* **487**, 94-98 (2012).

7.   N. Gaitonde, J. Joshi, K. Kunte, Evolution of ontogenic change in color defenses of

swallowtail butterflies. *Ecol Evol* **8**, 9751-9763 (2018).

8.   B. S. Tullberg, S. Merilaita, C. Wiklund, Aposematism and crypsis combined as a result

of distance dependence: functional versatility of the colour pattern in the swallowtail

butterfly larva. *P Roy Soc B-Biol Sci* **272**, 1315-1321 (2005).

9.   P. J. Wittkopp, B. L. Williams, J. E. Selegue, S. B. Carroll, *Drosophila* pigmentation

evolution: divergent genotypes underlying convergent phenotypes. *P Natl Acad Sci USA*

100, 1808-1813 (2003).

10.   R. Futahashi, Y. Banno, H. Fujiwara, Caterpillar color patterns are determined by a two-

phase melanin gene prepatterning process: new evidence from *tan* and *laccase2*. *Evol*

Dev **12**, 157-167 (2010).

11.   Y. Matsuoka, A. Monteiro, Melanin pathway genes regulate color and morphology of

butterfly wing scales. *Cell Rep* **24**, 56-65 (2018).

12.   C. Finet *et al.*, Multi-scale dissection of wing transparency in the clearwing butterfly. *J R*

Soc Interface **20**, (2023).

13.   A. Gürses, M. Açıkyıldız, K. Güneş, M. S. Gürses, in *Dyes and Pigments*. (Springer

International Publishing, 2016), pp. 13-29.

14.   V. J. Lloyd, N. J. Nadeau, The evolution of structural colour in butterflies. *Curr Opin*

Genet Dev **69**, 28-34 (2021).

15.   A. E. Seago, P. Brady, J. P. Vigneron, T. D. Schultz, Gold bugs and beyond: a review of

iridescence and structural colour mechanisms in beetles (Coleoptera). *J R Soc Interface* **6**,

S165-S184 (2009).

16.   Y. Oba *et al.*, Resurrecting the ancient glow of the fireflies. *Sci Adv* **6**, (2020).

17.   A. J. Syed, J. C. Anderson, Applications of bioluminescence in biotechnology and beyond.

Chem Soc Rev **50**, 5668-5705 (2021).

18.   X. L. Tong, L. Qiao, J. W. Luo, X. Ding, S. Y. Wu, The evolution and genetics of

lepidopteran egg and caterpillar coloration. *Curr Opin Genet Dev* **69**, 140-146 (2021).

19.   F. Y. Dai *et al.*, Mutations of an arylalkylamine-*N*-acetyltransferase, *Bm-iAANAT*, are

responsible for silkworm melanism mutant. *J Biol Chem* **285**, 19553-19560 (2010).

20.   P. J. Wittkopp, J. R. True, S. B. Carroll, Reciprocal functions of the *Drosophila* yellow

and ebony proteins in the development and evolution of pigment patterns. *Development*

129, 1849-1858 (2002).

21.   J. Q. Liu *et al.*, Lepidopteran wing scales contain abundant cross-linked film-forming

histidine-rich cuticular proteins. *Commun Biol* **4**, (2021).

22.   Z. W. Yan *et al.*, A blueprint of microstructures and stage-specific transcriptome

dynamics of cuticle formation in *Bombyx mori*. *Int J Mol Sci* **23**, (2022).

23.   R. Futahashi *et al.*, Genome-wide identification of cuticular protein genes in the

silkworm, *Bombyx mori*. *Insect Biochem Molec* **38**, 1138-1146 (2008).

24.   J. Yamaguchi *et al.*, Periodic Wnt1 expression in response to ecdysteroid generates twin-

spot markings on caterpillars. *Nat Commun* **4**, (2013).

25.   S. Yoda *et al.*, The transcription factor Apontic-like controls diverse colouration pattern in

caterpillars. *Nat Commun* **5**, (2014).

26.   H. Jin, T. Seki, J. Yamaguchi, H. Fujiwara, Prepatterning of *Papilio xuthus* caterpillar

camouflage is controlled by three homeobox genes: *clawless*, *abdominal-A*, and

Abdominal-B. *Sci Adv* **5**, (2019).

27.   W. A. Rogers *et al.*, A survey of the trans-regulatory landscape for *Drosophila*

melanogaster abdominal pigmentation. *Dev Biol* **385**, 417-432 (2014).

28.   S. Jeong, A. Rokas, S. B. Carroll, Regulation of body pigmentation by the Abdominal-B

Hox protein and its gain and loss in *Drosophila* evolution. *Cell* **125**, 1387-1399 (2006).

29.   H. D. Dufour, S. Koshikawa, C. Finet, Temporal flexibility of gene regulatory network

underlies a novel wing pattern in flies. *P Natl Acad Sci USA* **117**, 11589-11596 (2020).

30.   A. Prakash, C. Finet, T. Das Banerjee, V. Saranathan, A. Monteiro, *Antennapedia* and

optix regulate metallic silver wing scale development and cell shape in *Bicyclus anynana*

butterflies. *Cell Rep* **40**, (2022).

31.   R. D. Reed *et al.*, Drives the repeated convergent evolution of butterfly wing pattern

mimicry. *Science* **333**, 1137-1141 (2011).

32.   A. Kopp, I. Duncan, S. B. Carroll, Genetic control and evolution of sexually dimorphic

characters in *Drosophila*. *Nature* **408**, 553-559 (2000).

33.   L. Arnoult *et al.*, Emergence and diversification of fly pigmentation through evolution of

a gene regulatory module. *Science* **339**, 1423-1426 (2013).

34.   X. L. Tong *et al.*, High-resolution silkworm pan-genome provides genetic insights into

artificial selection and ecological adaptation. *Nat Commun* **13**, (2022).

35.   L. Cheng., D. Fangyin., X. Zhonghuai., Studies on the mutant systems of the *Bombyx*

mori gene bank. *Scientia agricultura sinica* **36**, 968-975 (2003).

36.   Y. Kawaguchi, T. Kusakabe, J. M. Lee, K. Koga, Manifestation of the *dilute black* (*bd*)

mutation and constitution of the *bd* Locus in *Bombyx mori*. *J Fac Agr Kyushu U* **52**, 355-

359 (2007).

37.   J. Duan *et al.*, SilkDB v2.0: a platform for silkworm (*Bombyx mori*) genome biology.

Nucleic Acids Res **38**, D453-D456 (2010).

38.   E. W. Sayers *et al.*, GenBank. *Nucleic Acids Res* **50**, D161-d164 (2022).

39.   M. Kawamoto, T. Kiuchi, S. Katsuma, SilkBase: an integrated transcriptomic and

genomic database for *Bombyx mori* and related species. *Database-Oxford* **2022**, (2022).

40.   V. J. Bardwell, R. Treisman, The POZ domain: a conserved protein-protein interaction

motif. *Gene Dev* **8**, 1664-1677 (1994).

41.   K. D. Huynh, V. J. Bardwell, The BCL-6 POZ domain and other POZ domains interact

with the co-repressors N-CoR and SMRT. *Oncogene* **17**, 2473-2484 (1998).

42.   P. Staller *et al.*, Repression of p15INK4b expression by Myc through association with

Miz-1. *Nat Cell Biol* **3**, 392-399 (2001).

43.   M. Mukai *et al.*, MAMO, a maternal BTB/POZ-Zn-finger protein enriched in germline

progenitors is required for the production of functional eggs in *Drosophila*. *Mech*

Develop **124**, 570-583 (2007).

44.   T. Ando, H. Fujiwara, Electroporation-mediated somatic transgenesis for rapid functional

analysis in insects. *Development* **140**, 454-458 (2013).

45.   C. O. Pabo, E. Peisach, R. A. Grant, Design and selection of novel Cys2His2 zinc finger

proteins. *Annu Rev Biochem* **70**, 313-340 (2001).

46.   S. A. Wolfe, L. Nekludova, C. O. Pabo, DNA recognition by Cys2His2 zinc finger

proteins. *Annu Rev Bioph Biom* **29**, 183-212 (2000).

47.   A. V. Persikov, M. Singh, De novo prediction of DNA-binding specificities for Cys2His2

zinc finger proteins. *Nucleic Acids Res* **42**, 97-108 (2014).

48.   S. Hira *et al.*, Binding of *Drosophila* maternal Mamo protein to chromatin and specific

DNA sequences. *Biochem Bioph Res Co* **438**, 156-160 (2013).

49.   P. J. Wittkopp, G. Kalay, *Cis*-regulatory elements: molecular mechanisms and

evolutionary processes underlying divergence. *Nat Rev Genet* **13**, 59-69 (2012).

50.   S. Preissl, K. J. Gaulton, B. Ren, Characterizing *cis*-regulatory elements using single-cell

epigenomics. *Nat Rev Genet*, (2022).

51.   E. Z. Kvon *et al.*, Genome-scale functional characterization of *Drosophila* developmental

enhancers *in vivo*. *Nature* **512**, 91-+ (2014).

52.   S. Y. Wu *et al.*, Comparative analysis of the integument transcriptomes of the *black dilute*

mutant and the wild-type silkworm *Bombyx mori*. *Sci Rep-Uk* **6**, (2016).

53.   G. Xiong *et al.*, Body shape and coloration of silkworm larvae are influenced by a novel

cuticular protein. *Genetics* **207**, 1053-1066 (2017).

54.   P. J. Wittkopp, S. B. Carroll, A. Kopp, Evolution in black and white: genetic control of

pigment patterns in *Drosophila*. *Trends Genet* **19**, 495-504 (2003).

55.   J. M. Gibert, Phenotypic plasticity in insects. *Biologie aujourd'hui* **214**, 33-44 (2020).

56.   R. Futahashi *et al.*, *yellow* and *ebony* are the responsible genes for the larval color

mutants of the silkworm *Bombyx mori*. *Genetics* **180**, 1995-2005 (2008).

57.   M. Y. Noh, S. Muthukrishnan, K. J. Kramer, Y. Arakane, Cuticle formation and

pigmentation in beetles. *Curr Opin Insect Sci* **17**, 1-9 (2016).

58.   N. N. Wang *et al.*, Abnormal overexpression of *SoxD* enhances melanin synthesis in the

Ursa mutant of *Bombyx mori*. *Insect Biochem Molec* **149**, (2022).

59.   K. J. Kramer *et al.*, Oxidative conjugation of catechols with proteins in insect skeletal

systems. *Tetrahedron* **57**, 385-392 (2001).

60.   M. Y. Noh, S. Muthukrishnan, K. J. Kramer, Y. Arakane, *Tribolium castaneum* RR-1

cuticular protein TcCPR4 is required for formation of pore canals in rigid cuticle. *Plos*

Genet **11**, (2015).

61.   M. Y. Noh *et al.*, Two major cuticular proteins are required for assembly of horizontal

laminae and vertical pore canals in rigid cuticle of *Tribolium castaneum. Insect Biochem*

Molec **53**, 22-29 (2014).

62.   S. Mun *et al.*, Cuticular protein with a low complexity sequence becomes cross-linked

during insect cuticle sclerotization and is required for the adult molt. *Sci Rep-Uk* **5**,

(2015).

63.   A. Prakash *et al.*, Nanoscale cuticle density variations correlate with pigmentation and

color in butterfly wing scales. *arXiv*, 2305.16628 (2023).

64.   R. Futahashi, H. Shirataki, T. Narita, K. Mita, H. Fujiwara, Comprehensive microarray-

based analysis for stage-specific larval camouflage pattern-associated genes in the

swallowtail butterfly, *Papilio xuthus. Bmc Biol* **10**, (2012).

65.   M. Y. Noh, S. Muthukrishnan, K. J. Kramer, Y. Arakane, Development and ultrastructure

of the rigid dorsal and flexible ventral cuticles of the elytron of the red flour beetle,

Tribolium castaneum. Insect Biochem Molec **91**, 21-33 (2017).

66.   R. Tajiri, N. Ogawa, H. Fujiwara, T. Kojima, Mechanical control of whole body shape by

a single cuticular protein Obstructor-E in *Drosophila melanogaster*. *Plos Genet* **13**,

(2017).

67.   X. M. Zhao *et al.*, The wing-specific cuticular protein LmACP7 is essential for normal

714   wing morphogenesis in the migratory locust. *Insect Biochem Molec* **112**, (2019).

715   68.   J. H. Willis, Structural cuticular proteins from arthropods: Annotation, nomenclature, and

716   sequence characteristics in the genomics era. *Insect Biochem Molec* **40**, 189-204 (2010).

717   69.   J. E. Rebers, L. M. Riddiford, Structure and expression of a *Manduca sexta* larval cuticle

718   gene homologous to *Drosophila* cuticle genes. *J Mol Biol* **203**, 411-423 (1988).

719   70.   M. V. Karouzou *et al.*, *Drosophila* cuticular proteins with the R&R Consensus:

720   Annotation and classification with a new tool for discriminating RR-1 and RR-2

721   sequences. *Insect Biochem Molec* **37**, 754-760 (2007).

722   71.   E. H. Chen, Q. L. Hou, Identification and expression analysis of cuticular protein genes

723   in the diamondback moth, *Plutella xylostella* (Lepidoptera: Plutellidae). *Pestic Biochem

724   Phys* **178**, (2021).

725   72.   X. Fu *et al.*, Genome-Wide Identification and Transcriptome-Based Expression Profile of

726   Cuticular Protein Genes in *Antheraea pernyi*. *Int J Mol Sci* **24**, (2023).

727   73.   C. H. Yang *et al.*, Identification, expression pattern, and feature analysis of cuticular

728   protein genes in the pine moth *Dendrolimus punctatus* (Lepidoptera: Lasiocampidae).

729   *Insect Biochem Molec* **83**, 94-106 (2017).

730   74.   X. Guan, B. W. Middlebrooks, S. Alexander, S. A. Wasserman, Mutation of TweedleD, a

731   member of an unconventional cuticle protein family, alters body shape in *Drosophila*. *P

732   Natl Acad Sci USA* **103**, 16794-16799 (2006).

733   75.   T. Togawa, W. A. Dunn, A. C. Emmons, J. H. Willis, CPF and CPFL, two related gene

734  families encoding cuticular proteins of *Anopheles gambiae* and other insects. *Insect*

735  *Biochem Molec* **37**, 675-688 (2007).

736 76. P. L. Pan *et al.*, A comprehensive omics analysis and functional survey of cuticular

737  proteins in the brown planthopper. *P Natl Acad Sci USA* **115**, 5175-5180 (2018).

738 77. K. Ito *et al.*, Positional cloning of a gene responsible for the *cts* mutation of the silkworm,

739  *Bombyx mori*. *Genome* **55**, 493-504 (2012).

740 78. M. Osanai-Futahashi *et al.*, Identification of the *Bombyx red egg* gene reveals

741  involvement of a novel transporter family gene in late steps of the insect ommochrome

742  biosynthesis pathway. *J Biol Chem* **287**, 17706-17714 (2012).

743 79. J. W. Luo *et al.*, Molecular basis of the silkworm mutant re^l causing red egg color and

744  embryonic death. *Insect Sci* **28**, 1290-1299 (2021).

745 80. J. J. Hanly *et al.*, Genetics of yellow-orange color variation in a pair of sympatric sulphur

746  butterflies. *Cell Rep* **42**, (2023).

747 81. G. Kalay, J. Lachowiec, U. Rosas, M. R. Dome, P. Wittkopp, Redundant and cryptic

748  enhancer activities of the *Drosophila yellow* gene. *Genetics* **212**, 343-360 (2019).

749 82. Y. Q. Xin *et al.*, Enhancer evolutionary co-option through shared chromatin accessibility

750  input. *P Natl Acad Sci USA* **117**, 20636-20644 (2020).

751 83. A. Jolma, J. Taipale, Methods for analysis of transcription factor DNA-binding specificity

752  in vitro. *Sub-cellular biochemistry* **52**, 155-173 (2011).

753 84. Y. F. Chan *et al.*, Adaptive evolution of pelvic reduction in sticklebacks by recurrent

754  deletion of a *Pitx1* enhancer. *Science* **327**, 302-305 (2010).

755   85.   K. P. Lu *et al.*, SilkMeta: a comprehensive platform for sharing and exploiting pan-

756   genomic and multi-omic silkworm data. *Nucleic Acids Res*, (2023).

757   86.   J. G. Monroe *et al.*, Mutation bias reflects natural selection in *Arabidopsis thaliana*.

758   *Nature* **602**, 101-105 (2022).

759   87.   E. J. Belfield *et al.*, Thermal stress accelerates *Arabidopsis thaliana* mutation rate.

760   *Genome Res* **31**, 40-50 (2021).

761   88.   R. C. Edgar, MUSCLE: multiple sequence alignment with high accuracy and high

762   throughput. *Nucleic Acids Res* **32**, 1792-1797 (2004).

763   89.   Y. L. Zou *et al.*, Bmmp influences wing morphology by regulating anterior-posterior and

764   proximal-distal axes development. *Insect Sci* **29**, 1569-1582 (2022).

765   90.   D. P. Long *et al.*, Genetic hybridization of highly active exogenous functional proteins

766   into silk-based materials using "light-clothing" strategy. *Matter-Us* **4**, 2039-2058 (2021).

Acknowledgments

Funding:

This work was supported by the National Natural Science Foundation of China (No. 32002230,

No. 32330102, No. U20A2058), the Fundamental Research Funds for the Central Universities in

China (No. SWU120024), National key research and development program (Project

No.2023YFD1600901, 2023YFF1103801), Natural Science Foundation of Chongqing, China

(No. cstc2021jcyj-cxtt0005) and High-level Talents Program of Southwest University (no.

SWURC2021001).

Author contributions:

FY Dai conceived and designed the experiments. SY Wu, CX Peng, JW Luo, KP Lu and CH

Zhang performed the study. SY Wu, XL Tong, KP Lu, CL Li, X Ding, YR Lu, XH Duan, D Tan

and H Hu analyzed the data. SY Wu wrote the paper. XL Tong and FY Dai edited and revised the

manuscript.

Competing interests:

The authors declare that they have no competing interests.

Data and material availability:

The authors confirm that the data supporting the findings of this study are available within the

article and its supplementary materials.

Figures and Tables

796 **Fig. 1. Phenotypes of *bd*, *bd*^{f}, and wild-type Dazao larvae.**

797 The epidermis of *bd* is dark gray, and the epidermis of *bd*^{f} is light gray at the 5th instar (day 3) of

798 silkworm larvae. The bar indicates 1 centimeter.

Fig. 2. Positional cloning of the *bd* locus.

(A) We used 532 BC1 individuals to map the *bd* locus between PCR markers c and i. The

numbers above the DNA markers indicate recombination events.

(B) A total of 1162 BC1 individuals were used to narrow the *bd* locus to an approximately 400

kb genomic region.

(C) Partial enlarged view of the region responsible for *bd*. This region contains 4 predicted genes,

BGIBMGA012517, *BGIBMGA012518*, *BGIBMGA012519* and *BGIBMGA014089*.

(D) Analysis of nucleotide differences in the region responsible for *bd*. The green block indicates

the deletion of the genome in *bd* mutants. The black vertical lines indicate the SNPs and indels of

bd^f mutants.

Fig. 3. Spatiotemporal expression of *Bm-mamo*.

(A) Temporal expression of *Bm-mamo*. In the molting stage and adult stage, this gene was

significantly upregulated. M: molting stage, W: wandering stage, P: pupal stage, P1: Day 1 of the

pupal stage, A: adult stage, 4th3d indicates the 4th instar on Day 3. 1st to 5th denote the first

instar of larvae to fifth instar of larvae, respectively.

(B) Tissue-specific expression of 4th-instar molting larvae. *Bm-mamo* had relatively high

expression levels in the midgut, head, and epidermis. AMSG: anterior division of silk gland and

middle silk gland, PSG: posterior silk gland.

(C) Detailed analysis of *Bm-mamo* at the 4th larval stage in the epidermis of the Dazao strain.

Bm-mamo expression is upregulated during the molting stage. HCS indicates the head capsule

stage. The "h" indicates the hour, 90 h: at 90 hours of the 4th instar.

(**A**) siRNA was introduced by microinjection followed by electroporation. "+" and "-" indicate

the positive and negative poles of the electrical current, respectively. Scale bars, 1 cm.

(**B**) Partial magnification of the siRNA experimental group and negative control group. Scale

bars, 0.2 cm.

(**C**) and (**D**) Relative expression levels of *Bm-mamo* in the negative control and RNAi groups

were determined by qPCR analysis. The means ± s.d.s. Numbers of samples are shown in the

upper right in each graph. **$P<0.01$, paired Student's t test (NS, not significant).

(E) Statistical analysis of the efficiency of the RNAi.

Fig. 5. *Bm-mamo* knockout

(**A**) Larval phenotype of G3 fourth-instar larvae of Dazao targeted for *Bm-mamo*. Scale bars, 1

cm.

(**B**) Partial magnification of the *Bm-mamo* knockout individual and control. Scale bars, 0.2 cm.

(**C**) After 48 hours of egg laying, the pigmentation of the eggs indicates that those produced by

knockout homozygous females cannot undergo normal pigmentation and development.

(**D**) Genomic structure of *Bm-mamo*. The open reading frame (blue) and untranslated region

(black) are shown. The gRNA 1 and gRNA 2 sequences are shown.

(**E**) Sequences of the *Bm-mamo* knockout individuals. Lines 1 and 2 indicate deletions of 15 and

71 bp, respectively.

Fig. 6. Melanin metabolism pathway and qPCR of related genes.

(**A**) The melanin metabolism pathway. Blue indicates amino acids and catecholamines, red

indicates the names of genes, and black indicates the names of enzymes.

(**B**) Relative expression levels of eight genes in the heterozygous *Bm-mamo* knockout group

(blue) and homozygous *Bm-mamo* knockout group (red) determined via qPCR analysis. The

means ± s.d.s. *P<0.05, paired Student's t test. 4th3d indicates the 4th instar on Day 3, M

indicates molting, and h indicates hours.

Fig. 7. Heatmap of 28 differentially expressed cuticular protein genes.

Differentially expressed cuticular protein genes between homozygous (*mamo⁻/mamo⁻*) and

heterozygous (*mamo⁻/+*) *Bm-mamo* knockout individuals. Red indicates upregulation in

homozygous individuals. Blue indicates downregulation in homozygous individuals. White

indicates no difference in expression level or no investigation. 4th1d indicates 4-instar Day 1, M

indicates molting, and h indicates hours.

Additional information

Supplementary Information is available for this paper.

Dazao   *+/bd*   *bd/bd*   *bd^f/bd^f*

A

B

C

A
RNAi   Control

Left (+)   Right (-)   Left (+)   Right (-)

B
RNAi   Control

Left (+)   Right (-)   Left (+)   Right (-)

C

RNAi (n=5)

Relative gene expression level

Left   Right

**

D

Control (n=3)

Relative gene expression level

ns

Left   Right

E

	Injected individuals	Melanism phenotype	Survival number	Efficiency
RNAi	48	23	48	47.92%
Control	21	0	17	0

A

Dazao   *Bm-mamo* knockout

B

Dazao   *Bm-mamo* knockout

C

Dazao   *Bm-mamo* knockout   *bd* muntant

D

Bm-mamo-L

Bm-mamo-S

1 kb

gRNA-1   gRNA-2

5'-ATGACGACTTTAAACGCAGCCGG-3' · · · 5'-CAAAGCCAAGACGCTAGAGAAGG-3'

3'-TACTGCTGAAATTTGCGTCGGCC-5' · · · 5'-GTTTCGGTTCTGCGATCTCTTCC-3'

E

Dazao   ATGACGACTTTAAACGCAGCCGGTATCGAAACAAGAAGTGTTTCGGAGCAAAGAGAGCAGGGCCAAAGCCAAGACGCTAGAGAAGG

line-1   ATGACGACTTTAAAC------------ACAAGAAGTGTTTCGGAGCAAAGAGAGCAGGGCCAAAGCCAAGACGCTAGAGAAGG

Deletion 15bp

line-2   ATGACGACTTTA---AGG

Deletion 71bp

A

B

TH

DDC

aaNATs

yellow

laccase2

tan

black

ebony

mamo⁻/+
mamo⁻/mamo⁻