The BTB-ZF gene *Bm-mamo* regulates pigmentation in silkworm caterpillars

Songyuan Wu, Xiaoling Tong, Chenxing Peng, Jiangwen Luo, Chenghao Zhang, Kunpeng Lu,

Chunlin Li, Xin Ding, Xiaohui Duan, Yaru Lu, Hai Hu, DuanTan, Fangyin Dai[*]

Affiliations

State Key Laboratory of Resource Insects, Key Laboratory of Sericultural Biology and Genetic

Breeding, Ministry of Agriculture and Rural Affairs, College of Sericulture, Textile and Biomass

Sciences, Southwest University, Chongqing 400715, China

Songyuan Wu fly_jungle@163.com

Xiaoling Tong xltong@swu.edu.cn

Chenxing Peng mxtaste@sina.com

Jiangwen Luo luo0510@email.swu.edu.cn

Chenghao Zhang hangsdo0452@163.com

Kunpeng Lu lukunpeng@swu.edu.cn

Chunlin Li lclin0522@swu.edu.cn

Xin Ding dingxin2019@swu.edu.cn

Xiaohui Duan duanxiaohui@email.swu.edu.cn

Yaru Lu 17723940349@163.com

Hai Hu huhaiswu@163.com

DuanTan tanzeduan@163.com

Fangyin Dai fydai@swu.edu.cn

*Corresponding author

Fangyin Dai

fydai@swu.edu.cn

86-023-68250793

Southwest University, No. 2 Tiansheng Road, Beibei District, Chongqing 400715, China

Abstract

The color pattern of insects is one of the most dazzling adaptive evolutionary phenotypes. However, the molecular regulation of this color pattern is not fully understood. In this study, we found that the transcription factor Bm-mamo is responsible for *black dilute* (*bd*) allele mutations in the silkworm. Bm-mamo belongs to the BTB zinc finger family and is orthologous to mamo in *Drosophila melanogaster*. This gene has a conserved function in gamete production in *Drosophila* and silkworms and evolved a pleiotropic function in the regulation of color patterns in caterpillars. We found that *Bm-mamo* can comprehensively regulate the expression of related pigment synthesis and cuticular protein genes to form color patterns. This indicates that insects have a genetic basis for coordinate regulation of the structure and shape of the cuticle, as well as color patterns. This genetic basis provides the possibility for constructing the complex appearances of some insects. This study provides new insight into the regulation of color patterns.

Teaser

The color patterns of insects are highly exquisite and significantly divergent. The metabolism of pigments is the material basis for insect coloration. However, the cuticle of insects plays an important role as a scaffold for carrying pigment particles. Cuticular proteins are some of the main components of the cuticle. This study showed that a BTB-ZF family transcription factor protein, Bm-mamo, can comprehensively regulate melanin synthesis and the expression of multiple cuticular protein genes. Hence, insects have a genetic basis for integrated control of cuticle and color pattern construction, which enables them to produce complex appearances.

Introduction

Insects often display stunning colors, and the patterns of these colors have been shown to be

involved in behavior(*1*); immunity(*2*); thermoregulation(*3*); UV protection(*4*); and especially

visual antagonism of predators via aposematism(*5*), mimicry(*6*), cryptic color patterns(*7*) or

some combination of the above (*8*). In addition, the color patterns are divergent and convergent

among populations(*9*). Due to these striking visual features and highly active adaptive

evolutionary phenotypes, the genetic basis and evolutionary mechanism of color patterns have

been a topic of interest for a long time.

Insect coloration can be pigmentary, structural, or bioluminescent. Pigments are mainly

synthesized by the insects themselves and form solid particles that are deposited in the cuticle of

the body surface and the scales of the wings(*10, 11*). Interestingly, recent studies have found that

bile pigments and carotenoid pigments synthesized through biological synthesis are incorporated

into body fluids and filled in the wing membranes of two butterflies (*Siproeta stelenes* and

Philaethria diatonica) via hemolymph circulation, providing color in the form of liquid

pigments(*12*). The pigments form colors by selective absorption and/or scattering of light

depending on their physical properties(*13*). However, structural color refers to colors, such as

metallic colors and iridescence, generated by optical interference and grating diffraction of the

microstructure/nanostructure of the body surface or appendages (such as scales)(*14, 15*). Pigment

color and structural color are widely distributed in insects and can only be observed by the naked

eye in illuminated environments. However, some insects, such as fireflies, exhibit colors (green

to orange) in the dark due to bioluminescence(*16*). Bioluminescence occurs when luciferase

catalyzes the oxidation of small molecules of luciferin(*17*). In conclusion, the color patterns of

insects have evolved to be highly sophisticated and are closely related to their living

environments. For example, cryptic color can deceive animals via high similarity to the

surrounding environment. However, the molecular mechanism by which insects form precise

color patterns to match their living environment is still unknown.

Recent research has identified the metabolic pathways of related pigments, such as melanins,

pterins, and ommochromes, in Lepidoptera(*18*). A deficiency of enzymes in the pigment

metabolism pathway can lead to changes in color patterns(*19, 20*). In addition to pigment

synthesis, the microstructure/nanostructure of the body surface and wing scales are important

factors that influence body color patterns. The body surface and wing scales of Lepidoptera are

mainly composed of cuticular proteins (CPs) (*21, 22*). There are multiple genes encoding CPs in

Lepidopteran genomes. For example, in *Bombyx mori,* over 220 genes encode CPs(*23*). However,

there are still many unknowns regarding the function of CPs and the molecular mechanisms

underlying their fine-scale localization in the cuticle.

In addition, some pleiotropic factors, such as *wnt1(24)*, *Apontic-like(25)*, *clawless(26)*,

abdominal-A(27), *abdominal-B(28)*, *engrailed(29)*, *antennapedia(30)*, *optix(31)*, *bric* à *brac*

(*bab*)(*32*), and *Distal-less* (*Dll*)(*33*), play an important role in the regulation of color patterns.

The molecular mechanism by which these factors participate in the regulation of body color

patterns needs further study. In addition, there are many undiscovered factors in the gene

regulatory network (GRN) involved in the regulation of insect color patterns. The identification

of new color pattern regulatory genes and the study of their molecular mechanisms would be

helpful for further understanding color patterns.

Silkworms (*B. mori*) have been completely domesticated for more than 5000 years and are

famous for their silk fiber production(*34*). Because of this long-term domestication, artificial

selection and genetic research, its genetic background is well resolved, and approximately 200

strains of mutants with variable color patterns have been preserved(*35*), which provide a good

resource for color pattern research. The *black dilute* (*bd*) mutant, which exhibits recessive

Mendelian inheritance, has a dark gray larval body color, and the female is sterile. bd^f (*black*

dilute fertile) is one of the alleles, leading to a lighter body color than that of *bd* and fertile

females (Fig. 1). The two mutations were mapped to a single locus at 22.9 centimorgans (cM) of

linkage group 9(*36*). No pigmentation-related genes have been reported at this locus. Thus,

research may reveal a new color pattern-related gene, which stimulated our interest.

Results

Candidate gene of the *bd* allele

To identify the genomic region responsible for the *bd* alleles, positional cloning of the *bd* locus

was performed. Due to the female infertility of the *bd* mutant and the fertility of females with the

bd^f allele, we used bd^f and the wild-type Dazao strain as the parents for mapping analysis.

The 1162 back-crossed filial 1st (BC1M) generation individuals from bd^f and Dazao were used

for fine mapping with molecular markers (Fig. 2). A genomic region of approximately 390

kilobases (kb) was responsible for the *bd* phenotype (Table S1). In the SilkDB database(*37*), this

region included five predicted genes (*BGIBMGA012516*, *BGIBMGA012517*, *BGIBMGA012518*,

BGIBMGA012519 and *BGIBMGA014089*). In addition, we analyzed the predictive genes for this

genomic region from the GenBank(*38*) and SilkBase(*39*) databases (Fig. S1). The number of

predicted genes varied among different databases. We performed sequence alignment analysis of

the predicted genes in the three databases to determine their correspondence. Then, Real-time

Quantitative polymerase chain reaction (qPCR) was performed, which showed that

BGIBMGA012517 and *BGIBMGA012518* were significantly downregulated on Day 3 of the fifth

instar of larvae in the *bd* phenotype individuals, while there was no difference in the expression

levels of other genes (Fig. S1). These two genes were predicted as a single locus

(*LOC101738295*) in GenBank. To determine the gene structure of *BGIBMGA012517* and

BGIBMGA012518, we used forward primers for the *BGIBMGA012517* gene and reverse primers

for the *BGIBMGA012518* gene to amplify cDNA from the wild-type Dazao strain. By gene

cloning, the predicted genes *BGIBMGA012517* and *BGIBMGA012518* were proven to be one

gene. For the convenience of description, we have temporarily called it the *12517/8* gene.

The *12517/8* gene produces two transcripts; the open reading frame (ORF) of the long transcript

is 2397 bp, and the ORF of the short transcript is 1824 bp, which shares the same 5'-terminus as

the wild-type Dazao strain (Fig. S2). The *12517/8* gene showed significantly lower expression in

the bd^f mutant (Fig. S3), and multiple variations were found in the nearby region of this gene by

comparative genomic analysis between bd^f and Dazao (Table S2). In addition, *12517/8* was

completely silenced due to the deletion of DNA fragments (approximately 168 kb) from the first

upstream intron and an insertion of 3560 bp in the *bd* mutant (Fig. S4).

To predict the function of the *12517*/8 gene, we performed a BLAST search using its full-length

sequence and found a transcription factor, the *maternal gene required for meiosis* (*mamo*) in *D.*

melanogaster, that had a high sequence similarity to that of the *12517/8* gene (Fig. S5).

Therefore, we named the *12517/18* gene *Bm-mamo*; the long transcript was designated as *Bm-*

mamo-L, and the short transcript was designated as *Bm-mamo-S*.

The *mamo* gene belongs to the Broad-complex, Tramtrack and bric à brac/poxvirus zinc finger

protein (BTB-ZF) family. In the BTB-ZF family, the zinc finger serves as a recognition motif for

DNA-specific sequences, and the BTB domain promotes oligomerization and the recruitment of

other regulatory factors(*40*). Most of these factors are typically transcription repressors, such as

nuclear receptor corepressor (N-CoR) and silencing mediator for retinoid and thyroid hormone

receptor (SMRT)(*41*), but some are activators, such as p300(*42*). Therefore, these features

commonly serve as regulators of gene expression. *mamo* is enriched in embryonic primordial

germ cells (PGCs) in *D. melanogaster*. Individuals deficient in *mamo* are able to undergo

oogenesis but fail to execute meiosis properly, leading to female infertility in *D.*

melanogaster(43). *Bm-mamo* was identified as an important candidate gene for further analysis.

Expression pattern analysis of *Bm-mamo*

To analyze the expression profiles of *Bm-mamo*, we performed quantitative PCR. The expression

levels of the *Bm-mamo* gene were investigated in the whole body at different developmental

stages, from the embryonic stage to the adult stage, in the Dazao strain. The gene was highly

expressed in the molting stage of caterpillars, and its expression was upregulated in the later

pupal and adult stages (Fig. 3A). This suggests that the *Bm-mamo* gene responds to the ecdysone

titer and participates in the processes of molting and metamorphosis in silkworms. In the

investigation of tissue-specific expression levels in the 5th-instar 3rd-day larvae of the Dazao

strain, the midgut, head, and epidermis had high expression levels; the trachea, nerves, silk

glands, testis, ovary, muscle, wing disc and Malpighian tubules had moderate expression levels;

and the blood and fat bodies had low expression levels (Fig. 3B). This suggests that *Bm-mamo* is

involved in the developmental regulation of multiple silkworm tissues. Due to the melanism of

the epidermis of the *bd* mutant and the high expression level of the *Bm-mamo* gene in the

epidermis, we measured the expression level of this gene in the epidermis of the 4th to 5th instar

of the Dazao strain. In the epidermis, the *Bm-mamo* gene was upregulated in the molting period,

and the highest expression was observed at the beginning of molting (Fig. 3C).

Functional analyses of *Bm-mamo*

To study the function of the *Bm-mamo* gene, we carried out an RNA interference (RNAi)

experiment. Short interfering RNA (siRNA) was injected into the hemolymph of silkworms, and

an electroporation experiment(*44*) was immediately conducted. We found significant melanin

pigmentation in the epidermis of the newly molted 5th-instar larvae. This indicates that *Bm-*

mamo deficiency can cause melanin pigmentation (Fig. 4). The melanism phenotype of RNAi

individuals was similar to that of *bd* mutants.

In addition, gene knockout was performed. Ribonucleoproteins (RNPs) generated by guide RNA

(gRNA) and recombinant Cas9 protein were injected into 450 silkworm embryos. In the G0

generation, individuals with a mosaic melanization phenotype were found. These melanistic

individuals were raised to moths and then crossed. The homozygous line with gene knockout

was obtained through generations of screening. The gene-edited individuals had a significantly

melanistic body color, and the female moths were sterile (Fig. 5).

These results indicated that the *Bm-mamo* gene negatively regulates melanin pigmentation in

caterpillars and participates in the reproductive regulation of female moths.

Downstream target gene analysis

Bm-mamo belongs to the zinc finger protein family, which specifically recognizes downstream

DNA sequences according to their zinc fingers. We identified homologous genes of *mamo* in

multiple species and conducted a phylogenetic analysis (Fig. S6). Sequence alignment revealed

that the amino acid residues of the zinc finger motif of the mamo-S protein were highly

conserved among 57 species. The insufficient prediction of alternative transcripts of mamo in

different species in GenBank may have resulted in the mamo-L protein being found in only 30

species (Fig. S7). As the mamo protein carries a tandem Cys_2His_2 zinc finger (C2H2-ZF) motif,

it can directly bind to DNA sequences. Previous research has suggested that the ZF-DNA binding

interface can be understood as a "canonical binding model", in which each finger contacts DNA

in an antiparallel manner. The binding sequence of the C2H2-ZF motif is determined by the

amino acid residue sequence of its α-helical component. Considering the first amino acid residue

in the α-helical region of the C2H2-ZF domain as position 1, positions -1, 2, 3, and 6 are key

amino acids in the recognition and binding of DNA. The residues at positions -1, 3, and 6

specifically interact with base 3, base 2, and base 1 of the DNA sense sequence, respectively,

while the residue at position 2 interacts with the complementary DNA strand(*45, 46*). To analyze

the downstream target genes of mamo, we first predicted its DNA binding motifs using online

software (http://zf.princeton.edu) based on the canonical binding model(*47*). In addition, the

DNA-binding sequence of mamo (<u>TGCGT</u>) in *Drosophila*, confirmed by electrophoretic

mobility shift assay (EMSA) experiments(*48*), has a consensus sequence with the predicted

binding site sequence of Bm-mamo-S (G<u>TGCGT</u>GGC), and the predicted sequence was longer.

This indicates that the predicted results of the DNA-binding site have good reliability.

Furthermore, the predicted DNA-binding sites of Bm-mamo-L and Bm-mamo-S were highly

consistent with those of mamo orthologs in different species (Fig. S8). This suggests that the

protein may regulate similar target genes between species.

C2H2-ZF transcription factors function by recognizing and binding to *cis*-regulatory sequences

in the genome, which harbor *cis*-regulatory elements (CREs)(*49*). CREs are broadly classified as

promoters, enhancers, silencers and insulators(*50*). CREs are often located near their target genes,

such as enhancers, which are typically located upstream (5') or downstream (3') of the gene they

regulate, or in introns, but approximately 12% of CREs are located far from their target gene(*51*).

Therefore, we first investigated the genome range 2 kb upstream and downstream of the

predicted genes in silkworms.

The predicted position weight matrices (PWMs) of the recognized sequences of Bm-mamo and

the Find Individual Motif Occurrences (FIMO) software of MEME were used to perform

silkworm whole-genome scanning for possible downstream target genes. The genomic regions 2

kb upstream and downstream of the 14,623 predicted genes in silkworms were investigated. A

total of 10,622 genes contained the recognition sites of the Bm-mamo protein in the silkworm

genome (Fig. S9).

Moreover, we compared transcriptome data of the integument tissue between homozygotes and

heterozygotes of the *bd* mutant at the 4[th] instar/beginning molting stage(*52*). In the integument

tissue, 10,072 genes (~69% of the total predicted genes of silkworm) were expressed in

heterozygotes, and 9,853 genes (~67% of the total predicted genes) were expressed in

homozygotes of the *bd* mutant. In addition, there were 191 genes with significant expression

differences between homozygotes (*bd*/*bd*) and heterozygotes (+/*bd*) by comparative

transcriptome analysis (Table S3)(*52*). Protein functional annotation was performed, and 19 CP

genes were significantly differentially expressed between heterozygotes and homozygotes of *bd*.

In addition, the orthologs of these CPs were analyzed in *Danaus plexippus*, *Papilio xuthu* and *D.*

melanogaster (Table S4). Furthermore, we identified 53 enzyme-encoding genes, 17

antimicrobial peptide genes, 6 transporter genes, 5 transcription factor genes, 5 cytochrome

genes, and others. CP genes were significantly enriched among the differentially expressed genes

(DEGs), and previous studies have found that CPs can affect pigmentation(*53*). Therefore, we

first conducted an investigation of the CP genes. Among them, 18 CP genes had Bm-mamo

binding sites in the upstream and downstream 2 kb genomic regions. In addition, we investigated

the expression level of the 18 CP genes in the integument from the 4[th] instar (day 1) to the

beginning of the 5[th] instar of *Bm-mamo* knockout lines (Fig. S10). The CP genes were

significantly upregulated at one or several time points in homozygous (*mamo*⁻/*mamo*⁻)

individuals compared with heterozygous (*mamo*⁻/+) *Bm-mamo* knockout line individuals.

Interestingly, the CP gene *BmorCPH24* was significantly upregulated at the feeding stage in *Bm-*

mamo knockout homozygotes. Previous studies have shown that BmorCPH24 deficiency can

lead to a marked decrease in pigmentation in silkworm larvae(*53*). Therefore, the expression of

some CP genes may be necessary for color patterns in caterpillars.

In addition, the synthesis of pigment substances is an important driver of color patterns(*54*).

Eight key genes (*TH*, *DDC*, *aaNAT*, *ebony*, *black*, *tan*, *yellow* and *laccase2*) involved in melanin

synthesis(*11*) were investigated in heterozygous and homozygous gene knockout lines of *Bm-*

mamo. Among them, *yellow*, *tan*, and *DDC* were significantly upregulated during the molting

period in the homozygous *Bm-mamo* knockout individuals (Fig. 6). The upstream and/or

downstream 2 kb genomic regions of *yellow*, *tan* and *DDC* contain the binding site of the Bm-

mamo protein. In addition, the expression of *yellow*, *DDC* and *tan* can promote the generation of

melanin(*55*).

To explore the interaction between the Bm-mamo protein and its binding sequence, EMSA

experiments were conducted. A binding site of Bm-mamo-S (CTGCGTGGT) was located

approximately 70 bp upstream of the transcription initiation site of the *Bm-yellow* gene. The

EMSA experiment showed that the Bm-mamo-S protein expressed in prokaryotes can bind to the

CTGCGTGGT sequence in vitro (Fig. S11). This suggests that the Bm-mamo-S protein can bind

to the upstream region of the *Bm-yellow* gene and regulate its transcription.

Therefore, the *Bm-mamo* gene may control the color pattern of caterpillars by regulating key

melanin synthesis genes and 18 CP genes.

Discussion

Insects have evolved many important phenotypes during the process of adapting to the

environment. Among these traits, color pattern is one of the most interesting. The biochemical

metabolic pathways of pigments, such as melanin, ommochromes, and pteridines, have been

identified in insects(*18*). However, the regulation of pigment metabolism-related genes and the

processes of the transport and deposition of pigment substances are not clear. In this study, we

discovered that the *Bm-mamo* gene negatively regulates melanin pigmentation in caterpillars.

When this gene is deficient, the body color of silkworms exhibits substantial melanism, and

changes in the expression of some melanin synthesis genes and some CP genes are also

significant.

Deficiency of some genes in the melanin synthesis pathway, such as *TH*, *yellow*, *ebony*, *tan*, and

aaNAT, can lead to variations in color patterns *(11)*. However, they often lead only to localized

pigmentation changes in later-instar caterpillars. For example, in *yellow*-mutant silkworms, the

eye spot, lunar spot, star spot, spiracle plate, head, and some sclerotized areas appear reddish

brown, and the other cuticle is consistent with that in the wild type in later instars of larvae (third

instar to fifth instar)(*56*). This situation is highly similar to that of the *tan* mutant in

silkworms(*57*). Why do the *yellow* and *tan* phenotypes appear only in a limited cuticle region in

later instars of silkworm larvae? One possible reason is that the expression of *yellow* and *tan* is

limited to a certain region by transcription regulators. Alternatively, other factors, such as CPs in

the cuticle, may limit pigmentation by interacting with pigment substances in the cuticle.

On the one hand, we investigated the expression levels of *yellow* and *tan* in the pigmented region

(lunar spot) and nonpigmented region of the epidermis during the 4th molting of the wild-type

Dazao strain. The expression level of *yellow* was significantly upregulated in the lunar spot (the

epidermis on the dorsum of the fifth body segment) compared with the nonpigmented region (the

epidermis on the dorsum of the sixth body segment) at 6 hours and 12 hours after molting.

Meanwhile, *tan* was significantly upregulated 18 hours after molting in the lunar spot region (Fig.

S12). This suggests that the upregulated expression of pigment synthesis genes at key time points

may be important for pigmentation. However, *yellow* and *tan* were still moderately expressed in

the nonpigmented epidermis, although they did not cause significant melanin pigmentation. This

indicates that pigment synthesis alone cannot dominantly determine the color pattern of the

cuticle in caterpillars.

On the other hand, synthesized pigment substances need to be transported from epidermal cells

and embedded into the cuticle to allow pigmentation. Therefore, the correct cuticle structure and

location of cuticular proteins in the cuticle may be important factors affecting pigmentation.

Previous studies have shown that the lack of expression of *BmorCPH24*, which encodes

important components of the endocuticle, can lead to dramatic changes in body shape and a

significant reduction in the pigmentation of caterpillars(*53*). We crossed *Bo* (*BmorCPH24* null

mutation) and *bd* to obtain $F_1(Bo/+^{Bo}, bd/+)$, then self-crossed F_1 and observed the phenotype of

F_2. The lunar spots and star spots decreased, and light-colored stripes appeared on the body

segments, but the other areas still had significant melanin pigmentation in double mutation (*Bo*,

bd) individuals (Fig. S13). However, in previous studies, introduction of *Bo* into *L* (ectopic

expression of *wnt1* results in lunar stripes generated on each body segment)(*24*) and *U*

(overexpression of *SoxD* results in excessive melanin pigmentation of the epidermis)(*58*) strains

by genetic crosses can remarkably reduce the pigmentation of *L* and *U* (*53*). Interestingly, there

was a more significant decrease in pigmentation in the double mutants (*Bo*, *L*) and (*Bo*, *U*) than

in (*Bo*, *bd*). This suggests that *Bm-mamo* has a stronger ability than *wnt1* and *SoxD* to regulate

pigmentation. On the one hand, mamo may be a stronger regulator of the melanin metabolic

pathway, and on the other hand, mamo may regulate other CP genes to reduce the impact of

BmorCPH24 deficiency.

How do CPs affect pigmentation? One study showed that some CPs can form "pore canals" to

transport some macromolecules(*59*). In addition, some CPs can be crosslinked with

catecholamines, which are synthesized in the melanin metabolism pathway(*60*). Because there

are no live cells in the cuticle, melanin precursor substances may be transported by the pore

canals formed by some CPs and fixed to specific positions through cross-linking with CPs. The

cuticular protein TcCPR4 is needed for the formation of pore canals of the cuticle in *Tribolium*

castaneum(61). In contrast, the vertical pore canal is lacking in the less pigmented cuticles of *T.*

castaneum(62). This suggests that the pore canals constructed by TcCPR4 may transport

pigments and contribute to cuticle pigmentation in *T. castaneum*. Moreover, the melanin

metabolites N-acetyldopamine (NADA) and N-β-alanyldopamine (NBAD) can target and

sclerotize the cuticle by cross-linking with specific cuticular proteins (*57*). This suggests that

pigments interact with specific CPs, thereby affecting pigmentation, hardening properties and the

structure of the cuticle. Interestingly, a study found that pigments, in addition to absorbing

specific wavelengths, can affect cuticle polymerization, density, and refractive index, which in

turn affects the reflected wavelengths that produce structural color in butterfly wing scales(*63*).

This implies that the interaction between pigments and CPs can be very subtle, resulting in the

formation of unique nanostructures, such as those on wing scales, that produce brilliant structural

colors.

Consequently, to maintain the accuracy of the color pattern, the localization of CPs in the cuticle

may be very important. In a previous study employing microarray analysis, different CPs were

found in differently colored areas of the epidermis in *Papilio xuthus* larvae(*64*). We investigated

the CP genes highly expressed in the black region of *Papilio xuthus* caterpillars. Thirteen genes

had orthologous genes in silkworms (Table S5). Among them, 11 genes (*BmorCPR67*,

BmorCPR71, *BmorCPR76*, *BmorCPR79*, *BmorCPR99*, *BmorCPR107*, *BmorCPT4*, *BmorCPH5*,

BmorCPFL4, *BmorCPG27* and *BmorCPG4*) were significantly upregulated in homozygous

(*mamo⁻/mamo⁻*) knockout individuals at some time points from the 3rd day of the 4th instar to the

5th instar; *BmorCPG4* was also among the 18 previously detected CP genes, and two genes

(*BmorCPG38* and *BmorCPR129*) were not different between homozygous (*mamo⁻/mamo⁻*) and

heterozygous (*mamo⁻/+*) individuals (Fig. S14).

The 28 CP genes mentioned above have significantly upregulated expression in homozygous

(*mamo⁻/mamo⁻*) gene knockout individuals at some stages. These CPs may be involved in the

transportation or cross-linking of melanin in the cuticle. However, among them, there were no

differences in their expression level in some periods compared with the control, and some genes

were significantly downregulated at some time points in melanic individuals (Fig. 7). This

suggests that the regulation of CP genes is complex and may involve other transcription factors

and feedback effects. CPs are essential components of the insect cuticle and are involved in

cuticular microstructure construction (*65*), body shape development(*66*), wing

morphogenesis(*67*), and pigmentation(*53*). CP genes usually account for over 1% of the total

genes in an insect genome and can be categorized into several families, including CPR, CPG,

CPH, CPAP1, CPAP3, CPT, CPF and CPFL(*68*). The CPR family is the largest group of CPs,

containing a chitin-binding domain called the Rebers and Riddiford motif (R&R)(*69*). The

variation in the R&R consensus sequence allows subdivision into three subfamilies (RR-1, RR-2,

and RR-3)(*70*). Among the 28 CPs, 11 RR-1 genes, 6 RR-2 genes, 4 hypothetical cuticular

protein (CPH) genes, 3 glycine-rich cuticular protein (CPG) genes, 3 cuticular protein Tweedle

motif (CPT) genes, and 1 CPFL (like the CPFs in a conserved C-terminal region) gene were

identified. The RR-1 consensus among species is usually more variable than RR-2, which

suggests that RR-1 may have a species-specific function. RR-2 often clustered into several

branches, which may be due to gene duplication events in co-orthologous groups and may result

in conserved functions between species (*71*). The classification of CPH is due to their lack of

known motifs. In the epidermis of Lepidoptera, the CPH genes often have high expression levels.

For example, *BmorCPH24* had a highest expression level, in silkworm larvae epidermis(*72*). The

CPG protein is rich in glycine. The CPH and CPG genes are less commonly found in insects

outside the order Lepidoptera (*73*). This suggests that they may provide species specific

functions for the Lepidoptera. CPT contains a Tweedle motif, and the *TweedleD1* mutation has a

dramatic effect on body shape in *D. melanogaster(74)*. The CPFL members are relatively

conserved in species and may be involved in the synthesis of larval cuticles(*75*). CPT and CPFL

may have relatively conserved functions among insects. The CP genes are a group of rapidly

evolving genes, and their copy numbers may undergo significant changes in different species. In

addition, RNAi experiments on 135 CP genes in brown planthopper (*Nilaparvata lugens*)

showed that deficiency of 32 CP genes leads to significant defective phenotypes, such as lethal,

developmental retardation, etc. It is suggested that the 32 CP genes are indispensable, and other

CP genes may have redundant and complementary functions (*76*). In previous studies, it was

found that the construction of the larval cuticle of silkworms requires the precise expression of

over two hundred CP genes(*22*). The production, interaction, and deposition of CPs and pigments

are complex and precise processes, and our research shows that *Bm-mamo* plays an important

regulatory role in this process in silkworm caterpillars. For further understanding of the role of

CPs, future work should aim to identify the function of important cuticular protein genes and the

deposition mechanism in the cuticle.

In addition, among the 191 DEGs found in the comparative transcriptome data, we also

discovered some interesting genes. For example, *BGIBMGA013242*, encoding a major facilitator

superfamily protein (MFS), named *BmMFS*, which is responsible for the *cheek and tail spot (cts)*

mutant, was significantly upregulated in the *bd* mutant. Deficiency of this gene (*BmMFS*) results

in chocolate-colored head and anal plates of the silkworm caterpillar(*77*). In the ommochrome

metabolic pathway, *Bm-re*, encoding an MFS protein, may function in the transportation of some

amino acids, such as cysteine or methionine, into pigment granules(*78*). Therefore, the encoded

product of *BmMFS* may participate in pigmentation by promoting the maturation of pigment

granules. Moreover, *BGIBMGA013576* and *BGIBMGA013656*, which encode MFS domain-

containing proteins belonging to the solute carrier family 22 (SLC22) and solute carrier family 2

(SLC2) families, respectively, were significantly upregulated in *bd*. These two genes may

participate in pigmentation in a similar manner to the *BmMFS* gene. In addition, the *red*

Malpighian tubules (*red*) gene encodes LysM domain-containing proteins. Deficiency of *red*

results in a significant decrease in orange pterin in the wing of *Colias* butterflies. Research

suggests that the product of *red* may interact with V-ATPase to modulate vacuolar pH across a

variety of endosomal organelles, thereby affecting the maturation of pigment granules in

cells(*79*). This indicates that the maturation of pigment granules plays an important role in the

coloring of Lepidoptera, and *Bm-mamo* may be involved in regulating the maturation of pigment

granules in the epidermal cells of silkworms.

Bm-mamo may affect the synthesis of melanin in epidermis cells by regulating *yellow*, *DDC*, and

tan; regulate the maturation of melanin granules in epidermis cells through *BmMFS*; and affect

the deposition of melanin granules in the cuticle by regulating CP genes, thereby

comprehensively regulating the color pattern in caterpillars.

Moreover, in *D. melanogaster*, *mamo* is needed for functional gamete production. In the

silkworm, *Bm-mamo* has a conserved function in female reproduction. However, *mamo* has

developed new functions in color pattern regulation in silkworm caterpillars. We found that in *D.*

melanogaster, the *mamo* gene is mainly expressed during the adult stage; it is not expressed

during the 1st instar or 2nd instar larval stage and has very low expression at the 3rd instar (Table

S6). In addition, several binding sites of mamo were found near the TTS of *yellow* in *D.*

melanogaster (Fig. S15). The *yellow* gene is a key melanin metabolic gene with upstream and

intron sequences that have been identified as multiple CREs, and it has been considered a

research model for CREs (*80, 81*). This may be due to a change in the expression pattern of

mamo, causing this gene to develop a new function of regulating coloration in silkworm

caterpillars.

It is generally believed that changes in gene expression patterns are the result of the evolution of

CREs. Because CREs have important functions in the spatiotemporal expression pattern

regulation of genes, many are found in the noncoding region, which is relatively prone to

sequence variation compared with the sequences of coding genes(*49*). However, the molecular

mechanism underlying the sequence change of CREs is not clear.

Transcription factors (TFs), because they recognize relatively short sequences, generally between

4 and 20 bp in length, can have many binding sites in the genome(*82*). Therefore, one member of

the TF family has the potential to regulate almost all genes in one genome. For example, in the

PWM sequence of the mamo protein recognition sequence, 10,622 genes (~73% of the total

number of genes) contain binding sites within 2 kb of their up/downstream region in silkworms

(Table S7 and Table S8). Divergence of CREs is currently considered an important cause of

phenotypic evolution(*83*). For example, the marine form of the three-spined stickleback

(*Gasterosteus aculeatus*) has thick armor, and the lake population (which was recently derived

from the marine form) does not. Research has shown that pelvic loss in different natural

populations of three-spined stickleback fish occurs by regulatory mutations deleting a tissue-

specific enhancer (*Pel*) of the *pituitary homeobox transcription factor 1* (*Pitx1*) gene. The

researchers genotyped 13 pelvic-reduced populations of three-spined stickleback from disparate

geographic locations. Nine of the 13 pelvic-reduced stickleback populations had sequence

deletions of varying lengths, all of which were located at the *Pel* enhancer. The author suggested

that the *Pitx1* locus of the stickleback genome may be prone to double-stranded DNA breaks that

are subsequently repaired by nonhomologous end joining (NHEJ) (*84*). There are also examples

of this in Lepidoptera. The *cortex* gene encodes a member of the Cdc20/fizzy family of cell cycle

regulators. The *cortex* gene locus is frequently mutated in Lepidoptera, such as the black

(*carbonaria*) form of the peppered moth (*Biston betularia*) in industrial melanism(*85*), the

diversification of the leaf wing of oakleaf butterflies(*86*), the mimicry wing patterning in

Heliconius butterflies(*87*), and the plastic wing pattern in *Junonia coenia (88)*. The explanation

of DNA fragility is very interesting, but it still cannot explain why DNA sequence mutations

occur frequently in a specially site corresponding to the adaptive phenotype. DNA fragility sites

are widely distributed in the genome(*89*). Under the stimulation of the same environmental

conditions, all fragile sites are thought to undergo frequent mutations, and this clearly brings

adverse burdens to the organism.

Some research suggests that common fragile sites (CFSs) are specific regions in the genome of

all individuals that are prone to DNA double stranded breaks (DSBs) and subsequent

rearrangements, and the DSB sites are correlated with epigenetic markers for chromatin

accessibility, including DNaseI HSSs, H3K4me3, and CTCF binding sites(*90, 91*).

Moreover, some studies have shown that there is a mutation bias in the genome; compared with

the intergenic region, the mutation frequency is reduced by half inside gene bodies and by two-

thirds in essential genes. In addition, they compared the mutation rates of genes with different

functions. The mutation rate in the coding region of essential genes (such as translation) is the

lowest, and the mutation rates in the coding region of specialized functional genes (such as

environmental response) are the highest. These patterns are mainly affected by the features of the

epigenome(*92*). Epigenetic modifications include DNA methylation, histone modifications,

remodeling of nucleosomes and higher order chromatin reorganization(*93*). TFs, such as BTB-

ZF proteins, can recruit histone modification factors such as DNA methyltransferase 1 (DMNT1),

cullin3 (CUL3), histone deacetylase 1 (HDAC1), and histone acetyltransferase 1 (HAT1) to

perform chromatin remodeling at specific genomic sites based on specific DNA recognition

capabilities(*94*). In addition, programmed events related to DSBs and repair at genome-specific

sites have been reported. For example, PRDM9 can determine recombination hotspots by H3K4

and H3K36 trimethylation (H3K4me3 and H3K36me3) of nucleosomes near its DNA-binding

site in meiosis recombination(*95*). This suggests that, on the basis of randomness, TFs and

epigenomic features can determine the mutation bias at unique positions of the genome.

460
Materials and Methods

Silkworm strains

The *bd* and *bd^f* mutant strains and wild-type strains Dazao and N4 were obtained from the bank

of genetic resources of Southwest University. Silkworms were reared on mulberry leaves or

artificial diets at 25 °C and 73% relative humidity under dark for 12 hours and light for 12 hours.

Positional cloning of the *bd* locus

For mapping of the *bd* locus, F1 heterozygous individuals were obtained from a cross between a

bd^f strain and a Dazao strain. Then, an F1 female was crossed with a *bd^f* male (BC1F), and an F1

male was backcrossed with a *bd^f* female (BC1M). A total of 1162 BC1M individuals were used

for recombination analysis. Genomic DNA was extracted from the parents (Dazao, *bd^f* and F1)

and each BC1 individual using the phenol chloroform extraction method. Available DNA

molecular markers were sought through polymorphism screening and linkage analysis. The

primers used for mapping are listed in Table S9.

Phylogenetic analysis

To determine whether *Bm-mamo* orthologs existed in other species, the BlastX program of the

National Center for Biotechnology Information (NCBI) (http://www.ncbi.nlm.nih.gov/BLAST/)

was used. The sequences of *Bm-mamo-L* and *Bm-mamo-S* were blasted against the nonredundant

protein sequence (nr) database. Sequences with a maximum score and E-value$\leq 10^{-4}$ were

downloaded. The sequences of multiple species were subjected to multiple sequence alignment

of the predicted amino acid sequences by MUSCLE(*96*). Then, a phylogenetic tree was

constructed using the neighbor-joining method with the MEGA7 program (Pearson model). The

confidence levels for various phylogenetic lineages were estimated by bootstrap analysis (2,000

replicates).

Quantitative PCR

Total RNA was isolated from the whole body and integument of the silkworms using TRIzol

reagent (Invitrogen, California, USA), purified by phenol chloroform extraction and then reverse

transcribed with a PrimeScript™ RT Reagent Kit (TAKARA, Dalian, China) according to the

manufacturers' protocol.

qPCR was performed using a CFX96™ Real-Time PCR Detection System (Bio-Rad, Hercules,

CA) with a Real-Time PCR System and an iTaq Universal SYBR Green Supermix System (Bio-

Rad). The cycling parameters were as follows: 95 °C for 3 min, followed by 40 cycles of 95 °C

for 10 s and annealing for 30 s. The primers used for target genes are listed in Table S9.

Expression levels for each sample were determined with biological replicates, and each sample

was analyzed in triplicate. The gene expression levels were normalized against the expression

levels of the ribosomal protein L3 (RpL3). The relative expression levels were analyzed using

the classical $R=2^{-\Delta\Delta Ct}$ method.

siRNA for gene knockdown

siRNAs for *Bm-mamo* were designed by the siDirect program (http://sidirect2.rnai.jp). The target

siRNAs and negative control were synthetized by Tsingke Biotechnology Company Limited. The

siRNA (5 μl, 1 μl/μg) was injected from the abdominal spiracle into the hemolymph at the fourth

instar (day 3) larval stage. Immediately after injection, phosphate-buffered saline (pH 7.3)

droplets were placed nearby, and a 20-voltage pulse for one second and pause for one second

were repeated 3 times. The phenotype was observed for fifth-instar larvae. The left and right

epidermis were separately dissected from the injected larvae, and RNA was extracted. Then,

cDNA was synthetized, and the expression level of the gene was detected by qPCR.

sgRNA synthesis and RNP complex assembly

CRISPRdirect (http://crispr.dbcls.jp/doc/) online software was used to screen appropriate single

guide RNA (sgRNA) target sequences. The gRNAs were synthetized by Beijing Genomics

Institute. sgRNA templates were transcribed using T7 polymerase with RiboMAX™ Large-Scale

RNA Production Systems (Promega, Beijing, China) according to the manufacturer's instructions.

The RNA transcripts were purified using 3 M sodium acetate (pH 5.2) and anhydrous ethanol

(1:30) precipitation, washed using 75% ethanol, and eluted in nuclease-free water. All injection

mixes contained 300 ng/µL Cas9 nuclease (Invitrogen, California, USA) and 300 ng/µL purified

sgRNA. Before injection, mixtures of Cas9 nuclease and gRNA were incubated for 15 min at

37 °C to reconstitute active ribonucleoproteins (RNPs).

Microinjection of embryos

For embryo microinjection, microcapillary needles were manufactured using a PC-10 model

micropipette puller (Narishige, Tokyo, Japan). Microinjection was performed using Eppendorf's

TransferMan NK 2 and a FemtoJet 4i system (Eppendorf, Hamburg, Germany). The eggs used

for microinjection came from the mating of female and male wild-type Dazao moths. Within 4

hours of being laid, the eggs were adhered onto a clean glass slide. CRISPR/Cas9-messenger

RNP mixtures with volumes of approximately 1 nL were injected into the middle of the eggs,

and the wound was sealed with glue. All injected embryos were allowed to develop in a climate

chamber at 25 °C and 80% humidity.

Comparative genomics

The reference genome of Dazao was downloaded from Silkbase (https://silkbase.ab.a.u-

tokyo.ac.jp/cgi-bin/index.cgi). The bd^f genome was obtained from the silkworm pangenome

project(*34*).

The short reads of the bd^f strains were mapped to the silkworm reference genome by BWA55

v0.7.17 mem with default parameters. The SAMtools56 v1.11 and Picard v2.23.5

(https://broadinstitute.github.io/picard/) programs were used to filter the unmapped and

duplicated reads. A GVCF file of the samples was obtained using GATK57 v4.1.8.1

HaplotypeCaller with the parameter -ERC = GVCF. The VCF files of insertion/deletions (indels)

and single-nucleotide polymorphisms (SNPs) were used for further analysis by eGPS software.

Downstream target gene screening

Online software (http://zf.princeton.edu) was used to predict the DNA-binding site for Cys2His2

zinc finger proteins. The confident ZF domains with scores higher than 17.7 were chosen. RF

regression on the B1H model was used to predict the DNA-binding sites. Then, the sequence

logo and position weight matrices (PWMs) of the DNA-binding sites were obtained. The

sequences 2 kb upstream and downstream of the predicted genes were extracted by a Perl script.

The FIMO package of the MEME suite was used to search for binding sites in the silkworm

genome.

Analysis of EMSA

Primers with binding sites and their flanking sequences were designed according to the FIMO

results. A biotin label was added to one end of the upstream primer (probe), and the downstream

primer was normal. Primers with the same sequence as the labeled probes were used as

competitive probes. The EMSA experiment was conducted with an EMSA reagent kit (Beyotime,

Shanghai, China) according to the manufacturers' protocol.

Parthenogenesis

Virgin moths were dissected in 0.9% bacterium-free physiological saline. The ovaries were

extracted, and water was removed by clean filter paper. The ovaries were stored at 25 °C and 40%

relative humidity for 10 hours. The eggs were treated in 46 °C warm water for 18 minutes and

then transferred to room temperature for 5-10 minutes. The activated egg was protected in an

environment of 15-17 °C and 18% relative humidity for 3 days. Then, it was immersed in acid

immediately (with a hydrochloric acid specific gravity of 1.075 at 46 °C for 5 minutes) or

transferred to the same conditions as the hibernating eggs for protection. The average hatching

rate of silkworm eggs treated in this way was 50-60%.

References

1.   G. A. Sword, S. J. Simpson, O. T. M. El Hadi, H. Wilps, Density-dependent aposematism

in the desert locust. *P Roy Soc B-Biol Sci* **267**, 63-68 (2000).

2.   A. I. Barnes, M. T. Siva-Jothy, Density-dependent prophylaxis in the mealworm beetle

Tenebrio molitor L. (Coleoptera: Tenebrionidae): cuticular melanization is an indicator of

investment in immunity. *P Roy Soc B-Biol Sci* **267**, 177-182 (2000).

3.   N. F. Hadley, A. Savill, T. D. Schultz, Coloration and its thermal consequences in the

New Zealand tiger beetle *Neocicindela perhispida*. *J Therm Biol* **17**, 55-61 (1992).

4.   Y. G. Hu, Y. H. Shen, Z. Zhang, G. Q. Shi, Melanin and urate act to prevent ultraviolet

damage in the integument of the silkworm, *Bombyx mori*. *Arch Insect Biochem* **83**, 41-55

(2013).

5.   M. Stevens, G. D. Ruxton, Linking the evolution and form of warning coloration in

nature. *P Roy Soc B-Biol Sci* **279**, 417-426 (2012).

6.   K. K. Dasmahapatra *et al.*, Butterfly genome reveals promiscuous exchange of mimicry

adaptations among species. *Nature* **487**, 94-98 (2012).

7.   N. Gaitonde, J. Joshi, K. Kunte, Evolution of ontogenic change in color defenses of

swallowtail butterflies. *Ecol Evol* **8**, 9751-9763 (2018).

8.   B. S. Tullberg, S. Merilaita, C. Wiklund, Aposematism and crypsis combined as a result

of distance dependence: functional versatility of the colour pattern in the swallowtail

butterfly larva. *P Roy Soc B-Biol Sci* **272**, 1315-1321 (2005).

9.   P. J. Wittkopp, B. L. Williams, J. E. Selegue, S. B. Carroll, *Drosophila* pigmentation

evolution: divergent genotypes underlying convergent phenotypes. *P Natl Acad Sci USA*

100, 1808-1813 (2003).

10.   R. Futahashi, Y. Banno, H. Fujiwara, Caterpillar color patterns are determined by a two-

phase melanin gene prepatterning process: new evidence from *tan* and *laccase2*. *Evol*

Dev **12**, 157-167 (2010).

11.   Y. Matsuoka, A. Monteiro, Melanin pathway genes regulate color and morphology of

butterfly wing scales. *Cell Rep* **24**, 56-65 (2018).

12.   C. Finet *et al.*, Multi-scale dissection of wing transparency in the clearwing butterfly. *J R*

Soc Interface **20**, (2023).

13.   A. Gürses, M. Açıkyıldız, K. Güneş, M. S. Gürses, in *Dyes and Pigments*. (Springer

International Publishing, 2016), pp. 13-29.

14.   V. J. Lloyd, N. J. Nadeau, The evolution of structural colour in butterflies. *Curr Opin*

Genet Dev **69**, 28-34 (2021).

15.   A. E. Seago, P. Brady, J. P. Vigneron, T. D. Schultz, Gold bugs and beyond: a review of

iridescence and structural colour mechanisms in beetles (Coleoptera). *J R Soc Interface* **6**,

S165-S184 (2009).

16.   Y. Oba *et al.*, Resurrecting the ancient glow of the fireflies. *Sci Adv* **6**, (2020).

17.   A. J. Syed, J. C. Anderson, Applications of bioluminescence in biotechnology and beyond.

Chem Soc Rev **50**, 5668-5705 (2021).

18.   X. L. Tong, L. Qiao, J. W. Luo, X. Ding, S. Y. Wu, The evolution and genetics of

lepidopteran egg and caterpillar coloration. *Curr Opin Genet Dev* **69**, 140-146 (2021).

19.   F. Y. Dai *et al.*, Mutations of an arylalkylamine-*N*-acetyltransferase, *Bm-iAANAT*, are

613   responsible for silkworm melanism mutant. *J Biol Chem* **285**, 19553-19560 (2010).

614   20.   P. J. Wittkopp, J. R. True, S. B. Carroll, Reciprocal functions of the *Drosophila* yellow

615   and ebony proteins in the development and evolution of pigment patterns. *Development*

616   **129**, 1849-1858 (2002).

617   21.   J. Q. Liu *et al.*, Lepidopteran wing scales contain abundant cross-linked film-forming

618   histidine-rich cuticular proteins. *Commun Biol* **4**, (2021).

619   22.   Z. W. Yan *et al.*, A blueprint of microstructures and stage-specific transcriptome

620   dynamics of cuticle formation in *Bombyx mori*. *Int J Mol Sci* **23**, (2022).

621   23.   R. Futahashi *et al.*, Genome-wide identification of cuticular protein genes in the

622   silkworm, *Bombyx mori*. *Insect Biochem Molec* **38**, 1138-1146 (2008).

623   24.   J. Yamaguchi *et al.*, Periodic *Wnt1* expression in response to ecdysteroid generates twin-

624   spot markings on caterpillars. *Nat Commun* **4**, (2013).

625   25.   S. Yoda *et al.*, The transcription factor Apontic-like controls diverse colouration pattern in

626   caterpillars. *Nat Commun* **5**, (2014).

627   26.   H. Jin, T. Seki, J. Yamaguchi, H. Fujiwara, Prepatterning of *Papilio xuthus* caterpillar

628   camouflage is controlled by three homeobox genes: *clawless*, *abdominal-A*, and

629   *Abdominal-B*. *Sci Adv* **5**, (2019).

630   27.   W. A. Rogers *et al.*, A survey of the trans-regulatory landscape for *Drosophila*

631   *melanogaster* abdominal pigmentation. *Dev Biol* **385**, 417-432 (2014).

28.   S. Jeong, A. Rokas, S. B. Carroll, Regulation of body pigmentation by the Abdominal-B

Hox protein and its gain and loss in *Drosophila* evolution. *Cell* **125**, 1387-1399 (2006).

29.   H. D. Dufour, S. Koshikawa, C. Finet, Temporal flexibility of gene regulatory network

underlies a novel wing pattern in flies. *P Natl Acad Sci USA* **117**, 11589-11596 (2020).

30.   A. Prakash, C. Finet, T. Das Banerjee, V. Saranathan, A. Monteiro, *Antennapedia* and

optix regulate metallic silver wing scale development and cell shape in *Bicyclus anynana*

butterflies. *Cell Rep* **40**, (2022).

31.   R. D. Reed *et al.*, Drives the repeated convergent evolution of butterfly wing pattern

mimicry. *Science* **333**, 1137-1141 (2011).

32.   A. Kopp, I. Duncan, S. B. Carroll, Genetic control and evolution of sexually dimorphic

characters in *Drosophila*. *Nature* **408**, 553-559 (2000).

33.   L. Arnoult *et al.*, Emergence and diversification of fly pigmentation through evolution of

a gene regulatory module. *Science* **339**, 1423-1426 (2013).

34.   X. L. Tong *et al.*, High-resolution silkworm pan-genome provides genetic insights into

artificial selection and ecological adaptation. *Nat Commun* **13**, (2022).

35.   L. Cheng., D. Fangyin., X. Zhonghuai., Studies on the mutant systems of the *Bombyx*

mori gene bank. *Scientia agricultura sinica* **36**, 968-975 (2003).

36.   Y. Kawaguchi, T. Kusakabe, J. M. Lee, K. Koga, Manifestation of the *dilute black* (*bd*)

mutation and constitution of the *bd* Locus in *Bombyx mori*. *J Fac Agr Kyushu U* **52**, 355-

359 (2007).

37.   J. Duan *et al.*, SilkDB v2.0: a platform for silkworm (*Bombyx mori*) genome biology.

Nucleic Acids Res **38**, D453-D456 (2010).

38.   E. W. Sayers *et al.*, GenBank. *Nucleic Acids Res* **50**, D161-d164 (2022).

39.   M. Kawamoto, T. Kiuchi, S. Katsuma, SilkBase: an integrated transcriptomic and

656   genomic database for *Bombyx mori* and related species. *Database-Oxford* **2022**, (2022).

657   40.   V. J. Bardwell, R. Treisman, The POZ domain: a conserved protein-protein interaction

658   motif. *Gene Dev* **8**, 1664-1677 (1994).

659   41.   K. D. Huynh, V. J. Bardwell, The BCL-6 POZ domain and other POZ domains interact

660   with the co-repressors N-CoR and SMRT. *Oncogene* **17**, 2473-2484 (1998).

661   42.   P. Staller *et al.*, Repression of p15INK4b expression by Myc through association with

662   Miz-1. *Nat Cell Biol* **3**, 392-399 (2001).

663   43.   M. Mukai *et al.*, MAMO, a maternal BTB/POZ-Zn-finger protein enriched in germline

664   progenitors is required for the production of functional eggs in *Drosophila*. *Mech*

665   *Develop* **124**, 570-583 (2007).

666   44.   T. Ando, H. Fujiwara, Electroporation-mediated somatic transgenesis for rapid functional

667   analysis in insects. *Development* **140**, 454-458 (2013).

668   45.   C. O. Pabo, E. Peisach, R. A. Grant, Design and selection of novel Cys2His2 zinc finger

669   proteins. *Annu Rev Biochem* **70**, 313-340 (2001).

670   46.   S. A. Wolfe, L. Nekludova, C. O. Pabo, DNA recognition by Cys2His2 zinc finger

671   proteins. *Annu Rev Bioph Biom* **29**, 183-212 (2000).

672   47.   A. V. Persikov, M. Singh, De novo prediction of DNA-binding specificities for Cys2His2

673   zinc finger proteins. *Nucleic Acids Res* **42**, 97-108 (2014).

48.   S. Hira *et al.*, Binding of *Drosophila* maternal Mamo protein to chromatin and specific

DNA sequences. *Biochem Bioph Res Co* **438**, 156-160 (2013).

49.   P. J. Wittkopp, G. Kalay, *Cis*-regulatory elements: molecular mechanisms and

evolutionary processes underlying divergence. *Nat Rev Genet* **13**, 59-69 (2012).

50.   S. Preissl, K. J. Gaulton, B. Ren, Characterizing *cis*-regulatory elements using single-cell

epigenomics. *Nat Rev Genet*, (2022).

51.   E. Z. Kvon *et al.*, Genome-scale functional characterization of *Drosophila* developmental

enhancers *in vivo*. *Nature* **512**, 91-+ (2014).

52.   S. Y. Wu *et al.*, Comparative analysis of the integument transcriptomes of the *black dilute*

mutant and the wild-type silkworm *Bombyx mori*. *Sci Rep-Uk* **6**, (2016).

53.   G. Xiong *et al.*, Body shape and coloration of silkworm larvae are influenced by a novel

cuticular protein. *Genetics* **207**, 1053-1066 (2017).

54.   P. J. Wittkopp, S. B. Carroll, A. Kopp, Evolution in black and white: genetic control of

pigment patterns in *Drosophila*. *Trends Genet* **19**, 495-504 (2003).

55.   J. M. Gibert, Phenotypic plasticity in insects. *Biologie aujourd'hui* **214**, 33-44 (2020).

56.   R. Futahashi *et al.*, *yellow* and *ebony* are the responsible genes for the larval color

mutants of the silkworm *Bombyx mori*. *Genetics* **180**, 1995-2005 (2008).

57.   M. Y. Noh, S. Muthukrishnan, K. J. Kramer, Y. Arakane, Cuticle formation and

pigmentation in beetles. *Curr Opin Insect Sci* **17**, 1-9 (2016).

58.   N. N. Wang *et al.*, Abnormal overexpression of *SoxD* enhances melanin synthesis in the

Ursa mutant of *Bombyx mori*. *Insect Biochem Molec* **149**, (2022).

59.   K. J. Kramer *et al.*, Oxidative conjugation of catechols with proteins in insect skeletal

systems. *Tetrahedron* **57**, 385-392 (2001).

60.   M. Y. Noh, S. Muthukrishnan, K. J. Kramer, Y. Arakane, *Tribolium castaneum* RR-1

cuticular protein TcCPR4 is required for formation of pore canals in rigid cuticle. *Plos*

Genet **11**, (2015).

61.   M. Y. Noh *et al.*, Two major cuticular proteins are required for assembly of horizontal

laminae and vertical pore canals in rigid cuticle of *Tribolium castaneum. Insect Biochem*

Molec **53**, 22-29 (2014).

62.   S. Mun *et al.*, Cuticular protein with a low complexity sequence becomes cross-linked

during insect cuticle sclerotization and is required for the adult molt. *Sci Rep-Uk* **5**,

(2015).

63.   A. Prakash *et al.*, Nanoscale cuticle density variations correlate with pigmentation and

color in butterfly wing scales. *arXiv*, 2305.16628 (2023).

64.   R. Futahashi, H. Shirataki, T. Narita, K. Mita, H. Fujiwara, Comprehensive microarray-

based analysis for stage-specific larval camouflage pattern-associated genes in the

swallowtail butterfly, *Papilio xuthus. Bmc Biol* **10**, (2012).

65.   M. Y. Noh, S. Muthukrishnan, K. J. Kramer, Y. Arakane, Development and ultrastructure

of the rigid dorsal and flexible ventral cuticles of the elytron of the red flour beetle,

Tribolium castaneum. Insect Biochem Molec **91**, 21-33 (2017).

66.   R. Tajiri, N. Ogawa, H. Fujiwara, T. Kojima, Mechanical control of whole body shape by

a single cuticular protein Obstructor-E in *Drosophila melanogaster. Plos Genet* **13**,

(2017).

67.   X. M. Zhao *et al.*, The wing-specific cuticular protein LmACP7 is essential for normal

718   wing morphogenesis in the migratory locust. *Insect Biochem Molec* **112**, (2019).

719   68.   J. H. Willis, Structural cuticular proteins from arthropods: Annotation, nomenclature, and

720   sequence characteristics in the genomics era. *Insect Biochem Molec* **40**, 189-204 (2010).

721   69.   J. E. Rebers, L. M. Riddiford, Structure and expression of a *Manduca sexta* larval cuticle

722   gene homologous to *Drosophila* cuticle genes. *J Mol Biol* **203**, 411-423 (1988).

723   70.   M. V. Karouzou *et al.*, *Drosophila* cuticular proteins with the R&R Consensus:

724   Annotation and classification with a new tool for discriminating RR-1 and RR-2

725   sequences. *Insect Biochem Molec* **37**, 754-760 (2007).

726   71.   E. H. Chen, Q. L. Hou, Identification and expression analysis of cuticular protein genes

727   in the diamondback moth, *Plutella xylostella* (Lepidoptera: Plutellidae). *Pestic Biochem

728   Phys* **178**, (2021).

729   72.   X. Fu *et al.*, Genome-Wide Identification and Transcriptome-Based Expression Profile of

730   Cuticular Protein Genes in *Antheraea pernyi*. *Int J Mol Sci* **24**, (2023).

731   73.   C. H. Yang *et al.*, Identification, expression pattern, and feature analysis of cuticular

732   protein genes in the pine moth *Dendrolimus punctatus* (Lepidoptera: Lasiocampidae).

733   *Insect Biochem Molec* **83**, 94-106 (2017).

734   74.   X. Guan, B. W. Middlebrooks, S. Alexander, S. A. Wasserman, Mutation of TweedleD, a

735   member of an unconventional cuticle protein family, alters body shape in *Drosophila*. *P

736   Natl Acad Sci USA* **103**, 16794-16799 (2006).

75.   T. Togawa, W. A. Dunn, A. C. Emmons, J. H. Willis, CPF and CPFL, two related gene

families encoding cuticular proteins of *Anopheles gambiae* and other insects. *Insect*

Biochem Molec **37**, 675-688 (2007).

76.   P. L. Pan *et al.*, A comprehensive omics analysis and functional survey of cuticular

proteins in the brown planthopper. *P Natl Acad Sci USA* **115**, 5175-5180 (2018).

77.   K. Ito *et al.*, Positional cloning of a gene responsible for the *cts* mutation of the silkworm,

Bombyx mori. *Genome* **55**, 493-504 (2012).

78.   M. Osanai-Futahashi *et al.*, Identification of the *Bombyx red egg* gene reveals

involvement of a novel transporter family gene in late steps of the insect ommochrome

biosynthesis pathway. *J Biol Chem* **287**, 17706-17714 (2012).

79.   J. J. Hanly *et al.*, Genetics of yellow-orange color variation in a pair of sympatric sulphur

butterflies. *Cell Rep* **42**, (2023).

80.   G. Kalay, J. Lachowiec, U. Rosas, M. R. Dome, P. Wittkopp, Redundant and cryptic

enhancer activities of the *Drosophila yellow* gene. *Genetics* **212**, 343-360 (2019).

81.   Y. Q. Xin *et al.*, Enhancer evolutionary co-option through shared chromatin accessibility

input. *P Natl Acad Sci USA* **117**, 20636-20644 (2020).

82.   A. Jolma, J. Taipale, Methods for analysis of transcription factor DNA-binding specificity

in vitro. *Sub-cellular biochemistry* **52**, 155-173 (2011).

83.   K. Mattioli *et al.*, *Cis* and *trans* effects differentially contribute to the evolution of

promoters and enhancers. *Genome biology* **21**, (2020).

84.   Y. F. Chan *et al.*, Adaptive evolution of pelvic reduction in sticklebacks by recurrent

deletion of a *Pitx1* enhancer. *Science* **327**, 302-305 (2010).

85.   A. E. van't Hof *et al.*, The industrial melanism mutation in British peppered moths is a

760   transposable element. *Nature* **534**, 102-+ (2016).

761   86.   S. T. Wang *et al.*, The evolution and diversification of oakleaf butterflies. *Cell* **185**, 3138-

762   + (2022).

763   87.   N. J. Nadeau *et al.*, The gene cortex controls mimicry and crypsis in butterflies and moths.

764   *Nature* **534**, 106-+ (2016).

765   88.   K. R. L. van der Burg *et al.*, Genomic architecture of a genetically assimilated seasonal

766   color pattern. *Science* **370**, 721-+ (2020).

767   89.   F. Ji *et al.*, Genome-wide high-resolution mapping of mitotic DNA synthesis sites and

768   common fragile sites by direct sequencing. *Cell Res* **30**, 1009-1023 (2020).

769   90.   S. Kodali *et al.*, Epigenomic signatures associated with spontaneous and replication

770   stress-induced DNA double strand breaks. *Front Genet* **13**, (2022).

771   91.   R. Mourad, K. Ginalski, G. Legube, O. Cuvier, Predicting double-strand DNA breaks

772   using epigenome marks or DNA at kilobase resolution. *Genome biology* **19**, (2018).

773   92.   J. G. Monroe *et al.*, Mutation bias reflects natural selection in *Arabidopsis thaliana*.

774   *Nature* **602**, 101-105 (2022).

775   93.   B. A. Nacev *et al.*, The epigenomics of sarcoma. *Nat Rev Cancer* **20**, 608-623 (2020).

776   94.   R. Mathew *et al.*, BTB-ZF factors recruit the E3 ligase cullin 3 to regulate lymphoid

777   effector programs. *Nature* **491**, 619-+ (2012).

778   95.   T. Bhattacharyya *et al.*, Prdm9 and meiotic cohesin proteins cooperatively promote DNA

779   double-strand break formation in mammalian spermatocytes. *Curr Biol* **29**, 1002-

780   1018.e1007 (2019).

781   96.   R. C. Edgar, MUSCLE: multiple sequence alignment with high accuracy and high

782   throughput. *Nucleic Acids Res* **32**, 1792-1797 (2004).

Acknowledgments

Funding:

This work was supported by the National Natural Science Foundation of China (No. 32002230), the Fundamental Research Funds for the Central Universities in China (No. SWU120024), the National Natural Science Foundation of China (No. 31830094, No. U20A2058), the China Agriculture Research System of MOF and MARA (No. CARS-18-ZJ0102, No. CARS-18-ZJ0103), the Natural Science Foundation of Chongqing, China (No. cstc2021jcyj-cxtt0005), the Yibin Academy of Southwest University (XNDX2022020015), the Southwest University Innovation Research 2035 Pilot Program (No. SWU-XDPY22011) and the High-level Talents Program of Southwest University (No. SWURC2021001).

Author contributions:

FY Dai conceived and designed the experiments. SY Wu, CX Peng, JW Luo, KP Lu and CH Zhang performed the study. SY Wu, XL Tong, KP Lu, CL Li, X Ding, YR Lu, XH Duan, D Tan and H Hu analyzed the data. SY Wu wrote the paper. XL Tong and FY Dai edited and revised the manuscript.

Competing interests:

The authors declare that they have no competing interests.

Data and material availability:

The authors confirm that the data supporting the findings of this study are available within the article and its supplementary materials.

805 **Figures and Tables**

813 **Fig. 1. Phenotypes of bd, bd^f, and wild-type Dazao larvae.**

814 The epidermis of bd is dark gray, and the epidermis of bd^f is light gray at the 5th instar (day 3) of

815 silkworm larvae. The bar indicates 1 centimeter.

Fig. 2. Positional cloning of the *bd* locus.

(A) We used 532 BC1 individuals to map the *bd* locus between PCR markers c and i. The

numbers above the DNA markers indicate recombination events.

(B) A total of 1162 BC1 individuals were used to narrow the *bd* locus to an approximately 400

kb genomic region.

(C) Partial enlarged view of the region responsible for *bd*. This region contains 4 predicted genes,

BGIBMGA012517, *BGIBMGA012518*, *BGIBMGA012519* and *BGIBMGA014089*.

(D) Analysis of nucleotide differences in the region responsible for *bd*. The green block indicates

the deletion of the genome in *bd* mutants. The black vertical lines indicate the SNPs and indels of

*bd^f^* mutants.

A

B

C

Fig. 3. Spatiotemporal expression of *Bm-mamo*.

(A) Temporal expression of *Bm-mamo*. In the molting stage and adult stage, this gene is

significantly upregulated. M: molting stage, W: wandering stage, P: pupal stage, P1: Day 1 of

pupal stage, A: adult stage, 4th3d indicates 4th instar day 3. 1st to 5th denote the first instar of

larvae to fifth instar of larvae, respectively.

(B) Tissue-specific expression of 4th-instar molting larvae. *Bm-mamo* had relatively high

expression levels in the midgut, head, and epidermis. AMSG: anterior division of silk gland and

middle silk gland, PSG: posterior silk gland.

(C) Detailed analysis of *Bm-mamo* at the 4th larval stage in the epidermis of the Dazao strain.

Bm-mamo expression is upregulated during the molting stage. HCS indicates the head capsule

stage. The "h" indicates the hour, 90 h: at 90 hours of the 4th instar.

(**A**) siRNA was introduced by microinjection followed by electroporation. "+" and "-" indicate

the positive and negative poles of the electrical current. Scale bars, 1 cm.

(**B**) Partial magnification of the siRNA experimental group and negative control group. Scale

bars, 0.2 cm.

(**C**) and (**D**) Relative expression levels of *Bm-mamo* in the negative control and RNAi groups

were determined by qPCR analysis. The means ± s.d.s. Numbers of samples are shown in the

upper right in each graph. **$P<0.01$, paired Student's t test (NS, not significant).

(E) Efficiency statistics of RNAi.

Fig. 5. Knockout of *Bm-mamo*

(**A**) Larval phenotype in G3 fourth-instar larvae of Dazao targeted for *Bm-mamo*. Scale bars, 1

cm.

(**B**) Partial magnification of the *Bm-mamo* knockout individual and control. Scale bars, 0.2 cm.

(**C**) After 48 hours of laying eggs, the pigmentation of the eggs indicates that those produced by

knockout homozygous females cannot undergo normal pigmentation and development.

943   (**D**) Genomic structure of *Bm-mamo*. Open reading frame (blue), untranslated region (black). The

944   gRNA 1 and gRNA 2 sequences are shown.

945   (**E**) Sequences of the *Bm-mamo* knockout individuals. Lines 1 and 2 indicate deletions of 15 and

946   71 bp, respectively.

Fig. 6. Melanin metabolism pathway and qPCR of related genes.

(**A**) The melanin metabolism pathway. Blue indicates amino acids and catecholamines, red

indicates the names of genes, and black indicates the names of enzymes.

(**B**) Relative expression levels of eight genes in the heterozygous *Bm-mamo* knockout group

(blue) and homozygous *Bm-mamo* knockout group (red) as determined by qPCR analysis. The

means ± s.d.s. *P<0.05, paired Student's t test. 4th3d indicates 4th instar day 3, M indicates

molting, and h indicates hours.

1000

Fig. 7. Heatmap of 28 differentially expressed cuticular protein genes.

Differentially expressed cuticular protein genes between homozygotes (*mamo⁻/mamo⁻*) and

heterozygotes (*mamo⁻/+*) of *Bm-mamo* knockout line individuals. Red indicates upregulation in

homozygous individuals. Blue indicates downregulation in homozygous individuals. White

indicates no difference in expression level or no investigation. 4th1d indicates 4-instar day one,

M indicates molting, and h indicates hours.

1007

1008 **Additional information**

1009 Supplementary Information is available for this paper.

1010

1011

1012

1013