Circadian rhythm disruptions associated with opioid use disorder in the synaptic proteomes of the human dorsolateral prefrontal cortex and nucleus accumbens

Stephanie Puig[1,2], Xiangning Xue[3], Ryan Salisbury[2], Micah A. Shelton[2], Sam-Moon Kim[2], Mariah A. Hildebrand[2], Jill R. Glausier[2], Zachary Freyberg[2,4], George C. Tseng[3], Anastasia K. Yocum[5], David A. Lewis[2], Marianne L. Seney[2], Matthew L. MacDonald[2*], and Ryan W. Logan[1,2,6,7*]

[1]Department of Pharmacology, Physiology and Biophysics, Boston University School of Medicine, Boston, MA, USA

[2]Department of Psychiatry, University of Pittsburgh School of Medicine, Pittsburgh, PA, USA

[3]Department of Biostatistics, University of Pittsburgh, Pittsburgh, PA, USA

[4]Department of Cell Biology, University of Pittsburgh, Pittsburgh, PA, USA

[5]A2IDEA, LLC, Ann Arbor, MI, USA

[6]Department of Psychiatry, University of Massachusetts Chan Medical School, Worcester, MA, USA

[7]Department of Neurobiology, University of Massachusetts Chan Medical School, Worcester, MA, USA

*Co-corresponding authors

Ryan W. Logan, PhD

364 Plantation Street, LRB Office 611

Worcester, MA 01605

ryan.logan@umassmed.edu

Matthew L. MacDonald

203 Lothrop St,

Thomas E. Starzl Biomedical Science Tower, Room W1655

Pittsburgh, PA 15213

macdonaldml@upmc.edu

Abstract

Opioid craving and relapse vulnerability is associated with severe and persistent sleep and circadian rhythm disruptions. Understanding the neurobiological underpinnings of circadian rhythms and opioid use disorder (OUD) may prove valuable for developing new treatments for opioid addiction. Previous work indicated molecular rhythm disruptions in the human brain associated with OUD, highlighting synaptic alterations in the dorsolateral prefrontal cortex (DLPFC) and nucleus accumbens (NAc)—key brain regions involved in cognition and reward, and heavily implicated in the pathophysiology of OUD. To provide further insights into the synaptic alterations in OUD, we used mass-spectrometry based proteomics to deeply profile protein expression alterations in bulk tissue and synaptosome preparations from DLPFC and NAc of unaffected and OUD subjects. We identified 55 differentially expressed (DE) proteins in DLPFC homogenates, and 44 DE proteins in NAc homogenates, between unaffected and OUD subjects. In synaptosomes, we identified 161 and 56 DE proteins in DLPFC and NAc, respectively, of OUD subjects. By comparing homogenate and synaptosome protein expression, we identified proteins enriched specifically in synapses that were significantly altered in both DLPFC and NAc of OUD subjects. Across brain regions, synaptic protein alterations in OUD subjects were primarily identified in glutamate, GABA, and circadian rhythm signaling. Using time-of-death (TOD) analyses, where the TOD of each subject is used as a time-point across a 24-hour cycle, we were able to map circadian-related changes associated with OUD in synaptic proteomes related to vesicle-mediated transport and membrane trafficking in the NAc and platelet derived growth factor receptor beta signaling in DLPFC.

58   Collectively, our findings lend further support for molecular rhythm disruptions in

59   synaptic signaling in the human brain as a key factor in opioid addiction.

Introduction

Rates of opioid use disorder (OUD) and deaths from opioid overdoses have continued to rise in the United States over the recent decade (2012-2022) [1]. Further understanding of the impact of long-term opioid use and opioid overdose on the human brain is important for considering new avenues of treatment and intervention. Investigating the molecular alterations in postmortem brains from subjects with OUD provide valuable insights into the neurobiological mechanisms and potential therapeutic targets for opioid addiction [2–8].

Recent work in postmortem brains of subjects with OUD from us [2, 3] and others [4, 6, 9, 10] have primarily focused on the dorsolateral prefrontal cortex (DLPFC) and nucleus accumbens (NAc), key brain regions involved in cognition and reward, which are significantly impacted in OUD [11–13]. An interplay between inflammation and synaptic remodeling associated with OUD in both DLPFC and NAc has been consistent findings across human postmortem brain studies [3, 9, 10]. Most recently, our work linked alterations in neuroinflammatory, dopaminergic, and GABAergic signaling in DLPFC and NAc of OUD subjects with significant disruptions in circadian rhythms of transcript expression [2]. In OUD, alterations in transcriptional rhythms were associated with sleep and circadian traits (*e.g.,* insomnia) [2], further supporting biological relationships between changes in sleep, circadian rhythms, and synaptic signaling in opioid addiction [14–22].

To date, much of the work has used transcriptomics approaches to investigate the molecular alterations in postmortem brains of subjects with OUD, with limited studies using proteomics [10]. Complementary to transcriptomics, proteomics can report

on functionally relevant information in cell bodies and neuropils, including synapses and

glial processes [23–25]. Recent advances have enabled reliable detection of numerous

peptides in synaptosomes isolated from postmortem human brains [24]. Deep profiling

of protein changes at the synaptic level may lead to functional insights into disease-

related mechanisms associated with psychiatric disorders.

Using quantitative mass spectrometry (MS) and tandem mass tags (TMT) [26,

27], we profiled both tissue homogenates and synaptosomes isolated from postmortem

DLPFC and NAc of unaffected subjects to compare tissue- and synapse-level protein

expression to subjects with OUD. Our approach allowed us to identify proteins that were

preferentially expressed, or enriched, specifically in synapses of DLPFC and NAc, and

assess synapse-specific alterations associated with OUD. Overall, pathways related to

neuroinflammation, and neurodegeneration were altered across brain regions in OUD,

accompanied by significant changes in proteins involved in GABAergic and

glutamatergic synaptic signaling. Notably, synaptic alterations associated with OUD

occurred in parallel to disruptions in pathways of circadian rhythm regulation. To further

explore circadian rhythms in human postmortem brain, we used time-of-death (TOD)

analysis to capture alterations associated with OUD in the diurnal variation of protein

expression in synaptosomes. In synapses, disruptions to molecular rhythms were

primarily related to endoplasmic reticulum functions, including protein trafficking and

vesicle-mediated transport, platelet derived growth factor receptor (PDGFR) signaling,

and proteins involved in GABA and glutamate synaptic signaling. Collectively, our

results suggest protein pathways related to neurodegeneration are altered at the tissue

level in DLPFC and NAc of OUD subjects, with implications for opioid-induced changes

in   circadian   regulation   of   inhibitory   and   excitatory   synaptic   signaling.

Materials and Methods

Human Subjects

Following consent from next-of-kin, postmortem brains were obtained during autopsies at Allegheny County (Pittsburgh, PA, USA; N=39) or Davidson Country (Nashville, TN, USA; N=1) Medical Examiner's Office. An independent committee of clinicians made consensus, lifetime DSM-IV diagnoses for each subject based on results from psychological autopsy, including structured family interviews, medical record reviews, and toxicological and neuropathological reports [28]. Similar procedures were used to confirm absence of lifetime psychiatric and neurological disorders in unaffected subjects. Procedures were approved by University of Pittsburgh Committee for Oversight of Research and Clinical Training Involving Decedents and Institutional Review Board for Biomedical Research. Cause of death in 19/20 OUD subjects was accidental, due to combined drug or opioid overdose. Accordingly, toxicology revealed that 19/20 OUD subjects had detectable opioids in their blood at time of death (Supplementary Table 1). Each OUD subject was matched with an unaffected comparison subject for sex, age, and postmortem interval (PMI) [2, 3]. Cause of death for unaffected subjects was either natural, accidental, or undetermined (Supplementary Table 1; opioids absent by blood toxicology). Cohorts differed by race (p=0.02) and brain pH (p=0.015, 0.2 pH units mean difference), and did not significantly differ in PMI, RNA integrity number (RIN), or TOD (p>0.25; Supplementary Table S1). TOD was determined from the death investigation report (Medical Examiner's Office). DLPFC (Brodmann Area 9) and NAc were anatomically identified and collected, as previously described [2, 3].

Brain Sample Preparation for Mass Spectrometry

Gray matter tissue (~20mg) from DLPFC and NAc were collected from fresh-frozen coronal tissue blocks via cryostat to minimize contamination from white matter and other subregions [29, 30]. Homogenate and synaptosome preparations were obtained using a variation of our enrichment protocol for postmortem human brain tissues [24, 31, 32] with SynPER reagent (ThermoFisher). From each sample, 10µg total protein (as measured by Micro BCA) was reduced, alkylated, and trypsin digested on S-Trap™ micro spin columns (ProtiFi). Subject pairs were randomly assigned to TMT blocks and labeled with TMTPro channels 1-10, with brain regions and preparations assigned to separate blocks [33]. Additional aliquots from each sample were used for a pooled control, digested separately with S-Trap Midi™ columns, divided then labeled with TMTPro channels 1 and 12. TMT labeled preparations from the same block were pooled with 10µg of the labeled pooled controls. The TMT labeled peptide pools were separated into eight fractions with the Pierce™ High pH Reversed-Phase Peptide Fractionation Kit (ThermoFisher Scientific), evaporated, and reconstituted in 20 µl 97% H2O, 3% ACN, 0.1% formic acid.

Mass Spectrometry

TMT labeled peptides (~1µg) were loaded onto a heated PepMap RSLC C18 column (2 µm, 100 angstrom, 75 µm × 50 cm; ThermoScientific), then eluted by gradients optimized for each high pH reverse-phase fraction [27]. Sample eluate was electrosprayed (2000 V) into an Orbitrap Eclipse Mass Spectrometer (MS;

ThermoFisher Scientific) for analysis. MS1 spectra were acquired at a resolving power

of 120,000. MS2 spectra were acquired in the Ion Trap with CID (35%) in centroid

mode. Real-time search (RTS) (max search time $=$ 34 s; max missed

cleavages $=$ 1; Xcorr $=$ 1; dCn $=$ 0.1; ppm $=$ 5) was used to select ions for SPS

for MS3. MS3 spectra were acquired in the Orbitrap with HCD (60%) with an isolation

window $=$ 0.7m/z and a resolving power of 60,000, and a max injection time of 400ms.

Data Processing

Raw MS files were processed in Proteome Discoverer (v. 2.5; ThermoFisher

Scientific). MS spectra were searched against the *Homo sapiens* SwissProt database.

SEQUEST search engine was used (enzyme=trypsin, maximum missed cleavage=2,

minimum peptide length=6, precursor tolerance=10ppm). Static modifications include

acetylation (N-term, +42.011 Da), Met-loss (N-term, -131.040 Da), Met-loss+Acetyl (N-

temr, -89.030 Da), and TMT labeling (N-term and K, +229.163 Da). Dynamic

modification, oxidation (M, +15.995 Da). PSMs were filtered by the Percolator node

(maximum Delta Cn=0.05, FDR =0.01). Reporter ion quantification was based on

intensity values with the following settings/filters: integration tolerance=20ppm,

method=most confident centroid, co-isolation threshold=100, and SPS mass matches

[%] threshold = 65. Peptide intensity values were normalized within and across TMT

plex runs with Normalization Mode=Total Peptide Amount and Scaling Mode=On All

Average in Proteome Discoverer. Peptides used to sum to protein measures were

determined using "Unique + Razor". Only proteins of high confidence were retained for

analyses and protein values were log2 transformed prior to analysis. The mass

spectrometry proteomics data have been deposited to the ProteomeXchange

Consortium via the PRIDE partner repository [34] with the dataset identifier PXD041333

and 10.6019/PXD041333.

Differential Expression of Proteins

Limma-Voom with covariate selection (TMT plex, sex, age, PMI, and peptide

expression) was used to detect differentially expressed (DE) proteins between OUD and

unaffected subjects in homogenate and synaptosome fractions [24]. Proteins were

considered DE if both $p \leq 0.05$ (unadjusted) and $\log_2$ fold-change (logFC) were greater

than or equal to ± 0.26 (20% change in expression), as previously used [3, 35, 36].

Over-representation pathway analyses were completed using clusterProfiler [37] and

ReactomePA [38] for DE proteins. Rank-rank hypergeometric overlap (RRHO) was

used to detect overlap of proteins expression changes between unaffected and OUD

subjects. To assess preferential enrichment of proteins in synapses, we compared

protein expression between homogenate and synaptosomes within unaffected subjects

and separately, within OUD subjects. Protein abundance in synaptosomes was

compared to abundance in homogenates to identify proteins with significantly greater

levels of expression in synaptosomes (Bonferroni-corrected $p < 0.05$). Proteins that

showed a difference of at least 1.25-fold in the synaptosome relative to homogenates

were classified as synapse-enriched proteins. All other proteins, by default, were

defined as non-enriched. Analysis of synaptosome differences between unaffected and

OUD subjects were limited to these enriched synaptic proteins, as described previously

[24].

Diurnal Rhythmicity Analysis

For each subject, TOD was converted to Zeitgeber Time (ZT) by using the times

of sunrise and sunset on the day of death (ZT0 is sunrise, negative ZTs reflect hours

prior to sunrise). Sinusoidal curves were fitted using nonlinear least-squares regression

with the coefficient of determination used as a proxy of goodness-of-fit (R^2). Estimates

of empirical p-values were determined using null distributions of R^2 generated from

1,000 TOD-randomized expression datasets. Diurnal rhythms of protein expression in

homogenates and synaptosomes were identified separately in unaffected and OUD

subjects. Rhythmic proteins were then compared between unaffected and OUD

subjects ($p<0.05$; Fisher's exact test). A complementary analysis determined which

proteins exhibited a significant change in rhythmicity between unaffected and OUD

subjects ($\Delta R^2 = R^2$ Unaffected - R^2 OUD). Change in rhythmicity analysis was restricted

to proteins that were significantly rhythmic in unaffected or OUD subjects. Proteins were

identified as significantly less rhythmic in OUD versus unaffected subjects if $\Delta R^2 > 0$ or

more rhythmic if $\Delta R^2 < 0$. Null distributions of ΔR^2 were permuted 1,000 times, whereby

the unaffected and OUD subjects were permuted independently to generate null

distributions (R^2 unaffected and R^2 OUD). Significantly less or more rhythmic in OUD

was calculated by comparing the ΔR^2 to the null ΔR^2 for unaffected and OUD subjects.

Differences in phase, amplitude, and base (mesor of fitted curve) were also calculated

and restricted to significantly rhythmic proteins from both unaffected and OUD subjects.

Pathway enrichments of rhythmic proteins were completed by Metascape (Gene

Ontology; Hallmark; Reactome; Canonical; metascape.org) [39].

Heatmaps represent the top rhythmic proteins in homogenates and

synaptosomes for each brain region, whereby each row represents a protein, and each

column represents a subject, ordered by ZT. Protein expression was Z-transformed and

ordered by phase (ZT of peak expression). Heatmaps were generated for: 1) top 200

rhythmic proteins in unaffected subjects; 2) top 200 rhythmic proteins identified in

unaffected subjects then plotted for OUD subjects to assess circadian proteomic

alterations in OUD; 3) top rhythmic proteins in OUD subjects; 4) top 200 rhythmic

proteins in OUD subjects then plotted for unaffected subjects.

Weighted Gene Co-Expression Network Analysis

To identify co-expression protein modules, we used weighted gene co-

expression network analysis (WGCNA) on protein expression from homogenates or

synaptosomes within each brain region and group [2, 3, 40–42]. Module differential

connectivity (MDC) analyses examined the impact of OUD on protein co-expression

network modules. For each module, the degree of connection was calculated in both

unaffected and OUD subjects. MDC was defined as the ratio of the protein co-

expression connections between unaffected and OUD subjects. To assess significance

of MDC, we permuted (1000 times) the samples and module members to generate null

distribution. When a module was built on unaffected subjects, a significant MDC greater

than 1 indicated a gain of connectivity of the modules in OUD, while a significant MDC

less than 1 indicated a loss of connectivity. Modules were tested for enrichment of DE

proteins and rhythmic proteins in unaffected and OUD. ARACne was used to identify

hub proteins in each module [43]. Proteins with an adjacency value higher than the 90[th]

quantile were considered neighbors. Hub proteins were identified with the highest

number of N-hob neighborhood nodes (NHNN) than the average. Hub proteins specific

to unaffected and OUD subjects were considered disease-specific hub proteins. Hub

gene networks were plotted using Cytoscape (3.9.1). We then identified enriched

pathways of synapse specific protein co-expression networks using SynGo [44].

Results

Brain Region-specific Protein Alterations in Tissue Homogenates and

Synaptosomes associated with OUD in NAc and DLPFC

Between unaffected and OUD subjects, we investigated protein expression

differences in tissue homogenates of DLPFC and NAc and isolated synaptosomes from

each brain region. We identified 43 DE ($p<0.05$ and logFC±0.26) proteins (14

upregulated and 29 downregulated; Fig. 1a, Supplementary Tables S2, S3) in the NAc

of subjects with OUD. In DLPFC, we identified 55 DE proteins, with most being

upregulated in OUD (46 upregulated and 9 downregulated; Fig. 1b, Supplementary

Tables S4, S5). In synaptosomes, we found 56 DE proteins (17 upregulated and 39

downregulated) in NAc (Fig. 1e, Supplementary Table S7, S8) and 161 DE proteins (80

upregulated and 81 downregulated) in DLPFC (Fig. 1f, Supplementary Table S9, S10).

To investigate the possible overlap of protein changes between brain regions, we

compared protein expression between NAc and DLPFC of OUD subjects using the

threshold-free approach, RRHO [45]. Overall, there was weak concordance of proteins

that were upregulated or downregulated between brain regions of OUD subjects (Fig.

1c), despite identifying overlapping proteins between these brain regions (Fig. 1d). In

contrast, we previously reported remarkable concordance between NAc and DLPFC in

both upregulated and downregulated transcripts associated with OUD of the same

subjects investigated here [2, 3]. In synaptosomes, we identified no significant overlap

of protein changes between brain regions (Fig. 1g,h). Therefore, protein alterations in

OUD were unique within brain region, and together with our previous results [3], may

reflect the impact of opioids and other factors on brain region-specific, post-

transcriptional and translational processing.

Alterations in Inflammatory and Neurodegeneration-related Pathways associated

with OUD in NAc and DLPFC Homogenates

To further define the biological significance of protein changes in OUD, we

conducted pathway enrichment analysis on DE proteins from tissue homogenates of

NAc and DLPFC (Fig. 2, Supplementary Table S6). In NAc homogenates, enrichment

analyses identified pathways related to immune modulatory pathways, including various

types of infection (*e.g.,* KRT14,16,17 [46]; HLA-B [47]) and leukocyte migration (*e.g.,*

JAM2 [48]; PLCG1 [49]). Neurodegeneration-related pathways were also enriched in

NAc homogenates (Fig. 2a; *e.g.,* CHRM1 [50, 51]; MAPT [52]; SQSTM1 [53]). Similarly,

several pathways related to neurodegeneration were highly enriched in DLPFC

homogenates from OUD subjects, such as Alzheimer (43 proteins; *e.g.,* SLC25A6 [54];

CHRM3 [55]; MTOR [56]; ITPR2 [57]), Huntington (38 proteins; *e.g.,* PSMD11 [58];

ADRM1 [59]; SOD1 [60]), Parkinson (41 proteins; PARK7 [61]; HTRA2 [62]), and prion

(42 proteins; *e.g.,* RYR2 [63]) disease-related pathways (Fig. 2b). Other pathways

enriched in DLPFC homogenates were related to cell stress including oxidative

phosphorylation and reactive oxygen species (Fig. 2b). Together, our findings highlight

a possible cascade of immune activation, diminished cellular health, and initiation of

neurodegenerative processes in human brain that are associated with OUD.

Differential Enrichment between Tissue Homogenates and Synaptosomes

Highlights Altered GABAergic and Glutamatergic Synaptic Signaling in NAc and

DLPFC associated with OUD

We investigated protein expression changes preferentially enriched in

synaptosomes to identify synapse-specific proteome alterations associated with OUD.

We identified proteins primarily expressed in homogenates and preferentially expressed

in synaptosomes within each brain region (Supplementary Tables S7-S12). Synapse

enriched proteins were then used for DE and pathway analyses comparing unaffected

and OUD subjects. Synaptosomes isolated from NAc were enriched for several

pathways involved in mitochondrial and endoplasmic reticulum functions (proximal

tubule bicarbonate reclamation, carbohydrate digestion and absorption), along with

second messenger signaling cascades (cGMP-PKG and cAMP: *e.g.,* GNAI2,3;

GABRA2; ACY1; Fig. 2a, Supplementary Table S6). Other pathways were specifically

enriched in DLPFC synaptosomes from OUD subjects, which included ribosome,

relaxin, and Rap1 signaling (Fig. 2b). OUD-associated alterations in serotonergic

synapses (*e.g.,* GNAO1; GNAS; GNAI2,3, GNB1,4) were also unique to DLPFC (Fig.

2b, Supplementary Table S6).

Additionally, we found similarly altered pathways in synaptosomes between brain

regions. The major pathways enriched across NAc and DLPFC synaptosomes included

morphine addiction and GABAergic signaling (*e.g.,* GNG3, GABBR2, GNB1 in NAc; and

GNAO1, GNB4, SLC6A11 in DLPFC), and glutamatergic synaptic signaling (*e.g.,*

GNG3, SLC1A3, and ADCY1 in NAc; and GNAO1, GNAS, and GNB14 in DLPFC; Fig.

2a,b).

Further investigation of the individual proteins enriched in synaptosomes

revealed THY1 [64, 65], CACNA2D1 [66–70], SLC3A2 [70], GSK3B [71–73], LSAMP

[73, 74], LY6H [75, 76], and NCAM1 [77–81], as the top synaptic proteins altered in NAc

of OUD subjects (Fig. 3). In DLPFC, proteins altered in OUD and enriched in

synaptosomes included RALB [82], CADM3 [83], GPC1 [84], MRAS, CNTFR [85],

IGLON5, IST1, and PLXNA4 [86, 87] (Fig. 3). Other synapse enriched proteins in

DLPFC were like those enriched in NAc (*e.g.,* GSK3B, LSAMP, NCAM1, CACNA2D1,

SLC3A2; Fig. 3). Consistent with our pathway enrichments in synaptosomes of OUD

subjects (Fig. 2), many of the synaptic proteins in both the NAc and DLPFC are involved

in GABAergic and glutamatergic synaptic functions. For example, each of the synaptic

proteins, CACNA2D1, SLC3A2, and GSK3B, altered in both brain regions, are known to

dynamically regulate the activity of glutamatergic-dependent synaptic activity and

plasticity [69, 70, 72].

Altered Circadian Rhythm Signaling associated with OUD in Synaptic Proteomes

of NAc and DLPFC

Previously, we reported transcriptional rhythm disruptions in NAc and DLPFC of

subjects with OUD [2]. In support of this, a significantly enriched pathway in DE synaptic

proteins in both brain regions were circadian entrainment (Fig. 2). Several preferentially

enriched synaptic proteins have integral roles in the regulation of circadian rhythms

(*e.g.,* CACNA2D1, GSK3B, NCAM1, SYT3 [88]; Fig. 3). To further explore potential

circadian rhythms in brain proteomes, we conducted TOD analysis on protein

expression profiles from homogenate (Supplementary Fig. S2, Supplementary Fig. S1,

Supplementary Tables S13-20) and synaptosomes (Fig. 4a and 4b) of each brain

region. In synaptosomes, we identified 300 and 77 significantly rhythmic proteins in NAc

of unaffected and OUD subjects, respectively (Fig. 4d). In DLPFC synaptosomes, we

identified 123 rhythmic proteins in unaffected and 182 rhythm proteins in OUD (Fig. 4h).

Top rhythmic synaptic proteins in unaffected subjects were significantly disrupted in

subjects with OUD across both brain regions in synaptosomes (Figs. 4a,e;

Supplementary Tables S17-S20 and S23-S24) and homogenates (Supplementary Fig

S2; Supplementary Tables S13-S16 and S21-S22). Vice-versa, we also found that

proteins were only rhythmic in OUD compared to unaffected subjects. Consequently,

top rhythmic proteins in OUD were different from rhythmic proteins in unaffected

subjects in synaptosomes (Fig 4a, NAc; Fig 4e, DLPFC; Supplementary Tables S17-

S20 and S23-S24) and homogenates (Supplementary Fig. S1; Supplementary Tables

S13-S16 and S21-S22). These findings highlighted a gain in rhythmicity in OUD

compared to unaffected subjects.

In NAc synaptosomes, the top rhythmic proteins of unaffected subjects included

the GABA transporter, GAT1 [89, 90] (Fig. 4b). In OUD, top rhythmic proteins included

neurofilament proteins (NEFL, NEFM [91]; Fig. 4b), consistent with enrichment of actin

and filament cytoskeleton pathways (Fig. 4c). Neuropeptide signaling pathways were

also enriched among rhythmic proteins in NAc of OUD subjects (Fig. 4c), a pathway that

involves upstream initiation by G protein-coupled receptor binding to opioid receptors

and downstream activation of intracellular pathways. Proteins encoding several opioid

peptides were rhythmic in synaptosomes of OUD subjects, including PDYN

(prodynorphin) and PENK (proenkephalin) [92, 93], and other proteins involved in opioid

receptor signaling (SCG5 [94] and PCSK1N [95]). SCG5 and PCSK1N are also

secretory proteins involved in preventing the aggregation of proteins involved in

neurodegenerative disorders.

In DLPFC synaptosomes, top rhythmic proteins of unaffected subjects included

brain abundant membrane attached signal protein 1, BASP1 (Fig.4f). Moreover, CHID1

[96] and SPCS2 [97] were among the top rhythmic synaptic proteins in DLPFC of OUD

subjects (Fig. 4e,f), both of which are linked to cytoskeletal pathology in

neurodegenerative disorders. Enrichment analyses of rhythmic synaptic proteins in

DLPFC in OUD subjects revealed pathways primarily associated with Golgi and

endoplasmic reticulum processing, autophagy, glycosylation, and synaptic vesicle

transport (Fig. 4g). Disrupted endoplasmic reticulum signaling, along with changes in

the process of protein glycosylation, may reflect the impact of opioids on local protein

translation and trafficking at postsynaptic sites, including dendritic spines, regulating

synaptic plasticity [98].

In NAc homogenates of OUD subjects, top enriched pathways included opioid

and synaptic signaling (Supplementary Fig. S2). DARRP32, a phosphoprotein critically

involved in dopaminergic synaptic functions, was among the top rhythmic proteins in

OUD (Supplementary Fig. S2). Rhythmic proteins specifically in NAc of OUD subjects

were associated with pathways in various vesicle endocytotic processes (*e.g.,*

endocytosis, synaptic vesicle cycle, and clathrin-mediated endocytosis; Supplementary

Fig. S2). In DLPFC, top rhythmic proteins in OUD subjects included the neurosecretory

protein, VGF [99], and the putative negative regulator of cannabinoid receptor 1 activity,

CNRIP1 (Supplementary Fig. S2). Top rhythmic pathways in OUD were related to glial

cell development, mitochondrion organization, and various components of lipid and

pyruvate metabolism (Supplementary Fig. S2). Rhythmicity in transcript abundance [2]

was more prevalent compared to rhythms in protein expression in NAc and DLPFC

(Supplementary Fig. S2). Notably, we observed very little overlap between rhythmic

transcripts and rhythmic synaptic proteins (Fig. 4d, NAc; Fig 4h, DLPFC) and

homogenates (Supplementary Figure S2) in either brain region of unaffected or OUD

subjects.

Altered Diurnal Rhythms of the Synaptic Proteome in DLPFC and NAc associated

with OUD

Rhythmic proteins were largely distinct between unaffected and OUD subjects

across brain regions. Given this, we examined whether proteins lost or gained rhythms

between unaffected and OUD subjects within each brain region and sample preparation

(Homogenates: Supplementary Figs. S1, S2, S3 and Supplementary Tables S21, S22;

Synaptosomes: NAc, Fig. 5a; DLPFC, Fig. 5d; Supplementary Tables S23, S24). In NAc

homogenates, 5 proteins lost rhythms and 19 proteins gained rhythms, primarily

involved in protein translation (Supplementary Fig. S3a, S3c, Supplementary Table

S21). Only 10 proteins were significantly altered in rhythmic expression in DLPFC

homogenates of OUD subjects, with 3 proteins losing rhythms and 7 proteins gaining

rhythms (Supplementary Fig. S3b). These proteins were enriched for pathways

involved in lipid homeostasis and neurotoxicity, potentially related to neurodegenerative

processes (Supplementary Fig. 3d; Supplementary table S22).

In synaptosomes, 23 proteins lost rhythmicity in NAc of OUD subjects (*e.g.,*

TRAPPC10, EHBP1, ZW10; Fig. 5a,b; Supplemental Table S23). Loss of protein

rhythms in NAc were enriched for endoplasmic reticulum to Golgi vesicle-mediated

transport pathways (Fig. 5c). In DLPFC, we found 7 synaptic proteins that lost

rhythmicity in OUD and 11 synaptic proteins that gained rhythmicity (Fig. 5d; *e.g.,*

USO1, NSDHL, PDE1C [100], Fig. 5e). Proteins that gained rhythmicity (*e.g.,* USO1,

NSDHL, PDE1C, CHID1, GRID1) were involved in chemical synaptic transmission and

organelle localization, while proteins that lost rhythmicity (*e.g.,* FAM177A1, ABI2,

TPD52L2, EIF2AK2, PRKCD) were involved in protein phosphorylation (Fig. 5f;

Supplemental Table S24). Processed together, altered rhythmicity of synaptic proteins

in DLPFC were associated with platelet-derived growth factor receptor beta (PDGFR-B)

signaling (Fig. 5f), implicated in neuroprotection in response to elevated glutamatergic

activity [101] and involved in opioid reward [102, 103]. Specifically, the 11 synaptic

proteins gaining rhythmicity in DLPFC were associated with modulation of chemical

synaptic transmission and organelle localization, while the 7 proteins losing rhythmicity

were associated with protein phosphorylation (Supplementary Fig. S4; Supplemental

Table S24). Collectively, our findings suggest circadian rhythm regulation of the

synaptic proteome that is significantly altered in a brain region-specific manner in OUD.

Further, circadian disruption of synaptic functions in OUD are associated with changes

in endoplasmic reticulum-mediated local protein translation, trafficking, and vesicle

endocytosis at the synapse, in addition to processes involved in neuroprotection and

neurodegeneration.

Protein Co-Expression Modules Identify Brain Region-Specific Alterations in

Circadian Rhythm Signaling and Synaptic Functions associated with OUD

Using weighted co-expression analyses, we identified highly connected protein

expression modules specific to NAc and DLPFC in homogenates (Supplementary Figs.

S5, S7, S9; Supplementary Table S25) and synaptosomes (Supplementary Figs. S6,

S8, S10; Supplementary Table S25). To further narrow modules relevant for OUD

associated protein alterations, we used MDC analysis, which examined the overall

connectivity of protein co-expression and network structure between unaffected and

OUD subjects (Supplementary Table S26).

Many modules in unaffected subjects displayed a loss of connectivity in NAc and

DLPFC homogenates (Supplementary Fig. S11; Supplementary Table S26) and

synaptosomes (Fig. 6). In synaptosomes, connectivity was significantly lost in 13/15

modules in the NAc of OUD subjects (Fig. 6a), along with 10/26 modules in DLPFC

(Fig. 6b). Loss of connectivity in protein co-expression modules indicates significant

dispersion of protein network structure related to OUD.

We then investigated whether modules with changes in connectivity were

enriched for hub proteins specific to unaffected and OUD subjects, and circadian

rhythms. Focusing on the modules that lost connectivity in OUD, we found significant

enrichment of both hub proteins specific to and/or rhythmic in unaffected subjects (Fig.

6). In NAc, there were three modules which lost connectivity in OUD that were

significantly enriched for hub proteins (turquoise, green-yellow, and green; Fig. 6a). In

the green module, proteins are associated with synaptic plasticity, dendritic spine

formation, and axon guidance (*e.g.,* ROCK2 [104], DPYSL5 [105, 106], PLCG1 [107,

108], GDAP1; Fig. 6a). In the green-yellow module, several of the rhythmic hub proteins

were related to glutamatergic (GRIA2 [109, 110]), GABAergic (GABRA2 [111–113]),

and calcium signaling (CAMK2B, CALB2) (Fig. 6a). Additionally, other hub proteins are

involved in calcium signaling, membrane potential, glutamatergic signaling, and

exocytosis of synaptic vesicles (rhythmic hub SCAMP5; hubs SYNGR3 and LRRC7;

Fig. 6a). Overall, protein networks that lost connectivity in the NAc of OUD subjects

were enriched for hub proteins that exhibited rhythmicity in unaffected subjects and

involved in GABAergic and glutamatergic neurotransmission (Fig. 6c).

In DLPFC, we identified two modules that lost connectivity in OUD and with

significant hub protein enrichments (Fig. 6b). The black module contained several highly

connected hub proteins involved in ECM formation (LAMC1 [114]), presynaptic

dopamine release (PLPP3 [115]), neuroprotection (NCEH1 [116], BCL2CL13),

inflammation (LRRFIP2 [117]), and pronociceptive receptor signaling in pain (PIP5KL1

[118]). The green module contained several rhythmic hub proteins involved neuronal

protection (NME3, SPHK2, RAB13), synaptic functions (MCTP1, NTNG2, GNG5), and

transposon activity in the brain (PGBD5 [119] (Fig. 6b). Enrichment analysis primarily

highlighted synaptic functions of vesicle cycling and structure (Fig. 6d). Collectively, our

analyses lend further support for a role of circadian rhythms in synaptic signaling within

human brain and suggest significant disruptions of circadian-dependent regulation of

synaptic functions in OUD.

Discussion

Our previous transcriptomic findings implicate circadian rhythm disruption in

proinflammatory signaling and synaptic remodeling in human NAc and DLPFC in OUD

[2, 3]. We extend these findings here by showing significant alterations in circadian

rhythm, GABAergic, and glutamatergic synaptic processes in both regions of OUD

subjects. Using high-density profiling of peptides in tissue homogenates and

synaptosomes within brain regions from the same subjects, we were able to identify and

parse significant OUD-associated changes in synaptic proteomes specific to and

common to NAc and DLPFC. In NAc synaptosomes specifically, pathways were

involved in inflammatory, mitochondria, and metabolic signaling. In DLPFC

synaptosomes from OUD subjects, pathways were mainly related to significant

alterations in inflammatory signaling and serotonergic, dopaminergic, cholinergic,

oxytocin neurotransmission. In both brain regions, pathways related to

neurodegeneration, GABAergic and glutamatergic synapses, and circadian rhythms

were associated with OUD, further supporting a role for circadian rhythms in synaptic

signaling in opioid addiction.

Alterations in protein pathways related to neurodegeneration in OUD subjects

were common in both NAc and DLPFC homogenates and synaptosomes. The

relationship between chronic opioid use and increased risk for neurodegenerative

disorders is complex. Several studies have reported increased risk of dementia and

higher incidence of neurodegenerative disorders in people with heavy, chronic opioid

use [120–122]. Chronic opioid use and dysfunction of the opioid system in the brain

may contribute to neuroinflammation and the hyperphosphorylation of tau, a key protein

involved in the pathogenesis of Alzheimer's disease [122]. For example, dysfunction of

opioid receptor signaling leads to increased amyloid beta expression and deposition

[120]. Accumulation of amyloid beta is involved in the neurotoxicity observed early in the

development of Alzheimer's disease.

Consistent with the emergence of early pathogenesis associated with

neurodegenerative disorders, several studies have shown significant elevation of early markers of neuropathology in Alzheimer's disease in subjects who chronically used heroin, including hyperphosphorylated tau and amyloid beta [122]. Neurodegenerative markers in the brain were positively correlated with neuroinflammatory markers and microglial activation. Intriguingly, levels of the enzyme that phosphorylates tau, GSK-3B, were also increased in the same subjects [122]. We identified GSK-3B as a synaptic-enriched protein significantly upregulated in NAc and to a lesser degree in DLPFC of OUD subjects. Other related proteins were changed in both regions, including reduced expression of CBR1, BIN1, and VPS29 in DLPFC synaptosomes and reduced expression of SIRT2 in NAc synaptosomes. CBR1 is usually elevated in response to neuroinflammation and neuronal injury, ultimately facilitating neuroprotection [123]. Reduced BIN1 expression and function is a known risk factor for late-onset Alzheimer's disease, having critical roles in presynaptic release of glutamate and the facilitation of learning and memory [124]. Further, VPS29 is involved in synaptic survival [125]. OUD-associated downregulation of each of these proteins in DLPFC may reflect the initiation of molecular signaling cascades resembling early pathogenesis of Alzheimer's disease. In addition, SIRT2 downregulation in NAc synapses could lead to augmented activity at excitatory synapses via dysfunction of AMPA and NMDA glutamate receptors [126, 126, 127]. Accumulation of phosphorylated tau and other markers of neurodegenerative diseases, accompanied by elevated neuroinflammation in the brain, likely have functional impacts on synaptic processes in NAc and DLPFC in OUD. In OUD, several synaptic proteins were enriched in both NAc and DLPFC that link neurotoxicity to augmented glutamatergic neurotransmission–CNTFR, PLXNA4, and SLC3A2. Although

speculative, alterations in synaptic processes we highlighted here in subjects with OUD,

particularly GABAergic and glutamatergic neurotransmission, may be associated with

cognitive, mood, and reward dysfunction associated with opioid addiction.

An imbalance of GABAergic and glutamatergic signaling in NAc and DLPFC has

been linked to OUD and other psychiatric disorders. Between NAc and DLPFC of OUD

subjects, our analyses identified several of the most impacted pathways in

synaptosomes including GABAergic and glutamatergic neurotransmission. Several

notable proteins involved in glutamatergic excitatory synapses were upregulated in

DLPFC synaptosomes–GRIN2A (glutamate ionotropic receptor NMDA type subunit 2A),

GRIN2B (glutamate ionotropic receptor NMDA type subunit 2A), GRM1 (glutamate

metabotropic receptor 1), and SHANK2 (SH3 and multiple ankyrin repeat domains 2).

GRIN proteins are subunits of the NMDA receptor complex, both of which are intimately

involved in opioid-induced excitatory synaptic plasticity, having integral roles in opioid

withdrawal and craving [128–130]. Synaptic elevation of SHANK2 may elevate NMDA

receptor activity and enhance bursting of parvalbumin-positive neurons in DLPFC [131].

Enhancing NMDA receptor activation impairs synaptic plasticity and leads to cognitive

impairments, potentially involved in OUD. We identified other synapse-specific proteins

in the NAc involved in NMDA-dependent glutamate signaling, such as FLOT2

(upregulated), HDAC11 (downregulated), and SLC18B1 (downregulated). Notably,

reduced HDAC11 expression in OUD and function may impede synaptic plasticity [132].

CACNA2D1, SLC3A2, and GSK3B, among others, were also enriched in both NAc and

DLPFC of OUD subjects, each of which have dynamic roles in the regulation of

glutamatergic-dependent synaptic plasticity. In addition, proteins involved in GABAergic

processes were also significantly altered in OUD. For example, GAD2 was reduced in

DLPFC synaptosomes in subjects with OUD. GAD2 is associated with synaptic vesicle

response during intense neuronal activity to release GABA into the synaptic cleft [133],

consistent with dysfunctional inhibitory neurotransmission in opioid addiction.

Additionally, we identified proteins specifically altered in NAc and DLPFC of OUD

subjects. For example, LSAMP and LY6H were more enriched in NAc synaptosomes,

both of which are involved in dendritic spine formation in response to excitatory inputs.

Other proteins related to neuroinflammation were upregulated in NAc (*e.g.,* HLA-B,

GSTT1, S100A9). In DLPFC, synaptic enriched proteins included IGLON5, a neuronal

adhesion molecule linked to neurological disorder characterized by sleep disorders and

cognitive impairments; EPHA4 [134], a receptor tyrosine kinase that modulates aberrant

synaptic functions in response to neuronal injury and neuroinflammation; and GPC1, a

glycoprotein involved in extracellular matrix (ECM) formation. Several proteins with the

highest fold-change in OUD subjects included keratin cytoskeletal, ECM-related

proteins in NAc [135]. Reduced expression and function of keratin proteins and

neurofilaments are associated with neuropsychiatric disorders [121, 122], including

stress-related disorders and substance use disorders [46]. The co-occurrence of

changes in proteins involved in the activation of neuroinflammatory processes,

degradation of ECM, and synaptic remodeling corroborates our previous transcriptomic

findings in opioid addiction [3].

Pathway analysis implicated circadian rhythm disruption primarily in

synaptosomes of OUD subjects. However, protein expression rhythms were markedly

disrupted in both homogenates and synaptosomes from NAc and DLPFC. For example,

DARPP-32, a protein highly expressed in striatal medium spiny neurons [136–138], was

highly rhythmic in NAc homogenates, suggesting opioid-induced circadian

reprogramming of molecular cascades in the striatum [139]. Notably, opioid signaling

was a top pathway that was rhythmic in NAc homogenates, where rhythms in opioid

neurotransmission may contribute to craving and relapse [140, 141]. Proteins that lost

rhythmicity in the NAc of OUD subjects were involved in membrane trafficking and

endoplasmic reticulum to Golgi vesicle-mediated transport (*e.g.,* TRAPPC5,

TRAPPC10, EHBP1, ZW10). Both TRAPPC5 and TRAPPC10 are transmembrane

proteins of the cis-Golgi complex that support vesicular transport from endoplasmic

reticulum to Golgi. TRAPP10 interacts with TRAPPC9 to form TRAPP II core proteins.

TRAPPC9/10 proteins regulate the endocytic receptor recycling of dopamine 1 and 2

receptors in postsynaptic striatal medium spiny neurons [142], integral to regulation of

medium spiny neuron physiology and motivated behaviors. Local translation at synaptic

sites may also occur through activity-dependent transport [143]. Glycoproteins THY1

and NCAM1 [144] were highly rhythmic in DLPFC homogenates. An area of future work

could explore the impact of opioids on protein glycosylation of synaptic proteins and

their impact of synaptic vesicle loading and release, particularly glutamate receptor

turnover and activation [145] in response to neuroinflammation [146, 147]. In addition,

PDGFR-B signaling was among the top pathways from altered rhythms in protein

expression of DLPFC synaptosomes in OUD. Importantly, PDGF-dependent signaling is

involved in numerous opioid actions, including tolerance and reward [148]. PDGF is a

receptor tyrosine kinase, suggesting interactions between tyrosine kinases and opioid

receptors may directly modulate glutamate receptor activity [149] depending on their

circadian regulation.

Co-expression analyses revealed further insights into the role of circadian rhythm

regulation of synaptic dysfunction in OUD. We identified several protein modules that

were enriched for rhythmic proteins and hub proteins involved in synaptic processes in

unaffected subjects (*e.g.,* CAMK2B, GSK-3B, ROCK2, PLCG1, SCAMP5, SYT17).

Each of which are involved in dendritic spine formation and synaptic plasticity [104].

Every highly connected module in unaffected subjects in NAc and DLPFC lost

connectivity in OUD subjects. Loss of connectivity in these protein networks in OUD

could resemble a lack of coordinated protein signaling at the synapse that is modulated

by circadian rhythms. OUD-associated upregulation of GSK-3B in synaptosomes of NAc

and DLPFC likely leads to significant molecular rhythm disruptions in synapses, with

potential consequences on activity-dependent synaptic plasticity, and thus, may be a

lynchpin in opioid and circadian actions at the synapse.

Overall, our findings implicate circadian rhythm disruption in altered functions at

GABAergic and glutamatergic synapses in both NAc and DLPFC. Considering the role

of sleep and circadian rhythms in opioid reward, craving, and relapse, treatments that

target circadian pathways may be an effective therapeutic strategy for OUD.

Acknowledgments

Thank you to the staff and technicians who work diligently as part of the Brain

Tissue Donation Program at the University of Pittsburgh. Postmortem human brain

tissue was provided by the University of Pittsburgh Brain Tissue Donation Program and

the National Institutes of Health NeuroBioBank at the University of Pittsburgh.

Author Contributions

MLS and RWL obtained funding for the project. JRG, DAL, MLS, MLM, and RWL

designed and coordinated the study. SP, RS, MAS, MAH, SMK, JRG, and MLM

processed samples for proteomics. SP, XX, ZF, GCT, AY, MLM, and RWL conducted

analyses and data interpretation. SP, XX, MLS, MLM, and RWL wrote the manuscript.

All authors read and approved the final version of the manuscript.

Funding

The research reported in this article was supported by: Hamilton Family Prize for

Basic Neuroscience Research in Psychiatry at the University of Pittsburgh School of

Medicine (RWL); National Heart, Lung, and Blood Institute (R01HL150432, RWL); and

National Institute on Drug Abuse (R01DA051390, MLS and RWL).

Competing Interests

The authors report no competing interests.

References

1. Abuse NI on D. Drug Overdose Death Rates. Natl Inst Drug Abuse. 2023. https://nida.nih.gov/research-topics/trends-statistics/overdose-death-rates. Accessed 23 March 2023.

2. Xue X, Zong W, Glausier JR, Kim S-M, Shelton MA, Phan BN, et al. Molecular rhythm alterations in prefrontal cortex and nucleus accumbens associated with opioid use disorder. Transl Psychiatry. 2022;12:1–13.

3. Seney ML, Kim S-M, Glausier JR, Hildebrand MA, Xue X, Zong W, et al. Transcriptional Alterations in Dorsolateral Prefrontal Cortex and Nucleus Accumbens Implicate Neuroinflammation and Synaptic Remodeling in Opioid Use Disorder. Biol Psychiatry. 2021;90:550–562.

4. Nagamatsu ST, Rompala G, Hurd YL, Núñez-Rios DL, Montalvo-Ortiz JL, Traumatic Stress Brain Research Group. CpH methylome analysis in human cortical neurons identifies novel gene pathways and drug targets for opioid use disorder. Front Psychiatry. 2022;13:1078894.

5. Jacobs MM, Ökvist A, Horvath M, Keller E, Bannon MJ, Morgello S, et al. Dopamine receptor D1 and postsynaptic density gene variants associate with opiate abuse and striatal expression levels. Mol Psychiatry. 2013;18:1205–1210.

6. Sillivan SE, Whittard JD, Jacobs MM, Ren Y, Mazloom AR, Caputi FF, et al. ELK1 transcription factor linked to dysregulated striatal mu opioid receptor signaling network and OPRM1 polymorphism in human heroin abusers. Biol Psychiatry. 2013;74:511–519.

7. Brown TG, Xu J, Hurd YL, Pan Y-X. Dysregulated expression of the alternatively

spliced variant mRNAs of the mu opioid receptor gene, OPRM1, in the medial

prefrontal cortex of male human heroin abusers and heroin self-administering

male rats. J Neurosci Res. 2022;100:35–47.

8.   Valeri J, O'Donovan SM, Wang W, Sinclair D, Bollavarapu R, Gisabella B, et al.

Altered expression of somatostatin signaling molecules and clock genes in the

hippocampus of subjects with substance use disorder. Front Neurosci.

2022;16:903941.

9.   Grimm SL, Mendez EF, Stertz L, Meyer TD, Fries GR, Gandhi T, et al. MicroRNA-

mRNA networks are dysregulated in opioid use disorder postmortem brain:

Further evidence for opioid-induced neurovascular alterations. Front Psychiatry.

2022;13:1025346.

10.   Mendez EF, Wei H, Hu R, Stertz L, Fries GR, Wu X, et al. Angiogenic gene

networks are dysregulated in opioid use disorder: evidence from multi-omics and

imaging of postmortem human brain. Mol Psychiatry. 2021;26:7803–7812.

11.   Baldo BA. Prefrontal Cortical Opioids and Dysregulated Motivation: A Network

Hypothesis. Trends Neurosci. 2016;39:366–377.

12.   Adinoff B, Rilling LM, Williams MJ, Schreffler E, Schepis TS, Rosvall T, et al.

Impulsivity, neural deficits, and the addictions: the 'oops' factor in relapse. J

Addict Dis. 2007;26 Suppl 1:25–39.

13.   Koob GF. Neurobiology of Opioid Addiction: Opponent Process, Hyperkatifeia,

and Negative Reinforcement. Biol Psychiatry. 2020;87:44–53.

14.   Mahfoud Y, Talih F, Streem D, Budur K. Sleep disorders in substance abusers:

how common are they? Psychiatry Edgmont Pa Townsh. 2009;6:38–42.

15. Huhn AS, Ellis JD, Dunn KE, Sholler DJ, Tabaschek P, Burns R, et al. Patient-
reported sleep outcomes in randomized-controlled trials in persons with
substance use disorders: A systematic review. Drug Alcohol Depend.
2022;237:109508.

16. Huhn AS, Finan PH, Gamaldo CE, Hammond AS, Umbricht A, Bergeria CL, et al.
Suvorexant ameliorated sleep disturbance, opioid withdrawal, and craving during
a buprenorphine taper. Sci Transl Med. 2022;14:eabn8238.

17. Eacret D, Veasey SC, Blendy JA. Bidirectional Relationship between Opioids and
Disrupted Sleep: Putative Mechanisms. Mol Pharmacol. 2020;98:445–453.

18. Eacret D, Manduchi E, Noreck J, Tyner E, Fenik P, Dunn AD, et al. Mu-opioid
receptor-expressing neurons in the paraventricular thalamus modulate chronic
morphine-induced wake alterations. Transl Psychiatry. 2023;13:78.

19. Oyefeso A, Sedgwick P, Ghodse H. Subjective sleep-wake parameters in
treatment-seeking opiate addicts. Drug Alcohol Depend. 1997;48:9–16.

20. Stinus L, Robert C, Karasinski P, Limoge A. Continuous quantitative monitoring of
spontaneous opiate withdrawal: locomotor activity and sleep disorders. Pharmacol
Biochem Behav. 1998;59:83–89.

21. Li S, Shi J, Epstein DH, Wang X, Zhang X, Bao Y, et al. Circadian alteration in
neurobiology during 30 days of abstinence in heroin users. Biol Psychiatry.
2009;65:905–912.

22. Cao M, Javaheri S. Effects of Chronic Opioid Use on Sleep and Wake. Sleep Med
Clin. 2018;13:271–281.

23. McClatchy DB, Liao L, Park SK, Venable JD, Yates JR. Quantification of the

synaptosomal proteome of the rat cerebellum during post-natal development.

Genome Res. 2007;17:1378–1388.

24. MacDonald ML, Garver M, Newman J, Sun Z, Kannarkat J, Salisbury R, et al.

Synaptic Proteome Alterations in the Primary Auditory Cortex of Individuals With

Schizophrenia. JAMA Psychiatry. 2020;77:86–95.

25. Carney KE, Milanese M, van Nierop P, Li KW, Oliet SHR, Smit AB, et al.

Proteomic analysis of gliosomes from mouse brain: identification and investigation

of glial membrane proteins. J Proteome Res. 2014;13:5918–5927.

26. Dumrongprechachan V, Salisbury RB, Butler L, MacDonald ML, Kozorovitskiy Y.

Dynamic proteomic and phosphoproteomic atlas of corticostriatal axons in

neurodevelopment. ELife. 2022;11:e78847.

27. Dumrongprechachan V, Salisbury RB, Soto G, Kumar M, MacDonald ML,

Kozorovitskiy Y. Cell-type and subcellular compartment-specific APEX2 proximity

labeling reveals activity-dependent nuclear proteome dynamics in the striatum.

Nat Commun. 2021;12:4855.

28. Glausier JR, Kelly MA, Salem S, Chen K, Lewis DA. Proxy measures of

premortem cognitive aptitude in postmortem subjects with schizophrenia. Psychol

Med. 2020;50:507–514.

29. Volk DW, Matsubara T, Li S, Sengupta EJ, Georgiev D, Minabe Y, et al. Deficits in

Transcriptional Regulators of Cortical Parvalbumin Neurons in Schizophrenia. Am

J Psychiatry. 2012;169:1082–1091.

30. Hashimoto T, Bazmi HH, Mirnics K, Wu Q, Sampson AR, Lewis DA. Conserved

regional patterns of GABA-related transcript expression in the neocortex of

subjects with schizophrenia. Am J Psychiatry. 2008;165:479–489.

31.   Hahn C-G, Banerjee A, Macdonald ML, Cho D-S, Kamins J, Nie Z, et al. The post-

synaptic density of human postmortem brain tissues: an experimental study

paradigm for neuropsychiatric illnesses. PloS One. 2009;4:e5251.

32.   MacDonald ML, Ciccimaro E, Prakash A, Banerjee A, Seeholzer SH, Blair IA, et

al. Biochemical Fractionation and Stable Isotope Dilution Liquid Chromatography-

mass Spectrometry for Targeted and Microdomain-specific Protein Quantification

in Human Postmortem Brain Tissue. Mol Cell Proteomics MCP. 2012;11:1670–

1681.

33.   Zecha J, Satpathy S, Kanashova T, Avanessian SC, Kane MH, Clauser KR, et al.

TMT Labeling for the Masses: A Robust and Cost-efficient, In-solution Labeling

Approach. Mol Cell Proteomics MCP. 2019;18:1468–1478.

34.   Perez-Riverol Y, Bai J, Bandla C, García-Seisdedos D, Hewapathirana S,

Kamatchinathan S, et al. The PRIDE database resources in 2022: a hub for mass

spectrometry-based proteomics evidences. Nucleic Acids Res. 2021;50:D543–

D552.

35.   Browne CJ, Futamura R, Minier-Toribio A, Hicks EM, Ramakrishnan A, Martínez-

Rivera F, et al. Transcriptional signatures of heroin intake and seeking throughout

the brain reward circuit. BioRxiv Prepr Serv Biol. 2023:2023.01.11.523688.

36.   Mews P, Cunningham AM, Scarpa J, Ramakrishnan A, Hicks EM, Bolnick S, et al.

Convergent abnormalities in striatal gene networks in human cocaine use disorder

and mouse cocaine administration models. Sci Adv. 2023;9:eadd8946.

37.   Yu G, Wang L-G, Han Y, He Q-Y. clusterProfiler: an R package for comparing

biological themes among gene clusters. Omics J Integr Biol. 2012;16:284–287.

38. Yu G, He Q-Y. ReactomePA: an R/Bioconductor package for reactome pathway

analysis and visualization. Mol Biosyst. 2016;12:477–479.

39. Zhou Y, Zhou B, Pache L, Chang M, Khodabakhshi AH, Tanaseichuk O, et al.

Metascape provides a biologist-oriented resource for the analysis of systems-level

datasets. Nat Commun. 2019;10:1523.

40. Langfelder P, Horvath S. WGCNA: an R package for weighted correlation network

analysis. BMC Bioinformatics. 2008;9:559.

41. Langfelder P, Zhang B, Horvath S. Defining clusters from a hierarchical cluster

tree: the Dynamic Tree Cut package for R. Bioinforma Oxf Engl. 2008;24:719–

720.

42. Zhang B, Horvath S. A general framework for weighted gene co-expression

network analysis. Stat Appl Genet Mol Biol. 2005;4:Article17.

43. Margolin AA, Nemenman I, Basso K, Wiggins C, Stolovitzky G, Dalla Favera R, et

al. ARACNE: an algorithm for the reconstruction of gene regulatory networks in a

mammalian cellular context. BMC Bioinformatics. 2006;7 Suppl 1:S7.

44. Koopmans F, van Nierop P, Andres-Alonso M, Byrnes A, Cijsouw T, Coba MP, et

al. SynGO: An Evidence-Based, Expert-Curated Knowledge Base for the

Synapse. Neuron. 2019;103:217-234.e4.

45. Cahill KM, Huo Z, Tseng GC, Logan RW, Seney ML. Improved identification of

concordant and discordant gene expression signatures using an updated rank-

rank hypergeometric overlap approach. Sci Rep. 2018;8:9588.

46. Morrison KE, Stenson AF, Marx-Rattner R, Carter S, Michopoulos V, Gillespie CF,

et al. Developmental Timing of Trauma in Women Predicts Unique Extracellular Vesicle Proteome Signatures. Biol Psychiatry. 2022;91:273–282.

47. Javitt A, Barnea E, Kramer MP, Wolf-Levy H, Levin Y, Admon A, et al. Pro-inflammatory Cytokines Alter the Immunopeptidome Landscape by Modulation of HLA-B Expression. Front Immunol. 2019;10.

48. Liang TW, Chiu HH, Gurney A, Sidle A, Tumas DB, Schow P, et al. Vascular endothelial-junctional adhesion molecule (VE-JAM)/JAM 2 interacts with T, NK, and dendritic cells through JAM 3. J Immunol Baltim Md 1950. 2002;168:1618–1626.

49. Koss H, Bunney TD, Behjati S, Katan M. Dysfunction of phospholipase Cγ in immune disorders and cancer. Trends Biochem Sci. 2014;39:603–611.

50. Sabbir MG, Speth RC, Albensi BC. Loss of Cholinergic Receptor Muscarinic 1 (CHRM1) Protein in the Hippocampus and Temporal Cortex of a Subset of Individuals with Alzheimer's Disease, Parkinson's Disease, or Frontotemporal Dementia: Implications for Patient Survival. J Alzheimers Dis JAD. 2022;90:727–747.

51. Zhu G, Fang Y, Cui X, Jia R, Kang X, Zhao R. Magnolol upregulates CHRM1 to attenuate Amyloid-β-triggered neuronal injury through regulating the cAMP/PKA/CREB pathway. J Nat Med. 2022;76:188–199.

52. Strang KH, Golde TE, Giasson BI. MAPT mutations, tauopathy, and mechanisms of neurodegeneration. Lab Invest. 2019;99:912–928.

53. Ma S, Attarwala IY, Xie X-Q. SQSTM1/p62: A Potential Target for Neurodegenerative Disease. ACS Chem Neurosci. 2019;10:2094–2114.

54. Aykac A, Sehirli AO. The Role of the SLC Transporters Protein in the

Neurodegenerative Disorders. Clin Psychopharmacol Neurosci. 2020;18:174–

187.

55. Chee LY, Cumming A. Polymorphisms in the Cholinergic Receptors Muscarinic

(CHRM2 and CHRM3) Genes and Alzheimer's Disease. Avicenna J Med

Biotechnol. 2018;10:196–199.

56. Querfurth H, Lee H-K. Mammalian/mechanistic target of rapamycin (mTOR)

complexes in neurodegeneration. Mol Neurodegener. 2021;16:44.

57. Foskett JK. Inositol trisphosphate receptor Ca2+ release channels in neurological

diseases. Pflugers Arch. 2010;460:481–494.

58. Lokireddy S, Kukushkin NV, Goldberg AL. cAMP-induced phosphorylation of 26S

proteasomes on Rpn6/PSMD11 enhances their activity and the degradation of

misfolded proteins. Proc Natl Acad Sci U S A. 2015;112:E7176–E7185.

59. Huang Z-N, Her L-S. The Ubiquitin Receptor ADRM1 Modulates HAP40-Induced

Proteasome Activity. Mol Neurobiol. 2017;54:7382–7400.

60. Benkler C, O'Neil AL, Slepian S, Qian F, Weinreb PH, Rubin LL. Aggregated

SOD1 causes selective death of cultured human motor neurons. Sci Rep.

2018;8:16393.

61. Heremans IP, Caligiore F, Gerin I, Bury M, Lutz M, Graff J, et al. Parkinson's

disease protein PARK7 prevents metabolite and protein damage caused by a

glycolytic metabolite. Proc Natl Acad Sci. 2022;119:e2111338119.

62. Plun-Favreau H, Klupsch K, Moisoi N, Gandhi S, Kjaer S, Frith D, et al. The

mitochondrial protease HtrA2 is regulated by Parkinson's disease-associated

kinase PINK1. Nat Cell Biol. 2007;9:1243–1252.

63. Shi Q, Li J-L, Ma Y, Gao L-P, Xiao K, Wang J, et al. Decrease of RyR2 in the

prion infected cell line and in the brains of the scrapie infected mice models and

the patients of human prion diseases. Prion. 2018;12:175–184.

64. Bradley JA, Ramirez G, Hagood JS. Roles and Regulation of Thy-1, a Context-

Dependent Modulator of Cell Phenotype. BioFactors Oxf Engl. 2009;35:258–265.

65. Jeng C-J, McCarroll SA, Martin TFJ, Floor E, Adams J, Krantz D, et al. Thy-1 Is a

Component Common to Multiple Populations of Synaptic Vesicles. J Cell Biol.

1998;140:685–698.

66. Dahimene S, von Elsner L, Holling T, Mattas LS, Pickard J, Lessel D, et al.

Biallelic CACNA2D1 loss-of-function variants cause early-onset developmental

epileptic encephalopathy. Brain J Neurol. 2022;145:2721–2729.

67. Grosso BJ, Kramer AA, Tyagi S, Bennett DF, Tifft CJ, D'Souza P, et al. Complex

effects on CaV2.1 channel gating caused by a CACNA1A variant associated with

a severe neurodevelopmental disorder. Sci Rep. 2022;12:9186.

68. Luo X, Rosenfeld JA, Yamamoto S, Harel T, Zuo Z, Hall M, et al. Clinically severe

CACNA1A alleles affect synaptic function and neurodegeneration differentially.

PLoS Genet. 2017;13:e1006905.

69. Chen J, Li L, Chen S-R, Chen H, Xie J-D, Sirrieh RE, et al. The α2δ-1-NMDA

Receptor Complex Is Critically Involved in Neuropathic Pain Development and

Gabapentin Therapeutic Actions. Cell Rep. 2018;22:2307–2321.

70. Lin C-H, Lin P-P, Lin C-Y, Lin C-H, Huang C-H, Huang Y-J, et al. Decreased

mRNA expression for the two subunits of system xc–, SLC3A2 and SLC7A11, in

WBC in patients with schizophrenia: Evidence in support of the hypo-

glutamatergic hypothesis of schizophrenia. J Psychiatr Res. 2016;72:58–63.

71. Peineau S, Bradley C, Taghibiglou C, Doherty A, Bortolotto ZA, Wang YT, et al.

The role of GSK-3 in synaptic plasticity. Br J Pharmacol. 2008;153:S428–S437.

72. Jaworski T, Banach-Kasper E, Gralec K. GSK-3β at the Intersection of Neuronal

Plasticity and Neurodegeneration. Neural Plast. 2019;2019:4209475.

73. Bregin A, Kaare M, Jagomäe T, Karis K, Singh K, Laugus K, et al. Expression and

impact of Lsamp neural adhesion molecule in the serotonergic neurotransmission

system. Pharmacol Biochem Behav. 2020;198:173017.

74. Bregin A, Mazitov T, Aug I, Philips M-A, Innos J, Vasar E. Increased sensitivity to

psychostimulants and GABAergic drugs in Lsamp-deficient mice. Pharmacol

Biochem Behav. 2019;183:87–97.

75. Moriwaki Y, Kubo N, Watanabe M, Asano S, Shinoda T, Sugino T, et al.

Endogenous neurotoxin-like protein Ly6H inhibits alpha7 nicotinic acetylcholine

receptor currents at the plasma membrane. Sci Rep. 2020;10:11996.

76. Puddifoot CA, Wu M, Sung R-J, Joiner WJ. Ly6h Regulates Trafficking of Alpha7

Nicotinic Acetylcholine Receptors and Nicotine-Induced Potentiation of

Glutamatergic Signaling. J Neurosci. 2015;35:3420–3430.

77. Brennaman LH, Zhang X, Guan H, Triplett JW, Brown A, Demyanenko GP, et al.

Polysialylated NCAM and EphrinA/EphA Regulate Synaptic Development of

GABAergic Interneurons in Prefrontal Cortex. Cereb Cortex N Y NY.

2013;23:162–177.

78. Chen HC, Loh HH. mu-Opioid receptor gene expression: the role of NCAM.

Neuroscience. 2001;108:7–15.

79. Duncan BW, Murphy KE, Maness PF. Molecular Mechanisms of L1 and NCAM Adhesion Molecules in Synaptic Pruning, Plasticity, and Stabilization. Front Cell Dev Biol. 2021;9:625340.

80. Nelson EC, Lynskey MT, Heath AC, Wray N, Agrawal A, Shand FL, et al. ANKK1, TTC12, and NCAM1 Polymorphisms and Heroin Dependence – importance of considering drug exposure. JAMA Psychiatry Chic Ill. 2013;70:325–333.

81. Polo-Parada L, Bose CM, Plattner F, Landmesser LT. Distinct Roles of Different Neural Cell Adhesion Molecule (NCAM) Isoforms in Synaptic Maturation Revealed by Analysis of NCAM 180 kDa Isoform-Deficient Mice. J Neurosci. 2004;24:1852–1864.

82. Bae Y-S, Chung W, Han K, Park KY, Kim H, Kim E, et al. Down-regulation of RalBP1 expression reduces seizure threshold and synaptic inhibition in mice. Biochem Biophys Res Commun. 2013;433:175–180.

83. Chen M-S, Kim H, Jagot-Lacoussiere L, Maurel P. Cadm3 (Necl-1) interferes with the activation of the PI3 kinase/Akt signaling cascade and inhibits Schwann cell myelination in vitro. Glia. 2016;64:2247–2262.

84. Kamimura K, Maeda N. Glypicans and Heparan Sulfate in Synaptic Development, Neural Plasticity, and Neurological Disorders. Front Neural Circuits. 2021;15:595596.

85. Beurrier C, Faideau M, Bennouar K-E, Escartin C, Kerkerian-Le Goff L, Bonvento G, et al. Ciliary Neurotrophic Factor Protects Striatal Neurons against Excitotoxicity by Enhancing Glial Glutamate Uptake. PLoS ONE. 2010;5:e8550.

86. Wang Q, Chiu S-L, Koropouli E, Hong I, Mitchell S, Easwaran TP, et al.
Neuropilin-2/PlexinA3 Receptors Associate with GluA1 and Mediate Sema3F-
dependent Homeostatic Scaling in Cortical Neurons. Neuron. 2017;96:1084-
1098.e7.

87. Yamashita N, Usui H, Nakamura F, Chen S, Sasaki Y, Hida T, et al. Plexin-A4-
dependent retrograde semaphorin 3A signalling regulates the dendritic
localization of GluA2-containing AMPA receptors. Nat Commun. 2014;5:3424.

88. Awasthi A, Ramachandran B, Ahmed S, Benito E, Shinoda Y, Nitzan N, et al.
Synaptotagmin-3 drives AMPA receptor endocytosis, depression of synapse
strength, and forgetting. Science. 2019;363:eaav1483.

89. Jin X-T, Galvan A, Wichmann T, Smith Y. Localization and Function of GABA
Transporters GAT-1 and GAT-3 in the Basal Ganglia. Front Syst Neurosci.
2011;5:63.

90. Savtchenko L, Megalogeni M, Rusakov DA, Walker MC, Pavlov I. Synaptic GABA
release prevents GABA transporter type-1 reversal during excessive network
activity. Nat Commun. 2015;6:6597.

91. Yuan A, Rao MV, Veeranna, Nixon RA. Neurofilaments at a glance. J Cell Sci.
2012;125:3257–3263.

92. Nikoshkov A, Drakenberg K, Wang X, Horvath MCs, Keller E, Hurd YL. Opioid
neuropeptide genotypes in relation to heroin abuse: Dopamine tone contributes to
reversed mesolimbic proenkephalin expression. Proc Natl Acad Sci.
2008;105:786–791.

93. Kieffer BL, Gavériaux-Ruff C. Exploring the opioid system by gene knockout. Prog

Neurobiol. 2002;66:285–306.

94. Chaplot K, Jarvela TS, Lindberg I. Secreted Chaperones in Neurodegeneration. Front Aging Neurosci. 2020;12:268.

95. Ram A, Edwards T, McCarty A, Afrose L, McDermott MV, Bobeck EN. GPR171 Agonist Reduces Chronic Neuropathic and Inflammatory Pain in Male, But Not Female Mice. Front Pain Res. 2021;2.

96. Castrogiovanni P, Sanfilippo C, Imbesi R, Maugeri G, Lo Furno D, Tibullo D, et al. Brain CHID1 Expression Correlates with NRGN and CALB1 in Healthy Subjects and AD Patients. Cells. 2021;10:882.

97. Manavalan A, Mishra M, Feng L, Sze SK, Akatsu H, Heese K. Brain site-specific proteome changes in aging-related dementia. Exp Mol Med. 2013;45:e39–e39.

98. Chanaday NL, Kavalali ET. Role of the endoplasmic reticulum in synaptic transmission. Curr Opin Neurobiol. 2022;73:102538.

99. Lewis JE, Brameld JM, Jethwa PH. Neuroendocrine Role for VGF. Front Endocrinol. 2015;6.

100. Rombaut B, Kessels S, Schepers M, Tiane A, Paes D, Solomina Y, et al. PDE inhibition in distinct cell types to reclaim the balance of synaptic plasticity. Theranostics. 2021;11:2080–2097.

101. Sil S, Periyasamy P, Thangaraj A, Chivero ET, Buch S. PDGF/PDGFR axis in the neural systems. Mol Aspects Med. 2018;62:63–74.

102. Puig S, Barker KE, Szott SR, Kann PT, Morris JS, Gutstein HB. Spinal Opioid Tolerance Depends upon Platelet-Derived Growth Factor Receptor-β Signaling, Not μ-Opioid Receptor Internalization. Mol Pharmacol. 2020;98:487–496.

103. Wang Y, Barker K, Shi S, Diaz M, Mo B, Gutstein HB. Blockade of PDGFR-β
activation eliminates morphine analgesic tolerance. Nat Med. 2012;18:385–387.

104. Zhou Z, Meng Y, Asrar S, Todorovski Z, Jia Z. A critical role of Rho-kinase
ROCK2 in the regulation of spine and synaptic function. Neuropharmacology.
2009;56:81–89.

105. Strawbridge RJ, Ward J, Lyall LM, Tunbridge EM, Cullen B, Graham N, et al.
Genetics of self-reported risk-taking behaviour, trans-ethnic consistency and
relevance to brain gene expression. Transl Psychiatry. 2018;8:1–11.

106. Jeanne M, Demory H, Moutal A, Vuillaume M-L, Blesson S, Thépault R-A, et al.
Missense variants in DPYSL5 cause a neurodevelopmental disorder with corpus
callosum agenesis and cerebellar abnormalities. Am J Hum Genet.
2021;108:951–961.

107. Gawden-Bone CM, Frazer GL, Richard AC, Ma CY, Strege K, Griffiths GM. PIP5
Kinases Regulate Membrane Phosphoinositide and Actin Composition for
Targeted Granule Secretion by Cytotoxic Lymphocytes. Immunity. 2018;49:427-
437.e4.

108. Schaffer TB, Smith JE, Cook EK, Phan T, Margolis SS. PKCε Inhibits Neuronal
Dendritic Spine Development through Dual Phosphorylation of Ephexin5. Cell
Rep. 2018;25:2470-2483.e8.

109. Russell SE, Puttick DJ, Sawyer AM, Potter DN, Mague S, Carlezon WA, et al.
Nucleus Accumbens AMPA Receptors Are Necessary for Morphine-Withdrawal-
Induced Negative-Affective States in Rats. J Neurosci. 2016;36:5748–5762.

110. Hearing M, Graziane N, Dong Y, Thomas MJ. Opioid and psychostimulant

plasticity: targeting overlap in nucleus accumbens glutamate signaling. Trends

Pharmacol Sci. 2018;39:276–294.

111. Mallard TT, Ashenhurst JR, Harden KP, Fromme K. GABRA2, alcohol, and illicit

drug use: an event-level model of genetic risk for polysubstance use. J Abnorm

Psychol. 2018;127:190–201.

112. Enoch M-A, Hodgkinson CA, Yuan Q, Shen P-H, Goldman D, Roy A. The

Influence of GABRA2, Childhood Trauma, and Their Interaction on Alcohol,

Heroin, and Cocaine Dependence. Biol Psychiatry. 2010;67:20–27.

113. Sun Y, Zhang Y, Zhang D, Chang S, Jing R, Yue W, et al. GABRA2 rs279858-

linked variants are associated with disrupted structural connectome of reward

circuits in heroin abusers. Transl Psychiatry. 2018;8:1–10.

114. Nirwane A, Yao Y. Laminins and their receptors in the CNS. Biol Rev Camb

Philos Soc. 2019;94:283–306.

115. Gómez-López S, Martínez-Silva AV, Montiel T, Osorio-Gómez D, Bermúdez-

Rattoni F, Massieu L, et al. Neural ablation of the PARK10 candidate Plpp3 leads

to dopaminergic transmission deficits without neurodegeneration. Sci Rep.

2016;6:24028.

116. Zhang S, Glukhova SA, Caldwell KA, Caldwell GA. NCEH-1 modulates

cholesterol metabolism and protects against α-synuclein toxicity in a C. elegans

model of Parkinson's disease. Hum Mol Genet. 2017;26:3823–3836.

117. Jin J, Yu Q, Han C, Hu X, Xu S, Wang Q, et al. LRRFIP2 negatively regulates

NLRP3 inflammasome activation in macrophages by promoting Flightless-I-

mediated caspase-1 inhibition. Nat Commun. 2013;4:2075.

118. Wright BD, Loo L, Street SE, Ma A, Taylor-Blake B, Stashko MA, et al. The lipid
kinase PIP5K1C regulates pain signaling and sensitization. Neuron. 2014;82:836–
847.

119. Pavelitz T, Gray LT, Padilla SL, Bailey AD, Weiner AM. PGBD5: a neural-specific
intron-containing piggyBac transposase domesticated over 500 million years ago
and conserved from cephalochordates to humans. Mob DNA. 2013;4:23.

120. Cai Z, Ratka A. Opioid system and Alzheimer's disease. Neuromolecular Med.
2012;14:91–111.

121. Kennedy GJ. Bad News and Good: Opioids are Associated with the Incidence of
Dementia, but the Effect is Substantial Only for Those 75-80 Years of Age. Am J
Geriatr Psychiatry Off J Am Assoc Geriatr Psychiatry. 2023;31:324–325.

122. Anthony IC, Norrby KE, Dingwall T, Carnie FW, Millar T, Arango JC, et al.
Predisposition to accelerated Alzheimer-related changes in the brains of human
immunodeficiency virus negative opiate abusers. Brain J Neurol. 2010;133:3685–
3698.

123. Mao L, Wang K, Zhang P, Ren S, Sun J, Yang M, et al. Carbonyl Reductase 1
Attenuates Ischemic Brain Injury by Reducing Oxidative Stress and
Neuroinflammation. Transl Stroke Res. 2021;12:711–724.

124. De Rossi P, Nomura T, Andrew RJ, Masse NY, Sampathkumar V, Musial TF, et
al. Neuronal BIN1 Regulates Presynaptic Neurotransmitter Release and Memory
Consolidation. Cell Rep. 2020;30:3520-3535.e7.

125. Ye H, Ojelade SA, Li-Kroeger D, Zuo Z, Wang L, Li Y, et al. Retromer subunit,
VPS29, regulates synaptic transmission and is required for endolysosomal

function in the aging brain. ELife;9:e51977.

126. Chen X, Lu W, Wu D. Sirtuin 2 (SIRT2): Confusing Roles in the Pathophysiology of Neurological Disorders. Front Neurosci. 2021;15:614107.

127. Ferguson D, Koo JW, Feng J, Heller E, Rabkin J, Heshmati M, et al. Essential Role of SIRT1 Signaling in the Nucleus Accumbens in Cocaine and Morphine Action. J Neurosci. 2013;33:16088–16098.

128. Zhao B, Zhu Y, Wang W, Cui H, Wang Y, Lai J. Analysis of Variations in the Glutamate Receptor, N-Methyl D-Aspartate 2A (GRIN2A) Gene Reveals Their Relative Importance as Genetic Susceptibility Factors for Heroin Addiction. PLoS ONE. 2013;8:e70817.

129. Elmasri M, Lotti JS, Aziz W, Steele OG, Karachaliou E, Sakimura K, et al. Synaptic Dysfunction by Mutations in GRIN2B: Influence of Triheteromeric NMDA Receptors on Gain-of-Function and Loss-of-Function Mutant Classification. Brain Sci. 2022;12:789.

130. Sheng Z, Liu Q, Cheng C, Li M, Barash J, Kofke WA, et al. Fentanyl induces autism-like behaviours in mice by hypermethylation of the glutamate receptor gene Grin2b. Br J Anaesth. 2022;129:544–554.

131. Lee E, Lee S, Shin JJ, Choi W, Chung C, Lee S, et al. Excitatory synapses and gap junctions cooperate to improve Pv neuronal burst firing and cortical social cognition in Shank2-mutant mice. Nat Commun. 2021;12:5116.

132. Bryant DT, Landles C, Papadopoulou AS, Benjamin AC, Duckworth JK, Rosahl T, et al. Disruption to schizophrenia-associated gene Fez1 in the hippocampus of HDAC11 knockout mice. Sci Rep. 2017;7:11900.

133. Glausier JR, Kimoto S, Fish KN, Lewis DA. Lower glutamic acid decarboxylase 65-kDa isoform messenger RNA and protein levels in the prefrontal cortex in schizoaffective disorder but not schizophrenia. Biol Psychiatry. 2015;77:167–176.

134. Kowalski EA, Chen J, Hazy A, Fritsch LE, Gudenschwager-Basso EK, Chen M, et al. Peripheral loss of EphA4 ameliorates TBI-induced neuroinflammation and tissue damage. J Neuroinflammation. 2019;16:210.

135. Bott CJ, Winckler B. Intermediate filaments in developing neurons: Beyond structure. Cytoskeleton. 2020;77:110–128.

136. Borgkvist A, Usiello A, Greengard P, Fisone G. Activation of the cAMP/PKA/DARPP-32 Signaling Pathway is Required for Morphine Psychomotor Stimulation but not for Morphine Reward. Neuropsychopharmacology. 2007;32:1995–2003.

137. Gould TD, Manji HK. DARPP-32: A molecular switch at the nexus of reward pathway plasticity. Proc Natl Acad Sci. 2005;102:253–254.

138. Mahajan SD, Aalinkeel R, Reynolds JL, Nair BB, Sykes DE, Hu Z, et al. Therapeutic targeting of 'DARPP-32': a key signaling molecule in the dopiminergic pathway for the treatment of opiate addiction. Int Rev Neurobiol. 2009;88:199–222.

139. Brami-Cherrier K, Lewis RG, Cervantes M, Liu Y, Tognini P, Baldi P, et al. Cocaine-mediated circadian reprogramming in the striatum through dopamine D2R and PPARγ activation. Nat Commun. 2020;11:4448.

140. Bumgarner JR, McCray EW, Nelson RJ. The disruptive relationship among circadian rhythms, pain, and opioids. Front Neurosci. 2023;17.

141. Gallerani M, Manfredini R, Dal Monte D, Calò G, Brunaldi V, Simonato M. Circadian differences in the individual sensitivity to opiate overdose. Crit Care Med. 2001;29:96–101.

142. Ke Y, Weng M, Chhetri G, Usman M, Li Y, Yu Q, et al. Trappc9 deficiency in mice impairs learning and memory by causing imbalance of dopamine D1 and D2 neurons. Sci Adv. 2020;6:eabb7781.

143. Carter SD, Hampton CM, Langlois R, Melero R, Farino ZJ, Calderon MJ, et al. Ribosome-associated vesicles: A dynamic subcompartment of the endoplasmic reticulum in secretory cells. Sci Adv. 2020;6:eaay9572.

144. Mauvoisin D, Wang J, Jouffe C, Martin E, Atger F, Waridel P, et al. Circadian clock-dependent and -independent rhythmic proteomes implement distinct diurnal functions in mouse liver. Proc Natl Acad Sci. 2014;111:167–172.

145. Ancona Esselmann SG, Díaz-Alonso J, Levy JM, Bemben MA, Nicoll RA. Synaptic homeostasis requires the membrane-proximal carboxy tail of GluA2. Proc Natl Acad Sci. 2017;114:13266–13271.

146. Weinhard L, di Bartolomei G, Bolasco G, Machado P, Schieber NL, Neniskyte U, et al. Microglia remodel synapses by presynaptic trogocytosis and spine head filopodia induction. Nat Commun. 2018;9:1228.

147. Calcia MA, Bonsall DR, Bloomfield PS, Selvaraj S, Barichello T, Howes OD. Stress and neuroinflammation: a systematic review of the effects of stress on microglia and the implications for mental illness. Psychopharmacology (Berl). 2016;233:1637–1650.

148. Gamble MC, Williams BR, Singh N, Posa L, Freyberg Z, Logan RW, et al. Mu-

opioid receptor and receptor tyrosine kinase crosstalk: Implications in

mechanisms of opioid tolerance, reduced analgesia to neuropathic pain,

dependence, and reward. Front Syst Neurosci. 2022;16:1059089.

149. Lee U, Choi C, Ryu SH, Park D, Lee S-E, Kim K, et al. SCAMP5 plays a critical

role in axonal trafficking and synaptic localization of NHE6 to adjust quantal size

at glutamatergic synapses. Proc Natl Acad Sci. 2021;118:e2011371118.

    Figure Legends

    Figure 1. Proteomic alterations in tissue homogenates from NAc and DLPFC in

    subjects with OUD. a) Log_2FC plotted relative to $-log_{10}$ *p* value by volcano plot for DE

    proteins in NAc homogenates. **b**) Log_2FC plotted relative to $-log_{10}$ *p* value by volcano

    plot for DE proteins in DLPFC homogenates. **c**) RRHO plot indicating weak overlap or

    concordance of protein alterations between DLPFC and NAc in OUD subjects. **d**) Venn

    diagrams of downregulated and upregulated proteins between NAc and DLPFC

    homogenates. **e**) Log_2FC plotted relative to $-log_{10}$ *p* value by volcano plot for DE

    proteins in NAc synaptosomes. **f**) Log_2FC plotted relative to $-log_{10}$ *p* value by volcano

    plot for DE proteins in DLPFC synaptosomes. **g**) RRHO plot indicating very weak

    discordance of protein expression alterations in OUD between DLPFC and NAc

    synaptosomes. **h**) Venn diagrams of downregulated and upregulated proteins between

    NAc and DLPFC synaptosomes. Horizontal red lines indicate significance cutoffs of

    p<0.05, with vertical red lines represent $log_2FC\pm0.26$ (panels a, b, e, and f). Proteins

    that reached both unadjusted p<0.05 and $log_2FC\pm0.26$ were identified as DE proteins,

    upregulated labeled as red circles and downregulated labeled as blue circles.

    Figure 2. Differential enrichment of synaptic proteins between NAc and DLPFC

    associated with OUD. Lollipop plot showing pathways enriched from DE proteins in

    synaptosomes relative to homogenates in **a**) NAc and **b**) DLPFC. Adjusted p-value by

    color. Size of the circle represents counts of proteins within pathways. The enrichment

    score is calculated as the count of proteins identified as DE in OUD subjects divided by

the count of proteins in the background of the respective ontological pathway. Bolded

text with stars represents pathways enriched in both brain regions. Bolded only text

represents pathways enriched in synaptosomes of the region analyzed. Also see

Supplementary Table S11 for Enrichment in NAc Synaptosomes; Supplementary Table

S12 for Enrichment in DLPFC Synaptosomes. NAc, Nucleus Accumbens; DLPFC,

Dorsolateral Prefrontal Cortex.

1104

Figure 3. Alterations in protein synaptic enrichments in DLPFC and NAc

associated with OUD. Heatmap highlighting top differentially expressed proteins in

homogenates and synaptosomes in NAc and DLPFC of OUD subjects compared to

unaffected controls. Warmer colors indicate increasing log_2FC and highly enriched

proteins in OUD. In contrast, cooler colors indicate decreasing log_2FC and negative

enrichment of proteins in OUD synaptosomes. Proteins are filtered for FDR < 0.10.

1111

Figure 4. Diurnal rhythms of protein expression in DLPFC and NAc synaptosomes

associated with OUD. a) Top left: Heatmap highlighting top 200 rhythmic proteins from

NAc synaptosomes of unaffected subjects. Top right: Rhythmic proteins in unaffected

subjects were plotted in OUD subjects to show disruption of protein rhythmicity. Bottom

left: Heatmap highlighting top 200 rhythmic proteins from NAc synaptosomes of OUD

subjects. Bottom right: rhythmic proteins in OUD subjects were plotted in unaffected

subjects to show gain of protein rhythmicity in OUD. Heatmaps were generated by

performing supervised clustering of expression of selected top 200 rhythmic proteins.

Subjects were ordered by TOD to visualize expression levels over a period of 24 hours.

Yellow color indicates increased Z-score and higher protein expression, while blue color

indicates decreased Z-score and lower protein expression. **b**) Scatterplots of top

rhythmic proteins in NAc synaptosomes from unaffected and OUD subjects.

Scatterplots were generated to represent expression rhythms for individual proteins.

The x-axis represents TOD on the ZT scale and protein expression level is on y-axis,

with each dot representing a subject. The red line is the fitted sinusoidal curve to reflect

temporal rhythms. **c**) Pathway enrichment analysis comparing rhythmic proteins in NAc

synaptosomes from unaffected and OUD subjects. Warmer colors indicate increasing

$-\log_{10}$ *p* value and highly rhythmic pathways in each group. **d**) Venn diagrams showing

low overlap of rhythmic proteins and genes in NAc synaptosomes from unaffected and

OUD subjects. **e**) Top left: Heatmap highlighting top 200 rhythmic proteins from DLPFC

synaptosomes of unaffected subjects. Top right: Rhythmic proteins in unaffected

subjects were plotted in OUD subjects to show disruption of protein rhythmicity. Bottom

left: heatmap highlighting top 200 rhythmic proteins from DLPFC synaptosomes of OUD

subjects. Bottom right: rhythmic proteins in OUD subjects were plotted in unaffected

subjects to show gain of protein rhythmicity in OUD**. f**) Scatterplots of top rhythmic

proteins in DLPFC synaptosomes from unaffected and OUD subjects. **g**) Pathway

enrichment analysis comparing top 200 rhythmic proteins found in DLPFC

synaptosomes from unaffected and OUD subjects. **h**) Venn diagrams showing low

overlap of rhythmic proteins and genes in DLPFC synaptosomes from unaffected and

OUD subjects. Also see Supplementary Figure S1 for protein expression rhythm

heatmaps in unaffected homogenates; Supplementary Figure S2 for protein expression

heatmaps in OUD homogenates; Supplementary table S13 for NAc Homogenates

Rhythms in Unaffected subjects; Supplementary table S14 for NAc Homogenates

Rhythms in OUD subjects; Supplementary table S15 for DLPFC Homogenates Rhythms

in Unaffected subjects; Supplementary table S16 for DLPFC Homogenates Rhythms in

OUD subjects; Supplementary table S17 for NAc Synaptosomes Rhythms in Unaffected

subjects; Supplementary table S18 for NAc Synaptosomes Rhythms in OUD subjects;

Supplementary table S19 for DLPFC Synaptosomes Rhythms in Unaffected subjects;

Supplementary table S20 for DLPFC Synaptosomes Rhythms in OUD subjects; TOD,

Time of Death; NAc, Nucleus Accumbens; DLPFC, Dorsolateral Prefrontal Cortex.

1152

Figure 5. Altered rhythmicity of the synaptic proteome in DLPFC and NAc

associated with OUD. Comparison of rhythmic proteins between unaffected and OUD

subjects in synaptosomes from NAc and DLPFC. **a**) Circoplot highlights few rhythmic

proteins that were identical (purple lines) and shared ontology (light blue lines) between

unaffected and OUD subjects in NAc synaptosomes. Change in rhythmicity analysis in

NAc Synaptosomes revealed that 0 proteins gained rhythmicity, while 23 lost rhythmicity

in OUD subjects. **b**) Scatterplots of top proteins that lost rhythmicity in NAc

synaptosomes between unaffected and OUD subjects. Scatterplots were generated to

represent expression rhythms for individual proteins. The x-axis represents TOD on the

ZT scale and protein expression level is on y-axis, with each dot representing a subject.

The red line is the fitted sinusoidal curve to reflect temporal rhythms. **c**) Pathway

enrichment on proteins that changed rhythmicity between unaffected and OUD subjects

in NAc synaptosomes. **d**) Circoplot highlights few rhythmic proteins that were identical

(purple lines) and shared ontology (light blue lines) between unaffected and OUD

subjects in DLPFC synaptosomes. Change in rhythmicity analysis in DLPFC

Synaptosomes revealed that 11 proteins gained rhythmicity, while 7 lost rhythmicity. **e**)

Scatterplots of top proteins that lost rhythmicity in DLPFC synaptosomes between

unaffected and OUD subjects. **f**) Pathway enrichment on proteins that changed

rhythmicity between unaffected and OUD subjects in DLPFC synaptosomes. Also see

Figure S3 for gain/loss rhythmicity proteins in NAc and DLPFC homogenates;

Supplementary Figure S4 for DLPFC synaptosomes enriched gain and lost rhythm

pathways; Table S21 for gain/loss rhythmicity protein list in NAc homogenates;

Supplementary Table S22 for gain/loss rhythmicity protein list in DLPFC homogenates;

Supplementary Table S23 for gain/loss rhythmicity protein list in NAc synaptosomes;

Supplementary Table S24 for gain/loss rhythmicity protein list in DLPFC synaptosomes.

TOD, time of death; NAc Nucleus Accumbens; DLPFC, DorsoLateral PreFrontal Cortex.

1179

Figure 6. Altered protein networks in NAc and DLPFC synaptosomes associated

with OUD. Weight Gene Correlation Network Analysis (WGCNA) was used to generate

protein coexpression modules from each brain region separately. The identified

modules that survived module preservation analysis were arbitrarily assigned colors.

Pie charts generated from module differential connectivity (MDC) analysis summarize

modules that gained or lost connectivity between unaffected and OUD subjects in NAc

(**a**) and DLPFC (**b**). **a**) Comparing module connectivity between unaffected and OUD

subjects in NAc synaptosomes identified 13 modules that lost connectivity in OUD,

while 1 remained unchanged. Modules Turquoise, Green Yellow and Green were

composed of several rhythmic protein hubs and showed loss in connectivity in OUD. **b**)

Comparing module connectivity between unaffected and OUD subjects in DLPFC

synaptosomes revealed all identified modules lost connectivity in OUD. Modules Green

and Black were composed of several rhythmic hubs and showed loss of connectivity in

OUD. **c**) Synaptic enrichment analysis of protein networks that lost connectivity in NAc

synaptosomes. **d**) Synaptic enrichment analysis of protein networks that lost

connectivity in DLPFC synaptosomes. Also see Figure S5 for WGCNA dendrograms on

NAc and DLPFC homogenates; Figure S6 for WGCNA dendrograms on NAc and

DLPFC synaptosomes; Figure S7 for modules from NAc homogenates; Figure S8 for

modules from NAc synaptosomes; Figure S9 for modules from DLPFC homogenates;

Figure S10 for modules from DLPFC synaptosomes; Figure S11 for additional MDC

summaries; Supplementary Table S25 for WGCNA module assignments and proteins;

Supplementary Table S26 for MDC analysis. DE, Differentially Expressed; Nucleus

Accumbens; DLPFC, Dorsolateral Prefrontal Cortex.

Proteins Altered in OUD and Enriched in Synaptosomes

a *NAc synaptosomes*

Unaffected — Top 200 Rhythmic Proteins
OUD — Top 200 Rhythmic Proteins from Unaffected Subjects
OUD — Top 200 Rhythmic Proteins
Unaffected — Top 200 Rhythmic Proteins from OUD Subjects

Subject Ordered by TOD

Z-score −2 to 2

b

Unaffected: GAT1, ATP1B2, ZW10
OUD: NEFL, NEFM, PLCL1

Expression vs ZT

c

Chemical synaptic trans. / Modulation of Chem. Syn. Trans. / Behavior / Neuron projection development / Pos. reg. of peptide hormone / Supramolecular fiber organization / Rho GTPase Sig. / Hemostasis / Cellular response to peptide hormone / Prostaglandin sig. / Synaptic sig. / Cellular sig. / Maintenance of cell polarity / Amino acid metabolism / Organic hydroxy metabolism / Neutrophil degranulation / Estrogen receptor sig. / Filament cytoskeleton organization / Actin filament / Neuropeptide sig.

Unaffected
OUD

−log10(p-value) 0 2 6 10 20

d

Unaffected
Proteins 324 · 6 · Genes 344
e.g., ATP9A, NDRG4, NPTX1

OUD
Proteins 69 · 8 · Genes 731
e.g., BAIAP3, GAT3, MPP2, SCN4B

e *DLPFC synaptosomes*

Unaffected — Top 200 Rhythmic Proteins
OUD — Top 200 Rhythmic Proteins from Unaffected Subjects
OUD — Top 200 Rhythmic Proteins
Unaffected — Top 200 Rhythmic Proteins from OUD Subjects

Subject Ordered by TOD

Z-score −2 to 2

f

Unaffected: BASP1, HMGN4, CD9
OUD: CHID1, SPCS2, NSHDL

Expression vs ZT

g

Gliogenesis / SRP-dependent sig. / Rho GTPases / Blood coagulation / Trans-synaptic sig. / Chemical synaptic transmission / Pos. reg. of cell development / Cell projection organization / Vesicle-mediated transport / Endomembrane organization / Secretion by cell / Trans-Golgi vesicle transport / Intracellular protein transport / Autophagy / Reg. of vesicle-mediated trans. / Small GTPase mediated sig. / Arp2/3 complex mediated actin nucleation / Protein nucleation / Glycosylation / Reg. of protein localization / Protein stability

Unaffected
OUD

−log10(p-value) 0 2 6 10 20

h

Unaffected
Proteins 117 · 6 · Genes 725
e.g., ABI2, FAT3, S100B

OUD
Proteins 180 · 2 · Genes 338
e.g., LMAN2, VPS33A

a Loss of Connectivity in Modules with Significant Synaptic Enrichment of Rhythmic Proteins in NAc

b Loss of Connectivity in Modules with Significant Synaptic Enrichment of Rhythmic Proteins in DLPFC