1 **DRMY1 promotes robust morphogenesis by sustaining the translation of cytokinin**
2 **signaling inhibitor proteins**
3

4 Shuyao Kong,[1,2,†] Mingyuan Zhu,[1,2,3,†] M. Regina Scarpin,[4] David Pan,[1,2] Longfei Jia,[5] Ryan E.
5 Martinez,[4] Simon Alamos,[6,7,8] Batthula Vijaya Lakshmi Vadde,[1,2] Hernan G. Garcia,[9,10,11,12,13] Shu-
6 Bing Qian,[5] Jacob O. Brunkard,[4] Adrienne H. K. Roeder [1,2,14,*]
7

[1] Weill Institute for Cell and Molecular Biology, Cornell University, Ithaca, NY 14853, USA
[2] Section of Plant Biology, School of Integrative Plant Science, Cornell University, Ithaca, NY 14853, USA
[3] Present address: Department of Biology, Duke University, Durham, NC 27708, USA
[4] Laboratory of Genetics, University of Wisconsin, Madison, WI 53706, USA
[5] Division of Nutritional Sciences, Cornell University, Ithaca, NY 14853, USA
[6] Environmental Genomics and Systems Biology Division, Lawrence Berkeley National Laboratory,
Berkeley, CA 94720, USA
[7] Feedstocks Division, Joint BioEnergy Institute, Emeryville, CA 94608, USA
[8] Department of Plant and Microbial Biology, University of California at Berkeley, Berkeley, CA 94720, USA
[9] Biophysics Graduate Group, University of California at Berkeley, Berkeley, CA 94720, USA
[10] Department of Physics, University of California at Berkeley, Berkeley, CA 94720, USA
[11] Institute for Quantitative Biosciences-QB3, University of California at Berkeley, Berkeley, CA 94720, USA
[12] Department of Molecular and Cell Biology, University of California at Berkeley, Berkeley, CA 94720, USA
[13] Chan Zuckerberg Biohub-San Francisco, San Francisco, CA 94158, USA
[14] Lead Contact
[†] These authors contributed equally
[*] Correspondence: ahr75@cornell.edu

25
26

27 **SUMMARY**
28

Robustness is the invariant development of phenotype despite environmental changes
and genetic perturbations. In the Arabidopsis flower bud, four sepals robustly initiate and grow to
constant size to enclose and protect the inner floral organs. We previously characterized the
mutant *development related myb-like1* (*drmy1*), where 3-5 sepals initiate variably and grow to
different sizes, compromising their protective function. The molecular mechanism underlying this
loss of robustness was unclear. Here, we show that *drmy1* has reduced TARGET OF
RAPAMYCIN (TOR) activity, ribosomal content, and translation. Translation reduction decreases
the protein level of ARABIDOPSIS RESPONSE REGULATOR7 (ARR7) and ARABIDOPSIS
HISTIDINE PHOSPHOTRANSFER PROTEIN 6 (AHP6), two cytokinin signaling inhibitors that
are normally rapidly produced before sepal initiation. The resultant upregulation of cytokinin
signaling disrupts robust auxin patterning and sepal initiation. Our work shows that the
homeostasis of translation, a ubiquitous cellular process, is crucial for the robust spatiotemporal
patterning of organogenesis.

Keywords: Robustness, TOR, translation, cytokinin, auxin, ribosomopathy, Arabidopsis, sepal,
morphogenesis

INTRODUCTION
46
Robustness, or canalization, is the invariant, reproducible development of phenotype,
unchanged by environmental fluctuations, genetic perturbations, or gene expression noise[1–4].
Commonly, within an individual, a given number of organs develop at well-defined positions to a
robust final size and shape, which is crucial for fitness under stabilizing selection[2]. For example,
transplanted eyes, limbs, and kidneys in mammals grow to a mature size similar to their donor,
irrespective of the mature size of the same type of organ in the recipient[5–7]. The pairs of wings
and halteres in Drosophila develop to robust, precisely coordinated final size and shape, which
are required for flight[8–11]. The characteristic cruciform flower in *Brassicaceae* consists of four
petals[12], a trait that can contribute to pollinator attraction[13]. The robust positioning of leaves
around the shoot apical meristem in plants, or phyllotaxis, ensures optimal light capture[14–16]. While
these examples of developmental robustness have been documented for a long time, the
underlying molecular mechanisms have just begun to be unveiled.
59
Earlier studies looking for genes involved in maintaining robustness have found *HEAT*
SHOCK PROTEIN 90 (*HSP90*). Mutations of *HSP90* cause a diverse array of phenotypic changes
in plants, fruit fly, zebrafish, worm, and humans[3,17,18]. Notably, the display and severity of these
changes vary between individuals and even between different parts of the same individual,
indicating that developmental robustness is disrupted[17,18]. *HSP90* encodes a protein chaperone
which has clients from nearly all developmental and signaling pathways[3]. *HSP90*, therefore, is a
hub gene that affects numerous other genes within the gene network[2]. Disruption of such a hub
gene would therefore trigger many defects in numerous developmental processes. Similarly,
genes involved in central cellular processes such as chromatin remodelling[19–21], transcription[19,20],
translation[22,23], and protein degradation[24,25] are also hub genes, and they have been found to be
important for developmental robustness in various systems including fungi, animals, and plants.
How these broad-acting hub genes contribute to the robustness of tissue-specific developmental
phenotypes is still largely unclear.
73
We have developed the *Arabidopsis* sepal as a system to elucidate the mechanisms
maintaining robustness in organ size and shape[26–28]. Sepals are the outermost floral organs
whose function is to enclose buds and protect the developing inner organs, i.e. petals, stamens,
and carpels, before the flower blooms. To fulfill this protective function, each flower robustly
develops four sepals of equal length, allowing them to close at the top (Figure 1A, top left); these
four sepals are of equal width and positioned 90° from each other, leaving no gap on the sides
(Figure 1A, middle left). This robustness in sepal size and shape stems from the robust initiation
of the four sepal primordia from the floral meristem with precisely coordinated spatiotemporal
patterns[27] (Figure 1A, bottom left). The initiated sepal primordia attain robust final size and shape
by spatiotemporal averaging of cellular growth variability during sepal elongation, and
synchronous progression of a whole-flower growth termination signal from tip to base[26]. In
addition, noise in gene expression must be kept low to ensure sepal size robustness[29]. We
previously characterized a mutant in *DEVELOPMENT RELATED MYB-LIKE 1* (*DRMY1*) that

develops flowers where the inner organs are exposed due to gaps between sepals[27]. The gaps
are caused by variability in sepal development. Specifically, some sepals are shorter than others,
leaving gaps on the top (Figure 1A, top right); the arrangement of sepals around the flower deviate
from the canonical form such that parts of the flower are not covered by a sepal, leaving gaps on
the side (Figure 1A, middle right). This variability in the size, number, and position of the mature
sepal originates from the earliest stages of floral development where the spatiotemporal pattern
of sepal initiation becomes variable (Figure 1A, bottom right). Variability in sepal initiation, in turn,
is driven by the loss of robust patterning of auxin and cytokinin[27], two plant hormones critical for
morphogenesis[30–32], in the floral meristem before sepal initiation. However, the molecular
mechanism through which DRMY1 maintains robust hormone patterning is still unknown.
97
In this study, we elucidate a mechanism through which DRMY1 maintains robust hormone
patterning and thus robust sepal initiation. Specifically, we find that DRMY1 maintains proper
activity of TARGET OF RAPAMYCIN (TOR), a crucial regulator of ribosome level and mRNA
translation[33,34], and thereby sustains translation *in vivo*. When *DRMY1* is mutated, the levels of
ARABIDOPSIS RESPONSE REGULATOR7 (ARR7) and ARABIDOPSIS HISTIDINE
PHOSPHOTRANSFER PROTEIN 6 (AHP6), two cytokinin inhibitor proteins that are rapidly
synthesized during hormone patterning prior to sepal initiation, are drastically reduced in the floral
meristem. Consequently, cytokinin signaling uniformly increases in the meristem periphery,
causing variability in auxin patterning and sepal initiation. We further propose that the increase in
cytokinin signaling may be a survival mechanism to alleviate the translation rate reduction when
ribosomal content is limited. In summary, our work shows that the hub processes of TOR signaling
and translation, which occur in every cell, have very tissue-specific roles in maintaining robust
organogenesis by sustaining the rapid synthesis of hormone signaling proteins.
111
112
113 **RESULTS**
114
115 **The *drmy1* mutant has reduced TOR activity, ribosome content, and translation rate**
116
DRMY1 encodes a MYB/SANT domain protein which may exert transcriptional
regulation[27]. To look for differentially expressed genes in *drmy1* which may be candidates
underlying variable sepal initiation, we performed RNA-sequencing (RNA-seq) in *drmy1* and wild
type (WT) of *apetala1* (*ap1*) *cauliflower* (*cal*) *AP1-GR* background[35,36]. The *ap1 cal AP1-GR*
inflorescence produces numerous tightly packed ball-shaped meristems, which, upon induction,
synchronously initiate sepal primordia, allowing us to collect large quantities of floral meristems
with sepal primordia initiating (Stage 3)[37] (Figure S1A). We crossed *drmy1* into *ap1 cal AP1-GR*
and performed RNA-seq on induced inflorescences of WT and *drmy1* in this background. We
detected transcripts from a total of 21,496 genes, of which 1,042 (4.8%) were differentially
expressed in *drmy1* (Figure S1B; Supplemental Dataset 1). We found that the 443 genes
downregulated at the transcript level in *drmy1* were most enriched in the gene ontology (GO) term
"Translation", a fundamental and ubiquitous cellular process that converts genetic information

from transcript to protein. Within this term, genes encoding ribosomal components were most
downregulated (Figure S1C). The 443 downregulated genes were also enriched in several other
ribosome-related GO terms (Figure 1B). We therefore hypothesized that ribosomal abundance
and translation are affected in *drmy1*, potentially altering the accumulation of proteins critical to
developmental robustness.

To determine whether and how ribosomal abundance and translation are affected in
drmy1, we performed polysome profiling in induced inflorescences of WT and *drmy1* in *ap1 cal*
AP1-GR background. Compared to WT, all peaks corresponding to 40S, 60S, monosomes, and
polysomes are drastically reduced in *drmy1* (Figure 1C; Supplemental Dataset 2). To see whether
this reduction in ribosomal content affected *de novo* protein synthesis rate *in vivo*, we performed
puromycin labeling. Samples were incubated with puromycin, an amino acid-tRNA analog that is
incorporated into nascent polypeptide chains and can be detected using an anti-puromycin
antibody to infer global translation rate[38,39]. In both young seedlings and induced *ap1 cal AP1-GR*
inflorescences, we found that the puromycin level detected in *drmy1* mutant samples was much
reduced compared to WT (Figure 1D), indicating a reduction in global translation rate. We
hypothesized that reduced global translation rate should likely result in globally decreased protein
levels. For this, we looked at a ubiquitously expressed membrane marker *UBQ10::mCherry-*
RCI2A, and found that it had a small (~25%) but significant decrease in fluorescence intensity in
the inflorescence meristem and young floral buds of *drmy1* compared with WT (Figure S1D, E).
We also measured its fluorescence intensity in the ribosomal mutant *ul4y* (*rpl4d*)[40] and we found
that the decrease in fluorescence intensity in *drmy1* is even greater than in *ul4y* (Figure S1F, G).
Overall, these results show that ribosomal content and translation are indeed reduced in the
drmy1 mutant.

To test how the global repression of translation in *drmy1* impacts its proteome, we
extracted total soluble protein from induced inflorescences of WT and *drmy1* in *ap1 cal AP-GR*
background and performed mass spectrometry. We identified a total of 5,077 proteins, of which
548 (10.8%) were differentially accumulated in *drmy1* (Figure S1B; Supplemental Dataset 1).
These differentially accumulated proteins were enriched in GO terms related to translation and
ribosomes (Figure 1E). Despite the overall reduction in ribosomes (Figure 1C), relative to other
proteins, ribosomal components are more abundant in *drmy1* (Figure S1H; Supplemental Dataset
1)*.* This is not true for all proteins involved in translation; poly-A binding proteins and tRNA
synthetases, for example, are relatively less abundant in *drmy1* than in WT. Moreover, the 26S
proteasome responsible for targeted protein degradation is much more abundant in *drmy1* than
in WT (Supplemental Dataset 1). In concert, these results demonstrate that the machinery
responsible for maintaining protein homeostasis is substantially dysregulated in *drmy1*.

A key signaling pathway that regulates protein homeostasis is TARGET OF RAPAMYCIN
(TOR)[41,42]. TOR is a hub that integrates information from light, sugars, nutrient availability, etc.,
to promote growth-related processes, including ribosome biogenesis and translation, and to
repress catabolic processes, including protein degradation by autophagy and the

proteasome[33,34,43–45]. TOR directly regulates the translation of specific mRNAs by promoting the
phosphorylation of proteins, including LARP1, eIF3h, RISP, eS6, and 4EBPs, that impact
translation initiation, reinitiation, or elongation of mRNAs with specific features, such as 5'TOP
motifs or short upstream open reading frames (uORFs) in the 5' leaders of mRNAs[34,42,46–52]. TOR
also indirectly increases overall protein synthesis rates by globally increasing ribosome levels.
We therefore hypothesize that the overall decrease in ribosomal content and protein synthesis in
drmy1 may reflect altered TOR signaling. To test for signatures of transcriptomic changes that
have been well defined in seedlings under TOR inhibition[34,53,54], we performed RNA-seq on
seedlings of WT, *drmy1*, WT treated with AZD-8055 (a potent TOR inhibitor), and mock-treated
WT (Supplemental Dataset 3). We found that the *drmy1* mutation causes transcriptomic changes
similar to TOR inhibition (Figure 1F). A significant portion of genes differentially expressed under
TOR inhibition vs. mock were also differentially expressed in *drmy1* vs. WT (466/2044 = 22.8%;
hypergeometric test, $p = 4.7 \times 10^{-108}$). Not only were these 466 genes differentially expressed in
both situations, but also most of them were coherently downregulated or upregulated (439/466 =
94.2%, Chi-square test, $p < 2.2 \times 10^{-16}$; Figure 1G, S1I). Genes coherently downregulated in both
situations were enriched in GO terms related to translation and ribosomes, and, most strikingly,
a quarter of them were under the GO term "translation" (Figure 1F, S1J). These similar
transcriptomic changes support our hypothesis that TOR activity is reduced in *drmy1*. To further
test this hypothesis, we measured TOR activity in WT and *drmy1* by assaying the phosphorylation
of its direct substrate, RIBOSOMAL PROTEIN eS6 KINASE (S6K)[55,56]. While the total protein
level of S6K did not change in *drmy1*, we found that S6K phosphorylation drastically decreased,
demonstrating reduced TOR activity (Figure 1H, I). Overall, these results are consistent with the
idea that *drmy1* has reduced TOR activity—a main pathway controlling ribosomal abundance and
translation—which causes reduced ribosomal content and global translation rate.
195

Defects in TOR activity, ribosome, and translation disrupt robust sepal initiation

We next asked whether defects in TOR activity, ribosome, or translation have any effects
on robust sepal initiation like the *drmy1* mutation does (Figure 2A, 2B; also see Zhu et al.[27]). In a
WT bud, initiation is robust in that four sepal primordia of similar size form evenly spaced around
the periphery of the floral meristem (Figure 2A, 2H). Angles between them vary little, i.e., they are
all at around 90° angles from each other (Figure 2I, 2J). By contrast, in *drmy1* buds, three to five
sepal primordia initiate and grow to different sizes (Figure 2B, 2H; also see Zhu et al.[27]). Sepal
primordia in *drmy1* buds are generally unevenly spaced, and angles between them have a high
coefficient of variation (CV) (Figure 2I, 2J). To determine whether defects in ribosomes can cause
the same sepal initiation defects, we imaged three ribosomal mutants, *ul4z* (*rpl4a*), *ul4y*, and *ul18z*
(*rpl5a*)[40], each mutated in a gene encoding a ribosomal component that is also downregulated in
drmy1 at the transcript level (Figure S1C). The *ul4z* mutant bud shows reduced size of the inner
sepal primordia relative to the outer sepal primordia (Figure 2C), and slightly more variable spatial
distribution of sepal primordia (Figure 2J), although it always develops four sepal primordia
(Figure 2H). This is a weaker phenotype than *drmy1* but has similar characteristics. The *ul4y* and
ul18z mutants show great variability in the number and position of sepal primordia (Figure 2D,

2E, 2H, 2J), more similar to *drmy1*. We also crossed these ribosomal mutants with *drmy1* to study
sepal variability in the double mutants (Figure S2A-H). In *drmy1 ul4z*, *drmy1 ul4y*, and *drmy1*
ul18z/+, on average, sepal initiation was as variable as in the *drmy1* single mutant (Figure S2I,
S2J). However, there were buds with no outgrowth in the adaxial or lateral regions of the bud
periphery (Figure S2B, S2E, S2G), buds with six sepal primordia (Figure S2C, S2F, S2H), and
buds with two outer sepal primordia (Figure S2D, S2H), which were not seen in the *drmy1* single
mutant. Note that we were unable to characterize the homozygous *drmy1 ul18z* double mutant
because they were embryo-lethal (Figure S2K), further supporting the idea that ribosomal
mutations enhance the phenotypic defects in *drmy1*.
222
We then imaged mutants with reduced TOR activity to determine whether sepal initiation
is also less robust. *lst8-1-1* is a T-DNA insertional mutant of the TOR complex component LST8-
1[57] and is weakly hypomorphic in TOR activity. We found that *lst8-1-1* shows variable sepal
initiation in a small proportion of buds (4/41, 9.8%) (Figure 2F, 2H, 2J). The *spaghetti-1* mutant
defective in TOR complex 1 (TORC1) assembly[58] showed a level of variability comparable to the
drmy1 mutant and the ribosomal mutants *ul4y* and *ul18z* (Figure 2G, 2H, 2J). Mutants with more
severe disruption of TOR activity are embryo lethal and could not be analyzed[58,59]. These results
show that reduction in TOR activity can cause variability in sepal initiation, similar to *drmy1*.
231
To corroborate these findings, we directly inhibited translation by *in vitro* culture of
dissected WT inflorescences on 2 µM cycloheximide (CHX, a chemical inhibitor of translation) for
9-10 days. This is a low concentration that does not completely block translation, as
inflorescences were still alive after 10 days in this condition. Compared with mock, CHX-treated
inflorescences develop buds that have 2 to 6 sepal primordia of variable sizes that are unevenly
spaced around the bud periphery (Figure 2K, 2M, 2N). These phenotypes are stronger than
drmy1. Similarly, we directly inhibited TOR activity by application of 2 nmol Torin2 to the growing
shoot apex twice a day for 15 days, and we observed variable sepal initiation (Figure 2L, 2M, 2N).
Overall, these data show that inhibition of TOR activity and translation can disrupt the robustness
of sepal initiation, in terms of sepal primordium number, position, and size.
242
We previously showed that *drmy1* buds develop sepals of different sizes because of
increased differences in the initiation timing of sepals within the same bud. The late-initiating
primordia remain smaller throughout development. They end up as smaller sepals relative to
those that initiated earlier, leaving gaps that expose the developing inner floral organs[27].
Moreover, different buds have different temporal patterns of sepal initiation, contributing to
between-bud variability of sepal phenotype[27]. We asked whether TOR or ribosomal defects
similarly disrupt the relative timing of sepal initiation, within-bud and between-bud. We live imaged
WT and *ul4y* every six hours during sepal initiation and quantified the amount of time the bud
takes to initiate the inner and lateral sepals after it initiates the outer sepal. In WT, after the
initiation of the outer sepal, most buds initiate the inner sepal within 6 hours and the lateral sepals
within 12 hours (Figure 3A, 3C; also see Zhu et al.[27]). In *ul4y*, the time differences between outer
and inner sepal initiation and between outer and lateral sepal initiation are longer (Figure 3B,

mean in Figure 3C). Moreover, these time differences are more variable across buds, indicating
a loss of robustness in organ initiation timing (SD in Figure 3C). Similarly, we compared the sepal
initiation timing in Torin2 vs mock-treated WT buds. While in most mock-treated buds, the inner
and lateral sepals initiate within 12 hours after the outer sepal (Figure 3D, 3F), in Torin2-treated
buds, sepals within the same bud initiates at vastly different times (Figure 3E, mean in Figure 3F),
and the temporal pattern of sepal initiation is more variable across buds than mock (SD in Figure
3F). These results show that TOR and ribosomal defects can disrupt the precisely orchestrated
initiation timing of sepal primordia.
263

Does the variability in initiation timing cause variable sizes and gaps in mature sepals, as
in *drmy1* (Figure S3A, B, G, H; also see Zhu et al. [27])? We imaged the mature sepals of the
ribosomal mutants *ul4z*, *ul4y*, *ul18z*, as well as the TOR component mutant *lst8-1-1*. Surprisingly,
unlike *drmy1*, the sepals in *ul4z*, *ul4y*, *ul18z* enclose the inner floral organs perfectly, leaving no
gaps, regardless of sepal number (Figure S3C-E). Small gaps still exist in buds of *lst8-1-1*,
although sepal size differences appear greatly reduced (Figure S3F). Further dissection shows
that in these mutants, sepals within the same flower are of similar sizes, although sepals from
different flowers can be of vastly different sizes, most conspicuously for *lst8-1-1* (Figure S3I-N).
This is unlike *drmy1*, where sepal size variability is equally high comparing sepals within the same
flower or from different flowers (Figure S3H, S3M-N). Upon closer examination, while sepals
initiating late in *drmy1* buds remain small, leaving a gap in the sepal whorl (Figure S3O-P), those
in *ul4y* were able to catch up with the other sepals and close the gap (Figure S3Q). Our results
suggest that there exists a size-coordinating mechanism independent of TOR or ribosomal
functions that allows sepals within the same bud to reach the same mature length, and that this
mechanism is disrupted in *drmy1*. Such a mechanism requires further investigation in the future.
279

280   **Inhibition of TOR activity and translation increase cytokinin signaling and disrupts the**
281   **robust spatial pattern of auxin and cytokinin signaling**
282

Auxin and cytokinin are two important plant hormones critical to many aspects of plant
development[30–32], and there is accumulating evidence that they act synergistically in the shoot
apical meristem to promote lateral organ initiation[16,60,61]. We previously showed that, in a WT floral
meristem prior to sepal initiation, auxin and cytokinin signaling are concentrated at the four
incipient primordia, which is required for robust sepal initiation from these regions (Figure 4A,
S4A; Zhu et al.[27]). In the *drmy1* mutant, cytokinin signaling becomes stronger and diffuse around
the bud periphery (Figure 4A-B). Auxin signaling also becomes more diffuse, forming irregular
auxin maxima that are less focused than those in WT, except at the incipient outer sepal where it
remains robust (Figure 4A, S4B; Zhu et al.[27]). These changes in hormone signaling correlate with
variable sepal initiation (Figure S4B)[27]. We wondered whether ribosomal mutations have similar
effects on auxin and cytokinin signaling. To this end, we imaged the auxin signaling reporter
DR5::3xVENUS-N7 and the cytokinin signaling reporter *TCS::GFP* in floral meristems of the
ribosomal mutant *ul4y*. Both reporters lose their robust spatial pattern except in the incipient outer
sepal (Figure 4A, S4C). The hormone signaling patterns were quantified by circular histogram

analysis (see Methods for details). For each of DR5 and TCS, WT buds showed four clear peaks
~90 degrees apart from each other, with very little signal in between, whereas in *drmy1* and *ul4y*,
peaks were barely seen except at the incipient outer sepal (at 45 degrees), and there was greater
noise and variation all around the bud (Figure 4C-D). Diffuse bands of auxin signaling that typically
occurs in the adaxial or lateral periphery of *drmy1* and *ul4y* buds (Figure S4B and S4C, brackets)
can later resolve into several distinct auxin maxima of various intensity and at various positions,
correlated with the initiation of sepal primordia of various sizes at these same positions (Figure
S4B and S4C, red arrowheads).
305
We also tested whether drug treatments that inhibit TOR activity or translation can disrupt
the robust hormone patterning. Buds treated *in vitro* with the translation inhibitor CHX (2 µM) for
3 days showed a 50% increase in cytokinin signaling, and both auxin and cytokinin signaling
became diffuse around the bud periphery (Figure 4E-H). By day 6, cytokinin signaling was still
diffuse all around, and increased to more than two-fold relative to mock (Figure 4I, 4J, 4L). Auxin
signaling formed maxima of variable number at variable positions (Figure 4I arrowheads, 4K),
correlated with the variable initiation of sepal outgrowth at these positions (Figure S4D-E). Similar
changes occurred in buds treated *in vitro* with the TOR inhibitor AZD-8055 (2 µM) for 6 days
(Figure 4I-L). For both CHX and AZD-8055, the disruptions of hormone signaling are similar to
drmy1. *In vivo* treatment using another TOR inhibitor Torin2 for 15 days increased cytokinin
signaling by 70%, although it did not make auxin and cytokinin signaling more diffuse (Figure
S4F-I). Overall, these results show that defects in TOR activity and translation increase cytokinin
signaling, and disrupt the precise spatial patterning of cytokinin and auxin signaling required for
robust sepal initiation.
320

An increase in cytokinin signaling is necessary and sufficient for variable auxin signaling
and sepal initiation under translation inhibition

Auxin is a critical hormone in organogenesis[62,63]. As shown above, variable patterning of
auxin signaling correlates with variable sepal initiation during inhibition of TOR activity and
translation. We wondered what caused auxin to lose its robust patterning under such conditions.
It was previously reported that the ribosomal mutants *ul4y*, *ul18z*, and *el24y* have reduced protein
levels of AUXIN RESPONSE FACTOR (ARF) 3, 5, and 7[64–66], key transcription factors that
mediate the auxin signaling response[67]. The transcripts of these ARFs contain upstream open
reading frames (uORFs), requiring translation reinitiation to translate their main open reading
frames[68,69], a process defective in the ribosomal mutants *ul4y*, *ul18z*, and *el24y*[64–66]. We therefore
hypothesized that *drmy1* loses robust auxin signaling pattern because of reduced translation of
uORF-containing transcripts, including those of certain ARFs. To begin, we utilized our
transcriptomics and proteomics data, and considered that the protein-transcript ratio of a gene
should reflect its level of translation, among other factors such as protein stability. Therefore,
following our hypothesis, genes containing uORFs should, in general, have a lower protein-
transcript ratio in *drmy1* than in WT. We calculated the difference of this ratio between *drmy1* and
WT for all 5,086 gene-protein pairs in our inflorescence dataset, and compared the ratio against

the number of uORFs in each transcript (Figure S5A; uORF data from von Arnim et al.[69]). We
found a small but significant decrease in the protein-transcript ratio in *drmy1* for the 724 genes
containing at least 2 uORFs in their transcripts, supporting the hypothesis that *drmy1* has reduced
translation reinitiation for uORF-containing transcripts, just like *ul4y*, *ul18z*, and *el24y*[64–66]. Then,
we examined whether the translation reinitiation of uORF-containing ARFs are indeed reduced in
the *drmy1* mutant. We selected *ARF3/ETTIN*, *ARF5/MONOPTEROS*, and *ARF6*, which have 2,
6, and 6 uORFs respectively, and as controls, *ARF8* and *ARF10* which do not contain uORFs.
None of these *ARF*s were differentially expressed in *drmy1* at the transcript level, except *ARF10*
which was slightly upregulated (Figure S5B). We utilized promoter-fluorescent protein fusion
reporters (Figure S5C) which have the same uORFs in the promoter region as the corresponding
ARF genes if the genes have them. These reporters reflect transcriptional and uORF-mediated
translational regulation. *pARF3::N3xGFP*, *pARF5::ER-EYFP-HDEL*, and *pARF6::N3xGFP*
contain uORFs and thus, following our hypothesis, are expected to drastically decrease in
fluorescence intensity in *drmy1* compared to WT. *pARF8::N3xGFP* and *pARF10::N3xGFP* do not
have uORFs and are thus expected to have comparable or higher fluorescence intensity in *drmy1*.
Surprisingly, we saw no correlation between the presence of uORFs and decrease in fluorescent
intensity in *drmy1* (Figure S5C-D). While it might arise from additional layers of regulation on
these ARFs, this result suggests that the decrease in translation reinitiation of uORF-containing
ARFs is not the main factor explaining the loss of robust auxin signaling pattern in *drmy1*.
358

It was previously reported that external application of cytokinin increases auxin
biosynthesis in actively growing tissue including the shoot apex, young leaves, and roots[70], and
cytokinin application also changes the expression and polarity of PIN-FORMED (PIN) polar auxin
transport carriers[71,72]. We previously noticed that external application of 6-benzylaminopurine
(BAP), a synthetic cytokinin, induced additional convergence points of PIN1 and increased
variability in auxin signaling, causing variability in sepal initiation (Zhu et al.[27], in this reference
see Fig. 4e, Extended Data Fig. 7e and 7f). Here, we confirmed this observation by circular
histogram analysis (Figure 5A-D). While the mock-treated WT buds showed four clear peaks of
DR5 signal with very little signal in between (Figure 5A-B), those treated with 5 µM BAP showed
a less robust spatial pattern, with less distinguishable peaks and larger variation all around the
bud (Figure 5C-D). Thus, excessive cytokinin is sufficient for the variable spatial pattern of auxin
signaling.
371

We then wondered whether an increase in cytokinin signaling (Figure 4) is the cause of
variable pattern of auxin signaling under translation-limited conditions such as *drmy1*. To test this
hypothesis, we crossed *drmy1* containing the DR5 reporter with a triple mutant of *ARABIDOPSIS*
RESPONSE REGULATOR (*ARR*) *1*, *10*, and *12*, the three most highly expressed B-type ARRs
in our RNA-seq (Supplementary Dataset 1) which are crucial for the activation of cytokinin-
responsive genes[73]. While buds of *arr1,10,12* did not show apparent phenotypic differences from
WT, the quadruple mutant *drmy1 arr1,10,12* largely rescued the *drmy1* phenotype, with much
less variability in sepal number and position (Figure 5E-G). While mature buds of *drmy1* have
sepals of variable sizes, leaving gaps and exposing the inner floral organs (Figure S6D vs. S6A-

C), those of *drmy1 arr1,10,12* have sepals of robust sizes that are able to close (Figure S6E).
Likewise, mutation in a cytokinin receptor *WOODEN LEG* (*WOL*)/*ARABIDOPSIS HISTIDINE*
KINASE 4 (*AHK4*) showed a similar rescue of the *drmy1* sepal phenotype (Figure 5E-G, S6F).
While the auxin signaling reporter DR5 was diffuse and variable in *drmy1* except in the incipient
outer sepal (Figure 5H-I), in *drmy1 arr1,10,12*, it was focused in all the four incipient sepals that
were robustly positioned, although the signal intensity in the incipient outer sepal was much higher
than others (Figure 5J-K). These results indicate that cytokinin signaling is required for the
increased variability in auxin signaling pattern and sepal initiation in *drmy1*.
389
Furthermore, we wanted to test whether cytokinin signaling is required for variability in
more general conditions where translation is inhibited. The translation inhibitor CHX disrupted
robustness in auxin signaling and sepal initiation in WT (Figure 2K, 4E, 4I), and we tested whether
these effects are still present in *arr1,10,12* and *wol* mutants. We found that, unlike WT, sepal
initiation remained mostly robust in *arr1,10,12* and *wol* after ten days of 2 µM CHX treatment
(Figure 5L-N). While DR5 in WT became diffuse and occurred in variable positions after three
days of CHX treatment (Figure 5O-P, arrow), DR5 in *arr1,10,12* remained robust and
concentrated at the four incipient sepal primordia (Figure 5Q-R). These results suggest that
elevated cytokinin signaling level is the primary cause for variability in auxin patterning under
translation-inhibited conditions. Thus, in WT, maintaining a low level and focused cytokinin
signaling is crucial for robust auxin patterning and sepal initiation.
401
402 **Upregulation of cytokinin signaling is required to sustain translation and fitness in *drmy1***
403
Under translation-inhibited conditions, why does the plant upregulate cytokinin signaling
at the cost of robust morphogenesis? Previous studies revealed that cytokinin signaling can
stimulate translation[74–78], by increasing transcription or protein abundance of ribosomal
components or biogenesis factors[79–81] and modification of initiation and elongation factors[82]. We
therefore hypothesized that an increase in cytokinin signaling under translation-inhibited
conditions (such as *drmy1*) sustains a survivable rate of translation in a feedback loop. We first
validated that, under our growth conditions, an increase in cytokinin signaling (*arr1 35S::ARR1*)
is sufficient to increase global translation (Figure 6A; also see Karunadasa et al.[74]) in 14-day-old
seedlings. We then tested whether cytokinin signaling is required to sustain global translation
(Figure 6B-C). Compared to WT, the cytokinin receptor single mutant *wol* has a mild reduction in
global translation rate at day 8 and a ~50% reduction at day 14. The *drmy1* single mutant shows
drastically reduced global translation rate at day 8, but by day 14, global translation rate in *drmy1*
increased and matched WT. In the *drmy1 wol* double mutant, however, translation rate was
unable to recover at day 14 and remained lower than *drmy1*. Our data suggest that, in *drmy1*
plants which has reduced TOR activity and ribosomal level (Figure 1), the upregulated cytokinin
signaling is required to sustain global translation at nearly WT levels.
420
We then hypothesized that an upregulation of cytokinin signaling in plants with reduced
TOR activity and ribosomal content such as *drmy1* and the consequent restoration of global

translation would provide fitness benefits. Thus, we expect that removal of the cytokinin receptor
WOL from *drmy1* and the consequent failure to sustain global translation should affect plant
vitality and reproduction. We found that at day 14, the *drmy1* single mutant is slightly smaller than
WT. In contrast, in the absence of WOL, growth of *drmy1 wol* plants were extremely retarded
compared to *wol*, with tiny and chlorotic cotyledons and true leaves (Figure 6D). In older plants,
the *drmy1* single mutant has similar rosette size and slightly shorter inflorescences compared to
WT. In contrast, in the absence of WOL, *drmy1 wol* produced tiny rosettes and stunted
inflorescences with a few chlorotic buds that develop into small, short siliques (Figure 6E, S6F).
Similarly, when B-type ARRs are mutated, the growth of the *drmy1 arr1,10,12* quadruple mutant
is much retarded compared to *arr1,10,12*. They produced slightly chlorotic and anthocyanin-rich
rosettes, a tiny inflorescence composed of very few buds (Figure 6F, S6E) and, in the end, siliques
in which all seeds had aborted (Figure 6G). Overall, these results show that *drmy1* requires the
cytokinin signaling pathway for normal growth and reproduction. While it remains possible that
unknown effects of the cytokinin signaling pathway other than promoting translation are critical
for the proper growth of *drmy1*, our results are consistent with our hypothesis that the upregulation
of global translation (Figure 6A) by increased cytokinin signaling (Figure 4) maintains a survivable
level of protein synthesis in plants with reduced ribosomal content such as *drmy1* (Figure 1C).
440
441 **TOR and translation inhibition decreases the protein level of cytokinin signaling inhibitors**
442 **ARR7 and AHP6**
443
What causes cytokinin signaling to increase in plants with reduced TOR activity and
translation (Figure 4)? It was previously shown that *cis*-type cytokinins can be synthesized from
tRNAs by the tRNA isopentenyltransferases (IPTs), IPT2 and IPT9[83]. We hypothesized that the
decrease in global translation may increase the availability of tRNAs as substrates for cytokinin
biosynthesis, increasing the level of cytokinins. To test this idea, we extracted cytokinins from
induced inflorescences of WT and *drmy1* in *ap1 cal AP1-GR* background (Figure S1A). We
measured the level of three cytokinin bases, *trans*-Zeatin (tZ), *cis*-Zeatin (cZ), and
isopentenyladenine (iP), and their corresponding nucleosides (tZR, cZR, and iPR), using liquid
chromatography-mass spectrometry. Surprisingly, we found no significant difference in their
levels between WT and *drmy1*, and notably, the amount of *cis*-Zeatin was barely detectable in all
samples (Figure S7A). This suggests that the increase in *cis*-type cytokinin synthesis is not the
mechanism underlying the increase in cytokinin signaling under our translation-inhibited
conditions.
457
We then considered the effects that a decrease in translation rate might have on the
protein components of the cytokinin signaling pathway. In particular, A-type *ARR*s, which encode
inhibitors of cytokinin signaling[84–86], are rapidly induced upon cytokinin application and serve to
dampen cytokinin response in the tissue[87–89]. Likewise, *AHP6* is highly expressed in lateral organ
primordia downstream of auxin signaling, which non-cell autonomously represses and restricts
cytokinin signaling to robust spatial patterns[16]. The rapid synthesis of the A-type ARR and AHP6
proteins may be crucial for maintaining the homeostasis of cytokinin signaling during

developmental processes. We therefore hypothesized that, during hormone patterning prior to
sepal initiation, translation defects in *drmy1* cause reduced synthesis of these proteins,
decreasing them to a level insufficient to repress cytokinin signaling (Figure 7A).

We started testing the level of protein reporters for a variety of cytokinin signaling
components, in *drmy1* vs. WT and/or under translation or TOR inhibition. We reasoned that, given
that the transcript level of most cytokinin signaling components do not significantly differ between
drmy1 and WT inflorescence tissue (Figure S7B), any changes in the level of these protein
reporters should reflect post-transcriptional regulation, including mRNA translation. For A-type
ARRs, we were unable to detect fluorescence in the inflorescence of a published *pARR4::ARR4-*
GFP line[90]. We reasoned that this was because A-type ARRs have low protein levels (none was
detected in our proteomics dataset) and short half-lives[91]. We therefore employed LlamaTagging,
a recently developed method to visualize the abundance of nuclear-localized proteins with short
half-lives[92]. Rapidly degraded proteins cannot be visualized through fusion with standard
fluorescent proteins, because fluorescent proteins take time to mature before they fluoresce, and
the protein of interest is degraded before the maturation of the fluorescent protein. On the other
hand, the LlamaTag folds immediately. A LlamaTag with a high affinity for GFP can be encoded
as a translational fusion with a nuclear-localized protein of interest. Soon after translation, the
fusion immediately binds cytoplasm-localized GFP and translocates it to the nucleus. Thus,
increased GFP fluorescence in the nucleus indicates higher abundance of the protein of interest
(Figure 7B).

We focused on ARR7, the most highly expressed A-type ARR in our inflorescence RNA-
seq (Figure S7B) which was also shown to be nuclear-localized[93,94]. We designed a construct with
ARR7 fused with GFP-specific LlamaTag by a short linker, driven by the ARR7 native promoter
(*pARR7::ARR7-linker-llama-ARR7ter*; *ARR7-llama* for short). This construct was co-transformed
with cytoplasm-localized GFP containing a nuclear exclusion signal (*pUBQ10::sfGFP-nes-*
UBQ3ter; *GFP-nes* for short; Figure 7C). As a proof of concept, we treated this reporter in WT
background with 200 μM BAP. We found that GFP signal became more nuclear-localized within
5 hours of the treatment (Figure S7C-D), agreeing with an increased expression and stability of
A-type ARR proteins upon cytokinin application as previously reported[91,95].

We then compared the localization of GFP signal in floral meristems of WT and *drmy1*
before sepal initiation. According to the null hypothesis, increased cytokinin signaling in *drmy1*
would cause an increase in ARR7 protein level and thus more nuclear-localized GFP signal in
drmy1 ARR7-llama GFP-nes than its WT counterpart. This is because cytokinin signaling
increases the gene expression and protein stability of A-type ARRs[88,91,95,96]. In contrast, according
to our hypothesis, insufficient protein synthesis of A-type ARRs contributes at least in part to
increased in cytokinin signaling in *drmy1*, so we expect to see reduced ARR7 protein level and
thus more cytoplasm-localized GFP signal in *drmy1 ARR7-llama GFP-nes*. We found that WT
buds had slightly more nuclear-localized GFP signal than cytoplasm-localized GFP signal, with
brighter spots corresponding to the nucleus surrounded by darker grooves in between

corresponding to the cytoplasm (Figure 7D,E). In contrast, in the periphery of *drmy1* buds, GFP
signal localizes more to the cytoplasm than to the nucleus, with darker spots surrounded by
brighter grooves (Figure 7D,E). More nuclear GFP was present near the center of *drmy1* buds.
This result indicates nuclear ARR7 protein concentration is reduced in the *drmy1* mutant,
particularly in the zone where sepals initiate. To see whether this conclusion holds in other
translation-inhibited conditions, we treated WT plants carrying the *ARR7-llama* and *GFP-nes*
reporters with the translation inhibitor CHX and the TOR inhibitor AZD-8055. 2 µM CHX treatment
for 24 hours drastically reduced the nuclear localization of the GFP signal and increased its
cytoplasmic localization (Figure 7F). 2 µM AZD-8055 treatment for 72 hours had a milder but
similar effect (Figure 7G). These treatments did not affect the localization of the GFP signal in
plants without *ARR7-llama* (Figure S7E-H). These results show that conditions that decrease
global protein synthesis greatly decrease the nuclear level of ARR7 protein.
519
We also tested whether TOR or translation inhibition alter the protein level of AHP6. To
this end, we imaged the *pAHP6::AHP6-VENUS*[16] protein reporter under mock, CHX, or AZD-8055
treatment. While mock-treated buds highly accumulate the AHP6 protein in the four incipient sepal
primordia, buds treated with CHX or AZD-8055 abolished AHP6 accumulation within 72 hours
(Figure 7H). The *pAHP6::GFP-ER*[97] transcriptional reporter does not change under these
treatments (Figure 7I), in agreement with our RNA-seq data of WT vs. *drmy1* (Figure S7B),
suggesting that the change in AHP6 protein level is due to post-transcriptional regulation such as
altered translation.
528
It is possible that inhibition of TOR and translation results in a general, uniform reduction
in the level of all proteins, not just for the cytokinin signaling inhibitors ARR7 and AHP6, but also
for the positive regulators of cytokinin signaling. For this, we tested whether AHP3, a component
of the cytokinin phosphorelay[98,99], respond to TOR and translation inhibition. CHX treatment did
not affect the level or spatial localization pattern of the *pAHP3::AHP3-GFP* reporter[100], while AZD-
8055 treatment only mildly decreased its level (Figure S7I). We also tested how the level of other,
more generic proteins respond to TOR and translation inhibition. Unexpectedly, the level of
pUBQ10::mCherry-RCI2A increased upon 72 hours of CHX or AZD-8055 treatments (Figure
S7J). Overall, these results show that TOR and translation inhibition does not result in a uniform
reduction in the level of all proteins, but specific proteins such as ARR7 and AHP6 are dramatically
decreased. Further, these results are consistent with our hypothesis that depletion of cytokinin
signaling inhibitor proteins, including ARR7 and AHP6, may underlie the upregulation of cytokinin
signaling when the floral meristem is under TOR or translation inhibition (Figure 4).
542
Having shown a reduction in the level of ARR7 and AHP6 greater than other proteins
under TOR and translation inhibition, we next tested whether such a reduction contributes to the
variability in sepal initiation also seen under such conditions (Figure 2). High-order mutant of A-
type *ARR*s (*arr3,4,5,6,7,8,9,15*[101]) shows reduced size of the inner sepal primordium (Figure 7J,K)
and a minor but significant increase in the positional variability of sepal primordia compared to
WT (Figure 7Q), although sepal primordium number remains robust (Figure 7P). The *ahp6*

mutant[16] shows great variability of sepal primordium number, position, and size, although to a
lower extent than *drmy1* (Figure 7J,L,P,Q). Similarly, if the reduction of ARR7 level in *drmy1*
contributes to variability in sepal initiation, increasing ARR7 expression should restore sepal
initiation robustness. While *ARR7-llama GFP-nes* plants phenocopy WT (Figure 7M), introducing
the *ARR7-llama GFP-nes* constructs into *drmy1* plants partially restores robustness in sepal
initiation, particularly in the position of sepal primordia (Figure 7N-Q). In older buds, while *drmy1*
buds show great variability in sepal number and size resulting in gaps, some buds of *drmy1 ARR7-*
llama GFP-nes have robustly sized sepals that are able to close properly (Figure S7K-N). Overall,
these results show that reducing the level of cytokinin signaling inhibitor proteins ARR7 and AHP6
create variability in sepal primordium initiation, and increasing their level in *drmy1* partially
restores robustness. We propose that, during hormone patterning prior to sepal initiation, the rapid
synthesis of these inhibitor proteins in response to auxin and cytokinin signaling is crucial for
maintaining the homeostasis of cytokinin signaling and thus the robustness in sepal initiation.
562

We also considered other hormone-related proteins that are dynamically regulated during
organogenesis and thus may be depleted under translation defects. AUXIN/INDOLE-3-ACETIC
ACID INDUCIBLE (Aux/IAA) proteins are auxin signaling inhibitors that are rapidly induced by
auxin[102,103]. They bind auxin and are rapidly degraded by the ubiquitin E3 ligase SKP1, CUL1, F-
BOX PROTEIN (SCF) complex involving TRANSPORT INHIBITOR RESPONSE1/AUXIN
SIGNALING F-BOX (TIR1/AFB)[104–107]. Degradation is dependent on the Short Linear Motif (SLiM)
degron contained within Domain II (DII)[106–108]. We hypothesized that the level of DII-containing
proteins including Aux/IAAs would be drastically decreased in translation-inhibited conditions
such as *drmy1* because they are unable to be rapidly synthesized to keep up with their
degradation upon auxin signaling. To test this, we used the R2D2 reporter[109], which contains a
DII fused with 3xVENUS (*pUS7Y::DII-n3xVENUS*), and as a control, a mutated non-degraded DII
fused with tdTomato (*pUS7Y::mDII-ntdTomato*). We compared this reporter in *drmy1* vs. WT. The
ratio of *VENUS* to *tdTomato* was not reduced in *drmy1*, but instead slightly but significantly
elevated (Figure S7O-P). In addition, *drmy1* has stochastic patches of DII-VENUS degradation,
consistent with its often mislocalized auxin maxima (Figure 4A, S4B), unlike WT which had four
patches of degradation corresponding to the four incipient sepal primordia where auxin maxima
robustly form (Figure 4A, S4A). Overall, these results suggest that the level of DII-containing
Aux/IAA proteins is not reduced in *drmy1*, despite the high requirement for synthesis due to their
rapid turnover. They also indicate that not all proteins that are dynamically regulated in response
to hormone signaling are equally affected by translation inhibition, which may result in different
changes in hormone signaling output under such condition.
584

DISCUSSION

Robustness, the strikingly reproducible development of phenotype, has fascinated
biologists for decades[2]. The Arabidopsis flower robustly develops four sepals of equal size. This
stems from the robust initiation of four sepal primordia from the floral meristem, which is in turn

dictated by the robust patterning of auxin and cytokinin controlled by DRMY1[27]. Here we
elucidated how DRMY1 controls robust hormone patterning and thus robust sepal initiation. We
show that DRMY1 sustains TOR activity, ribosomal content, and translation. We further show that
inhibition of TOR activity or translation is sufficient to cause variability in the timing, position, and
number of sepal primordia, mimicking the *drmy1* phenotype. Our findings are in concert with
previous studies that have shown robustness is often maintained by genes involved in central
cellular processes[2]. In our case, the rate of translation in wild type maintains proper levels of
ARR7 and AHP6, two cytokinin signaling inhibitor proteins, which need to be rapidly synthesized
to dampen cytokinin signaling. Homeostasis of cytokinin signaling ensures robustness in auxin
signaling patterns, and thus robustness in sepal initiation (Figure 7R, top). In the *drmy1* mutant,
the reduced TOR activity, ribosomal content, and translation rate causes inability to rapidly
synthesize these inhibitor proteins. Consequently, cytokinin signaling is elevated, disrupting the
robust spatial pattern of auxin signaling, leading to variable sepal initiation (Figure 7R, bottom).
Blocking cytokinin signaling in *drmy1* is sufficient to restore robust initiation of four sepal
primordia, but has severe consequences on the overall fitness of the plant. Our results reveal how
defects in hub cellular processes such as TOR signaling and translation can have tissue-specific
phenotypic effects.
608

Although we propose that reduced TOR activity and mRNA translation affects
developmental robustness through reduced synthesis of ARR7 and AHP6, we do not exclude
other potential mechanisms that could contribute to the drmy1 phenotype. For example, we
observed that several subunits of the 26S proteasome are more abundant in drmy1 than in WT
(Supplemental Dataset 1), which could reflect or influence the accumulation of proteotoxic
peptides and disrupt protein homeostasis[110–114]. This accumulation of 26S proteasomes could
function upstream and/or downstream of the defect in TOR activity that impacts robust
organogenesis[34,115–120]. Substantial future research will be needed to comprehensively define
how DRMY1 participates in the complex interactions among TOR, mRNA translation, and
proteolysis.
619

It was discovered long ago that extrinsic cytokinin application to plant tissue or cell-free
extracts can promote mRNA translation[75–78]. Recent studies further confirmed that the up-
regulation of translation by cytokinin is at least in part mediated by the cytokinin signaling
pathway[74,82]. Here, we show that cytokinin signaling in floral buds is upregulated in translation-
inhibited conditions, such as *drmy1*, AZD-8055 treatment, or CHX treatment (Figure 4), through
reduced level of cytokinin inhibitor proteins (Figure 7; also potentially through other untested
mechanisms). The enhanced cytokinin signaling maintains translation rate at a level necessary
for the survival and reproduction of the plant (Figure 6). We propose that this represents a
homeostasis mechanism where plants leverage increased cytokinin signaling to rescue the
translation rate reduction caused by deficient TOR activity and ribosomal content (Figure 7R,
bottom). It remains to be tested how widely this mechanism is applicable to other mutants with
ribosomal defects, or whether parallel mechanisms operate in other species across kingdoms.
632

633   While translation-inhibited plants likely upregulate cytokinin signaling to maintain protein
634 synthesis, this upregulation negatively affects developmental robustness. We have previously
635 shown that exogenous cytokinin application to the WT floral meristem increases variability in PIN1
636 convergence and auxin signaling patterns, and consequently, in sepal initiation. These effects are
637 more pronounced in the *drmy1* mutant, which by itself has increased and diffuse cytokinin
638 signaling[27]. Here, we provide additional evidence that increased and diffuse cytokinin signaling is
639 necessary for such variability. While *drmy1* and CHX-treated WT floral meristems are variable in
640 auxin signaling pattern and sepal initiation (Figure 2, 4), mutations in *wol* and *arr1,10,12*, which
641 decreases cytokinin signaling, largely restore robustness (Figure 5). Robustness is also restored
642 in the mature sepals of *drmy1 wol* and *drmy1 arr1,10,12*, enabling sepal closure (Figure S6).
643 Similar effects in restoring robustness are seen when an extra functional transgene of *ARR7*
644 (*pARR7::ARR7-llama*) is introduced to the *drmy1* mutant (Figure 7J-Q, S7K-N). Our results
645 suggest that cytokinin upregulation is necessary and sufficient for variability in auxin patterning
646 and sepal initiation, indicating that the cytokinin signaling changes are primary defects in *drmy1*,
647 and the auxin signaling changes are secondary. Our results suggest a mechanism different from
648 that previously reported in *ul4y*, *ul18z*, and *el24y*, where ribosomal mutations affect auxin
649 signaling through reduced translation reinitiation of uORF-containing mRNAs, including those of
650 AUXIN RESPONSE FACTOR (ARF) 3, 5, and 7[64–66]. While we found that uORF-containing
651 mRNAs generally have reduced protein-transcript ratio in *drmy1* suggestive of reduced
652 translation, we did not see a consensus reduction in the level of uORF-containing promoter
653 reporters of ARFs (Figure S5). This suggests that the variable auxin signaling pattern in *drmy1* is
654 unlikely to result from changes in uORF-mediated translational regulation of ARFs. Overall, our
655 results suggest that homeostasis in cytokinin signaling is crucial for maintaining robust patterns
656 of auxin signaling and robust morphogenesis in the floral meristem.

658   Mutations affecting ribosome abundance or translation have long attracted interest due to
659 the surprisingly tissue-specific phenotypes they cause[121]. In humans, these mutations have been
660 associated with diseases collectively known as ribosomopathies, where patients show various
661 abnormalities in blood, skeleton, hair, teeth, and pancreas, as well as intellectual disability and
662 increased risk of cancer[122–127]. Ribosomal protein mutants have been characterized in numerous
663 other species with similarly diverse impacts. They display a range of specific phenotypic changes,
664 such as altered pigmentation and skeletal structure in mouse[128–130] and zebrafish[131], shorter
665 bristles and notched wing margins in fruit fly[23,132], abnormal gonad development in worm[133], and
666 pointed leaves and abnormal vascular patterning in Arabidopsis[64,66,134–136]. Here, we show that
667 the Arabidopsis mutant *drmy1* has reduced TOR activity, ribosomal content, and translation rate,
668 causing variable sepal initiation which phenocopies the ribosomal mutants *ul4y* and *ul18z* and the
669 TORC1 assembly mutant *spaghetti-1* (Figure 2, 3). We therefore propose that *drmy1* is an
670 Arabidopsis ribosomopathy mutant like those previously characterized[135].

672   Several mechanisms have been proposed to explain why ribosomopathies do not usually
673 cause a general reduction in growth, but rather affect development in tissue-specific ways. These
674 include extra-ribosomal functions of certain ribosomal proteins[137–141], altered translation behavior

of ribosomal variants on certain mRNAs[142], different competitiveness of mRNAs for scarce
ribosomes[64–66,143–146], and high translation rate requirement for certain proteins[147,148]. For example,
neurotransmitter release in animals relies on constant synthesis of the synaptic vesicle protein
Syt1[149]. A Drosophila *Minute* mutant, *uS15/+,* shows reduced synthesis of Syt1, which in turn
reduces ecdysone secretion in 5-HT neurons, causing delayed larval-to-pupal transition[147].
Similarly, the human apoptosis inhibitor Mcl-1 has a half-life of ~30 min and thus requires a high
translation rate to maintain its proper level. Under translation inhibition, the synthesis of Mcl-1 is
unable to keep up with its degradation, causing apoptosis[148]. Here, we show that the levels of
ARR7 and AHP6, which are rapidly induced by cytokinin and auxin signaling, respectively[150,151],
are drastically reduced under translation inhibition, which underlies the upregulation of cytokinin
signaling and loss of robustness in auxin signaling and morphogenesis (Figure 7R). This
mechanism parallels those previously found in animal systems[147,148], highlighting how
downregulation of proteins with high translational requirements can underlie the tissue-specificity
of ribosomopathy. Outside the floral meristem, the *drmy1* mutant shows other phenotypic changes
such as enlarged shoot apical meristem, reduced apical dominance, phyllotaxy defects, and
reduced root system, all of which are related to altered cytokinin/auxin signaling activity[27]. Thus,
our work highlights how defects in translation, which occurs in every cell, can have tissue-specific
effects on how cells robustly arrange into organs.
693

In addition, we note that not all proteins are equally reduced under broad translation
inhibition. Our data suggest that the cytokinin signaling inhibitor proteins ARR7 and AHP6 are
drastically reduced, compared to others such as AHP3, RCI2A, and GFP-nes. This suggests that
the observed increase in cytokinin signaling under translation inhibition may be due to an
imbalance in the relative levels of activators and inhibitors, which may further suggest that the
inhibitor proteins are more temporally dynamic and thus have higher translational requirements
during development. These hypotheses remain to be tested in future studies.
701

ACKNOWLEDGEMENTS
We thank Bella Burda, Frances Clark, Byron Rusnak, Erich Schwarz, Avilash Yadav, and Maura
Zimmermann for comments and suggestions on the manuscript. We thank Frank Wellmer for the
ap1 cal 35S::AP1-GR (Ler) seeds, Elliot Meyerowitz and Arnavaz Garda for the *DR5::3xVENUS-*
N7/*PIN1::GFP* (Ler) seeds, Teva Vernoux and Géraldine Brunoud for the *TCS::GFP*, *TCS-DR5*
(Col), *ahp6*, *pAHP6::AHP6-VENUS*, and *pAHP6::GFP-ER* seeds, Jan Smalle for the *arr1-1*
35S::ARR1 seeds, Joseph Kieber and Jamie Winshell for the *arr3,4,5,6,7,8,9,15* and
pARR4::ARR4-GFP seeds, Thomas Greb and Min-Hao Chiang for the *pARF5::ER-ARF5-HDEL*
seeds, and Jianru Zuo for the *pAHP3::AHP3-GFP* plasmid. We thank Georg Jander for advice
and protocol for cytokinin extraction. We thank Brian Curtis and Frank Schroeder for help on mass
spectrometry, and Sheng Zhang and Qin Fu for help on Xcalibur. We thank Richie Ragas, Yanã
Rizzieri, and Ziqing Wei for assistance on experiments and data analysis. We thank Vicky
Spencer and Minsung Kim for sharing the *in vivo* Torin2 treatment protocol. We thank Arabidopsis
Biological Resource Center for providing seed stocks and plasmids used in this research.

Research reported in this publication was supported by the National Institute of General Medical
Sciences of the National Institutes of Health (NIH) under award numbers R01GM134037,
DP5OD023072, and R01GM145814; Cornell Graduate School new student fellowship (S.K.); and
in part by a Schmittau-Novak Grant from the School of Integrative Plant Science, Cornell
University (M.Z.). H.G.G. was supported by NIH Director's New Innovator Award (DP2
OD024541-01) and NSF CAREER Award (1652236), NIH R01 Award (R01GM139913), and the
Koret-UC Berkeley-Tel Aviv University Initiative in Computational Biology and Bioinformatics.
H.G.G. is also a Chan Zuckerberg Biohub Investigator. We thank the Metabolomics Workbench[152]
for providing the data repository for our cytokinin mass spectrometry data, which is funded by NIH
under award numbers U2C-DK119886 and OT2-OD030544. We thank the Biotechnology
Resource Center (BRC) Genomics Facility (RRID:SCR_021727) of Cornell University for
performing RNA-seq on *ap1 cal AP1-GR* inflorescence samples. We thank Novogene for
performing seedling RNA-Seq library synthesis and sequencing. We thank the BRC Proteomics
Facility (RRID:SCR_021743) of Cornell University for performing mass spectrometry on *ap1 cal*
AP1-GR inflorescence samples, and NIH SIG Grant 1S10 OD017992-01 for supporting the
Orbitrap Fusion mass spectrometer. The content is solely the responsibility of the authors and
does not necessarily represent the official views of the National Institutes of Health.
734
735

AUTHOR CONTRIBUTIONS

S.K.: Conceptualization, Investigation, Formal Analysis, Visualization, Writing - Original Draft,
Writing - Review & Editing. M.Z.: Conceptualization, Investigation, Formal Analysis, Writing -
Review & Editing. M.R.S: Investigation, Visualization. D.P.: Investigation, Formal Analysis,
Visualization. L.J.: Investigation. R.E.M.: Investigation, Formal Analysis. S.A.: Methodology,
Resources. B.V.L.V.: Methodology, Resources, Writing - Review & Editing. H.G.: Methodology,
Resources, Writing - Review & Editing, Supervision. S.B.Q: Writing - Review & Editing,
Supervision. J.O.B.: Resources, Writing - Review & Editing, Supervision. A.H.K.R:
Conceptualization, Writing - Review & Editing, Supervision, Project administration, Funding
acquisition.
746
747

DECLARATION OF INTERESTS

749 The authors declare no competing interests.

750 **FIGURE TITLES AND LEGENDS**
751

Figure 1. *drmy1* has reduced ribosome abundance, translation rate, and TOR activity.
(A) Top row, stage 12 buds of WT (left) and *drmy1* (right) viewed from the side. Asterisk shows
the gap between sepals with petals and carpels exposed. Middle row, stage 12 buds of WT (left)
and *drmy1* (right) viewed from the top. Arrowheads point to sepals. Note that the *drmy1* bud has
5 sepals of unequal size and unevenly spaced, exposing the stamens and carpels. Bottom row,
stage 5 buds of WT (left) and *drmy1* (right) containing *35S::mCitrine-RCI2A* (plasma membrane
marker). Arrowheads point to sepal primordia. Note that the *drmy1* bud has 5 sepal primordia of
different sizes. Scale bars are 0.5 mm for stage 12 bud images and 25 µm for stage 5 bud images.
(B) Gene ontology (GO) enrichment of downregulated genes in *drmy1* compared to WT, in the
ap1 cal AP1-GR background. Shown are the top 8 GO terms and their enrichment p-values. A
complete list can be found in Supplemental Dataset 1. Arrowheads highlight terms related to
ribosome biogenesis or translation.
(C) Polysomal profiles of WT (blue) and *drmy1* (red) in the *ap1 cal AP1-GR* background,
representative of 3 biological replicates each. Additional replicates are in Supplemental Dataset
2. M, monosomes. P, polysomes.
(D) Puromycin labeling of WT vs *drmy1*. Left, WT and *drmy1* seedlings. From left to right: WT pre-
treated with CHX, two biological replicates of WT pre-treated with mock, and two biological
replicates of *drmy1* pre-treated with mock. All groups were then treated with puromycin. For
seedlings to match in size, WT seedlings were 8 days old and *drmy1* seedlings were 10 days old.
Right, WT and *drmy1* inflorescences of induced *ap1 cal AP1-GR* background. From left to right:
WT co-treated with puromycin and CHX, three biological replicates of WT treated with puromycin,
and three biological replicates of *drmy1* treated with puromycin. In both experiments, RuBisCO
large subunit on Ponceau S-stained membrane is shown as a loading control (bottom). Ratio
between puromycin and Ponceau S signals, normalized by the mean of WT, is shown on the
bottom (p-values are from two-sided Student's t-test).
(E) Gene ontology (GO) enrichment of differentially accumulated proteins in *drmy1* compared to
WT, in the *ap1 cal AP1-GR* background. Shown are the top 8 GO terms and their enrichment p-
values. A complete list can be found in Supplemental Dataset 1. Arrowheads highlight terms
related to ribosome biogenesis or translation.
(F-G) Coherent alteration of gene expression by *drmy1* and AZD-8055 TOR inhibitor treatment.
(F) Scatterplot of RNA log2 fold change in *drmy1* vs WT (x-axis), and WT+AZD vs WT+Mock (y-
axis), in 7-day-old seedlings. Genes are color-coded based on the following categories: genes in
"Structural constituents of the ribosome" (GO:0003735) and its offspring terms (magenta); all
other genes in "Translation" (GO:0006412) and its offspring terms (orange); all other genes (gray).
Blue line shows a linear regression of all points (R^2 = 0.1446, p < 2.2×10^{-16}). **(G)** Of the 466 genes
that are differentially expressed under both conditions, 439 (94%) are coherently altered by AZD-
8055 treatment and the *drmy1* mutation.
(H-I) Phosphorylation of the direct TOR substrate, S6K-pT449, in WT and *drmy1*. **(H)** A
representative blot. Top, S6K-pT449. Middle, total S6K protein. Bottom, Ponceau S staining. Ratio
between S6K-pT449 signal and Ponceau S signal is shown above the blots. **(I)** Ratio between

S6K-pT449 and Ponceau S signals normalized by WT, quantified across in three experiments,

shows that TOR activity decreased by half in *drmy1*. (mean ± SD; *, p<0.05).

Figure 2. Defects in TOR activity, ribosome, and translation disrupt robust sepal initiation.
(A-G) Representative images of stage 5 buds in WT (A), *drmy1* (B), *ul4z* (C), *ul4y* (D), *ul18z* (E),
lst8-1-1 (F), and *spaghetti-1* (G). Tissue morphology is visualized by either propidium iodide (a
cell wall-staining dye) or a plasma membrane marker. Arrowheads indicate sepal primordia that
are variable in number, position, and size. Note that *ul4z* flowers always develop four sepal
primordia, although of different sizes; *lst8-1-1* occasionally (4/41, 9.8%) develops buds with more
than four sepal primordia.
(H) Quantification of sepal primordium number, comparing *drmy1* (n = 67 buds), *ul4z* (n = 52
buds), *ul4y* (n = 53 buds), *ul18z* (n = 52 buds), *lst8-1-1* (n = 41 buds), and *spaghetti-1* (n = 84
buds) with WT (n = 51 buds). Asterisks indicate statistically significant ($p < 0.05$) differences from
WT in Fisher's contingency table tests.
(I) Illustration of robust versus variable positioning of sepal primordia. Primordia are considered
robustly positioned if they are evenly distributed around the edge of the bud. Within each bud,
angles between adjacent primordia with respect to the center of the bud are measured, and
coefficient of variation (CV) is calculated. A bud with robustly positioned primordia would have
similar angular values and a low CV value. A bud with variably positioned primordia would have
very different angular values and a high CV value.
(J) Quantification of variability in primordium positioning (CV) in the same buds as in (H), following
illustration in (I). Asterisks indicate statistically significant ($p < 0.05$) differences from WT in
Wilcoxon's rank sum tests.
(K) Representative images of buds from *in vitro*-cultured WT inflorescences treated with mock or
2 µM CHX for 9-10 days. Arrowheads indicate sepal primordia that are variable in number,
position, and size.
(L) Representative images of buds from WT plants treated with mock or 2 nmol Torin2 for 15
days. Arrowheads indicate sepal primordia that are variable in number, position, and size.
(M-N) Quantification of sepal primordium number (M) and positional variability (N) similar to (H,J),
comparing CHX-treated (n = 31 buds), CHX-mock (n = 42 buds), Torin2-treated (n = 51 buds)
and Torin2-mock buds (n = 56 buds). Scale bars in all micrographs, 25 µm.

Figure 3. TOR and ribosomal defects cause variability in the timing of sepal initiation.

(A-C) 6h-interval live imaging of the sepal initiation process in WT (A) and *ul4y* (B), which is
quantified in (C). n = 48 buds for WT; n = 40 buds for *ul4y*.

(D-F) 6h-interval live imaging of the sepal initiation process in buds from WT plants treated with
mock or 2 nmol Torin2 twice a day for 15 days, which is quantified in (F). n = 31 buds for mock; n
= 15 buds for Torin2.

In **(A,B,D,E)**, top rows show the *35S::mCitrine-RCI2A* membrane marker, and bottom rows show
Gaussian curvature heatmaps of the same image stacks. Asterisks indicate sepal initiation
events, defined as a dark red band (primordium with positive curvature) separated from the floral
meristem by a dark blue band (boundary with negative curvature) in the heatmap. Scale bars, 25
μm.

In **(C,F)**, the amount of time between outer and inner sepal initiation (left) and between outer and
lateral sepal initiation (right) were calculated for each bud. Bar plot shows mean ± SD which is
also shown on top of each plot. Asterisks indicate statistically significant differences ($p < 0.05$) in
Wilcoxon's rank sum test (for mean) or Levene's test (for SD).

Figure 4. Inhibition of TOR activity and translation cause variability in auxin and cytokinin signaling.

(A-D) The ribosomal mutant *ul4y* loses robustness in auxin and cytokinin signaling. (A) Representative images of late stage 2 buds of WT, *drmy1*, and *ul4y*, showing the auxin signaling reporter *DR5::3xVENUS-N7* in yellow, the cytokinin signaling reporter *TCS::GFP* in cyan, and both merged with Chlorophyll (in WT) or *UBQ10::mCherry-RCI2A* (in *drmy1* and *ul4y*) in magenta. (B) Quantification of TCS intensity (integrated density divided by area) from maximum intensity projection images, normalized to mean of WT. Shown are mean ± SD. Asterisks show statistically significant differences from WT in two-tailed Student's t-tests (*drmy1*, p = 2.1×10^{-6}; *ul4y*, p = 3.4×10^{-5}). (C) Circular histogram of DR5 signal distribution. Each bud was divided into 360 sectors of 1° each. Within each sector, DR5 signal measured in pixel intensity units (0-255 range) was summed. This sum was plotted along the x-axis starting from the sector at 1:30 position (between the incipient outer sepal and incipient right sepal) going counterclockwise. I.e., in WT, the outer sepal is near 45°, the inner sepal near 225°, and the lateral sepals near 45° and 135° (vertical dotted lines). The mean was plotted as a solid line, and mean ± SD was plotted as a shaded area. (D) Circular histogram of TCS signal distribution. Sample size for (A-D): WT, n = 12 buds; *drmy1*, n = 15 buds; *ul4y*, n = 10 buds.

(E-H) 3 days of translation inhibition causes increased and diffuse cytokinin signaling, and diffuse auxin signaling. (E) Representative images of late stage 2 WT buds treated *in vitro* with mock or 2 µM CHX for 3 days. Shown are *DR5::3xVENUS-N7* in yellow, *TCS::GFP* in cyan, and both merged with Chlorophyll in magenta. (F) Quantification of TCS intensity from maximum intensity projection images, normalized to mean of WT mock day 3. Shown are mean ± SD. Asterisk shows statistically significant difference in a two-tailed Student's t-test (p = 2.0×10^{-4}). (G) Circular histogram of DR5 signal distribution. (H) Circular histogram of TCS signal distribution. Sample size for (E-H): WT mock day 3, n = 10 buds; WT CHX day 3, n = 12 buds.

(I-L) 6 days of TOR or translation inhibition causes increased and diffuse cytokinin signaling, and randomly positioned auxin signaling maxima. (I) Representative images of late stage 2 WT buds treated *in vitro* with mock, 2 µM CHX, or 2 µM AZD for 6 days. Shown are *DR5::3xVENUS-N7* in yellow, *TCS::GFP* in cyan, and both merged with Chlorophyll in magenta. Arrowheads point to randomly positioned auxin maxima. (J) Quantification of TCS intensity from maximum intensity projection images, normalized to mean of WT mock day 6. Shown are mean ± SD. Asterisks show statistically significant differences from mock in two-tailed Student's t-tests (CHX, p = 1.0×10^{-3}; AZD, p = 1.2×10^{-4}). (K) Circular histogram of DR5 signal distribution. (L) Circular histogram of TCS signal distribution. Sample size for (I-L): WT mock day 6, n = 12 buds; WT CHX day 6, n = 11 buds; WT AZD day 6, n = 10 buds. Scale bars in all micrographs, 25 µm.

Figure 5. Cytokinin signaling is required for increased variability in auxin signaling and
sepal initiation under translation inhibition.
(A-D) Cytokinin treatment makes auxin signaling diffuse. Shown are late stage 2 WT buds under
mock (A,B) or 5 µM cytokinin (BAP) treatment (C,D) for 4 days. (A,C) Auxin signaling reporter
DR5 in yellow, and *DR5* merged with Chlorophyll in magenta. (B,D) Circular histograms of the
DR5 signal, showing mean (solid line) and mean ± SD (shaded area). Arrows point to DR5 signal
in variable positions. Sample size: WT Mock n = 10, WT BAP n = 10. Also see Zhu et al. (2020),
in this reference see Extended Data Figure 7e.
(E-G) Cytokinin signaling is required for variable sepal initiation in *drmy1*. (E) Stage 5 buds. Sepal
primordia in *drmy1* are variable (arrowheads), which does not occur in *drmy1 arr1,10,12* and
drmy1 wol mutants. (F,G) Quantification of sepal primordium number (F) and positional variability
(G), comparing WT (n = 58) with *drmy1* (n = 31), *arr1,10,12* (n = 24) with *drmy1 arr1,10,12* (n =
20), and *wol* (n = 36) with *drmy1 wol* (n = 39). Asterisks indicate statistically significant (p < 0.05)
differences in Fisher's contingency table tests (F) and Wilcoxon's rank sum tests (G) respectively.
(H-K) Cytokinin signaling is required for variable patterning of auxin signaling in *drmy1*. Shown
are late stage 2 buds of WT vs *drmy1* (H,I), and *arr1,10,12* vs *drmy1 arr1,10,12* (J,K). (H,J) Auxin
signaling reporter *DR5* in yellow, and *DR5* merged with propidium iodide in magenta. Arrows point
to diffuse DR5 signal in variable positions of the *drmy1* bud. Arrowheads show four robust DR5
maxima in the *drmy1 arr1,10,12* bud. (I,K) Circular histograms of the *DR5* signal, showing mean
(solid line) and mean ± SD (shaded area). For ease of visualization, circular histograms of *drmy1*
and *drmy1 arr1,10,12* between 90 and 360 degrees are enlarged and shown as insets (y-axis
range 0-0.4). Note the presence of DR5 signal in inter-sepal regions in *drmy1* (black arrow) which
is largely suppressed in *drmy1 arr1,10,12*. Sample size: WT n = 19, *drmy1* n = 16, *arr1,10,12* n =
13, *drmy1 arr1,10,12* n = 9.
(L-N) Cytokinin signaling is required for variable sepal initiation under translation inhibition. (L)
Stage 6 buds of WT, *arr1,10,12*, and *wol*, treated with Mock or 2 µM CHX for 10 days. WT initiates
sepal primordia at variable positions when treated with CHX (arrowheads), which does not occur
in *arr1,10,12* and *wol*. (M,N) Quantification of sepal primordium number (M) and positional
variability (N), comparing mock and CHX within each genotype. Sample size: WT Mock n = 29,
WT CHX n = 19, *arr1,10,12* Mock n = 18, *arr1,10,12* CHX n = 19, *wol* Mock n = 15, *wol* CHX n =
19. Asterisks indicate statistically significant (p < 0.05) differences in Fisher's contingency table
tests (M) and Wilcoxon's rank sum tests (N) respectively.
(O-R) Cytokinin signaling is required for diffuse auxin signaling under translation inhibition. Shown
are late stage 2 buds of WT (O,P) and *arr1,10,12* (Q,R), treated with Mock or 2 µM CHX for 3
days. (O,Q) Auxin signaling reporter *DR5* in yellow, and *DR5* merged with Chlorophyll in magenta.
Arrows point to diffuse DR5 signal in variable positions in CHX-treated WT. Arrowheads show
four robust DR5 maxima in CHX-treated *arr1,10,12*. (P,R) Circular histograms of the *DR5* signal,
showing mean (solid line) and mean ± SD (shaded area). Sample size: WT Mock n = 17, WT CHX
n = 18, *arr1,10,12* Mock n = 7, *arr1,10,12* CHX n = 7. Scale bars in all micrographs, 25 µm.
913

Figure 6. Upregulation of cytokinin signaling is required to maintain translation and fitness
in *drmy1*.
(A) Puromycin labeling of WT seedlings with 4 h CHX pre-treatment (control), and three biological
replicates each of WT and *arr1 35S::ARR1* seedlings with 4 h mock pre-treatment. All seedlings
are 14 days old. RuBisCO large subunit in Ponceau S-stained membrane is shown as a loading
control. Signal ratio between puromycin and Ponceau S, normalized to mean of WT, is show on
the bottom. P-value is from a two-sided Student's t-test. Also see Karunadasa et al. (2020).
(B,C) Puromycin labeling of WT seedlings with 4 h CHX pre-treatment (control), and two biological
replicates of WT, *drmy1*, *wol*, and *drmy1 wol* seedlings with 4 h mock pre-treatment. Seedlings
are 8 days old in (B) and 14 days old in (C). RuBisCO large subunit in Ponceau S-stained
membrane is shown as a loading control. Letters show compact letter display of a Tukey's all-pair
comparison in a one-way ANOVA model.
(D) Representative 14 days old seedling images of WT, *drmy1*, *wol*, and *drmy1 wol* used in (C).
Notice that *drmy1 wol* is very small and pale. Scale bars, 5 mm.
(E) Representative aerial part images of 42 days old plants of WT, *drmy1*, *wol*, and *drmy1 wol*.
Inset shows the zoomed-in *drmy1 wol* plant, which has a tiny rosette and a short inflorescence.
Scale bars, 5 cm. See also Figure S6F.
(F) Representative aerial part images of 74 days old plants of WT, *drmy1*, *arr1*,*10*,*12*, and *drmy1*
arr1,*10*,*12*. Inset shows the zoomed-in *drmy1 arr1*,*10*,*12* plant, which has pale leaves
accumulating anthocyanin and a short inflorescence. Scale bars, 5 cm. See also Figure S6E.
(G) Dissected siliques of *arr1,10,12* (left) and *drmy1 arr1,10,12* (right) showing developing
seeds. Notice that while *arr1,10,12* occasionally have aborted seeds, all seeds in the *drmy1*
arr1,10,12 silique were aborted. Scale bars, 0.2 mm.

Figure 7. ARR7 and AHP6 protein levels are reduced upon inhibition of TOR and
translation.
(A) The hypothesis. During hormone patterning prior to sepal initiation in the WT floral meristem,
A-type ARR and AHP6 proteins are rapidly produced to dampen cytokinin signaling to a normal
level. In *drmy1*, reduced protein synthesis causes reduced levels of these cytokinin signaling
inhibitor proteins, resulting in an upregulation of cytokinin signaling.
(B) Illustration of Llama Tag. Plants were co-transformed with *ARR7-llama* (*pARR7::ARR7-linker-*
llama-ARR7ter) and *GFP-nes* (*pUBQ10::sfGFP-nes-UBQ3ter*). Without ARR7-llama, GFP
localizes to the cytosol due to the nuclear export sequence (nes). ARR7-llama is produced in the
cytoplasm, C, and translocates into the nucleus, N. The Llama Tag on ARR7-llama binds to GFP
and drags GFP into the nucleus (note that from our observation it is excluded from the nucleolus,
NL). Thus, at low ARR7-llama levels, GFP signal is mainly in the cytoplasm. At intermediate
ARR7-llama levels, GFP is at comparable levels between the cytoplasm and the nucleus, and no
clear pattern can be seen. At high ARR7-llama levels, GFP is mainly seen in the nucleus.
(C) A *GFP-nes* bud showing localization of the GFP signal to the cytoplasm.
(D,E) GFP images of buds from two independent transgenic lines of *ARR7-llama GFP-nes*, 7-4
(D) and 7-6 (E), of WT (top) vs *drmy1* (bottom) genotypes. Images are representative of n = 17
(line 7-4, WT), n = 40 (line 7-4, *drmy1*), n = 9 (line 7-6, WT), and n = 6 (line 7-6, *drmy1*) buds.
Note that GFP is more cytoplasm-localized in *drmy1* than WT, indicating reduced ARR7-llama
protein level.
(F) GFP images of WT *ARR7-llama GFP-nes* buds treated with mock (top) or 2 µM CHX (bottom)
for 24 hours. The mock image is representative of n = 20 buds from three independent lines. The
CHX image is representative of n = 19 buds from these same lines.
(G) GFP images of WT *ARR7-llama GFP-nes* buds treated with mock (top) or 2 µM AZD-8055
(bottom) for 72 hours. The mock image is representative of n = 13 buds from two independent
lines. The AZD-8055 image is representative of n = 11 buds from these same lines. For (C-G),
each image was brightened to reveal subcellular localization patterns of GFP. A square region
taken from the image containing 5-10 cells is enlarged and shown on the top right. Within the
square, GFP intensity was quantified along the dotted line and plotted on the bottom right. X-axis,
pixels (range 0-238). Y-axis, GFP intensity (smoothened by taking the average intensity of 11-
pixel neighborhoods; range 90-210 in gray value).
(H-I) Response of the AHP6 protein reporter (H) and transcriptional reporter (I) to mock, CHX,
and AZD-8055 treatments for 72 hours. For (H), images are representative of n = 29 (mock), n =
29 (CHX), and n = 34 (AZD-8055) buds in three experiments. For (I), images are representative
of n = 11 (mock), n = 9 (CHX), and n = 12 (AZD-8055) buds in two experiments.
(J-Q) Reduction of A-type ARR and AHP6 protein levels contribute to the variability in sepal
initiation. (J-O) Stage 5-6 buds of indicated genotype stained with propidium iodide. Arrowheads
indicate sepal primordia that are variable in number, position, and/or size. Note that the
arr3,4,5,6,7,8,9,15 bud has an inner sepal that is slightly smaller than its outer sepal and
positioned slightly right-skewed (K). The *ahp6* bud develops five sepal primordia of variable sizes
and unevenly positioned (L). The *ARR7-llama GFP-nes* constructs partially rescue the *drmy1*
phenotype in some buds (O, left) but not others (O, right). (P) Quantification of sepal primordium

number. Asterisk indicates statistically significance difference in a Fisher's exact test (WT vs
ahp6, p = 3.026×10^{-7}; *drmy1* vs *drmy1 ARR7-llama GFP-nes*, p = 0.4389). (Q) Quantification of
variability in sepal primordium position. Asterisk indicates statistically significant difference in a
Wilcoxon rank sum test (WT vs. arr3,4,5,6,7,8,9,15, p = 2.948×10^{-4}; WT vs *ahp6*, p = 2.137×10^{-}
^{11}; WT vs *ARR7-llama GFPnes*, p = 1; *drmy1* vs *drmy1 ARR7-llama GFPnes*, p = 1.538×10^{-7}).
Data for *drmy1* were reused from Figure 2H, 2J. Data for *ARR7-llama GFP-nes* and *drmy1 ARR7-*
llama GFP-nes were pooled from two independent lines (7-4 and 7-6). Sample size: WT, n = 78;
arr3,4,5,6,7,8,9,15, n = 28; *ahp6*, n = 106; *ARR7-llama GFPnes*, n = 16; *drmy1*, n = 67; *drmy1*
ARR7-llama GFP-nes, n = 20.
(R) Working model. In WT, DRMY1 maintains TOR activity and translation, which sustains the
rapid production of cytokinin signaling inhibitors (ARR7 and AHP6) in response to cytokinin
signaling. These inhibitors maintain cytokinin signaling at a normal level, allowing auxin and
cytokinin signaling to interact and form robust spatial patterns. Robust patterning of auxin and
cytokinin signaling gives rise to robustly initiated sepal primordia. In *drmy1*, due to decreased
TOR signaling and translation rate, the meristem cannot rapidly produce cytokinin signaling
inhibitor proteins in response to cytokinin signaling. As a result, cytokinin signaling is upregulated,
which rescues the translation rate reduction in a homeostatic mechanism. This upregulation of
cytokinin signaling disrupts the robust spatial pattern of both cytokinin and auxin signaling, which
in turn causes variability in sepal initiation. Scale bars in all micrographs, 25 μm.
998

999 **MATERIALS AND METHODS**

1000

Plant material

Most Arabidopsis plants were in Col-0 background (WT). *ap1 cal 35S::AP1-GR* was in Ler
background. *drmy1* (Col-0) was backcrossed to Ler twice and then crossed with *ap1 cal*
35S::AP1-GR to obtain *drmy1 ap1 cal 35S::AP1-GR*. R2D2 was originally in Col-Utrecht
background and was backcrossed twice into WT (Col-0) and *drmy1* (Col-0). The following mutants
and reporters were previously described: *drmy1-2*[27], *wol-1*[153], *spaghetti-1* (*tpr5-1*)[154], *ahp6*[16],
arr3,4,5,6,7,8,9,15[101], *ap1 cal 35S::AP1-GR* (Ler)[35,36], *arr1-1 35S::ARR1*[74], *DR5::3xVENUS-*
N7[155], *TCS::GFP*[156], *pARF5::ER-EYFP-HDEL*[157], *pUS7Y-mDII-NtdTomato-pUS7Y-DII-*
N3xVENUS (R2D2)[109], *35S::mCirtine-RCI2A*[27], *UBQ10::mCherry-RCI2A*[27], *pAHP3::AHP3-*
GFP[100], *pAHP6::AHP6-VENUS*[16], and *pAHP6::GFP-ER*[16,97]. The following mutants and reporter
lines were obtained from Arabidopsis Biological Resource Center (ABRC): *ul4z* (SALK_130595),
ul4y (SALK_029203), *ul18z* (SALK_089798), *arr1-3 arr10-5 arr12-1*[158] (CS39992), *lst8-1-1*
(SALK_002459), *pARF3::N3xGFP*[159] (CS67072), *pARF6::N3xGFP*[159] (CS67078),
pARF8::N3xGFP[159] (CS67082), *pARF10::N3xGFP*[159] (CS67086).

1015

Llama-tagged ARR7 construct

For the LlamaTag system, we first generated plasmid *pVV13* containing linker-llama. We
amplified the LlamaTag (from a plasmid containing vhhGFP4[160]) and added a linker sequence of
tccggagcagctgcggctgccgctgcggcagcggccactagt at its 5' end by two rounds of overlap PCRs.
Primers for the first round were oVV64 and oVV53, and primers for the second round were oVV35
and oVV53. After the second round, we A-tailed the PCR product according to the Promega
manufacturer's protocol. A-tailed product was ligated to the pGEMTeasy vector according to the
Promega ligation protocol, to create the plasmid *pVV13*.

To make *pARR7::ARR7-llama*, a genomic fragment of *pARR7::ARR7* minus the stop
codon and terminator was amplified from the Arabidopsis (Col-0) genome using the primers
oSK197 and oSK198. The linker-llama fragment was PCR-amplified from *pVV13* using the
primers oSK199 and oSK200. The *ARR7* stop codon, 3' UTR, and terminator was amplified from
the Arabidopsis (Col-0) genome using the primers oSK201 and oSK202. *pMLBART* backbone
was digested with NotI, and all fragments were assembled into *pMLBART* using NEBuilder
according to the manufacturer's protocol.

To make *pUBQ10::sfGFP-NES:UBQ3ter*, sfGFP sequence was amplified from the *35S-*
sfGFP-nosT plasmid[161] (Addgene # 80129) using primers UsfGM-F1 and UsfGnes-R1. The
UBQ10 promoter was amplified from the *UPG* plasmid[162] (Addgene # 161003) using primers
OutALFd and UsfGM-R1. The UBQ3 terminator was amplified from the *UPG* plasmid[162] (Addgene
161003) using primers UsfGnes-F1 and OutALRb. Primer overhangs spanning the junction
between sfGFP and the UBQ3 terminator contain the sequence of the mouse PKIα NES.
pCambia1300 backbone was digested with BamHI and KpnI, and all fragments were Gibson-
assembled into the backbone. Sequences of primers, *pARR7::ARR7-llama*, and *pUBQ10::sfGFP-*
NES:UBQ3ter can be found in Supplemental Dataset 5.

Col-0 plants were co-transformed with *pARR7::ARR7-llama* and *pUBQ10::sfGFP-*
NES:UBQ3ter, and selected with Basta (for *pARR7::ARR7-llama*) + Hygromycin (for
pUBQ10::sfGFP-NES:UBQ3ter). Surviving T1 plants were screened for clear nuclear signal in the
inflorescence, and 5 independent T1 plants were selected and crossed into *drmy1*. F2 plants from
each line were again selected with Basta + Hygromycin and genotyped. One line showed co-
segregation with the *DRMY1* locus. Two lines showed severe silencing in the F2 and could not
be used. Two lines (7-4 and 7-6), though with minor silencing in F2, were used for imaging and
image analysis. F3 plants of 7-4 and 7-6 had severe silencing, and therefore only F2 were imaged.
1048

Plant growth conditions
For most experiments, seeds were sown in wetted Lamber Mix LM-111 soil and stratified
at 4°C for 3-5 days. For experiments including *drmy1 wol* and *drmy1 arr1,10,12*, all seeds were
sown onto ½ MS plates with 0.05% (w/v) MES, 1% (w/v) sucrose, 1.2% (w/v) agar, pH 5.7, and
stratified at 4°C for a week. They were grown for 7-10 days before being transplanted to soil (for
imaging of inflorescence or aerial part of the plant) or left on the plates until desired time of the
experiment (for seedling imaging or puromycin labeling).
Most plants were grown under 16 h – 8 h light-dark cycles (fluorescent light, ~100 µmol
m^{-1} s^{-1}) at 22°C in a Percival walk-in growth chamber. We found that the *drmy1* phenotype is more
pronounced in this condition than under continuous light. The *ap1 cal 35S::AP1-GR* and *drmy1*
ap1 cal 35S::AP1-GR plants were grown in soil under continuous light at 16°C to prevent
premature floral induction.
1061

Flower staging
Flower buds were staged as previously described[37]. Briefly, stage 1 is when the floral
meristem emerges, but not yet separated, from the inflorescence meristem. Stage 2 is when the
floral meristem separates from the inflorescence meristem but with no floral organs initiated.
Stage 3 is when sepal primordia initiate. Stage 4 is when sepal primordia bend to cover part of
the floral meristem. Stage 5 is when stamen primordia initiate. Stage 6 is when sepal primordia
completely cover the floral meristem.
1069

RNA-seq data collection and analysis
For RNA-seq in the inflorescence, bolting *ap1 cal 35S::AP1-GR* and *drmy1 ap1 cal*
35S::AP1-GR plants were induced daily with an aqueous solution containing 10 µM
dexamethasone (Sigma-Aldrich), 0.01% (v/v) ethanol, and 0.015% (v/v) Silwet L-77
(Rosecare.com). When sepals initiated from the floral meristems, usually on the fourth day after
three daily inductions, three inflorescence samples per genotype (including inflorescence
meristems and buds under stage 6) were collected and immediately put into liquid nitrogen. RNA
extraction, library preparation, RNA-seq, and data analysis for inflorescence samples were done
as previously described[27] with a few changes. After read mapping, genes with at least two raw
reads in at least two biological replicates in either WT or *drmy1* were kept for downstream
analysis. For differentially expressed genes, we set a log2 fold change threshold of ±1 and a BH-
adjusted p-value threshold of 0.05. For GO term enrichment, gene-GO mapping data was

obtained   from   TAIR   (https://www.arabidopsis.org

/download_files/GO_and_PO_Annotations/Gene_Ontology_Annotations/ATH_GO_GOSLIM.txt)

. The R package "topGO"[163] (version 2.38.1) was used for the enrichment, with statistic "fisher",

algorithm "weight01", annotation function "annFUN.gene2GO", and minimum node size 10. The

results were ranked by their p-value, and the first 8 terms were plotted.

For RNA-seq in seedlings, WT and *drmy1* seedlings were grown to quiescence (7 days)

in ½ MS liquid media as previously described[34]. After 7 days, the media was replaced with ½ MS

liquid media containing 15 mM glucose and incubated for 24 hours to activate TOR. Seedlings

were then incubated with or without AZD-8055 in addition to 15 mM glucose in ½ MS liquid media

for 2 hours before collecting tissue. RNA was extracted from 100 mg pooled seedlings using the

Spectrum Plant Total RNA Kit (Sigma). This RNA was used as a template for RNA-Seq library

synthesis and sequencing, which was performed by Novogene. RNA-seq data for AZD-8055

treated WT and *drmy1* seedlings were preprocessed with fastp (v. 0.22.0) using default

parameters. Preprocessed reads were then mapped to the TAIR10 reference genome using

STAR (v. 2.7.10z_alpha_220314). Following alignment, BAM output files from STAR were used

to generate feature counts for transcripts using subread-featureCounts (v. 2.0.3) and the

Araport11 transcriptome. TPMs were generated using TPMCalculator (v. 0.0.3). Differential

expression analysis was performed using feature count data and DESeq2 (v. 1.36.0).

A list of genes with uORFs based on gene models of the TAIR10 Arabidopsis genome

assembly were downloaded from von Arnim et al.[69]. For each gene, within each genotype, protein-

transcript ratio was calculated as the ratio between mean protein abundance and mean transcript

TPM across all bio-reps in our proteomics and RNA-seq datasets, respectively. This was log2-

transformed, and the difference between *drmy1* and WT was calculated. This was used as an

indicator of translation rate difference between *drmy1* and WT, although we acknowledge that

other factors such as protein stability may affect this number. This was plotted against the number

of uORFs in each gene model (0, 1, or ≥ 2).

1108

Proteomics

Five induced inflorescence samples of WT and *drmy1* in *ap1 cal AP1-GR* background

were collected as described above. Samples were ground in liquid nitrogen. Total soluble proteins

were extracted in ice-cold extraction buffer (50 mM PBS-HCl (pH 8.0) buffer with 150 mM NaCl,

2% NP-40, 1 mM PMSF, 1x Roche cOmplete protease inhibitor cocktail (Sigma 11697498001),

and 1x Halt TM Phosphatase inhibitor cocktail (ThermoFisher 78420)) and filtered through

Pierce™ Micro-Spin Columns (30 μm pore size; Thermo Scientific 89879). Extracts were

RuBisCO-depleted using Seppro Bubisco Kit (Sigma SEP070-1KT), concentrated, denatured,

reduced, cysteine blocked, trypsin-digested, and TMT 10-plex labeled. Then, mass spectrometry

was done using an UltiMate 3000 RSLCnano / Orbitrap Fusion system (Thermo Scientific). Raw

data was searched against the NCBI protein database using PD 2.3 (Thermo Scientific) with

Sequest HT searching engine. Precursor-based protein identification and relative quantification

was done using the standard processing workflow in PD 2.3, with an additional node of Minora

Feature Detector. Proteins with at least 2 supporting peptides were kept for downstream analysis.

For each protein, data was fit with an ANOVA model and a p-value was calculated. Proteins with

a p-value < 0.05 were considered differentially accumulated in *drmy1*. GO term enrichment was
done as above, using genes corresponding to the differentially accumulated proteins.
1126
Polysome extraction and profiling
Three induced inflorescence samples of WT and *drmy1* in *ap1 cal AP1-GR* background
were collected as described above, and polysomes were extracted as previously described[164].
Briefly, samples were ground in liquid nitrogen, mixed with an extraction buffer twice the volume
of pulverized tissue (0.2 M Tris pH 9.0, 0.2 M KCl, 0.025 M EGTA, 0.035 M $MgCl_2$, 1% (w/v) Brij-
35, 1% (v/v) Triton X-100, 1% (v/v) Igepal CA-630, 1% (v/v) Tween-20, 1% (w/v) Sodium
deoxycholate, 1% (v/v) Polyoxyethylene 10 tridecyl ether, 5 mM Dithiothreitol, 1 mM
Phenylmethylsulfonyl fluoride, 100 µg/ml cycloheximide, 100 µg/ml chloramphenicol, 40 U/ml
RNasin, 10 U/ml DNase I), and let sit on ice for 10 min. Samples were centrifuged at 4°C 4,000 g
for 5 min, supernatant was transferred to a new tube, centrifuged at 4°C 16,000 g for 15 min, and
supernatant was filtered through Miracloth.
Polysome extracts were profiled as previously described[165]. Briefly, 600 µl of each sample
were loaded onto a 15%-45% sucrose density gradient and centrifuged at 4°C 32,000 rpm in a
SW41 rotor. Separated samples were fractionated at a rate of 0.375 mL/min in an Isco
fractionation system, and absorbance at 254 nm was recorded.
1142
Puromycin labeling
Puromycin labeling was done as previously described[39], with slight modifications.
In seedlings, when comparing WT and *drmy1*, in order to control for plant size, WT
seedlings were grown for 8 days and *drmy1* seedlings were grown for 10 days (Figure 1E). When
comparing WT, *drmy1*, *wol*, and *drmy1 wol*, we were unable to control for plant size because
drmy1 wol seedlings were too small. We therefore controlled for plant age, and seedlings were
grown to specified age (8 days for Figure 6B and 14 days for Figure 6C). Seedlings were
harvested from plates and incubated with an incubation buffer (½ MS, 0.05% (w/v) MES, 1% (w/v)
sucrose, 0.1% (v/v) Tween-20, 0.1% (v/v) DMSO, 1x Gamborg vitamin mix, pH 5.7), with or
without 50 µM CHX, for 4 hours in an illuminated growth chamber. Then, the buffer was replaced
with a fresh incubation buffer (which is same as above, but contains 50 µM puromycin (GoldBio
P-600-100)), and incubation continued for another 45 min.
In inflorescences of WT and *drmy1* in *ap1 cal AP1-GR* background, inflorescences were
DEX-induced as described above. Inflorescence samples were collected and put in an incubation
buffer (½ MS, 1% (w/v) sucrose, 0.02% (v/v) Silwet L-77, 0.1% (v/v) DMSO, 50 µM puromycin,
1x Gamborg vitamin mix, pH 5.7), with or without 100 µM CHX. Samples were vacuum infiltrated
for 15 minutes and then put on a rocking shaker in an illuminated growth chamber for 45 minutes.
In both cases, at the end of the incubation, samples were washed three times with water,
blot dry, weighed, and frozen in liquid nitrogen. Soluble proteins were extracted as described
above. Puromycin incorporated into the proteins were detected in a Western blot using a mouse-
origin anti-puromycin monoclonal antibody (12D10, Sigma MABE343, lot # 3484967) and a goat-
anti-mouse HRP-conjugated secondary antibody (Abcam ab6789, lot # 3436981). RuBisCO large
subunit in Ponceau S-stained membrane was used as a loading control. Quantification was done

in ImageJ. A background signal was determined using blank regions, and subtracted from all
quantified signals (separately for puromycin and Ponceau S).
1168

TOR activity assay

WT and *drmy1* seedlings were grown in a six-well plate containing ½ MS liquid media.
After seven days, the media were replaced with half-strength MS liquid media plus 15 mM glucose
and incubated for 24 hours. At least 120 quiescent seedlings per sample were collected and
frozen in liquid nitrogen. Protein was then extracted from the plant tissue in 100 mM MOPS (pH
7.6), 100 mM NaCl, 5% SDS, 0.5% b-mercaptoethanol, 10% glycerin, 2 mM PMSF, and 1x
PhosSTOP phosphatase inhibitor (Sigma). S6K-pT449 was detected by Western blot using a
phosphospecific antibody (Abcam ab207399) and an HRP-conjugated goat anti-rabbit IgG
secondary antibody (Jackson Immuno Research 111-035-003). Total S6K was detected using a
custom monoclonal antibody described by Busche et al.[166]. Total protein visualized in Ponceau
S-stained membrane was used as a loading control.
1180

Confocal microscopy

Confocal imaging of reporter lines in the inflorescence were done as previously
described[27]. Briefly, main inflorescences (not side branches) were cut and dissected with a
Dumont tweezer (Electron Microscopy Sciences, style 5, no. 72701-D) to remove buds older than
stage 9 or 10. The inflorescences were then inserted upright into a small petri dish (VWR, 60 x
15 mm) containing inflorescence culture medium (1/2 MS, 1% (w/v) sucrose, 1x Gamborg vitamin
mixture, 0.1% (v/v) plant preservative mixture (Plant Cell Technology) 1% (w/v) agarose, pH 5.8),
leaving most of the stem inside the medium and the buds outside. They were then further
dissected to reveal stage 6 and younger buds, immersed with water, and imaged under a
Zeiss710 upright confocal microscope with a 20x Plan-Apochromat water-dipping lens (1.0 NA).
For live imaging experiments, inflorescence samples were put in a continuous-light growth
chamber between time points. To prevent bacterial growth, samples were transferred onto fresh
media every 2 to 3 days, and for live imaging experiments lasting longer than 6 days, once in the
middle, plants were incubated with an aqueous solution of 100 µg/ml Carbenicillin (GoldBio, C-
103-5, lot # 0129.091814A) for 30 minutes.
To visualize tissue morphology of inflorescence samples without a reporter, samples were
stained for 5 minutes with an aqueous solution of 0.1 mg/ml propidium iodide (PI) and 0.1% (v/v)
Tween-20, washed three times with water, and imaged.
The following laser and wavelength were used in confocal imaging. Chlorophyll, excitation
488 nm, emission 647-721 nm. PI, excitation 514 nm, emission 566-659 nm. mCherry, excitation
594 nm, emission 600-659 nm. tdTomato, excitation 561 nm, emission 566-595 nm. For
EYFP/VENUS/mCitrine, in *35S::mCirtine-RCI2A*, excitation 514 nm, emission 519-580 nm; in
DR5::3xVENUS-N7, excitation 514 nm, emission 519-569 nm; in *pARF5::ER-EYFP-HDEL*,
excitation 514 nm, emission 519-550 nm; in *R2D2*, excitation 488 nm, emission 493-551 nm. For
GFP/sfGFP, in *pARR7::ARR7-llama UBQ10::sfGFP-NES*, excitation 488 nm, emission 493-569
nm; in *pARF3::N3xGFP*, *pARF6::*N3xGFP, *pARF8::N3xGFP*, and *pARF10::N3xGFP*, excitation
488 nm, emission 493-564 nm; in *TCS::GFP*, excitation 488 nm, emission 493-513 nm.

1208

Visualization of tissue morphology

For single-channel image stacks intended for the visualization of tissue morphology
(*35S::mCitrine-RCI2A* or PI), stacks were 3D-rendered using the ZEN confocal software
(Processing -> 3D). Parameters were set to best visualize tissue morphology, typically, minimum
5-10, ramp 60-80, maximum 100. Buds were rotated to desired orientation, and screenshots were
taken using the "Create Image" button. For fluorophores that are dimmer, less sharp, or have a
noisy background (*UBQ10::mCherry-RCI2A* or Chlorophyll), stacks were converted from LSM to
TIF using ImageJ[167,168], loaded into MorphoGraphX[169], and screenshots were taken using the
built-in screenshot function in MorphoGraphX.

To aid visualizing tissue morphology and determine the timing of sepal initiation, each
stack was fitted with a surface, and a Gaussian curvature heatmap was calculated from the
surface (see below). We consider a sepal primordium as initiated when we see a dark red band
of positive Gaussian curvature (primordium) separated from the center of the floral meristem by
a dark blue band of negative Gaussian curvature (boundary)[27].

Gaussian curvature heatmaps were calculated as previously described[27], with slight
modifications. Briefly, stacks underwent the following processes in MorphoGraphX: Gaussian blur
(3 times; X/Y/Z sigma = 1 µm for the first 2 times, and 2 µm for the third time), edge detection
(threshold = 2000-8000 depending on the brightness of the stack, multiplier = 2.0, adapt factor =
0.3, fill value = 30000), marching cube surface (cube size = 8 µm, threshold = 20000), subdivide
mesh, smooth mesh (passes = 5), subdivide mesh, smooth mesh (passes = 5), project mesh
curvature (type = Gaussian, neighborhood = 10 µm, autoscale = no, min curv = -0.0015, max curv
= 0.0015). For ease of visualization, the lookup table "jet" was applied to the mesh.

1231

Quantification of sepal initiation robustness

For sepal primordium number, screenshots were taken of stage 3-6 buds of indicated
genotypes, in either ZEN or MorphoGraphX. The number of sepal primordia initiated were
counted from these screenshots.

For variability in sepal primordium positioning, within each bud, an angular distance was
measured between each pair of adjacent sepal primordia (with vertex at the center of the bud),
using ImageJ. Note that the last pair was not measured – the angular distance was calculated as
the sum of all other angular distances subtracted from 360°. A CV value (standard deviation
divided by mean) was calculated from all the measured or calculated angular distances. Buds
with sepal primordia evenly distributed around the bud periphery should have a small CV value,
i.e. all angles are around 90° for four-sepal buds (or 72° for five-sepal buds, etc.). Buds whose
sepal primordia distributed variably or randomly around the bud periphery will have widely varying
angular distances between adjacent sepal primordia, and thus large CV values.

Relative sepal initiation timing was quantified as previously described[27]. Briefly, dissected
inflorescence samples were live-imaged every 6 hours. A Gaussian curvature heatmap was
generated for each sample at each time point and was used to determine the time point at which
a sepal primordium initiates. A sepal primordium is considered initiated at time point Tn if it is
absent at time point T(n-1) but becomes present at time point Tn. Within the same bud, we

counted the number of time points between outer and inner sepal initiation, and between outer
and lateral sepal initiation, and multiplied them by 6 hours to get the relative initiation timing of
these sepals.
1253
Quantification of fluorescent reporters
For *TCS::GFP*, *pARF3::N3xGFP*, *pARF5::ER-YFP-HDEL*, *pARF6::N3xGFP*,
pARF8::N3xGFP, *pARF10::N3xGFP*, *pUS7Y::mDII-NtdTomato, pUS7Y::DII-N3xVENUS*, and
UBQ10::mCherry-RCI2A, total signal (integrated density) was quantified from maximum intensity
projection images using ImageJ[167,168]. Fluorescence intensity was measured in pixel intensity
units (0-255 range). Signal intensity was calculated as total signal divided by area.
For both *TCS::GFP* and *DR5:3xVENUS-N7,* circular histogram analysis was done as
previously described[27]. Briefly, individual buds were cropped out of image stacks, channels were
split using FIJI and saved in TIF format, and TIF stacks were imported into MorphoGraphX. Signal
from outside the buds (e.g. inflorescence meristem, parts of other buds within the same image)
was manually removed using the Voxel Edit function. Buds were positioned so that the incipient
sepal primordia are in the XY plane: the incipient outer sepal is at 45°, the incipient inner sepal
and the inflorescence meristem are at 225°, and the incipient lateral sepals are at 135° and 315°,
respectively. Fluorescence intensity was measured in pixel intensity units (0-255 range). A circular
histogram of bin width 1° centered around the Z axis was exported for each replicate expressing
DR5 and/or TCS. Multiple circular histograms of the same reporter and genotype were pooled
and mean ± SD were plotted.
For GFP signal in plants carrying *pUBQ10::sfGFP-nes-UBQ3ter* and *pARR7::ARR7-*
linker-llama-ARR7ter reporters, screenshots were taken in MorphoGraphX as described above.
Screenshots were subtracted of a background determined using blank regions with no tissue, and
brightened to the same level to reveal differences in GFP distribution patterns. A square region
containing 5-10 cells were taken from each screenshot, and GFP intensity (in gray value ranging
from 0 to 255) along a straight line of 239 pixels in length was quantified using ImageJ[167,168]
(Analyze -> Plot profile). For ease of visualization, the curves were smoothed by taking the
average of the gray value of 11 neighboring pixels (including itself) as the value of each pixel.
For VENUS or GFP signal in *pAHP6::AHP6-VENUS*, *pAHP6::GFP-ER*, *pAHP3::AHP3-*
GFP, and *pUBQ10::mCherry-RCI2A* under mock, CHX, or AZD-8055 treatment, total signal at 24
hours (for CHX) or 72 hours (AZD-8055) was normalized by bud area in the 2D projection to get
the signal intensity. To account for bud-to-bud differences in signal intensity prior to treatment,
the signal intensity was normalized to the 0-hour time point (pre-treatment). Relative level
between treatment and mock was calculated by normalizing this value to the mean of mock.
1285
In vitro **drug treatments on inflorescence samples**
For cycloheximide (CHX) treatment, a stock solution of 10 mM CHX was made from
powder (Sigma C1988) in pH 4.0 water. The stock was filter-sterilized and stored in -20°C, and
added to autoclaved inflorescence culture medium to a final concentration of 2 µM just before
use. For AZD-8055 treatment, a stock solution of 16 mM AZD-8055 was prepared from powder
(Cayman Chemical 16978) in DMSO within days of use, and stored in -80°C. The stock was

added to autoclaved inflorescence culture medium to a final concentration of 2 µM. 0.0125% (v/v)

DMSO was added to the mock medium. For 6-benzylaminopurine (BAP) treatment, a stock

solution of 50 mM BAP was prepared from powder (Alfa Aesar A14678) in DMSO, and stored in

-80°C. The stock was added to autoclaved inflorescence culture medium to a final concentration

of 5 µM. 0.01% (v/v) DMSO was added to the mock medium.

Inflorescences were dissected and inserted into regular inflorescence culture medium

without drugs, and pre-treatment image stacks were captured. Then, they were transferred into

specified treatment or mock media, and imaged at the specified time points. For live imaging,

inflorescence samples were transferred onto new medium after each imaging session.

1301

***In vivo* Torin2 treatment**

Starting at 14 days after germination, twice each day for 15 days, 2 nmol of Torin2

(Cayman Chemical 14185) in 20 µl of aqueous solution containing 0.5% DMSO and 0.5% Tween-

20 was applied to the center of the rosette using a pipette. For mock, 20 µl aqueous solution

containing 0.5% DMSO and 0.5% Tween-20 was applied. At the end of the 15-day treatment

period, inflorescences were dissected and put in the inflorescence culture medium for imaging.

To prevent Torin2 degradation, throughout the duration of this experiment, the Torin2

stock solution in DMSO was kept in -80°C and replaced each week, and the treatment and mock

solutions were kept in 4°C and replaced each day.

1311

Imaging of whole plant, whole inflorescence, silique, and mature sepals

For whole-plant imaging, aerial parts of the plants were removed from the pots, flattened,

put on a dark cloth, and imaged with a cell phone (iPhone 12, iOS 16.2).

For whole-inflorescence imaging, inflorescences consisting of open flowers and unopened

buds were removed from the plant and held with forceps. Images were taken under a Zeiss Stemi

2000-C Stereo Microscope with a cell phone (iPhone 12, iOS 16.2).

For silique imaging, siliques on inflorescences sufficiently distant from the shoot apex that

were developed and started to ripen were picked with forceps, opened with a razor blade, and

imaged under a Zeiss Stemi 2000-C Stereo Microscope with a cell phone (iPhone 12, iOS 16.2).

Mature sepal imaging was done as previously described[26,27]. Briefly, mature sepals from

stage 15 flowers (10th to 25th flower on the inflorescence) were dissected and sandwiched

between two slides to flatten. Images were taken using a Canon Powershot A640 camera

attached to a Zeiss Stemi 2000-C Stereo Microscope. Minor damages were manually fixed, and

undesired objects such as pollen grains were manually removed from these images. Sepals with

major damages were discarded. Then, a contour was extracted from each sepal using custom

python scripts[26]. This gave us measurements such as length, width, area, etc. of each sepal. To

measure between-flower variability of length, within each genotype and for each of outer, inner,

and lateral positions, a CV (standard deviation divided by mean) of all sepals was calculated (for

example, a CV of length of all outer sepals in WT). To determine statistical significance, genotypes

were compared pairwise using permutation tests. To measure within-flower variability of length,

a CV was calculated for all sepals within each flower (for example, a CV of length of outer, inner,

and two lateral sepals in WT bud #10). For accurate calculation of CV, flowers with length data of

at least four sepals were included in the analysis. To determine statistical significance, genotypes
were compared pairwise using Wilcoxon rank sum tests.
1336

Cytokinin extraction and measurement
Cytokinin extraction was based on a previously published protocol[170] with modifications.
Briefly, five inflorescence samples of induced *ap1 cal 35S::AP1-GR*, and six inflorescence
samples of induced *drmy1 ap1 cal 35S::AP1-GR* were collected as described above. Samples
were ground in liquid nitrogen and twice extracted in methanol : water : formic acid (15:4:1). 200
pg of BAP per sample was added as an internal control. Extracts were centrifuged at 14,650 rpm
in -4°C for 30 min, and supernatant was evaporated of methanol and reconstituted in 1% (v/v)
acetic acid. Samples were passed through an Oasis MCX SPE column (Waters 186000252),
washed with 1% acetic acid, washed with methanol, and eluted with 0.35 M ammonia in 70%
methanol. Eluents were evaporated to complete dryness, reconstituted in 5% acetonitrile, and
sent for LC-MS.
LC-MS was done as previously described[171], with modifications. Briefly, 1 µl of each
sample was injected into a Thermo Fisher Scientific Vanquish Horizon UHPLC System coupled
with a Thermo Q Exactive HF hybrid quadropole-orbitrap high-resolution mass spectrometer
equipped with a HESI ion source. Samples were separated on a C18 ODS column (AQUITY
UPLC BEH C18, 1.7 µm, 2.1 × 100 mm, Waters), at a flow rate of 0.3 ml/min, with linear gradients
of solvent A (0.1% formic acid) and solvent B (0.1% formic acid in methanol) according to the
following profile: 0 min, 99.0% A + 1.0% B; 4.0 min, 55.0% A + 45.0% B; 7 min, 30.0% A + 70.0%
B; and then with isocratic conditions: 8 min, 1.0% A + 99.0% B; 12 min, 99.0% A + 1.0% B.
Cytokinins were detected using the positive ion mode.
For tZ, tZR, iP, iPR, and the internal control BAP, peaks were identified from an external
standard mix composed of 0.1 µg/ml each of BAP (Alfa Aesar A14678), tZ (Sigma Z0876), tZR
(Sigma Z3541), iP (Cayman Chemical 17906), and iPR (Cayman chemical 20522) in 5%
acetonitrile. For cZ and cZR, peaks were identified based on previously reported precursor m/z
and retention time[172]. Using Xcalibur (Thermo Scientific), peak area was quantified for each
cytokinin in each sample, normalized against the peak area of BAP (internal control) and sample
fresh weight, and then normalized against the average abundance of tZ in WT samples.
1364

Software
Image processing was done in ImageJ (version 2.9.0/1.53t, build a33148d777)[167,168] and
MorphoGraphX (version 2.0, revision 1-294, CUDA version 11.40)[169].
Data processing was done in RStudio (R version 4.0.5 "Shake and Throw" (2021-03-
31))[173]. Graphs were made using the package ggplot2 (version 3.3.3)[174]. Fisher's contingency
table tests were done using the function fisher.test in R. Wilcoxon rank sum tests were done using
the function wilcox.test in R. Hypergeometric tests were done using the function phyper in R. Data
fitting with ANOVA was done using the function aov in R.
Figures were assembled in Adobe Illustrator (version 25.4.1). An RGB color profile "Image
P3" was used for all the figures.
1375

Accession numbers

RNA-seq data for *ap1 cal AP1-GR* and *drmy1 ap1 cal AP1-GR* inflorescence tissue were
deposited in Gene Expression Omnibus (GEO) under accession number GSE230100. RNA-seq
data for WT and *drmy1* seedlings treated with mock or AZD-8055 were deposited in NIH
BioProject under accession number PRJNA961813. Mass spectrometry data for proteomics were
deposited in the ProteomeXchange Consortium via the PRIDE[175] partner repository under
accession number PXD041723 (reviewer username: reviewer_pxd041723@ebi.ac.uk, and
password: 8pl3ZD1l). Mass spectrometry data for cytokinins were deposited in NIH's National
Metabolomics Data Repository (NMDR) website, the Metabolomics Workbench[152], under
accession number ST002571.
1386

1387 **SUPPLEMENTAL INFORMATION TITLES AND LEGENDS**
1388

Supplemental Figure 1. Evidence that the *drmy1* mutant has ribosomal and translation
defects, associated with Figure 1.
(A) The *drmy1* phenotype is reproduced in the *ap1 cal AP1-GR* system (Ler background). Shown
are representative buds of *ap1 cal AP1-GR* (top row) *and drmy1 ap1 cal AP1-GR* (bottom row) at
day 0 (before DEX induction), day 3 (after 3 DEX inductions, when tissue is collected for RNA,
protein, or cytokinin extraction), and day 5 (after 5 DEX inductions). Arrowheads show sepal
primordia that are of variable number, position, and sizes. Asterisks indicate periphery of the floral
meristem that has limited or no sepal outgrowth. Scale bars, 25 μm.
(B) Summary of the inflorescence RNA-seq and proteomics datasets. Shown are numbers of
genes in each category. Down, downregulated in *drmy1*; NS, not significantly changed between
drmy1 and WT; Up, upregulated in *drmy1*; NA, not available. Note that in the combined dataset
(gene-protein pairs), different genes encoding for the same protein were separately counted, so
were different proteins encoded by the same gene. See also Supplemental Dataset 1.
(C) Violin and box plots of log2 fold change in RNA between *drmy1* and WT in induced *ap1 cal*
AP1-GR inflorescence, for genes encoding ribosomal components ("Structural constituents of the
ribosome" GO:0003735, and its offspring terms) and all other genes involved in translation
("Translation" GO:0006412, and its offspring terms). The following genes are labeled on the
graph: *UL4Z* (AT3G09630), log2FC = -0.492; *UL4Y* (AT5G02870), log2FC = -0.509; *UL18Z*
(AT3G25520), log2FC = -0.459.
(D-G) Fluorescence of a constitutively expressed marker supports the hypothesis that *drmy1* has
reduced translation rate. (D-F) Representative confocal images of *UBQ10::mCherry-RCI2A* in
dissected inflorescences of WT (D), *drmy1* (E), and *ul4y* (F). Numbers show how the signal is
divided based on the stage of floral meristem when quantified (IM+1, inflorescence meristem plus
stage 1; 2, stage 2; 3, stage 3). Scale bars, 25μm. (G) Signal intensity (i.e. integrated density
divided by area) in all images divided as in (D-F). Mean $\pm$ SD are shown. Data was fit using a
two-way ANOVA model with genotype and stage as two additive factors. Asterisks show
statistically significant pairwise contrasts between WT and *drmy1* ($p < 2\times10^{-16}$) and between WT
and *ul4y* ($p = 2.1\times10^{-15}$). Sample sizes: WT IM+1, n = 30; *drmy1* IM+1, n = 22; *ul4y* IM+1, n = 18;
WT stage 2, n = 99; *drmy1* stage 2, n = 100; *ul4y* stage 2, n = 52; WT stage 3, n = 39; *drmy1*
stage 3, n = 27; *ul4y* stage 3, n = 26.
(H) Violin and box plots of log2 fold change in protein level between *drmy1* and WT in induced
ap1 cal AP1-GR inflorescence, for genes in the same categories as in (C). The following genes
are labeled on the graph: *UL4Z* (AT3G09630), log2FC = 0.352; *UL4Y* (AT5G02870), log2FC =
0.811; *UL18Z* (AT3G25520), log2FC = 0.742.
(I) Coherent regulation of gene expression by *drmy1* and AZD-8055. Shown is a contingency
table of genes downregulated (Down), not significantly changed (NS), and upregulated (Up) in
drmy1 vs WT (columns), and in AZD-8055-treated WT vs mock-treated WT (rows). Bold font
shows the number of genes in each category, and gray font shows the expected number of genes
if there were no correlation between two conditions (calculated as row margin × column margin /
total number of genes). Categories where the number of genes is above expectation are

highlighted blue, and categories where the number of genes is below expectation are highlighted

red. Chi-square test $p < 2.2 \times 10^{-16}$.

(J) Gene ontology enrichment of genes coherently downregulated by both *drmy1* and AZD-8055.

Shown are the top 8 terms and their enrichment p-values. Note that the first 7 terms are all related

to ribosome and translation. A complete list can be found in Supplementary Dataset 3.

1434

Supplemental Figure 2. Ribosomal mutations enhance the *drmy1* phenotype, associated

with Figure 2.

(A-H) Examples of stage 5 buds from *drmy1* (A), *drmy1 ul4z* (B-D), *drmy1 ul4y* (E-F), and *drmy1*

ul18z/+ (G-H). In (B,E,G) sepal primordia within each bud have bigger size differences than typical

drmy1 single mutant buds; asterisks show giant outer sepal primordia and brackets show bud

peripheral regions with little or no primordium outgrowth. In (C,F,H), arrowheads show 6 sepal

primordia within each bud, which does not occur in *drmy1*. In (D,H), asterisks show the presence

of two outer sepal primordia within a bud, instead of one in *drmy1*. Scale bars, 25µm.

(I-J) Quantification of sepal primordium number (I) and positional variability (J), comparing each

of *drmy1 ul4z* (n = 60), *drmy1 ul4y* (n = 61), and *drmy1 ul18z/+* (n = 69) with *drmy1* (n = 67). "ns"

indicates no significant difference in Fisher's contingency table tests (I) and Wilcoxon's rank sum

tests (J) respectively. Data for *drmy1* is reused from Figure 2H, 2J.

(K) Dissected young silique of a *drmy1 ul18z/+* plant. Arrowheads point to aborted ovules. Scale

bar, 200 µm.

1449

Supplemental Figure 3. Sepal primordia in ribosome and TOR mutants catch up in growth
to form uniformly sized mature sepals within the bud, associated with Figure 3.
(A-F) Inflorescences (left) of WT (A), *drmy1* (B), *ul4z* (C), *ul4y* (D), *ul18z* (E), and *lst8-1-1* (F), with
boxed regions enlarged (right). Blue arrowheads show sepals of regular length, and red
arrowheads show sepals shorter than others. Note that sepals in *drmy1* were unable to close due
to unequal lengths, while sepals in *ul4z*, *ul4y*, and *ul18z*, and close like in WT. Sepals in *lst8-1-1*
were unable to close although there is no apparent variation in length. Scale bars, 0.5 mm.
(G-L) Dissected sepals from a bud of WT (G), *drmy1* (H), *ul4z* (I), *ul4y* (J), *ul18z* (K), and two
buds of *lst8-1-1* (L). Note that sepals in the *drmy1* bud are of different sizes. Sepals within each
bud of *ul4z*, *ul4y*, *ul18z*, and *lst8-1-1* are of similar sizes, although there can be variation between
different buds of the same genotype. O, outer sepal. I, inner sepal. L, lateral sepal. Scale bars,
200 μm.
(M) Quantification of between-flower variability of sepal length. Length was measured from all
imaged sepals of each genotype and each position (outer, inner, lateral), and coefficient of
variation (CV) was calculated. A two-sided permutation test (100,000 permutations) for CV
difference not equating to zero was done for each pair of genotypes, and results were represented
by letters. Sample size: Outer sepal, WT n = 35, *drmy1* n = 43, *ul4z* n = 37, *ul4y* n = 42, *ul18z* n
= 39, *lst8-1-1* n = 43. Inner sepal, WT n = 34, *drmy1* n = 46, *ul4z* n = 38, *ul4y* n = 44, *ul18z* n =
37, *lst8-1-1* n = 44. Lateral sepal, WT n = 65, *drmy1* n = 84, *ul4z* n = 81, *ul4y* n = 89, *ul18z* n =
76, *lst8-1-1* n = 82.
(N) Quantification of within-flower variability of length. Flowers with length data from at least four
sepals were analyzed. A CV of length from all sepals within each flower was calculated, and mean
± SD was plotted, grouped by genotype. A Wilcoxon rank sum test was done for each pair of
genotypes, and results were represented by letters. Sample size: WT n = 31 buds, *drmy1* n = 38
buds, *ul4z* n = 33 buds, *ul4y* n = 36 buds, *ul18z* n = 32 buds, *lst8-1-1* n = 39 buds.
(O-Q) Live imaging of sepal development from stage 3 to 6 in WT (O), *drmy1* (P), and *ul4y* (Q),
showing chlorophyll or propidium iodide channel, and Gaussian curvature of the surface. Note
that both *drmy1* and *ul4y* have inner sepals that initiate late (day 2, asterisk). The *drmy1* inner
sepal develops slowly, and leaves the bud open at day 3 (red arrowhead). The *ul4y* inner sepal
catches up with the rest of the sepals and closes the bud (blue arrowhead). Scale bars, 25 μm.
1480

Supplemental Figure 4. Inhibition of TOR activity and translation causes auxin maxima
formation at variable positions, correlated with variable positions of sepal primordia,
associated with Figure 4.
(A-E) Variable patterning of auxin signaling in *drmy1*, *ul4y*, and CHX-treated WT buds
corresponds to variable sepal initiation. During the time course, top rows show *DR5::3xVENUS-*
N7 (yellow), middle rows show composites of *DR5* (yellow) and Chlorophyll (magenta), and
bottom rows show Gaussian curvature of the Chlorophyll channel. In the last time point, buds
were stained with propidium iodide (top), and Gaussian curvature of the propidium iodide channel
is shown on the bottom.
(A) In WT, four robustly positioned auxin maxima at day 1 correlates with four robustly positioned
sepal primordia at day 4 (blue arrowheads). (B) In *drmy1*, at day 1 there are three robustly
positioned auxin maxima (blue arrowheads). At day 2, a diffuse band of auxin signaling occurs in
the adaxial periphery of the bud, joining with one of the lateral auxin maxima (red bracket). At day
3, this diffuse band splits into three auxin maxima (red arrowheads), making a total of 5. The
maxima correlate with the five sepal primordia at day 4, three at robust positions (blue
arrowheads) and two at irregular positions (red arrowheads). (C) In *ul4y*, at day 1 there are two
auxin maxima at robust positions (blue arrowheads), one at robust lateral position but much
weaker (red arrowhead), and a band of weak auxin signaling in the adaxial periphery of the bud
(red bracket). At day 2, the weak auxin maxima at the lateral position got stronger, and the weak
band at the adaxial position split into two auxin maxima (red arrowheads). These five auxin
maxima correspond to the five sepal primordia at day 3, three in robust positions (blue
arrowheads) and two in irregular positions (red arrowheads). (D) In a mock-treated WT bud, four
robust auxin maxima at day 6 of the treatment grow into four robust sepal primordia at day 9 (blue
arrowheads). (E) In a CHX-treated WT bud, at day 6 there are three stronger auxin maxima (blue
arrowheads) and two weaker ones (red arrowheads), corresponding to three bigger regions of
outgrowth (blue arrowheads) and two smaller ones (red arrowheads) at day 9. For ease of display,
the DR5 channel in CHX-treated WT was brightened three times relative to mock.
(F-I) TOR inhibition using Torin2 causes increased cytokinin signaling, and occasional spatial
variability in auxin and cytokinin signaling. (F) Late stage 2 buds of WT treated *in vivo* with mock
or 2 nmol Torin2 twice a day for 15 days. Shown are *DR5::3xVENUS-N7* in yellow, *TCS::GFP* in
cyan, and both merged with propidium iodide in magenta. Note that 3/16 (19%) buds had variable
number and position of DR5 and TCS maxima, and 13/16 (81%) had robust DR5 and TCS
maxima, although TCS intensity is higher than mock in both cases. (G) Quantification of TCS
intensity from maximum intensity projection images, normalized to the mean of WT mock. Shown
are mean ± SD. Asterisk shows statistical significance in a two-tailed Student's t-test ($p = 1.2 \times 10^{-4}$). (H) Circular histograms of DR5 signal distribution (mean ± SD). (I) Circular histograms of TCS
signal distribution (mean ± SD). For calculation of circular histograms, please see Figure 4
legends and Materials and Methods. Sample size: WT mock, n = 11 buds; WT Torin2, n = 16
buds. Scale bars in all micrographs, 25 μm.

Supplemental Figure 5. Translation of uORF-containing ARFs is not universally
downregulated in _drmy1_, associated with Figure 5.
(A) _drmy1_ has a lower protein-transcript ratio than WT for genes with at least 2 uORFs. 5,086
transcript-protein pairs in our inflorescence dataset were grouped according to the maximum
number of uORFs in all transcript isoforms (0, n = 3,485; 1, n = 874; ≥ 2, n = 724) (von Arnim et
al., 2014). For each pair, protein-transcript ratio was calculated, log-transformed, and the
difference between _drmy1_ and WT was plotted. A negative value means this gene has less protein
per transcript in _drmy1_ than WT, and could indicate reduced translation or protein stability.
Medians for each group: 0 uORF, -0.00367; 1 uORF, -0.00808; ≥ 2 uORFs, -0.0243. Asterisk
show statistically significant difference from Group 0 in a Wilcoxon rank sum test (p = $3.167×10^{-4}$
), while ns means no significant difference from Group 0 (p = 0.167).
(B-D) There is no universal decrease in the expression of uORF-containing ARF reporters. (B)
Transcript level of three activator _ARF_s (_ARF5_, _ARF6_, _ARF8_) and two repressor _ARF_s (_ARF3_,
ARF10) in inflorescence RNA-seq (n = 3 per genotype). _ARF3_, _ARF5_, and _ARF6_ contain
uORFs before the main ORF, and _ARF8_ and _ARF10_ do not. Asterisk indicates statistically
significant differences between WT and _drmy1_ from DESeq2 output. p values: _ARF3_, p = 0.583;
ARF5, p = 0.497; _ARF6_, p = 0.603; _ARF8_, p = 0.058; _ARF10_ p = 0.019. (C) Transcriptional
reporters for these ARFs in stage 2 buds of WT and _drmy1_ (cyan, GFP or YFP; magenta,
propidium iodide). Note that the _pARF3_, _pARF5_, and _pARF6_ reporters contain the same uORFs
as the genes, reflecting a combination of transcriptional and uORF regulations. Scale bars, 20
μm. (D) Quantification of GFP intensity. Sample size: _pARF3_ WT, n = 22; _pARF3 drmy1_, n = 25;
pARF5 WT, n = 22; _pARF5 drmy1_, n = 22; _pARF6_ WT, n = 19; _pARF6 drmy1_, n = 28; _pARF8_
WT, n = 25; _pARF8 drmy1_, n = 31; _pARF10_ WT, n = 20; _pARF10 drmy1_, n = 29. Asterisks show
statistically significant differences between WT and _drmy1_ in Wilcoxon rank sum tests. p values:
pARF3, p = 0.3797; _pARF5_, p = $6.22×10^{-5}$; _pARF6_, p = $2.868×10^{-13}$; _pARF8_, p = 0.5127;
pARF10 p = $7.073×10^{-14}$.

Supplemental Figure 6. Cytokinin signaling causes variability in mature sepal number and
size in *drmy1*, associated with Figure 5.
Shown are top-view inflorescence images of WT (A), *arr1,10,12* (B), *wol* (C), *drmy1* (D), *drmy1*
arr1,10,12 (E), and *drmy1 wol* (F), with boxed areas of individual buds enlarged and shown on
the right. In the enlarged views, blue arrowheads point to sepals of regular size, and red
arrowheads point to sepals that are much smaller. Scale bars, 0.5 mm.
1552

Supplemental Figure 7. Investigating other mechanisms that may explain the observed
changes in hormone signaling.
(A) Cytokinin abundance does not significantly change in *drmy1*. Shown is mean ± SD of levels
of trans-zeatin (tZ), cis-Zeatin (cZ), N^6-(Δ^2-Isopentenyl)adenine (iP), trans-Zeatin riboside (tZR),
cis-Zeatin riboside (cZR), and N^6-(Δ^2-Isopentenyl)adenosine (iPR) quantified by LC-MS in
induced WT and *drmy1* inflorescences of *ap1 cal AP1-GR* background. Levels are normalized to
the mean tZ level in WT. Sample size: n = 5 for WT; n = 6 for *drmy1*. ns, no significant difference
between WT and *drmy1* in two-sided Wilcoxon rank sum tests. P-values: tZ, p = 0.2468; cZ, p =
0.7922; iP, p = 0.2468; tZR, p = 0.1775; cZR, p = 0.6623; iPR, p = 0.6623.
(B) Expression of cytokinin signaling components in WT vs *drmy1* inflorescences of *ap1 cal AP1-*
GR background (from Supplemental Dataset 1).
(C-D) The *ARR7-llama GFP-nes* reporter responds to externally applied cytokinin. Shown are
GFP images of the same bud before (C) or after (D) 5 hours of 200 µM BAP treatment. Images
are representative of n = 9 buds from two independent lines.
(E-H) CHX (E-F) or AZD-8055 (G-H) treatments do not change the subcellular localization of GFP-
nes. Images are representative of n = 10 buds (E), n = 9 buds (F), n = 10 buds (G), and n = 11
buds (H). For (C-H), each image was brightened to reveal patterns of GFP distribution. A square
region taken from the image containing 5-10 cells is enlarged and shown on the top right. Within
the square, GFP intensity was quantified along the dotted line and plotted on the bottom right. X-
axis, pixels (range 0-238). Y-axis, GFP intensity (smoothened by taking the average intensity of
11-pixel neighborhoods; range 90-175 in gray value). Scale bars, 25 µm.
(I-J) Response of *pAHP3::AHP3-GFP* (I) and *pUBQ10::mCherry-RCI2A* (J) to mock, CHX, and
AZD-8055 treatments for 72 hours. Scale bars, 25 µm. For (I), images are representative of n =
13 (mock), n = 15 (CHX), and n = 16 (AZD-8055) buds in two experiments. For (J), images are
representative of n = 12 (mock), n = 9 (CHX), and n = 10 (AZD-8055) buds in two experiments.
(K-N) The *ARR7-llama GFP-nes* construct partially rescues the mature sepal variability in *drmy1*.
Shown are inflorescence images of WT (K), *ARR7-llama GFP-nes* (L), *drmy1* (M), and *drmy1*
ARR7-llama GFP-nes (N). The boxed regions were enlarged and shown on the right of each
panel. Note that while *drmy1* buds have normal-sized (blue arrowheads) and smaller (red
arrowheads) sepals, some buds in *drmy1 ARR7-llama GFP-nes* have robustly sized sepals (N,
middle) while others still show variability (N, right). Scale bars, 0.5 mm.
(O-P) *drmy1* has decreased and disrupted pattern of DII degradation. **(O)** Representative images
of WT and *drmy1* showing *DII-n3xVENUS* (cyan), *mDII-ntdTomato* (magenta), and merge. For
ease of display, the *VENUS* channel was brightened 3 times relative to the *tdTomato* channel.
Scale bars, 25 µm. **(P)** Quantification of VENUS/tdTomato ratio. A background of 6 gray value
per pixel (determined in blank regions) were subtracted before calculation of ratios. Sample size:
WT, n = 8 buds; *drmy1*, n = 19 buds. Asterisk shows statistically significant difference in a
Wilcoxon rank sum test (p = 0.01335).
1591

1592   **Supplemental Dataset 1. Inflorescence RNA-seq and proteomics.**

1593

1594   **Supplemental Dataset 2. Unprocessed ribosome profiles.**

1595

1596   **Supplemental Dataset 3. Seedling RNA-seq.**

1597

1598   **Supplemental Dataset 4. Data used in graphs.**

1599

1600   **Supplemental Dataset 5. DNA sequences.**

1601

REFERENCES

1602

1. Kaneko, K. (2007). Evolution of robustness to noise and mutation in gene expression
dynamics. PLoS One *2*, e434. 10.1371/journal.pone.0000434.
2. Lachowiec, J., Queitsch, C., and Kliebenstein, D.J. (2016). Molecular mechanisms
governing differential robustness of development and environmental responses in plants.
Ann. Bot. *117*, 795–809. 10.1093/aob/mcv151.
3. Zabinsky, R.A., Mason, G.A., Queitsch, C., and Jarosz, D.F. (2019). It's not magic –
Hsp90 and its effects on genetic and epigenetic variation. Semin. Cell Dev. Biol. *88*, 21–
35. 10.1016/j.semcdb.2018.05.015.
4. Waddington, C.H. (1942). Canalization of Development and the Inheritance of Acquired
Characters. Nature *150*, 563–656.
5. Twitty, V.C., and Schwind, J.L. (1931). The growth of eyes and limbs transplanted
heteroplastically between two species of Amblystoma. J. Exp. Zool. *59*, 61–86.
10.1002/jez.1400590105.
6. Silber, S.J. (1976). Growth of Baby Kidneys Transplanted Into Adults. Arch. Surg. *111*,
75–77. 10.1001/archsurg.1976.01360190077014.
7. Summerbell, D., and Lewis, J.H. (1975). Time, place and positional value in the chick
limb bud. J. Embryol. Exp. Morphol. *33*, 621–643. 10.1242/dev.33.3.621.
8. Parker, J., and Struhl, G. (2020). Control of *Drosophila* wing size by morphogen range
and hormonal gating. Proc. Natl. Acad. Sci. U. S. A. *117*, 31935–31944.
10.1073/pnas.2018196117.
9. Garelli, A., Gontijo, A.M., Miguela, V., Caparros, E., and Dominguez, M. (2012). Imaginal
discs secrete insulin-like peptide 8 to mediate plasticity of growth and maturation.
Science *336*, 579–582. 10.1126/science.1216735.
10. Crickmore, M.A., and Mann, R.S. (2006). Hox control of organ size by regulation of
morphogen production and mobility. Science *313*, 63–68. 10.1126/science.1128650.
11. Recasens-Alvarez, C., Ferreira, A., and Milán, M. (2017). JAK/STAT controls organ size
and fate specification by regulating morphogen production and signalling. Nat. Commun.
8, 10–12. 10.1038/ncomms13815.
12. Irish, V.F. (2008). The Arabidopsis petal: a model for plant organogenesis. Trends Plant
Sci. *13*, 430–436. 10.1016/j.tplants.2008.05.006.
13. Fenster, C.B., Armbruster, W.S., Wilson, P., Dudash, M.R., and Thomson, J.D. (2004).
Pollination syndromes and floral specialization. Annu. Rev. Ecol. Evol. Syst. *35*, 375–403.
10.1146/annurev.ecolsys.34.011802.132347.
14. King, S., Beck, F., and Lüttge, U. (2004). On the mystery of the golden angle in
phyllotaxis. Plant, Cell Environ. *27*, 685–695. 10.1111/j.1365-3040.2004.01185.x.
15. Smith, R.S., Guyomarc'h, S., Mandel, T., Reinhardt, D., Kuhlemeier, C., and
Prusinkiewicz, P. (2006). A plausible model of phyllotaxis. Proc. Natl. Acad. Sci. U. S. A.
103, 1301–1306. 10.1073/pnas.0510457103.
16. Besnard, F., Refahi, Y., Morin, V., Marteaux, B., Brunoud, G., Chambrier, P., Rozier, F.,
Mirabet, V., Legrand, J., Lainé, S., et al. (2014). Cytokinin signalling inhibitory fields
provide robustness to phyllotaxis. Nature *505*, 417–421. 10.1038/nature12791.
17. Queitsch, C., Sangster, T.A., and Lindquist, S. (2002). Hsp90 as a capacitor of
phenotypic variation. Nature *417*, 618–624. 10.1038/nature749.
18. Rutherford, S.L., and Lindquist, S. (1998). Hsp90 as a capacitor for morphological
evolution. Nature *396*, 336–342. 10.1038/24550.
19. Lehner, B., Crombie, C., Tischler, J., Fortunato, A., and Fraser, A.G. (2006). Systematic
mapping of genetic interactions in *Caenorhabditis elegans* identifies common modifiers of
diverse signaling pathways. Nat. Genet. *38*, 896–903. 10.1038/ng1844.
20. Levy, S.F., and Siegal, M.L. (2008). Network hubs buffer environmental variation in

Saccharomyces cerevisiae. PLoS Biol. *6*, 2588–2604. 10.1371/journal.pbio.0060264.

21.   Folta, A., Severing, E.I., Krauskopf, J., van de Geest, H., Verver, J., Nap, J.P., and
Mlynarova, L. (2014). Over-expression of Arabidopsis AtCHR23 chromatin remodeling
ATPase results in increased variability of growth and gene expression. BMC Plant Biol.
14, 76. 10.1186/1471-2229-14-76.

22.   Hintze, M., Katsanos, D., Shahrezaei, V., and Barkoulas, M. (2021). Phenotypic
Robustness of Epidermal Stem Cell Number in *C. elegans* Is Modulated by the Activity of
the Conserved N-acetyltransferase nath-10/NAT10. Front. Cell Dev. Biol. *9*, 640856.
10.3389/fcell.2021.640856.

23.   Marygold, S.J., Roote, J., Reuter, G., Lambertsson, A., Ashburner, M., Millburn, G.H.,
Harrison, P.M., Yu, Z., Kenmochi, N., Kaufman, T.C., et al. (2007). The ribosomal protein
genes and Minute loci of *Drosophila melanogaster*. Genome Biol. *8*, R216. 10.1186/gb-
2007-8-10-r216.

24.   Wang, S., Kurepa, J., and Smalle, J.A. (2009). The *Arabidopsis* 26S proteasome subunit
RPN1a is required for optimal plant growth and stress responses. Plant Cell Physiol. *50*,
1721–1725. 10.1093/pcp/pcp105.

25.   Gallois, J.L., Guyon-Debast, A., Lecureuil, A., Vezon, D., Carpentier, V., Bonhomme, S.,
and Guerche, P. (2009). The *Arabidopsis* proteasome RPT5 subunits are essential for
gametophyte development and show accession-dependent redundancy. Plant Cell *21*,
442–459. 10.1105/tpc.108.062372.

26.   Hong, L., Dumond, M., Tsugawa, S., Sapala, A., Routier-Kierzkowska, A.-L., Zhou, Y.,
Chen, C., Kiss, A., Zhu, M., Hamant, O., et al. (2016). Variable Cell Growth Yields
Reproducible Organ Development through Spatiotemporal Averaging. Dev. Cell *38*, 15–
32. 10.1016/j.devcel.2016.06.016.

27.   Zhu, M., Chen, W., Mirabet, V., Hong, L., Bovio, S., Strauss, S., Schwarz, E.M.,
Tsugawa, S., Wang, Z., Smith, R.S., et al. (2020). Robust organ size requires robust
timing of initiation orchestrated by focused auxin and cytokinin signalling. Nat. Plants *6*,
686–698. 10.1038/s41477-020-0666-7.

28.   Roeder, A.H.K. (2021). Arabidopsis sepals: A model system for the emergent process of
morphogenesis. Quant. Plant Biol. *2*, e14. 10.1017/qpb.2021.12.

29.   Trinh, D.-C., Martin, M., Bald, L., Maizel, A., Trehin, C., and Hamant, O. (2023).
Increased gene expression variability hinders the formation of regional mechanical
conflicts leading to reduced organ shape robustness. Proc. Natl. Acad. Sci. *120*,
e2302441120. 10.1073/pnas.2302441120.

30.   Jing, H., and Strader, L.C. (2019). Interplay of auxin and cytokinin in lateral root
development. Int. J. Mol. Sci. *20*, 486. 10.3390/ijms20030486.

31.   Hussain, S., Nanda, S., Zhang, J., Rehmani, M.I.A., Suleman, M., Li, G., and Hou, H.
(2021). Auxin and Cytokinin Interplay during Leaf Morphogenesis and Phyllotaxy. Plants
10, 1732. 10.3390/plants10081732.

32.   Schaller, G.E., Bishopp, A., and Kieber, J.J. (2015). The yin-yang of hormones: Cytokinin
and auxin interactions in plant development. Plant Cell *27*, 44–63.
10.1105/tpc.114.133595.

33.   Ingargiola, C., Duarte, G.T., Robaglia, C., Leprince, A.-S., and Meyer, C. (2020). The plant
target of rapamycin: A conduc TOR of nutrition and metabolism in photosynthetic
organisms. Genes (Basel). *11*, 1285. 10.3390/genes11111285.

34.   Scarpin, M.R., Leiboff, S., and Brunkard, J.O. (2020). Parallel global profiling of plant tor
dynamics reveals a conserved role for LARP1 in translation. eLife *9*, e58795.
10.7554/eLife.58795.

35.   Yu, H., Ito, T., Wellmer, F., and Meyerowitz, E.M. (2004). Repression of AGAMOUS-LIKE
24 is a crucial step in promoting flower development. Nat. Genet. *36*, 157–161.
10.1038/ng1286.

36. Wellmer, F., Alves-Ferreira, M., Dubois, A., Riechmann, J.L., and Meyerowitz, E.M.
(2006). Genome-wide analysis of gene expression during early *Arabidopsis* flower
development. PLoS Genet. *2*, 1012–1024. 10.1371/journal.pgen.0020117.
37. Smyth, D.R., Bowman, J.L., and Meyerowitz, E.M. (1990). Early flower development in
Arabidopsis. Plant Cell *2*, 755–767. 10.1105/tpc.2.8.755.
38. Goodman, C.A., and Hornberger, T.A. (2013). Measuring protein synthesis with SUnSET:
A valid alternative to traditional techniques? Exerc. Sport Sci. Rev. *41*, 107–115.
10.1097/JES.0b013e3182798a95.
39. Van Hoewyk, D. (2016). Use of the non-radioactive SUnSET method to detect decreased
protein synthesis in proteasome inhibited Arabidopsis roots. Plant Methods *12*, 20.
10.1186/S13007-016-0120-Z.
40. Scarpin, M.R., Busche, M., Martinez, R.E., Harper, L.C., Reiser, L., Szakonyi, D.,
Merchante, C., Lan, T., Xiong, W., Mo, B., et al. (2023). An updated nomenclature for
plant ribosomal protein genes. Plant Cell *35*, 640–643. 10.1093/plcell/koac333.
41. Schepetilnikov, M., and Ryabova, L.A. (2018). Recent discoveries on the role of tor
(Target of rapamycin) signaling in translation in plants. Plant Physiol. *176*, 1095–1105.
10.1104/pp.17.01243.
42. Scarpin, M.R., Simmons, C.H., and Brunkard, J.O. (2022). Translating across kingdoms:
target of rapamycin promotes protein synthesis through conserved and divergent
pathways in plants. J. Exp. Bot. *73*, 7016–7025. 10.1093/jxb/erac267.
43. Brunkard, J.O. (2020). Exaptive Evolution of Target of Rapamycin Signaling in
Multicellular Eukaryotes. Dev. Cell *54*, 142–155. 10.1016/j.devcel.2020.06.022.
44. Battaglioni, S., Benjamin, D., Wälchli, M., Maier, T., and Hall, M.N. (2022). mTOR
substrate phosphorylation in growth control. Cell *185*, 1814–1836.
10.1016/j.cell.2022.04.013.
45. Valvezan, A.J., and Manning, B.D. (2019). Molecular logic of mTORC1 signalling as a
metabolic rheostat. Nat. Metab. *1*, 321–333. 10.1038/s42255-019-0038-7.
46. Schepetilnikov, M., Kobayashi, K., Geldreich, A., Caranta, C., Robaglia, C., Keller, M.,
and Ryabova, L.A. (2011). Viral factor TAV recruits TOR/S6K1 signalling to activate
reinitiation after long ORF translation. EMBO J. *30*, 1343–1356. 10.1038/emboj.2011.39.
47. Schepetilnikov, M., Dimitrova, M., Mancera-Martínez, E., Geldreich, A., Keller, M., and
Ryabova, L.A. (2013). TOR and S6K1 promote translation reinitiation of uORF-containing
mRNAs via phosphorylation of eIF3h. EMBO J. *32*, 1087–1102. 10.1038/emboj.2013.61.
48. Schepetilnikov, M., Makarian, J., Srour, O., Geldreich, A., Yang, Z., Chicher, J.,
Hammann, P., and Ryabova, L.A. (2017). GTPase ROP2 binds and promotes activation
of target of rapamycin, TOR, in response to auxin. EMBO J. *36*, 886–903.
10.15252/embj.201694816.
49. Chen, G.H., Liu, M.J., Xiong, Y., Sheen, J., and Wu, S.H. (2018). TOR and RPS6
transmit light signals to enhance protein translation in deetiolating Arabidopsis seedlings.
Proc. Natl. Acad. Sci. U. S. A. *115*, 12823–12828. 10.1073/pnas.1809526115.
50. Mancera-Martínez, E., Dong, Y., Makarian, J., Srour, O., Thiébeauld, O., Jamsheer, M.,
Chicher, J., Hammann, P., Schepetilnikov, M., and Ryabova, L.A. (2021).
Phosphorylation of a reinitiation supporting protein, RISP, determines its function in
translation reinitiation. Nucleic Acids Res. *49*, 6908–6924. 10.1093/nar/gkab501.
51. Dong, Y., Srour, O., Lukhovitskaya, N., Makarian, J., Baumberger, N., Galzitskaya, O.,
Elser, D., Schepetilnikov, M., and Ryabova, L.A. (2023). Functional analogs of
mammalian 4E-BPs reveal a role for TOR in global plant translation. Cell Rep. *42*,
112892. 10.1016/j.celrep.2023.112892.
52. Dasgupta, A., Camacho, R.A.U., Enganti, R., Cho, S.K., Tucker, L.L., Torreverde, J.S.,
Abraham, P.E., and Arnim, A.G. von (2024). A phosphorylation-deficient ribosomal
protein eS6 is largely functional in Arabidopsis thaliana, rescuing mutant defects from

1754    global translation and gene expression to photosynthesis and growth. Plant Direct *8*,
1755    e566. 10.1002/pld3.566.

1756 53. Xiong, Y., McCormack, M., Li, L., Hall, Q., Xiang, C., and Sheen, J. (2013). Glucose-TOR
1757    signalling reprograms the transcriptome and activates meristems. Nature *496*, 181–186.
1758    10.1038/nature12030.

1759 54. Riegler, S., Servi, L., Scarpin, M.R., Godoy Herz, M.A., Kubaczka, M.G., Venhuizen, P.,
1760    Meyer, C., Brunkard, J.O., Kalyna, M., Barta, A., et al. (2021). Light regulates alternative
1761    splicing outcomes via the TOR kinase pathway. Cell Rep. *36*, 109676.
1762    10.1016/j.celrep.2021.109676.

1763 55. Obomighie, I., Lapenas, K., Murphy, B.E., Bowles, A.M.C., Bechtold, U., and Prischi, F.
1764    (2021). The Role of Ribosomal Protein S6 Kinases in Plant Homeostasis. Front. Mol.
1765    Biosci. *8*, 636560. 10.3389/fmolb.2021.636560.

1766 56. Xiong, Y., and Sheen, J. (2012). Rapamycin and glucose-target of rapamycin (TOR)
1767    protein signaling in plants. J. Biol. Chem. *287*, 2836–2842. 10.1074/jbc.M111.300749.

1768 57. Moreau, M., Azzopardi, M., Clé ment, G., Dobrenel, T., Marchive, C., Renne, C., Martin-
1769    Magniette, M.L., Taconnat, L., Renou, J.P., Robaglia, C., et al. (2012). Mutations in the
1770    *Arabidopsis* homolog of LST8/GβL, a partner of the target of Rapamycin kinase, impair
1771    plant growth, flowering, and metabolic adaptation to long days. Plant Cell *24*, 463–481.
1772    10.1105/tpc.111.091306.

1773 58. Brunkard, J.O., Xu, M., Regina Scarpin, M., Chatterjee, S., Shemyakina, E.A., Goodman,
1774    H.M., and Zambryski, P. (2020). TOR dynamically regulates plant cell-cell transport. Proc.
1775    Natl. Acad. Sci. U. S. A. *117*, 5049–5058. 10.1073/pnas.1919196117.

1776 59. Menand, B., Desnos, T., Nussaume, L., Bergert, F., Bouchez, D., Meyer, C., and
1777    Robaglia, C. (2002). Expression and disruption of the *Arabidopsis TOR* (target of
1778    rapamycin) gene. Proc. Natl. Acad. Sci. U. S. A. *99*, 6422–6427.
1779    10.1073/pnas.092141899.

1780 60. Zhao, Z., Andersen, S.U., Ljung, K., Dolezal, K., Miotk, A., Schultheiss, S.J., and
1781    Lohmann, J.U. (2010). Hormonal control of the shoot stem-cell niche. Nature *465*, 1089–
1782    1092. 10.1038/nature09126.

1783 61. Yoshida, S., Mandel, T., and Kuhlemeier, C. (2011). Stem cell activation by light guides
1784    plant organogenesis. Genes Dev. *25*, 1439–1450. 10.1101/gad.631211.

1785 62. Du, F., Guan, C., and Jiao, Y. (2018). Molecular Mechanisms of Leaf Morphogenesis.
1786    Mol. Plant *11*, 1117–1134. 10.1016/j.molp.2018.06.006.

1787 63. Taylor-Teeples, M., Lanctot, A., and Nemhauser, J.L. (2016). As above, so below:
1788    Auxin's role in lateral organ development. Dev. Biol. *419*, 156–164.
1789    10.1016/j.ydbio.2016.03.020.

1790 64. Nishimura, T., Wada, T., Yamamoto, K.T., and Okada, K. (2005). The *Arabidopsis* STV1
1791    protein, responsible for translation reinitiation, is required for auxin-mediated gynoecium
1792    patterning. Plant Cell *17*, 2940–2953. 10.1105/tpc.105.036533.

1793 65. Zhou, F., Roy, B., and von Arnim, A.G. (2010). Translation reinitiation and development
1794    are compromised in similar ways by mutations in translation initiation factor eIF3h and the
1795    ribosomal protein RPL24. BMC Plant Biol. *10*, 193. 10.1186/1471-2229-10-193.

1796 66. Rosado, A., Li, R., Van De Ven, W., Hsu, E., and Raikhel, N. V. (2012). *Arabidopsis*
1797    ribosomal proteins control developmental programs through translational regulation of
1798    auxin response factors. Proc. Natl. Acad. Sci. U. S. A. *109*, 19537–19544.
1799    10.1073/pnas.1214774109.

1800 67. Cancé, C., Martin-Arevalillo, R., Boubekeur, K., and Dumas, R. (2022). Auxin response
1801    factors are keys to the many auxin doors. New Phytol. *235*, 402–419.
1802    10.1111/nph.18159.

1803 68. Kim, B.H., Cai, X., Vaughn, J.N., and Von Arnim, A.G. (2007). On the functions of the h
1804    subunit of eukaryotic initiation factor 3 in late stages of translation initiation. Genome Biol.

8, R60. 10.1186/gb-2007-8-4-r60.

69.   Von Arnim, A.G., Jia, Q., and Vaughn, J.N. (2014). Regulation of plant translation by
upstream open reading frames. Plant Sci. *214*, 1–12. 10.1016/j.plantsci.2013.09.006.

70.   Jones, B., Ljung, K., Gunnerås, S.A., Petersson, S. V., Tarkowski, P., Graham, N., May,
S., Dolezal, K., and Sandberg, G. (2010). Cytokinin regulation of auxin synthesis in
Arabidopsis involves a homeostatic feedback loop regulated via auxin and cytokinin
signal transduction. Plant Cell *22*, 2956–2969. 10.1105/tpc.110.074856.

71.   Marhavý, P., Duclercq, J., Weller, B., Feraru, E., Bielach, A., Offringa, R., Friml, J.,
Schwechheimer, C., Murphy, A., and Benková, E. (2014). Cytokinin controls polarity of
PIN1-dependent Auxin transport during lateral root organogenesis. Curr. Biol. *24*, 1031–
1037. 10.1016/j.cub.2014.04.002.

72.   Růžička, K., Šimášková, M., Duclercq, J., Petrášek, J., Zažímalová, E., Simon, S., Friml,
J., Van Montagu, M.C.E., and Benková, E. (2009). Cytokinin regulates root meristem
activity via modulation of the polar auxin transport. Proc. Natl. Acad. Sci. U. S. A. *106*,
4284–4289. 10.1073/pnas.0900060106.

73.   Müller, B., and Sheen, J. (2007). Advances in cytokinin signaling. Science *318*, 68–69.
10.1126/science.1145461.

74.   Karunadasa, S.S., Kurepa, J., Shull, T.E., and Smalle, J.A. (2020). Cytokinin-induced
protein synthesis suppresses growth and osmotic stress tolerance. New Phytol. *227*, 50–
64. 10.1111/nph.16519.

75.   Szweykowska, A., Gwoidi, E., and Spychala, M. (1980). The Cytokinin Control of Protein
Synthesis in Plants. In Metabolism and Molecular Activities of Cytokinins, J. Guern and C.
Peaud-Lenoel, eds. (Springer-Verlag Berlin Heidelberg), pp. 212–217. 10.1007/978-3-
642-68035-9.

76.   Woźny, A., and Gwóźdź, E.A. (1980). The effect of cytokinin on the polyribosome
formation in cucumber cotyledons. Biochem. und Physiol. der Pflanz. *175*, 476–480.
10.1016/s0015-3796(80)80032-1.

77.   Klyachko, N.L., Yakovleva, L.A., Shakirova, F.M., and Kulaeva, O.N. (1982). Cell-free
translation of polyribosomes from detached pumpkin cotyledons: Effects of starvation and
cytokinin. Biol. Plant. *24*, 374–380. 10.1007/BF02909106.

78.   Short, K.C., Tepfer, D.A., and Fosket, D.E. (1974). Regulation of polyribosome formation
and cell division in cultured soybean cells by cytokinin. J. Cell Sci. *15*, 75–87.

79.   Brenner, W.G., and Schmülling, T. (2012). Transcript profiling of cytokinin action in
Arabidopsis roots and shoots discovers largely similar but also organ-specific responses.
BMC Plant Biol. *12*, 112. 10.1186/1471-2229-12-112.

80.   Kiba, T., Naitou, T., Koizumi, N., Yamashino, T., Sakakibara, H., and Mizuno, T. (2005).
Combinatorial microarray analysis revealing Arabidopsis genes implicated in cytokinin
responses through the His→Asp phosphorelay circuitry. Plant Cell Physiol. *46*, 339–355.
10.1093/pcp/pci033.

81.   Černý, M., Kuklová, A., Hoehenwarter, W., Fragner, L., Novák, O., Rotková, G., Jedelský,
P.L., Žáková, K., Šmehilová, M., Strnad, M., et al. (2013). Proteome and metabolome
profiling of cytokinin action in Arabidopsis identifying both distinct and similar responses
to cytokinin down- and up-regulation. J. Exp. Bot. *64*, 4193–4206. 10.1093/jxb/ert227.

82.   Cerný, M., Dycka, F., Bobál'Ová, J., and Brzobohatý, B. (2011). Early cytokinin response
proteins and phosphoproteins of Arabidopsis thaliana identified by proteome and
phosphoproteome profiling. J. Exp. Bot. *62*, 921–937. 10.1093/jxb/erq322.

83.   Miyawaki, K., Tarkowski, P., Matsumoto-Kitano, M., Kato, T., Sato, S., Tarkowska, D.,
Tabata, S., Sandberg, G., and Kakimoto, T. (2006). Roles of Arabidopsis ATP/ADP
isopentenyltransferases and tRNA isopentenyltransferases in cytokinin biosynthesis.
Proc. Natl. Acad. Sci. U. S. A. *103*, 16598–16603. 10.1073/pnas.0603522103.

84.   Ren, B., Liang, Y., Deng, Y., Chen, Q., Zhang, J., Yang, X., and Zuo, J. (2009). Genome-

1856    wide comparative analysis of type-A Arabidopsis response regulator genes by
1857    overexpression studies reveals their diverse roles and regulatory mechanisms in
1858    cytokinin signaling. Cell Res. *19*, 1178–1190. 10.1038/cr.2009.88.
1859 85. Ferreira, F.J., and Kieber, J.J. (2005). Cytokinin signaling. Curr. Opin. Plant Biol. *8*, 518–
1860    525. 10.1016/j.pbi.2005.07.013.
1861 86. Kakimoto, T. (2003). Perception and Signal Transduction of Cytokinins. Annu. Rev. Plant
1862    Biol. *54*, 605–627. 10.1146/annurev.arplant.54.031902.134802.
1863 87. Jain, M., Tyagi, A.K., and Khurana, J.P. (2006). Molecular characterization and
1864    differential expression of cytokinin-responsive type-A response regulators in rice (*Oryza*
1865    *sativa*). BMC Plant Biol. *6*, 1. 10.1186/1471-2229-6-1.
1866 88. To, J.P.C., Haberer, G., Ferreira, F.J., Deruère, J., Mason, M.G., Schaller, G.E., Alonso,
1867    J.M., Ecker, J.R., and Kieber, J.J. (2004). Type-A Arabidopsis response regulators are
1868    partially redundant negative regulators of cytokinin signaling. Plant Cell *16*, 658–671.
1869    10.1105/tpc.018978.
1870 89. Kiba, T., Yamada, H., Sato, S., Kato, T., Tabata, S., Yamashino, T., and Mizuno, T.
1871    (2003). The type-A response regulator, ARR15, acts as a negative regulator in the
1872    cytokinin-mediated signal transduction in *Arabidopsis thaliana*. Plant Cell Physiol. *44*,
1873    868–874. 10.1093/pcp/pcg108.
1874 90. Acheampong, A.K., Shanks, C., Cheng, C.Y., Eric Schaller, G., Dagdas, Y., and Kieber,
1875    J.J. (2020). EXO70D isoforms mediate selective autophagic degradation of type-A ARR
1876    proteins to regulate cytokinin sensitivity. Proc. Natl. Acad. Sci. U. S. A. *117*, 27034–
1877    27043. 10.1073/pnas.2013161117.
1878 91. To, J.P.C., Deruère, J., Maxwell, B.B., Morris, V.F., Hutchison, C.E., Ferreira, F.J.,
1879    Schaller, G.E., and Kieber, J.J. (2007). Cytokinin regulates type-A Arabidopsis response
1880    regulator activity and protein stability via two-component phosphorelay. Plant Cell *19*,
1881    3901–3914. 10.1105/tpc.107.052662.
1882 92. Bothma, J.P., Norstad, M.R., Alamos, S., and Garcia, H.G. (2018). LlamaTags: A
1883    Versatile Tool to Image Transcription Factor Dynamics in Live Embryos. Cell *173*, 1810-
1884    1822.e16. 10.1016/j.cell.2018.03.069.
1885 93. Imamura, A., Yoshino, Y., and Mizuno, T. (2001). Cellular Localization of the Signaling
1886    Components of Arabidopsis His-to-Asp Phosphorelay. Biosci. Biotechnol. Biochem. *65*,
1887    2113–2117. 10.1271/bbb.65.2113.
1888 94. Lee, D.J., Kim, S., Ha, Y.M., and Kim, J. (2008). Phosphorylation of Arabidopsis
1889    response regulator 7 (ARR7) at the putative phospho-accepting site is required for ARR7
1890    to act as a negative regulator of cytokinin signaling. Planta *227*, 577–587.
1891    10.1007/s00425-007-0640-x.
1892 95. Buechel, S., Leibfried, A., To, J.P.C., Zhao, Z., Andersen, S.U., Kieber, J.J., and
1893    Lohmann, J.U. (2010). Role of A-type Arabidopsis Response Regulators in meristem
1894    maintenance and regeneration. Eur. J. Cell Biol. *89*, 279–284.
1895    10.1016/j.ejcb.2009.11.016.
1896 96. Xie, M., Chen, H., Huang, L., O'Neil, R.C., Shokhirev, M.N., and Ecker, J.R. (2018). A B-
1897    ARR-mediated cytokinin transcriptional network directs hormone cross-regulation and
1898    shoot development. Nat. Commun. *9*, 1604. 10.1038/s41467-018-03921-6.
1899 97. Mähönen, A.P., Bishopp, A., Higuchi, M., Nieminen, K.M., Kinoshita, K., Törmäkangas,
1900    K., Ikeda, Y., Oka, A., Kakimoto, T., and Helariutta, Y. (2006). Cytokinin signaling and its
1901    inhibitor AHP6 regulate cell fate during vascular development. Science *311*, 94–98.
1902    10.1126/science.1118875.
1903 98. Kieber, J.J., and Schaller, G.E. (2018). Cytokinin signaling in plant development.
1904    Development *145*, dev149344. 10.1242/dev.149344.
1905 99. Hwang, I., Chen, H.C., and Sheen, J. (2002). Two-component signal transduction
1906    pathways in Arabidopsis. Plant Physiol. *129*, 500–515. 10.1104/pp.005504.

1907 100. Feng, J., Wang, C., Chen, Q., Chen, H., Ren, B., Li, X., and Zuo, J. (2013). S-
1908   nitrosylation of phosphotransfer proteins represses cytokinin signaling. Nat. Commun. *4*,
1909   1529. 10.1038/ncomms2541.
1910 101. Zhang, W., To, J.P.C., Cheng, C.Y., Eric Schaller, G., and Kieber, J.J. (2011). Type-A
1911   response regulators are required for proper root apical meristem function through post-
1912   transcriptional regulation of PIN auxin efflux carriers. Plant J. *68*, 1–10. 10.1111/j.1365-
1913   313X.2011.04668.x.
1914 102. Abel, S., Nguyen, M.D., and Theologis, A. (1995). The *PS-IAA4/5-like* Family of Early
1915   Auxin-inducible mRNAs in *Arabidopsis thaliana*. J. Mol. Biol. *251*, 533–549.
1916   10.1006/jmbi.1995.0454.
1917 103. Park, J.Y., Kim, H.J., and Kim, J. (2002). Mutation in domain II of IAA1 confers diverse
1918   auxin-related phenotypes and represses auxin-activated expression of Aux/IAA genes in
1919   steroid regulator-inducible system. Plant J. *32*, 669–683. 10.1046/j.1365-
1920   313X.2002.01459.x.
1921 104. Abel, S., Oeller, P.W., and Theologis, A. (1994). Early auxin-induced genes encode
1922   short-lived nuclear proteins. Proc. Natl. Acad. Sci. U. S. A. *91*, 326–330.
1923   10.1073/pnas.91.1.326.
1924 105. Zenser, N., Ellsmore, A., Leasure, C., and Callis, J. (2001). Auxin modulates the
1925   degradation rate of Aux/IAA proteins. Proc. Natl. Acad. Sci. U. S. A. *98*, 11795–11800.
1926   10.1073/pnas.211312798.
1927 106. Worley, C.K., Zenser, N., Ramos, J., Rouse, D., Leyser, O., Theologis, A., and Callis, J.
1928   (2000). Degradation of Aux/IAA proteins is essential for normal auxin signalling. Plant J.
1929   *21*, 553–562. 10.1046/j.1365-313X.2000.00703.x.
1930 107. Ramos, J.A., Zenser, N., Leyser, O., and Callis, J. (2001). Rapid degradation of
1931   auxin/indoleacetic acid proteins requires conserved amino acids of domain II and is
1932   proteasome dependent. Plant Cell *13*, 2349–2360. 10.1105/tpc.13.10.2349.
1933 108. Figueiredo, M.R.A. de, and Strader, L.C. (2022). Intrinsic and extrinsic regulators of
1934   Aux/IAA protein degradation dynamics. Trends Biochem. Sci. *47*, 865–874.
1935   10.1016/j.tibs.2022.06.004.
1936 109. Liao, C.Y., Smet, W., Brunoud, G., Yoshida, S., Vernoux, T., and Weijers, D. (2015).
1937   Reporters for sensitive and quantitative measurement of auxin response. Nat. Methods
1938   *12*, 207–210. 10.1038/nmeth.3279.
1939 110. Radhakrishnan, S.K., Lee, C.S., Young, P., Beskow, A., Chan, J.Y., and Deshaies, R.J.
1940   (2010). Transcription Factor Nrf1 Mediates the Proteasome Recovery Pathway after
1941   Proteasome Inhibition in Mammalian Cells. Mol. Cell *38*, 17–28.
1942   10.1016/j.molcel.2010.02.029.
1943 111. Gladman, N.P., Marshall, R.S., Lee, K.H., and Vierstra, R.D. (2016). The proteasome
1944   stress regulon is controlled by a pair of NAC transcription factors in arabidopsis. Plant
1945   Cell *28*, 1279–1296. 10.1105/tpc.15.01022.
1946 112. Njomen, E., and Tepe, J.J. (2019). Proteasome Activation as a New Therapeutic
1947   Approach to Target Proteotoxic Disorders. J. Med. Chem. *62*, 6469–6481.
1948   10.1021/acs.jmedchem.9b00101.
1949 113. Anderson, R.T., Bradley, T.A., and Smith, D.M. (2022). Hyperactivation of the
1950   proteasome in Caenorhabditis elegans protects against proteotoxic stress and extends
1951   lifespan. J. Biol. Chem. *298*, 102415. 10.1016/j.jbc.2022.102415.
1952 114. Kandel, R., Jung, J., and Neal, S. (2024). Proteotoxic stress and the ubiquitin proteasome
1953   system. Semin. Cell Dev. Biol. *156*, 107–120. 10.1016/j.semcdb.2023.08.002.
1954 115. Quy, P.N., Kuma, A., Pierres, P., and Mizushima, N. (2013). Proteasome-dependent
1955   activation of mammalian target of rapamycin complex 1 (mTORC1) is essential for
1956   autophagy suppression and muscle remodeling following denervation. J. Biol. Chem. *288*,
1957   1125–1134. 10.1074/jbc.M112.399949.

116. Zhao, J., Zhai, B., Gygi, S.P., and Goldberg, A.L. (2015). MTOR inhibition activates
overall protein degradation by the ubiquitin proteasome system as well as by autophagy.
Proc. Natl. Acad. Sci. U. S. A. *112*, 15790–15797. 10.1073/pnas.1521919112.
117. Zhang, Y., Nicholatos, J., Dreier, J.R., Ricoult, S.J.H., Widenmaier, S.B., Hotamisligil,
G.S., Kwiatkowski, D.J., and Manning, B.D. (2014). Coordinated regulation of protein
synthesis and degradation by mTORC1. Nature *513*, 440–443. 10.1038/nature13492.
118. Zhao, J., Garcia, G.A., and Goldberg, A.L. (2016). Control of proteasomal proteolysis by
mTOR. Nature *529*, E1–E2. 10.1038/nature16472.
119. Zhang, Y., and Manning, B.D. (2016). Zhang & Manning reply. Nature *529*, E2–E3.
10.1038/nature16473.
120. Rousseau, A., and Bertolotti, A. (2016). An evolutionarily conserved pathway controls
proteasome homeostasis. Nature *536*, 184–189. 10.1038/nature18943.
121. Mills, E.W., and Green, R. (2017). Ribosomopathies: There's strength in numbers.
Science *358*, eaan2755. 10.1126/science.aan2755.
122. Warren, A.J. (2018). Molecular basis of the human ribosomopathy Shwachman-Diamond
syndrome. Adv. Biol. Regul. *67*, 109–127. 10.1016/j.jbior.2017.09.002.
123. Hawer, H., Mendelsohn, B.A., Mayer, K., Kung, A., Malhotra, A., Tuupanen, S., Schleit,
J., Brinkmann, U., and Schaffrath, R. (2020). Diphthamide-deficiency syndrome: a novel
human developmental disorder and ribosomopathy. Eur. J. Hum. Genet. *28*, 1497–1508.
10.1038/s41431-020-0668-y.
124. Armistead, J., Patel, N., Wu, X., Hemming, R., Chowdhury, B., Basra, G.S., Del Bigio,
M.R., Ding, H., and Triggs-Raine, B. (2015). Growth arrest in the ribosomopathy, bowen-
conradi syndrome, is due to dramatically reduced cell proliferation and a defect in mitotic
progression. Biochim. Biophys. Acta - Mol. Basis Dis. *1852*, 1029–1037.
10.1016/j.bbadis.2015.02.007.
125. Zhou, C., Zang, D., Jin, Y., Wu, H., Liu, Z., Du, J., and Zhang, J. (2011). Mutation in
ribosomal protein L21 underlies hereditary hypotrichosis simplex. Hum. Mutat. *32*, 710–
714. 10.1002/humu.21503.
126. Dixon, M.J. (1995). Treacher Collins syndrome. J. Med. Genet. *32*, 806–808.
10.1136/jmg.32.10.806.
127. Gazda, H.T., Sheen, M.R., Vlachos, A., Choesmel, V., O'Donohue, M.F., Schneider, H.,
Darras, N., Hasman, C., Sieff, C.A., Newburger, P.E., et al. (2008). Ribosomal Protein L5
and L11 Mutations Are Associated with Cleft Palate and Abnormal Thumbs in Diamond-
Blackfan Anemia Patients. Am. J. Hum. Genet. *83*, 769–780. 10.1016/j.ajhg.2008.11.004.
128. Oliver, E.R., Saunders, T.L., Tarlé, S.A., and Glaser, T. (2004). Ribosomal protein L24
defect in belly spot and tail (*Bst*), a mouse *Minute*. Development *131*, 3907–3920.
10.1242/dev.01268.
129. McGowan, K.A., Li, J.Z., Park, C.Y., Beaudry, V., Tabor, H.K., Sabnis, A.J., Zhang, W.,
Fuchs, H., De Angelis, M.H., Myers, R.M., et al. (2008). Ribosomal mutations cause p53-
mediated dark skin and pleiotropic effects. Nat. Genet. *40*, 963–970. 10.1038/ng.188.
130. Oristian, D.S., Slooíman, L.G., Zhou, X., Wang, L., Farach-Carson, M.C., and Kirn-
Safran, C.B. (2009). Ribosomal protein L29/HIP deficiency delays osteogenesis and
increases fragility of adult bone in mice. J. Orthop. Res. *27*, 28–35. 10.1002/jor.20706.
131. Uechi, T., Nakajima, Y., Nakao, A., Torihara, H., Chakraborty, A., Inoue, K., and
Kenmochi, N. (2006). Ribosomal protein gene knockdown causes developmental defects
in zebrafish. PLoS One *1*, e37. 10.1371/journal.pone.0000037.
132. Marygold, S.J., Coelho, C.M.A., and Leevers, S.J. (2005). Genetic analysis of RpL38 and
RpL5, two minute genes located in the centric heterochromatin of chromosome 2 of
Drosophila melanogaster. Genetics *169*, 683–695. 10.1534/genetics.104.034124.
133. Lee, C.C., Tsai, Y.T., Kao, C.W., Lee, L.W., Lai, H.J., Ma, T.H., Chang, Y.S., Yeh, N.H.,
and Lo, S.J. (2014). Mutation of a Nopp140 gene *dao-5* alters rDNA transcription and

increases germ cell apoptosis in *C. elegans*. Cell Death Dis. *5*, e1158.
10.1038/cddis.2014.114.

134. Ito, T., Kim, G.T., and Shinozaki, K. (2000). Disruption of an Arabidopsis cytoplasmic
ribosomal protein S13-homologous gene by transposon-mediated mutagenesis causes
aberrant growth and development. Plant J. *22*, 257–264. 10.1046/j.1365-
313X.2000.00728.x.

135. Horiguchi, G., Mollá-Morales, A., Pérez-Pérez, J.M., Kojima, K., Robles, P., Ponce, M.R.,
Micol, J.L., and Tsukaya, H. (2011). Differential contributions of ribosomal protein genes
to *Arabidopsis thaliana* leaf development. Plant J. *65*, 724–736. 10.1111/j.1365-
313X.2010.04457.x.

136. Zhao, H., Lü, S., Li, R., Chen, T., Zhang, H., Cui, P., Ding, F., Liu, P., Wang, G., Xia, Y.,
et al. (2015). The Arabidopsis gene *DIG6* encodes a large 60S subunit nuclear export
GTPase 1 that is involved in ribosome biogenesis and affects multiple auxin-regulated
development processes. J. Exp. Bot. *66*, 6863–6875. 10.1093/jxb/erv391.

137. Takagi, M., Absalon, M.J., McLure, K.G., and Kastan, M.B. (2005). Regulation of p53
translation and induction after DNA damage by ribosomal protein L26 and nucleolin. Cell
123, 49–63. 10.1016/j.cell.2005.07.034.

138. Deisenroth, C., and Zhang, Y. (2010). Ribosome biogenesis surveillance: Probing the
ribosomal protein-Mdm2-p53 pathway. Oncogene *29*, 4253–4260.
10.1038/onc.2010.189.

139. Warner, J.R., and McIntosh, K.B. (2009). How Common Are Extraribosomal Functions of
Ribosomal Proteins? Mol. Cell *34*, 3–11. 10.1016/j.molcel.2009.03.006.

140. Dutt, S., Narla, A., Lin, K., Mullally, A., Abayasekara, N., Megerdichian, C., Wilson, F.H.,
Currie, T., Khanna-Gupta, A., Berliner, N., et al. (2011). Haploinsufficiency for ribosomal
protein genes causes selective activation of p53 in human erythroid progenitor cells.
Blood *117*, 2567–2576. 10.1182/blood-2010-07-295238.

141. Danilova, N., Sakamoto, K.M., and Lin, S. (2008). Ribosomal protein S19 deficiency in
zebrafish leads to developmental abnormalities and defective erythropoiesis through
activation of p53 protein family. Blood *112*, 5228–5237. 10.1182/blood-2008-01-132290.

142. Girardi, T., Vereecke, S., Sulima, S.O., Khan, Y., Fancello, L., Briggs, J.W., Schwab, C.,
De Beeck, J.O., Verbeeck, J., Royaert, J., et al. (2018). The T-cell leukemia-associated
ribosomal RPL10 R98S mutation enhances JAK-STAT signaling. Leukemia *32*, 809–819.
10.1038/leu.2017.225.

143. Ludwig, L.S., Gazda, H.T., Eng, J.C., Eichhorn, S.W., Thiru, P., Ghazvinian, R., George,
T.I., Gotlib, J.R., Beggs, A.H., Sieff, C.A., et al. (2014). Altered translation of GATA1 in
Diamond-Blackfan anemia. Nat. Med. *20*, 748–753. 10.1038/nm.3557.

144. Khajuria, R.K., Munschauer, M., Ulirsch, J.C., Fiorini, C., Ludwig, L.S., McFarland, S.K.,
Abdulhay, N.J., Specht, H., Keshishian, H., Mani, D.R., et al. (2018). Ribosome Levels
Selectively Regulate Translation and Lineage Commitment in Human Hematopoiesis.
Cell *173*, 90-103.e19. 10.1016/j.cell.2018.02.036.

145. In, K., Zaini, M.A., Müller, C., Warren, A.J., Von Lindern, M., and Calkhoven, C.F. (2016).
Shwachman-Bodian-Diamond syndrome (SBDS) protein deficiency impairs translation re-
initiation from C/EBPα and C/EBPβ mRNAs. Nucleic Acids Res. *44*, 4134–4146.
10.1093/nar/gkw005.

146. Kondrashov, N., Pusic, A., Stumpf, C.R., Shimizu, K., Hsieh, A.C., Xue, S., Ishijima, J.,
Shiroishi, T., and Barna, M. (2011). Ribosome-mediated specificity in Hox mRNA
translation and vertebrate tissue patterning. Cell *145*, 383–397.
10.1016/j.cell.2011.03.028.

147. Deliu, L.P., Turingan, M., Jadir, D., Lee, B., Ghosh, A., and Grewal, S.S. (2022).
Serotonergic neuron ribosomal proteins regulate the neuroendocrine control of
Drosophila development. PLoS Genet. *18*, e1010371. 10.1371/journal.pgen.1010371.

148. Adams, K.W., and Cooper, G.M. (2007). Rapid turnover of Mcl-1 couples translation to
cell survival and apoptosis. J. Biol. Chem. *282*, 6192–6200. 10.1074/jbc.M610643200.
149. Truckenbrodt, S., Viplav, A., Jähne, S., Vogts, A., Denker, A., Wildhagen, H., Fornasiero,
E.F., and Rizzoli, S.O. (2018). Newly produced synaptic vesicle proteins are preferentially
used in synaptic transmission. EMBO J. *37*, e98044. 10.15252/embj.201798044.
150. D'Agostino, I.B., Deruère, J., and Kieber, J.J. (2000). Characterization of the response of
the Arabidopsis response regulator gene family to cytokinin. Plant Physiol. *124*, 1706–
1717. 10.1104/pp.124.4.1706.
151. Bishopp, A., Help, H., El-Showk, S., Weijers, D., Scheres, B., Friml, J., Benková, E.,
Mähönen, A.P., and Helariutta, Y. (2011). A mutually inhibitory interaction between auxin
and cytokinin specifies vascular pattern in roots. Curr. Biol. *21*, 917–926.
10.1016/j.cub.2011.04.017.
152. Sud, M., Fahy, E., Cotter, D., Azam, K., Vadivelu, I., Burant, C., Edison, A., Fiehn, O.,
Higashi, R., Nair, K.S., et al. (2016). Metabolomics Workbench: An international
repository for metabolomics data and metadata, metabolite standards, protocols, tutorials
and training, and analysis tools. Nucleic Acids Res. *44*, D463–D470.
10.1093/nar/gkv1042.
153. Mähönen, A.P., Bonke, M., Kauppinen, L., Riikonen, M., Benfey, P.N., and Helariutta, Y.
(2000). A novel two-component hybrid molecule regulates vascular morphogenesis of the
Arabidopsis root. Genes Dev. *14*, 2938–2943. 10.1101/gad.189200.
154. Sotta, N., Shantikumar, L., Sakamoto, T., Matsunaga, S., and Fujiwara, T. (2016). TPR5
is involved in directional cell division and is essential for the maintenance of meristem cell
organization in *Arabidopsis thaliana*. J. Exp. Bot. *67*, 2401–2411. 10.1093/jxb/erw043.
155. Heisler, M.G., Ohno, C., Das, P., Sieber, P., Reddy, G. V., Long, J.A., and Meyerowitz,
E.M. (2005). Patterns of auxin transport and gene expression during primordium
development revealed by live imaging of the Arabidopsis inflorescence meristem. Curr.
Biol. *15*, 1899–1911. 10.1016/j.cub.2005.09.052.
156. Müller, B., and Sheen, J. (2008). Cytokinin and auxin interaction in root stem-cell
specification during early embryogenesis. Nature *453*, 1094–1097. 10.1038/nature06943.
157. Brackmann, K., Qi, J., Gebert, M., Jouannet, V., Schlamp, T., Grünwald, K., Wallner,
E.S., Novikova, D.D., Levitsky, V.G., Agustí, J., et al. (2018). Spatial specificity of auxin
responses coordinates wood formation. Nat. Commun. *9*, 875. 10.1038/s41467-018-
03256-2.
158. Argyros, R.D., Mathews, D.E., Chiang, Y.H., Palmer, C.M., Thibault, D.M., Etheridge, N.,
Argyros, D.A., Mason, M.G., Kieber, J.J., and Schallera, G.E. (2008). Type B response
regulators of Arabidopsis play key roles in cytokinin signaling and plant development.
Plant Cell *20*, 2102–2116. 10.1105/tpc.108.059584.
159. Rademacher, E.H., Möller, B., Lokerse, A.S., Llavata-Peris, C.I., Van Den Berg, W., and
Weijers, D. (2011). A cellular expression map of the *Arabidopsis AUXIN RESPONSE
FACTOR* gene family. Plant J. *68*, 597–606. 10.1111/j.1365-313X.2011.04710.x.
160. Saerens, D., Pellis, M., Loris, R., Pardon, E., Dumoulin, M., Matagne, A., Wyns, L.,
Muyldermans, S., and Conrath, K. (2005). Identification of a universal VHH framework to
graft non-canonical antigen-binding loops of camel single-domain antibodies. J. Mol. Biol.
352, 597–607. 10.1016/j.jmb.2005.07.038.
161. Fujii, Y., and Kodama, Y. (2015). *In planta* comparative analysis of improved green
fluorescent proteins with reference to fluorescence intensity and bimolecular fluorescence
complementation ability. Plant Biotechnol. *32*, 81–87.
10.5511/plantbiotechnology.15.0120a.
162. Alamos, S., Reimer, A., Niyogi, K.K., and Garcia, H.G. (2021). Quantitative imaging of
RNA polymerase II activity in plants reveals the single-cell basis of tissue-wide
transcriptional dynamics. Nat. Plants *7*, 1037–1049. 10.1038/s41477-021-00976-0.

163. Alexa, A., Rahnenführer, J., and Lengauer, T. (2006). Improved scoring of functional groups from gene expression data by decorrelating GO graph structure. Bioinformatics *22*, 1600–1607. 10.1093/bioinformatics/btl140.

164. Mustroph, A., Juntawong, P., and Bailey-Serres, J. (2009). Isolation of Plant Polysomal mRNA by Differential Centrifugation and Ribosome Immunopurification Methods. In Methods in molecular biology (vol. 553) (Humana Press, a part of Springer Science+Business Media, LLC), pp. 109–126. 10.1007/978-1-60327-563-7_6.

165. Gao, X., Wan, J., Liu, B., Ma, M., Shen, B., and Qian, S.B. (2015). Quantitative profiling of initiating ribosomes *in vivo*. Nat. Methods *12*, 147–153. 10.1038/nmeth.3208.

166. Busche, M., Regina Scarpin, M., Hnasko, R., and Brunkard, J.O. (2021). TOR coordinates nucleotide availability with ribosome biogenesis in plants. Plant Cell *33*, 1615–1632. 10.1093/plcell/koab043.

167. Schindelin, J., Rueden, C.T., Hiner, M.C., and Eliceiri, K.W. (2015). The ImageJ ecosystem: An open platform for biomedical image analysis. Mol. Reprod. Dev. *82*, 518–529. 10.1002/mrd.22489.

168. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: An open-source platform for biological-image analysis. Nat. Methods *9*, 676–682. 10.1038/nmeth.2019.

169. de Reuille, P.B., Routier-Kierzkowska, A.L., Kierzkowski, D., Bassel, G.W., Schüpbach, T., Tauriello, G., Bajpai, N., Strauss, S., Weber, A., Kiss, A., et al. (2015). MorphoGraphX: A platform for quantifying morphogenesis in 4D. eLife *4*, e05864. 10.7554/eLife.05864.

170. Schäfer, M., Brütting, C., Baldwin, I.T., and Kallenbach, M. (2016). High-throughput quantification of more than 100 primary- and secondary-metabolites, and phytohormones by a single solid-phase extraction based sample preparation with analysis by UHPLC-HESI-MS/MS. Plant Methods *12*, 30. 10.1186/s13007-016-0130-x.

171. Kojima, M., Kamada-Nobusada, T., Komatsu, H., Takei, K., Kuroha, T., Mizutani, M., Ashikari, M., Ueguchi-Tanaka, M., Matsuoka, M., Suzuki, K., et al. (2009). Highly sensitive and high-throughput analysis of plant hormones using ms-probe modification and liquid chromatographytandem mass spectrometry: An application for hormone profiling in oryza sativa. Plant Cell Physiol. *50*, 1201–1214. 10.1093/pcp/pcp057.

172. Novák, O., Antoniadi, I., and Ljung, K. (2017). High-resolution cell-type specific analysis of cytokinins in sorted root cell populations of *Arabidopsis thaliana*. Methods Mol. Biol. *1497*, 231–248. 10.1007/978-1-4939-6469-7_1.

173. R Core Team (2021). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

174. Wickham, H. (2009). ggplot2: Elegant Graphics for Data Analysis R. Gentleman, K. Hornik, and G. Parmigiani, eds. (Springer) 10.1007/978-0-387-98141-3.

175. Perez-Riverol, Y., Bai, J., Bandla, C., García-Seisdedos, D., Hewapathirana, S., Kamatchinathan, S., Kundu, D.J., Prakash, A., Frericks-Zipper, A., Eisenacher, M., et al. (2022). The PRIDE database resources in 2022: A hub for mass spectrometry-based proteomics evidences. Nucleic Acids Res. *50*, D543–D552. 10.1093/nar/gkab1038.

Figure 1. *drmy1* has reduced ribosome abundance, translation rate, and TOR activity.

(A) Top row, stage 12 buds of WT (left) and *drmy1* (right) viewed from the side. Asterisk shows the gap between sepals with petals and carpels exposed. Middle row, stage 12 buds of WT (left) and *drmy1* (right) viewed from the top. Arrowheads point to sepals. Note that the *drmy1* bud has 5 sepals of unequal size and unevenly spaced, exposing the stamens and carpels. Bottom row, stage 5 buds of WT (left) and *drmy1* (right) containing *35S::mCitrine-RCI2A* (plasma membrane marker). Arrowheads point to sepal primordia. Note that the *drmy1* bud has 5 sepal primordia of different sizes. Scale bars are 0.5 mm for stage 12 bud images and 25 μm for stage 5 bud images.

(B) Gene ontology (GO) enrichment of downregulated genes in *drmy1* compared to WT, in the *ap1 cal AP1-GR* background. Shown are the top 8 GO terms and their enrichment p-values. A complete list can be found in Supplemental Dataset 1. Arrowheads highlight terms related to ribosome biogenesis or translation.

(C) Polysomal profiles of WT (blue) and *drmy1* (red) in the *ap1 cal AP1-GR* background, representative of 3 biological replicates each. Additional replicates are in Supplemental Dataset 2. M, monosomes. P, polysomes.

(D) Puromycin labeling of WT vs *drmy1*. Left, WT and *drmy1* seedlings. From left to right: WT pre-treated with CHX, two biological replicates of WT pre-treated with mock, and two biological replicates of *drmy1* pre-treated with mock. All groups were then treated with puromycin. For seedlings to match in size, WT seedlings were 8 days old and *drmy1* seedlings were 10 days old.

Right, WT and *drmy1* inflorescences of induced *ap1 cal AP1-GR* background. From left to right:
WT co-treated with puromycin and CHX, three biological replicates of WT treated with puromycin,
and three biological replicates of *drmy1* treated with puromycin. In both experiments, RuBisCO
large subunit on Ponceau S-stained membrane is shown as a loading control (bottom). Ratio
between puromycin and Ponceau S signals, normalized by the mean of WT, is shown on the
bottom (p-values are from two-sided Student's t-test).
(E) Gene ontology (GO) enrichment of differentially accumulated proteins in *drmy1* compared to
WT, in the *ap1 cal AP1-GR* background. Shown are the top 8 GO terms and their enrichment p-
values. A complete list can be found in Supplemental Dataset 1. Arrowheads highlight terms
related to ribosome biogenesis or translation.
(F-G) Coherent alteration of gene expression by *drmy1* and AZD-8055 TOR inhibitor treatment.
(F) Scatterplot of RNA log2 fold change in *drmy1* vs WT (x-axis), and WT+AZD vs WT+Mock (y-
axis), in 7-day-old seedlings. Genes are color-coded based on the following categories: genes in
"Structural constituents of the ribosome" (GO:0003735) and its offspring terms (magenta); all
other genes in "Translation" (GO:0006412) and its offspring terms (orange); all other genes (gray).
Blue line shows a linear regression of all points (R^2 = 0.1446, p < $2.2{\times}10^{-16}$). **(G)** Of the 466 genes
that are differentially expressed under both conditions, 439 (94%) are coherently altered by AZD-
8055 treatment and the *drmy1* mutation.
(H-I) Phosphorylation of the direct TOR substrate, S6K-pT449, in WT and *drmy1*. **(H)** A
representative blot. Top, S6K-pT449. Middle, total S6K protein. Bottom, Ponceau S staining. Ratio
between S6K-pT449 signal and Ponceau S signal is shown above the blots. **(I)** Ratio between
S6K-pT449 and Ponceau S signals normalized by WT, quantified across in three experiments,
shows that TOR activity decreased by half in *drmy1*. (mean ± SD; *, p<0.05).
45

Figure 2. Defects in TOR activity, ribosome, and translation disrupt robust sepal initiation.
(A-G) Representative images of stage 5 buds in WT (A), *drmy1* (B), *ul4z* (C), *ul4y* (D), *ul18z* (E), *lst8-1-1* (F), and *spaghetti-1* (G). Tissue morphology is visualized by either propidium iodide (a cell wall-staining dye) or a plasma membrane marker. Arrowheads indicate sepal primordia that are variable in number, position, and size. Note that *ul4z* flowers always develop four sepal primordia, although of different sizes; *lst8-1-1* occasionally (4/41, 9.8%) develops buds with more than four sepal primordia.
(H) Quantification of sepal primordium number, comparing *drmy1* (n = 67 buds), *ul4z* (n = 52 buds), *ul4y* (n = 53 buds), *ul18z* (n = 52 buds), *lst8-1-1* (n = 41 buds), and *spaghetti-1* (n = 84 buds) with WT (n = 51 buds). Asterisks indicate statistically significant (p < 0.05) differences from WT in Fisher's contingency table tests.
(I) Illustration of robust versus variable positioning of sepal primordia. Primordia are considered robustly positioned if they are evenly distributed around the edge of the bud. Within each bud, angles between adjacent primordia with respect to the center of the bud are measured, and coefficient of variation (CV) is calculated. A bud with robustly positioned primordia would have similar angular values and a low CV value. A bud with variably positioned primordia would have very different angular values and a high CV value.

(J) Quantification of variability in primordium positioning (CV) in the same buds as in (H), following

illustration in (I). Asterisks indicate statistically significant ($p < 0.05$) differences from WT in

Wilcoxon's rank sum tests.

(K) Representative images of buds from *in vitro*-cultured WT inflorescences treated with mock or

2 µM CHX for 9-10 days. Arrowheads indicate sepal primordia that are variable in number,

position, and size.

(L) Representative images of buds from WT plants treated with mock or 2 nmol Torin2 for 15 days.

Arrowheads indicate sepal primordia that are variable in number, position, and size.

(M-N) Quantification of sepal primordium number (M) and positional variability (N) similar to (H,J),

comparing CHX-treated (n = 31 buds), CHX-mock (n = 42 buds), Torin2-treated (n = 51 buds)

and Torin2-mock buds (n = 56 buds). Scale bars in all micrographs, 25 µm.

Figure 3. TOR and ribosomal defects cause variability in the timing of sepal initiation.

(A-C) 6h-interval live imaging of the sepal initiation process in WT (A) and *ul4y* (B), which is quantified in (C). n = 48 buds for WT; n = 40 buds for *ul4y*.

(D-F) 6h-interval live imaging of the sepal initiation process in buds from WT plants treated with mock or 2 nmol Torin2 twice a day for 15 days, which is quantified in (F). n = 31 buds for mock; n = 15 buds for Torin2.

In **(A,B,D,E)**, top rows show the *35S::mCitrine-RCI2A* membrane marker, and bottom rows show Gaussian curvature heatmaps of the same image stacks. Asterisks indicate sepal initiation events, defined as a dark red band (primordium with positive curvature) separated from the floral meristem by a dark blue band (boundary with negative curvature) in the heatmap. Scale bars, 25 μm.

In **(C,F)**, the amount of time between outer and inner sepal initiation (left) and between outer and lateral sepal initiation (right) were calculated for each bud. Bar plot shows mean ± SD which is

90 also shown on top of each plot. Asterisks indicate statistically significant differences ($p < 0.05$) in

91 Wilcoxon's rank sum test (for mean) or Levene's test (for SD).

Figure 4. Inhibition of TOR activity and translation cause variability in auxin and cytokinin signaling.

(A-D) The ribosomal mutant *ul4y* loses robustness in auxin and cytokinin signaling. (A) Representative images of late stage 2 buds of WT, *drmy1*, and *ul4y*, showing the auxin signaling reporter *DR5::3xVENUS-N7* in yellow, the cytokinin signaling reporter *TCS::GFP* in cyan, and both merged with Chlorophyll (in WT) or *UBQ10::mCherry-RCI2A* (in *drmy1* and *ul4y*) in magenta. (B) Quantification of TCS intensity (integrated density divided by area) from maximum intensity projection images, normalized to mean of WT. Shown are mean ± SD. Asterisks show statistically significant differences from WT in two-tailed Student's t-tests (*drmy1*, p = 2.1×10^{-6}; *ul4y*, p = 3.4×10^{-5}). (C) Circular histogram of DR5 signal distribution. Each bud was divided into 360 sectors of 1° each. Within each sector, DR5 signal measured in pixel intensity units (0-255 range) was summed. This sum was plotted along the x-axis starting from the sector at 1:30 position (between the incipient outer sepal and incipient right sepal) going counterclockwise. I.e., in WT, the outer

sepal is near 45°, the inner sepal near 225°, and the lateral sepals near 45° and 135° (vertical

dotted lines). The mean was plotted as a solid line, and mean ± SD was plotted as a shaded area.

(D) Circular histogram of TCS signal distribution. Sample size for (A-D): WT, n = 12 buds; *drmy1*,

n = 15 buds; *ul4y*, n = 10 buds.

(E-H) 3 days of translation inhibition causes increased and diffuse cytokinin signaling, and diffuse

auxin signaling. (E) Representative images of late stage 2 WT buds treated *in vitro* with mock or

2 µM CHX for 3 days. Shown are *DR5::3xVENUS-N7* in yellow, *TCS::GFP* in cyan, and both

merged with Chlorophyll in magenta. (F) Quantification of TCS intensity from maximum intensity

projection images, normalized to mean of WT mock day 3. Shown are mean ± SD. Asterisk shows

statistically significant difference in a two-tailed Student's t-test (p = 2.0×10^{-4}). (G) Circular

histogram of DR5 signal distribution. (H) Circular histogram of TCS signal distribution. Sample

size for (E-H): WT mock day 3, n = 10 buds; WT CHX day 3, n = 12 buds.

(I-L) 6 days of TOR or translation inhibition causes increased and diffuse cytokinin signaling, and

randomly positioned auxin signaling maxima. (I) Representative images of late stage 2 WT buds

treated *in vitro* with mock, 2 µM CHX, or 2 µM AZD for 6 days. Shown are *DR5::3xVENUS-N7* in

yellow, *TCS::GFP* in cyan, and both merged with Chlorophyll in magenta. Arrowheads point to

randomly positioned auxin maxima. (J) Quantification of TCS intensity from maximum intensity

projection images, normalized to mean of WT mock day 6. Shown are mean ± SD. Asterisks show

statistically significant differences from mock in two-tailed Student's t-tests (CHX, p = 1.0×10^{-3};

AZD, p = 1.2×10^{-4}). (K) Circular histogram of DR5 signal distribution. (L) Circular histogram of

TCS signal distribution. Sample size for (I-L): WT mock day 6, n = 12 buds; WT CHX day 6, n =

11 buds; WT AZD day 6, n = 10 buds. Scale bars in all micrographs, 25 µm.

Figure 5. Cytokinin signaling is required for increased variability in auxin signaling and sepal initiation under translation inhibition.

(A-D) Cytokinin treatment makes auxin signaling diffuse. Shown are late stage 2 WT buds under mock (A,B) or 5 µM cytokinin (BAP) treatment (C,D) for 4 days. (A,C) Auxin signaling reporter

DR5 in yellow, and *DR5* merged with Chlorophyll in magenta. (B,D) Circular histograms of the
DR5 signal, showing mean (solid line) and mean ± SD (shaded area). Arrows point to DR5 signal
in variable positions. Sample size: WT Mock n = 10, WT BAP n = 10. Also see Zhu et al. (2020),
in this reference see Extended Data Figure 7e.
(E-G) Cytokinin signaling is required for variable sepal initiation in *drmy1*. (E) Stage 5 buds. Sepal
primordia in *drmy1* are variable (arrowheads), which does not occur in *drmy1 arr1,10,12* and
drmy1 wol mutants. (F,G) Quantification of sepal primordium number (F) and positional variability
(G), comparing WT (n = 58) with *drmy1* (n = 31), *arr1,10,12* (n = 24) with *drmy1 arr1,10,12* (n =
20), and *wol* (n = 36) with *drmy1 wol* (n = 39). Asterisks indicate statistically significant ($p < 0.05$)
differences in Fisher's contingency table tests (F) and Wilcoxon's rank sum tests (G) respectively.
(H-K) Cytokinin signaling is required for variable patterning of auxin signaling in *drmy1*. Shown
are late stage 2 buds of WT vs *drmy1* (H,I), and *arr1,10,12* vs *drmy1 arr1,10,12* (J,K). (H,J) Auxin
signaling reporter *DR5* in yellow, and *DR5* merged with propidium iodide in magenta. Arrows point
to diffuse DR5 signal in variable positions of the *drmy1* bud. Arrowheads show four robust DR5
maxima in the *drmy1 arr1,10,12* bud. (I,K) Circular histograms of the *DR5* signal, showing mean
(solid line) and mean ± SD (shaded area). For ease of visualization, circular histograms of *drmy1*
and *drmy1 arr1,10,12* between 90 and 360 degrees are enlarged and shown as insets (y-axis
range 0-0.4). Note the presence of DR5 signal in inter-sepal regions in *drmy1* (black arrow) which
is largely suppressed in *drmy1 arr1,10,12*. Sample size: WT n = 19, *drmy1* n = 16, *arr1,10,12* n =
13, *drmy1 arr1,10,12* n = 9.
(L-N) Cytokinin signaling is required for variable sepal initiation under translation inhibition. (L)
Stage 6 buds of WT, *arr1,10,12*, and *wol*, treated with Mock or 2 μM CHX for 10 days. WT initiates
sepal primordia at variable positions when treated with CHX (arrowheads), which does not occur
in *arr1,10,12* and *wol*. (M,N) Quantification of sepal primordium number (M) and positional
variability (N), comparing mock and CHX within each genotype. Sample size: WT Mock n = 29,
WT CHX n = 19, *arr1,10,12* Mock n = 18, *arr1,10,12* CHX n = 19, *wol* Mock n = 15, *wol* CHX n =
19. Asterisks indicate statistically significant ($p < 0.05$) differences in Fisher's contingency table
tests (M) and Wilcoxon's rank sum tests (N) respectively.
(O-R) Cytokinin signaling is required for diffuse auxin signaling under translation inhibition. Shown
are late stage 2 buds of WT (O,P) and *arr1,10,12* (Q,R), treated with Mock or 2 μM CHX for 3
days. (O,Q) Auxin signaling reporter *DR5* in yellow, and *DR5* merged with Chlorophyll in magenta.
Arrows point to diffuse DR5 signal in variable positions in CHX-treated WT. Arrowheads show
four robust DR5 maxima in CHX-treated *arr1,10,12*. (P,R) Circular histograms of the *DR5* signal,
showing mean (solid line) and mean ± SD (shaded area). Sample size: WT Mock n = 17, WT CHX
n = 18, *arr1,10,12* Mock n = 7, *arr1,10,12* CHX n = 7. Scale bars in all micrographs, 25 μm.
170

Figure 6. Upregulation of cytokinin signaling is required to maintain translation and fitness in *drmy1*.

(A) Puromycin labeling of WT seedlings with 4 h CHX pre-treatment (control), and three biological replicates each of WT and *arr1 35S::ARR1* seedlings with 4 h mock pre-treatment. All seedlings are 14 days old. RuBisCO large subunit in Ponceau S-stained membrane is shown as a loading control. Signal ratio between puromycin and Ponceau S, normalized to mean of WT, is show on the bottom. P-value is from a two-sided Student's t-test. Also see Karunadasa et al. (2020).

(B,C) Puromycin labeling of WT seedlings with 4 h CHX pre-treatment (control), and two biological replicates of WT, *drmy1*, *wol*, and *drmy1 wol* seedlings with 4 h mock pre-treatment. Seedlings are 8 days old in (B) and 14 days old in (C). RuBisCO large subunit in Ponceau S-stained membrane is shown as a loading control. Letters show compact letter display of a Tukey's all-pair comparison in a one-way ANOVA model.

(D) Representative 14 days old seedling images of WT, *drmy1*, *wol*, and *drmy1 wol* used in (C). Notice that *drmy1 wol* is very small and pale. Scale bars, 5 mm.

(E) Representative aerial part images of 42 days old plants of WT, *drmy1*, *wol*, and *drmy1 wol*. Inset shows the zoomed-in *drmy1 wol* plant, which has a tiny rosette and a short inflorescence. Scale bars, 5 cm. See also Figure S6F.

(F) Representative aerial part images of 74 days old plants of WT, *drmy1*, *arr1,10,12*, and *drmy1 arr1,10,12*. Inset shows the zoomed-in *drmy1 arr1,10,12* plant, which has pale leaves accumulating anthocyanin and a short inflorescence. Scale bars, 5 cm. See also Figure S6E.

(G) Dissected siliques of *arr1,10,12* (left) and *drmy1 arr1,10,12* (right) showing developing seeds. Notice that while *arr1,10,12* occasionally have aborted seeds, all seeds in the *drmy1 arr1,10,12* silique were aborted. Scale bars, 0.2 mm.

195
196

Figure 7. ARR7 and AHP6 protein levels are reduced upon inhibition of TOR and translation.

(A) The hypothesis. During hormone patterning prior to sepal initiation in the WT floral meristem, A-type ARR and AHP6 proteins are rapidly produced to dampen cytokinin signaling to a normal

level. In *drmy1*, reduced protein synthesis causes reduced levels of these cytokinin signaling
inhibitor proteins, resulting in an upregulation of cytokinin signaling.
(B) Illustration of Llama Tag. Plants were co-transformed with *ARR7-llama* (*pARR7::ARR7-linker-*
llama-ARR7ter) and *GFP-nes* (*pUBQ10::sfGFP-nes-UBQ3ter*). Without ARR7-llama, GFP
localizes to the cytosol due to the nuclear export sequence (nes). ARR7-llama is produced in the
cytoplasm, C, and translocates into the nucleus, N. The Llama Tag on ARR7-llama binds to GFP
and drags GFP into the nucleus (note that from our observation it is excluded from the nucleolus,
NL). Thus, at low ARR7-llama levels, GFP signal is mainly in the cytoplasm. At intermediate
ARR7-llama levels, GFP is at comparable levels between the cytoplasm and the nucleus, and no
clear pattern can be seen. At high ARR7-llama levels, GFP is mainly seen in the nucleus.
(C) A *GFP-nes* bud showing localization of the GFP signal to the cytoplasm.
(D,E) GFP images of buds from two independent transgenic lines of *ARR7-llama GFP-nes*, 7-4
(D) and 7-6 (E), of WT (top) vs *drmy1* (bottom) genotypes. Images are representative of n = 17
(line 7-4, WT), n = 40 (line 7-4, *drmy1*), n = 9 (line 7-6, WT), and n = 6 (line 7-6, *drmy1*) buds.
Note that GFP is more cytoplasm-localized in *drmy1* than WT, indicating reduced ARR7-llama
protein level.
(F) GFP images of WT *ARR7-llama GFP-nes* buds treated with mock (top) or 2 μM CHX (bottom)
for 24 hours. The mock image is representative of n = 20 buds from three independent lines. The
CHX image is representative of n = 19 buds from these same lines.
(G) GFP images of WT *ARR7-llama GFP-nes* buds treated with mock (top) or 2 μM AZD-8055
(bottom) for 72 hours. The mock image is representative of n = 13 buds from two independent
lines. The AZD-8055 image is representative of n = 11 buds from these same lines. For (C-G),
each image was brightened to reveal subcellular localization patterns of GFP. A square region
taken from the image containing 5-10 cells is enlarged and shown on the top right. Within the
square, GFP intensity was quantified along the dotted line and plotted on the bottom right. X-axis,
pixels (range 0-238). Y-axis, GFP intensity (smoothened by taking the average intensity of 11-
pixel neighborhoods; range 90-210 in gray value).
(H-I) Response of the AHP6 protein reporter (H) and transcriptional reporter (I) to mock, CHX,
and AZD-8055 treatments for 72 hours. For (H), images are representative of n = 29 (mock), n =
29 (CHX), and n = 34 (AZD-8055) buds in three experiments. For (I), images are representative
of n = 11 (mock), n = 9 (CHX), and n = 12 (AZD-8055) buds in two experiments.
(J-Q) Reduction of A-type ARR and AHP6 protein levels contribute to the variability in sepal
initiation. (J-O) Stage 5-6 buds of indicated genotype stained with propidium iodide. Arrowheads
indicate sepal primordia that are variable in number, position, and/or size. Note that the
arr3,4,5,6,7,8,9,15 bud has an inner sepal that is slightly smaller than its outer sepal and
positioned slightly right-skewed (K). The *ahp6* bud develops five sepal primordia of variable sizes
and unevenly positioned (L). The *ARR7-llama GFP-nes* constructs partially rescue the *drmy1*
phenotype in some buds (O, left) but not others (O, right). (P) Quantification of sepal primordium
number. Asterisk indicates statistically significance difference in a Fisher's exact test (WT vs *ahp6*,
p = 3.026×10^{-7}; *drmy1* vs *drmy1 ARR7-llama GFP-nes*, p = 0.4389). (Q) Quantification of
variability in sepal primordium position. Asterisk indicates statistically significant difference in a
Wilcoxon rank sum test (WT vs. arr3,4,5,6,7,8,9,15, p = 2.948×10^{-4}; WT vs *ahp6*, p = $2.137 \times 10^{-}$

[11]; WT vs *ARR7-llama GFPnes*, p = 1; *drmy1* vs *drmy1 ARR7-llama GFPnes*, p = 1.538×10^{-7}).

Data for *drmy1* were reused from Figure 2H, 2J. Data for *ARR7-llama GFP-nes* and *drmy1 ARR7-*

llama GFP-nes were pooled from two independent lines (7-4 and 7-6). Sample size: WT, n = 78;

arr3,4,5,6,7,8,9,15, n = 28; *ahp6*, n = 106; *ARR7-llama GFPnes*, n = 16; *drmy1*, n = 67; *drmy1*

ARR7-llama GFP-nes, n = 20.

(R) Working model. In WT, DRMY1 maintains TOR activity and translation, which sustains the

rapid production of cytokinin signaling inhibitors (ARR7 and AHP6) in response to cytokinin

signaling. These inhibitors maintain cytokinin signaling at a normal level, allowing auxin and

cytokinin signaling to interact and form robust spatial patterns. Robust patterning of auxin and

cytokinin signaling gives rise to robustly initiated sepal primordia. In *drmy1*, due to decreased

TOR signaling and translation rate, the meristem cannot rapidly produce cytokinin signaling

inhibitor proteins in response to cytokinin signaling. As a result, cytokinin signaling is upregulated,

which rescues the translation rate reduction in a homeostatic mechanism. This upregulation of

cytokinin signaling disrupts the robust spatial pattern of both cytokinin and auxin signaling, which

in turn causes variability in sepal initiation. Scale bars in all micrographs, 25 µm.