A statistical framework for quantifying the nuclear export rate of influenza viral mRNAs.

Michi Miura[a#], Naho Kiuchi[a], Siu-Ying Lau[b], Bobo Wing-Yee Mok[b], Hiroshi Ushirogawa[a], Tadasuke Naito[a], Honglin Chen[b], Mineki Saito[a#]

[a] Department of Microbiology, Kawasaki Medical School, Japan
[b] State Key Laboratory for Emerging Infectious Diseases, Department of
  Microbiology, The University of Hong Kong, Hong Kong SAR, People's Republic of
  China

[#]Correspondence

Michi Miura: michi.miura@med.kawasaki-m.ac.jp; michi.miura.res@gmail.com

Mineki Saito: mineki@med.kawasaki-m.ac.jp

Abstract

Influenza A virus acutely transcribes viral mRNAs from the eight segmented viral genome when it infects. The kinetics of viral transcription, nuclear export of viral transcripts, and their potential variation between the eight segments are poorly characterised. Here we introduce a statistical framework for estimating the nuclear export rate of each segment from a snapshot of mRNA *in situ* localisation at a single time point, exploiting the cell-to-cell variation observed by an imaging-based *in situ* transcriptome assay. A mathematical modelling indicated that the relationship between the nuclear ratio and the total count of mRNAs in single cells is dictated by a proxy for the nuclear export rate. Using this model, we showed that the two influenza viral antigens hemagglutinin and neuraminidase were the slowest segments in the nuclear export, suggesting that influenza A virus uses the nuclear retention of viral transcripts to delay the expression of antigenic molecules. Our framework presented in this study can be widely used for investigating the nuclear retention of nascent transcripts produced in a transcription burst.

Introduction

Temporal gene expression is of vital importance for viral pathogens to proliferate in the host. Human exogenous retroviruses insert their RNA genome into the host chromatin and use host factors to minimise the viral protein production to evade the immune response and persist in the host [1]. In contrast, influenza A virus acutely infects the respiratory tract and produces thousands of viral progenitors in a single round of infection to spread in the surrounding epithelial cells by transcribing and replicating its RNA genome [2]. Influenza A virus genome is segmented into eight single-stranded RNAs, and expression of each gene is essential for productive infection [3]. Unlike the human exogenous retroviruses, influenza A virus does not seem to have a sophisticated mechanism of controlling the viral gene expression at the transcription level. However, it has been known for decades that hemagglutinin (HA) and neuraminidase (NA), the two influenza viral antigens targeted by the host immune response, are produced late during the course of infection [4, 5, 6], a mechanism of which is currently unclear.

Eight viral RNA segments are transcribed by the same set of molecular machinery [7]. Viral genome is contained in the viral envelope in association with viral proteins. The 5′ and 3′ ends of each viral genomic RNA hybridise to form an intramolecular loop [8]. The RNA-dependent RNA polymerase of influenza A virus, comprising PB1 (polymerase basic 1), PB2 (polymerase basic 2), and PA (polymerase acidic) subunits, is associated with the 5′ and 3′ joint end; the rest of the loop is bound by the oligomeric viral nucleoprotein (NP) [9]. The viral ribonucleoprotein complex (vRNPs) initiates the viral transcription in the nucleus by deriving a 5′-capped short RNA fragment from the host nascent RNA on the stalled RNA polymerase II (Pol II) [10]. At the end of transcription, the viral polymerase adds a poly-A tail to the viral mRNA by stuttering on a short uridine repeat in the viral RNA template [11]. Thus, eight viral

segments sharing the common mechanism of viral transcription with no additional layers of transcriptional controls e.g., transcription-factor binding, the transcription of eight influenza viral genes is likely to occur at the same rate across the segments that is determined by the collision of the vRNP and the stalled Pol II.

Another possible factor that would determine the rate of transcription is the length of viral RNA template [12, 13], given that the duration of transcription is rate-limiting. However, this does not agree with the order of protein expression in the influenza virus infection: PB2 (2.4 kb), PB1 (2.3 kb) and PA (2.2 kb), the three longest viral genes, and NP (1.8 kb) are expressed first, followed by the expression of three segments HA (1.8 kb), NA (1.5 kb) and M (1.0 kb) [4, 6].

A third possibility is the nuclear retention of viral mRNAs. Nuclear retention of mRNAs has gained an attention as a point of control on the protein expression in the context of transcription burst [14]. Previous studies reported the accumulation of M mRNAs in the nuclear speckles [15], NXF1-dependent nuclear export of HA, NA and M mRNAs [16, 17], and the nuclear export inhibition by the host factor hnRNPAB for M, HA, NP and NS mRNAs [18]. It is becoming clear that the influenza viral segments show differential dependencies on the nuclear export factors; and this dependency is not accounted for by the RNA structure (i.e., whether they are intron-less, unspliced or spliced). Because an incomplete set of viral segments was studied in each of these studies, the influenza viral mRNA transport remains elusive.

In this paper, we introduce a statistical framework for systematically quantifying the nuclear transport across all the eight viral segments. We localised eight viral mRNAs simultaneously in single cells *in situ*. Intracellular distribution of viral mRNAs indicated that the nuclear export rate of viral transcripts varied according to the segments. A statistical model, exploiting cell-to-cell variation in the abundance of transcripts due to the stochastic nature of multiple virological

processes prior to the onset of viral transcription, allowed for the estimation of

nuclear export rate from the intracellular distribution of mRNAs observed at a

single time point. With this model, we show that hemagglutinin and neuraminidase

mRNAs were the slowest in exiting the nucleus, suggesting that influenza A virus

uses nuclear retention of mRNAs to delay the production of antigenic molecules in

the course of viral infection. This study demonstrates that the nuclear export rate

of nascent transcripts can be estimated from a snapshot of RNA *in situ*

measurement in single cells, and our result serves as a reference for the nuclear

export rate of viral transcripts across the segments, shedding light on the

comprehensive understanding on the nuclear export of viral transcripts.

Results

Diffraction-limited, single-colour imaging of viral mRNAs.

First, we used single-molecule RNA FISH (smFISH) to study the viral transcription of

PB1, HA, NP and NS. The lung carcinoma cell line A549 was infected with the WSN

strain of influenza A virus at the multiplicity of infection (M.O.I.) of ~2 for two

hours, and the viral mRNAs were visualised with a tile of short oligonucleotide

probes, each labelled with a single fluorophore [19] (Fig. 1A). Viral mRNAs were

detected as single diffraction-limited spots in the three-dimensional image stacks,

allowing for absolute mRNA quantification (Fig. 1B).

We localised the mRNAs in 3D and counted the total number of mRNAs in each cell

(Fig. 1C). The median number of mRNAs per cell was 145, 231, 310 and 303

molecules for PB1, HA, NP and NS, respectively (Fig. 1D), in line with the theoretical

prediction that the rate of mRNA production is partly determined by the gene

length.

Figure 1

108
Fig. 1 Viral mRNA detection by single-colour RNA FISH. **(A)** A schematic diagram of mRNA
labelling for the diffraction-limited imaging. Each mRNA was hybridised with a tile of oligonucleotide,
each labelled with a single fluorophore at the 3′ end. **(B)** A representative field-of-view of the
microscope image in which NP mRNAs were stained with Quasar 670. White, smFISH signal; blue,
nucleus stained with DAPI. The red square indicates the area for the sequential close-up of the
image. The signal intensity is shown on the same grey scale across the three panels, indicated on
the right. **(C)** Spot detection in 3D. Spots are projected to either the XZ or YZ plane to indicate the
distribution of spots in a typical 3D cell volume. The colour of each spot indicates the Z position of
that spot. Spot projection to the XY plane at the bottom (magenta) highlights the XY positions of
spots relative to the nucleus boundary (blue), that is, nucleus projection on the XY plane. The grid
interval on the X and Y axis is ca. 18.3 μm. **(D)** Total count of mRNAs in each cell. The sample size
(n) indicates the number of cells segmented in the microscope images. **(E)** Intracellular distribution of
viral mRNAs. Spots in the 3D volume were projected on the XY plane, together with the cytoplasmic
boundary (black solid line) and the nucleus (blue solid line). The grid interval indicates ca. 18.3 μm.
(F) Spot segmentation within cells. The panel on the left shows the entire field-of-view (measures
~294 μm), and the panel on the right shows the close-up of the area indicated in the red square (~73
μm). Spots within the 2D nuclear projection are indicated in blue (nuclear); spots within the expanded
nuclear boundary twofold from the centre of mass of the nucleus in 2D are indicated in orange
(perinuclear); and spots outside this expansion are indicated in green (peripheral). **(G)** Distribution of

spots on the Z axis in the three segments (nuclear, perinuclear and peripheral) presented in panel F.
The red dashed line shows the median Z coordinates of the peripheral spots, used to indicate the cell
baseline. The number in the histogram for nuclear spots (0.731) indicates the fraction of spots that
were above the cell base line in the field-of-view presented in panel F. **(H)** Fraction of spots above
the cell baseline in the nuclear segment. The black dots indicate the fraction obtained in each field-of-
view, and the blue bar indicates the mean. **(I)** Fraction of spots within the 2D boundary of the nucleus.
The sample size is the same as in panel D.

What came intriguingly was the intracellular distribution of mRNAs of the four

segments. The intracellular distribution of mRNAs projected on the XY plane (Fig.

1C) indicated that the HA mRNAs were more abundant in the nucleus compared to

the other three segments (Fig. 1E).

We wished to quantify the nuclear ratio of mRNAs in each cell. The mRNA count

within the 2D nuclear projection is considered to be a good approximation for the

number of mRNAs contained in the nucleus [20, 21]. We validated this approximation

for the lung epithelial A549 cells used in this study by looking at the intracellular Z

positions of mRNAs: The mRNAs within the nuclear projection would have higher Z

positions over the cell baseline, considering that the nucleus has a diameter of

several microns, much thicker than the cytoplasm in the cell periphery. To this end,

mRNAs were segmented into three regions in the 2D projection, that is, nuclear,

perinuclear and peripheral, depending on the relative location to the nucleus (Fig.

1F). In contrast to the perinuclear (orange) and peripheral (green) mRNAs, mRNAs

within the nuclear projection (blue) were distributed around the nuclear focal plane

Z = 0 (Fig. 1G); and about 80% of mRNAs in the nuclear projection (blue) were above

the cell baseline (Fig. 1H), defined by the median of the Z distribution of peripheral

mRNAs (Fig. 1G, red dashed line). Thus, in this paper, we consider the spots within

the nuclear 2D projection to be the nuclear fraction for the sake of simplicity.

With this approximation, about 30% of the PB1, NP and NS mRNAs were nuclear,

while ~70% of HA mRNAs were in the nuclear fraction (Fig. 1I), confirming the visual

inspection that the HA mRNAs are more abundant in the nucleus at the early time

point of infection (Fig. 1E). This result indicates that the velocity of viral mRNA

export from the nucleus varies according to the viral segments. Note that our

fluorescent probes could also detect complementary RNA (cRNA), an intermediate

RNA that is produced for the viral genome replication, as it has the same polarity as

viral mRNA. However, the fluorescent signals detected at 2 hours post-infection are

deemed to be from the mRNAs, because cRNA production occurs much later than

the mRNA synthesis [13, 22].

Mapping eight viral mRNAs in single cells by multiplex RNA FISH.

Next, we wish to simultaneously quantify all the eight viral segments *in situ.* To this

end, we applied MERFISH (multiplex error-robust RNA FISH) [23], an imaging-based

transcriptome assay in which multiple rounds of probe hybridisation and imaging

are performed to identify hundreds to thousands of mRNA species *in situ* by

decoding binary fluorescent signals. For example, hundreds of distinct mRNAs can

be encoded in 16-bit binary signals, in which case eight rounds of hybridisation and

imaging would be required to read out the signals with two fluorophores [23]. In our

case of detecting eight influenza A virus segments, 6 bits were sufficient to

accommodate all the eight mRNAs, with each code being separated by two or more

binary flips from others (Fig. 2A). Thus, only three rounds of hybridisation and

two-colour imaging were required. We sought for an economical way of performing

sequential rounds of hybridisation and imaging, instead of setting up a custom-

built, automated liquid pumping system used elsewhere [23]. Inspired by an *in vitro*

reconstitution assay of molecular motors [24], we assembled a flow cell on a slide

glass, comprising two stripes of double-sided tape and the coverslip, with cells on

the coverslip facing inwards (Fig. 2A). Liquids were introduced from the right open

side and wicked from the left. During image acquisition, the two open sides were

Figure 2

184
Fig. 2 Viral mRNA detection using multiplex RNA FISH. (A) An overview of the assay. The viral
mRNAs were hybridised with the encoding probes, and then the amplification probes were annealed
(not shown in the diagram for simplicity). A flow cell was assembled in which the readout probes
were hybridised and the cells were imaged in the three sequential rounds of hybridisation and imaging.
Subsequently fluorescent signals were decoded according to the encodings listed on the right. **(B)** A
representative mapping of identified viral mRNAs on the segmented cell and nuclear boundary in the
entire field-of-view (left) and the close-up in the red square (right). The mesh grids indicate ~34-µm

sealed with rubber cement. This easy implementation of flow cell allowed us to

comfortably perform three rounds of hybridisation and two-colour imaging to read

out the 6-bit binary tags; and by decoding the 6-bit encodings, the eight viral

segments were identified in each cell (Fig. 2B). On average, about 100-200

molecules of each segment were contained per cell (Fig. 2C, top row). The

difference in the total number of spots from the result presented in Fig. 1D is likely

due to the batch effect (experimental variation).

The nuclear fraction of mRNAs varied between the viral segments (Fig. 2C, bottom

row): While ~30% of mRNAs were nuclear for PB1, NP and NS segments, about 70%

of mRNAs remained in the nucleus for HA and NA segments; for the other

segments (PB2, PA and M), around half of mRNAs were in the nucleus. This

reiterates the idea that the velocity of viral mRNA export from the nucleus varies

according to the viral segments. These results were confirmed by smFISH using

probes   for   each   segment   that   were   synthesised   economically   in-house

(Supplementary figure 1A, B and C).

Single-cell clustering according to the abundance of eight viral segments revealed

that the cell population was heterogeneous in terms of the abundance of viral

mRNAs (Fig. 2D). A pair-wise analysis of the abundance of viral segments indicated

inequal distribution of viral mRNAs, regardless of the segments (Fig. 2E): Cells that

carry abundant mRNAs from one segment tend to also carry abundant mRNAs from

the other segments. These observations suggest that the viral transcription is

susceptible to the stochastic variation in each infection [25, 26], which we subsequently

used in our statistical model to estimate the nuclear export rate.

Estimating the nuclear export rate using population statistics.

We suspected that the higher nuclear ratio of HA and NA mRNAs is due to the

slower nuclear export rate for these mRNAs compared to other segments such as

NP and NS. Thus, we wished to quantify the nuclear export rate of eight viral mRNA

species from the observed nuclear-to-cytoplasmic distribution. To this end, we

conceptualised a statistical model using the cell-to-cell variation in the abundance

of viral transcripts of each segment (Fig. 3A). In this model, influenza viral particles

are added to the cell population at time 0 h, and are allowed to undergo viral

attachment to the cell surface, membrane fusion, and vRNP migration to the

Figure 3

231
Fig. 3 Cell-to-cell heterogeneity for time-course inference. **(A)** A theoretical diagram of how viral
mRNAs emerge in single cells in the population. The font runners designate ones in which the viral
transcription was initiated earlier than others, and the slow starters are ones that started transcription
lately. The bottom plot shows the theoretical relationship between the nuclear ratio and the
abundance of transcripts in which the front runners and slow starters (and many more along the
spectrum in between) compose the clusters as indicated. **(B)** Scatter plots of nuclear fraction against
the total count in each cell, derived from the dataset presented in Fig. 2C. **(C)** Time-course of the
distribution of NP mRNA count per cell, measured by smFISH, at the indicated time points (top row);
and the change in the nuclear fraction size, plotted against the total number of transcripts (bottom
row). The sample size (n) indicates the number of cells observed at each time point.

nucleus. The waiting time for these processes vary in each infection due to the

stochastic nature of biochemical events [25, 27], and therefore, the onset of viral

transcription varies from cell to cell in the population. In other words, each cell will

have a varying duration for viral transcription by the time the observation is made

at time 2 h (Fig. 3A, top). As a result, the population contains front runners in

which the viral transcription was initiated early, and therefore abundant viral

transcripts are carried; slow starters in which the transcription began late,

containing fewer transcripts; and many other cells that lie along the spectrum

between the two. The cell population at the single point in time, therefore, forms a

trajectory of the time development of viral transcription and subsequent nuclear

export on the scale of nuclear fraction against the abundance of transcripts (Fig. 3A,

bottom): In a spontaneous transcription burst, the nascent transcripts emerge in

the nucleus at the early stage of the burst, and they are exported to the cytoplasm

as more transcripts are being synthesised [20].

Indeed, when the nuclear ratio of viral mRNA was plotted against the total quantity

of mRNA molecules observed by multiplex RNA FISH in each cell, the plots formed a

curve that originated from top left (i.e., nearly 100% of the nuclear ratio when a cell

contained a few mRNA molecules), and descended as the number of mRNA

molecules increased (30% to 70%, depending on the segments) (Fig. 3B and

Supplementary figure 2).

We further validated this concept by performing smFISH measurements along the

time-course with 40-min interval (Fig. 3C). A549 was infected with WSN at the M.O.I

of ~0.02. At each time point, the NP mRNAs were stained and quantified. At the

beginning (0 and 40 min), the mRNA count was almost zero. As the mRNA count

increased (80 min to 200 min post-infection) (Fig. 3C, top), the nuclear ratio against

the total mRNA count derived from the single cells progressed along a curve, from

the top left to the bottom right (Fig. 3C, bottom), supporting the concept of our

statistical model.

The curvature of the trace appeared to vary between segments (Fig. 3B). We

explored to see if the curvature could represent a proxy for the nuclear export rate.

Thus, we formulated a simple kinetic model in which mRNAs are synthesised at the

rate k in the nucleus, and are exported with the reaction constant ε (Fig. 4A). In this

model, the molar concentrations of mRNAs in the nuclear and cytoplasmic

compartments are described, respectively,

Figure 4

278
Fig. 4 Kinetic model for estimating the nuclear export rate. (A) A schematic diagram representing
the production of viral transcripts in the nucleus and the export to the cytoplasm. The two spherical
compartments are for the nucleus and the cytoplasm. Viral transcripts (magenta) are produced at a
rate constant k in the nucleus, and exported at a rate constant ε to the cytoplasm. **(B)** The
relationship between the nuclear fraction and the total count in the kinetic model, according to the
formula presented in panel A. The parameter α determines the curvature of the plot. **(C)** The
parameter fit to the observed datapoints. The parameter α was estimated by fitting the equation to
the observed datapoints presented in Fig. 3B. The colour representation in each line is the same as
the ones used in panel B, according to the parameter α. **(D)** Estimated nuclear export rate. The
parameter ε was estimated from the three independent assays (indicated in black for each segment);
the blue bar indicates the mean of the three measurements.

$$290 \qquad \frac{d[Nuclear]}{dt} = k - \varepsilon[Nuclear] \qquad (1)$$

$$291 \qquad \frac{d[Cytoplasmic]}{dt} = \varepsilon[Nuclear] \qquad (2)$$

292 Omitting the degradation of mRNAs, the analytical solutions of these equations

293 were:

$$294 \qquad [Nuclear] = \frac{k}{\varepsilon}(1 - e^{-\varepsilon t}) \qquad (3)$$

$$295 \qquad [Cytoplasmic] = kt - \frac{k}{\varepsilon}(1 - e^{-\varepsilon t}) \qquad (4)$$

296 Thus, the mRNA nuclear fraction φ was described as a function of total mRNA

297 molecules N_{total},

$$298 \qquad Nuclear\ fraction\ \varphi = \frac{1}{\alpha N_{total}}(1 - e^{-\alpha N_{total}}) \qquad (5)$$

299 where

$$300 \qquad \alpha = \frac{\varepsilon}{k} \qquad (6)$$

The nuclear fraction φ described in equation (5) is time-independent, and the

parameter α determines the curvature of the line that the single cells would trace in

the time-course (Fig. 4B). Thus, we estimated the parameter α by fitting to the

observed data points derived at the time in which cells were fixed (Fig. 4C). By

obtaining the synthesis rate k from the total number of mRNAs produced by the

time of fixation (Fig. 2C, top row), the nuclear export rate ε for each segment was

derived according to equation (6) (Fig. 4D). The nuclear export rate ε agreed with

the historical classification of the early and late influenza A virus genes [4, 6],

indicating that the timing of influenza A virus protein expression in the initial stage

of infection is determined by the velocity of the nuclear export. Notably, the HA

and NA mRNAs had the slowest nuclear export rate, suggesting that influenza A

virus delays the viral antigen production by nuclear retention of viral transcripts.

Discussion

Based on the stochastic nature of influenza viral transcription studied by single-cell sequencing [25, 26], we introduced a quantitative framework for estimating the nuclear export rate of viral mRNAs to advanced our current understanding on the kinetics of viral gene expression. This was achieved by *in situ* localisation of eight influenza A virus segments simultaneously, coupled with the statistical model that allowed for estimating the nuclear export rate from the intracellular distribution of mRNAs at a single time point. This is based on the concept that the cells in the population are at various stages of transcription at the time the observation is made, due to the multiple stochastic processes the virus needs to overcome before it begins transcription. We demonstrated that the initiation of each viral transcription is concurrent in the eight viral segments; it is instead the nuclear export rate that controls the temporal protein expression at the initial phase of infection. The early production of the polymerase subunits (PB2, PB1 and PA) and nucleoprotein would facilitate the formation of vRNP, and hence the viral transcription and genome replication. On the other hand, pushing back the expression of viral antigens HA and NA would be beneficial for the virus to delay the host immune response against the infected cells in which the virus is being replicated.

Instead of time-series measurements, the estimation of mRNA export rate was made from a snapshot of *in situ* RNA localisation data taken at a single time point. We anticipate that the statistical framework for quantifying the nuclear export rate proposed in our study can be widely used and benefits the research area on the regulation of gene expression, especially since nuclear retention has been recognised as a point of control against the transcription burst [14]. For example, it is applicable for the cell population in which *de novo* synthesis of mRNAs occurs concurrently, such as a stimuli-induced activation of immediate-early genes [21, 28]. On

the contrary, this approach is not appropriate for scenarios in which asynchronous

transcription bursts are taking place randomly in each single cell [20, 29]; cells at the

late stage of transcription burst, having residual amount of transcripts in the

cytoplasm [20], would obscure the time-course trajectory of the nuclear export.

The experiment in this study was conducted at the early stage of infection (up to

200 min post-infection). This conferred two advantages over investigating

influenza viral mRNA export at the late stage of infection (e.g., 6-12 hours post-

infection). First, our study allowed for the absolute quantification of viral

transcripts at the single-molecule level in each cell, and thereby enabled us to

formulate the model and fit the equation to the numerical data points. Otherwise

at the late stage of infection, accurate quantification of mRNAs by the imaging-

based single-molecule detection would not be feasible due to the high density of

mRNAs. Second, at the late stage of infection, NS1 protein would be expressed in

the cytoplasm. NS1 protein would then migrate into the nucleus and alter the

nuclear export of viral mRNAs [30], obscuring the inherent nature of nuclear export of

each viral segment. Thus, in our study, we elucidated the quantitative measure for

the influenza viral mRNA nuclear export that is intrinsic to each viral segment.

Host factors that relay the influenza viral mRNAs to the nuclear pore remain elusive.

NXF1 and CRM are two major cellular mRNA export pathways for the 5′-capped

mRNAs [31]. Of these two pathways, influenza A virus mRNAs, also carrying 5′ cap

structure derived from the host nascent mRNA and the poly-A tail as a result of

viral polymerase stuttering the uridine repeat, are dependent on the NXF1 pathway

[16, 17]. However, the eight viral segments show differential dependencies on this

export pathway. The segments HA, M and NS is exported by NXF1 and UAP56,

whereas PB2, PB1 and PA appear to be independent of the NXF1 pathway [17]. The

nuclear export rate derived in our study does not agree with the dependencies on

NXF1. According to the previous study, both HA and NS are dependent on NXF1

and UAP56 [16]; however, in our study, HA and NS had significantly different nuclear

export rate.

Two viral segments M and NS are subject to splicing, yielding unspliced and spliced

transcripts. Our current system lacks the ability of discriminating between the

unspliced and spliced forms of these segments; although we infer that the majority

of NS segments that we detected in this study were in the unspliced form because

we obtained similar result with probes exclusively detecting the unspliced form of

NS (Supplementary figure 3). An observation that the unspliced form of M and NS

dominates throughout the infection supports this [13]. In either case, the RNA

structure does not appear to accord with the NXF1 dependency, nor the nuclear

export rate, either.

Our study suggests that the nuclear export of influenza viral transcripts is intrinsic

to the nucleotide sequence of each segment. Recently, the human nuclear factor

hnRNPAB was shown to associate with the nucleoprotein of influenza A virus, and

inhibit the NXF1 pathway. It is tempting to speculate that the host factors (e.g.,

hnRNPAB) have differential binding affinities to each segment, thus determining the

velocity of nuclear export (or the degree of nuclear export inhibition). It is currently

unknown what factors binds to the early genes such as PB2, PB1 and PA. Further

studies will be required to identify the host factors that bind to these transcripts

for the nuclear export. Our study serves as a reference for the nuclear export rate

of each segment, and thus sheds light to comprehensive understanding on the

nuclear export of viral segments. Having revealed the nuclear retention of

influenza viral transcripts at the unprecedented resolution, this study opens up a

new research direction on how the nuclear retention is regulated and its implication

in viral pathogenesis.

Materials and Methods

Cells and virus preparation

The human lung carcinoma cell line A549 was maintained in Dulbecco's Modified Eagle's Medium (DMEM) (Sigma, D5796) supplemented with 10% FBS (Nichirei Biosciences, 175012), 100 U/ml penicillin and 100 μg/ml streptomycin (Gibco, 15140-122) at 37°C in 5% CO_2. Madin-Darby canine kidney (MDCK) cell line was maintained in Minimum Essential Medium (MEM) (Sigma, M4655) supplemented with 10% FBS, 100 U/ml penicillin and 100 μg/ml streptomycin at 37°C in 5% CO_2. Influenza A virus strain WSN was propagated in MDCK cells in a 75 cm^2 flask. The plaque-forming unit of the harvested virus was determined using MDCK cells grown confluent in a 6-well plate.

Coverslip preparation

An 18-mm round coverslip (0.17±0.005 mm thickness, Matsunami Glass) was used. For MERFISH, two straight sides were created by cutting two edges off the glass using a laser-wheel glass cutter (Nikken Dia, NC-X03 LASER 110°). The coverslip was rinsed in 70% ethanol and placed in a 12-well plate. A549 cells were seeded and attached onto the coverslip overnight. On the following day cells were rinsed twice with PBS and infected with WSN at the indicated multiplicity of infection in MEM for 2 hours (unless indicated otherwise in Fig. 3C) in the absence of FBS. After the infection, cells were fixed with 4% paraformaldehyde (Electron Microscopy Sciences, 15714-S) in PBS for 10 min and then rinsed three times with PBS. The coverslip was immersed in 70-75% ethanol and stored at −20°C until use.

smFISH

The probes used in Figs. 1, 3C, Supplementary figure 1 (probes for NP) and Supplementary figure 3 were obtained from LGC Biosearch Technologies. Stellaris RNA FISH Probe Designer (https://www. biosearchtech.com/stellaris-designer) was used to design the probes against the coding region of WSN mRNAs.

The coverslip was equilibrated in 1 ml of Wash buffer A (Stellaris) for >5 min in a 12-well plate prior to hybridisation. Cells were hybridised to 125 nM probes in Hybridization buffer (Stellaris) on a Hybrislip (Sigma, GBL712222) at 37°C overnight in a humidified chamber. On the following day the coverslip was rinsed twice in 1 ml of Wash buffer A (Stellaris), each for 30 min. Cytoplasmic stain (HCS Cellmask Green Stain, Invitrogen, H32714) was included in the second wash at the concentration of 50 ng/ml in Figs 1, 3C and Supplementary figure 3. Then the coverslip was rinsed in 1 ml of Wash buffer B (Stellaris) for 5 min. The coverslip was mounted in anti-fade medium containing DAPI (Vector Laboratories, H-1200) on a slide glass and sealed with nail polish.

For an economical reason and to allow for more flexible design of probes to increase the number of fluorophores attached per mRNA molecule, the probes used in Supplementary figure 1 (except the NP probes) were synthesised in-house (Supplementary figure 1A). (The NP probe used in Supplementary figure 1 was purchased from LGC Biosearch.) Candidate probe-binding sites in the coding region of each viral segment were identified using OligoMiner [32], and were further selected against the human RefSeq. The sequences were flanked by P5 (5′ AAT GAT ACG GCG ACC ACC GA 3′) and P7 (5′ CAA GCA GAA GAC GGC ATA CGA GAT 3′), with P7 at the 5′ end and P5 at the 3′ end of the probe sequence (Supplementary figure 1A). For the probes against HA, NA, M and NS, additional 20-nt sequences RS0332 (5′ GGG AGA ATG AGG TGT AAT GT 3′) and RS0406 (5′ GAT GAT GTA GTA GTA AGG

GT 3′) [23] were inserted between P7 and the probe, the probe and P5, respectively

(Supplementary figure 1A). These two insertions were used for additional probes to

bind to increase the number of fluorophores per molecule. An oligonucleotide pool

comprising the set of probe-binding sites with these additional nucleotide

sequences was synthesised at Integrated DNA Technologies (IDT). The

oligonucleotide pool was subjected to PCR amplification (98°C for 30 sec; 18 cycles

of 98°C for 5 sec, 64°C for 10 sec and 72°C for 20 sec; and 72°C for 5 min) (Phusion

High-Fidelity DNA Polymerase, NEB, M0530S) using the P7 and a T7-fused P5 primer

(5′ TAA TAC GAC TCA CTA TAG GGA ATG ATA CGG CGA CCA CCG A 3′)

(Supplementary figure 1A). RNA was synthesised using T7 RNA polymerase

(HiScribe T7 High Yield RNA Synthesis Kit, NEB, E2040S). Then RNA was reverse-

transcribed using a recombinant Moloney leukemia virus reverse transcriptase

(GeneAce cDNA Synthesis Kit, Nippon Gene, 319-08881) with an ATTO550-

conjugated P7 primer at the 5′ end. The single-stranded DNA probe was purified

using a spin column (Zymo Research, D7010) after RNA digestion with a cocktail of

RNase H and RNase A. The probe was further purified by Urea-PAGE to remove any

residual RNA and the fluorescent-labelled primer.

The coverslip was equilibrated in 1 ml of 30% formamide buffer composed of 2×SSC

and 30% deionised formamide (Sigma, F9037) in a 12-well plate for >5 min prior to

hybridisation. Cells were hybridised with the probes at the concentration of 5.6 nM

per probe in 100 μl Hybridization buffer composed of 2×SSC, 30% deionised

formamide, 0.1% yeast tRNA (Invitrogen, 15401-011), 1% (v/v) RNase inhibitor (NEB,

M0314S) and 10% (w/v) Dextran Sulfate (Merck, S4030) [23] on a Hybrislip at 37°C

overnight in a humidified chamber. On the following day the coverslip was rinsed

twice in 1 ml of 30% formamide buffer, each for 30 min. DAPI was included at 5

ng/ml in the second wash if additional staining was not performed with the

external probes (for PB2, PB1 and PA). In this case the coverslip was rinsed in 1 ml

of PBS after the DAPI staining, mounted in anti-fade medium (Vector Laboratories,

H-1900) on a slide glass and sealed with nail polish. For staining HA, NA, M and NS

with the extra probes, the coverslip was first equilibrated with 10% formamide

buffer composed of 2×SSC, 10% deionised formamide and 0.05% (v/v) RNase

inhibitor, and then hybridised with 10 nM each of probes for RS0332 and RS0406 [23],

both labelled with Quasar 570 (LGC Biosearch) at the 3′ end, in Stellaris

hybridisation buffer (containing 10% formamide) at 37°C in the humidified chamber

for 1 hour. Cells were washed in 10% formamide buffer twice, with DAPI included

in the second wash. The coverslip was rinsed in PBS, mounted in the anti-fade

medium (Vector Laboratories, H-1900), and sealed with nail polish.

Sequential hybridisation and imaging in the flow cell

An oligonucleotide pool for synthesising the encoding probes used in Fig. 2,

comprising the target-binding sequence, readout tags compatible with the two-step

amplification probes [23, 33], and the P5 and P7 PCR tags, were obtained from IDT. The

target-binding sites were identified in the coding region of each segment using

OligoMiner [32] and selected against human RefSeq. The readout tags RS0332, RS0343,

RS0406, RS0255, RS0015 and RS038 [23, 33] were inserted adjacent to the upstream or

downstream of the target-binding sites for the 6-bit encoding of each segment (Fig.

2A). Probes were synthesised by PCR, T7 transcription and reverse-transcription as

described above.

A coverslip was equilibrated in 1 ml of 30% formamide buffer (2×SSC and 30%

deionised formamide) [23]. Cells were hybridised with 1.6 μM encoding probes in

Hybridisation buffer composed of 2×SSC, 30% deionised formamide, 0.1% yeast

tRNA, 1% RNase inhibitor, 10% dextran sulfate [23] on a Hybrislip at 37°C overnight in

a humidified chamber. Cells were washed twice in 1 ml of 30% formamide buffer at

37°C, and then equilibrated in 1 ml of Readout-probe hybridisation and wash buffer composed of 2×SSC, 10% deionised formamide and 20 units/ml Murine RNase inhibitor [23]. The first and second amplification probes for RS0332, RS0255, RS0343, RS0406, RS0015 and RS0384, described previously [33], were annealed sequentially at 10 nM of each probe in Readout-probe hybridisation and wash buffer for 2 hours at 37°C on a Hybrislip. The coverslip was washed for 30 min twice in Readout-probe hybridisation and wash buffer at each of the first and second probe hybridisation. At the final wash, 0.10 μm green-fluorescent carboxyl polystyrene beads (Bangs Laboratories, FCDG002) were included (~0.1% v/v) as a fiducial marker for image registration. The cells and beads were fixed with 3.2% paraformaldehyde in PBS for 10 min. The coverslip was rinsed in PBS three times and stored in 2×SSC supplemented with 40 units/ml RNase inhibitor at 4°C overnight.

On the following day a flow cell was assembled on a slide glass, composed of two parallel stripes of double-sided tape (0.23 mm thickness, ~5 mm apart) and the coverslip on top, with cells facing inwards (Fig. 2A). The cells at the top and bottom end of the coverslip were scraped using a cell scraper before the assembly and any trace amount of liquid remaining on the exposed surface was removed with filter paper to ensure that the coverslip is firmly attached to the tape. The chamber was immediately flushed with 2×SSC to avoid cells drying up.

Readout-probe hybridisation and wash buffer was flushed in the chamber to equilibrate the cells. Readout probes were introduced at 10 nM in Readout-probe hybridisation and wash buffer and incubated at 37°C in a humidified chamber for 30 min. Subsequently, Readout-probe hybridisation and wash buffer was flushed to wash away the probes. Imaging buffer, composed of 2×SSC, 50 mM Tris-HCl pH 8, 10% glucose (Fujifilm Wako, 049-31165), 2 mM Trolox (Sigma, 238813), 0.5 mg/ml glucose oxidase (Sigma, G2133), 40 μg/ml catalase (Sigma, C30) and 0.1% RNase

inhibitor [23], was flushed to the flow cell prior to the image acquisition. The flow cell

was sealed at the both sides with Fixogum rubber cement ejected through a 200-μl

pipette tip during the image acquisition.

After the imaging, the disulfide bond linking the probe and the fluorophore was

cleaved with Cleavage buffer comprising 2×SSC and 50 mM TCEP (Sigma, 646547)

for 15 min. The chamber was flushed with Readout-probe hybridisation and wash

buffer to wash away the free fluorophores prior to introducing the readout probes

for the next round. Readout probes used in this study are listed in Table 1.

Table 1

ID	Round	Channel	Sequence (5′ to 3′) and fluorescent dye	Manufacturer
RS0332amp	1	1	TTC TTC CCT CAA TCT TCA TC -S-S- Q570	LGC Biosearch
RS0343amp	1	2	TAC TAC AAA CCC ATA ATC CC -S-S- Q670	LGC Biosearch
RS0406amp	2	1	AAT CTC ACC TTC CAC TTC AC -S-S Q570	LGC Biosearch
RS0255amp	2	2	TCA CCT CTA ACT CAT TAC CT -S-S- Q670	LGC Biosearch
RS0015amp	3	1	TCT CAC ACC ACT TTC CTC AT -S-S- ATTO550	Generay
RS0384amp	3	2	TCC TCA TCT TAC TCC CTC TA -S-S- ATTO647N	Generay

Imaging platform

Images were acquired using a Nikon Ti2-E with a Nikon Plan Apo Lambda 60x (NA

1.40) (Fig. 1, Supplementary figure 3) or 100x (NA 1.45) (Fig. 3C) objective lens

equipped with a Prime 95B 25MM back-illuminated sCMOS camera (Photometrics).

Otherwise, an Olympus IX83 equipped with an Olympus UPlan XApo 60x (NA 1.42)

objective lens and a cooled monochrome CCD camera (DP80, Olympus) was used

(Fig. 2 and Supplementary figure 1).

Spot quantification

Fluorescent spots detected by smFISH were quantified using either FISH-QUANT [34]

(Figs. 1, 3C and Supplementary figure 3) or a more recent Python package Big-FISH [35]

(Supplementary figure 1). Cytoplasmic and nuclear segmentation was performed

using Cellpose [36, 37]. Objects that were on the boundary of the image was eliminated.

Cytoplasmic regions with no nucleus therein due to the erroneous segmentation

was also eliminated. The identification numbers assigned to each object by

Cellpose were rearranged from scratch so that the cytoplasm and nucleus of the

same cell correspond for the downstream analyses. In order to make the Cellpose

mask readable in FISH-QUANT, the Cellpose mask were converted into x and y

coordinates by tracing each object in the mask using the Moore neighbourhood

edge-finding algorithm.

For MERFISH signal decoding, signals from the green-fluorescent carboxyl beads

were used to register the images. Images at the corresponding field-of-view from

the three rounds of imaging were registered using phase_cross_correlation from the

Python package skimage.registration. The registered images were filtered using

Laplacian-of-Gaussian implemented in Big-FISH, and the image intensity was

manually normalised. Fluorescent signals were then decoded using

PixelSpotDecoder in DetectPixels implemented in starfish (https://github.com/

spacetx/starfish).

Data analyses and presentation

Microscopy images were presented using the Python package microfilm

(https://github.com/guiwitz/microfilm) (Fig. 1B). Cell clustering was performed

with the method centroid implemented in scipy.cluster.hierarchy.linkage from SciPy

(https://scipy.org/) (Fig. 2D). The parameter α in equation (5) was estimated using

curve_fit implemented in scipy.optimize from SciPy (Fig. 4C).

Acknowledgements

We thank Imaging and Flow Cytometry Core, The Centre for PanorOmic Sciences (CPOS), LKS Faculty of Medicine of The University of Hong Kong for the use of Nikon Ti2-E widefield imaging system; and Central Research Institute of Kawasaki Medical School for the use of Olympus IX83 imaging system. This study was supported by Research Project Grant R04S-003 and R03S-001 from Kawasaki Medical School, The KAWASAKI Foundation of Medical Science and Medical Welfare, and Grant-in-Aid for Scientific Research (C) (22K07091) from Japan Society for the Promotion of Science (JSPS) to MM; Grant-in-Aid for Scientific Research (B) (21H03188) from JSPS to TN; the General Research Fund (17107019) of the Research Grants Council, Hong Kong SAR to HC.

Author contributions

Michi Miura: Conceptualization, Methodology, Software, Validation, Formal analysis, Investigation, Resources, Data Curation, Writing - Original Draft, Writing - Review & Editing, Visualization, Supervision, Project administration, Funding acquisition. Naho Kiuchi: Software, Validation, Investigation, Writing - Review & Editing. Siu-Ying Lau: Resources, Writing - Review & Editing. Bobo Wing-Yee Mok: Resources, Writing - Review & Editing. Hiroshi Ushirogawa: Resources, Writing - Review & Editing. Tadasuke Naito: Resources, Writing - Review & Editing, Funding acquisition. Honglin Chen: Resources, Data Curation, Writing - Review & Editing, Supervision, Project administration, Funding acquisition. Mineki Saito: Resources, Data Curation, Writing - Review & Editing, Supervision, Project administration, Funding acquisition.

Declaration of interests

The authors declare no competing interests.

Data and code availability

The probe sequences, original images, and the custom codes used in this study are available upon request.

597 **Reference**

598 1.   Bangham CRM, Miura M, Kulkarni A, Matsuoka M. Regulation of Latency in
599   the Human T Cell Leukemia Virus, HTLV-1. *Annu Rev Virol* **6**, 365-385 (2019).

600
601 2.   Einav T, Gentles LE, Bloom JD. SnapShot: Influenza by the Numbers. *Cell* **182**,
602   532-532 e531 (2020).

603
604 3.   Dou D, Revol R, Ostbye H, Wang H, Daniels R. Influenza A Virus Cell Entry,
605   Replication, Virion Assembly and Movement. *Front Immunol* **9**, 1581 (2018).

606
607 4.   Lamb RA, Choppin PW. The gene structure and replication of influenza virus.
608   *Annu Rev Biochem* **52**, 467-506 (1983).

609
610 5.   Shapiro GI, Gurney T, Jr., Krug RM. Influenza virus gene expression: control
611   mechanisms at early and late times of infection and nuclear-cytoplasmic
612   transport of virus-specific RNAs. *J Virol* **61**, 764-773 (1987).

613
614 6.   Hatada E, Hasegawa M, Mukaigawa J, Shimizu K, Fukuda R. Control of
615   influenza virus gene expression: quantitative analysis of each viral RNA
616   species in infected cells. *J Biochem* **105**, 537-546 (1989).

617
618 7.   Te Velthuis AJ, Fodor E. Influenza virus RNA polymerase: insights into the
619   mechanisms of viral RNA synthesis. *Nat Rev Microbiol* **14**, 479-493 (2016).

620
621 8.   Fodor E, Pritlove DC, Brownlee GG. Characterization of the RNA-fork model
622   of virion RNA in the initiation of transcription in influenza A virus. *J Virol* **69**,
623   4012-4019 (1995).

624
625 9.   Arranz R*, et al.* The structure of native influenza virion ribonucleoproteins.
626   *Science* **338**, 1634-1637 (2012).

627
628 10.   Walker AP, Fodor E. Interplay between Influenza Virus and the Host RNA
629   Polymerase II Transcriptional Machinery. *Trends Microbiol* **27**, 398-407 (2019).

630
631 11.   Poon LL, Pritlove DC, Fodor E, Brownlee GG. Direct evidence that the poly(A)
632   tail of influenza A virus mRNA is synthesized by reiterative copying of a U
633   track in the virion RNA template. *J Virol* **73**, 3473-3476 (1999).

634
635 12.   Enami M, Fukuda R, Ishihama A. Transcription and replication of eight RNA
636   segments of influenza virus. *Virology* **142**, 68-77 (1985).

637
638 13.   Phan T, Fay EJ, Lee Z, Aron S, Hu WS, Langlois RA. Segment-specific kinetics
639   of mRNA, cRNA and vRNA accumulation during influenza infection. *J Virol*
640   **95**, (2021).

14. Bahar Halpern K, *et al.* Nuclear Retention of mRNA in Mammalian Tissues. *Cell Rep* **13**, 2653-2662 (2015).

15. Mor A, *et al.* Influenza virus mRNA trafficking through host nuclear speckles. *Nat Microbiol* **1**, 16069 (2016).

16. Read EK, Digard P. Individual influenza A virus mRNAs show differential dependence on cellular NXF1/TAP for their nuclear export. *J Gen Virol* **91**, 1290-1301 (2010).

17. Larsen S, Bui S, Perez V, Mohammad A, Medina-Ramirez H, Newcomb LL. Influenza polymerase encoding mRNAs utilize atypical mRNA nuclear export. *Virol J* **11**, 154 (2014).

18. Wang X, *et al.* Cellular hnRNPAB binding to viral nucleoprotein inhibits flu virus replication by blocking nuclear export of viral mRNA. *iScience* **24**, 102160 (2021).

19. Raj A, van den Bogaard P, Rifkin SA, van Oudenaarden A, Tyagi S. Imaging individual mRNA molecules using multiple singly labeled probes. *Nat Methods* **5**, 877-879 (2008).

20. Billman MR, Rueda D, Bangham CRM. Single-cell heterogeneity and cell-cycle-related viral gene bursts in the human leukaemia virus HTLV-1. *Wellcome Open Res* **2**, 87 (2017).

21. Miura M, Dey S, Ramanayake S, Singh A, Rueda DS, Bangham CRM. Kinetics of HTLV-1 reactivation from latency quantified by single-molecule RNA FISH and stochastic modelling. *PLoS Pathog* **15**, e1008164 (2019).

22. Kawakami E, *et al.* Strand-specific real-time RT-PCR for distinguishing influenza vRNA, cRNA, and mRNA. *J Virol Methods* **173**, 1-6 (2011).

23. Moffitt JR, Hao J, Wang G, Chen KH, Babcock HP, Zhuang X. High-throughput single-cell gene-expression profiling with multiplexed error-robust fluorescence in situ hybridization. *Proc Natl Acad Sci U S A* **113**, 11046-11051 (2016).

24. Miura M, Matsubara A, Kobayashi T, Edamatsu M, Toyoshima YY. Nucleotide-dependent behavior of single molecules of cytoplasmic dynein on microtubules in vitro. *FEBS Lett* **584**, 2351-2355 (2010).

25. Heldt FS, Kupke SY, Dorl S, Reichl U, Frensing T. Single-cell analysis and stochastic modelling unveil large cell-to-cell variability in influenza A virus infection. *Nat Commun* **6**, 8938 (2015).

688   26.   Russell AB, Trapnell C, Bloom JD. Extreme heterogeneity of influenza virus
689   infection in single cells. *Elife* **7**, (2018).

691   27.   Heldt FS, Frensing T, Reichl U. Modeling the intracellular dynamics of
692   influenza virus replication to understand the control of viral RNA synthesis.
693   *J Virol* **86**, 7806-7817 (2012).

695   28.   Kulkarni A, Taylor GP, Klose RJ, Schofield CJ, Bangham CR. Histone H2A
696   monoubiquitylation and p38-MAPKs regulate immediate-early gene-like
697   reactivation of latent retrovirus HTLV-1. *JCI Insight* **3**, (2018).

699   29.   Kiik H, Ramanayake S, Miura M, Tanaka Y, Melamed A, Bangham CRM. Time-
700   course of host cell transcription during the HTLV-1 transcriptional burst.
701   *PLoS Pathog* **18**, e1010387 (2022).

703   30.   Pereira CF, Read EKC, Wise HM, Amorim MJ, Digard P. Influenza A Virus NS1
704   Protein Promotes Efficient Nuclear Export of Unspliced Viral M1 mRNA. *J
705   Virol* **91**, (2017).

707   31.   Palazzo AF, Lee ES. Sequence Determinants for Nuclear Retention and
708   Cytoplasmic Export of mRNAs and lncRNAs. *Front Genet* **9**, 440 (2018).

710   32.   Beliveau BJ*, et al.* OligoMiner provides a rapid, flexible environment for the
711   design of genome-scale oligonucleotide in situ hybridization probes. *Proc
712   Natl Acad Sci U S A* **115**, E2183-e2192 (2018).

714   33.   Xia C, Babcock HP, Moffitt JR, Zhuang X. Multiplexed detection of RNA using
715   MERFISH and branched DNA amplification. *Sci Rep* **9**, 7721 (2019).

717   34.   Mueller F*, et al.* FISH-quant: automatic counting of transcripts in 3D FISH
718   images. *Nat Methods* **10**, 277-278 (2013).

720   35.   Imbert A*, et al.* FISH-quant v2: a scalable and modular tool for smFISH image
721   analysis. *RNA* **28**, 786-795 (2022).

723   36.   Stringer C, Wang T, Michaelos M, Pachitariu M. Cellpose: a generalist
724   algorithm for cellular segmentation. *Nat Methods* **18**, 100-106 (2021).

726   37.   Pachitariu M, Stringer C. Cellpose 2.0: how to train your own model. *Nat
727   Methods* **19**, 1634-1641 (2022).

Supplementary figure 1: smFISH for eight segments

Supplementary figure 1 Independent validation of the total mRNA count and the nuclear ratio of each segment by smFISH. (A) A schematic diagram of in-house probe synthesis. The pool of single-stranded DNA templates were PCR-amplified, T7-transcribed, and reverse-transcribed using the fluorescent-labelled primer. The number of distinct probes contained in the pool is indicated for each segment. For HA, NA, M and NS segments, two additional fluorescent probes bind to each probe. As a result, the number of fluorophores bound per mRNA molecule is theoretically three times as many as the probes for those segments. **(B)** Total count of mRNAs per cell for each segment. The sample size (n) indicates the number of cells analysed. **(C)** The nuclear ratio of mRNAs in each cell. The sample size is the same as the one indicated in panel B.

Supplementary figure 2: Nuclear ratio vs. total count

Supplementary figure 2 Independent validation of the nuclear ratio against the total count of mRNAs for each segment by smFISH. The nuclear ratio of mRNAs in each cell was plotted against the total number of mRNAs in that cell. The original data were derived from Supplementary figure 1B and C.

Supplementary figure 3: Unspliced form of NS segment

Supplementary figure 3   Exclusive detection of unspliced NS transcripts.   (A) Probe-binding sites on the NS segment. The exon probes, used in Fig. 1, target the second exon of NS2 (spliced form), and the intron probes are against the region that is spliced out. Thus, exon probes detect both NS1 and NS2, while intron probes exclusively detect NS1. The black line represents the NS gene. The protein coding region is indicated with the wide horizontal bar. **(B)** The total spot count (left column) detected using either exon probes or intron probes. The spot nuclear ratio in each cell is indicated on the right. Note that the plots of the exon probes (orange) are the reproduction from Fig. 1 and are presented for comparison with those of the intron probes (magenta).