Genetic variation in *Loudetia simplex* supports the presence of ancient grasslands in Madagascar

George P. Tiley[1,†], Andrew A. Crowl[2], Tchana O. M. Almary[3], W. R. Quentin Luke[4], Cédrique L. Solofondranohatra[5], Guillaume Besnard[6], Caroline E.R. Lehmann[7,8], Anne D. Yoder[2], Maria S. Vorontsova[1]

[1]Royal Botanic Gardens Kew, Richmond, UK

[2]Department of Biology, Duke University, Durham, North Carolina, 27708 USA

[3]Kew Madagascar Conservation Centre, Antananarivo 101, Madagascar

[4]East African Herbarium, National Museums of Kenya, P.O. Box 45166, Nairobi, 00100, Kenya

[5]Madagascar Biodiversity Center, Parc Botanique et Zoologique de Tsimbazaza, Antananarivo 101, Madagascar

[6]CNRS-UPS-IRD, UMR5174, Laboratoire Évolution & Diversité Biologique, Université Paul Sabatier, Toulouse, France

[7]School of GeoSciences, University of Edinburgh, Edinburgh, UK

[8]Royal Botanic Garden Edinburgh, Edinburgh, UK

[†]Author for Correspondence: g.tiley@kew.org

Summary

1) Research Aims — The extent of Madagascar's grasslands prior to human colonization is unresolved. We used population genetic analyses of a broadly dominant C_4 fire-adapted grass, *Loudetia simplex*, as a proxy for estimating grassland change through time. In the absence of population genomic resources, we used target-enrichment data. We carefully examined the utility of these data for population genetics to make recommendations on cost-effective strategies for conservation genetics. We explored the potential of estimating individual ploidy levels from target-enrichment data and how assumptions about ploidy could affect analyses.

2) Methods — We developed a novel bioinformatic pipeline based to estimate ploidy and genotypes from target-enrichment data. We estimated standard population genetic summary statistics in addition to species trees and population structure. Extended Bayesian skyline plots provided estimates of population size through time for empirical and simulated data.

3) Key Result — All Malagasy *Loudetia simplex* individuals sampled in this study formed a clade and possibly indicated an ancestral high-altitude distribution near the present-day Itremo protected area. Demographic models suggested grassland expansions occurred prior to the Last Interglacial Period and supported extensive grasslands prior to human colonization. Though there are limitations to target-enrichment data for population genetic studies, we find that analyses of population structure are reliable.

4) Key Point —Genetic variation in *Loudetia simplex* supports widespread grasslands in Madagascar prior to the more recent periods of notable paleoclimatic change. However, the methods explored here could not differentiate between paleoclimatic change near the Last Glacial Maximum and anthropogenic effects. Target-enrichment data can be a valuable tool for analyses of population structure in the absence of reference genomic resources.

Keywords

Effective Population Size, Target-Enrichment, HybSeq, Polyploidy, Anthropogenic Climate Change, Conservation Genetics

Societal Impact Statement

Recognizing *Loudetia* dominated grasslands were widespread prior to human colonization highlights that open ecosystems were and continue to be an important component to Madagascar's biodiversity. Urgently required are biodiversity inventories and risk assessments for unique grassland flora and fauna under present day environmental conditions to recognize and quantify modern human impacts within ecosystems historically regarded as wastelands. Substantial financial and logistic barriers exist to implementing conservation studies using contemporary genomic tools that we seek to ameliorate by developing computational resources to leverage a cost-effective data generation strategy that requires no prior genetic knowledge of the target species.

Introduction

The degree to which anthropogenic activities have transformed the ancestral vegetative composition of Madagascar's ecosystems is heavily debated. While the narrative that most of central Madagascar was once covered in closed-canopy forests has been rejected (Bond et al., 2008; Vorontsova et al., 2016; Hackel et al., 2018 Solofondranohatra et al., 2020; Bond et al., 2022; Lehmann et al., 2022), the degree to which a former mosaic of grassland, shrubland, and forests has become more dominated by grasses is contentious (reviewed in Joseph et al., 2021). Understanding population dynamics of individual ecologically important species and plant communities will have profound effects on conservation policy decisions and the economic productivity of grasslands for local communities. However, it is difficult to determine where grasslands were in the past and how extensive they were. Stratigraphic sediment records from central Madagascar indicate an increased relative abundance of grasses and fire through the Holocene (Burney, 1987; Gasse & Van Campo, 1998; Virah-Swamy et al., 2010), especially near the expected age of human colonization [c. two thousand years ago (ka); Dewar & Wright, 1993; Dewar et al., 2013; Pierron et al., 2017]. Megafaunal isotope records also support Holocene expansions of C_4 grasslands but show a vegetative mosaic of C_4 grasses and C_3 vegetation was at least present (Crowley et al., 2021) near the last glacial maximum (LGM; c. 26.5-19 ka). Reconstructing spatio-temporal vegetative dynamics beyond the LGM begin to approach limitations of paleoecological data, but molecular evolutionary approaches can fill the gaps by providing probabilistic inferences through time. Population genetic approaches for reconstruct demographic histories has been successfully applied to lemurs to reconstruct patterns of past forest connectivity (Quéméré et al., 2012; Yoder et al., 2016; Salmona et al., 2017; Teixeira et al., 2021; Tiley et al., 2022) and may similarly provide insights into the Late Pleistocene dynamics of grasses between the LGM and Last Interglacial period (LIG; c. 132-112 ka) or older. This strategy has been used to reconstruct vegetative dynamics of temperate Japanese grasslands from forb species (Yamaura et al., 2019).

An array of tools now exists to estimate the effective population size (N_e) of species through time. As generating genomic data for non-model groups has become more accessible, population genomic methods that utilize precisely estimated site frequency spectra (SFS; Nielsen et al., 2000; Gutenkunst et al., 2009; Excoffier et al., 2013; Liu & Fu, 2020; Blischack et al., 2022) have become a valuable tool for reconstructing the demographic history of natural populations. There are still barriers to generating population genomic data though, either due to the lack of suitable tissue, an appropriate reference genome, or costs. Target-enrichment

libraries have been transformative for the analysis of plant phylogenetic relationships (e.g.

Breinholt et al., 2021; Baker et al., 2022), but the population genetic utility of such data are not

well-explored. A survey of taxa with low sample sizes suggested population genetic parameters

estimated from target-enrichment data could be unreliable, in addition to violating assumptions

of the neutral coalescent model that underpins most tests of selection and demographic change

(Slimp et al., 2021). However, the relatively long loci recovered by target-enrichment (or

HybSeq) methods could be appropriate for a demographic modeling approach, the extended

Bayesian skyline plot (EBSP; Heled & Drummond, 2008), which has been successfully applied

to a range of conservation genetic questions despite some model violations. We are interested

in exploring the potential of target-enrichment data for population or conservation genetic

insights from plant groups that may be lacking genomic resources.

Here, we examined the population genetics of the grass species *Loudetia simplex*

(Tristachyideae: Panicoideae: Poaceae). It is largely dominant in Madagascar's Central

Highlands (Koechlin, 1993; Hagl et al., 2021; Fig. 1a) and is notable for its fire- and grazing-

adapted functional traits (Solofondranohatra et al., 2018). Population genetic analyses of

microsatellites suggest Malagasy *L. simplex* has been isolated from mainland African

populations with no detectable level of gene flow, and that there is additional population

structure between the northern and southern extents of the species range across the Central

Highlands of Madagascar (Hagl et al., 2021). Here, we use a subset of individuals analyzed by

Hagl et al. (2021) along with a new sample from the center of the *L. simplex* distribution to

explore the population genetic utility of target-enrichment data and its implications for the natural

history of Madagascar's grasslands. Because Malagasy *L. simplex* are putative polyploids

(tetraploids and hexaploids; Hagl et al., 2021), we developed novel bioinformatic tools for the

processing and analysis of polyploid data, integrated into the PATÉ allele phasing pipeline (Tiley

et al., 2021).

Materials and Methods

Sampling and Sequencing

Target-enrichment data was generated for 24 individuals of *Loudetia simplex* and one outgroup,

Tristachya nodiglumus (Supplementary Table S1). Twenty individuals were selected to best

represent Madagascar from available collections as a single unstructured population and avoid

complications of population substructure and low sample sizes in downstream analyses (Fig.

1b). Genomic DNA samples were used from Hagl et al. (2021) while new extracts used a

modified CTAB protocol (Doyle & Doyle, 1987) from silica-dried leaf tissue. Genomic DNA was

then sent to Rapid Genomics (Gainesville, FL, USA) where sample enrichment and sequencing

was performed with a bait kit developed for angiosperms following Breinholt et al. (2021). All

sample metadata is provided in Supplementary Table S1.

Sequence Assembly and Processing

Initial contigs were assembled with HybPiper (Johnson et al., 2016) with default settings. We

then used HybPiper's supercontigs as reference sequences for PATÉ (Tiley et al., 2021; Crowl

et al., 2022) to genotype individuals, filter low-quality variants with GATK v4.2.0 (McKenna et

al., 2010; DePristo et al., 2011), and phase haplotype sequences using HPOP-G (Xie et al.,

2016). Ploidy information for individuals (Supplementary Table S1) was based on previous

microsatellite allele count data (Hagl et al., 2021); however, we also implemented a statistical

test to infer ploidy level in PATÉ based on mixtures of distributions from allele balance data,

similar to other approaches that use the number of reads supporting an alternate allele over

total reads to infer ploidy (Weiß et al., 2018). We extended the mixture models of Tiley et al.

(2018) to determine the ploidy of a sample with the model weights as a selection criterion

(Supplementary Methods: *Ploidy Test*). PATÉ control files with all filtering options for GATK are

provided in the Supplementary Data (Dryad X).

To investigate nucleotide variation and the potential population genetic value of non-coding

versus coding regions of target-enrichment loci, we identified the following regions:

1) *Left Flank* – Non-coding sequence upstream of the 5' start codon or splice site

2) *Core* – The coding region targeted by probes for sample enrichment

3) *Right Flank* – Non-coding sequence downstream of the 3' stop codon or splice site

Identification of individual regions was automated (https://github.com/gtiley/Locus-Splitter)

based on the tblasx program from BLAST v2.10.0 (Camacho et al., 2009) to find in-frame

alignments between our assembled sequences and the original transcriptome loci used for

probe development. Individual regions were aligned with MAFFT v7.471 (Katoh & Standley,

2013). The aligned regions were then concatenated together to make whole-locus alignments.

Because we are interested in investigating the analytic impact of ploidy in sequence data, we

created three datasets from the phased ploidy-aware data:

1) *One Haplotype (OH)* – Only one phased haplotype sequence was randomly selected from

each individual for each locus.

2) *Ambiguous Genotype (AG)* – The genotypic information for an individual is represented as a

single sequence with IUPAC ambiguity codes for biallelic sites. Analyses that treat IUPAC

codes as missing data were randomly resolved into two sequences.

3) *Phased Haplotypes (PH)* – All phased haplotype sequences for an individual are used.

We expect the OH data to be representative of majority consensus sequences (e.g. HybPiper

supercontigs) since our target-enrichment loci are anchored in single exons and we anticipate

that any risk of assembling chimeric sequences from such data without paralog warnings is low.

Any site genotyped as polyallelic was assumed to be an error and treated as missing sequence

across all data sets. All expected copies of a locus are represented even if they are invariant.

For example, some individuals in the PH data may have six identical sequences at a locus if

they are hexaploids and every site is homozygous. While the ploidy at an individual locus may

vary, this is a simplifying assumption to preserve expected allele frequency information.

Phylogeny estimation

A species tree was estimated for all *L. simplex* individuals using whole-locus sequences for the

OH, AG, and PH datasets with BPP v4.6.2 (Flouri et al., 2020). We chose a full-likelihood

multispecies coalescent (MSC; Rannala & Yang, 2003) model since it is robust to low-

information gene tree estimates (Xu & Yang, 2016). Markov chain Monte Carlo (MCMC)

analyses collected 10,000 posterior samples, saving every 10 samples after a 10,000

generation burnin. Model and prior specifications are provided in the supplementary information

(Supplementary Methods: *BPP Prior Specification*). The starting tree for our Bayesian species

tree analyses was based on a concatenated maximum likelihood (ML) estimate from the OH

data with IQTREE v2.1.2 (Minh et al., 2020). We used unpartitioned analyses with

ModelFinderPlus (Kalyaanamoorthy et al., 2017) for the OH, AG, and PH data to estimate ML

evolutionary distances between individuals, with random concatenation of haplotypes within

individuals.

Population structure

Potential population structure within our sample of *L. simplex* from Madagascar was explored

with STRUCTURE v2.3.4 (Pritchard et al., 2000). For each dataset, we sampled only one

biallelic SNP per locus, if present, and required a minimum minor allele count of three with no

more than 25% missing data. To explore uncertainty due to sampling error of sites in a target-

enrichment dataset that is relatively small compared to contemporary population genomic data,

we generated 10 datasets with random SNP sampling. The number of clusters that best satisfy

Hardy-Weinberg assumptions was determined with the ΔK method (Evanno et al., 2005).

Principal component analyses (PCAs) were conducted on the OH, AG, and PH datasets with

adegenet v2.1.3 (Jombart & Ahmed, 2011) by compressing allele counts to the same dimension

as the lowest ploidy level present. For example, a hexaploid with three reference and three

alternate alleles would be scored as a 2 on a scale from 0 to 4, to be comparable to tetraploids

in the dataset. Additional details on allelic imbalances would be lost as, for example, hexaploids

with either one or two alternate alleles would be represented as a 1 on the tetraploid scale.

Scripts for generating PCA or other clustering-ready matrices from fasta files are available on

GPT's GitHub (https://github.com/gtiley/fasta-conversion). Additional details about compression

of PCA data matrices can be found in the Supplementary Material (Supplementary Methods:

PCA Input Data). The presence of isolation-by-distance as an explanation for observed patterns

of genetic structure and ordination was tested with the patristic distances from the ML analyses

or the uncorrected pairwise distances (p-distances) regressed against the natural logarithm of

the great circle distance between individuals. A Mantel test was performed for isolation-by-

distance analyses in the ape v5.6.2 R package (Paradis & Schliep, 2019).

Population genetic summary statistics

The pairwise nucleotide diversity (π), number of segregating sites (S), and Tajima's D were

calculated across locus regions and datasets with the PopGenome v2.7.5 R package (Pfeifer et

al., 2014). The significance of Tajima's D statistics was determined by 100 standard coalescent

simulations with ms (Hudson, 2002). To estimate 95% confidence intervals on summary

statistics, we generated 200 bootstrap replicates of each alignment. The relative estimation

error between regions and datasets was compared using the root squared mean error (RSME)

from the bootstrap sample. The scaling and interpretation of parameters are described in the

supplementary material (Supplementary Methods: *Summary Statistics*). We tested for

differences in the distribution of summary statistics among locus regions (left flanks, cores, and

right flanks) and data types (OH, AG, and PH) with Wilcoxon Rank-Sum tests. We tested for

biases in Tajima's D results among locus regions and data types with X^2 tests.

Demographic History

We chose the extended Bayesian skyline plot (EBSP, Heled & Drummond, 2008) as the most

appropriate model of N_e change through time for our sample of target enrichment data. The

model assumes N_e changes coincide with coalescent events, and although an unrealistic

assumption (Yang, 2014), is common. Here we assumed a strict molecular clock underlying all

lineages but allowed rates of evolution to vary among loci. One locus was fixed to the clock rate

and all other locus rates were relative. Rates of molecular evolution among plants are especially

complicated, as there can be life histories with variable reproductive strategies, unknown

generation times, and overlapping generations. Thus, we relied on an estimate of the neutral

substitution rate at third-position synonymous sites from grass genomes (Christin et al., 2014)

and constructed a vague prior around this mean rate, such that any biologically plausible

uncertainty in the rate of evolution is reflected in our posterior sample. It is a tenuous rate

estimate at best, since it remains unknown whether polyploid *L. simplex* largely propagates

clonally, through apomixis, or through sexual reproduction. The MCMC analyses for the OH,

AG, and PH data collected 20,000 samples after a burnin of 50,000,000 and sample interval of

10,000. Two replicates were run for each analysis to evaluate convergence and mixing was

assessed with Tracer v1.7 (Rambaut et al., 2017). All model and prior specifications are

provided in the supplementary information (Supplementary Methods: *EBSP Prior Specification*)

and are replicable from XML files on Dryad (X).

Adequacy of Demographic Models

To explore if EBSP models could reliably differentiate between Late Pleistocene versus

Holocene population size increases, we evaluated the accuracy of EBSP inferences with

simulations that captured some features of our target-enrichment data. We simulated 57 loci

that were 1000 bp in length for 20 haplotype individuals 100 times with ms (Hudson, 2002) and

seq-gen (Rambaut & Grass, 1997) for four scenarios: 1) a constant N_e of 100,000, 2) an N_e

increase from 10,000 to 100,000 individuals roughly coinciding with the LGM at 20,000 years in

the past, 3) the same ten-fold increase but at 2,000 years in the past to reflect anthropogenic

expansion, and 4) a scenario combining both LGM and anthropogenic effects by exponentially

increasing N_e from 10,000 to 100,000 between 20,000 years ago and the present. Each

simulated dataset was then evaluated with the same EBSP model using a fixed clock used for

our empirical analyses, except the MCMC algorithm collected 10,000 posterior samples with a

sample interval of 1,000 after a 100,000 sample burnin. Commands for simulations are given in

the supplementary information (Supplementary Methods: *Simulating Demographic Histories*).

Results

Sequencing statistics

The probe set targeted 408 loci from which an average of 327 loci were assembled by HybPiper

(Supplementary Table S2). These were supported by an average of 217 read pairs per locus.

Although there was variable read depth across loci and individuals, the number of assembled

loci was significantly correlated with read depth (Pearson's correlation coefficient = 0.57, p-

value = 0.002). After joint-genotyping all individuals as diploids, normal mixture models

suggested all individuals were hexaploids, with the exception of two individuals from South

Africa (Supplementary Table S3). Some individuals were inferred as pentaploids, but since this

was unexpected based on previous observations from microsatellites (Supplementary Table S1)

these cases were assumed to be hexaploids. Although available evidence leaves the ploidy of

some individuals ambiguous and raises skepticism of the inference methods, genotyping and

phasing of individuals proceeded with the ploidy suggested by our mixture models. We inferred

an average of 247 loci with allelic variation, which had a heterozygosity of 0.012 variants per

base. After reducing the number of loci with BLAST clustering to avoid tight linkage, a final 301

loci were retained from the OH, AG, and PH datasets for downstream analyses.

Phylogeny Estimation

Species trees revealed all Malagasy *L. simplex* form a well-supported clade (Supplementary

Fig. S1), but there was disagreement in the biogeographic history within Madagascar based on

the treatment of ploidy. Individuals in northern Madagascar formed a clade with a posterior

probability of 0.9 or greater across the OH, AG, and PH analyses. However, the individuals from

central and southern Madagascar were paraphyletic in the OH and AG analyses

(Supplementary Figs. 1a and 1b). A clade of central and southern individuals was recovered in

the PH analysis, but with a weak posterior probability of 0.64 (Fig. 1c; Supplementary Fig. 1c).

Individuals from the center of the *L. simplex* distribution appear to be the earliest diverging

lineages in the OH and AG analyses (Supplementary Figs. 1a and 1b), or with respect to the

sample of central and southern individuals in the PH analysis (Fig. 1c; Supplementary Fig. 1c).

Population sub-structure in Madagascar

STRUCTURE analyses recovered a variable number of Hardy-Weinberg groups across the ten

datasets that randomly sampled one non-singleton biallelic SNP per locus (Supplementary

Table S4). For the OH data, we found a *K* of two, three, or four in eight, one, and one of the

respective replicates. For AG data, the optimal number of clusters was more ambiguous with a

K of two, three, and four recovered for four, five, and one replicates. PH data found *K* of two and

three for eight and two replicates. We note that the method of Evanno et al. (2005) cannot infer

if one panmictic population is the best model, and while there is reasonable geographic

structure at K of two weighted across biological and technical replicates (Fig. 2a) the non-zero

admixture between individuals at the most extreme ends could indicate a single population with

a cline of genetic variation. This is further supported by the lack of distinct cluster assignments

at higher values of K despite some evidence for additional population structure (Supplementary

Fig. S2). A weak signal of isolation-by-distance was detected when regressing genetic distance

against spatial distance (Fig. 2b). Notably, the AG data inferred much higher pairwise distances

between individuals; although, the relative differences and overall trend were similar. The same

patterns can be observed when using patristic distances from ML phylogenetic estimates,

although the disparity between AG versus OH or PH distances is reduced, as is the percent of

genetic variation explained by spatial distances (Supplementary Fig. S3).

PCAs reflected findings from STRUCTURE analyses and reinforce the presence of a genetic

cline (Fig. 3). For OH data, where only one reference or alternate allele was observed per

individual, Malagasy *L. simplex* is clearly distinguished from mainland African individuals (Fig.

3a), but there is no clear separation of individuals based on geographic location (Fig. 3b). PCAs

of data that accounts for heterozygosity but not dosage recovered a single Malagasy versus

mainland African pattern (Fig. 3c), PCAs reflect a north-to-south gradient of genetic variation

regardless of whether allelic dosage was treated as hexaploid (Fig. 3d), tetraploid (Fig. 3e), or

diploid (Fig. 3f). The total percent of variance explained by principal component axis 1 (PC1)

and axis 2 (PC2) increased as allelic dosage was compressed from a hexaploid to a diploid

state.

Variation in summary statistics

There were significant differences in the distributions of π, S, and Tajima's D among locus

regions and between the OH, AG, and PH data (Fig. 4). For π, there was always a difference

between flanking regions and the exon core (Supplementary Table S5), with the cores

surprisingly having more variation. Differences were also observed between the flanking regions

and cores for S and Tajima's D, except when comparing the left flanks and cores for OH data

(Supplementary Table S5). The only case of a difference between the left and right flanks was

when estimating π for AG data (Supplementary Table S5). The OH and PH data resulted in

similar estimates of π; however, the AG and PH data had similar distributions of S

(Supplementary Table S5). Any differences in Tajima's D statistics were largely explained by an

over-representation of values less than negative two, indicating positive selection or population

size expansion, in the OH data and among core regions across datasets (Table 1).

We attempted to explore biases in summary statistic estimates due to sampling error of sites

through bootstrapping the OH (Supplementary Fig. S4), AG (Supplementary Fig. S5), and PH

data (Supplementary Fig. S6). Since mutations are assumed to arise through a Poisson process

and exponential distributions of measures of genetic variation, such as π and S are often

observed, the bootstrap confidence intervals are not a reasonable measure of uncertainty.

However, we used the bias, difference between the bootstrap mean and observed estimate, and

the RSME of bootstrap replicates with respect to the observed estimate, to gain some general

insights. Notably, the bias of π is larger and much noisier in the AG data (Supplementary Fig.

S5) compared to the OH (Supplementary Fig. S4) and PH (Supplementary Fig. S6) data. Even if

the bias for a whole-locus is low, there can be high variability between regions. For the

observed values, estimates are not always proportional among regions either. The RSME of an

estimate tends to decrease along with genetic variation, implying some role of genotyping error

or rare variants and singletons inflating values at the higher end of the spectrum. Loci with very

low levels of genetic variation can also be problematic though, as the RSME for flanking regions

increases towards the tail of the distribution, creating opportunity for sampling error to wildly

swing estimates regardless of ploidy (Supplementary Figs. S4-S6). The number of segregating

sites increased with ploidy of the data, but π was relatively stable. However, differences

between π and S will affect estimates of Tajima's D, and this could affect interpretations of

demography or selection acting on loci; although, the majority of loci and their regions were not

significantly different from neutral expectations given the simulation-based test statistics

(Supplementary Table S6).

Population Size Change through time

After filtering for longer loci with low amounts of missing data, 57 loci were retained for EBSP

analyses. Reducing our data to a subset of more informative loci was done for computational

convenience, because EBSP models are parameter-rich and require long MCMC runs for

convergence. Runs converged for each data type for both analyses with a fixed rate of

molecular evolution (Fig. 5) and variable rate (Supplementary Fig. S7). All analyses suggested

the *L. simplex* N_e increased through the Late Pleistocene, well before the LIG, approximately

one million years ago. Between the LGM and period of potential human colonization of

Madagascar in the Holocene, N_e had already reached present-day estimates. N_e estimates

through the Holocence were only possible for the PH data, as accounting for the within-

individual variation should have captured more recent coalescences. Simulations cast doubt on

the seriousness with which EBSP results can be interpreted through the Holocene though. For

example, when N_e was truly constant through time, some individual EBSP analyses would

suggest recent N_e increases; although, the mean estimate across simulations recovered the

constant population scenario (Fig. 6a). This problem was exaggerated in the presence of model

over-fitting (Supplementary Fig. S8). In scenarios with recent N_e increases, the contemporary N_e

was underestimated but a trend indicating a recent N_e increase was detectable (Figs. 6b-6d).

However, simulations indicated it would not be possible to differentiate between N_e increases

that occurred during the LGM (Fig. 6b) versus the Late Holocene (Fig. 6c), let alone some

combination of paleoclimate and anthropogenic effects (Fig. 6d). Interpreting these recent N_e

increases from EBSP analyses becomes impossible with model over-fitting (Supplementary Fig.

S8).

Discussion

Robustness of analyses across loci and ploidy

Although transformative for species-level phylogenomic studies (Breinholt et al., 2021; Baker et

al., 2022), the robustness of population genetic estimates for a sample of closely-related

individuals was largely unexplored (but see Slimp et al., 2021). Some of the most important

measures of genetic variation for populations were susceptible to sampling error, especially in

the flanking intron regions (Supplementary Figs. S4-S6). While estimates from cores showed

less error, they will lead to some violations of population genetic methods that assume evolution

consistent with neutral theory, given potential signatures of purifying selection (Table 1). This

creates some conflict, since most of the information is contained within the exon cores of target-

enrichment loci (Fig. 4), presumably due to read coverage. Because Tajima's *D* alone cannot

disentangle demography from selection, some judgment calls and tolerance for downstream

model violations are necessary. The majority of loci examined were consistent with neutral

expectations based on simulations (Supplementary Table S6), and the proportion of loci with

significant Tajima's *D* statistics were exaggerated when ignoring heterozygosity (Table 1).

Predicting the effects of assumptions about genotype on estimates of genetic variation and

downstream analyses are not always straightforward. For example, the pairwise distances (Fig.

2b) or evolutionary distances (Supplementary Fig. S3) between individuals was higher when

assuming all individuals were diploid (AG) compared to assuming all individuals were haploid

(OH) or genotyped at their anticipated ploidy level (PH). If we were willing to ignore the absolute

estimates of genetic distance, the interpretation of isolation-by-distance analyses was the same

across the OH, AG, and PH data. This implies that biologically meaningful conclusions can arise

even with uncertainty in ploidy and phasing. Interestingly, treating all individuals as diploid was

sufficient for STRUCTURE to recover northern and southern clusters with some admixed central

individuals of varying proportions (Fig. 2a; Supplementary Fig. S2). Similarly, PCAs reflected the

geographic distribution of our *L. simplex* samples in the AG and PH data with, surprisingly, the

AG data explaining more of the observed genetic variance (Fig. 3f) than the PH data (Fig. 3d).

This may be due to genotyping error at higher ploidy levels caused by read stochasticity and a

flat genotype likelihood surface (Blischak et al., 2016) or caused by incorrect ploidy inferences.

A similar mixture model approach to the one implemented here (Weiß et al., 2018) has been

shown to have good performance based on comparisons of allele balance distributions from

target-enrichment data and flow cytometry-based ploidy estimates (Viruel et al., 2019). We are

cautiously optimistic about the prospects of direct ploidy estimation from sequence data,

because likelihood-based criteria can be prone to overfitting (Tiley et al., 2018). However, our

ploidy estimates are generally consistent with the previous inferences from microsatellite allele

counts (Hagl et al., 2021; Supplementary Table S1; Supplementary Table S3). A positive insight

from our comparisons across ploidy levels is that if unknown individuals are genotyped as

diploid, there will be sufficient information about population structure, and that ploidy estimation

should not be a barrier to investigations of natural plant populations with little *a priori* cytological

or genetic knowledge.

Limits to differentiating anthropogenic and paleoclimate N_e change

Large population genomic studies in plants that can leverage demographic estimates from SFS

(e.g. Excoffier et al., 2013; Liu & Fu, 2020) are commonly focused on temperate regions (e.g.

Yamaura et al., 2019) and species exploited for energy or agriculture (e.g. Huang et al., 2014;

Wang et al., 2017). Thus, it is tempting to leverage models that use gene trees as data such as

the EBSP (Heled & Drummond, 2008) or other extensions of skyline plots (Pybus et al., 2000),

since they could be paired with relatively low-cost target-enrichment strategies that yield multi-

locus nuclear data while stretching material quality.

While our analyses provide valuable insights into the ancient origins of *L. simplex* dominated

grasslands, both our empirical and simulated analyses show we cannot differentiate between N_e

increases that occurred during human colonization or paleoclimate change near the LGM. The

change in vegetative composition towards more C_4 grass species in central Madagascar is well-

documented (Burney, 1987; Gasse & Van Campo, 1998; Virah-Swamy et al., 2010; Crowley et

al., 2021). It is unlikely that these increases in C_4 grasses did not involve *L. simplex*; although,

an increase in census size may not be reflected by N_e (Fig. 5; Supplementary Fig. S7), the

expected population size under a neutral coalescent model. Our simulations show that detecting

these recent increases is difficult, and that we cannot differentiate between a ten-fold increase

that occurred 200 generations (2 ka) or 2000 generations (20 ka; Fig. 6). Diagnosing a true N_e

change becomes impossible in the presence of model overfitting (Supplementary Fig. S8) and

diagnosing these problems in empirical data can be difficult aside from convergence

assessments.

These limitations do not necessarily disagree with an assessment of EBSP models with a larger

genomic dataset from endemic palm species in Madagascar (Helmstetter et al., 2021).

Helmstetter et al. (2021) found that the estimated N_e trends were valuable predictors of IUCN

risk assessments, and that population decreases could be detected at anthropogenic time

scales. However, it should be easier to detect population declines than population increases in a

system that already had a large degree of standing genetic variation, such as *L. simplex*. This is

because the time until the most recent common ancestor of a genealogy will be much more

recent in a small population than a large one (e.g. see equation 2.13 from Gillespie 2004).

Analyses of demographic models with genomic SFS should be able to increase resolution at

recent anthropogenic time scales (Patton et al., 2019) and differentiate anthropogenic from

paleoclimatic demographic effects (Tiley et al., 2022), but this would require much more

extensive sampling and sequencing effort. Additionally, our analyses treat all individuals as a

single population, and since we detected some level of population structure, we cannot rule out

their confounding effects on analyses of N_e through time (Heller et al., 2013; Mazet et al., 2016)

or other confounding affects such as clonal growth and overlapping populations. Population-

level sampling of multiple sites throughout a species range should be used to control for

population structure, while accounting for localized effects such as agriculture and grazing

versus regional climate effects.

New insights into Loudetia simplex evolution and policy implications

Our analyses yielded new findings about the evolution of *L. simplex* that are potentially

consequential for the broader understanding of Malagasy grasslands. First, our species trees

(Supplementary Fig. S1) indicate that individuals from what we call central Madagascar – a

loose term meant only to indicate individuals from the center of the sampled *L. simplex*

distribution – represent the ancestral distribution. It is likely that *L. simplex* dominated

grasslands have origins in some of the higher elevation regions of the Central Highlands. For

example, MSV1564 comes from the Itremo protected area (Fig. 1b; Supplementary Table S1),

and further investigations of this region should be productive for characterizing natural

grasslands in Madagascar. The hypothesis of central high-altitude origins is also supported by

our analyses of population structure (Fig. 2a; Supplementary Fig. S2) and the weak but

significant pattern of isolation-by-distance (Fig. 2b; Supplementary Fig. S3).

Our models of N_e change through suggests that *L. simplex* grasslands were already extensive

(Fig. 5; Supplementary Fig. S7) well before any hypothesized human presence in Madagascar

(c. 10 ka; Hansford et al., 2018) let alone human colonization (c. 2 ka; Dewar and Wright, 1993;

Dewar et al., 2013; Pierron et al., 2017). We found an approximate ten-fold increase between

one MA and the LIG which is beyond the limitations of available stratigraphic records. Increases

in grass and especially C_4 grass relative abundance through the Holocene are well-

characterized (Burney, 1987; Gasse & Van Campo, 1998; Virah-Swamy et al., 2010; Railsback

et al., 2020; Crowley et al., 2021); however, we showed that limitations of our model do not

permit us to make inferences at these very recent timescales (Fig. 6). Strong assumptions about

the mutation rate and generation time to calibrate analyses to absolute time and N_e were

necessary, but our interpretations of results are within the biologically plausible uncertainty of

these parameters. Even if the per-year substitution rate was overestimated by a factor of ten,

the N_e increase would still be constrained to the Late Pleistocene, occurring between the LIG

and LGM. These analyses, especially in the context of a previous investigation of population

structure (Hagl et al., 2021), suggest that *L. simplex* dominated grasslands have pre-human

origins and likely experienced some N_e increase coincident with Pleistocene climate change.

Tree planting programs have been deployed to combat the measurable habitat loss and

deforestation in Madagascar since the 1950s (Green & Sussman, 1990; Harper et al., 2007;

Vieilledent et al., 2018). The results of our study strongly imply that these efforts should be

cautious of treating *L. simplex* dominated grasslands as degraded forests, and at least a mosaic

of *L. simplex* and native woody species would be more appropriate.

Potential use cases of target-enrichment data for conservation genetics

One application of target-enrichment data going forward could be analyses of population

structure and population assignment of unknown individuals. For example, many conservation

programs operating at scale rely on reduced-SNP panels to determine provenance of an

individual or sample (e.g. Bourgeois et al., 2018; Stronen et al., 2022). Developing SNP panels

requires at least some *a priori* genomic resources, so target-enrichment libraries could provide

feasible approaches in the absence of such references and practical improvements over

existing barcoding strategies for identifying, for example, Madagascar's rosewoods (Hassold et

al., 2016); although acquiring viable DNA samples from woody tissue would remain a barrier

(Jiao et al., 2018). We anticipate limited use cases of target-enrichment data in population

genetics going forward as reference genome assembly is becoming more accessible through

decreasing long-read costs, and less complicated library preparation protocols for whole-

genome resequencing will become preferential to any target-enrichment strategy. However, the

population genetic inferences from target-enrichment data should be biologically meaningful

when available.

A practical need for plant population genomics in Madagascar

There are few studies of plant population genetics from the Central Highlands and those are

limited to AFLP (Gardiner et al., 2017) or microsatellite data (Salmona et al., 2020; Hagl et al.,

2021). Demographic analyses from the endemic olive tree *Noronhia lowryi* suggested recent

population declines of the woody species, but it is unclear if these declines represent

anthropogenic or paleoclimatic change near the LGM (Salmona et al., 2020). While our

analyses of *L. simplex* are also ambiguous regarding this window of anthropogenic effects on

the extent of grasslands, we can at least conclude that the *L. simplex* N_e was quite large.

Contemporary population genomic analyses will be necessary to provide statistical power to

differentiate between competing demographic hypotheses, not only for grasses such as *L.*

simplex but also species representing plant diversity across families and life-history. A better

reconstruction of the natural history of Madagascar's grasslands is not only necessary for

understanding diversity within and outside of the Central Highlands (Yoder et al., 2016; Burbrink

et al., 2019; Everson et al., 2020), but also to make effective land management decisions for

tree planting programs.

Acknowledgements

GPT was supported by a Marie Curie Individual Fellowship (MSCA-IF-EF-ST 101026923). GB is

supported by LabEx TULIP (ANR-10-LABX-0041). CERL was supported by a The Royal Society

International Collaboration Award. Thank you to MEDD Madagascar, Madagascar National

Parks, Isalo MNP, and Itremo NPA for supporting our research and exportation permits in 2014

and 2017, and also to COSTECH Tanzania for their support for the 2014 collection permits for

MSV and WRQL. This work would not have been possible without Stuart Cable and everyone at

the Kew Madagascar Conservation Centre. We thank Emily Lockwood for DNA extractions, and

Lorena Endara and J. Gordon Burleigh for coordinating library preparation and sequencing.

Author Contributions

GPT, ADY, and MSV conceived the study. WRQL, CLS, CERL, GB and MSV collected

samples. GPT and TOMA curated data. GPT and AAC analyzed data. GPT wrote the first draft.

All authors revised the manuscript and approved the final version.

Data Availability Statement

Scripts for analyses and input files are available through the Dryad Digital Repository:

http://dx.doi.org/10.5061/dryad.[NNNN]. Additional scripts are available on GPT's GitHub for

identifying exon-intron boundaries from target enrichment data (https://github.com/gtiley/Locus-

Splitter) generating matrices for PCAs from fasta files (https://github.com/gtiley/fasta-

conversion) and the ploidy test implemented through PATÉ (https://github.com/gtiley/Phasing).

Raw sequence data is available via the NCBI short read archive (SRA) database, with individual

SRA and BioSample identifiers in Supplementary Table S1. Raw sequence data is associated

with NCBI BioProject PRJNA952819.

Conflict of Interest Statement

The authors declare no conflict of interest.

References

Baker, W. J., Bailey, P., Barber, V., Barker, A., Bellot, S., Bishop, D., ... & Forest, F. (2022). A comprehensive phylogenomic platform for exploring the angiosperm tree of life. *Systematic Biology*, *71*(2), 301-319. https://doi.org/10.1093/sysbio/syab035

Blischak, P.D., Kubatko, L.S., Wolfe, A.D. (2016_. Accounting for genotype uncertainty in the estimation of allele frequencies in autopolyploids. *Molecular Ecology Resources, 16*(3), 742-754. https://doi.org/10.1111/1755-0998.12493

Blischak, P.D., Sajan, M., Barker, M.S., Gutenkunst, R.N. (2022). Demographic history inference and the polyploid continuum. *bioRxiv*, https://doi.org/10.1101/2022.09.15.508148

Bond, W. J., Silander Jr, J. A., Ranaivonasy, J., & Ratsirarson, J. (2008). The antiquity of Madagascar's grasslands and the rise of C4 grassy biomes. *Journal of Biogeography*, *35*(10), 1743-1758. https://doi.org/10.1111/j.1365-2699.2008.01923.x

Bond, W. J., Silander, J. A., & Ratsirarson, J. (2023). Madagascar's grassy biomes are ancient and there is much to learn about their ecology and evolution. *Journal of Biogeography*, *50*(3), 614-621. https://doi.org/10.1111/jbi.14494

Bourgeois, S., Senn, H., Kaden, J., Taggart, J. B., Ogden, R., Jeffery, K. J., ... & McEwing, R. (2018). Single-nucleotide polymorphism discovery and panel characterization in the African forest elephant. *Ecology and Evolution*, *8*(4), 2207-2217. https://doi.org/10.1002/ece3.3854

Breinholt, J. W., Carey, S. B., Tiley, G. P., Davis, E. C., Endara, L., McDaniel, S. F., ... & Burleigh, J. G. (2021). A target enrichment probe set for resolving the flagellate land plant tree of life. *Applications in Plant Sciences*, *9*(1), e11406. https://doi.org/10.1002/aps3.11406

Burbrink, F. T., Ruane, S., Kuhn, A., Rabibisoa, N., Randriamahatantsoa, B., Raselimanana, A. P., ... & Raxworthy, C. J. (2019). The origins and diversification of the exceptionally rich gemsnakes (Colubroidea: Lamprophiidae: Pseudoxyrhophiinae) in Madagascar. *Systematic Biology*, *68*(6), 918-936. https://doi.org/10.1093/sysbio/syz026

Burney, D. A. (1987). Late Holocene vegetational change in central Madagascar. *Quaternary Research*, *28*(1), 130-143. https://doi.org/10.1016/0033-5894(87)90038-X

Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., & Madden, T. L. (2009). BLAST+: architecture and applications. *BMC bioinformatics*, *10*, 1-9. https://doi.org/10.1186/1471-2105-10-421

Christin, P.-A., Spriggs, E., Osborne, C. P., Strömberg, C. A., Salamin, N., & Edwards, E. J. (2014). Molecular dating, evolutionary rates, and the age of the grasses. *Systematic biology*, *63*(2), 153-165. https://doi.org/10.1093/sysbio/syt072

Crowl, A. A., Fritsch, P. W., Tiley, G. P., Lynch, N. P., Ranney, T. G., Ashrafi, H., & Manos, P. S. (2022). A first complete phylogenomic hypothesis for diploid blueberries (Vaccinium section Cyanococcus). *American Journal of Botany*, *109*(10), 1596-1606. https://doi.org/10.1002/ajb2.16065

Crowley, B. E., Godfrey, L. R., Hansford, J. P., & Samonds, K. E. (2021). Seeing the forest for the trees—and the grasses: revisiting the evidence for grazer-maintained grasslands in Madagascar's Central Highlands. *Proceedings of the Royal Society B*, *288*(1950), 20201785. https://doi.org/10.1098/rspb.2020.1785

DePristo, M. A., Banks, E., Poplin, R., Garimella, K. V., Maguire, J. R., Hartl, C., ... & Daly, M. J. (2011). A
framework for variation discovery and genotyping using next-generation DNA sequencing data. *Nature*
genetics, *43*(5), 491-498. https://doi.org/10.1038/ng.806

Dewar, R. E., Radimilahy, C., Wright, H. T., Jacobs, Z., Kelly, G. O., & Berna, F. (2013). Stone tools and
foraging in northern Madagascar challenge Holocene extinction models. *Proceedings of the National*
Academy of Sciences, *110*(31), 12583-12588. https://doi.org/10.1073/pnas.1306100110

Dewar, R. E., & Wright, H. T. (1993). The culture history of Madagascar. *Journal of World Prehistory*, *7*,
417-466. https://doi.org/10.1007/BF00997802

Doyle, J.J., Doyle, J.L. (1987). A rapid DNA isolation procedure for small quantities of fresh leaf tissue.
Phytochemical Bulletin, 19(1), 11-15.

Evanno, G., Regnaut, S., & Goudet, J. (2005). Detecting the number of clusters of individuals using the
software structure: A simulation study. Molecular Ecology, 14(8), 2611–2620.
https://doi.org/10.1111/J.1365-294X.2005.02553.X

Everson, K. M., Jansa, S. A., Goodman, S. M., & Olson, L. E. (2020). Montane regions shape patterns of
diversification in small mammals and reptiles from Madagascar's moist evergreen forest. *Journal of*
Biogeography, *47*(10), 2059-2072. https://doi.org/10.1111/jbi.13945

Excoffier, L., Dupanloup, I., Huerta-Sánchez, E., Sousa, V. C., & Foll, M. (2013). Robust demographic
inference from genomic and SNP data. *PLoS genetics*, *9*(10), e1003905.
https://doi.org/10.1371/journal.pgen.1003905

Flouri, T., Jiao, X., Rannala, B., & Yang, Z. (2020). A Bayesian implementation of the multispecies
coalescent model with introgression for phylogenomic analysis. *Molecular Biology and Evolution*, *37*(4),
1211-1223. https://doi.org/10.1093/molbev/msz296

Gardiner, L. M., Rakotoarinivo, M., Rajaovelona, L. R., & Clubbe, C. (2017). Population genetics data
help to guide the conservation of palm species with small population sizes and fragmented habitats in
Madagascar. *PeerJ*, *5*, e3248. https://doi.org/10.7717/peerj.3248

Gasse, F., & Van Campo, E. (1998). A 40,000-yr pollen and diatom record from Lake Tritrivakely,
Madagascar, in the southern tropics. *Quaternary Research*, *49*(3), 299-311.
https://doi.org/10.1006/qres.1998.1967

Gillespie, J.H. (2004). The coalescent. In, *Population Genetics: A Concise Guide* (pp.40-47). Baltimore,
MD. John Hopkins University Press.

Green, G. M., & Sussman, R. W. (1990). Deforestation history of the eastern rain forests of Madagascar
from satellite images. *Science*, *248*(4952), 212-215. https://doi.org/10.1126/science.248.4952.212

Gutenkunst, R. N., Hernandez, R. D., Williamson, S. H., & Bustamante, C. D. (2009). Inferring the joint
demographic history of multiple populations from multidimensional SNP frequency data. *PLoS*
genetics, *5*(10), e1000695. https://doi.org/10.1371/journal.pgen.1000695

Hackel, J., Vorontsova, M.S., Nanjarisoa, O.P., Hall, R.C., Razanatsoa, J., Malakasi, P., & Besnard, G.
(2018). Grass diversification in Madagascar: In situ radiation of two large C3 shade clades and support
for a Miocene to Pliocene origin of C4 grassy biomes. *Journal of Biogeography, 45*(4), 750–761.
https://doi.org/10.1111/jbi.13147

Hagl, P. A., Gargiulo, R., Fay, M. F., Solofondranohatra, C., Salmona, J., Suescun, U., ... & Vorontsova,
M. S. (2021). Geographical structure of genetic diversity in Loudetia simplex (Poaceae) in Madagascar

and South Africa. *Botanical Journal of the Linnean Society*, *196*(1), 81-99.
https://doi.org/10.1093/botlinnean/boaa098

Hansford, J., Wright, P. C., Rasoamiaramanana, A., Pérez, V. R., Godfrey, L. R., Errickson, D., ... & Turvey, S. T. (2018). Early Holocene human presence in Madagascar evidenced by exploitation of avian megafauna. *Science Advances*, *4*(9), eaat6925. https://doi.org/10.1126/sciadv.aat6925

Harper, G. J., Steininger, M. K., Tucker, C. J., Juhn, D., & Hawkins, F. (2007). Fifty years of deforestation and forest fragmentation in Madagascar. *Environmental conservation*, *34*(4), 325-333.
https://doi.org/10.1017/S0376892907004262

Heled, J., & Drummond, A. J. (2008). Bayesian inference of population size history from multiple loci. *BMC evolutionary biology*, *8*(1), 1-15. https://doi.org/10.1186/1471-2148-8-289

Heller, R., Chikhi, L., & Siegismund, H. R. (2013). The confounding effect of population structure on Bayesian skyline plot inferences of demographic history. *PloS one*, *8*(5), e62992.
https://doi.org/10.1371/journal.pone.0062992

Helmstetter, A. J., Cable, S., Rakotonasolo, F., Rabarijaona, R., Rakotoarinivo, M., Eiserhardt, W. L., ... & Papadopulos, A. S. (2021). The demographic history of Madagascan micro-endemics: have rare species always been rare?. *Proceedings of the Royal Society B*, *288*(1959), 20210957.
https://doi.org/10.1098/rspb.2021.0957

Huang, P., Feldman, M., Schroder, S., Bahri, B. A., Diao, X., Zhi, H., ... & Kellogg, E. A. (2014). Population genetics of Setaria viridis, a new model system. *Molecular Ecology*, *23*(20), 4912-4925.
https://doi.org/10.1111/mec.12907

Hudson, R. R. (2002). Generating samples under a Wright–Fisher neutral model of genetic variation. *Bioinformatics*, *18*(2), 337-338. https://doi.org/10.1093/bioinformatics/18.2.337

Jiao, L., Yu, M., Wiedenhoeft, A. C., He, T., Li, J., Liu, B., ... & Yin, Y. (2018). DNA barcode authentication and library development for the wood of six commercial Pterocarpus species: the critical role of xylarium specimens. *Scientific Reports*, *8*(1), 1945. https://doi.org/10.1038/s41598-018-20381-6

Johnson, M. G., Gardner, E. M., Liu, Y., Medina, R., Goffinet, B., Shaw, A. J., ... & Wickett, N. J. (2016). HybPiper: Extracting coding sequence and introns for phylogenetics from high-throughput sequencing reads using target enrichment. *Applications in plant sciences*, *4*(7), 1600016.
https://doi.org/10.3732/apps.1600016

Jombart, T., & Ahmed, I. (2011). adegenet 1.3–1: New tools for the analysis of genome-wide SNP data. *Bioinformatics, 27(*21), 3070–3071. https://doi.org/10.1093/bioin forma tics/btr521

Katoh, K., & Standley, D. M. (2013). MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Molecular Biology and Evolution*, *30*(4), 772-780.
https://doi.org/10.1093/molbev/mst010

Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K., Von Haeseler, A., & Jermiin, L. S. (2017). ModelFinder: fast model selection for accurate phylogenetic estimates. *Nature methods*, *14*(6), 587-589.
https://doi.org/10.1038/nmeth.4285

Koechlin, J. (1993). Grasslands of Madagascar. In R. T. Coupland (Ed.), *Natural Grasslands: Eastern Hemisphere and Résumé* (pp. 291-301). Amsterdam, Netherlands. Elsevier.

Lehmann, C. E. R., Solofondranohatra, C., Morton, J. A., Phelps, L. N., Ralimanana, H., Razanatsoa, J., Rakotoarimanana, V., Vorontsova, M.S. (2022). The grassy ecosystems of Madagascar. In S. M.

725   Goodman (Ed.), *The new natural history of Madagascar* (pp. 152-168). Princetonm NJ. Princeton
726   University Press.
727
728   Liu, X., & Fu, Y. X. (2020). Stairway Plot 2: demographic history inference with folded SNP frequency
729   spectra. *Genome biology*, *21*(1), 1-9. https://doi.org/10.1186/s13059-020-02196-9
730
731   Mazet, O., Rodríguez, W., Grusea, S., Boitard, S., & Chikhi, L. (2016). On the importance of being
732   structured: instantaneous coalescence rates and human evolution—lessons for ancestral population size
733   inference?. *Heredity*, *116*(4), 362-371. https://doi.org/10.1038/hdy.2015.104
734
735   McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., ... & DePristo, M. A.
736   (2010). The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA
737   sequencing data. *Genome research*, *20*(9), 1297-1303. https://doi.org/10.1101/gr.107524.110
738
739   Minh, B. Q., Schmidt, H. A., Chernomor, O., Schrempf, D., Woodhams, M. D., Von Haeseler, A., &
740   Lanfear, R. (2020). IQ-TREE 2: new models and efficient methods for phylogenetic inference in the
741   genomic era. *Molecular Biology and Evolution*, *37*(5), 1530-1534.
742   https://doi.org/10.1093/molbev/msaa015
743
744   Nielsen, R. (2000). Estimation of population parameters and recombination rates from single nucleotide
745   polymorphisms. *Genetics*, *154*(2), 931-942. https://doi.org/10.1093/genetics/154.2.931
746
747   Paradis, E., & Schliep, K. (2019). ape 5.0: an environment for modern phylogenetics and evolutionary
748   analyses in R. *Bioinformatics*, *35*(3), 526-528. https://doi.org/10.1093/bioinformatics/bty633
749
750   Patton, A. H., Margres, M. J., Stahlke, A. R., Hendricks, S., Lewallen, K., Hamede, R. K., ... & Storfer, A.
751   (2019). Contemporary demographic reconstruction methods are robust to genome assembly quality: A
752   case study in Tasmanian devils. *Molecular Biology and Evolution*, *36*(12), 2906-2921.
753   https://doi.org/10.1093/molbev/msz191
754
755   Pfeifer, B., Wittelsbürger, U., Ramos-Onsins, S. E., & Lercher, M. J. (2014). PopGenome: an efficient
756   Swiss army knife for population genomic analyses in R. *Molecular Biology and Evolution*, *31*(7), 1929-
757   1936. https://doi.org/10.1093/molbev/msu136
758
759   Pierron, D., Heiske, M., Razafindrazaka, H., Rakoto, I., Rabetokotany, N., Ravololomanga, B., ... &
760   Letellier, T. (2017). Genomic landscape of human diversity across Madagascar. *Proceedings of the*
761   *National Academy of Sciences*, *114*(32), E6498-E6506. https://doi.org/10.1073/pnas.1704906114
762
763   Pritchard, J. K., Stephens, M., & Donnelly, P. (2000). Inference of population structure using multilocus
764   genotype data. *Genetics*, *155*(2), 945-959. https://doi.org/10.1093/genetics/155.2.945
765
766   Pybus, O. G., Rambaut, A., & Harvey, P. H. (2000). An integrated framework for the inference of viral
767   population history from reconstructed genealogies. *Genetics*, *155*(3), 1429-1437.
768   https://doi.org/10.1093/genetics/155.3.1429
769
770   Quéméré, E., Amelot, X., Pierson, J., Crouau-Roy, B., & Chikhi, L. (2012). Genetic data suggest a natural
771   prehuman origin of open habitats in northern Madagascar and question the deforestation narrative in this
772   region. *Proceedings of the National Academy of Sciences*, *109*(32), 13028-13033.
773   https://doi.org/10.1073/pnas.1200153109
774
775   Railsback, L. B., Dupont, L. A., Liang, F., Brook, G. A., Burney, D. A., Cheng, H., & Edwards, R. L.
776   (2020). Relationships between climate change, human environmental impact, and megafaunal extinction
777   inferred from a 4000-year multi-proxy record from a stalagmite from northwestern
778   Madagascar. *Quaternary Science Reviews*, *234*, 106244. https://doi.org/10.1016/j.quascirev.2020.106244
779

780 Rambaut, A., Drummond, A. J., Xie, D., Baele, G., & Suchard, M. A. (2018). Posterior summarization in
781 Bayesian phylogenetics using Tracer 1.7. *Systematic biology*, *67*(5), 901-904.
782 https://doi.org/10.1093/sysbio/syy032
783
784 Rambaut, A., & Grass, N. C. (1997). Seq-Gen: an application for the Monte Carlo simulation of DNA
785 sequence evolution along phylogenetic trees. *Bioinformatics*, *13*(3), 235-238.
786 https://doi.org/10.1093/bioinformatics/13.3.235
787
788 Rannala, B., & Yang, Z. (2003). Bayes estimation of species divergence times and ancestral population
789 sizes using DNA sequences from multiple loci. *Genetics*, *164*(4), 1645-1656.
790 https://doi.org/10.1093/genet ics/164.4.1645
791
792 Salmona, J., Heller, R., Quéméré, E., & Chikhi, L. (2017). Climate change and human colonization
793 triggered habitat loss and fragmentation in Madagascar. *Molecular Ecology*, *26*(19), 5203-5222.
794 https://doi.org/10.1111/mec.14173
795
796 Salmona, J., Olofsson, J. K., Hong-Wa, C., Razanatsoa, J., Rakotonasolo, F., Ralimanana, H., ... &
797 Besnard, G. (2020). Late Miocene origin and recent population collapse of the Malagasy savanna olive
798 tree (Noronhia lowryi). *Biological Journal of the Linnean Society*, *129*(1), 227-243.
799 https://doi.org/10.1093/biolinnean/blz164
800
801 Solofondranohatra, C. L., Vorontsova, M. S., Hackel, J., Besnard, G., Cable, S., Williams, J., ... &
802 Lehmann, C. E. (2018). Grass functional traits differentiate forest and savanna in the Madagascar central
803 highlands. *Frontiers in Ecology and Evolution*, *6*, 184. https://doi.org/10.3389/fevo.2018.00184
804
805 Solofondranohatra, C. L., Vorontsova, M. S., Hempson, G. P., Hackel, J., Cable, S., Vololoniaina, J., &
806 Lehmann, C. E. (2020). Fire and grazing determined grasslands of central Madagascar represent ancient
807 assemblages. *Proceedings of the Royal Society B*, *287*(1927), 20200598.
808 https://doi.org/10.1098/rspb.2020.0598
809
810 Stronen, A. V., Mattucci, F., Fabbri, E., Galaverni, M., Cocchiararo, B., Nowak, C., ... & Caniglia, R.
811 (2022). A reduced SNP panel to trace gene flow across southern European wolf populations and detect
812 hybridization with other Canis taxa. *Scientific Reports*, *12*(1), 4195. https://doi.org/10.1038/s41598-022-
813 08132-0
814
815 Teixeira, H., Montade, V., Salmona, J., Metzger, J., Bremond, L., Kasper, T., ... & Radespiel, U. (2021).
816 Past environmental changes affected lemur population dynamics prior to human impact in
817 Madagascar. *Communications biology*, *4*(1), 1084. https://doi.org/10.1038/s42003-021-02620-1
818
819 Tiley, G. P., Barker, M. S., & Burleigh, J. G. (2018). Assessing the performance of Ks plots for detecting
820 ancient whole genome duplications. *Genome biology and evolution*, *10*(11), 2882-2898.
821 https://doi.org/10.1093/gbe/evy200
822
823 Tiley, G.P., Crowl, A.A., Manos, P.S., Sessa, E.B., Solís-Lemus, C., Yoder, A.D., Burleigh, J.G. 2021.
824 Phasing alleles improves network inference with allopolyploids. *bioRxiv*
825 https://doi.org/10.1101.2021.05.04.442457
826
827 Tiley, G. P., van Elst, T., Teixeira, H., Schüßler, D., Salmona, J., Blanco, M. B., ... & Yoder, A. D. (2022).
828 Population genomic structure in Goodman's mouse lemur reveals long-standing separation of
829 Madagascar's Central Highlands and eastern rainforests. *Molecular ecology*, *31*(19), 4901-4918.
830 https://doi.org/10.1111/mec.16632
831
Vieilledent, G., Grinand, C., Rakotomalala, F. A., Ranaivosoa, R., Rakotoarijaona, J. R., Allnutt, T. F., &
Achard, F. (2018). Combining global tree cover loss data with historical national forest cover maps to look
at six decades of deforestation and forest fragmentation in Madagascar. *Biological Conservation*, *222*,
189-197. https://doi.org/10.1016/j.biocon.2018.04.008

Virah-Sawmy, M., Willis, K. J., & Gillson, L. (2010). Evidence for drought and forest declines during the
recent megafaunal extinctions in Madagascar. *Journal of Biogeography*, *37*(3), 506-519.
https://doi.org/10.1111/j.1365-2699.2009.02203.x

Viruel, J., Conejero, M., Hidalgo, O., Pokorny, L., Powell, R. F., Forest, F., ... & Leitch, I. J. (2019). A
target capture-based method to estimate ploidy from herbarium specimens. *Frontiers in Plant Science*,
937. https://doi.org/10.3389/fpls.2019.00937

Vorontsova, M. S., Besnard, G., Forest, F., Malakasi, P., Moat, J., Clayton, W. D., ... & Linder, H. P.
(2016). Madagascar's grasses and grasslands: anthropogenic or natural?. *Proceedings of the Royal
Society B: Biological Sciences*, *283*(1823), 20152262. https://doi.org/10.1098/rspb.2015.2262

Wang, L., Beissinger, T. M., Lorant, A., Ross-Ibarra, C., Ross-Ibarra, J., & Hufford, M. B. (2017). The
interplay of demography and selection during maize domestication and expansion. *Genome
biology*, *18*(1), 1-13. https://doi.org/10.1186/s13059-017-1346-4

Weiß, C. L., Pais, M., Cano, L. M., Kamoun, S., & Burbano, H. A. (2018). nQuire: a statistical framework
for ploidy estimation using next generation sequencing. *BMC bioinformatics*, *19*(1), 1-8.
https://doi.org/10.1186/s12859-018-2128-z

Xie, M., Wu, Q., Wang, J., & Jiang, T. (2016). H-PoP and H-PoPG: heuristic partitioning algorithms for
single individual haplotyping of polyploids. *Bioinformatics*, *32*(24), 3735-3744.
https://doi.org/10.1093/bioinformatics/btw537

Yamaura, Y., Narita, A., Kusumoto, Y., Nagano, A. J., Tezuka, A., Okamoto, T., ... & Lindenmayer, D.
(2019). Genomic reconstruction of 100 000-year grassland history in a forested country: population
dynamics of specialist forbs. *Biology letters*, *15*(5), 20180577. https://doi.org/10.1098/rsbl.2018.0577

Yang, Z. (2014). Population demographic process. In, *Molecular Evolution: A Statistical Approach*
(pp.323-325). Croyden, UK. Oxford University Press.

Yoder, A. D., Campbell, C. R., Blanco, M. B., Dos Reis, M., Ganzhorn, J. U., Goodman, S. M., ... &
Weisrock, D. W. (2016). Geogenetic patterns in mouse lemurs (genus Microcebus) reveal the ghosts of
Madagascar's forests past. *Proceedings of the National Academy of Sciences*, *113*(29), 8049-8056.
https://doi.org/10.1073/pnas.1601081113

Figure 1 — *Loudetia simplex* distribution and single Madagascar origin. a) A field of *L. simplex* near Antsirabe Madagascar planted with eucalyptus (photo credit: George P. Tiley). b) Our sampling of *L. simplex* was based on collections made throughout Madagascar's Central Highland Plateau, with an elevation between 800 and 1800 meters. c) Phylogenetic relationships of individuals based on the phased (PH) data. Numbers next to nodes are posterior probabilities.

Figure 2 — Patterns of genetic structure and variation in Madagascar. a) Aggregated

STRUCTURE results across runs for an *a priori* number of clusters of two (K=2) for the different

data types. c) A plot of natural-log-scaled geographic distance against the pairwise distance (p-

distance). Lines are from a least-squares linear regression with the coefficient of determination

(R^2) shown for each data type. Statistical significance was determined with a Mantel test using

9999 permutations. Significance levels are *p*-value < 0.05 (*), *p*-value < 0.01 (**), and *p*-value <

0.001 (***). Colors represent different data types: one haplotype (OH), ambiguous genotypes

(AG), and phased haplotypes (PH).

Figure 3 — Principal component analyses across data types. a) All individuals using haploid (OH) data. b) Only Malagasy individuals using OH data. c) All individuals treating all sites as diploid. d) Only Malagasy individuals treating all sites as hexaploid. e) Only Malagasy individuals compressing unbalanced allelic dosage to single categories. f) Only Malagasy individuals treating all sites as diploid.

Figure 4 — Distributions of summary statistics across locus regions. Box-and-whisker plots show distribution medians and quartiles for the expected pairwise nucleotide divergence (π), the number of segregating sites (S), and a normalized measure of the difference between two estimators of nucleotide divergence, indicating deviations from neutral expectations (Tajima's D).

911 **Figure 5 — Extended Bayesian Skyline Plot results with fixed clock rates.** Two
912 independent runs are shown with mean population sizes shown by solid lines. Polygons show
913 the 95% HPD intervals with the overlap between runs in purple. Results are shown for the a)
914 OH, b) AG, and c) PH data. Relevant time periods are shown in gray, which includes likely
915 human colonization (LHC), the last glacial maximum (LGM), and last interglacial period (LIG).
916

921 **Figure 6 — Simulation results from EBSP analyses for an appropriate number of change
922 points.** Individual red lines are the mean N_e estimates from 100 simulated data sets. Blue points
923 and vertical lines are the mean of mean N_e estimates and their 95% HPD intervals. The dashed
924 black line is the true demographic history of the population.

926 **Tables**

Table 1 — X^2 test result from Tajima's D statistics across datasets and locus regions

Tested Effect	Fixed Effect	X^2	p-value	Largest Residual
Region	OH	7.72	0.27	Under-representation of Tajima's D < -2 in core regions
	AG	37.77	0.0001	Over-representation of Tajima's D < -2 in core regions
	PH	83.47	0.0001	Over-representation of Tajima's D < -2 in core regions
Dataset	Left Flank	12.65	0.046	Over-representation of Tajima's D < -2 in OH dataset
	Right Flank	8.61	0.19	Over-representation of Tajima's D < -2 in OH dataset
	Core	57.15	0.0001	Over-representation of Tajima's D < -2 in OH dataset