1   **Genetic variation in *Loudetia simplex* supports the presence of ancient grasslands in**
2   **Madagascar**
3

4   George P. Tiley[1,†], Andrew A. Crowl[2], Tchana O. M. Almary[3], W. R. Quentin Luke[4], Cédrique L.
5   Solofondranohatra[5], Guillaume Besnard[6], Caroline E.R. Lehmann[7,8], Anne D. Yoder[2], Maria S.
6   Vorontsova[1]
7   [1]Royal Botanic Gardens Kew, Richmond, UK
8   [2]Department of Biology, Duke University, Durham, North Carolina, 27708 USA
9   [3]Kew Madagascar Conservation Centre, Antananarivo 101, Madagascar
10   [4]East African Herbarium, National Museums of Kenya, P.O. Box 45166, Nairobi, 00100, Kenya
11   [5]Madagascar Biodiversity Center, Parc Botanique et Zoologique de Tsimbazaza, Antananarivo
12   101, Madagascar
13   [6]CNRS-UPS-IRD, UMR5174, Laboratoire Évolution & Diversité Biologique, Université Paul
14   Sabatier, Toulouse, France
15   [7]School of GeoSciences, University of Edinburgh, Edinburgh, UK
16   [8]Royal Botanic Garden Edinburgh, Edinburgh, UK
17
18   [†]Author for Correspondence: g.tiley@kew.org
19

Summary

1) Research Aims — The extent of Madagascar's grasslands prior to human colonization is unresolved. We used population genetic analyses of a broadly dominant C_4 fire-adapted grass, *Loudetia simplex*, as a proxy for estimating grassland change through time. We carefully examined the utility of target-enrichment data for population genetics to make recommendations for conservation genetics. We explored the potential of estimating individual ploidy levels from target-enrichment data and how assumptions about ploidy could affect analyses.

2) Methods — We developed a novel bioinformatic pipeline to estimate ploidy and genotypes from target-enrichment data. We estimated standard population genetic summary statistics in addition to species trees and population structure. Extended Bayesian skyline plots provided estimates of population size through time for empirical and simulated data.

3) Key Result — All Malagasy *Loudetia simplex* individuals sampled in this study formed a clade and possibly indicated an ancestral Central Highland distribution of 800m in altitude and above. Demographic models suggested grassland expansions occurred prior to the Last Interglacial Period and supported extensive grasslands prior to human colonization. Though there are limitations to target-enrichment data for population genetic studies, we find that analyses of population structure are reliable.

4) Key Point —Genetic variation in *Loudetia simplex* supports widespread grasslands in Madagascar prior to the more recent periods of notable paleoclimatic change. However, the methods explored here could not differentiate between paleoclimatic change near the Last Glacial Maximum and anthropogenic effects. Target-enrichment data can be a valuable tool for analyses of population structure in the absence a reference genome.

Keywords

Effective Population Size, Target-Enrichment, HybSeq, Polyploidy, Anthropogenic Climate Change, Conservation Genetics

Societal Impact Statement

Recognizing *Loudetia* dominated grasslands were widespread prior to human colonization

highlights that open ecosystems were and continue to be an important component of

Madagascar's biodiversity. Urgently required are biodiversity inventories and integrative

taxonomic treatments of grassland flora and fauna to asses risks to understudied ecosystems

historically regarded as wastelands. Substantial financial and logistical barriers exist to

implementing conservation studies using contemporary genomic tools. We ameliorated some of

the challenges for population genetic analyses of non-model polyploids lacking reference

genomes by developing computational resources to leverage a cost-effective data generation

strategy that requires no prior genetic knowledge of the target species.

Résumé

1) Les objectifs de la recherche — L'étendue des écosystèmes ouverts de Madagascar

avant la colonisation humaine reste à éclaircir. Nous avons utilisé une analyse de la

population génétique d'une graminée C_4 adaptée au feu, largement dominante, Loudetia

simplex, comme référence pour estimer les changements au niveau de ces biomes au fil

du temps. Nous avons examiné attentivement l'utilité des données d'enrichissement

ciblé pour la génétique de population afin de formuler des recommandations pour la

conservation génétique. Nous avons exploré le potentiel de l'estimation du niveau des

ploidies individuelles à partir des données d'enrichissement ciblé et comment les

hypothèses à propos de ces ploidies pourraient affecter les analyses.

2) Les méthodes — Nous avons développé un nouveau canal bioinformatique pour estimer

les ploidies et les génotypes à partir des données d'enrichissement ciblé. Nous avons

estimé les statistiques standard de la population génétique, en plus des arbres des

espèces et de la structure de la population. L'utilisation des tracés étendus du ciel

bayésien a fourni une estimation de la taille de la population au fil du temps pour des

données empiriques et simulées.

3) Résultat clé — Tous les individus Malagasy de *Loudetia simplex* échantillonnés dans

cette étude ont formé un clade, indiquant une éventuelle ancienne distribution dans les

hauts plateaux. Les modèles démographiques suggèrent une expansion des prairies

bien avant la dernière période interglaciaire et soutiennent l'existence d'une vaste

distribution avant la colonisation humaine. Bien qu'il y ait des limites à l'enrichissement

des données cibles pour l'étude de la génétique des populations, nous constatons que

l'analyse des structures des populations est fiable.

4) Les points clés — La variation génétique de *Loudetia simplex* soutient l'existence de

vastes prairies à Madagascar avant les périodes plus récentes de changements

paléoclimatiques notables. Cependant, les méthodes explorées ici n'ont pas permis de

faire la différence entre les changements paléoclimatiques près du dernier maximum

glaciaire et les effets anthropogènes. Les données d'enrichissement ciblé peuvent être

un outil précieux pour les analyses de la structure des populations en l'absence d'un

génome de référence.

Déclaration d'impact societal

Reconnaître que les prairies dominées par Loudetia étaient répandues avant la colonisation

humaine souligne que les écosystèmes ouverts étaient et continuent d'être un composant

important de la biodiversité de Madagascar. Il est urgent de réaliser des inventaires de la

biodiversité et une taxonomie intégrée pour le traitement de la flore et de la faune des

écosystèmes ouverts afin d'évaluer les risques pour les écosystèmes sous-étudiés considérés

historiquement comme des terres en friches. Des barrières financières et logistiques existent

pour mettre en œuvre l'étude de la conservation en utilisant les outils génomiques

contemporains. Nous avons amélioré certains des défis liés aux analyses génétiques de

populations de polyploïdes non modèles, sans génomes de référence, en développant des

ressources informatiques pour exploiter une stratégie pouvant générer des données rentables

ne nécessitant aucune connaissance génétique préalable de l'espèce cible.

104
105

Famintinana

1) Ny tanjon'ny fikarohana — Mbola tsy fantatra mazava tsara ny fivelaran'ny hivoka teto
Madagasikara talohan'ny fahatongava'ny olombelona. Mba ahafantarana ny fihovana
nitranga nandritra ny fotoana naharitra teo amin'ireo hivoka ireo dia nanao
famakafakahana ara-genetika amin'ny ahitra C_4 miompana amin'ny afo iray antsoina
Loudetia simplex ara-tsiantifika na Berambo na Hara amin'ny teny malagasy izahay.
Nandinika tsara ny maha-zava-dehibe ny fampitomboana ny antotan-kevitra mba
ahafahana manolo-kevitra momba ny fiarovana ny fototarazo genetika. Nandinika ny
mety mampiavaka ny fanombanana an'ny ploidy tsirairay amin'ny fampitomboana
antotan-kevitra sy ny mety ho fiantraikan'ny fiheverana momba ireo ploidy ireo amin'ny
fikarohana.

2) Fomba Fiasa — Namorona fantsona bioinformatika vaovao mba ahafahana manombana
ny ploidy sy ny « genotypes » avy amin'ny antotan-kevitra nokendrena izahay.
Notombanana ny antontan'isa famintinana ny fototarazo ara-genetikan'ireo
vondron'ahitra ireo, miampy ny karazana hazo sy ny firafitry ny vondrona na koa hoe
mponina. Nanome tombantombana ny haben'ny mponina amin'ny alàlan'ny fotoana ny
antontan-kevitra voavinavina azo tamin'ny fikarohana. Fikarohana izay azo tamin'ny
alalan'ny « Bayesina Skuline Plots ».

3) Vokam-pikarohana fototra — Ny vondrona *Loudetia simplex* eto Madagasikara izay
niasana dia namorona « clade » na fikambanana iray, izay manondro ny mety maha ela
netezana sy tranainy an'io ahitra io eny amin'ny faritra avo. Ny modely demografika dia
manoro hevitra amin'ny naha be velarana ny hivoka izay efa ela talohan'ny vanim-
potoana « interglacial » farany ary manohana ny fivelarana midadasika an'ireo kijana
ireo alohan'ny fonenan'ny olombelona. Na dia misy fetrany aza ny fampitomboana ny
antotan-kevitra kendrena amin'ny fandalinana ny fototarazo genetika momban'ny
mponina, dia hita fa azo itokisana ny fikarohana natao momban'ny firafitry ny mponina.

4) Hevi-dehibe — Ny fahasamihafana ara-genetika ao amin'ny *Loudetia simplex* dia
manohana ny fisian'ny hivoka na kijana midadasika eto Madagasikara talohan'ny
vanim-potoanan'ny fiovana paleoclimatika nisongadina. Na izany aza, ny fombam-
pikarohana nampiasana teto dia tsy nahavita nanavaka ny fiovan'ny paleoclimatika
akaikin'ny vanim-potoana lehibe nangatsiaka farany sy ny vokatry ny fitrandrahana
nataon'ny olombelona. Mety ho fitaovana manan-danja amin'ny famakafakana ny firafitry
ny mponina ny antotan-kevitra nampitombona na dia tsy misy fitaovana genomika
iangaina aza.

Fanambarana fiantraika ara-tsosialy

Ny fanekena fa niely patrana ny hivoka itoeran'ny *Loudetia* talohan'ny fanjanahan'ny

olombelona dia manamarika fa ireo hivoka ireo dia singa manan-danja amin'ny zavamananaina

eto Madagasikara. Ilaina maika ny fahafantarana ara biolojika sy taxononomique ny zavamaniry

sy ny biby amin'ny hivoka mba hanombanana ny loza mety hitranga amin'ny hivoka izay tsy

ananana fahalalana maro sady heverina ho tany maina. Misy sakana ara-bola sy ara-pitaovana

amin'ny fampiharana ny fandalinana momba ny fiarovana izay nampiasana fitaovana génomika

ankehitriny. Nohatsarainay ny sasany amin'ireo fanamby mifandraika amin'ny famakafakana

ara-genetika ny mponina manana ploidy maro tsy modely, izay tsy misy fitaovana genomika

iaingana , amin'ny alàlan'ny fampivoarana loharanon-kevitra kajy mba hitrandrahana paikady

izay mety hiteraka angon-drakitra mahomby tsy mitaky fahalalana mahakasika ny fototarazo

ara-genetika ny zava-maniry izay tiana karohina.

152
153

Introduction

The degree to which anthropogenic activities have transformed the ancestral vegetative composition of Madagascar's ecosystems is heavily debated. While the narrative that most of central Madagascar was once covered in closed-canopy forests has been rejected (Bond et al., 2008; Vorontsova et al., 2016; Hackel et al., 2018; Solofondranohatra et al., 2020; Bond et al., 2022; Lehmann et al., 2022), the degree to which a former mosaic of grassland, shrubland, and forests has become more dominated by grasses is contentious (reviewed in Joseph et al., 2021). Understanding population dynamics of individual ecologically important species and plant communities will have profound effects on conservation policy decisions and the economic productivity of grasslands for local communities. However, it is difficult to determine where grasslands, which we consider to be an ecosystem where the ground layer is a majority of tussock-forming grass with little tree cover that experiences a dry season with regular fire regimes (e.g. Dixon et al., 2014), were in the past and how extensive they were. Stratigraphic sediment records from Central Madagascar indicate an increased relative abundance of grasses and fire through the Holocene (Burney, 1987; Gasse & Van Campo, 1998; Virah-Swamy et al., 2010), especially near the expected age of human colonization [c. two thousand years ago (ka); Dewar & Wright, 1993; Dewar et al., 2013; Pierron et al., 2017]. Megafaunal isotope records also support Holocene expansions of C_4 grasslands but show a vegetative mosaic of C_4 grasses and C_3 vegetation was at least present (Crowley et al., 2021) near the last glacial maximum (LGM; c. 26.5-19 ka). Reconstructing spatio-temporal vegetative dynamics beyond the LGM begin to approach limitations of paleoecological data, but molecular evolutionary approaches can fill the gaps by providing probabilistic inferences through time. Population genetic approaches for reconstruct demographic histories has been successfully applied to lemurs to reconstruct patterns of past forest connectivity (Quéméré et al., 2012; Yoder et al., 2016; Salmona et al., 2017; Teixeira et al., 2021; Tiley et al., 2022) and may similarly provide insights into the Late Pleistocene dynamics of grasses between the LGM and Last Interglacial period (LIG; c. 132-112 ka) or older. This strategy has been used to reconstruct vegetative dynamics of temperate Japanese grasslands from forb species (Yamaura et al., 2019).

An array of tools now exists to estimate the effective population size (N_e) of species through time. As generating genomic data for non-model groups has become more accessible, population genomic methods that utilize precisely estimated site frequency spectra (SFS; Nielsen et al., 2000; Gutenkunst et al., 2009; Excoffier et al., 2013; Liu & Fu, 2020; Blischack et

al., 2022) have become a valuable tool for reconstructing the demographic history of natural

populations. There are still barriers to generating population genomic data though, either due to

the lack of suitable tissue, an appropriate reference genome, or costs. Target-enrichment

libraries have been transformative for the analysis of plant phylogenetic relationships (e.g.

Breinholt et al., 2021; Baker et al., 2022), but the population genetic utility of such data is not

well-explored. A survey of taxa with low sample sizes suggested population genetic parameters

estimated from target-enrichment data could be unreliable, in addition to violating assumptions

of the neutral coalescent model that underpins most tests of selection and demographic change

(Slimp et al., 2021). However, the relatively long loci recovered by target-enrichment (or

HybSeq) methods could be appropriate for a demographic modeling approach, the extended

Bayesian skyline plot (EBSP; Heled & Drummond, 2008), which has been successfully applied

to a range of conservation genetic questions despite some model violations. We are interested

in exploring the potential of target-enrichment data for population or conservation genetic

insights from plant groups that may be lacking genomic resources.

Here, we examined the population genetics of the grass species *Loudetia simplex*

(Tristachyideae: Panicoideae: Poaceae). It is largely dominant in Madagascar's Central

Highlands (Koechlin, 1993; Hagl et al., 2021; Fig. 1a) and is notable for its fire- and grazing-

adapted functional traits (Solofondranohatra et al., 2018). Population genetic analyses of

microsatellites suggest Malagasy *L. simplex* has been isolated from mainland African

populations with no detectable level of gene flow, and that there is additional population

structure between the northern and southern extents of the species range across the Central

Highlands of Madagascar (Hagl et al., 2021). Here, we use a subset of individuals analyzed by

Hagl et al. (2021) along with a new sample from the center of the *L. simplex* distribution to

explore the population genetic utility of target-enrichment data and its implications for the natural

history of Madagascar's grasslands. Because Malagasy *L. simplex* are putative polyploids

(tetraploids and hexaploids; Hagl et al., 2021), we developed novel bioinformatic tools for the

processing and analysis of polyploid data, integrated into the PATÉ allele phasing pipeline (Tiley

et al., 2021).

Materials and Methods

Sampling and Sequencing

Target-enrichment data was generated for 24 individuals of *Loudetia simplex* and one outgroup,

Tristachya nodiglumus (Supplementary Table S1). Twenty individuals were selected to best

represent Madagascar from available collections as a single unstructured population and avoid

complications of population substructure and low sample sizes in downstream analyses (Fig.

1b). Genomic DNA samples were used from Hagl et al. (2021) while new extracts used a

modified CTAB protocol (Doyle & Doyle, 1987) from silica-dried leaf tissue. Genomic DNA was

then sent to Rapid Genomics (Gainesville, FL, USA) where sample enrichment and sequencing

was performed with a bait kit developed for angiosperms following Breinholt et al. (2021). All

sample metadata is provided in Supplementary Table S1.

Sequence Assembly and Processing

Initial contigs were assembled with HybPiper (Johnson et al., 2016) with default settings. We

then used HybPiper's supercontigs as reference sequences for PATÉ (Tiley et al., 2021; Crowl

et al., 2022) to genotype individuals, filter low-quality variants with GATK v4.2.0 (McKenna et

al., 2010; DePristo et al., 2011), and phase haplotype sequences using HPOP-G (Xie et al.,

2016). Ploidy information for individuals (Supplementary Table S1) was based on previous

microsatellite allele count data (Hagl et al., 2021); however, we also implemented a statistical

test to infer ploidy level in PATÉ based on mixtures of distributions from allele balance data,

similar to other approaches that use the number of reads supporting an alternate allele over

total reads to infer ploidy (Weiß et al., 2018). We extended the mixture models of Tiley et al.

(2018) to determine the ploidy of a sample with the model weights as a selection criterion

(Supplementary Methods: *Ploidy Test*). PATÉ control files with all filtering options for GATK are

provided in the Supplementary Data (https://doi.org/10.5061/dryad.905qfttqx). Because target-

enrichment probes may represent multiple exons from the same gene with overlapping flanking

regions, we were concerned that some assembled loci might overlap and violate some

independence assumptions in downstream analyses. Therefore, we performed an all-by-all

blastn search (Camacho et al., 2009) with an e-value of 1×10^{-5} among the supercontig

assemblies followed by single-linkage clustering (https://github.com/gtiley/HybSeq-Cleaning).

Only the longest locus per cluster was retained for downstream analyses.

To investigate nucleotide variation and the potential population genetic value of non-coding

versus coding regions of target-enrichment loci, we identified the following regions:

1) *Left Flank* – Non-coding sequence upstream of the 5' start codon or splice site.

2) *Core* – The coding region targeted by probes for sample enrichment.

3) *Right Flank* – Non-coding sequence downstream of the 3' stop codon or splice site.

Identification of individual regions was automated (https://github.com/gtiley/Locus-Splitter)

based on the tblasx program from BLAST v2.10.0 (Camacho et al., 2009) to find in-frame

alignments between our assembled sequences and the original transcriptome loci used for

probe development. Individual regions were aligned with MAFFT v7.471 (Katoh & Standley,

2013). The aligned regions were then concatenated together to make whole-locus alignments.

Because we are interested in investigating the analytic impact of ploidy in sequence data, we

created three datasets from the phased ploidy-aware data:

1) *One Haplotype (OH)* – Only one phased haplotype sequence was randomly selected from

each individual for each locus.

2) *Ambiguous Genotype (AG)* – The genotypic information for an individual is represented as a

single sequence with IUPAC ambiguity codes for biallelic sites. Analyses that treat IUPAC

codes as missing data were randomly resolved into two sequences.

3) *Phased Haplotypes (PH)* – All phased haplotype sequences for an individual are used.

We expect the OH data to be representative of majority consensus sequences (e.g. HybPiper

supercontigs) since our target-enrichment loci are anchored in single exons and we anticipate

that any risk of assembling chimeric sequences from such data without paralog warnings is low.

Any site genotyped as polyallelic, having more than one alternate allele, was assumed to be an

error and treated as missing sequence across all data sets. All expected copies of a locus are

represented even if they are invariant. For example, some individuals in the PH data may have

six identical sequences at a locus if they are hexaploids and every site is homozygous. While

the ploidy at an individual locus may vary due to differential gene loss among chromosomes,

this is a simplifying assumption to preserve expected allele frequency information.

Phylogeny Estimation

A species tree was estimated for all *L. simplex* individuals using whole-locus sequences for the

OH, AG, and PH datasets with BPP v4.6.2 (Flouri et al., 2020). We chose a full-likelihood

multispecies coalescent (MSC; Rannala & Yang, 2003) model since it is robust to low-

information gene tree estimates (Xu & Yang, 2016). Markov chain Monte Carlo (MCMC)

analyses collected 10,000 posterior samples, saving every 10 samples after a 10,000

generation burnin. Model and prior specifications are provided in the Supplementary Material

(Supplementary Methods: *BPP Prior Specification*). The starting tree for our Bayesian species

tree analyses was based on a concatenated maximum likelihood (ML) estimate from the OH

data with IQTREE v2.1.2 (Minh et al., 2020). We used unpartitioned analyses with

ModelFinderPlus (Kalyaanamoorthy et al., 2017) for the OH, AG, and PH data to estimate ML

evolutionary distances between individuals, with random concatenation of haplotypes within

individuals.

Population Structure

Potential population structure within our sample of *L. simplex* from Madagascar was explored

with STRUCTURE v2.3.4 (Pritchard et al., 2000). For each dataset, we sampled only one

biallelic SNP per locus, if present, and required a minimum minor allele count of three with no

more than 25% missing data. To explore uncertainty due to sampling error of sites in a target-

enrichment dataset that is relatively small compared to contemporary population genomic data,

we generated 10 datasets with random SNP sampling. The number of clusters that best satisfy

Hardy-Weinberg assumptions was determined with the ΔK method (Evanno et al., 2005).

Principal component analyses (PCAs) were conducted on the OH, AG, and PH datasets with

adegenet v2.1.3 (Jombart & Ahmed, 2011) by compressing allele counts to the same dimension

as the lowest ploidy level present. For example, a hexaploid with three reference and three

alternate alleles would be scored as a 2 on a scale from 0 to 4, to be comparable to tetraploids

in the dataset. Additional details on allelic imbalances would be lost as, for example, hexaploids

with either one or two alternate alleles would be represented as a 1 on the tetraploid scale.

Scripts for generating PCA or other clustering-ready matrices from fasta files are available on

GPT's GitHub (https://github.com/gtiley/fasta-conversion). Additional details about compression

of PCA data matrices can be found in the Supplementary Material (Supplementary Methods:

PCA Input Data). The presence of isolation-by-distance as an explanation for observed patterns

of genetic structure and ordination was tested with the patristic distances from the ML analyses

or the uncorrected pairwise distances (p-distances) regressed against the natural logarithm of

the great circle distance between individuals. A Mantel test was performed for isolation-by-

distance analyses in the ape v5.6.2 R package (Paradis & Schliep, 2019).

Population Genetic Summary Statistics

The pairwise nucleotide diversity (π), number of segregating sites (*S*), and Tajima's *D* were

calculated across locus regions and datasets with the PopGenome v2.7.5 R package (Pfeifer et

al., 2014). The significance of Tajima's *D* statistics was determined by 100 standard coalescent

simulations with ms (Hudson, 2002). We generated 200 bootstrap replicates of each alignment

to calculate a percentile bootstrap 95% confidence interval and compare the relative estimation

error between regions and datasets using the root mean square error (RMSE) from the

bootstrap samples. The scaling and interpretation of parameters are described in the

Supplementary Material (Supplementary Methods: *Summary Statistics*). We tested for

differences in the distribution of summary statistics among locus regions (left flanks, cores, and

right flanks) and data types (OH, AG, and PH) with Wilcoxon Rank-Sum tests. We tested for

biases in Tajima's *D* results among locus regions and data types with X^2 tests.

Demographic History

We chose the extended Bayesian skyline plot (EBSP, Heled & Drummond, 2008) as the most

appropriate model of N_e change through time for our sample of target enrichment data. Here we

assumed a strict molecular clock underlying all lineages but allowed rates of evolution to vary

among loci. One locus was fixed to the clock rate and all other locus rates were relative. Rates

of molecular evolution among plants are especially complicated, as there can be life histories

with variable reproductive strategies, unknown generation times, and overlapping generations.

Thus, we relied on an estimate of the neutral substitution rate at third-position synonymous sites

from grass genomes (Christin et al., 2014) and constructed a vague prior around this mean rate,

such that any biologically plausible uncertainty in the rate of evolution is reflected in our

posterior sample. It is a tenuous rate estimate at best, since it remains unknown whether

polyploid *L. simplex* largely propagates clonally, through apomixis, or through sexual

reproduction. The MCMC analyses for the OH, AG, and PH data collected 20,000 samples after

a burnin of 50,000,000 and sample interval of 10,000. Two replicates were run for each analysis

to evaluate convergence and mixing was assessed with Tracer v1.7 (Rambaut et al., 2017). All

model and prior specifications are provided in the Supplementary Material (Supplementary

Methods: *EBSP Prior Specification*) and are replicable from XML files on Dryad

(https://doi.org/10.5061/dryad.905qfttqx).

Adequacy of Demographic Models

To explore if EBSP models could reliably differentiate between Late Pleistocene versus

Holocene population size increases, we evaluated the accuracy of EBSP inferences with

simulations that captured some features of our target-enrichment data. We simulated 57 loci

that were 1000 bp in length for 20 haploid individuals 100 times with ms (Hudson, 2002) and

seq-gen (Rambaut & Grass, 1997) for four scenarios: 1) a constant N_e of 100,000, 2) an N_e

increase from 10,000 to 100,000 individuals roughly coinciding with the LGM at 20,000 years in

the past, 3) the same ten-fold increase but at 2,000 years in the past to reflect anthropogenic

expansion, and 4) a scenario combining both LGM and anthropogenic effects by exponentially

increasing N_e from 10,000 to 100,000 between 20,000 years ago and the present. Each

simulated dataset was then evaluated with the same EBSP model using a fixed clock used for

our empirical analyses, except the MCMC algorithm collected 10,000 posterior samples with a

sample interval of 1,000 after a 100,000 sample burnin. Commands for simulations are given in

the Supplementary Material (Supplementary Methods: *Simulating Demographic Histories*).

Results

Sequencing Statistics

The probe set targeted 408 loci from which an average of 327 loci were assembled by HybPiper

(Supplementary Table S2). These were supported by an average of 217 read pairs per locus.

Although there was variable read depth across loci and individuals, the number of assembled

loci was significantly correlated with read depth (Pearson's correlation coefficient = 0.57, *p*-

value = 0.002). After joint-genotyping all individuals as diploids, normal mixture models

suggested all individuals were hexaploids, with the exception of two individuals from South

Africa (Supplementary Table S3). Some individuals were inferred as pentaploids, but since this

was unexpected based on previous observations from microsatellites (Supplementary Table S1)

these cases were assumed to be hexaploids. Although available evidence leaves the ploidy of

some individuals ambiguous and raises skepticism of the inference methods, genotyping and

phasing of individuals proceeded with the ploidy suggested by our mixture models. We inferred

an average of 247 loci with allelic variation, which had a heterozygosity of 0.012 variants per

base. After reducing the number of loci with BLAST clustering to avoid tight linkage, a final 301

loci were retained from the OH, AG, and PH datasets for downstream analyses. The average

number of haplotype blocks across all loci was 1.005, meaning that in very few cases there was

insufficient data to phase across the entire locus (Supplementary Table S2). In cases where

multiple variants within a locus could not be phased, only the variants from the longest block

were retained and variants on the shorter block were replaced with missing data.

Phylogeny Estimation

Species trees revealed all Malagasy *L. simplex* form a well-supported clade (Supplementary

Fig. S1), but there was disagreement in the biogeographic history within Madagascar based on

the treatment of ploidy. Individuals in northern Madagascar formed a clade with a posterior

probability of 0.9 or greater across the OH, AG, and PH analyses. However, the individuals from

central and southern Madagascar were paraphyletic in the OH and AG analyses

(Supplementary Figs. S1a and S1b). A clade of central and southern individuals was recovered

in the PH analysis, but with a weak posterior probability of 0.64 (Fig. 1c; Supplementary Fig.

S1c). Individuals from the center of the *L. simplex* distribution appear to be the earliest diverging

lineages in the OH and AG analyses (Supplementary Figs. S1a and S1b), or with respect to the

sample of central and southern individuals in the PH analysis (Fig. 1c; Supplementary Fig. S1c).

Population Sub-Structure in Madagascar

STRUCTURE analyses recovered a variable number of Hardy-Weinberg groups across the ten

datasets that randomly sampled one non-singleton biallelic SNP per locus (Supplementary

Table S4). For the OH data, we found a *K* of two, three, or four in eight, one, and one of the

respective replicates. For AG data, the optimal number of clusters was more ambiguous with a

K of two, three, and four recovered for four, five, and one replicates. PH data found *K* of two and

three for eight and two replicates. We note that the method of Evanno et al. (2005) cannot infer

if one panmictic population is the best model, and while there is reasonable geographic

structure at *K* of two weighted across biological and technical replicates (Fig. 2a) the non-zero

admixture between individuals at the most extreme ends could indicate a single population with

a cline of genetic variation. This is further supported by the lack of distinct cluster assignments

at higher values of *K* despite some evidence for additional population structure (Supplementary

Fig. S2). A weak signal of isolation-by-distance was detected when regressing genetic distance

against spatial distance (Fig. 2b). Notably, the AG data inferred much higher pairwise distances

between individuals; although, the relative differences and overall trend were similar. The same

patterns can be observed when using patristic distances from ML phylogenetic estimates,

although the disparity between AG versus OH or PH distances is reduced, as is the percent of

genetic variation explained by spatial distances (Supplementary Fig. S3).

PCAs reflected findings from STRUCTURE analyses and reinforce the presence of a genetic

cline (Fig. 3). For OH data, where only one reference or alternate allele was observed per

individual, Malagasy *L. simplex* is clearly distinguished from mainland African individuals (Fig.

3a), but there is no clear separation of individuals based on geographic location (Fig. 3b). PCAs

of data that accounts for heterozygosity but not dosage recovered a single Malagasy versus

mainland African pattern (Fig. 3c), PCAs reflect a north-to-south gradient of genetic variation

regardless of whether allelic dosage was treated as hexaploid (Fig. 3d), tetraploid (Fig. 3e), or

diploid (Fig. 3f). The total percent of variance explained by principal component axis 1 (PC1)

and axis 2 (PC2) increased as allelic dosage was compressed from a hexaploid to a diploid

state.

Variation in Summary Statistics

There were significant differences in the distributions of π, S, and Tajima's D among locus

regions and between the OH, AG, and PH data (Fig. 4). For π, there was always a difference

between flanking regions and the exon core (Supplementary Table S5), with the cores

surprisingly having more variation. Differences were also observed between the flanking regions

and cores for S and Tajima's D, except when comparing the left flanks and cores for OH data

(Supplementary Table S5). The only case of a difference between the left and right flanks was

when estimating π for AG data (Supplementary Table S5). The OH and PH data resulted in

similar estimates of π; however, the AG and PH data had similar distributions of S

(Supplementary Table S5). Any differences in Tajima's D statistics were largely explained by an

over-representation of values less than negative two, indicating positive selection or population

size expansion, in the OH data and among core regions across datasets (Table 1).

We attempted to explore biases in summary statistic estimates due to sampling error of sites

through bootstrapping the OH (Supplementary Fig. S4), AG (Supplementary Fig. S5), and PH

data (Supplementary Fig. S6). Our observed summary statistics typically fell outside of their

percentile bootstrap 95% confidence intervals, but other confidence interval approximation

techniques (e.g. Efron & Narasimhan, 2020) may be more appropriate for population genetic

summary statistics (Leblois et al., 2003). However, we used the bias, difference between the

bootstrap mean and observed estimate, and the RMSE of bootstrap replicates with respect to

the observed estimate, to gain some general insights. Notably, the bias of π is larger and much

noisier in the AG data (Supplementary Fig. S5) compared to the OH (Supplementary Fig. S4)

and PH (Supplementary Fig. S6) data. Even if the bias for a whole-locus is low, there can be

high variability between regions. For the observed values, estimates are not always proportional

among regions either. The RMSE of an estimate tends to decrease along with genetic variation,

implying some role of genotyping error or rare variants and singletons inflating values at the

higher end of the spectrum. Loci with very low levels of genetic variation can also be

problematic though, as the RMSE for flanking regions increases towards the tail of the

distribution, creating opportunity for sampling error to wildly swing estimates regardless of ploidy

(Supplementary Figs. S4-S6). The number of segregating sites increased with ploidy of the

data, but π was relatively stable. However, differences between π and S will affect estimates of

Tajima's D, and this could affect interpretations of demography or selection acting on loci;

although, the majority of loci and their regions were not significantly different from neutral

expectations given the simulation-based test statistics (Supplementary Table S6).

Population Size Change Through Time

After filtering for longer loci with low amounts of missing data, 57 loci were retained for EBSP

analyses. Reducing our data to a subset of more informative loci was done for computational

convenience, because EBSP models are parameter-rich and require long MCMC runs for

convergence. Runs converged for each data type for both analyses with a fixed rate of

molecular evolution (Fig. 5) and variable rate (Supplementary Fig. S7). All analyses suggested

the *L. simplex* N_e increased through the Late Pleistocene, well before the LIG, approximately

one million years ago. Between the LGM and period of potential human colonization of

Madagascar in the Holocene, N_e had already reached present-day estimates. N_e estimates

through the Holocene were only possible for the PH data, as accounting for the within-individual

variation should have captured more recent coalescences. Simulations cast doubt on the

seriousness with which EBSP results can be interpreted through the Holocene though. For

example, when N_e was truly constant through time, some individual EBSP analyses would

suggest recent N_e increases; although, the mean estimate across simulations recovered the

constant population scenario (Fig. 6a). This problem was exaggerated in the presence of model

over-fitting (Supplementary Fig. S8). In scenarios with recent N_e increases, the contemporary N_e

was underestimated but a trend indicating a recent N_e increase was detectable (Figs. 6b-6d).

However, simulations indicated it would not be possible to differentiate between N_e increases

that occurred during the LGM (Fig. 6b) versus the Late Holocene (Fig. 6c), let alone some

combination of paleoclimate and anthropogenic effects (Fig. 6d). Interpreting these recent N_e

increases from EBSP analyses becomes impossible with model over-fitting (Supplementary Fig.

S8).

Discussion

Robustness of Analyses Across Loci and Ploidy

Although target-enrichment data has been transformative for species-level phylogenomic

studies (Breinholt et al., 2021; Baker et al., 2022), the robustness of population genetic

estimates for a sample of closely related individuals was largely unexplored (but see Slimp et

al., 2021). Some of the most important measures of genetic variation for populations were

susceptible to sampling error, especially in the flanking intron regions (Supplementary Figs. S4-

S6). While estimates from cores showed less error, they will lead to some violations of

population genetic methods that assume evolution consistent with neutral theory, given potential

signatures of purifying selection (Table 1). This creates some conflict, since most of the

information is contained within the exon cores of target-enrichment loci (Fig. 4), presumably due

to read coverage. Because Tajima's *D* alone cannot disentangle demography from selection,

some judgment calls and tolerance for downstream model violations are necessary. The

majority of loci examined were consistent with neutral expectations based on simulations

(Supplementary Table S6), and the proportion of loci with significant Tajima's *D* statistics were

exaggerated when ignoring heterozygosity (Table 1). Although, the estimates of nucleotide

diversity and simulations under the standard neutral coalescent used here should be reasonable

for polyploids in cases of perfect multisomic inheritance in outcrossers (Arnold et al., 2012;

Blischak et al., 2023), we note that the presence of subgenome structure and selfing could

cause deviations in expectations of allele frequencies (e.g. Ronfort et al., 1998; Meirmans &

Van Tienderen, 2013; Blischak et al., 2023). The mechanism of polyploidization is unknown in *L.*

simplex, but our results do provide some expectations of how genotyping choices for polyploids

can impact standard summary statistic estimates.

Predicting the effects of assumptions about genotype on estimates of genetic variation and

downstream analyses are not always straightforward. For example, the pairwise distances (Fig.

2b) or evolutionary distances (Supplementary Fig. S3) between individuals was higher when

assuming all individuals were diploid (AG) compared to assuming all individuals were haploid

(OH) or genotyped at their anticipated ploidy level (PH). If we were willing to ignore the absolute

estimates of genetic distance, the interpretation of isolation-by-distance analyses was the same

across the OH, AG, and PH data. This implies that biologically meaningful conclusions can arise

even with uncertainty in ploidy and phasing. Interestingly, treating all individuals as diploid was

sufficient for STRUCTURE to recover northern and southern clusters with some admixed central

individuals of varying proportions (Fig. 2a; Supplementary Fig. S2). Similarly, PCAs reflected the

geographic distribution of our *L. simplex* samples in the AG and PH data with, surprisingly, the

AG data explaining more of the observed genetic variance (Fig. 3f) than the PH data (Fig. 3d).

This may be due to genotyping error at higher ploidy levels caused by read stochasticity and a

flat genotype likelihood surface (Blischak et al., 2016) or caused by incorrect ploidy inferences.

A similar mixture model approach to the one implemented here (Weiß et al., 2018) has been

shown to have good performance based on comparisons of allele balance distributions from

target-enrichment data and flow cytometry-based ploidy estimates (Viruel et al., 2019). We are

cautiously optimistic about the prospects of direct ploidy estimation from sequence data,

because likelihood-based criteria can be prone to overfitting (Tiley et al., 2018). However, our

ploidy estimates are generally consistent with the previous inferences from microsatellite allele

counts (Hagl et al., 2021; Supplementary Table S1; Supplementary Table S3). A positive insight

from our comparisons across ploidy levels is that if unknown individuals are genotyped as

diploid, there will be sufficient information about population structure, and that ploidy estimation

should not be a barrier to investigations of natural plant populations with little *a priori* cytological

or genetic knowledge.

Limits to Differentiating Anthropogenic and Paleoclimate N_e Change

Large population genomic studies in plants that can leverage demographic estimates from SFS

(e.g. Excoffier et al., 2013; Liu & Fu, 2020; Blischak et al., 2023) are commonly focused on

temperate regions (e.g. Yamaura et al., 2019) and species exploited for energy or agriculture

(e.g. Huang et al., 2014; Wang et al., 2017). Thus, it is tempting to leverage models that use

gene trees as data such as the EBSP (Heled & Drummond, 2008) or other extensions of skyline

plots (Pybus et al., 2000), since they could be paired with relatively low-cost target-enrichment

strategies that yield multi-locus nuclear data while stretching material quality.

While our analyses provide valuable insights into the ancient origins of *L. simplex* dominated

grasslands, both our empirical and simulated analyses show we cannot differentiate between N_e

increases that occurred during human colonization or paleoclimate change near the LGM. The

change in vegetative composition towards more C_4 grass species in Central Madagascar is well-

documented (Burney, 1987; Gasse & Van Campo, 1998; Virah-Swamy et al., 2010; Crowley et

al., 2021). It is likely that these increases in C_4 grasses included *L. simplex*; although, an

increase in census size may not be reflected by N_e (Fig. 5; Supplementary Fig. S7), the

expected population size under a neutral coalescent model. Our simulations showed that

detecting these recent increases is difficult, and that we cannot differentiate between a ten-fold

increase that occurred 200 (2 ka) or 2000 generations ago (20 ka; Fig. 6). Diagnosing a true N_e

change becomes impossible in the presence of model overfitting (Supplementary Fig. S8) and

diagnosing these problems in empirical data can be difficult aside from convergence

assessments.

These limitations do not necessarily disagree with an assessment of EBSP models from a larger

genomic dataset of endemic palm species in Madagascar (Helmstetter et al., 2021). Helmstetter

et al. (2021) found that the estimated N_e trends were valuable predictors of IUCN risk

assessments, and that population decreases could be detected at anthropogenic time scales.

However, it should be easier to detect population declines than population increases in a

system that already had a large degree of standing genetic variation, such as *L. simplex*. This is

because the time until the most recent common ancestor of a genealogy will be much more

recent in a small population than a large one (e.g. see equation 2.13 from Gillespie, 2004).

Analyses of demographic models with genomic SFS should be able to increase resolution at

recent anthropogenic time scales (Patton et al., 2019) and differentiate anthropogenic from

paleoclimatic demographic effects (Tiley et al., 2022), but this would require more extensive

sampling and sequencing. Additionally, our analyses treat all individuals as a single population,

and since we detected some level of population structure, we cannot rule out their confounding

effects on analyses of N_e through time (Heller et al., 2013; Mazet et al., 2016) or other

confounding affects such as clonal growth and overlapping generations. Population-level

sampling of multiple sites throughout a species range should be used to control for population

structure, while accounting for localized effects such as agriculture and grazing versus regional

climate effects.

New Insights into Loudetia simplex *Evolution and Policy Implications*

Our analyses yielded new findings about the evolution of *L. simplex* that are potentially

consequential for the broader understanding of Malagasy grasslands. First, our species trees

(Supplementary Fig. S1) indicate that individuals from what we call Central Madagascar – a

loose term meant only to indicate individuals from the center of the sampled *L. simplex*

distribution – represent the ancestral distribution. It is likely that *L. simplex* dominated

grasslands have origins in some of the higher elevation regions of the Central Highlands. For

example, MSV1564 comes from the Itremo protected area (Fig. 1b; Supplementary Table S1),

and further investigations of this region should be productive for characterizing natural

grasslands in Madagascar. The hypothesis of Central Highland origins, which we consider

800m in altitude and above, is also supported by our analyses of population structure (Fig. 2a;

Supplementary Fig. S2) and the weak but significant pattern of isolation-by-distance (Fig. 2b;

Supplementary Fig. S3).

Our models of N_e change through time suggests that *L. simplex* grasslands were already

extensive (Fig. 5; Supplementary Fig. S7) well before any hypothesized human presence in

Madagascar (c. 10 ka; Hansford et al., 2018) let alone human colonization (c. 2 ka; Dewar and

Wright, 1993; Dewar et al., 2013; Pierron et al., 2017). We found an approximate ten-fold

increase between one MA and the LIG which is beyond the limitations of available

paleoecological records (Burney, 1987; Gasse & Van Campo, 1998; Virah-Swamy et al., 2010;

Railsback et al., 2020; Crowley et al., 2021). Strong assumptions about the mutation rate and

generation time to calibrate analyses to absolute time and N_e were necessary, but our

interpretations of results are within the biologically plausible uncertainty of these parameters.

Even if the per-year substitution rate was overestimated by a factor of ten, the N_e increase

would still be constrained to the Late Pleistocene, occurring between the LIG and LGM. These

analyses, especially in the context of a previous investigation of population structure (Hagl et al.,

2021), suggest that *L. simplex* dominated grasslands have pre-human origins and likely

experienced some N_e increase coincident with Pleistocene climate change. Tree planting

programs have been deployed to combat the devastating habitat loss and deforestation in

Madagascar since the 1950s (Green & Sussman, 1990; Harper et al., 2007; Vieilledent et al.,

2018; Schüßler et al., 2018), with country-wide forest loss from 1953 to 2014 at 44% (Vieilledent

et al., 2018). The results of our study strongly imply that tree planting efforts should be cautious

of treating *L. simplex* dominated grasslands as degraded forests, and we suggest that focusing

resources on forest edges with native species will be more productive (Miandrimanana et al.,

2019).

Potential Use Cases of Target-Enrichment Data for Conservation Genetics

One application of target-enrichment data going forward could be analyses of population

structure and population assignment of unknown individuals. For example, many conservation

programs performing long-term biodiversity monitoring rely on reduced-SNP panels to

determine provenance of an individual or sample (e.g. Bourgeois et al., 2018; Stronen et al.,

2022). Developing SNP panels requires at least some *a priori* genomic resources, so target-

enrichment libraries could provide feasible approaches in the absence of such references and

practical improvements over existing barcoding strategies for identifying, for example,

Madagascar's rosewoods (Hassold et al., 2016); although acquiring viable DNA samples from

woody tissue would remain a barrier (Jiao et al., 2018). We anticipate limited use cases of

target-enrichment data in population genetics going forward as reference genome assembly is

becoming more accessible through decreasing long-read costs, and less complicated library

preparation protocols for whole-genome resequencing will become preferential to any target-

enrichment strategy. However, the population genetic inferences from target-enrichment data

should be biologically meaningful when available.

A Practical Need for Plant Population Genomics in Madagascar

There are few studies of plant population genetics from the Central Highlands and those are

limited to AFLP (Gardiner et al., 2017) or microsatellite data (Salmona et al., 2020; Hagl et al.,

2021). Demographic analyses from the endemic olive tree *Noronhia lowryi* suggested recent

population declines of the woody species, but it is unclear if these declines represent

anthropogenic or paleoclimatic change near the LGM (Salmona et al., 2020). While our

analyses of *L. simplex* are also ambiguous regarding this window of anthropogenic effects on

the extent of grasslands, we can at least conclude that the *L. simplex* N_e was quite large.

Interpreting what a large N_e means in the context of the *L. simplex* species distribution is difficult

though; it could be a single large distribution or multiple smaller isolated distributions that yield

the expected measures of genetic variation. Contemporary population genomic analyses will be

necessary to provide statistical power to differentiate between competing demographic

hypotheses, not only for grasses such as *L. simplex* but also species representing plant

diversity across families and life-history. A better reconstruction of the natural history of

Madagascar's grasslands is not only necessary for understanding diversity within and outside of

the Central Highlands (Yoder et al., 2016; Burbrink et al., 2019; Everson et al., 2020), but also

to make effective land management decisions for tree planting programs.

Acknowledgements

This project has received funding from the European Union's Horizon 2020 research and

innovation programme under the Marie Sklodowska-Curie grant agreement No 101026923,

awarded to GPT and MSV. GB is supported by LabEx TULIP (ANR-10-LABX-0041). CERL was

supported by a The Royal Society International Collaboration Award. Thank you to MEDD

Madagascar, Madagascar National Parks, Isalo MNP, and Itremo NPA for supporting our

research and exportation permits in 2014 and 2017, and also to COSTECH Tanzania for their

support for the 2014 collection permits for MSV and WRQL. Sequence data were generated

with research funds from Duke University to ADY. This work would not have been possible

without Stuart Cable and everyone at the Kew Madagascar Conservation Centre. We thank

Emily Lockwood for DNA extractions, and Lorena Endara and J. Gordon Burleigh for

coordinating library preparation and sequencing. Colleen Seymour provided helpful feedback on

an earlier version of this manuscript. We thank two anonymous reviewers and the editor Andrew

Young for helpful comments that improved the manuscript.

Author Contributions

GPT, ADY, and MSV conceived the study. WRQL, CLS, CERL, GB and MSV collected

samples. GPT and TOMA curated data. GPT and AAC analyzed data. GPT wrote the first draft.

TOMA translated the abstract to French and Malagasy. All authors revised the manuscript and

approved the final version.

Data Availability Statement

Scripts for analyses and input files are available through the Dryad Digital Repository:

https://doi.org/10.5061/dryad.905qfttqx. Additional scripts are available on GPT's GitHub for

identifying exon-intron boundaries from target enrichment data (https://github.com/gtiley/Locus-

Splitter) generating matrices for PCAs from fasta files (https://github.com/gtiley/fasta-

conversion) and the ploidy test implemented through PATÉ (https://github.com/gtiley/Phasing).

Raw sequence data is available via the NCBI short read archive (SRA) database, with individual

SRA and BioSample identifiers in Supplementary Table S1. Raw sequence data is associated

with NCBI BioProject PRJNA952819.

Conflict of Interest Statement

The authors declare no conflict of interest.

References

Arnold, B., Bomblies, & K.,Wakeley, J. (2012). Extending coalescent theory to autotetraploids. *Genetics, 192*(1), 195-204. https://doi.org/10.1534/genetics.112.140582

Baker, W. J., Bailey, P., Barber, V., Barker, A., Bellot, S., Bishop, D., ... & Forest, F. (2022). A comprehensive phylogenomic platform for exploring the angiosperm tree of life. *Systematic Biology*, *71*(2), 301-319. https://doi.org/10.1093/sysbio/syab035

Blischak, P.D., Kubatko, L.S., Wolfe, A.D. (2016). Accounting for genotype uncertainty in the estimation of allele frequencies in autopolyploids. *Molecular Ecology Resources, 16*(3), 742-754. https://doi.org/10.1111/1755-0998.12493

Blischak, P.D., Sajan, M., Barker, M.S., Gutenkunst, R.N. (2023). Demographic history inference and the polyploid continuum. *Genetics,* iyad107. https://doi.org/10.1093/genetics/iyad107

Bond, W. J., Silander Jr, J. A., Ranaivonasy, J., & Ratsirarson, J. (2008). The antiquity of Madagascar's grasslands and the rise of C_4 grassy biomes. *Journal of Biogeography*, *35*(10), 1743-1758. https://doi.org/10.1111/j.1365-2699.2008.01923.x

Bond, W. J., Silander, J. A., & Ratsirarson, J. (2023). Madagascar's grassy biomes are ancient and there is much to learn about their ecology and evolution. *Journal of Biogeography*, *50*(3), 614-621. https://doi.org/10.1111/jbi.14494

Bourgeois, S., Senn, H., Kaden, J., Taggart, J. B., Ogden, R., Jeffery, K. J., ... & McEwing, R. (2018). Single-nucleotide polymorphism discovery and panel characterization in the African forest elephant. *Ecology and Evolution*, *8*(4), 2207-2217. https://doi.org/10.1002/ece3.3854

Breinholt, J. W., Carey, S. B., Tiley, G. P., Davis, E. C., Endara, L., McDaniel, S. F., ... & Burleigh, J. G. (2021). A target enrichment probe set for resolving the flagellate land plant tree of life. *Applications in Plant Sciences*, *9*(1), e11406. https://doi.org/10.1002/aps3.11406

Burbrink, F. T., Ruane, S., Kuhn, A., Rabibisoa, N., Randriamahatantsoa, B., Raselimanana, A. P., ... & Raxworthy, C. J. (2019). The origins and diversification of the exceptionally rich gemsnakes (Colubroidea: Lamprophiidae: Pseudoxyrhophiinae) in Madagascar. *Systematic Biology*, *68*(6), 918-936. https://doi.org/10.1093/sysbio/syz026

Burney, D. A. (1987). Late Holocene vegetational change in Central Madagascar. *Quaternary Research*, *28*(1), 130-143. https://doi.org/10.1016/0033-5894(87)90038-X

Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., & Madden, T. L. (2009). BLAST+: architecture and applications. *BMC Bioinformatics*, *10*, 421. https://doi.org/10.1186/1471-2105-10-421

Christin, P.-A., Spriggs, E., Osborne, C. P., Strömberg, C. A., Salamin, N., & Edwards, E. J. (2014). Molecular dating, evolutionary rates, and the age of the grasses. *Systematic Biology*, *63*(2), 153-165. https://doi.org/10.1093/sysbio/syt072

Crowl, A. A., Fritsch, P. W., Tiley, G. P., Lynch, N. P., Ranney, T. G., Ashrafi, H., & Manos, P. S. (2022). A first complete phylogenomic hypothesis for diploid blueberries (*Vaccinium* section *Cyanococcus*). *American Journal of Botany*, *109*(10), 1596-1606. https://doi.org/10.1002/ajb2.16065

Crowley, B. E., Godfrey, L. R., Hansford, J. P., & Samonds, K. E. (2021). Seeing the forest for the trees—and the grasses: revisiting the evidence for grazer-maintained grasslands in Madagascar's Central Highlands. *Proceedings of the Royal Society B: Biological Sciences*, *288*(1950), 20201785. https://doi.org/10.1098/rspb.2020.1785

738
739 DePristo, M. A., Banks, E., Poplin, R., Garimella, K. V., Maguire, J. R., Hartl, C., ... & Daly, M. J. (2011). A
740 framework for variation discovery and genotyping using next-generation DNA sequencing data. *Nature*
741 *Genetics*, *43*(5), 491-498. https://doi.org/10.1038/ng.806
742
743 Dewar, R. E., Radimilahy, C., Wright, H. T., Jacobs, Z., Kelly, G. O., & Berna, F. (2013). Stone tools and
744 foraging in northern Madagascar challenge Holocene extinction models. *Proceedings of the National*
745 *Academy of Sciences*, *110*(31), 12583-12588. https://doi.org/10.1073/pnas.1306100110
746
747 Dewar, R. E., & Wright, H. T. (1993). The culture history of Madagascar. *Journal of World Prehistory*, *7*,
748 417-466. https://doi.org/10.1007/BF00997802
749
750 Dixon, A. P., Faber-Langerndoen, D., Josse, C., Morrison, J., & Loucks, C. J. (2014). Distribution
751 mapping of world grassland types. *Journal of Biogeography, 41*(11), 2003-2019.
752 https://doi.org/10.1111/jbi.12381
753
754 Doyle, J.J., Doyle, J.L. (1987). A rapid DNA isolation procedure for small quantities of fresh leaf tissue.
755 *Phytochemical Bulletin, 19*(1), 11-15.
756
757 Efron, B., & Narasimhan, B. (2020). The automatic construction of bootstrap confidence intervals. *Journal*
758 *of Computational and Graphical Statistics, 29*(3), 608-619.
759 https://doi.org/10.1080/10618600.2020.1714633
760
761 Evanno, G., Regnaut, S., & Goudet, J. (2005). Detecting the number of clusters of individuals using the
762 software structure: A simulation study. *Molecular Ecology*, 14(8), 2611–2620.
763 https://doi.org/10.1111/J.1365-294X.2005.02553.X
764
765 Everson, K. M., Jansa, S. A., Goodman, S. M., & Olson, L. E. (2020). Montane regions shape patterns of
766 diversification in small mammals and reptiles from Madagascar's moist evergreen forest. *Journal of*
767 *Biogeography*, *47*(10), 2059-2072. https://doi.org/10.1111/jbi.13945
768
769 Excoffier, L., Dupanloup, I., Huerta-Sánchez, E., Sousa, V. C., & Foll, M. (2013). Robust demographic
770 inference from genomic and SNP data. *PLoS Genetics*, *9*(10), e1003905.
771 https://doi.org/10.1371/journal.pgen.1003905
772
773 Flouri, T., Jiao, X., Rannala, B., & Yang, Z. (2020). A Bayesian implementation of the multispecies
774 coalescent model with introgression for phylogenomic analysis. *Molecular Biology and Evolution*, *37*(4),
775 1211-1223. https://doi.org/10.1093/molbev/msz296
776
777 Gardiner, L. M., Rakotoarinivo, M., Rajaovelona, L. R., & Clubbe, C. (2017). Population genetics data
778 help to guide the conservation of palm species with small population sizes and fragmented habitats in
779 Madagascar. *PeerJ*, *5*, e3248. https://doi.org/10.7717/peerj.3248
780
781 Gasse, F., & Van Campo, E. (1998). A 40,000-yr pollen and diatom record from Lake Tritrivakely,
782 Madagascar, in the southern tropics. *Quaternary Research*, *49*(3), 299-311.
783 https://doi.org/10.1006/qres.1998.1967
784
785 Gillespie, J.H. (2004). The coalescent. In, *Population Genetics: A Concise Guide* (pp.40-47). Baltimore,
786 MD. John Hopkins University Press.
787
788 Green, G. M., & Sussman, R. W. (1990). Deforestation history of the eastern rain forests of Madagascar
789 from satellite images. *Science*, *248*(4952), 212-215. https://doi.org/10.1126/science.248.4952.212
790
791 Gutenkunst, R. N., Hernandez, R. D., Williamson, S. H., & Bustamante, C. D. (2009). Inferring the joint
792 demographic history of multiple populations from multidimensional SNP frequency data. *PLoS*
793 *Genetics*, *5*(10), e1000695. https://doi.org/10.1371/journal.pgen.1000695

794
795 Hackel, J., Vorontsova, M.S., Nanjarisoa, O.P., Hall, R.C., Razanatsoa, J., Malakasi, P., & Besnard, G.
796 (2018). Grass diversification in Madagascar: In situ radiation of two large C_3 shade clades and support for
797 a Miocene to Pliocene origin of C_4 grassy biomes. *Journal of Biogeography, 45*(4), 750–761.
798 https://doi.org/10.1111/jbi.13147
799
800 Hagl, P. A., Gargiulo, R., Fay, M. F., Solofondranohatra, C., Salmona, J., Suescun, U., ... & Vorontsova,
801 M. S. (2021). Geographical structure of genetic diversity in *Loudetia simplex* (Poaceae) in Madagascar
802 and South Africa. *Botanical Journal of the Linnean Society*, *196*(1), 81-99.
803 https://doi.org/10.1093/botlinnean/boaa098
804
805 Hansford, J., Wright, P. C., Rasoamiaramanana, A., Pérez, V. R., Godfrey, L. R., Errickson, D., ... &
806 Turvey, S. T. (2018). Early Holocene human presence in Madagascar evidenced by exploitation of avian
807 megafauna. *Science Advances*, *4*(9), eaat6925. https://doi.org/10.1126/sciadv.aat6925
808
809 Harper, G. J., Steininger, M. K., Tucker, C. J., Juhn, D., & Hawkins, F. (2007). Fifty years of deforestation
810 and forest fragmentation in Madagascar. *Environmental Conservation*, *34*(4), 325-333.
811 https://doi.org/10.1017/S0376892907004262
812
813 Heled, J., & Drummond, A. J. (2008). Bayesian inference of population size history from multiple
814 loci. *BMC Evolutionary Biology*, *8*(1), 289. https://doi.org/10.1186/1471-2148-8-289
815
816 Heller, R., Chikhi, L., & Siegismund, H. R. (2013). The confounding effect of population structure on
817 Bayesian skyline plot inferences of demographic history. *PLoS One*, *8*(5), e62992.
818 https://doi.org/10.1371/journal.pone.0062992
819
820 Helmstetter, A. J., Cable, S., Rakotonasolo, F., Rabarijaona, R., Rakotoarinivo, M., Eiserhardt, W. L., ... &
821 Papadopulos, A. S. (2021). The demographic history of Madagascan micro-endemics: have rare species
822 always been rare?. *Proceedings of the Royal Society B: Biological Sciences*, *288*(1959), 20210957.
823 https://doi.org/10.1098/rspb.2021.0957
824
825 Huang, P., Feldman, M., Schroder, S., Bahri, B. A., Diao, X., Zhi, H., ... & Kellogg, E. A. (2014).
826 Population genetics of *Setaria viridis*, a new model system. *Molecular Ecology*, *23*(20), 4912-4925.
827 https://doi.org/10.1111/mec.12907
828
829 Hudson, R. R. (2002). Generating samples under a Wright–Fisher neutral model of genetic
830 variation. *Bioinformatics*, *18*(2), 337-338. https://doi.org/10.1093/bioinformatics/18.2.337
831
832 Jiao, L., Yu, M., Wiedenhoeft, A. C., He, T., Li, J., Liu, B., ... & Yin, Y. (2018). DNA barcode authentication
833 and library development for the wood of six commercial *Pterocarpus* species: the critical role of Xylarium
834 specimens. *Scientific Reports*, *8*(1), 1945. https://doi.org/10.1038/s41598-018-20381-6
835
836 Johnson, M. G., Gardner, E. M., Liu, Y., Medina, R., Goffinet, B., Shaw, A. J., ... & Wickett, N. J. (2016).
837 HybPiper: Extracting coding sequence and introns for phylogenetics from high-throughput sequencing
838 reads using target enrichment. *Applications in Plant Sciences*, *4*(7), 1600016.
839 https://doi.org/10.3732/apps.1600016
840
841 Jombart, T., & Ahmed, I. (2011). adegenet 1.3–1: New tools for the analysis of genome-wide SNP data.
842 *Bioinformatics, 27(*21), 3070–3071. https://doi.org/10.1093/bioin forma tics/btr521
843
844 Katoh, K., & Standley, D. M. (2013). MAFFT multiple sequence alignment software version 7:
845 improvements in performance and usability. *Molecular Biology and Evolution*, *30*(4), 772-780.
846 https://doi.org/10.1093/molbev/mst010
847

848 Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K., Von Haeseler, A., & Jermiin, L. S. (2017). ModelFinder:
849 fast model selection for accurate phylogenetic estimates. *Nature Methods*, *14*(6), 587-589.
850 https://doi.org/10.1038/nmeth.4285
851
852 Koechlin, J. (1993). Grasslands of Madagascar. In R. T. Coupland (Ed.), *Natural Grasslands: Eastern*
853 *Hemisphere and Résumé* (pp. 291-301). Amsterdam, Netherlands. Elsevier.
854
855 Leblois, R., Estoup, A., & Rousset, F. (2003). Influence of mutational and sampling factors on the
856 estimation of demographic parameters in a "continuous" population under isolation by distance. *Molecular*
857 *Biology and Evolution*, 20(4), 491-502. https://doi.org/10.1093/molbev/msg034
858
859 Lehmann, C. E. R., Solofondranohatra, C., Morton, J. A., Phelps, L. N., Ralimanana, H., Razanatsoa, J.,
860 Rakotoarimanana, V., Vorontsova, M.S. (2022). The grassy ecosystems of Madagascar. In S. M.
861 Goodman (Ed.), *The New Natural History of Madagascar* (pp. 152-168). Princetonm NJ. Princeton
862 University Press.
863
864 Liu, X., & Fu, Y. X. (2020). Stairway Plot 2: demographic history inference with folded SNP frequency
865 spectra. *Genome Biology*, *21*(1), 280. https://doi.org/10.1186/s13059-020-02196-9
866
867 Mazet, O., Rodríguez, W., Grusea, S., Boitard, S., & Chikhi, L. (2016). On the importance of being
868 structured: instantaneous coalescence rates and human evolution—lessons for ancestral population size
869 inference?. *Heredity*, *116*(4), 362-371. https://doi.org/10.1038/hdy.2015.104
870
871 McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., ... & DePristo, M. A.
872 (2010). The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA
873 sequencing data. *Genome Research*, *20*(9), 1297-1303. https://doi.org/10.1101/gr.107524.110
874
875 Meirmans P. G., & Van Tienderen P. H. (2013). The effects of inheritance in tetraploids on genetic
876 diversity and population divergence. *Heredity,* *110*(2), 131-137. https://doi.org/10.1038/hdy.2012.80
877
878 Miandrimanana, C., Reid, J. L., Rivoharison, T., & Birkinshaw, C. (2018). Planting position and shade
879 enchance native seedling performance in forest restoration for an endangered Malagasy plant. *Plant*
880 *Diversity,* *41*(2), 118-123. https://doi.org/10.1016/j.pld.2018.09.005
881
882 Minh, B. Q., Schmidt, H. A., Chernomor, O., Schrempf, D., Woodhams, M. D., Von Haeseler, A., &
883 Lanfear, R. (2020). IQ-TREE 2: new models and efficient methods for phylogenetic inference in the
884 genomic era. *Molecular Biology and Evolution*, *37*(5), 1530-1534.
885 https://doi.org/10.1093/molbev/msaa015
886
887 Nielsen, R. (2000). Estimation of population parameters and recombination rates from single nucleotide
888 polymorphisms. *Genetics*, *154*(2), 931-942. https://doi.org/10.1093/genetics/154.2.931
889
890 Paradis, E., & Schliep, K. (2019). ape 5.0: an environment for modern phylogenetics and evolutionary
891 analyses in R. *Bioinformatics*, *35*(3), 526-528. https://doi.org/10.1093/bioinformatics/bty633
892
893 Patton, A. H., Margres, M. J., Stahlke, A. R., Hendricks, S., Lewallen, K., Hamede, R. K., ... & Storfer, A.
894 (2019). Contemporary demographic reconstruction methods are robust to genome assembly quality: A
895 case study in Tasmanian devils. *Molecular Biology and Evolution*, *36*(12), 2906-2921.
896 https://doi.org/10.1093/molbev/msz191
897
898 Pfeifer, B., Wittelsbürger, U., Ramos-Onsins, S. E., & Lercher, M. J. (2014). PopGenome: an efficient
899 Swiss army knife for population genomic analyses in R. *Molecular Biology and Evolution*, *31*(7), 1929-
900 1936. https://doi.org/10.1093/molbev/msu136
901

902 Pierron, D., Heiske, M., Razafindrazaka, H., Rakoto, I., Rabetokotany, N., Ravololomanga, B., ... &
903 Letellier, T. (2017). Genomic landscape of human diversity across Madagascar. *Proceedings of the*
904 *National Academy of Sciences*, *114*(32), E6498-E6506. https://doi.org/10.1073/pnas.1704906114
905
906 Pritchard, J. K., Stephens, M., & Donnelly, P. (2000). Inference of population structure using multilocus
907 genotype data. *Genetics*, *155*(2), 945-959. https://doi.org/10.1093/genetics/155.2.945
908
909 Pybus, O. G., Rambaut, A., & Harvey, P. H. (2000). An integrated framework for the inference of viral
910 population history from reconstructed genealogies. *Genetics*, *155*(3), 1429-1437.
911 https://doi.org/10.1093/genetics/155.3.1429
912
913 Quéméré, E., Amelot, X., Pierson, J., Crouau-Roy, B., & Chikhi, L. (2012). Genetic data suggest a natural
914 prehuman origin of open habitats in northern Madagascar and question the deforestation narrative in this
915 region. *Proceedings of the National Academy of Sciences*, *109*(32), 13028-13033.
916 https://doi.org/10.1073/pnas.1200153109
917
918 Railsback, L. B., Dupont, L. A., Liang, F., Brook, G. A., Burney, D. A., Cheng, H., & Edwards, R. L.
919 (2020). Relationships between climate change, human environmental impact, and megafaunal extinction
920 inferred from a 4000-year multi-proxy record from a stalagmite from northwestern
921 Madagascar. *Quaternary Science Reviews*, *234*, 106244. https://doi.org/10.1016/j.quascirev.2020.106244
922
923 Rambaut, A., Drummond, A. J., Xie, D., Baele, G., & Suchard, M. A. (2018). Posterior summarization in
924 Bayesian phylogenetics using Tracer 1.7. *Systematic Biology*, *67*(5), 901-904.
925 https://doi.org/10.1093/sysbio/syy032
926
927 Rambaut, A., & Grass, N. C. (1997). Seq-Gen: an application for the Monte Carlo simulation of DNA
928 sequence evolution along phylogenetic trees. *Bioinformatics*, *13*(3), 235-238.
929 https://doi.org/10.1093/bioinformatics/13.3.235
930
931 Rannala, B., & Yang, Z. (2003). Bayes estimation of species divergence times and ancestral population
932 sizes using DNA sequences from multiple loci. *Genetics*, *164*(4), 1645-1656.
933 https://doi.org/10.1093/genet ics/164.4.1645
934
935 Ronfort, J., Jenczewski E., Bataillon, T., & Rousset, F. (1998). Analysis of population structure in
936 autotetraploid species. *Genetics,* *150*(2), 921-930. https://doi.org/10.1093/genetics/150.2.921
937
938 Salmona, J., Heller, R., Quéméré, E., & Chikhi, L. (2017). Climate change and human colonization
939 triggered habitat loss and fragmentation in Madagascar. *Molecular Ecology*, *26*(19), 5203-5222.
940 https://doi.org/10.1111/mec.14173
941
942 Salmona, J., Olofsson, J. K., Hong-Wa, C., Razanatsoa, J., Rakotonasolo, F., Ralimanana, H., ... &
943 Besnard, G. (2020). Late Miocene origin and recent population collapse of the Malagasy savanna olive
944 tree (*Noronhia lowryi*). *Biological Journal of the Linnean Society*, *129*(1), 227-243.
945 https://doi.org/10.1093/biolinnean/blz164
946
947 Solofondranohatra, C. L., Vorontsova, M. S., Hackel, J., Besnard, G., Cable, S., Williams, J., ... &
948 Lehmann, C. E. (2018). Grass functional traits differentiate forest and savanna in the Madagascar central
949 highlands. *Frontiers in Ecology and Evolution*, *6*, 184. https://doi.org/10.3389/fevo.2018.00184
950
951 Solofondranohatra, C. L., Vorontsova, M. S., Hempson, G. P., Hackel, J., Cable, S., Vololoniaina, J., &
952 Lehmann, C. E. (2020). Fire and grazing determined grasslands of central Madagascar represent ancient
953 assemblages. *Proceedings of the Royal Society B: Biological Sciences*, *287*(1927), 20200598.
954 https://doi.org/10.1098/rspb.2020.0598
955
Stronen, A. V., Mattucci, F., Fabbri, E., Galaverni, M., Cocchiararo, B., Nowak, C., ... & Caniglia, R.
(2022). A reduced SNP panel to trace gene flow across southern European wolf populations and detect

hybridization with other *Canis* taxa. *Scientific Reports*, *12*(1), 4195. https://doi.org/10.1038/s41598-022-
08132-0

Teixeira, H., Montade, V., Salmona, J., Metzger, J., Bremond, L., Kasper, T., ... & Radespiel, U. (2021).
Past environmental changes affected lemur population dynamics prior to human impact in
Madagascar. *Communications Biology*, *4*(1), 1084. https://doi.org/10.1038/s42003-021-02620-1

Tiley, G. P., Barker, M. S., & Burleigh, J. G. (2018). Assessing the performance of Ks plots for detecting
ancient whole genome duplications. *Genome Biology and Evolution*, *10*(11), 2882-2898.
https://doi.org/10.1093/gbe/evy200

Tiley, G.P., Crowl, A.A., Manos, P.S., Sessa, E.B., Solís-Lemus, C., Yoder, A.D., Burleigh, J.G. 2021.
Phasing alleles improves network inference with allopolyploids. *bioRxiv*
https://doi.org/10.1101.2021.05.04.442457

Tiley, G. P., van Elst, T., Teixeira, H., Schüßler, D., Salmona, J., Blanco, M. B., ... & Yoder, A. D. (2022).
Population genomic structure in Goodman's mouse lemur reveals long-standing separation of
Madagascar's Central Highlands and eastern rainforests. *Molecular Ecology*, *31*(19), 4901-4918.
https://doi.org/10.1111/mec.16632

Vieilledent, G., Grinand, C., Rakotomalala, F. A., Ranaivosoa, R., Rakotoarijaona, J. R., Allnutt, T. F., &
Achard, F. (2018). Combining global tree cover loss data with historical national forest cover maps to look
at six decades of deforestation and forest fragmentation in Madagascar. *Biological Conservation*, *222*,
189-197. https://doi.org/10.1016/j.biocon.2018.04.008

Virah-Sawmy, M., Willis, K. J., & Gillson, L. (2010). Evidence for drought and forest declines during the
recent megafaunal extinctions in Madagascar. *Journal of Biogeography*, *37*(3), 506-519.
https://doi.org/10.1111/j.1365-2699.2009.02203.x

Viruel, J., Conejero, M., Hidalgo, O., Pokorny, L., Powell, R. F., Forest, F., ... & Leitch, I. J. (2019). A
target capture-based method to estimate ploidy from herbarium specimens. *Frontiers in Plant Science*,
937. https://doi.org/10.3389/fpls.2019.00937

Vorontsova, M. S., Besnard, G., Forest, F., Malakasi, P., Moat, J., Clayton, W. D., ... & Linder, H. P.
(2016). Madagascar's grasses and grasslands: anthropogenic or natural?. *Proceedings of the Royal
Society B: Biological Sciences*, *283*(1823), 20152262. https://doi.org/10.1098/rspb.2015.2262

Wang, L., Beissinger, T. M., Lorant, A., Ross-Ibarra, C., Ross-Ibarra, J., & Hufford, M. B. (2017). The
interplay of demography and selection during maize domestication and expansion. *Genome
Biology*, *18*(1), 215. https://doi.org/10.1186/s13059-017-1346-4

Weiß, C. L., Pais, M., Cano, L. M., Kamoun, S., & Burbano, H. A. (2018). nQuire: a statistical framework
for ploidy estimation using next generation sequencing. *BMC Bioinformatics*, *19*(1), 122.
https://doi.org/10.1186/s12859-018-2128-z

Xie, M., Wu, Q., Wang, J., & Jiang, T. (2016). H-PoP and H-PoPG: heuristic partitioning algorithms for
single individual haplotyping of polyploids. *Bioinformatics*, *32*(24), 3735-3744.
https://doi.org/10.1093/bioinformatics/btw537

Yamaura, Y., Narita, A., Kusumoto, Y., Nagano, A. J., Tezuka, A., Okamoto, T., ... & Lindenmayer, D.
(2019). Genomic reconstruction of 100 000-year grassland history in a forested country: population
dynamics of specialist forbs. *Biology Letters*, *15*(5), 20180577. https://doi.org/10.1098/rsbl.2018.0577

Yoder, A. D., Campbell, C. R., Blanco, M. B., Dos Reis, M., Ganzhorn, J. U., Goodman, S. M., ... &
Weisrock, D. W. (2016). Geogenetic patterns in mouse lemurs (genus *Microcebus*) reveal the ghosts of

1013 Madagascar's forests past. *Proceedings of the National Academy of Sciences*, *113*(29), 8049-8056.
1014 https://doi.org/10.1073/pnas.1601081113

1015

Figures

Figure 1 — *Loudetia simplex* distribution and single Madagascar origin. a) A field of *L.*

simplex near Antsirabe Madagascar planted with eucalyptus (photo credit: George P. Tiley). b)

Our sampling of *L. simplex* was based on collections made throughout Madagascar's Central

Highland Plateau, which is typically considered an altitude between 800 and 1800 meters. c)

Phylogenetic relationships of individuals based on the phased (PH) data. Numbers next to

nodes are posterior probabilities. ZA = South Africa, TZ = Tanzania, BI = Burundi.

1025

1026

1027

1028

1029

Figure 2 — Patterns of genetic structure and variation in Madagascar. a) Aggregated

STRUCTURE results across runs for an *a priori* number of clusters of two (*K* = 2) for the

different data types. c) A plot of natural-log-scaled geographic distance against the pairwise

distance (p-distance). Lines are from a least-squares linear regression with the coefficient of

determination (R^2) shown for each data type. Statistical significance was determined with a

Mantel test using 9999 permutations. Significance levels are *p*-value < 0.05 (*), *p*-value < 0.01

(**), and *p*-value < 0.001 (***). Colors represent different data types: one haplotype (OH),

ambiguous genotypes (AG), and phased haplotypes (PH).

1038

1039

1040 **Figure 3 — Principal component analyses across data types.** a) All individuals using haploid

1041 (OH) data. b) Only Malagasy individuals using OH data. c) All individuals treating all sites as

1042 diploid. d) Only Malagasy individuals treating all sites as hexaploid. e) Only Malagasy

1043 individuals compressing unbalanced allelic dosage to single categories. f) Only Malagasy

1044 individuals treating all sites as diploid.

1045

1046

1047

1048

1049 **Figure 4 — Distributions of summary statistics across locus regions.** Box-and-whisker

1050 plots show distribution medians and quartiles for the expected pairwise nucleotide divergence

1051 (π), the number of segregating sites (S), and a normalized measure of the difference between

1052 two estimators of nucleotide divergence, indicating deviations from neutral expectations

1053 (Tajima's D).

1054

1055

1056

1057

1058 **Figure 5 — Extended Bayesian Skyline Plot results with fixed clock rates.** Two

1059 independent runs are shown with mean population sizes shown by solid lines. Polygons show

1060 the 95% HPD intervals with the overlap between runs in purple. Results are shown for the a)

1061 OH, b) AG, and c) PH data. Relevant time periods are shown in gray, which includes likely

1062 human colonization (LHC), the last glacial maximum (LGM), and last interglacial period (LIG).

1063

1064

1065

1066

1067

1068 **Figure 6 — Simulation results from EBSP analyses for an appropriate number of change**

1069 **points.** Individual red lines are the mean N_e estimates from 100 simulated data sets. Blue points

1070 and vertical lines are the mean of mean N_e estimates and their 95% HPD intervals. The dashed

1071 black line is the true demographic history of the population.

1072

1073 **Tables**

1074

Table 1 — X^2 test result from Tajima's D statistics across datasets and locus regions

Tested Effect	Fixed Effect	X^2	p-value	Largest Residual
Region	OH	7.72	0.27	Under-representation of Tajima's D < -2 in core regions
	AG	37.77	0.0001	Over-representation of Tajima's D < -2 in core regions
	PH	83.47	0.0001	Over-representation of Tajima's D < -2 in core regions
Dataset	Left Flank	12.65	0.046	Over-representation of Tajima's D < -2 in OH dataset
	Right Flank	8.61	0.19	Over-representation of Tajima's D < -2 in OH dataset
	Core	57.15	0.0001	Over-representation of Tajima's D < -2 in OH dataset

1075

1076