Dynamical BLUP modeling with

Robertson-Price updating of mean reaction norm parameter values

Running head: BLUP modeling with Robertson-Price updating

Rolf Ergon

University of South-Eastern Norway

Porsgrunn, Norway

April 12, 2023

rolf.ergon@usn.no

https://orcid.org/0000-0003-2393-2997

Keywords Dynamical BLUP; Mean reaction norm parameter updating; Microevolution vs. plasticity disentanglement; Random effect variance estimation; Robertson-Price identity; Robertson's secondary theorem of natural selection.

Abstract

A general reaction norm model of any order can be formulated as a linear mixed model. From

this follows that estimates of mean phenotypic traits in a population (fixed effects), and

predictions of individual additive genetic deviations from mean reaction norm parameter

values (random effects), can be found from the best linear unbiased predictions (BLUP)

equation in matrix form. The resulting BLUP model is dynamical in the sense that the

incidence matrix varies with time. This leads to a straightforward and multivariate alternative

to infinite-dimensional and random regression modeling of reaction norms.

Based on such a BLUP model, the within-generation changes in predicted mean reaction

norm parameter values can be found by use of the Robertson-Price identity, applied on the

predicted random effects. From this follows that the between-generation change in the mean

values are found from Robertson's secondary theorem of natural selection, applied on the

predicted random effects. This explains why and to which extent the variances of BLUP

random effects are underestimated, which is a well-known observation.

The dynamical BLUP model will thus produce the mean reaction norms over time, which

makes it possible to disentangle the microevolutionary and plasticity components in for

example climate change responses. The BLUP responses will depend on the additive genetic

relationship matrix $\boldsymbol{A}_t$. When $\boldsymbol{A}_t$ is an identity matrix, the results will be identical to the

results from a variant of the multivariate breeder's equation, based on the selection gradient

with respect to the individual phenotypic trait values.

Parameters are assumed to be known and constant, but it is discussed how they can be

estimated by means of a prediction error method. Generations are assumed to be non-

overlapping, but adjustments for overlapping generations can easily be done.

1. Introduction

A reaction norm describes the phenotypes that a genotype can produce across a range of environments. Mean reaction norms in a population can evolve, and this evolution can be modelled by use of several methods. In its simplest form a mean reaction norm is characterized by an intercept value and a plasticity slope value, but it may also be appropriate to use models with a multiple of reaction norms, not necessarily linear.

Reaction norms may evolve, and a common evolutionary model is the multivariate breeder's equation (Lande, 1979), where the mean reaction norm parameters may be treated as traits in their own right, as in, e.g., Lande (2009). For applications on field data from studies of wild populations, there are two problems with such models. First, the individual reaction norm parameters are not available, and second, additive genetic relationships in the population cannot be taken into account. The first problem can be solved by a linear transformation as shown in Ergon (2022a), and as used for comparison purposes in Section 2, but the second problem will still exist.

An alternative approach for reaction norm evolutionary modeling is to use infinite-dimensional characters, as in a method introduced by Kirkpatrick and Heckman (1989), and as applied on reaction norms by Gomulkiewicz and Kirkpatrick (1992). It is then necessary to obtain smooth covariance function estimates (Kingsolver et al, 2001), and one method for that purpose is random regression (Shaeffer, 2004), where individual response curves are modelled as relatively simple weighted sums of basis functions. Additive genetic relationship matrices can be included in such models (Oliviera et al., 2019). Obvious difficulties are here the need to find for example polynomial basis functions that fit the data reasonably well over time, and that the reaction norms in multivariate cases are correlated, such that they cannot be modelled independently.

The present article introduces a multivariate modeling approach based on best linear

unbiased predictions (BLUP), allowing for multiple (and potentially correlated) traits to have

joint norms of reaction, which can be linear or approximated by power series (Gavrilets and

Scheiner, 1993). This is a dynamical BLUP model in the sense that the incidence matrix for

the random effects is a function of the environment. We will here develop the theory under

the assumption that the parameters in the model are known, but as will be shown separately

they can also be identified by a prediction error method, as introduced in a microevolutionary

context in Ergon (2022a,b).

Best linear unbiased predictions (BLUP) utilizing linear mixed models with fixed and

random effects, are extensively used in domestic animal and plant breeding (Robinson, 1991;

Ch. 26, Lynch and Walsh, 1998; Arnold et al., 2019). These methods may also be applied on

wild populations (Kruuk, 2004; Nussey et al., 2007), although such uses have been criticized

owing to errors in estimated variances of the random effects (Hadfield et al., 2010). An

important application is the disentanglement of microevolutionary and plasticity components

in for example climate change responses (Merilä and Hendry, 2014; Ergon, 2022a,b). The

basic BLUP equations was first developed in summation form (Henderson, 1950), but as done

here it is more convenient to use matrix formulations.

In this article we will study how mean reaction norm parameter values, and thus mean

phenotypic traits, evolve under the influence of environmental cues and changes of the fitness

landscape. Such evolution of reaction norms and traits seeks to maximize the mean fitness of

a given population, and changes in the location of fitness peaks in the phenotypic space are

therefore the driving force. We will thus study the dynamics of microevolutionary systems,

and we will mainly do so by use of BLUP, but also with reference to the well-known

multivariate breeder's equation (Lande, 1979). Although fitness can be defined as the long-

run growth rate (Sæther and Engan, 2015), or for non-overlapping generations the expected

geometric mean fitness (Autzen and Okasha, 2022), we will in simulations simply use the number of surviving descendants as a measure of individual fitness (Ch. 6, Rice, 2004).

It is well known that BLUP underestimates the variances of the random effects in linear mixed models (Ch. 26, Lynch and Walsh, 1998; Hadfield et al., 2010). Here, we will see why and to which extent that is necessary in order to obtain the correct incremental changes in mean reaction norm parameter values from parents to offspring (within-generation changes), and that these changes may be found from the well-known Roberson-Price identity (Robertson, 1966; Price, 1970) applied on the predicted random effects. From this follows that the between-generation changes in the mean values follow from Robertson's secondary theorem of natural selection (Robertson, 1966; Ch. 6, Walsh and Lynch, 2018), applied on the predicted random effects. This also applies to non-plastic organisms, where the mean reaction norms degenerate into mean phenotypic trait values (Ergon, 2022c). It is worth noting that the Robinson-Price identity and Robertson's secondary theorem in this context are applied on the predicted random effects, and not on the true additive genetic effects as normally done (Ch. 6, Walsh and Lynch, 2018).

The dynamical BLUP model with Robertson-Price updating of mean reaction norm parameter values, will make use of the additive genetic relationship matrix $\boldsymbol{A}_t$ in a standard way (Ch. 26, Lynch and Walsh, 1998). The theoretical treatment is limited to cases where only mean phenotypic traits are included in the fixed effects, and it assumes that generations are non-overlapping. It is, however, also shown how cases with overlapping generations can be handled. In cases with sexual reproduction, we will assume a hypothetical single parent (mid-parent) occupying an intermediate phenotypic position between the two parents (Ch. 7, Rice, 2004).

The theoretical development will be general, i.e., for any number of phenotypic traits and any number of environmental cues in the model. For clarity of presentation, however, some

details will be given for a system with only two phenotypic traits and only two environmental

cues. A similar limited system will also be used in simulations.

The additive genetic and phenotypic covariance matrices, G and P, are here assumed to be

constant and known. They may, however, be estimated by means of a prediction error method

(PEM), utilizing information contained in environmental cues and individual phenotypic trait

values over many generations, as well as fitness information (Ergon, 2022a,b). With plastic

traits in the dynamical BLUP model, restricted maximum likelihood (REML) methods

applied on data from a single generation cannot be used for this purpose. The simple reason

for this is that each element in the linear mixed model residual matrix is a function of several

non-additive effects, such that the REML equations become indeterminate.

As developed theoretically, and verified in simulations, the dynamical BLUP model with

an additive genetic relationship matrix equal to an identity matrix, will give the same results

as a selection gradient prediction method (GRAD) based on the multivariate breeder's

equation (Ergon, 2022a,b). For large populations, these results will asymptotically also be the

same as from the multivariate breeder's equation directly.

After this introduction, Theory and Methods follow in Section 2, Simulations in Section 3,

and Summary and Discussion in Section 4. Proofs of two theorems are given in Appendices A

and B. MATLAB code for the simulations are archived as Supplementary Material on

bioRxiv.

2. Theory and Methods

2.1 Notation

Mathematical symbols with descriptions in the order they appear in equations are shown in

Table 1.

Table 1. Mathematical symbols with description

Symbol	Description
$\Delta \bar{z}_t$	Incremental change in mean trait from generation t to generation $t+1$
$w_{i,t}$	Relative individual fitness, $w_{i,t} = W_{i,t}/\overline{W}_t$, as function of time
$z_{i,t}, y_{i,t}, \mathbf{z}_{i,t}, \mathbf{y}_{i,t}$	Individual traits, and vectors of individual traits, as functions of time
$\mathbf{G}$	Additive genetic covariance matrix, with block elements $\mathbf{G}_{aa}, \mathbf{G}_{ab}, \mathbf{G}_{bb}$
$\mathbf{P}$	Phenotypic covariance matrix, with block elements $\mathbf{P}_{aa}, \mathbf{P}_{ab}, \mathbf{P}_{bb}$
$\boldsymbol{\beta}_t$	Selection gradient as function of time
$a'_{i,t} + v_{i,t}$	Individual intercept deviations around mean values $\bar{a}_t$ and 0, respectively, with additive and non-additive terms
$b'_{i,t} + \eta_{i,t}$	Individual plasticity slope deviations around mean values $\bar{b}_t$ and 0, respectively, with additive and non-additive terms
$\Delta \bar{a}_t, \Delta \bar{b}_t$	Incremental changes in mean reaction norm parameter values
$\sigma_v^2, \sigma_\eta^2$	Variances of non-additive effects in reaction norm models
$u_t, \mathbf{u}_t$	Environmental variable, and vector of environmental variables
$P_{yy,t}$	Variance of $y_{i,t}$
$\mathbf{X}$	Design matrix in linear mixed model
$\widetilde{\mathbf{Z}}_t = \mathbf{Z}_t \otimes \mathbf{I}_n$	Incident matrix in linear mixed model, with $\mathbf{Z}_t = f(\mathbf{u}_t)$
$\mathbf{x}_t$	Random effects in linear mixed model, for special case with 2 traits and 2 env. variables, $\mathbf{x}_t = \begin{bmatrix} \mathbf{a}'^T_{1,t} & \mathbf{a}'^T_{2,t} & \mathbf{b}'^T_{11,t} & \mathbf{b}'^T_{12,t} & \mathbf{b}'^T_{21,t} & \mathbf{b}'^T_{22,t} \end{bmatrix}^T$
$\mathbf{e}_t$	Residual vector in mixed model
$\mathbf{A}_t$	Additive genetic relationship matrix
$\widetilde{\mathbf{G}}_t = \mathbf{G} \otimes \mathbf{A}_t$	Kronecker covariance matrix for BLUP model
$\widetilde{\mathbf{R}}_t = \mathbf{R}_t \otimes \mathbf{I}_n$	Kronecker residual matrix for BLUP model, with $\mathbf{R}_t = E[\mathbf{e}_t \mathbf{e}_t^T]$
$\widehat{z'}_{i,t}$	Predicted individual intercept or slope deviations around mean values
$\bar{z}_t^{parents}, \bar{z}_t^{offspring}$	Mean reaction norm parameter values for parents and offspring

2.2 Introductory example

For a simple toy example, intended to ease the readers into the concepts used below, consider a single trait $y_{i,t}$ measured on a single individual. In this case, the BLUP estimate of the true additive genetic value for that individual (the deviation from zero) is $\widehat{a}'_{i,t} = h^2(y_{i,t} - mean)$, where h^2 is the heritability, while $(y_{i,t} - mean)$ denotes the phenotypic deviation from the mean. If we substitute this estimate into Robertson's secondary theorem of natural selection, we find the between-generation response $R = cov(w_{i,t}, \widehat{a}'_{i,t}) = h^2 cov(w_{i,t}, y_{i,t}) = h^2 S$, where $w_{i,t}$ is the relative fitness, while S is the within-generation change in the mean. In this way we recover the standard univariate breeder's equation.

What is done below is to consider a much more complicated phenotype (an observed vector $\boldsymbol{y}_t$ of individual focal traits with reaction norms) and use BLUP to estimate the vector of additive effects associated with the norm of reaction functions, with these BLUPs then substituted into the expression for Robertson's secondary theorem of natural selection. The incremental changes in the mean reaction norm parameter values thus follow from $cov(w_{i,t}, \widehat{z}'_{i,t})$, where $\widehat{z}'_{i,t}$ stands for the BLUP estimates of the the true additive genetic values involved (Eq. (11) below).

For the special case with an additive genetic relationship matrix $\boldsymbol{A}_t = \boldsymbol{I}_n$, the BLUP estimates yield a matrix-based inheritance expression (using the correlated nature of the random effects) to replace h^2, and a Robertson-Price term $cov(w_{i,t}, y_{i,t})$ to measure phenotypic selection (Eq. (12) below). From a practical point of view, Eq. (12) is unnecessary, but it is included for the purpose of comparisons with results from a variant of the multivariate breeder's equation.

2.3 Background theory

For the development of the dynamical BLUP matrix equation that follows, we need some background theory. First, the Price equation for selection in a population with n individuals says that the evolution of the mean trait of an $n \times 1$ vector $\mathbf{z}_t$ of individual quantitative traits is described by (Price, 1970, 1972)

$$\Delta \bar{z}_t = cov(w_{i,t}, z_{i,t}) + E\left[w_{i,t}\left(z_{i,t}^{descendants} - z_{i,t}\right)\right], \tag{1}$$

where $\Delta \bar{z}_t = \bar{z}_{t+1} - \bar{z}_t$ is the incremental change in mean trait value from generation to generation, and where $z_{i,t}$ is an individual trait. Here, $w_{i,t}$ is the relative individual fitness, i.e., individual fitness divided by the mean fitness in the population, while $z_{i,t}^{descendants}$ is the mean trait of the descendants (and the parent if it survives) of individual i in generation t. The trait $z_{i,t}$ may be any property we can assign a numerical value to, not necessarily biological. In a biological context the trait may be a behavioral, morphological or physiological characteristic, but it may also be a parameter in a reaction norm model that describes a plastic organism. Disregarding the second term on the righthand side of Eq. (1), we find the Robertson-Price identity, $\Delta \bar{z}_t = cov(w_{i,t}, z_{i,t})$, which we will use in connection with Assumption 7 below. If we here replace $z_{i,t}$ with its additive genetic component, $x_{i,t}$, we find Robertson's secondary theorem of natural selection, which we will use in Theorem 1 below.

Second, we need to see how the multivariate breeder's equation (Lande, 1979; Lande and Arnold, 1983),

$$\Delta \bar{\mathbf{z}}_t = \mathbf{G}\mathbf{P}^{-1}cov(w_{i,t}, \mathbf{z}_{i,t}) = \mathbf{G}\boldsymbol{\beta}_t, \tag{2}$$

where $\boldsymbol{\beta}_t$ is the selection gradient, can be applied on the parameters in a reaction norm model. Eq. (2) was derived from a multivariate version of Eq. (1), which requires several assumptions, as detailed in Ergon (2019,2022c):

1. The vector $\boldsymbol{z}_{i,t}$ of individual phenotypic traits is the sum of independent additive

genetic effects $\boldsymbol{x}_{i,t}$ and non-additive environmental and genetic effects $\boldsymbol{e}_{i,t}$, i.e.,

$\boldsymbol{z}_{i,t} = \boldsymbol{x}_{i,t} + \boldsymbol{e}_{i,t}.$

2. The non-additive effects $\boldsymbol{e}_{i,t}$ are zero mean, independent and identically distributed

(iid) random variables.

3. There are no expected fitness weighted changes in the individual additive genetic

effects $\boldsymbol{x}_{i,t}$ from one generation to the next besides selection, i.e.,

$E\left[w_{i,t}\left(\boldsymbol{x}_{i,t}^{descendants} - \boldsymbol{x}_{i,t}\right)\right] = \boldsymbol{0}.$

4. The additive genetic effects $\boldsymbol{x}_{i,t}$, and the environmental effects $\boldsymbol{e}_{i,t}$ and $\boldsymbol{e}_{i,t}^{descendants}$,

are multivariate normal.

5. The additive genetic effects $\boldsymbol{x}_{i,t}$ and non-additive effects $\boldsymbol{e}_{i,t}$ and $\boldsymbol{e}_{i,t}^{descendants}$

influence individual fitness only through $\boldsymbol{z}_{i,t}$.

6. All individuals in the population are genetically unrelated, which means that the

additive genetic relationship matrix $\boldsymbol{A}_t$ is a unity matrix.

In what follows, we will make use of Assumptions 1, 2, and 3, while Assumptions 4, 5, and 6

will be used only indirectly when the BLUP results with $\boldsymbol{A}_t = \boldsymbol{I}_n$ are compared with results

based on the multivariate breeder's equation.

In order to see how Eq. (2) can be applied on the parameters in a reaction norm model, we

may use an individual intercept-slope model based on Assumptions 1 and 2 above (Lande,

2009),

$$y_{i,t} = \bar{a}_t + a'_{i,t} + v_{i,t} + \left(\bar{b}_t + b'_{i,t} + \eta_{i,t}\right)u_t = \bar{y}_t + a'_{i,t} + b'_{i,t}u_t + v_{i,t} + \eta_{i,t}u_t, \qquad (3)$$

where u_t is an environmental cue, and where the mean reaction norm is $\bar{y}_t = \bar{a}_t + \bar{b}_t u_t$. Here,

$a_{i,t} = \bar{a}_t + a'_{i,t} + v_{i,t}$ and $b_{i,t} = \bar{b}_t + b'_{i,t} + \eta_{i,t}u_t$ are the individual intercept and slope

parameters, where $v_{i,t}$ and $\eta_{i,t}$ are iid and zero mean random variables, with variances σ_v^2 and

σ_η^2, respectively. We thus use $a'_{i,t} + v_{i,t}$ and $b'_{i,t} + \eta_{i,t}$ to denote individual deviations from

mean values $\bar{a}_t$ and $\bar{b}_t$, respectively, where $a'_{i,t}$ and $b'_{i,t}$ are the additive genetic components

of these deviations. Such additive genetic deviations will be the random effects in the linear

mixed model developed below, and thus the random effects that are predicted by use of the

BLUP equations.

When the reaction norm parameters in Eq. (3) are treated as quantitative traits in their own

right, Eq. (2) leads to

$$\begin{bmatrix} \Delta\bar{a}_t \\ \Delta\bar{b}_t \end{bmatrix} = \boldsymbol{GP}^{-1} \begin{bmatrix} cov(w_{i,t}, a_{i,t}) \\ cov(w_{i,t}, b_{i,t}) \end{bmatrix}, \tag{4}$$

with $\boldsymbol{G}$ and $\boldsymbol{P}$ are given by $\boldsymbol{G} = E\left[\begin{bmatrix} a'_{i,t} \\ b'_{i,t} \end{bmatrix} [a'_{i,t} \quad b'_{i,t}] \right] = \begin{bmatrix} G_{aa} & G_{ab} \\ G_{ab} & G_{bb} \end{bmatrix}$ and

$\boldsymbol{P} = \begin{bmatrix} G_{aa} + \sigma_v^2 & G_{ab} \\ G_{ab} & G_{bb} + \sigma_\eta^2 \end{bmatrix}$. The additive genetic and phenotypic covariance matrices $\boldsymbol{G}$ and

$\boldsymbol{P}$ may by time-varying, but for simplicity we will here assume that they are constant. As

discussed in Ergon (2022a), it is essential that the environmental input in Eq. (3) has a proper

reference value, and for simplicity we here assume an environmental scale such that the

reference environment is zero.

For comparisons with BLUP results, we finally need a modified version of the multivariate

breeder's equation. As shown in Ergon (2022a), Eq. (4) can by use of a linear transformation

of the vector $[a_{i,t} \quad b_{i,t}]^T$ onto the vector $[a_{i,t} \quad b_{i,t} \quad y_{i,t}]^T$ be reformulated into the selection

gradient (GRAD) form,

$$\begin{bmatrix} \Delta\bar{a}_t \\ \Delta\bar{b}_t \end{bmatrix} = \begin{bmatrix} G_{aa} & G_{ab} \\ G_{ab} & G_{bb} \end{bmatrix} \begin{bmatrix} 1 \\ u_t \end{bmatrix} P_{yy,t}^{-1} cov(w_{i,t}, y_{i,t}), \tag{5}$$

where selection with respect to $a_{i,t}$ and $b_{i,t}$, as in Eq. (4), is replaced by selection with respect

to $y_{i,t}$. Here, $P_{yy,t} = P_{aa} + 2G_{ab}u_t + P_{bb}u_t^2$, while $P_{yy,t}^{-1} cov(w_{i,t}, y_{i,t})$ is the selection

gradient. It is essential to note that Eqs. (4) and (5) give identical results only asymptotically,

when the population size $n \to \infty$, and the reason for that is the differences in how the

covariance functions are used in the two equations. As we will see, extensions of Eq. (5) are

possible for more complex reaction norm models. This is interesting because with an additive

genetic relationship matrix $\boldsymbol{A}_t = \boldsymbol{I}_n$, the dynamical BLUP model we will develop results in

incremental changes in mean reaction norm parameter values, that are identical to those found

from an extended version of Eq. (5). Here, we should finally note that since Eq. (5) is derived

from the multivariate breeder's equation (2), it is valid only under Assumptions 1 to 6 above.

2.4 Development of the dynamical BLUP model

233   For clarity of presentation, some details will here be limited to a system with $p = 2$

234   phenotypic traits, and $q = 2$ environmental cues, and a similar simplified system will also be

235   used in the simulations. The theory will, however, be developed in such a way that extensions

236   to higher dimensions are obvious.

237   For a specific trait j, the individual reaction norm model with $q = 2$ environmental cues is

$$y_{j,i,t} = \bar{a}_{j,t} + a'_{j,i,t} + v_{j,i,t} + \left(\bar{b}_{j1,,t} + b'_{j1,i,t} + \eta_{j1,i,t}\right)u_{1,t}$$

$$+ \left(\bar{b}'_{j2,t} + b'_{j2,i,t} + \eta_{j2,i,t}\right)u_{2,t}$$

$$= \bar{y}_{j,t} + a'_{j,i,t} + b'_{j1,i,t}u_{1,t} + b'_{j2,i,t}u_{2,t} + v_{j,i,t} + \eta_{j1,i,t}u_{1,t} + \eta_{j2,i,t}u_{2,t}, \tag{6a}$$

238   where $\bar{a}_{j,t} + a'_{j,i,t} + v_{j,i,t}$, $\bar{b}_{j1,t} + b'_{j1,i,t} + \eta_{j1,i,t}u_{1,t}$ and $\bar{b}_{j2,t} + b'_{j2,i,t} + \eta_{j2,i,t}u_{2,t}$ are the

239   individual parameter values, while

$$\bar{y}_{j,t} = \bar{a}_{j,t} + \bar{b}_{j1,t}u_{1,t} + \bar{b}_{j2,t}u_{2,t} \tag{6b}$$

240   is the mean trait value. For a population with n individuals we may collect $y_{j,i,t}$, $a'_{j,i,t}$, etc., in

241    $n \times 1$ vectors, and obtain the individual trait vector for trait j,

$$\boldsymbol{y}_{j,t} = \boldsymbol{1}_n\bar{y}_{j,t} + \boldsymbol{a}'_{j,t} + \boldsymbol{b}'_{j1,t}u_{1,t} + \boldsymbol{b}'_{j2,t}u_{2,t} + \boldsymbol{v}_t + \boldsymbol{\eta}_{j1,t}u_{1,t} + \boldsymbol{\eta}_{j2,t}u_{2,t}. \tag{6c}$$

242 With $p = 2$ traits and $q = 2$ environmental cues, we thus obtain the linear mixed model in

243 general form (Ch. 26, Lynch and Walsh 1998) (with $\bar{\boldsymbol{y}}_t$ as fixed effects vector)

$$\boldsymbol{y}_t = \boldsymbol{X}\bar{\boldsymbol{y}}_t + \tilde{\boldsymbol{Z}}_t\boldsymbol{x}_t + \boldsymbol{e}_t, \tag{7}$$

with $\boldsymbol{y}_t = \begin{bmatrix} \boldsymbol{y}_{1,t}^T & \boldsymbol{y}_{2,t}^T \end{bmatrix}^T$, $\bar{\boldsymbol{y}}_t = \begin{bmatrix} \bar{y}_{1,t} & \bar{y}_{2,t} \end{bmatrix}^T$,

$\boldsymbol{x}_t = \begin{bmatrix} \boldsymbol{a}'^T_{1,t} & \boldsymbol{a}'^T_{2,t} & \boldsymbol{b}'^T_{11,t} & \boldsymbol{b}'^T_{12,t} & \boldsymbol{b}'^T_{21,t} & \boldsymbol{b}'^T_{22,t} \end{bmatrix}^T$, and $\boldsymbol{e}_t = \begin{bmatrix} \boldsymbol{e}_{1,t}^T & \boldsymbol{e}_{2,t}^T \end{bmatrix}^T$, where

$\boldsymbol{e}_{1,t} = \boldsymbol{v}_{1,t} + \boldsymbol{\eta}_{11,,t}\boldsymbol{u}_{1,t} + \boldsymbol{\eta}_{12,t}\boldsymbol{u}_{2,t}$, and $\boldsymbol{e}_{2,t} = \boldsymbol{v}_{2,t} + \boldsymbol{\eta}_{21,,t}\boldsymbol{u}_{1,t} + \boldsymbol{\eta}_{22,t}\boldsymbol{u}_{2,t}$. Here,

$\boldsymbol{X} = \begin{bmatrix} \boldsymbol{1}_n & \boldsymbol{0} \\ \boldsymbol{0} & \boldsymbol{1}_n \end{bmatrix}$, with dimension $pn \times p = 2n \times 2$, while $\tilde{\boldsymbol{Z}}_t$ has dimension

$pn \times p(1 + q)n = 2n \times 6n$. In more detail we have $\tilde{\boldsymbol{Z}}_t = \boldsymbol{Z}_t \otimes \boldsymbol{I}_n$, where

$\boldsymbol{Z}_t = \begin{bmatrix} 1 & 0 & u_{1,t} & u_{2,t} & 0 & 0 \\ 0 & 1 & 0 & 0 & u_{1,t} & u_{2,t} \end{bmatrix}$, and where $\otimes$ is the Kronecker product operator, which

means that all elements in $\boldsymbol{Z}_t$ should be multiplied by $\boldsymbol{I}_n$. For use in Appendix B we may note

that $\boldsymbol{Z}_t = \begin{bmatrix} \boldsymbol{I}_p & \boldsymbol{U}_t^T \end{bmatrix}$, with $\boldsymbol{U}_t = \begin{bmatrix} u_{1,t} & u_{2,t} & 0 & 0 \\ 0 & 0 & u_{1,t} & u_{2,t} \end{bmatrix}^T$. It is an essential feature of the

model in Eq. (7) that $E[\boldsymbol{a}'_{j,t}] = E[\boldsymbol{e}_{j,t}] = \boldsymbol{0}$, such that $E[a'_{j,i,t}] = E[e_{j,i,t}] = 0$ for $j = 1$ to p,

and that $E[\boldsymbol{b}'_{jk,t}] = \boldsymbol{0}$, such that $E[b'_{jk,i,t}] = 0$ for $j = 1$ to p and $k = 1$ to q (Ch. 26, Lynch

and Walsh 1998). We thus also have that $E[y_{j,i,t}] = \bar{y}_{j,t}$.

The random effects in Eq. (7) may all be correlated, with an additive genetic covariance

matrix $\boldsymbol{G} = E[\boldsymbol{x}_t \boldsymbol{x}_t^T] = \begin{bmatrix} \boldsymbol{G}_{aa} & \boldsymbol{G}_{ab} \\ \boldsymbol{G}_{ab}^T & \boldsymbol{G}_{bb} \end{bmatrix}$. The residuals $\boldsymbol{e}_{1,t}$ and $\boldsymbol{e}_{2,t}$ in Eq. (7) are assumed to be

uncorrelated, with a covariance matrix $\boldsymbol{R}_t = E[\boldsymbol{e}_t \boldsymbol{e}_t^T] = \begin{bmatrix} r_{1,t} & 0 \\ 0 & r_{2,t} \end{bmatrix}$, where

$r_{1,t} = \sigma_{v_1}^2 + u_{1,t}^2 \sigma_{\eta_{11}}^2 + u_{2,t}^2 \sigma_{\eta_{12}}^2$ and $r_{2,t} = \sigma_{v_2}^2 + u_{1,t}^2 \sigma_{\eta_{21}}^2 + u_{2,t}^2 \sigma_{\eta_{22}}^2$, with $\sigma_{v_j}^2$, $\sigma_{\eta_{j1}}^2$ and $\sigma_{\eta_{j2}}^2$

as the variances of the non-additive effects $v_{j,i,t}$, $\eta_{j1,i,t}$ and $\eta_{j2,i,t}$ according to Eq. (6a). Also

$\boldsymbol{G}$ may in general be a function of time, but we will here assume that it is constant. Also $\sigma_{v_1}^2$,

$\sigma_{v_2}^2$, $\sigma_{\eta_{11}}^2$, $\sigma_{\eta_{12}}^2$, $\sigma_{\eta_{21}}^2$ and $\sigma_{\eta_{22}}^2$ are assumed to be constant. We will assume that the parameters

in G and R_t are known, although they may in practice be estimated by use of restricted

maximum likelihood (REML) (Ch. 27, Lynch and Walsh 1998), or a prediction error method

(Ergon, 2022a,b). Note, however, that REML equations for a single generation will be

indeterminate, i.e., we can find $r_{1,t}$ and $r_{2,t}$, but not all the variances in the expressions for $r_{1,t}$

and $r_{2,t}$. For use in a general multivariate BLUP equation, we also need the $n \times n$ additive

genetic relationship matrix A_t (Ch. 26, Lynch and Walsh, 1998), and the Kronecker

covariance matrices $\widetilde{G}_t = G \otimes A_t$ and $\widetilde{R}_t = R_t \otimes I_n$.

For comparisons with predictions based on selection gradients (GRAD), as in Eq. (5), we

also need the phenotypic covariance matrix $P = \begin{bmatrix} P_{aa} & G_{ab} \\ G_{ab}^T & P_{bb} \end{bmatrix}$, where

$P_{aa} = G_{aa} + diag([\sigma_{v_1}^2 \quad \sigma_{v_2}^2])$ and $P_{bb} = G_{bb} + diag([\sigma_{\eta_{11}}^2 \quad \sigma_{\eta_{12}}^2 \quad \sigma_{\eta_{21}}^2 \quad \sigma_{\eta_{22}}^2])$, with

$diag([\cdot])$ denoting diagonal matrices. Also P is assumed to be constant.

As will be shown in detail below, the fixed effects $\bar{y}_{j,t}$ and random effects $a'_{j,t}$ and $b'_{jk,t}$ in

Eq. (7) may be estimated by use of BLUP. In line with established language in the breeding

literature, $\hat{\bar{y}}_{j,t}$ will here be called estimations, while $\widehat{a}'_{j,t}$ and $\widehat{b}'_{jk,t}$ will be called predictions.

See Robinson (1991) for a discussion of the historical background for this terminology. From

the mixed model (7) follows the multivariate BLUP equation in matrix form (Henderson,

1950; Ch. 26, Lynch and Walsh, 1998),

$$\begin{bmatrix} X^T \widetilde{R}_t^{-1} X & X^T \widetilde{R}_t^{-1} \widetilde{Z}_t \\ \widetilde{Z}_t^T \widetilde{R}_t^{-1} X & \widetilde{Z}_t^T \widetilde{R}_t^{-1} \widetilde{Z}_t + \widetilde{G}_t^{-1} \end{bmatrix} \begin{bmatrix} \hat{\bar{y}}_t \\ \widehat{x}_t \end{bmatrix} = \begin{bmatrix} X^T \widetilde{R}_t^{-1} y_t \\ \widetilde{Z}_t^T \widetilde{R}_t^{-1} y_t \end{bmatrix}, \tag{8}$$

279   where, for $p = 2$ and $q = 2$, $\hat{\bar{y}}_t = [\hat{\bar{y}}_{1,t} \quad \hat{\bar{y}}_{2,t}]^T$ and

280    $\widehat{x}_t = [\widehat{a}'^T_{1,t} \quad \widehat{a}'^T_{2,t} \quad \widehat{b}'^T_{11,t} \quad \widehat{b}'^T_{12,t} \quad \widehat{b}'^T_{21,t} \quad \widehat{b}'^T_{22,t}]^T$. Here, $E[\hat{\bar{y}}_{j,i,t}] = \bar{y}_{j,t}$ and

281    $E[\widehat{a}'_{1,i,t}] = E[\widehat{a}'_{2,i,t}] = E[\widehat{b}'_{11,i,t}] = E[\widehat{b}'_{12,i,t}] = E[\widehat{b}'_{21,i,t}] = E[\widehat{b}'_{22,i,t}] = 0$. Note that the

282   derivation of Eq. (8) does not necessarily require an assumption of normal data (Robinson

283   1991).

2.5 Updating of mean reaction norm parameter values

When Eq. (8) is applied on any given parent generation, the expected mean values of the

vector elements in $\hat{\mathbf{x}}_t$ will be zero, i.e., $E\left[\bar{\bar{a}}'_{1,t}\right] = E\left[\bar{\bar{a}}'_{2,t}\right] = \cdots = E\left[\bar{\bar{b}}'_{22,t}\right] = 0$. However,

owing to different fitness (number of descendants) among the individuals in the parent

generation, the corresponding expected mean values in the offspring generation, before new

reproduction, will be different from $E\left[\bar{\bar{a}}'_{1,t}\right]$, etc., and the within-generation differences in

$\bar{\bar{a}}'_{1,t}$, etc., from parents to offspring, may be used for updating of the mean reaction norm

parameters $\bar{a}_{1,t}, \bar{a}_{2,t}, \bar{b}_{11,t}, \bar{b}_{12,t}, \bar{b}_{21,t}$, and $\bar{b}_{22,t}$ in Eq. (6a,b). After this updating the offspring

are ready to become new parents, again with $E\left[\bar{\bar{a}}'_{1,t}\right] = E\left[\bar{\bar{a}}'_{2,t}\right] = \cdots = E\left[\bar{\bar{b}}'_{22,t}\right] = 0$.

Selection will thus result in within-generation incremental changes in the mean values of

the predicted random effects from parents to offspring before reproduction, generally given by

Eq. (1),

$$\bar{z}_t^{offspring} - \bar{z}_t^{parents} = cov\left(w_{i,t}, \hat{z}'_{i,t}\right) + \left[w_{i,t}\left(\hat{z}'^{descendants}_{i,t} - \hat{z}'_{i,t}\right)\right], \qquad (9)$$

where $\hat{z}'_{i,t}$ is any predicted individual value $\hat{a}'_{1,i,t}, \hat{a}'_{2,i,t}, \hat{b}'_{11,i,t}, \hat{b}'_{12,i,t}, \hat{b}'_{21,i,t}$, or $\hat{b}'_{22,i,t}$ in

the random effects vector $\hat{\mathbf{x}}_t$ in Eq. (8), while $\hat{z}'^{descendants}_{i,t}$ is the corresponding mean value

for the descendants of individual i in generation t. In Eq. (9), $\bar{z}_t^{parents}$ stands for the mean

value of any one of the predicted random effects from Eq. (8), i.e., $\bar{\bar{a}}'_{1,t}, \bar{\bar{a}}'_{2,t}, \bar{\bar{b}}'_{11,t}, \bar{\bar{b}}'_{12,t},$

$\bar{\bar{b}}'_{21,t}$, or $\bar{\bar{b}}'_{22,t}$, while $\bar{z}_t^{offspring}$ is the corresponding mean value for the offspring after

selection but before reproduction. These incremental changes should thus at each generation

be used for updating of the mean reaction norm parameter values before the offspring become

new parents. For this purpose, we need an additional assumption, which follows from

Assumption 3 above, because $\hat{z}'_{i,t}$ and $\hat{z}'^{descendants}_{i,t}$ are predictions of an additive genetic

component of a reaction norm parameter, and thus have no non-additive components:

Assumption 7 *There are no expected fitness weighted changes in predicted random effects*

from individual parents to their descendants after selection but before reproduction, i.e.,

$E\left[w_{i,t}\left(\widehat{z'}_{i,t}^{descendants} - \widehat{z'}_{i,t}\right)\right] = 0.$

When Assumption 7 is applied on Eq. (9), we obtain the Robertson-Price identity for the

within-generation changes in the mean values (Ch. 6, Walsh and Lynch, 2018), but note that

$\widehat{z'}_{i,t}$ are BLUP predictions. Here, the incremental changes $\bar{z}_t^{offspring} - \bar{z}_t^{parents}$ will be

entirely determined by the predicted additive genetic values $\widehat{z'}_{i,t}$ and individual fitness, and

when these changes are used for updating we thus obtain between-generation changes in the

mean values as given by Robertson's secondary theorem of natural selection, but also here

based on BLUP predictions. When Eq. (8) is applied on a given parent generation, and when

the changes in mean values of predicted random effects from the parent to the offspring

generation are used for updating, the incremental changes in those values under Assumption 7

thus follow from the following theorem:

Theorem 1 *In a population that is adequately described by Eq. (7), the incremental changes*

in mean reaction norm parameter values from generation to generation are found from

Robertson's secondary theorem of natural selection, applied on the BLUP predictions,

$$\Delta\bar{z}_t = cov\left(w_{i,t}, \widehat{z'}_{i,t}\right), \tag{10}$$

322 *where $\bar{z}_t$ is any mean parameter value $\bar{a}_{1,t}, \bar{a}_{2,t}, \bar{b}_{11,t}, \bar{b}_{12,t}, \bar{b}_{21,t},$ or $\bar{b}_{22,t}$, while $\widehat{z'}_{i,t}$ is the*

323 *corresponding predicted individual value $\widehat{a'}_{1,i,t}, \widehat{a'}_{2,i,t}, \widehat{b'}_{11,i,t}, \widehat{b'}_{12,i,t}, \widehat{b'}_{21,i,t},$ or $\widehat{b'}_{22,i,t}$, in the*

324 *random effects vector $\widehat{\mathbf{x}}_t$ in Eq. (8).*

325   From Theorem 1 follows the incremental changes in mean traits according to

$$\begin{bmatrix} \Delta\bar{a}_{1,t} \\ \Delta\bar{a}_{2,t} \\ \Delta\bar{b}_{11,t} \\ \Delta\bar{b}_{12,t} \\ \Delta\bar{b}_{21,t} \\ \Delta\bar{b}_{22,t} \end{bmatrix} = \begin{bmatrix} cov\left(w_{i,t}, \widehat{a}'_{1,i,t}\right) \\ cov\left(w_{i,t}, \widehat{a}'_{2,i,t}\right) \\ cov\left(w_{i,t}, \widehat{b}'_{11,i,t}\right) \\ cov\left(w_{i,t}, \widehat{b}'_{12,i,t}\right) \\ cov\left(w_{i,t}, \widehat{b}'_{21,i,t}\right) \\ cov\left(w_{i,t}, \widehat{b}'_{22,i,t}\right) \end{bmatrix}. \tag{11}$$

With correct initial mean reaction norm parameter values, this gives $\bar{y}_{1,t}$ and $\bar{y}_{2,t}$ according to

Eq. (6b), and this can be generalized to higher dimensions, with $p > 2$ and $q > 2$.

For finite population sizes there will be drift in the mean reaction norm parameter values,

owing to random errors in the covariance computations according to Eq. (11). However, as we

will see in the simulations, there are also other sources of drift.

From the BLUP theory above follows the following theorem:

Theorem 2 *For a reaction norm evolutionary system with $p = 2$ phenotypic traits and $q = 2$*

environmental cues, Eq. (8) and Theorem 1 will with $\boldsymbol{A}_t = \boldsymbol{I}_n$ result in incremental changes in

mean reaction norm parameter values according to

$$\begin{bmatrix} \Delta\bar{a}_{1,t} \\ \Delta\bar{a}_{2,t} \\ \Delta\bar{b}_{11,t} \\ \Delta\bar{b}_{12,t} \\ \Delta\bar{b}_{21,t} \\ \Delta\bar{b}_{22,t} \end{bmatrix} = \boldsymbol{G}(\boldsymbol{Z}_t^T \boldsymbol{R}_t^{-1} \boldsymbol{Z}_t \boldsymbol{G} + \boldsymbol{I}_6)^{-1} \boldsymbol{Z}_t^T \boldsymbol{R}_t^{-1} \begin{bmatrix} cov(w_{i,t}, y_{1,i,t}) \\ cov(w_{i,t}, y_{2,i,t}) \end{bmatrix}. \tag{12}$$

This can be generalized to higher dimensions, with $p > 2$ and $q > 2$.

See Appendix A for proof, and simulation results in Section 3.

For comparisons with the multivariate breeder's equation, the results from Eq. (12) can also

be found by a generalization of the GRAD incremental parameter changes according to Eq.

(5):

Theorem 3. *Eq. (12) can be reformulated as an extension of Eq. (5),*

$$
\begin{bmatrix}
\Delta\bar{a}_{1,t} \\
\Delta\bar{a}_{2,t} \\
\Delta\bar{b}_{11,t} \\
\Delta\bar{b}_{12,t} \\
\Delta\bar{b}_{21,t} \\
\Delta\bar{b}_{22,t}
\end{bmatrix}
=
\begin{bmatrix}
G_{aa} & G_{ab} \\
G_{ab}^T & G_{bb}
\end{bmatrix}
\begin{bmatrix}
I_2 \\
U_t
\end{bmatrix}
P_{yy,t}^{-1}
\begin{bmatrix}
cov(w_{i,t}, y_{1,i,t}) \\
cov(w_{i,t}, y_{2,i,t})
\end{bmatrix},
\tag{13}
$$

where $\begin{bmatrix} I_2 \\ U_t \end{bmatrix} = Z_t^T$*, while* $P_{yy,t} = P_{aa} + 2U_t^T G_{ab} + U_t^T P_{bb} U_t$.

See Appendix B for proof, and simulation results in Section 3, where Eqs. (12) and (13) give

identical results for population sizes $n \geq 2$. The simulations also show that the results from

Eq. (13) are close to the results from a corresponding version of the multivariate breeder's

equation (4), with declining differences for increasing population size.

2.6 Heritability influences on the BLUP results

The BLUP results found from Eq. (8) are affected by the heritabilities

$h_{aa,jj}^2 = G_{aa,jj}/\left(G_{aa,jj} + \sigma_{v_j}^2\right)$ for j from 1 to p, and $h_{bb,jk}^2 = G_{bb,jk}/\left(G_{bb,jk} + \sigma_{\eta_{jk}}^2\right)$ for j

from 1 to p and k from 1 to q. A closer look at this leads to the following theorem:

Theorem 4 *When all heritability values are close to one, i.e., when* $\sigma_{v_j}^2 \to 0$ *and* $\sigma_{\eta_{jk}}^2 \to 0$, *the*

incremental changes in mean reaction norm parameter values with $A_t \neq I_n$ *are*

asymptotically given by Eqs. (12) and (13), i.e., as with $A_t = I_n$.

Proof*:* We may reformulate Eq. (8) as

$$
\begin{bmatrix}
X^T \widetilde{R}_t^{-1} X & X^T \widetilde{R}_t^{-1} \widetilde{Z}_t \\
\widetilde{Z}_t^T \widetilde{R}_t^{-1} X & \widetilde{Z}_t^T \widetilde{R}_t^{-1} \widetilde{Z}_t + \left(G \otimes I_n + G \otimes (A_t - I_n)\right)^{-1}
\end{bmatrix}
\begin{bmatrix}
\widehat{\bar{y}}_t \\
\widehat{x}_t
\end{bmatrix}
=
\begin{bmatrix}
X^T \widetilde{R}_t^{-1} y_t \\
\widetilde{Z}_t^T \widetilde{R}_t^{-1} y_t
\end{bmatrix}.
\tag{14}
$$

356 When all the diagonal elements in $\widetilde{R}_t^{-1}$ have large values, the influence of the matrix

357 $G \otimes (A_t - I_n)$ vanishes, such as we approximately obtain Eq. (8), with $A_t = I_n$.

2.7 Special case without plasticity

When all plasticity slope parameter values are zero, Eq. (6a) gives $\bar{y}_{j,t} = \bar{a}_{j,t}$, and the individual phenotypic traits

$$y_{j,i,t} = \bar{y}_{j,t} + a'_{j,i,t} + v_{j,i,t}. \tag{15}$$

For p traits this leads to Eq. (8) with $\hat{x}_t = \begin{bmatrix} \hat{a}'^T_{1,t} & \hat{a}'^T_{2,t} & \cdots & \hat{a}'^T_{p,t} \end{bmatrix}^T$, $\widetilde{Z}_t = I_{pn}$,

$$X = \begin{bmatrix} 1_n & 0 & \cdots & 0 \\ 0 & 1_n & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 1_n \end{bmatrix}, \widetilde{G} = G_{aa} \otimes A_t, \text{ and } \widetilde{R}_t = \begin{bmatrix} r_1 & 0 & \cdots & 0 \\ 0 & r_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & r_p \end{bmatrix} \otimes I_N, \text{ where } r_j \text{ is the}$$

variance of $v_{j,i,t}$. For $A_t = I_n$ this leads to the following theorem:

Theorem 5 *With individual phenotypic traits according to Eq. (15), and with $A_t = I_n$, the dynamical BLUP model above and the multivariate breeder's equation (2) give identical results.*

A proof of Theorem 5 is given in Ergon (2022c), although then relying on a comparison with the multivariate breeder's equation. Here, Theorem 1 makes it into an independent proof. Note that in this case Eq. (13) degenerates into Eq. (2).

2.8 Errors in estimated random effects variances

The fact that updated mean reaction norm parameter values, and thus also updated mean phenotypic trait values, are found by use of Robertson's secondary theorem of natural selection applied on predicted random effects, as stated in Theorem 1, quite generally shows that the variances of these effects are underestimated. This is easily seen in the case without plasticity, with individual traits given by Eq. (15), in which case Eq. (2) can be formulated as

$$\begin{bmatrix} \Delta\bar{y}_{1,t} \\ \Delta\bar{y}_{2,t} \\ \vdots \\ \Delta\bar{y}_{p,t} \end{bmatrix} = cov\left(w_{i,t}, GP^{-1} \begin{bmatrix} a'_{1,i,t} + v_{1,i,t} \\ a'_{2,i,t} + v_{2,i,t} \\ \vdots \\ a'_{p,i,t} + v_{p,i,t} \end{bmatrix} \right), \tag{16a}$$

376   which should be compared with the result following from Theorem 1,

$$\begin{bmatrix} \Delta\bar{y}_{1,t} \\ \Delta\bar{y}_{2,t} \\ \vdots \\ \Delta\bar{y}_{p,t} \end{bmatrix} = cov\left(w_{i,t}, \begin{bmatrix} \widehat{a}'_{1,i,t} \\ \widehat{a}'_{2,i,t} \\ \vdots \\ \widehat{a}'_{p,i,t} \end{bmatrix} \right).$$

(16b)

377   The underestimation of the variances $G_{jj} = E\left[a'^2_{j,i,t}\right]$ becomes especially transparent with

378   diagonal G and P matrices, where we from Eqs. (16a,b), with use of $\sigma^2_{v_j} = E\left[v^2_{j,i,t}\right]$, find

379    $\widehat{a}'_{j,i,t} = G_{jj}\left(G_{jj} + \sigma^2_{v_j}\right)^{-1}\left(a'_{j,i,t} + v_{j,i,t}\right)$. In this case we thus find

(16c)

$$var(\widehat{a}'_{j,i,t}) = \hat{G}_{jj} = var\left(\frac{G_{jj}}{G_{jj} + \sigma^2_{v_j}}(a'_{j,i,t} + v_{j,i,t}) \right) = \frac{G^2_{jj}}{G_{jj} + \sigma^2_{v_j}},$$

380   i.e., $var(\widehat{a}'_{j,i,t}) = \hat{G}_{jj} < G_{jj}$ for $\sigma^2_{v_j} > 0$. See Ergon (2022c) for simulation results.

381   ## 2.9 Adjustments for overlapping generations

382   With surviving parents, only a fraction $f_t < 1$ of a given generation are offspring from the

383   previous generation. The incremental changes in mean reaction norm parameter values from

384   one generation to the next are then reduced accordingly, and by use of Eq. (11) we thus obtain

$$\Delta\bar{z}_t = f_t \times cov\left(w_{i,t}, \widehat{z}'_{1,i,t}\right),$$

(17)

where $\bar{z}_t$ is any one of the mean reaction norm parameters, while $\widehat{z}'_{1,i,t}$ is the corresponding

predicted individual random effect. As verified in the simulations in Section 3, this will slow

down responses on environmental changes, with reduced mean fitness as consequence.

3. Simulations

3.1 The aim of the simulations

The aim of the simulations is to verify the theoretical BLUP results by means of a toy

example, and the purpose is fivefold. First, it is verified that mean reaction norm parameter

values can be updated from generation to generation by means of Robertson's secondary

theorem of natural selection, as applied on BLUP predictions (Theorem 1). Second, it is

shown that it is possible to disentangle the microevolutionary and plasticity components of for

example climate change acclimations as shown in Eq. (11) in general, and in Eq. (12) for the

special case with $A_t = I_n$. Third, it is verified that the dynamical BLUP and GRAD results

for the incremental changes in mean reaction norm parameter values are identical for

population sizes $n \geq 2$, provided that $A_t = I_n$ (Theorems 2 and 3). Fourth, it is shown that

the GRAD results are erroneous for populations with genetic relatedness between the

individuals, i.e., for $A_t \neq I_n$. Fifth, it is verified that heritability values close to one lead to

prediction results as with $A_t = I_n$, i.e., independent of the actual relationship values in A_t

(Theorem 4).

3.2 Description of toy example

In a toy example in Ergon (2022a), the environmental input was a noisy positive trend in

spring temperature, starting in 1970, and resulting in a noisy negative trend in mean breeding

(clutch-initiation) date for a certain bird species, approximately as in Fig. 2 in Bowers et al.

(2016).

Here, the example is extended to include a second environmental variable with a noisy

positive trend in mean value, and with variations from year to year that are somewhat

positively correlated with the variations in spring temperature. This input may for example be

a measure of spring rainfall. The example also includes a second adaptive phenotype, that

might be the breeding habitat, as discussed in Chalfoun and Schmidt (2012). We thus have a

microevolutionary system with two environmental cues and two phenotypic traits, similar to

the theoretical example case.

The individual (mid-parent) fitness values are integers from 0 to 4, with number of

descendants as unit, and generations are assumed to be non-overlapping. The population size

is assumed to be constant, which implies that not all descendants survive until reproduction. A

constant population size is not essential for the principal results, but it simplifies the

simulations.

In the simulations, the two environmental reference values are assumed to be known from

historical data, i.e., it is assumed that the population was fully adapted to the stationary

stochastic environment before the onset of anthropogenic and global climate change around

1970.

3.3 Environmental inputs and reaction norm model

Assume a population that is fully adapted to a stationary stochastic environment with mean

spring temperature $u_{1,ref} = 10\ °C$ (the reference value for temperature), and mean spring

rainfall $u_{2,ref} = 2$ mm/day (the reference value for rainfall). Also assume phenotypic scales

such that the phenotypic values that maximize fitness in the reference environment are given

by $\theta_{1,ref} = \theta_{2,ref} = 0$. Further assume environmental cues $u_{1,t} = \mu_{U_1,t} - 10 + u_{1,s,t}$, and

$u_{2,t} = \mu_{U_2,t} - 2 + u_{2,s,t}$, where the mean values $\mu_{U_1,t}$ and $\mu_{U_2,t}$ are ramp functions as shown

in Fig. 1, while $u_{1,s,t}$ and $u_{2,s,t}$ are zero mean and white random variables, i.e., without

autocorrelation. In a corresponding way assume that $\theta_{1,t} = \mu_{\Theta_1,t} + \theta_{1,s,t}$ and

$\theta_{2,t} = \mu_{\Theta_2,t} + \theta_{2,s,t}$, where $\mu_{\Theta_1,t}$ and $\mu_{\Theta_2,t}$ are ramp functions as shown in Fig. 1, while $\theta_{1,s,t}$

and $\theta_{2,s,t}$ are zero mean and white random variables.

Assume that $u_{1,s,t}, u_{2,s,t}, \theta_{1,s,t}$ and $\theta_{2,s,t}$ have a joint normal distribution with variances

$\sigma_{U_{1,s}}^2, \sigma_{U_{2,s}}^2, \sigma_{\Theta_{1,s}}^2$ and $\sigma_{\Theta_{2,s}}^2$, and covariances $\sigma_{U_{1,s}U_{2,s}}, \sigma_{U_{1,s}\Theta_{1,s}} = -2\rho_\tau \sigma_{U_{1,s}}^2$, and

$\sigma_{U_{2,s}\Theta_{2,s}} = -2\rho_\tau \sigma_{U_{2,s}}^2$, with numerical values as given in Fig. 1. Here, ρ_τ is the autocorrelation

of background environmental fluctuations, as described in more detail in Lande (2009). Data

were generated for 60 generations, with typical input data as shown in Fig. 1 (as mean values

in breeding season). See Supplementary Material for MATLAB code.

Figure 1. Typical input data for simulation example, with mean values shown by

dashed lines, and with ramp functions starting at generation $t = 10$ (1970). Numerical

values are $\mu_{\Theta_1,t} = -2\left(\mu_{U_1,t} - 10\right)$, $\mu_{\Theta_2,t} = -2\left(\mu_{U_2,t} - 2\right)$, $\sigma^2_{U_{1,s}} = 0.5$,

$\sigma^2_{U_{2,s}} = 0.1875$, $\sigma^2_{\Theta_{1,s}} = 2$, $\sigma^2_{\Theta_{2,s}} = 1$, $\sigma_{U_{1,s}U_{2,s}} = 0.25$, $\sigma_{U_{1,s}\Theta_{1,s}} = -0.25$, and

$\sigma_{U_{2,s}\Theta_{2,s}} = -0.09375$.

Also assume an individual reaction norm model with two phenotypic traits, according to

$$y_{1,i,t} = \bar{a}_{1,t} + a'_{1,i,t} + v_{1,i,t} + \left(\bar{b}_{11,t} + b'_{11,i,t} + \eta_{11,i,t}\right)u_{1,t}, \tag{18a}$$

$$y_{2,i,t} = \bar{a}_{2,t} + a'_{2,i,t} + v_{2,i,t} + \left(\bar{b}_{22,t} + b'_{22,i,t} + \eta_{22,i,t}\right)u_{2,t}, \tag{18b}$$

448 with parameters $\boldsymbol{G} = \begin{bmatrix} \boldsymbol{G}_{aa} & \boldsymbol{G}_{ab} \\ \boldsymbol{G}_{ab}^T & \boldsymbol{G}_{bb} \end{bmatrix}$ and $\boldsymbol{P} = \begin{bmatrix} \boldsymbol{P}_{aa} & \boldsymbol{G}_{ab} \\ \boldsymbol{G}_{ab}^T & \boldsymbol{P}_{bb} \end{bmatrix}$, and parameter values

449 $\boldsymbol{G}_{aa} = \begin{bmatrix} 0.2 & 0.1 \\ 0.1 & 0.2 \end{bmatrix}$, $\boldsymbol{G}_{ab} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$, $\boldsymbol{G}_{bb} = \begin{bmatrix} 0.05 & 0.025 \\ 0.025 & 0.05 \end{bmatrix}$, $\boldsymbol{P}_{aa} = \begin{bmatrix} 0.4 & 0.1 \\ 0.1 & 0.4 \end{bmatrix}$ and

450    $P_{bb} = \begin{bmatrix} 0.1 & 0.025 \\ 0.025 & 0.1 \end{bmatrix}$. Note that the two traits in Eqs. (18a,b) are correlated, with the

451   covariance $\sigma_{y_1,y_2} = G_{aa,12} + G_{bb,12}\sigma_{U_{1,s}U_{2,s}}$.

3.4 Fitness function and initial mean reaction norm values

453   The individual fitness function is assumed to be rounded values of

$$W_{i,t} = 4 \cdot exp\left(-\left((y_{1,i,t} - \theta_{1,t})^2 + (y_{2,i,t} - \theta_{2,t})^2\right)/2\omega^2\right), \tag{19}$$

where $\theta_{1,t}$ and $\theta_{2,t}$ are the phenotypic values that maximize fitness, while $\omega^2 = 10$. The

discrete values of $W_{i,t}$ (number of descendants) are thus integers from 0 to 4.

In the simulations it is essential that the mean reaction norm parameters are given correct

initial values at generation $t = 1$. We will assume that the phenotypic values are scaled such

that the initial mean trait values are $\bar{y}_{1,1} = \bar{y}_{2,1} = 0$, and since $u_{1,1} = u_{2,1} = 0$ this gives the

initial mean intercept values $\bar{a}_{1,1} = \bar{a}_{2,1} = 0$. We also assume that the initial mean slope

values are the optimal values in a stationary stochastic environment. These values are the ones

that maximize the expected individual fitness according to Eq. (19), in a stationary stochastic

environment, and thus minimize the criterion functions $J_1 = E\left[(y_{1,i,t} - \theta_{1,t})^2\right]$ and

$J_2 = E\left[(y_{2,i,t} - \theta_{2,t})^2\right]$. With $E[y_{1,i,t}] = 0$, and substituting $E[y_{1,i,t}^2] = \bar{b}_{11,t}^2\sigma_{U_{1,s}}^2$ and

$E[y_{1,i,t}\theta_{1,t}] = \bar{b}_{11,t}\sigma_{U_{1,s}\Theta_{1,s}}$, we find $J_1 = \bar{b}_{11,t}^2\sigma_{U_{1,s}}^2 - 2\bar{b}_{11,t}\sigma_{U_{1,s}\Theta_{1,s}} + \sigma_{\Theta_{1,s}}^2$, and setting

$\frac{dJ_1}{d\bar{b}_{11,t}} = 2\bar{b}_{11,t}\sigma_{U_{1,s}}^2 - 2\sigma_{U_{1,s}\Theta_{1,s}} = 0$, we thus find the optimal mean slope value

$\bar{b}_{11,opt} = \sigma_{U_{1,s}\Theta_{1,s}}/\sigma_{U_{1,s}}^2 = -0.5$. In the same way we find that $\bar{b}_{22,opt} = -0.5$ will minimize

J_2.

3.5 Simulation results

Simulation results with population size $n = 100$, and additive genetic relationship matrix

$A_t = I_n$, are shown in Fig. 2. New additive genetic (random) effects, and non-additive effects

(residuals), were at each generation drawn from normal distributions in accordance with the

given G and P matrices. The BLUP results given by Eq. (12) (green lines), and the GRAD

results given by Eq. (13) (dashed blue lines), are identical for population sizes $n \geq 2$. Results

given by the multivariate breeder's equation (4) (dotted magenta lines), are somewhat

different from the BLUP results. These differences are clearly smaller for a population size of

$n = 1,000$. Note that $\bar{y}_{1,t}$ and $\bar{y}_{2,t}$ lag behind $\theta_{1,t}$ and $\theta_{2,t}$, as shown in Fig. 1, which is typical

for ramp responses from dynamical systems with time constants.

Figure 2. Simulation results with population size $n = 100$, and an additive genetic

relationship matrix $A_t = I_n$. Mean trait values $\bar{y}_{1,t}$ and $\bar{y}_{2,t}$ are shown by solid blue

lines. Mean reaction norm parameter values are shown by solid green lines (BLUP),

dashed blue lines (GRAD), and dotted magenta lines (the multivariate breeder's

equation).

As a test, the simulations were repeated with a constant additive genetic relationship matrix

for a population with a high degree of relatedness among individuals, as shown for $n = 6$ in

Eq. (20),

$$A = \begin{bmatrix} 1 & 1/2 & 1/4 & 1/4 & 1/4 & 1/4 \\ 1/2 & 1 & 1/2 & 1/4 & 1/4 & 1/4 \\ 1/4 & 1/2 & 1 & 1/2 & 1/4 & 1/4 \\ 1/4 & 1/4 & 1/2 & 1 & 1/2 & 1/4 \\ 1/4 & 1/4 & 1/4 & 1/2 & 1 & 1/2 \\ 1/4 & 1/4 & 1/4 & 1/4 & 1/2 & 1 \end{bmatrix}. \tag{20}$$

As shown in Fig. 3, this gave different results for the BLUP and GRAD methods. In this case

new additive genetic (random) effects at each new generation were found as $\boldsymbol{z'}_t = \sqrt[M]{\boldsymbol{A}}\boldsymbol{z'}_{0,t}$,

where $\boldsymbol{z'}_t$ stands for $\boldsymbol{a'}_{1,t}$, $\boldsymbol{a'}_{2,t}$, $\boldsymbol{b'}_{11,t}$, $\boldsymbol{b'}_{12,t}$, $\boldsymbol{b'}_{21,t}$, or $\boldsymbol{b'}_{22,t}$, and where the different data

vectors $\boldsymbol{z'}_{0,t}$ were drawn from normal distributions in accordance with the given $\boldsymbol{G}$ matrix.

Here, $\sqrt[M]{\boldsymbol{A}}$ is the matrix square root, i.e., $\sqrt[M]{\boldsymbol{A}}\sqrt[M]{\boldsymbol{A}} = \boldsymbol{A}$. This reduced the variances of the

random effects to approximately 75% of the nominal values. The random effects were also

mean centred.

Figure 3. Simulation results with population size $n = 100$, and an additive genetic

relationship matrix $\boldsymbol{A}$ according to Eq. (20). Mean trait values are shown by solid blue

lines. Mean reaction norm parameter values are shown by solid green lines (BLUP)

and dashed blue lines (GRAD).

Responses with $\boldsymbol{A}$ according to Eq. (20), and heritability values close to one, were tested

with use of $\sigma_{v_1}^2 = 10^{-5}$, $\sigma_{v_2}^2 = 10^{-6}$, and $\sigma_{\eta_{11}}^2 = \sigma_{\eta_{12}}^2 = \sigma_{\eta_{21}}^2 = \sigma_{\eta_{22}}^2 = 0$. For population size

$n \geq 2$, the mean reaction norm parameter results based on Eq. (11), given with four decimals,

were in this case equal to the GRAD results given by Eq. (13). The response plots were

similar to those in Fig. 2, except that the plots of $\bar{y}_{1,t}$ and $\bar{y}_{2,t}$ were somewhat less noisy.

As a final test, the simulations with results as in Fig. 3 were repeated for a case with

surviving parents, where only a fraction $f_t = 0.5$ of any given generation are offspring from

the previous generation, i.e., with use of Eq. (17). As seen in Fig. 4, panels (a) and (c), and as

must be expected, the mean reaction norm parameter responses are slowed down. As can also

be seen in panels (a) and (c), the fraction $f_t < 1$ causes the mean trait values $\bar{y}_{1,t}$ and $\bar{y}_{2,t}$ to

lag further behind the phenotypic values $\theta_{1,t}$ and $\theta_{2,t}$ that maximize fitness. The result of this

is lower mean fitness values, as can be seen in the (identical) plots in panels (b) and (d).

Figure 4. Simulation results with population size $n = 100$, and additive genetic relationship matrix $\boldsymbol{A}_t \neq \boldsymbol{I}_n$ according to Eq. (20). In panels (a) and (c), mean trait values $\bar{y}_{1,t}$ and $\bar{y}_{2,t}$ and mean reaction norm parameter values $\bar{a}_{1,t}$ and $\bar{a}_{2,t}$ for $f_t = 1$ are shown by solid blue and solid green lines, respectively. Dashed blue and magenta lines show the corresponding responses with a fraction $f_t = 0.5$ of new offspring in the population at all generations. Phenotypic values $\theta_{1,t}$ and $\theta_{2,t}$ that maximize fitness are added as weak dotted blue lines. Panels (b) and (d) show identical plots of mean fitness for $f_t = 1$ (solid green lines) and $f_t = 0.5$ (dashed magenta lines).

The changes in mean values over 60 generations in Figs. 2 and 3, will to some degree be caused by drift. As a test, the total change in $\bar{a}_{1,t}$ over 60 generations in a stationary stochastic environment as before $t = 10$ in Fig. 1, was computed. For population sizes from $n = 10$ to 400, and based on 100 repeated simulations, this change had a mean value of approximately zero, and a standard error of approximately 0.1, i.e., around 10% of the

corresponding changes in panel (a) in Figs. 2 and 3. With $n = 2$, this standard error due to drift increased to 15%. There were no noticeable differences between cases with $A_t \neq I_n$ and $A_t = I_n$. In order to find an explanation of the drift, the environmental cue $u_{1,t}$ over 60 generations in a stationary environment was approximated by a least squares straight line, denoted $\hat{u}_{1,t}$. Based on 100 repeated simulations with $n = 100$, the resulting change in $\hat{u}_{1,t}$ over 60 generations had a mean value of approximately zero, and a standard error of approximately 0.30, i.e., around 12% of the corresponding changes of $u_{1,t}$ in Fig. 1, panel (a). This result indicates that the major part of the drift is caused by the stochastic nature of the environmental cues and the phenotypic values that maximize fitness.

4. Summary and Discussion

As shown in Section 2, a general reaction norm model with p phenotypic traits and q environmental cues, i.e., with $p(1 + q)$ mean reaction norm parameters, can be formulated as a linear mixed model with fixed and random effects. In this model, the fixed effects will be the mean trait values, while the random effects will be the additive genetic components of individual deviations from mean reaction norm parameter values. From this follows that for any parent generation, estimates of the mean trait values, and predictions of the additive genetic components of individual deviations from mean reaction norm parameter values, can be found from BLUP equations. Because the incidence matrix in the linear mixed model is a function of the environmental cues, and thus of time, we have introduced the concept of dynamical BLUP. Note that the elements in the random effects vector in Eq. (7) could be ordered differently, for example as $x_t = \begin{bmatrix} a'^T_{1,t} & b'^T_{11,t} & b'^T_{12,t} & a'^T_{2,t} & b'^T_{21,t} & b'^T_{22,t} \end{bmatrix}^T$, but then also the G, P, U_t and Z_t matrices must be reorganized accordingly.

The development of the dynamical BLUP model in Eq. (8) relies on Assumptions 1, 2, and 3, as given in Section 2, while Assumptions 4, 5, and 6 are needed only for the comparison

with results based on the multivariate breeder's equation. Updating of the mean reaction norm

parameters according to Theorem 1, requires an additional Assumption 7. Most importantly,

Assumption 4 implies that applications of the multivariate breeder's equation on non-normal

data in general will produce incorrect microevolutionary results. This problem was

acknowledged by Lande and Arnold (1983), who in their Appendix proposed a possible

correction for errors caused by skewness. The fundamental reason for such errors is that the

multivariate breeder's equation involves only mean and variance values of the additive

genetic and non-additive effects, while influences from third and higher order statistical

moments are ignored. See for example discussions in Bonamour et al. (2017) and Pick et al.

(2022).

Although the development of the dynamical BLUP model in Eq. (8) does not rely on

normal data (Robinson, 1991), it still follows from Theorems 2 and 3 that the BLUP results

will be incorrect in cases with non-normal data and an additive relationship matrix $A_t = I_n$. A

natural conclusion is therefore that non-normal data will give erroneous BLUP results also

when $A_t \neq I_n$, at least for the dynamical BLUP model used here.

In the derivation of the linear mixed model in Eq. (7), we assumed linear reaction norms as

functions of environmental cues $u_{1,t}$, $u_{2,t}$, etc., but there is nothing in the theory that prevents

us from use of non-linear reaction norms that are also functions of $u_{1,t}u_{2,t}$, $u_{1,t}^2$, $u_{2,t}^2$, etc., as

discussed in Gavrilets and Scheiner (1993) and Ergon (2018).

As shown by Theorem 1, updating of the mean reaction norm parameter values from

generation to generation can be done by applying Robertson's secondary theorem of natural

selection on the vector elements $\widehat{a}'^T_{1,t}$, etc., in the predicted random effects. This shows that

the well-known underestimation of the variances of the random effects, is just what is needed,

in order to find the correct incremental changes in mean reaction norm parameter values. The

resulting dynamics will depend on the additive genetic and residual covariance matrices $\boldsymbol{G}$ and

$\boldsymbol{R}_t$, as well as of the additive genetic relationship matrix $\boldsymbol{A}_t$ for the parent generation.

It is worth noticing that the additive genetic relationship matrix for the offspring generation

does not affect the updating of mean reaction norm parameter values. This is shown in the

theoretical derivations, but it is also a natural consequence of the fact that the dynamical

BLUP model, just as the multivariate breeder's equation, is an evolutionary state-space model

(Ergon, 2018).

For cases with $\boldsymbol{A}_t \neq \boldsymbol{I}_n$, it should be noted that $\boldsymbol{A}_t$ is included in the BLUP matrix equation

(8) via $\widetilde{\boldsymbol{G}}_t$, just as in the standard equation used in breeding (Ch. 26, Lynch and Walsh, 1998).

It is also reassuring to know that the BLUP equation asymptotically, for $\boldsymbol{A}_t \rightarrow \boldsymbol{I}_n$, gives

results that are identical to the GRAD results based on the multivariate breeder's equation

(Theorems 2 and 3). It is interesting to note that there are no estimated parameter or variable

values involved in Eqs. (12) and (13), i.e., the mean trait prediction results hold as long as the

assumptions in Subsection 2.3 hold. This implies that BLUP predictions corrected for the

underestimated variance of the random effects would give un-correct predictions of the mean

reaction norm parameters. It is also interesting to note that the GRAD and BLUP results for

$\boldsymbol{A}_t \neq \boldsymbol{I}_n$ are asymptotically identical when all heritability values $h^2 \rightarrow 1$ (Theorem 4). Note

that results from use of the multivariate breeder's equation are equal to the GRAD results only

asymptotically, i.e., for population size $n \rightarrow \infty$. For cases without plasticity, and with

$\boldsymbol{A}_t = \boldsymbol{I}_n$, the results from all three methods are identical, independent of population size

(Theorem 5).

Theorems 1, 2, and 3 were verified in simulations with use of $\boldsymbol{A}_t = \boldsymbol{I}_n$, and population sizes

down to $n = 2$, while Theorem 4 was verified by use of $\boldsymbol{A}_t \neq \boldsymbol{I}_n$, and all heritability values

close to one. Theorem 5 was verified by simulations in Ergon (2022c). Simulation results with

$A_t \neq I_n$ were found by use of a constant and possibly unrealistic additive genetic relationship

matrix, but these results still serve the purpose of showing that the BLUP and GRAD results

with $A_t \neq I_n$ are different for heritability values $h^2 < 1$.

The changes in mean trait and mean reaction norm parameter values over 60 generations

that can be seen in Figs. 2 and 3, are partly caused by drift. As indicated at the end of Section

3, the major sources of drift are the stochastic nature of the input environmental variables $u_{1,t}$

and $u_{2,t}$, as well as of the phenotypic values $\theta_{1,t}$ and $\theta_{2,t}$ that maximize fitness. In the

simulations, the population size plays a vital role only when $n < 10$.

In the simulations, the G and P parameters are assumed to be known, although they are

normally not, and this is the case also for the reference environment and the initial mean

reaction norm parameters. However, as shown in Ergon (2022a,b), these parameters can be

found by a prediction error method (PEM) using all available input-output experimental or

field data, including individual fitness data. Ergon (2022a,b) combined PEM with the GRAD

model, i.e., with extensions of Eq. (5), but as will be reported separately, PEM works just as

well when combined with the BLUP model in Eq. (8). BLUP/PEM will in fact be much better

than GRAD/PEM when it comes to identification of the reference environment. As will be

reported separately, the essential feature of PEM in the present setting is to predict the

reaction norms as well as possible, while parameter values may be very much influenced by

measurement noise and modeling errors.

Parameter estimation by means of restricted maximum likelihood (REML) applied on data

from a single generation is not possible for plastic organisms. A simple reason for this is that

the variance components $r_{j,t}$ in the diagonal residual covariance matrix R_t are functions of

several unknown variances of the non-additive effects in the reaction norm model, i.e., we can

find $r_{1,t}, r_{2,t}$, etc., but not all the residual covariances involved. The reason behind this

problem is that we have confounding variables in the reaction norm equations. In for example

Eq. (3), all the terms in $a'_{i,t} + b'_{i,t}u_t + v_{i,t} + \eta_{i,t}u_t$ are confounded, which implies that use of

REML at a single generation will not give details of the $\boldsymbol{G}$ and $\boldsymbol{R}_t$ matrices, and a solution is

to use a prediction error method (PEM) where data from all generations are used. Note that

this confounding is not a problem in the modeling process, where $\bar{a}_t$ and $\bar{b}_t$ in

$a'_{i,t} = a_{i,t} - \bar{a}_t$ and $b'_{i,t} = b_{i,t} - \bar{b}_t$, respectively, are found from Eq. (11), and where $a'_{i,t}$,

$b'_{i,t}$, $v_{i,t}$ and $\eta_{i,t}$ are samples with given variances.

With surviving parents, only a fraction f_t of the population will be offspring from the

previous generation, and as shown in Eq. (17), the incremental changes in mean reaction norm

parameter values will then be reduced accordingly. As verified in simulations, and as must be

expected, the responses on environmental changes will then be slowed down, with reduced

mean fitness as consequence. Note that the last term on the right-hand side of Eq. (1), the

Price equation, in general is affected by surviving parents, but that this term according to

Assumption 7 does not affect the dynamical BLUP updating from generation to generation.

Acknowledgements

636 The GRAD model in Eq. (5) was inspired by a comment by Michael B. Morrissay in a review

637 of a quite different manuscript. I thank Tormod Ådnøy for input regarding matrix details in

638 BLUP modeling, and I thank University of South-Eastern Norway for support and funding.

Author contribution

640 Rolf Ergon is the sole author of this article.

Competing interests

642 There are no competing interests.

Data availability statement

MATLAB code for simulations is given in Supplementary Material archived on bioRxiv

doi: https://doi.org/10.1101/2023.04.09.536146.

ORCID

Rolf Ergon https://orcid.org/0000-0003-2393-2997

References

Arnold PA, Kruuk LEB, Nicotra AB (2019) How to analyse plant phenotypic plasticity in response to a changing climate. New Phytologist 222:1235-1241. doi: 10.1111/nph.15656

Autzen B, Okasha S (2022) On geometric mean fitness: a reply to Takacs and Bourrat. Biology & Philisophy 37:37. https://doi.org/10.1007/s10539-022-09874-x

Bonamour S, Teplitsky C, Charmantier A, Crochet P-A, Chevin L-M (2017) Selection on skewed characters and the paradox of stasis. Evolution 71(11):2703–2713. doi: 10.1111/evo.13368

Bowers EK, Grindstaff JL, Soukup SS, Drilling NE, Eckerle KP, Sakaluk SK, Thompson CF (2016) Spring temperatures influence selection on breeding data and the potential for phenological mismatch in a migratory bird. Ecology 97(10): 2880-2891. doi: 10.1002/ecy.1516

Ergon R (2018) The environmental zero-point problem in evolutionary reaction norm modeling. Ecology and Evolution 8(8):4031-4041. doi: 10.1002/ece3.3929

Ergon R (2019) Quantitative genetics state-space modeling of phenotypic plasticity and evolution. Modeling, Identification and Control 40/1:51–69. doi:10.4173/mic.2019.1.5

Ergon R (2022a) The important choice of reference environment in microevolutionary

climate response predictions. Ecology and Evolution 12(4):e8836.

https://doi.org/10.1002/ece3.8836

Ergon R (2022b) Microevolutionary system identification and climate response predictions.

Modeling, Identification and Control 43(3):91–99. doi: 10.4173/mic.2022.3.1

Ergon R (2022c) A BLUP derivation of the multivariate breeder's equation, with an

elucidation of errors in BLUP variance estimates, and a prediction method for inbred

populations. Modeling, Identification and Control 43/(4):131–140.

doi:10.4173/mic.2022.4.2

Gavrilets, S., & Scheiner, S. M. (1993). The genetics of phenotypic plasticity V. Evolution of

reaction norm shape. J. Evol. Biol. *6*, 31–48.

https://doi.org/10.1046/j.1420-9101.1993.6010031.x

Gomulkiewicz R, Kirkpatrick M (1992) Quantitative Genetics and the Evolution of Reaction

Norms. Evolution, 46/22:390-411.

Hadfield JD, Wilson AJ, Garant D, Sheldon BC, Kruuk LEB (2010) The Misuse of BLUP in

Ecology and Evolution. Am. Nat. 175:116–125. doi: 10.1086/648604

Henderson CR (1950) Estimation of genetic parameters (abstract).

Ann. Math. Statist. 21:309-3010

Kingsolver JG, Gomulkiewicz R, Carter PA (2001) Variation, selection and evolution of

function-valued traits, Genetica 112–113: 87–104.

Kirkpatrick M, Heckman N (1989) A quantitative genetic model for growth, shape, reaction

norms, and other infinite-dimensional characters. J. Math. Biol. 27:429-450.

Kruuk LEB (2004) Estimating genetic parameters in natural populations using the 'animal

model'. Phil. Trans. R. Soc. Lond. B 359:873–890. doi: 10.1098/rstb.2003.1437

Lande R (1979) Quantitative genetic analysis of multivariate evolution, applied to

brain:body size allometry. Evolution 33:402-416

Lande R (2009) Adaptation to an extraordinary environment by evolution of phenotypic

plasticity and genetic assimilation. J. Evol. Biol. 22:1435-1446.

doi: 10.1111/j.1420-9101.2009.01754

Lande R, Arnold SJ (1983) Measurement of selection on correlated characters.

Evolution 37(6):210-1226

Lynch M, Walsh B (1998) Genetics and Analysis of Quantitative Traits. Sinauer Associates,

Sunderland, MA

Merilä J, Hendry AP (2014). Climate change, adaptation, and phenotypic

plastcity: the problem and the evidence. Evol. Appl. 7, 1-14.

doi: 10.1111/eva. 12137.

Nussey DH, Wilson AJ, Brommer JE (2007) The evolutionary ecology of individual

phenotypic plasticity in wild populations. J. Evol. Biol. 20(3):831-844.

doi: 10.1111/j.1420-9101.2007.01300.x

Oliviera HR, Brito LF, Lourenco DAL, Silva FF, Jamrozik J, Schaeffer LR, Schenkel1 FS

(2019) J. Dairy Sci. 102:7664–7683. https://doi.org/10.3168/jds.2019-16265

Pick JL, Lemon HE, Thomson CE, Hadfield JD (2022) Decomposing phenotypic skew and its

effects on the predicted response to strong selection. Nature Ecology & Evolution 6:774–785.

https://www.nature.com/articles/s41559-022-01694-2

Price GR (1970) Selection and covariance. Nature 227:529-31

Price GR (1972) Extension of covariance selection mathematics. Annals of Human

Genetics 35 (4):485–490. doi: 10.1111/j.1469-1809.1957.tb01874.x.

Rice SH (2004) Evolutionary Theory: Mathematical and Conceptual Foundations. Sinauer

Ass., Mass.

Robertson A (1966) A mathematical model of the culling process in dairy cattle. Animal

Science 8/1:95-108. doi: https://doi.org/10.1017/S0003356100037752

Robinson GK (1991) That BLUP Is a Good Thing: The Estimation of Random Effects.

Statistical Science 6/1:15-51

Sæther B-E, Engan S (2015) The concept of fitness in fluctuating environments. Trends in

Ecology & Evolution, 30/5. http://dx.doi.org/10.1016/j.tree.2015.03.007

Shaeffer LR (2004) Application of random regression models in animal breeding. Livestock

Production Science 86:35–45. doi:10.1016/S0301-6226(03)00151-9

Appendix A Proof of Theorem 2

Lemma 1 *Assuming an additive genetic relationship matrix $A_t = I_n$, the mean values of the*

estimated fixed effects in Eq. (8), for j from 1 to p, are $\hat{\bar{y}}_{j,t} = \bar{y}_{j,t}$, while the mean values of

the predicted random effects, for j from 1 to p and k from 1 to q, are $\overline{\hat{a}}'_{j,t} = 0$ and

$\overline{\hat{b}}'_{jk,t} = 0.$

Proof: Eq. (8) will after elimination of $X^T \tilde{R}_t^{-1}$ give the following equation for a specific trait

$j,$

$$\hat{a}'_{j,t} + u_{1,t}\hat{b}'_{j1,t} + u_{2,t}\hat{b}'_{j2,t} + \cdots + u_{r,t}\hat{b}'_{jq,t} = y_{j,t} - \mathbf{1}_n\hat{\bar{y}}_{j,t}, \quad\quad \text{(A1a)}$$

728 Taking mean values on both sides of Eq. (A1a) gives

$$\overline{\hat{a}}'_{j,t} + u_{1,t}\overline{\hat{b}}'_{j1,t} + u_{2,t}\overline{\hat{b}}'_{j2,t} + \cdots + u_{r,t}\overline{\hat{b}}'_{jq,t} = \bar{y}_{j,t} - \hat{\bar{y}}_{j,t}, \quad\quad \text{(A1b)}$$

729 Eq. (8) also gives

$$(Z_t^T R_t^{-1} Z_t + G^{-1}) \otimes I_n \hat{x}_t = (Z_t^T R_t^{-1}) \otimes I_n \begin{bmatrix} y_{1,t} - 1_n \hat{\bar{y}}_{1,t} \\ \vdots \\ y_{j,t} - 1_n \hat{\bar{y}}_{j,t} \\ \vdots \\ y_{p,t} - 1_n \hat{\bar{y}}_{p,t} \end{bmatrix}, \tag{A2a}$$

730   from which with use of the mixed model property $(A \otimes B)(C \otimes D) = (AC) \otimes (BD)$

731   follows that

$$\tilde{Q} \hat{x}_t = \begin{bmatrix} y_{1,t} - 1_n \hat{\bar{y}}_{1,t} \\ \vdots \\ y_{j,t} - 1_n \hat{\bar{y}}_{j,t} \\ \vdots \\ y_{p,t} - 1_n \hat{\bar{y}}_{p,t} \end{bmatrix}, \tag{A2b}$$

732   where $\tilde{Q} = Q \otimes I_n = ((Z_t^T R_t^{-1})^T Z_t^T R_t^{-1})^{-1} (Z_t^T R_t^{-1})^T (Z_t^T R_t^{-1} Z_t + G^{-1}) \otimes I_n$ is an

733    $pn \times pn(1 + q)$ matrix, with block row vectors $\tilde{Q}_j = Q_j \otimes I_n$, for j from 1 to p. Since all

734   scalar elements in Q_j are multiplied with I_n, Eq. (A2b) gives

$$Q_j \hat{x}_t = y_{j,t} - 1_n \hat{\bar{y}}_{j,t}. \tag{A2c}$$

735   By taking mean values of both sides of Eq. (A2c) we thus find

$$Q_j \bar{\hat{x}}_t = \bar{y}_{j,t} - \hat{\bar{y}}_{j,t}. \tag{A2d}$$

Eqs. (A1b) and (A2d) thus give two expressions for $\bar{y}_{j,t} - \hat{\bar{y}}_{j,t}$, and since in general

$Q_j \bar{\hat{x}}_t \neq \bar{\bar{a}}'_{j,t} + u_{1,t} \bar{\bar{b}}'_{j1,t} + u_{2,t} \bar{\bar{b}}'_{j2,t} + \cdots + u_{r,t} \bar{\bar{b}}'_{jq,t}$, these expressions are equal if and only

if $\hat{\bar{y}}_{j,t} = \bar{y}_{j,t}$ and $\bar{\bar{a}}'_{j,t} = u_{1,t} \bar{\bar{b}}'_{j1,t} = u_{2,t} \bar{\bar{b}}'_{j2,t} = \cdots = u_{r,t} \bar{\bar{b}}'_{jq,t} = 0$.

Having established that $A_t = I_n$ results in $\bar{\bar{a}}'_{j,t} = 0$, $\bar{\bar{b}}'_{jk,t} = 0$ and $\hat{\bar{y}}_{j,t} = \bar{y}_{j,t}$, it is time to

find how fitness affects the incremental changes in mean reaction norm parameter values. For

clarity of presentation, we here limit the number of phenotypic traits to $p = 2$, and the number

of environmental cues to $q = 2$, as we also did in parts of Section 2. We also introduce the

diagonal $n \times n$ matrix $F_t = diag(w_{i,t})$, and the following two lemmas:

Lemma 2 *The mean value of the product of $\boldsymbol{F}_t$ and a predicted random effect vector $\widehat{\boldsymbol{z}}'_t$ in*

Eq. (8), where $\widehat{\boldsymbol{z}}'_t = \widehat{\boldsymbol{a}}'_{j,t}$ for j from 1 to p, or $\widehat{\boldsymbol{z}}'_t = \widehat{\boldsymbol{b}}'_{jk,t}$ for j from 1 to p and k from 1 to q,

is $\overline{\boldsymbol{F}_t \widehat{\boldsymbol{z}}'_t} = \Delta \bar{z}_t$, i.e., the incremental change in the mean reaction norm parameter value $\bar{z}_t$.

Proof: Since $\bar{w}_t = 1$ and $\overline{\widehat{\boldsymbol{z}}'}_t = 0$ (from Lemma 1), we find $\overline{\boldsymbol{F}_t \widehat{\boldsymbol{z}}'}_t = \frac{1}{n} \sum_{i=1}^{n} w_{i,t} \widehat{\boldsymbol{z}}'_{i,t} -$

$\bar{w}_t \overline{\widehat{\boldsymbol{z}}'}_t = cov(w_{i,t}, \widehat{\boldsymbol{z}}'_{i,t})$, and thus according to Theorem 1, $\overline{\boldsymbol{F}_t \widehat{\boldsymbol{z}}'}_t = \Delta \bar{z}_t$.

Lemma 3 *The mean value of the sum $\boldsymbol{F}_t \boldsymbol{y}_{j,t} - \boldsymbol{F}_t \mathbf{1}_n \bar{y}_{j,t}$, for j from 1 to p, is $cov(w_{i,t}, y_{j,i,t})$.*

Proof: $\overline{\boldsymbol{F}_t \boldsymbol{y}_{j,t}} - \overline{\boldsymbol{F}_t \mathbf{1}_n \bar{y}_{j,t}} = \frac{1}{n} \sum_{i=1}^{n} w_{i,t} y_{j,i,t} - \bar{w}_t \bar{y}_{j,t} = cov(w_{i,t}, y_{j,i,t})$.

Assuming $p = 2$ and $q = 2$, for simplicity, and using that $\hat{\bar{y}}_{j,t} = \bar{y}_{j,t}$ (Lemma 1),

multiplication of Eq. (A2a) from the left with the block diagonal matrix $\widetilde{\boldsymbol{F}}_t = \boldsymbol{I}_6 \otimes \boldsymbol{F}_t$, and

insertion of $\widetilde{\boldsymbol{F}}_t^{-1} \widetilde{\boldsymbol{F}}_t$ between $(\boldsymbol{Z}_t^T \boldsymbol{R}_t^{-1} \boldsymbol{Z}_t + \boldsymbol{G}^{-1})$ and $\otimes \boldsymbol{I}_n \widehat{\boldsymbol{x}}_t$, we find by use of the mixed

model property $(\boldsymbol{A} \otimes \boldsymbol{B})(\boldsymbol{C} \otimes \boldsymbol{D}) = (\boldsymbol{AC}) \otimes (\boldsymbol{BD})$ that

$$(\boldsymbol{Z}_t^T \boldsymbol{R}_t^{-1} \boldsymbol{Z}_t + \boldsymbol{G}^{-1}) \otimes \boldsymbol{F}_t \begin{bmatrix} \widehat{\boldsymbol{a}}'_{1,t} \\ \widehat{\boldsymbol{a}}'_{2,t} \\ \widehat{\boldsymbol{b}}'_{11,t} \\ \widehat{\boldsymbol{b}}'_{12,t} \\ \widehat{\boldsymbol{b}}'_{21,t} \\ \widehat{\boldsymbol{b}}'_{22,t} \end{bmatrix} = (\boldsymbol{Z}_t^T \boldsymbol{R}_t^{-1}) \otimes \boldsymbol{F}_t \begin{bmatrix} \boldsymbol{y}_{1,t} - \mathbf{1}_n \bar{y}_{1,t} \\ \boldsymbol{y}_{2,t} - \mathbf{1}_n \bar{y}_{2,t} \end{bmatrix}. \quad (A3a)$$

755 Since all elements in $\boldsymbol{Z}_t^T \boldsymbol{R}_t^{-1} \boldsymbol{Z}_t + \boldsymbol{G}^{-1}$ and $\boldsymbol{Z}_t^T \boldsymbol{R}_t^{-1}$ are multiplied by $\boldsymbol{F}_t$, Eq. (A3a) can be

756 reformulated as

$$(\boldsymbol{Z}_t^T \boldsymbol{R}_t^{-1} \boldsymbol{Z}_t + \boldsymbol{G}^{-1}) \begin{bmatrix} \boldsymbol{F}_t \widehat{\boldsymbol{a}}'_{1,t} \\ \boldsymbol{F}_t \widehat{\boldsymbol{a}}'_{2,t} \\ \boldsymbol{F}_t \widehat{\boldsymbol{b}}'_{11,t} \\ \boldsymbol{F}_t \widehat{\boldsymbol{b}}'_{12,t} \\ \boldsymbol{F}_t \widehat{\boldsymbol{b}}'_{21,t} \\ \boldsymbol{F}_t \widehat{\boldsymbol{b}}'_{22,t} \end{bmatrix} = \boldsymbol{Z}_t^T \boldsymbol{R}_t^{-1} \begin{bmatrix} \boldsymbol{F}_t \boldsymbol{y}_{1,t} - \boldsymbol{F}_t \mathbf{1}_n \bar{y}_{1,t} \\ \boldsymbol{F}_t \boldsymbol{y}_{2,t} - \boldsymbol{F}_t \mathbf{1}_n \bar{y}_{2,t} \end{bmatrix}. \quad (A3a)$$

757 Taking mean values of all vector elements, will according to Lemma 2 and 3 finally give

$$\begin{bmatrix} \Delta\bar{a}_{1,t} \\ \Delta\bar{a}_{2,t} \\ \Delta\bar{b}_{11,t} \\ \Delta\bar{b}_{12,t} \\ \Delta\bar{b}_{21,t} \\ \Delta\bar{b}_{22,t} \end{bmatrix} = G(Z_t^T R_t^{-1} Z_t G + I_6)^{-1} Z_t^T R_t^{-1} \begin{bmatrix} cov(w_{i,t}, y_{1,i,t}) \\ cov(w_{i,t}, y_{2,i,t}) \end{bmatrix}. \tag{A4}$$

Appendix B Proof of Theorem 3

Since $\begin{bmatrix} I_2 \\ U_t \end{bmatrix} = Z_t^T$, we must in order to prove Theorem 3 show that

$(Z_t^T R_t^{-1} Z_t G + I_6)^{-1} Z_t^T R_t^{-1} = Z_t^T P_{yy,t}^{-1}$. We can do that by use of the matrix inversion lemma

$(A + BCD)^{-1} = A^{-1} - A^{-1}B(C^{-1} + DA^{-1}B)^{-1}DA^{-1}$, with use of $A = I_6$,

$B = Z_t^T$, $C = R_t^{-1}$, and $D = Z_t G$. This gives

$$(Z_t^T R_t^{-1} Z_t G + I_6)^{-1} Z_t^T R_t^{-1} = (I_6 - Z_t^T (R_t + Z_t G Z_t^T)^{-1} Z_t G) Z_t^T R_t^{-1}. \tag{B1}$$

Here, $R_t + Z_t G Z_t^T = \begin{bmatrix} r_{1,t} & 0 \\ 0 & r_{2,t} \end{bmatrix} + [I_2 \quad U_t^T] \begin{bmatrix} G_{aa} & G_{ab} \\ G_{ab}^T & G_{bb} \end{bmatrix} \begin{bmatrix} I_2 \\ U_t \end{bmatrix}$, whitch with

$r_{1,t} = \sigma_{v_1}^2 + u_{1,t}^2 \sigma_{\eta_{11}}^2 + u_{1,t}^2 \sigma_{\eta_{12}}^2$ and $r_{2,t} = \sigma_{v_2}^2 + u_{1,t}^2 \sigma_{\eta_{21}}^2 + u_{2,t}^2 \sigma_{\eta_{22}}^2$ gives

$R_t + Z_t G Z_t^T = P_{aa} + 2U_t^T G_{ab} + U_t^T P_{bb} U_t = P_{yy,t}$, and thus $Z_t G Z_t^T = P_{yy,t} - R_t$. Eq.

(B1) thus finally gives

$$(Z_t^T R_t^{-1} Z_t G + I_6)^{-1} Z_t^T R_t^{-1} = Z_t^T R_t^{-1} - Z_t^T P_{yy,t}^{-1} Z_t G Z_t^T R_t^{-1}$$

$$= Z_t^T R_t^{-1} - Z_t^T P_{yy,t}^{-1} (P_{yy,t} - R_t) R_t^{-1} = Z_t^T P_{yy,t}^{-1}. \tag{B2}$$