A novel and highly specific Forssman antigen-binding protein from sheep polyomavirus

Nils H. Rustmeier[1], Lisete M. Silva[2], Antonio Di Maio[2], Joshua C. Müller[1], Alexander Herrmann[1], Ten Feizi[2], Yan Liu[2], Thilo Stehle[1]

[1]Interfaculty Institute of Biochemistry, University of Tübingen, Tübingen, Germany

[2]Glycosciences Laboratory, Faculty of Medicine, Imperial College London, London, UK

Corresponding author:

Thilo Stehle: thilo.stehle@uni-tuebingen.de

Phone: +49-7071-2978090

Abstract

Polyomaviruses are small, non-enveloped double-stranded DNA viruses of humans and other mammals, birds, and fish. Infections are usually asymptomatic and result in latency, however, some polyomaviruses can induce severe diseases, including cancer, in immunocompromised individuals. Established cellular receptors for polyomavirus infection are sialylated glycolipids (such as gangliosides), membrane proteins, and glycosaminoglycans. Polyomaviruses are usually highly host specific, but the exact

principles that govern host tropism remain unknown in many cases. Here, glycan array

screening shows that the major capsid protein VP1 of sheep polyomavirus (ShPyV)

binds to the Forssman glycolipid, an antigen of many vertebrates and a potential tumor

marker in humans. Following closer investigation, we can report for the first time that

a neutral, non-sialylated glycolipid acts as a polyomavirus receptor. Concurrently, we

present the first report of a viral protein that specifically engages the Forssman antigen.

We demonstrate that ShPyV VP1 binds to Forssman-positive erythrocytes but not

those of human A, B and O blood groups, which is a clear distinction from features

thus far described for Forssman lectins. X-ray crystallography and structural analysis

of the VP1-Forssman glycan complex define the terminal Forssman disaccharide as

the determinant of this protein-receptor interaction. These results strongly suggest that

the sheep polyomavirus can use Forssman antigen for infectious cell entry.

Furthermore, the ability of ShPyV VP1 to distinguish Forssman-positive from -negative

cells may prove useful for monitoring the Forssman-'status' of normal, preneoplastic

and neoplastic cells and tissues and establishing the antigen level as a biomarker.

Author summary

Elucidation of host cell receptor specificities of viral infection is crucial to understand

the pathobiology of associated diseases and develop treatments. However, for many

polyomaviruses the receptor engagement as the initial event in infection is poorly

understood. In only a few cases polyomavirus tropism has been pinned down to a

single type of glycan receptor. While many polyomaviruses utilize sialyl glycans to

attach to host cells, the role of non-sialylated glycans as receptors is so far

underestimated. Here, we show for the first time that a glycan of neutral charge, in this

case the carbohydrate portion of the Forssman antigen, acts as an attachment receptor

for a polyomavirus capsid protein and may thus contribute to host tropism and infective

cell entry. These results represent a significant addition to knowledge on polyomavirus-glycan interactions and complement general principles of carbohydrate engagement by viruses. Furthermore, as a specific binding protein of Forssman antigen, VP1 may help to determine levels of this antigen in healthy and malignant tissues in humans.

Introduction

The Forssman antigen (FA), a neutral globo-series glycosphingolipid (GSL, see Table 1), is produced by the glycosyltransferase activity of Forssman synthetase (FS)[1,2]. Encoded by the *GBGT1* gene, FS catalyzes the addition of αGalNAc to the non-reducing end of globoside (see Table 1). This reaction results in the key immunogenic motif of the Forssman antigen, the terminal GalNAcα1-3GalNAcβ Forssman disaccharide[3]. With few exceptions in humans, *GBGT1* is transcribed but encodes for an inactive FS, resulting in a predominantly Forssman-negative human population[4,5]. So far, only a small number of healthy individuals worldwide were found to carry an activating mutation within *GBGT1.* In these individuals, the resulting allele gives rise to a Forssman antigen-positive (FA+) erythrocyte phenotype, which is now recognized as the human FORS blood group[6–8]. Other reports of healthy human FA+ tissues are scarce, and human sera often contain natural anti-FA antibodies[9–11]. More frequently, Forssman antigenicity was described in cancerogenesis and a few other diseases of humans[12–21]. Yet, both the genetics and the precise role of FA expression in disease remain poorly understood.

The Forssman glycolipid also serves as a receptor for different bacterial infections in animals[22–28], however, it is so far unknown if viruses also utilize FA to target their hosts. In this study, by means of glycan microarray screening, NMR, surface plasmon

resonance (SPR), and cell binding assays, we show that FA, which is the predominant

glycolipid in ovine erythrocytes[29], serves as a cellular attachment receptor for the VP1

of a polyomavirus originally identified in sheep[30]. This designated sheep polyomavirus

(ShPyV) is a member of a non-enveloped, double-stranded DNA virus family with

icosahedral capsids, scaffolded by the major capsid protein VP1. To initiate an infection

many polyomaviruses rely on the interaction between their VP1 and sialic acid-

equipped glycolipids in the target cell membranes[31–36]. Although other infection

receptors such as glycoproteins and glycosaminoglycans have also been reported for

some polyomaviruses[37–42], detailed structural information is so far only available for

the interactions between VP1s and sialylated oligosaccharides. Thus, our studies

present the first example of a polyomavirus VP1 protein interacting with a non-sialyl

neutral glycan, complementing the principles of how polyomaviruses recognize their

targets.

Results

<u>Sheep polyomavirus VP1 binds the Forssman epitope with a strong preference over</u>

<u>other glycans</u>

To identify potential attachment factors of ShPyV, we prepared recombinant VP1

pentamers as previously described[43]. The protein was analyzed using a neoglycolipid-

based glycan screening array encompassing a broad spectrum of lipid-linked glycan

probes[44]. A full list of 672 glycan probes is in Table S1, and these include major types

of mammalian sequences found on glycoproteins (N- and O-linked), glycolipids, blood

group and Lewis antigen related, oligosaccharide fragments of glycosaminoglycan

heparin, as well as those derived from polysaccharides of bacteria, fungi, and plants.

Among these are 204 sialylated glycan probes variously bound by several

polyomavirus VP1s described previously[45–49]. We detected a signal for the Forssman

glycolipid that surpasses all other signals by at least one order of magnitude (Fig. 1).

A dominance of the terminal GalNAcα1-3GalNAcβ Forssman disaccharide is apparent

as the VP1 did not give binding signals with globoside glycolipid (P-antigen, position

25, Table S1), the precursor of the Forssman glycolipid which lacks the terminal

α-linked GalNAc residue. Furthermore, there were no signals with other GalNAc-

terminating glycan probes, including the blood group A related glycan probes, and the

two Tn antigen probes GalNAcα-Ser and -Thr (positions 221 and 222, Table S1). The

disaccharide probe GalNAcα1-3GalNAc (position 223) with a ring-opened GalNAc

core (reduced by reductive amination) was also not bound. These suggest the

importance of adjoining monosaccharides beyond the αGalNAc residue for ShPyV VP1

binding. Among the sialylated glycans analyzed only one short α2,6-linked sialyl probe

with a 9-O-acetylated N-acetylneuraminic acid residue (Neu5,9Ac$_2$) yielded a weak

signal (position 607). No binding signals were detected in the negative control

experiment with detection antibodies alone in the absence of ShPyV VP1 (Table S1).

To assess if the glycan binding revealed by the glycan microarray also occurs in

solution, we evaluated the interactions between ShPyV VP1 and the Forssman

pentaose (F$_P$) in the presence of α2,6-sialyllactose (6'SL) by saturation transfer

difference NMR (STD NMR) spectroscopy, which reveals protein-to-ligand

magnetization transfer in the case of short-term binding (10^{-8} M < K$_D$ < 10^{-3} M)[50,51]. In

the first instance, the ^{1}H reference resonances of isolated F$_P$ and 6'SL were collected

(Fig. 2a and 2b), which allowed us to unambiguously assign the oligosaccharide

signals[43,52] (except for the overlaps within 3.5-4.0 ppm) in the ^{1}H spectrum of a sample

comprising VP1, F$_P$, and 6'SL (Fig. 2c). Subsequently, we used the same sample to

perform the STD NMR experiment. Evaluation of the resulting spectrum reveals

magnetization transfer to both F_P and 6'SL, which infers that VP1 interacts with both

glycans in the solution (Fig. 2b). However, while resonances derived from all

monosaccharides in F_P (two GalNAc, two Gal, and a Glc) received magnetization

transfer, only the sialic acid methyl group of 6'SL re-emerged in the STD spectrum.

Yet, these results are in good agreement with the findings from the glycan array

analysis of ShPyV VP1, which indicated a much higher affinity for F_P than sialylated

glycans (Fig. 1). Strikingly, the methyl group of the αGalNAc in F_P gives a particularly

strong peak in the STD spectrum (Fig. 2b), which implies a prominent (and potentially

specificity-conferring) role in the VP1-F_P interaction.

<u>ShPyV VP1 has a nanomolar affinity towards Forssman pentaose-decorated surfaces</u>

After assessing the complex formation between ShPyV VP1 and F_P in the context of

the solid-supported glycan array and solution NMR, we sought to evaluate the

underlying binding kinetics by surface plasmon resonance (SPR) and compare the

performance of VP1 with the albumen gland agglutinin of *Helix pomatia* (HPA), an

αGalNAc-specific lectin with high affinity towards the Forssman antigen[53,54]. Therefore,

we prepared an SPR chip decorated with F_P (see Fig. S3) and measured binding

responses using different dilutions of VP1 and HPA. The resulting response curves

show that within the given association time the pentavalent VP1 and hexavalent HPA

bind to the immobilized F_P without reaching a complex formation equilibrium, which

suggests that the proteins do not readily dissociate from the surface (Fig. 3a). Indeed,

fitting of the kinetic parameters results in very low dissociation rates (k_{off}) for both

proteins, which is a common phenomenon for multivalent binding systems. However,

the curves of VP1 result in a higher rate for complex formation (k_{on}), determining an

overall approximately three-fold lower dissociation constant (K_D) compared to HPA

(Fig. 3a).

<u>ShPyV VP1 specifically binds to Forssman-positive erythrocytes</u>

To confirm the functional relevance of the VP1-F_P complex *in vitro*, we performed cell

binding assays using different erythrocytes and compare the cell binding of VP1 with

commercial reagents for Forssman antigen detection, i.e., HPA and the monoclonal

anti-FA IgM antibody M1/87[55]. We incubated VP1 with Forssman-positive red blood

cells (RBCs) from sheep and human RBCs of the different ABO phenotypes and found

that ShPyV VP1 bound sheep RBCs with saturation but did not attach to any of the

human erythrocytes (Fig. 3b left). In contrast, the αGalNAc-specific lectin HPA bound

to the FA on sheep RBCs and also to the antigens on human blood group A1 and A2

RBCs[56] (Fig. 3b left). To confirm that the carbohydrate portions of the antigens are

responsible for binding, we incubated the two proteins with excessive amounts of the

terminal trisaccharides (trioses) of FA (GalNAcα1-3GalNAcβ1-3Gal) and A substance

(GalNAcα1-3[Fucα1-2]Gal) and assessed the remaining sheep RBCs attachment. As

predicted, FA triose blocked VP1 and HPA, whereas A antigen triose could only inhibit

HPA attachment (Fig. 3b right). Lastly, we determined the effective protein

concentrations (EC50, see Methods) for sheep RBCs attachment using different

dilutions of ShPyV VP1, HPA, and M1/87. Evaluation of the response curves yielded

the lowest EC50 for ShPyV VP1 with a value of 1.2 nM, followed by M1/87 and HPA

with 3.3 nM and 5.2 nM, respectively (Fig. 3c). Collectively, these results demonstrate

the remarkable specificity and affinity of ShPyV VP1 towards the Forssman antigen.

<u>Crystal structure of Forssman pentaose bound to VP1 reveals an intricate network of</u>

<u>interactions</u>

To characterize the binding of Forssman antigen by ShPyV VP1 with atomic detail, we

solved the crystal structure of the VP1 in complex with 10 mM of Forssman pentaose

(F_P) at 1.92 Å resolution (Table S4). Initial electron density calculation using

unliganded ShPyV VP1 model (PDB 6Y61) shows that five molecules of the

oligosaccharide (GalNAcα1-3GalNAcβ1-3Galα1-4Galβ1-4Glc) associate with the

outer surface of the pentameric VP1, at a position that does not overlap with the

previously described sialic acid binding site[43] (Fig. S5), in a five-fold symmetric array.

All five Forssman binding sites are identical and located between two neighboring VP1

monomers (Fig. 4a). In the binding sites without artificial crystal contacts, electron

density provides an unambiguous outline of the last four monosaccharides of F_P (Fig.

4b). The intermolecular contacts show that the terminal trisaccharide of F_P contributes

to binding, whereas the remaining βGal and Glc divert away from VP1 and do not

interact with the protein surface. Closer inspection of the binding site reveals that the

methyl group of the terminal αGalNAc invades a hydrophobic surface depression

constituted by phenylalanine residues of two neighboring VP1 monomers, establishing

van der Waals interactions (Fig. 4c). Additionally, the *N*-acetyl moiety of αGalNAc

interacts with the Q266 and F66cw residues via hydrogen bonds (Fig. 4c). In βGalNAc,

the carbonyl oxygen forms two hydrogen bonds with the side chains of Q266 and N268

(Fig. 4c). Apart from these contacts, numerous interactions involve the hydroxyl groups

of F_P that are directed towards the protein surface. An overview of all contacts and their

distances is provided in Table 2. We also solved the crystal structure of VP1 in complex

with the FA glycan precursor, globo-*N*-tetraose, which reveals negligible binding only

(Fig. S5). We therefore conclude that the terminal GalNAcα1-3GalNAc disaccharide is

essential for the high affinity of the ShPyV VP1-FP complex.

Discussion

Sheep polyomavirus (ShPyV) was originally identified as a genomic contaminant in

processed lamb meat and it remains unknown if sheep are the natural hosts of this

virus[30]. Our functional data show that ShPyV engages Forssman antigen (FA)-

decorated cells with high affinity and specificity, making it plausible that FA initiates

sheep polyomavirus infection. However, in sheep the FA was thus far only detected on

erythrocytes but not on leukocytes, bone marrow or other tissues[29,57,58]. However, as

the studies about the distribution of Forssman antigen in sheep tissues are rather old,

the more sensitive techniques available today could help to elucidate the potential

targets cells for viral replication as well as the precise role of Forssman antigen during

ShPyV infection.

The antigenic determinant of Forssman antigen is the GalNAcα1-3GalNAcβ

disaccharide. In sheep erythrocytes it is predominantly linked to globotriasolyceramide,

giving rise to the linear Forssman pentaosylceramide (see Table 1), but it also appears

on the (unbranched) glycans of neolacto-series or Galili glycolipids[29]. In other animals,

GalNAcα1-3GalNAcβ occurs on branched structures such as extended H-like antigens

(dog gastric mucosa and mucus)[59,60], GM1 ganglioside (English sole fish)[61], isoglobo-

ganglio-neolacto glycans (swine)[62], as well as in the form of mucin core 5 (GalNAcα1-

3[NeuAca2-6]GalNAc) in bovine submaxillary mucin[63]. We believe that ShPyV VP1

can bind a range of different Forssman glycans, providing that the structural deviations

do not result in clashes with the protein. By the assessment of our ShPyV VP1-

Forssman pentaose (F$_P$) complex structure, the designated minimal glycan binding

motif is the GalNAcα1-3GalNAc disaccharide with few additional contacts to the

subsequent αGal. From structural perspective, this epitope recognition clearly

disallows branching of the glycan at certain positions, e.g., O3 of αGalNAc, N2 of

βGalNAc or O4 of αGal. For example, in the blood group A glycan fucose replaces the

N-acetyl group of the βGalNAc, which would result in overlaps with the protein and

accounts for the lack of binding (Fig. S5b). This feature distinguishes VP1 from many

other Forssman glycan-binding proteins such as Galectin-9, *Helix pomatia* agglutinin

(HPA), *Sinularia lochmodes* lectin-2, and *Dolichos biflorus* agglutinin, where

recognition is based on the subterminal or terminal GalNAc (see Fig. S6), resulting in

a much broader glycan binding capacity[64–67].

Previously we have shown that ShPyV VP1 interacts with sialylated trisaccharides in

crystal structures[43]. Also, our newly presented results from STD NMR and

crystallography reveal that Forssman and sialyl glycan binding occurs on separate

sites of the protein surface (Fig. S5b), most likely independently from each other. Yet,

in our flow cytometry assays we did not detect any binding to human erythrocytes

although they are known to be decorated with sialic acids. However, the preferred form

of sialic acid for ShPyV VP1, as evinced by our glycan array data, is 9-*O*-acetylated

N-acetylneuraminic acid (Neu5,9Ac$_2$), which in human erythrocytes is present in too

low amounts to result in detectable binding[68]. These findings imply that human cells

are generally incompatible targets for the sheep virus.

Humans are considered a largely Forssman-negative species and the presence of

Forssman antigenicity is mostly associated with human cancer[69]. During the last

decade, the elucidation of the genetic basis behind FA reactivation and the

establishment of the human FORS blood group underline the continual relevance of

Forssman antigen[5,6]. We propose that ShPyV VP1 complements and expands the

available resources for the comprehensive detection of Forssman antigen in cellular

and histological samples of healthy or malignant provenance. We also anticipate

refining mutant VP1 protein to precisely target the Forssman antigen for biomedical

use.

Summary

We show that the Forssman antigen serves as a cellular attachment receptor for the

major capsid protein VP1 of ShPyV. After initial detection by glycan array screening,

in-solution NMR spectroscopy and cell binding assays demonstrate functional

relevance. Structural analysis of VP1 in complex with the carbohydrate moiety of FA,

Forssman pentaose (F_P), characterizes the determinants of the underlying interactions

with high accuracy. Our findings have significance for the fields of structural virology,

glycobiology, and pathobiology as follows:

(i) We report the first structural and functional data of a viral entity to specifically

interact with the Forssman antigen.

(ii) The ShPyV VP1 – F_P complex provides the first detailed information on how

a polyomavirus binds to a non-sialylated ligand.

(iii) The high specificity and affinity of ShPyV VP1 toward Forssman antigen

establishes this protein as a potential biomedical reagent.

Materials and methods

<u>Protein expression and purification</u>

The VP1 of sheep polyomavirus was recombinantly produced as previously.[43] Briefly,

transformed *E. coli* BL21 (DE3) were grown in LB (Miller) medium supplemented with

100 µg/ml ampicillin at 37 °C. Expression was induced at an OD_{600} of 0.7 with 400 µM

IPTG and carried out at 20 °C for 16 h. ShPyV VP1 was extracted from bacterial pellets

by sonication and subsequent centrifugation (17,000 rpm at 4 °C). The His_6-tagged

VP1 was purified by immobilized-metal affinity chromatography (IMAC) and

subsequent gel filtration (20 mM HEPES pH 7.5, 150 mM NaCl). For crystallization of

ShPyV VP1 the His_6-tag was cleaved off by incubation with 10 U/mg thrombin

overnight at room temperature. Tags and thrombin were separated from ShPyV VP1

by IMAC and gel filtration. If not immediately used, the protein was stored at -20 °C.

Glycan array

The His_6-tagged ShPyV VP1 was analyzed in the neoglycolipid (NGL)-based

microarray system[70]. A broad-spectrum screening glycan microarray set

encompassing 672 sequence-defined lipid-linked glycan probes was used (the glycan

probes included, and their sequences are given in Supplemental Table S1). Details of

the preparation of the glycan probes and the generation of the microarrays are in

Supplementary Glycan Microarray Document (Supplemental Table S2) in accordance

with the MIRAGE (Minimum Information Required for A Glycomics Experiment)

guidelines for reporting of glycan microarray-based data[71]. The microarray analyses

were performed essentially as described[47,48]. In brief, after blocking the slides for 1h

with HBS buffer (10 mM HEPES, pH 7.4, 150 mM NaCl) containing 0.33% (w/v) blocker

Casein (Pierce), 0.3% (w/v) Bovine Serum Albumin (Sigma-Aldrich) and 5 mM $CaCl_2$,

the microarray was overlaid with the VP1 protein for 90 minutes as a protein-antibody

complex that was prepared by preincubating VP1 with mouse monoclonal anti-

polyhistidine and biotinylated anti-mouse IgG antibodies (both from Sigma) at a ratio

of 4:2:1 (by weight) and diluted in the blocking solution to provide a final VP1

concentration of 150 μg/mL. Binding was detected with Alexa Fluor-647-labelled

streptavidin (Molecular Probes) at 1 μg/mL for 30 minutes. Apart from the protein-

antibody precomplexation step, which was performed on ice, all other steps were

carried out at ambient temperature. Microarray imaging and data analysis are

described in the Supplementary MIRAGE document (Supplemental Table S2).

NMR

All nuclear magnetic resonance (NMR) data were collected in a Bruker AVIII-600 MHz

spectrometer equipped with a TXI-z probe head at 298 K using 3 mm NMR tubes.

Samples were provided in 99 % D_2O-PBS containing 20 mM KH_2PO_4/K_2HPO_4 (pH 7.4)

and 150 mM NaCl. [1]H reference spectra of isolate oligosaccharides samples were

recorded using pure solutions of 4 mM of Forssman pentaose and 2 mM of 6'SL.

Subsequently, the [1]H oligosaccharide resonances of a sample containing 1 mM

Forssman pentaose and 1 mM 6'SL in the presence of 20 μM ShPyV VP1 were

collected under suppression of protein signals. From the same sample, the saturation

difference transfer (STD) NMR experiment was performed using previously published

pulse protocols[43]. NMR spectra were processed using TOPSPIN 4 (Bruker). The

carbohydrates used here were purchased from Biosynth, UK.

VP1 mutagenesis

Site-directed mutagenesis of VP1 was performed in samples of 20 μl containing

1x ReproFast amplification buffer (Genaxxon), 20 ng template vector DNA, 50 ng

primers, 2 mM dNTP, and 1 unit of ReproFast polymerase (Genaxxon). Initial

denaturation was performed at 95 °C for 2 minutes, followed by 18 cycles of 95 °C for

1 minute (denaturation), 55-70 °C for 1 minute (annealing), and 72 °C for 6 minutes

(elongation). Subsequently, DNA synthesis was completed at 72 °C for 10 minutes.

Parental template DNA was digested with 2 units of DpnI for two hours at 37 °C. The

PCR product was directly transformed into *E. coli* DH5α.

DNA primers for the site-directed mutagenesis of ShPyV VP1 were ordered from

ThermoFisher Scientific and are displayed from 5'- to 3' ends:

S95C_fwd: GCTGAACCAAGACATGACCTGCGATACCATCCTGATGTGGGAGG
S95C_rev: CCTCCCACATCAGGATGGTATCGCAGGTCATGTCTTGGTTCAGC

<u>Fluorescence labeling</u>

VP1 was expressed and purified as described above. Surficial cysteine residues were

reduced in degassed SEC buffer (20 mM HEPES pH 7.4, 150 mM NaCl) supplemented

with 80 mM DTT at 4 °C overnight. DTT was removed from the protein solution using

a PD-10 desalting column (Cytiva) with DTT-free SEC buffer. A 20-fold molar excess

of cysteine reactive AFDye 488 Maleimide (Fluoroprobes) was added to the protein

solutions (0.5 - 0.7 mg/ml) and the reaction was incubated in the dark at 4 °C for 16-

20 h. Excess dye was quenched and removed by desalting in SEC buffer

supplemented with 10 mM DTT. VP1-488 conjugate was eluted in the same buffer.

The degree of labeling was photometrically determined according to the

manufacturer's protocol.

<u>Flow cytometry</u>

Defibrinated blood from sheep was purchased from TCS Biosciences Ltd and

BioTrading Benelux B.V., and human reference erythrocytes from Immucor. Prior to

analysis, all blood was diluted to 0.25 % in 1x PBS + 2 % fetal calf serum (FC buffer).

For staining, labeled protein was added to samples of ~1×10^6 cells in a total volume of

100 µL. After incubation in the dark for 15 minutes, cells were precipitated by

centrifugation (1.5 min at 1900 rpm) and resuspended in fresh FC buffer. Unstained

reference samples were prepared accordingly without the addition of labeled protein.

For cell attachment inhibition assays, proteins were incubated with oligosaccharides

(Forssman antigen trisaccharide: Elicityl, France; Blood group A trisaccharide:

Biosynth, UK) for 5 minutes prior to their addition to the cellular samples (~1×10^6 cells

in 100 µL). To determine the concentration for half maximal cell attachment (effective

concentration, EC50), samples comprising serially diluted concentrations of labeled

protein, starting from 10 µg/mL for M1/87-FITC (Santa Cruz Biotechnology) and HPA

Alexa Fluor^{TM} 488 conjugate (Invitrogen), and 5 µg/mL for VP1-488, were prepared as

described above. Here, the wash step was omitted to guarantee consistent protein

concentrations and to prevent agglutination. All samples were transferred to the flow

cytometer (LSR II, BD Biosciences) and fluorescent fractions were captured at 520 nM.

100,000 events were recorded for all samples except those with agglutination, in which

case 10,000-20,000 events were recorded. All data were analyzed using FACSdiva

software (BD Biosciences). EC50 values were calculated using R[72].

<u>Surface plasmon resonance</u>

Assessment of the kinetic binding parameters were performed via surface plasmon

resonance (SPR) in a Biacore X-100 system. Using an adapted protocol[73], BSA

decorated with Forssman pentaose (~76 kDa) as the ligand was prepared via reductive

amination (see Fig. S3). After rebuffering into 10 mM acetate buffer at pH 4,

approximately 100 response units (RU) of ligand were immobilized onto the CM5

sensor chip surface using amine coupling as described in the manufacturer's protocol

(Cytiva). For the reference flow cell, native BSA (Sigma-Aldrich) was immobilized

similarly. Purified VP1 (162 kDa) and native HPA (79 kDa, Sigma-Aldrich) buffered in

SPR assay buffer (20 mM HEPES pH 7.4, 150 mM NaCl, 0.005 % Polysorbate 20)

were used as analytes in different concentrations between 0.2 nM and 20 µM. All

measurements were performed at 25 °C. During binding (120 s) and dissociation

(180 s), a flow rate of 30 µL/min was applied. After each cycle, the surfaces were

regenerated applying 10 mM glycine pH 1.5 for 120 s. Experimental curves were

interpreted using BIAevaluation software (Cytiva).

<u>Crystallization and data collection</u>

Purified ShPyV VP1 was concentrated to 3.5 mg/ml. Crystallization droplets comprised

1 µl protein and 1 µl reservoir solution (150 mM KSCN and 20 % (w/v) PEG 3,350).

The droplets were equilibrated against the 500 µl reservoir using the hanging drop

vapor diffusion method at 20 °C. For ligand derivatization, VP1 crystals were soaked

in crystallization solution supplemented with the respective oligosaccharides (10 mM

Forssman pentaose (F$_P$), 10 mM globo-*N*-tetraose, 5 mM F$_P$ + 20 mM 3'SLN, 5 mM F$_P$

+ 20 mM 6'SLN, 20 mM A antigen trisaccharide (all purchased from Biosynth, UK), and

20 mM Tf antigen (TCI Europe), at 20 °C for half an hour. Prior to flash freezing, crystals

were transferred to oligosaccharide soaking solutions supplemented with 20 % (v/v)

MPD for cryo-protection. X-ray diffraction experiments were performed at the Swiss

Light Source (SLS) beamline X06DA (PSI in Villigen, Switzerland). Data were collected

at a wavelength of 1 Å with an excitation time and angle of 0.1 s and 0.1° per image.

<u>Structure determination and refinement</u>

X-ray diffraction images were integrated and scaled using *XDS*[74]. For each data set,

2,000-3,000 reflections were reserved as a test set for structure validation. Phases

were determined by molecular replacement using the native sheep polyomavirus VP1

structure as search model (PDB 6Y61) in Phaser (CCP4)[75]. Phase refinement and

model building were performed until validation parameters converged using Phenix and Refmac5, and Coot, respectively[76–78]. Oligosaccharides conformations are in accordance with the restraints from the CCP4 libraries. All protein structure representations were rendered in PyMOL (Schrödinger).

Data availability

Atomic coordinates of ShPyV VP1 in complex with carbohydrate ligands are deposited in the RCSB protein data bank (PDB) with the accession codes 7B6S (Forssman pentaose, F$_P$), 7B6T (globo-*N*-tetraose), 7B6U (F$_P$ + 6'SLN), and 7B6V (F$_P$ + 3'SLN). Other data can be provided by the authors upon request.

References

1. B Siddiqui, S. H. A revised structure for the Forssman glycolipid hapten. *J Biol Chem* **246**, 5766–5769 (1971).
2. Fraser, B. A. & Mallette, M. F. Structure of forssman hapten glycosphingolipid from sheep erythrocytes. *Immunochemistry* **11**, 581–593 (1974).
3. Haslam, D. B., Baenziger, J. U., Baenzigert, J. U. & Baenziger, J. U. Expression cloning of Forssman glycolipid synthetase: A novel member of the histo-blood group ABO gene family. *Proc. Natl. Acad. Sci. U. S. A.* **93**, 10697–10702 (1996).
4. Xu, H., Storch, T., Yu, M., Elliott, S. P. & Haslam, D. B. Characterization of the human Forssman synthetase gene. An evolving association between glycolipid synthesis and host-microbial interactions. *J. Biol. Chem.* **274**, 29390–29398 (1999).
5. Yamamoto, M., Cid, E. & Yamamoto, F. Molecular genetic basis of the human Forssman glycolipid antigen negativity. *Sci. Rep.* **2**, 975 (2012).
6. Svensson, L. *et al.* Forssman expression on human erythrocytes: Biochemical and genetic evidence of a new histo-blood group system. *Blood* **121**, 1459–1468 (2013).
7. Storry, J. R. *et al.* International Society of Blood Transfusion Working Party on red cell immunogenetics and blood group terminology: Cancun report (2012). *Vox Sang.* **107**, 90–96 (2014).
8. Hult, A. K. & Olsson, M. L. The FORS awakens: review of a blood group system reborn. *Immunohematology* **33**, 64–72 (2017).
9. Ono, K. *et al.* Expression of Forssman antigen in human large intestine. *J. Histochem. Cytochem.* **42**, 659–665 (1994).
10. Jesus, C. *et al.* Prevalence of antibodies to a new histo-blood system: the FORS system. *Blood Transfus.* **16**, 178 (2018).
11. Hult, A. K., McSherry, E., Möller, M. & Olsson, M. L. GBGT1 is allelically diverse but dispensable in humans and naturally occurring anti-FORS1 shows an ABO-restricted pattern. *Transfusion* **58**, 2036–2045 (2018).
12. Kawanami, J. The appearance of Forssman hapten in human tumor. *J. Biochem.* **72**, 783–785 (1972).
13. Yoda, Y., Ishibashi, T. & Makita, A. Isolation, Characterization, and Biosynthesis of Forssman Antigen in Human Lung and Lung Carcinoma. *J. Biochem.* **88**, 1887–1894 (1980).
14. Taniguchi, N., Yokosawa, N., Narita, M., Mitsuyama, T. & Makita, A. Expression of forssman antigen synthesis and degradation in human lung cancer. *J. Natl. Cancer Inst.* **67**, 577–583 (1981).

15. Yokota, M., Warner, G. A. & Hakomori, S. Blood Group A-like Glycolipid and a Novel Forssman
Antigen in the Hepatocarcinoma of a Blood Group O Individual. *Cancer Res.* **41**, (1981).
16. Mori, T., Sudo, T. & Kano, K. Expression of Heterophil Forssman Antigen on Cultured
Malignant Cell Lines. *JNCI J. Natl. Cancer Inst.* **70**, 811–818 (1983).
17. Hakomori, S. *Tumor-Associated Carbohydrate Antigens. Annual review of immunology* vol. 2
103–126 (Annual Reviews 4139 El Camino Way, P.O. Box 10139, Palo Alto, CA 94303-0139,
USA , 1984).
18. Uemura, K., Hattori, H., Ono, K., Ogata, H. & Taketomi, T. Expression of Forssman glycolipid
and blood group-related antigens A, Le(x) and Le(y) in human gastric cancer and in fetal
tissues. *Jpn. J. Exp. Med.* **59**, 239–249 (1989).
19. Säljö, K., Thornell, A., Jin, C., Norlén, O. & Teneberg, S. Characterization of human medullary
thyroid carcinoma glycosphingolipids identifies potential cancer markers. *Int. J. Mol. Sci.* **22**,
10463 (2021).
20. Ariga, T., Yoshida, T., Mimori, T. & Yu, R. K. Autoantibodies against Forssman glycolipids in
Graves' disease and Hashimoto's thyroiditis. *Clin. Exp. Immunol.* **86**, 483–488 (1991).
21. Hakomori, S., Wang, S. M. & Young, W. W. Isoantigenic expression of Forssman glycolipid in
human gastric and colonic mucosa: Its possible identity with 'A like antigen' in human cancer.
Proc. Natl. Acad. Sci. U. S. A. **74**, 3023–3027 (1977).
22. Stromberg, N. *et al.* Host-specificity of uropathogenic Escherichia coli depends on differences
in binding specificity to Galα1-4Gal-containing isoreceptors. *EMBO J.* **9**, 2001–2010 (1990).
23. Stromberg, N., Nyholm, P. G., Pascher, I. & Normark, S. Saccharide orientation at the cell
surface affects glycolipid receptor function. *Proc. Natl. Acad. Sci. U. S. A.* **88**, 9340–9344
(1991).
24. Johanson, I., Lindstedt, R. & Svanborg, C. Roles of the pap- and prs-encoded adhesins in
Escherichia coli adherence to human uroepithelial cells. *Infect. Immun.* **60**, 3416–3422 (1992).
25. Lanne, B. *et al.* Glycoconjugate receptors for P-fimbriated Escherichia coli in the mouse: An
animal model of urinary tract infection. *J. Biol. Chem.* **270**, 9017–9025 (1995).
26. Elliott, S. P., Yu, M., Xu, H. & Haslam, D. B. Forssman synthetase expression results in
diminished Shiga toxin susceptibility: A role for glycolipids in determining host-microbe
interactions. *Infect. Immun.* **71**, 6543–6552 (2003).
27. Müthing, J. *et al.* Promiscuous Shiga toxin 2e and its intimate relationship to Forssman.
Glycobiology **22**, 849–862 (2012).
28. Nasir, W., Nilsson, J., Olofsson, S., Bally, M. & Rydell, G. E. Parvovirus B19 VLP recognizes
globoside in supported lipid bilayers. *Virology* **456–457**, 364–369 (2014).
29. Santos, L. *et al.* Characterization of sheep erythrocyte glycosphingolipids recognized by human
anti-Forssman antibodies. *Glycobiology* (2020) doi:10.1093/glycob/cwaa032.
30. Buck, C. B. *et al.* The ancient evolutionary history of polyomaviruses. *PLoS Pathog.* **12**,
1005574 (2016).
31. Fried, H., Cahan, L. D. & Paulson, J. C. Polyoma virus recognizes specific
sialyloligosaccharide receptors on host cells. *Virology* **109**, 188–192 (1981).
32. Tsai, B. *et al.* Gangliosides are receptors for murine polyoma virus and SV40. *EMBO J.* **22**,
4346–4355 (2003).
33. Smith, A. E., Lilie, H. & Helenius, A. Ganglioside-dependent cell attachment and endocytosis of
murine polyomavirus-like particles. *FEBS Lett.* (2003) doi:10.1016/S0014-5793(03)01220-1.
34. Erickson, K. D., Garcea, R. L. & Tsai, B. Ganglioside GT1b is a putative host cell receptor for
the Merkel cell polyomavirus. *J. Virol.* **83**, 10275–10279 (2009).
35. Gilbert, J. *et al.* Ganglioside GD1a Restores Infectibility to Mouse Cells Lacking Functional
Receptors for Polyomavirus. *J. Virol.* **79**, 615–618 (2005).
36. Blaum, B. S. & Stehle, T. Sialic acids in nonenveloped virus infections. in *Advances in
Carbohydrate Chemistry and Biochemistry* vol. 76 65–111 (Academic Press Inc., 2018).
37. Elphick, G. F. *et al.* The human polyomavirus, JCV, uses serotonin receptors to infect cells.
Science (80-.). **306**, 1380–1383 (2004).
38. Assetta, B. *et al.* 5-HT2 receptors facilitate JC polyomavirus entry. *J. Virol.* **87**, 13490 (2013).
39. Schowalter, R. M., Pastrana, D. V. & Buck, C. B. Glycosaminoglycans and sialylated glycans
sequentially facilitate merkel cell polyomavirus infectious entry. *PLoS Pathog.* **7**, 1002161
(2011).
40. Hurdiss, D. L., Frank, M., Snowden, J. S., Macdonald, A. & Ranson, N. A. The structure of an
infectious human polyomavirus and its interactions with cellular receptors. *Structure* **26**, 839
(2018).
41. Geoghegan, E. M. *et al.* Infectious entry and neutralization of pathogenic JC polyomaviruses.
Cell Rep. **21**, 1169–1179 (2017).
42. Assetta, B. *et al.* Genetic and functional dissection of the role of individual 5-HT2 receptors as

entry receptors for JC polyomavirus. *Cell Rep.* **27**, 1960-1966.e6 (2019).

43.   Ströh, L. J. *et al.* Structural basis and evolution of glycan receptor specificities within the
polyomavirus family. *MBio* **11**, 1–21 (2020).

44.   Palma, A. S., Feizi, T., Childs, R. A., Chai, W. & Liu, Y. The neoglycolipid (NGL)-based
oligosaccharide microarray system poised to decipher the meta-glycome. *Curr. Opin. Chem.*
Biol. **18**, 87–94 (2014).

45.   Campanero-Rhodes, M. A. *et al.* N-glycolyl GM1 ganglioside as a receptor for simian virus 40.
J. Virol. **81**, 12846–12858 (2007).

46.   Neu, U. *et al.* Structure-function analysis of the human JC polyomavirus establishes the LSTc
pentasaccharide as a functional receptor motif. *Cell Host Microbe* **8**, 309–319 (2010).

47.   Neu, U. *et al.* A structure-guided mutation in the major capsid protein retargets BK
polyomavirus. *PLoS Pathog.* **9**, 1–13 (2013).

48.   Khan, Z. M. *et al.* Crystallographic and glycan microarray analysis of human polyomavirus 9
VP1 identifies N-glycolyl neuraminic acid as a receptor candidate. *J. Virol.* **88**, 6100–6111
(2014).

49.   Sorin, M. N. *et al.* Structural and functional analysis of natural capsid variants suggests sialic
acid-independent entry of BK polyomavirus. *CellReports* **42**, 112114 (2023).

50.   Mayer, M. & Meyer, B. Group epitope mapping by saturation transfer difference NMR to identify
segments of a ligand in direct contact with a protein receptor. *J. Am. Chem. Soc.* (2001)
doi:10.1021/ja0100120.

51.   Blaum, B. S., Neu, U., Peters, T. & Stehle, T. Spin ballet for sweet encounters: Saturation-
transfer difference NMR and X-ray crystallography complement each other in the elucidation of
protein-glycan interactions. *Acta Crystallographica Section F: Structural Biology*
Communications vol. 74 451–462 (2018).

52.   Grönberg, G., Nilsson, U., Bock, K. & Magnusson, G. Nuclear magnetic resonance and
conformational investigations of the pentasaccharide of the Forssman antigen and overlapping
di-, tri-, and tetra-saccharide sequences. *Carbohydr. Res.* **257**, 35–54 (1994).

53.   Wu, A. M. & Sugii, S. Coding and classification of D-galactose, N-acetyl-D-galactosamine, and
β-D-Galp-[1→3(4)]-β-D-GlcpNAc, specificities of applied lectins. *Carbohydr. Res.* **213**, 127–
143 (1991).

54.   Sanchez, J. F. *et al.* Biochemical and structural analysis of Helix pomatia agglutinin. A
hexameric lectin with a novel fold. **281**, 20171–20180 (2006).

55.   Springer, T., Galfrè, G., Secher, D. S. & Milstein, C. Monoclonal xenogeneic antibodies to
murine cell surface antigens: identification of novel leukocyte differentiation antigens. *Eur. J.*
Immunol. **8**, 539–551 (1978).

56.   Hauser, R., Fechner, G. & Brinkmann, B. A1 and A2 blood group substances: are there
structural differences? *Z. Rechtsmed.* **103**, 587–591 (1990).

57.   Tanaka, N. & Leduc, E. H. A study of the cellular distribution of forssman antigen in various
species. *J. Immunol.* **77**, 198–212 (1956).

58.   Koizumi, N., Hara, A., Uemura, K. & Taketomi, T. Glycosphingolipids in sheep liver, kidney,
and various blood cells. *Jpn. J. Exp. Med.* **58**, 21–31 (1988).

59.   Slomiany, B. L. & Slomiany, A. Forssman glycolipid variants of dog gastric mucosa. Structure
of a branched ceramide octasaccharide. *Eur. J. Biochem.* **83**, 105–111 (1978).

60.   Slomiany, B. L., Banas-Gruszka, Z., Zdebska, E. & Slomiany, A. Characterization of the
Forssman-active oligosaccharides from dog gastric mucus glycoprotein isolated with the use of
a monoclonal antibody. *J. Biol. Chem.* **257**, 9561–9565 (1982).

61.   Ostrander, G. K., Levery, S. B., Hakomori, S. & Holmes, E. H. Isolation and characterization of
the major acidic glycosphingolipids from the liver of the English sole (Parophrys vetulus).
Presence of a novel ganglioside with a Forssman antigen determinant. *J. Biol. Chem.* **263**,
3103–3110 (1988).

62.   Yamamoto, H. *et al.* Isolation and characterization of a novel Forssman active acidic
glycosphingolipid with branched isoglobo-, ganglio-, and neolacto-series hybrid sugar chains. *J.*
Biochem. **125**, 923–930 (1999).

63.   Savage, A. V., Donoghue, C. M., D'Arcy, S. M., Koeleman, C. A. M. & van den Eijnden, D. H.
Structure determination of five sialylated trisaccharides with core types 1, 3 or 5 isolated from
bovine submaxillary mucin. *Eur. J. Biochem.* **192**, 427–432 (1990).

64.   Hamelryck, T. W. *et al.* Carbohydrate binding, quaternary structure and a novel hydrophobic
binding site in two legume lectin oligomers from Dolichos biflorus. *J. Mol. Biol.* **286**, 1161–1177
(1999).

65.   Lescar, J. *et al.* Structural basis for recognition of breast and colon cancer epitopes Tn antigen
and Forssman disaccharide by Helix pomatia lectin. *Glycobiology* **17**, 1077–1083 (2007).

66. Nagae, M. *et al.* Structural analysis of the human galectin-9 N-terminal carbohydrate
recognition domain reveals unexpected properties that differ from the mouse orthologue. *J.*
Mol. Biol. **375**, 119–135 (2008).
67. Kita, A. *et al.* Crystal structure of octocoral lectin SLL-2 complexed with Forssman antigen
tetrasaccharide. *Glycobiology* **27**, 696–700 (2017).
68. Bulai, T., Bratosin, D., Pons, A., Montreuil, J. & Zanetta, J. P. Diversity of the human
erythrocyte membrane sialic acids in relation with blood groups. *FEBS Lett.* **534**, 185–189
(2003).
69. Cid, E., Yamamoto, M. & Yamamoto, F. Mixed-Up Sugars: Glycosyltransferase Cross-
Reactivity in Cancerous Tissues and Their Therapeutic Targeting. *ChemBioChem* **23**,
e202100460 (2022).
70. Liu, Y. *et al.* Neoglycolipid-based oligosaccharide microarray system: Preparation of ngls and
their noncovalent immobilization on nitrocellulose-coated glass slides for microarray analyses.
Methods Mol. Biol. **808**, 117–136 (2012).
71. Liu, Y. *et al.* The minimum information required for a glycomics experiment (MIRAGE) project:
improving the standards for reporting glycan microarray-based data. *Glycobiology* **27**, 280–284
(2017).
72. Ritz, C., Baty, F., Streibig, J. C. & Gerhard, D. Dose-response analysis using R. *PLoS One* **10**,
e0146021 (2015).
73. Gildersleeve, J. C., Oyelaran, O., Simpson, J. T. & Allred, B. Improved procedure for direct
coupling of carbohydrates to proteins via reductive amination. *Bioconjug. Chem.* **19**, 1485–
1490 (2008).
74. Kabsch, W. *et al.* XDS. *Acta Crystallogr. Sect. D Biol. Crystallogr.* (2010).
75. McCoy, A. J. *et al. Phaser* crystallographic software. *J. Appl. Crystallogr.* **40**, 658–674 (2007).
76. Liebschner, D. *et al.* Macromolecular structure determination using X-rays, neutrons and
electrons: Recent developments in Phenix. *Acta Crystallogr. Sect. D Struct. Biol.* (2019)
doi:10.1107/S2059798319011471.
77. Murshudov, G. N. *et al.* REFMAC5 for the refinement of macromolecular crystal structures.
Acta Crystallogr. Sect. D Biol. Crystallogr. (2011) doi:10.1107/S0907444911001314.
78. Emsley, P. & Cowtan, K. Coot: Model-building tools for molecular graphics. *Acta Crystallogr.*
Sect. D Biol. Crystallogr. (2004) doi:10.1107/S0907444904019158.

614
615
616

Acknowledgements

The glycan microarray studies were performed in the Carbohydrate Microarray Facility

at the Imperial College Glycosciences Laboratory, which is supported by Wellcome

Trust biomedical resource grants (099197/Z/12/Z, 108430/Z/15/Z, and 218304/Z/19/Z)

and in part by the March of Dimes Prematurity research center grant (22-FY18-82).

The sequence defined glycan microarrays contain many saccharides provided by

collaborators whom we thank, as well as members of the Glycosciences Laboratory

for their contribution in the establishment of the NGL-based microarray system. We

also thank the German Research Foundation for funding (FOR2327 – VIROCARB,

Project no. 269564371). Crystallographic data collection was performed at the

macromolecular beam lines X06SA and X06DA of the Swiss Light Source, access to

which was generously granted by the Paul Scherrer Institute in Villigen, Switzerland. We thank the Max-Planck Institute for Biology in Tübingen for the access to the NMR facility and Dr. Vincent Truffault for his assistance in NMR data acquisition. We also thank Prof. Barbara Mulloy (Imperial College London) for critical review of the manuscript. We thank the lab of Prof. Schulze-Osthoff at the University of Tübingen for the access to the flow cytometry appliances and especially Dr. Anja Schmitt for her help with the experiments.

Author information

Authors and affiliations

Interfaculty Institute for Biochemistry, University of Tübingen, Tübingen, Germany

Nils H. Rustmeier, Alexander Herrmann, Joshua Müller, Thilo Stehle

Glycosciences Laboratory, Faculty of Medicine, Imperial College London, London, UK

Lisete S. Silva, Antonio Di Maio, Ten Feizi, Yan Liu

Contributions

N.R. and T.S. conceptualized the study. A.H., J.M., and N.R. prepared and crystallized the protein. L.S.S and A.D.M. performed the glycan array screening experiments. A.D.M., Y.L. and T.F. performed the glycan array analyses and interpreted the glycan interaction data. N.R. performed and analyzed the NMR, SPR and flow cytometry experiments. A.H., J.M., and N.R. collected the crystallographic data and solved the structures. N.R. prepared the first draft of the manuscript and all contributed to the final revision. T.S. provided funding. All authors approved publication of the manuscript.

Ethics declarations

653   The authors declare no competing interests. Animal blood was acquired from vendors

654   guaranteeing animal wellbeing. Human erythrocytes were derived from anonymous

655   donors consenting research purposes.

Table 1 Carbohydrate antigens cited in this report.

Class	Name	Symbolic structure	Sequence
Globoseries glycosphingolipids	LacCer		Galβ1-4Glcβ-Cer
	Gb3Cer		Galα1-4Galβ1-4Glcβ-Cer
	Gb4Cer (P antigen)		GalNAcβ1-3Galα1-4Galβ1-4Glcβ-Cer
	Forssman pentaosylceramide		GalNAcα1-3GalNAcβ1-3Galα1-4Galβ1-4Glcβ-Cer
Histo blood group antigens	H antigen		Fucα1-2Galβ1-R
	A antigen		GalNAcα1-3(Fucα1-2)Galβ1-R
	B antigen		Galα1-3(Fucα1-2)Galβ1-R
O-Glycans	Tn antigen		GalNAcα-Ser/Thr
	TF antigen		Galβ1-3GalNAcα-Ser/Thr

Lac - lactose, Cer - ceramide, Gb3 - globotriose, Gb4 - globotetraose, R - glycan backbone linked to protein or lipid carriers

○ - Gal   ▢ - GalNAc   ● - Glc   ◼ - GlcNAc   ▲ - Fuc

Figure 1 Glycan microarray screening analysis of ShPyV VP1 reveals selective engagement of Forssman glycolipid. The results are the means of fluorescence intensities of duplicate spots, printed at 5 fmol per spot. The error bars represent half of the difference between the two values. In the glycan array the 672 lipid-linked probes are grouped into non-sialylated and sialylated glycans as annotated by the colored panels. The list of glycan probes, their sequences and binding scores are in Table S1.

Figure 2 Assessment of the sheep polyomavirus VP1 – Forssman pentaose

interaction in solution by STD NMR. [1]H spectra of **a** Forssman pentaose (F$_P$) and **b**

6'sialyllactose (6'SL) in their isolated forms. Proton resonances are labeled according

to the carbon positions in the monosaccharides. **c** 1H spectrum of the test sample

comprising 20 μM ShPyV VP1, 1 mM F$_P$, and 1 mM 6'SL collected under protein

suppression. **d** The STD NMR spectrum recorded from the test sample.

Figure 3 Flow cytometry and surface plasmon resonance analyses of Forssman antigen engagement. a Determination of the binding kinetics towards a Forssman-decorated surface using ShPyV VP1 and HPA. Response curves are shown as triplicates for each protein concentration. Kinetic parameters and the dissociation constant are provided in the table below. **b** Absolute cell attachment of ShPyV VP1 and *Helix pomatia* agglutinin (HPA) to Forssman antigen-positive red blood cells (RBCs) from sheep (Sh) and human erythrocytes of the blood groups A1, A2, B, and O (left). Effects of Forssman and blood group A antigen trioses on ShPyV VP1 and HPA attachment to sheep RBCs. Results are shown as concatenated and normalized histograms (right). **c** EC50 determination using ShPyV VP1, HPA, and anti-Forssman IgM M1/87 to sheep blood cells. Significance is indicated with *** for $p < 0.001$.

Figure 4 Forssman pentaose binds to the external surface of ShPyV VP1 with numerous contacts. **a** The crystal structure of a single VP1 pentamer is shown with five molecules of Forssman pentaose (F_P) associated to its outer surface. The bias-reduced Fo-Fc omit electron density maps of F_P are displayed as blue meshes around the oligosaccharides, with a contour level of 2.5 σ and a radius of 1.6 Å. The individual F_P binding sites are each constituted by adjoining protein chains, highlighted by the orange coloring of one VP1 monomer. **b** Magnified view of a bound F_P molecule (above) with its symbol structure (below). The glucose of F_P, which is not visible in the crystal structure, is grayed out in the symbol representation. **c** Molecular interactions between VP1 and F_P are displayed with the hydrogen bonds indicated as dashed lines.

698 VP1 coloring corresponds to subfigure **a**. Water molecules are shown as spheres.

699 Throughout the figure, oxygen and nitrogen atoms are colored in red and blue,

700 respectively.

Table 2 Intermolecular contacts between ShPyV VP1 and Forssman pentaose.

Monosaccharide		VP1 contacts	
		Van der Waals	
αGalNAc	Methyl	F53, F56, F66cw, A122, F264	
		Hydrogen-bonds	Distance (Å)
αGalNAc	O3	*D64cw*	2.74
	Amide N	Q266	2.70
	Acetyl O	*F66cw*	2.90
βGalNAc	O4	H₂O-H₂O-*S57*	2.95-2.90-2.63
	O4	N59	3.06
	O4	Q266	2.55
	O6	Q47	2.95
	Acetyl O	Q266	2.96
	Acetyl O	N268	2.78
αGal	O2	K270	3.31
	O4	D58	2.67
	O4	N59	3.45
	O4	H₂O-*N59*	2.61-2.83
	O6	D58	2.54

Residues using their backbone atoms to establish contacts are printed in italic. Distance values are derived from a single binding site and may alter between chains.