Characterization of Target Gene Regulation by the Two Epstein-Barr Virus Oncogene LMP1 Domains Essential for B-cell Transformation

Bidisha Mitra[1-3], Nina Rose Beri[1-3], Rui Guo[1-3], Eric M. Burton[1-3], Laura A. Murray-Nerger[1-3] and Benjamin E. Gewurz[1-3*]

[1] Division of Infectious Diseases, Department of Medicine, Brigham and Women's Hospital, 181 Longwood Avenue, Boston MA 02115, USA
[2] Center for Integrated Solutions in Infectious Disease, Broad Institute of Harvard and MIT, Cambridge, MA 02142, USA
[3] Department of Microbiology, Harvard Medical School, Boston, Massachusetts 02115, USA
*Correspondence: bgewurz@bwh.harvard.edu

Abstract:

The Epstein-Barr virus (EBV) oncogene latent membrane protein 1 (LMP1) mimics CD40 signaling and is expressed by multiple malignancies. Two LMP1 C-terminal cytoplasmic tail regions, termed transformation essential sites (TES) 1 and 2, are critical for EBV transformation of B lymphocytes into immortalized lymphoblastoid cell lines (LCL). However, TES1 versus TES2 B-cell target genes have remained incompletely characterized, and whether both are required for LCL survival has remained unknown. To define LCL LMP1 target genes, we profiled transcriptome-wide effects of acute LMP1 CRISPR knockout (KO) prior to cell death. To then characterize specific LCL TES1 and TES2 roles, we conditionally expressed wildtype, TES1 null, TES2 null or double TES1/TES2 null LMP1 alleles upon endogenous LMP1 KO. Unexpectedly, TES1 but not TES2 signaling was critical for LCL survival. The LCL dependency factor cFLIP, which plays obligatory roles in blockade of LCL apoptosis, was highly downmodulated by loss of TES1 signaling. To further characterize TES1 vs TES2 roles, we conditionally expressed wildtype, TES1 and/or TES2 null LMP1 alleles in two Burkitt models. Systematic RNAseq analyses revealed gene clusters that responded more strongly to TES1 versus TES2, that respond strongly to both or that are oppositely regulated. Robust TES1 effects on cFLIP induction were again noted. TES1 and 2 effects on expression of additional LCL dependency factors, including BATF and IRF4, and on EBV super-enhancers were identified. Collectively, these studies suggest a model by which LMP1 TES1 and TES2 jointly remodel the B-cell transcriptome and highlight TES1 as a key therapeutic target.

Key Words: gammaherpesvirus; lymphoproliferative disease; tumor virus; B-cell oncogenesis; interferon regulatory factor; BATF; NF-kB; super-enhancer; dependency factor, apoptosis.

Short title: The LMP1 TES1 and TES2 host target gene landscape

Importance:

Epstein-Barr virus (EBV) causes multiple human cancers, including B-cell lymphomas. In cell culture, EBV converts healthy human B-cells into immortalized ones that grow continuously, which model post-transplant lymphomas. Constitutive signaling from two cytoplasmic tail domains of the EBV oncogene Latent Membrane Protein 1 (LMP1) is required for this transformation, yet there has not been systematic analysis of their host gene targets. We identified that only signaling from the membrane proximal domain is required for survival of these EBV-immortalized cells and that its loss triggers apoptosis. We identified key LMP1 target genes, whose abundance changed significantly with loss of LMP1 signals, or that were instead upregulated in response to switching on signaling by one or both LMP1 domains in an EBV-uninfected human B-cell model. These included major anti-apoptotic factors necessary for EBV-infected B-cell survival. Bioinformatics analyses identified clusters of B-cell genes that respond differently to signaling by either or both domains.

Introduction:

Epstein-Barr virus (EBV) is a gamma-herpesvirus that persistently infects most adults worldwide. EBV causes 200,000 cancers per year, including Burkitt lymphoma, Hodgkin lymphoma, post-transplant lymphoproliferative disease (PTLD), and HIV/AIDS associated lymphomas. EBV also causes a range of epithelial cell tumors, including gastric and nasopharyngeal carcinomas, as well as T and NK cell lymphomas(1). The key EBV oncogene latent membrane protein 1 (LMP1) is expressed in most of these tumors, where it drives growth and survival pathway signaling.

To colonize the B-cell compartment and establish lifelong infection, EBV uses a series of viral latency genome programs, in which different combinations of latency genes are expressed. These include six Epstein-Barr nuclear antigens (EBNA) and the membrane oncoproteins LMP1, LMP2A and LMP2B. LMP1 mimics aspects of signaling by the B-cell co-receptor CD40 (2-5), whereas LMP2A rewires surface B-cell immunoglobulin receptor signaling (6). All nine latency oncoproteins are expressed in the EBV B-cell transforming latency III program, which are expressed in immunoblastic lymphomas of immunosuppressed hosts. These include PTLD and primary central nervous system lymphoma. The latency II program is observed in EBV+ Hodgkin lymphoma, where the Reed-Sternberg tumor cells express EBNA1, LMP1 and LMP2A. Latency II is also frequently observed in T and NK cell lymphomas and in nasopharyngeal carcinoma. Host genome NF-κB activating mutations are frequently observed in EBV-negative Hodgkin lymphoma, but to a much lesser extent in EBV+ tumors, underscoring LMP1's key role in activating growth and survival signaling (7).

LMP1 localizes to lipid rafts, where it signals constitutively in a ligand-independent fashion to activate NF-κB, MAP kinase, STAT3, PI3K, interferon and P62 pathways. LMP1 is comprised of a short N-terminal cytoplasmic tail, six transmembrane (TM) domains and a 200 residue C-terminal cytoplasmic tail (2-5). LMP1 TM domains drive homotypic aggregation, lipid raft association and constitutive signaling (8, 9). The LMP1 C-terminal tail functionally mimics signaling from activated CD40 receptors, to the point that the CD40 tail can essentially be replaced by that of LMP1 in transgenic mice studies. However, while CD40/LMP1 knockin mice had relatively normal B-cell development and evidence of intact CD40 function, including germinal center formation and class switch recombination, T-cell independent B-cell activation was also observed (10). These experiments suggest that the LMP1 C-terminal tail mimics CD40 signaling, but has also evolved additional functions.

Reverse genetic studies identified two LMP1 C-terminal cytoplasmic tail domains that are critical for EBV-mediated conversion of primary human B-cells into immortalized, continuously growing lymphoblastoid cell lines (LCL). Transformation effector site 1 (TES1), also called C-terminal activating region 1 (CTAR1), spans LMP1 residues 186-231. TES1/CTAR1 contains a PXQXT motif that engages tumor necrosis factor receptor associated factors (TRAFs). TES1 activates canonical NF-κB, non-canonical NF-κB, MAP kinase, PI3K and STAT3 pathways (3, 4, 11-16). TES2, which spans residues 351-386 and is also referred to as CTAR2, activates canonical NF-κB, MAPK, IRF7 and P62 pathways (3-5, 16-20). TRAF6 is critical for LMP1 TES2-driven canonical NF-κB, MAPK and p62 pathway activation (21-26). Canonical NF-κB signaling is critical for TES2/CTAR2 driven target gene regulation in a 293 cell conditional expression model (27). Signaling from a third LMP1 C-terminal tail region, CTAR3, activates JAK/STAT and SUMOylation pathways (28-30) potentially important *in vivo* but that

are not essential for EBV-driven B-cell transformation (31). ChIP-seq analyses demonstrated a
complex NF-κB binding landscape in LCLs, in which constitutive LMP1 signaling stimulates
different combinations of the NF-κB transcription factors RelA, RelB, cRel, p50 and p52 to bind
B-cell enhancers and promoters (32).

LMP1 is the only EBV oncogene that can independently transform rodent fibroblasts,
driving anchorage independent growth and loss of contact inhibition (33-35). Notably, CTAR1
signaling is sufficient for LMP1-mediated fibroblast transformation, whereas CTAR2 was
dispensable(36). LMP1 expression drives aberrant B-cell growth in transgenic B-cell models,
particularly in combination with LMP2 upon disruption of cell mediated immunity (12, 37-40).
While not critical for the first 8 days of EBV-driven B-cell outgrowth (41), LMP1 is critical for
EBV-mediated conversion of primary human B-cells into immortalized LCLs (42, 43). A
longstanding question has remained why TES1 and TES2 are each essential for EBV-mediated
LCL establishment. Whether either or both are required for LCL survival is also unknown.
Experiments using the EBV second site mutagenesis method (44) demonstrated that TES1 is
critical for initiation of EBV-infected lymphoblastoid cell outgrowth (45). By contrast, TES2 is
critical for long-term LCL growth, although TES2 null EBV infected lymphoblastoid cells could be
propagated on epithelial feeders (45, 46). However, it remains incompletely understood the
extent to which TES1 and TES2 play overlapping versus non-redundant roles.

While LMP1 B-cell target genes have been analyzed on small scales through qPCR and
limited microarray analysis, unbiased genome-wide approaches have yet to be applied. Little is
presently known about TES1 and TES2 shared versus non-redundant roles in transformed B-
cells. To gain insights into LMP1 targets in the latency III LCL context, we therefore profiled
transcriptome-wide changes in response to acute CRISPR LMP1 knockout (KO). These studies,
performed at an early timepoint prior to apoptosis, identified that LMP1 strongly controls the LCL
transcriptome, with expression levels of nearly 3400 host genes significantly altered by LMP1
KO. To then characterize specific LCL TES1 and TES2 roles, we conditionally expressed
wildtype, TES1 null or TES2 null LMP1 rescue cDNAs at physiological levels upon endogenous
LMP1 KO. This approach unexpectedly highlighted that signaling by TES1, but not TES2, is
critical for LCL growth and survival. Loss of TES1 but not TES2 signaling rapidly triggered
apoptosis, and strongly impaired expression of the LCL dependency factor cFLIP, which is
required to block TNFα-driven apoptosis(25). Transcriptomic profiling further highlighted six
clusters of LCL gene responses wildtype, TES1 or TES2 mutant LMP1, newly identifying
independent, additive, or antagonistic roles in LCL target gene regulation. As multiple latency III
genes often target the same host cell targets, we also constructed Burkitt B-cell models with
conditional expression of wildtype, TES1 and/or TES2 null LMP1. These studies extended the
LCL findings by further identifying shared versus distinct LMP1 roles in B-cell target gene
regulation. Collectively, these studies highlight a complex landscape of TES1 and TES2 target
gene regulation, in which each controls expression levels of large numbers of B-cell targets.

Results:

CRISPR analysis of LCL LMP1 Target Genes

To characterize LMP1 target genes in the latency III context, we used CRISPR to knockout (KO) LMP1 in the well-characterized LCL GM12878, a Tier 1 Encode project cell line that we have used extensively for CRISPR analyses, and which we confirmed to have the latency III program (25, 47). GM12878 with stable Cas9 expression were transduced with lentivirus expressing a control single guide RNA (sgRNA) targeting a human genome intergenic region or LMP1. Immunoblot confirmed efficient LMP1 depletion by 48 hours post-puromycin selection of transduced LCLs (**Fig. 1A**). CRISPR LMP1 editing rapidly downmodulated the LMP1/NF-κB target genes TRAF1 and IRF4 and decreased non-canonical pathway processing of the p100 NF-κB precursor into the active p52 transcription factor subunit, suggesting successful on-target effects of LMP1 knockout (KO) (**Fig. 1A**). At this early timepoint post-CRISPR editing, LCLs remained viable (**Fig. S1A**). However, LMP1 KO triggers LCL growth arrest and cell death shortly thereafter. We therefore used this early 2-day post-puromycin selection timepoint to perform systematic RNAseq analyses of control vs LMP1 KO LCLs. At a multiple hypothesis testing adjusted p-value <0.05 and fold change of >2 cutoff, acute LMP1 KO significantly altered the levels of around 3400 host genes.

Genes differentially expressed in LMP1 KO vs control cells could be broadly characterized into two k-means clusters, in which LMP1 KO either downregulated 1,476 or upregulated 1926 host genes (**Fig. 1B, Table S7**). Kyoto Encyclopedia of Genetic Elements (KEGG) pathway Enrichr analysis (48) identified that cytokine receptor signaling, NF-κB signaling and tumor necrosis factor (TNF) signaling as enriched amongst genes rapidly downmodulated by LMP1 KO (**Fig. 1C**). As examples of cluster 1 NF-κB target genes, interferon regulatory factor 4 (IRF4) and CFLAR, which encodes c-FLIP, were strongly downmodulated by LMP1 KO (**Fig. 1D**). By contrast, KEGG highlighted that autophagy, p53 signaling and protein-processing in the endoplasmic reticulum as enriched amongst cluster 2 genes (**Fig. 1C**). As examples from these enriched pathways, LMP1 KO highly induced expression of the autophagy suppressor DEPP1 and the p53 target and tumor suppressor cyclin dependent kinase inhibitor 2A (CDKN2A) (**Fig. 1D-F**). Quantitative real-time PCR (qRT-PCR) validated RNAseq results for the CCL22, EBI3 and IRF4 mRNAs, each of which were significantly downmodulated by LMP1 KO (**Fig. S1B-D**).

We next integrated our RNAseq dataset with published CRISPR analysis of host dependency factors essential for EBV+ LCL, but not Burkitt B-cell proliferation (25), to gain insights into key LMP1 roles in LCL growth and survival. This analysis identified that mRNA abundances of 37 of the 87 CRISPR-defined LCL selective dependency factors significantly changed upon LMP1 KO, suggesting multiple LMP1 roles in support of LCL survival (**Fig. 1G and S1E**). Of these, it is notable that multiple key suppressors of LCL intrinsic and extrinsic apoptotic pathways were rapidly lost upon LCL LMP1 KO. For instance, our published CRISPR analyses highlighted non-redundant roles for the transcription factors IRF4 and BATF in blockade of the intrinsic apoptosis pathway and for CFLAR-encoded cFLIP in extrinsic apoptosis pathway inhibition (25), each of whose mRNAs rapidly decreased upon LCL LMP1 KO. Likewise, LMP1 KO strongly downmodulated expression of MDM2, an LCL-selective dependency factor (25) that targets p53 for proteasomal degradation and that prevents LCL p53-dependent apoptosis(49). Furthermore, NF-κB blockade triggers LCL apoptosis(50) and the LCL dependency factor NFKB2, which encodes the NF-κB transcription factor subunit p52, was

also highly downmodulated by LMP1 KO (**Fig. 1F-G**). STRING network analysis also
underscored that each of these assemble into a network with 23 other LMP1-regulated LCL
dependency factors (**Fig. 1H**).

Since LMP1 is highly expressed in Hodgkin lymphoma Reed-Sternberg tumor cells, we next
analyzed effects of LMP1 KO on Hodgkin lymphoma KEGG pathway genes (**Fig. S1F**).
Interestingly, LMP1 KO strongly downmodulated expression of the T-cell tropic chemokines
CCL22 and CCL17, consistent with several prior reports linking LMP1 to their expression (51,
52). These findings raise the possibility that LMP1-driven chemokine expression may contribute
to the striking enrichment of T-cells characteristic of the Hodgkin Reed-Sternberg
microenvironment. However, volcano plot analysis also highlighted that LMP1 KO increased
expression of CD274, which encodes the checkpoint inhibitor PD-L1, further implicating LMP1 in
T-cell regulation.

Given widespread effects of LMP1 KO on LCL host gene expression, we next characterized
effects of LMP1 KO on viral latency III genes. Mapping of RNAseq reads onto the GM12878
EBV transcriptome identified that LMP1 depletion significantly downmodulated mRNAs
encoding EBNA3A, 3C and EBNA-LP, though interestingly not those encoding EBNA2 or
EBNA3B (**Fig. 1I**). While it has been reported that LMP1 regulates its own mRNA expression
(53, 54), we did not observe changes in LMP1 mRNA abundance upon LMP1 CRISPR KO. We
note that CRISPR editing often results in insertions or deletions, causing functional protein
knockout without necessarily changing mRNA levels of the edited gene. However, it is plausible
a compensatory response to LMP1 knockout occurred on the mRNA level at this early
timepoint, potentially balancing loss of NF-κB induced LMP1. Taken together, our RNAseq
analyses raise the possibility that secondary effects of LMP1 KO on Epstein-Barr nuclear
antigens may also contribute to changes in the host transcriptome and cell death upon LMP1
KO.

TES1 but not TES2 signaling is critical for LCL survival

While TES1 and TES2 signaling are each critical for B-cell transformation, it has remained
unknown whether either or both are necessary for proliferation of fully transformed LCLs.
Likewise, knowledge has remained incomplete about shared versus non-redundant TES1 and
TES2 roles in LCL host gene regulation. To gain insights into these key questions, we
engineered Cas9+ GM12878 LCLs with conditional expression of wildtype (WT) LMP1, or with
well characterized point mutants that abrogate signaling from the TES1 TRAF binding domain
(TES1m, residues 204PQQAT208 -> **A**Q**A**AT), from TES2 (TES2m, 384YYD386 -> ID) (46, 55,
56) (**Fig. 2A**). A silent mutation in the CRISPR protospacer adjacent motif (PAM) was used to
abrogate CRISPR editing of these LMP1 rescue cDNA constructs. For cross-comparison, we
also established conditional TES1/TES2 double mutant (DM) cell lines with both mutations, to
profile responses to other LMP1 regions, potentially including CTAR3 or unfolded protein
responses induced by LMP1 induction (57, 58) (**Fig. 2A**). LCLs were then transduced with
lentivirus expressing a control sgRNA targeting a human intergenic region or LMP1. Conditional
LMP1 expression was then induced by addition of 400 ng/ml doxycycline, such that the rescue
cDNA was induced as endogenous EBV-encoded LMP1 was depleted. We confirmed similar
levels of LMP1 expression across this series and achieved similar LMP1 levels as in unedited
GM12878 LCLs (**Fig. 2B**). Importantly, we validated that WT LMP1 rescued physiological levels

of LMP1 target TRAF1 expression and p100/p52 processing (**Fig. 2B**). TES1 is responsible for
the majority of LMP1-mediated TRAF1 induction and p100/p52 processing(55, 59) and as
expected, conditional TES1m and DM expression failed to rescue physiological levels of TRAF1
or p100/p52 processing in GM12878 with endogenous LMP1 KO. By contrast, TES2m induced
levels of TRAF1 and p100/p52 processing in LMP1 KO levels approaching those in unedited
GM12878 (**Fig. 2B**), validating our LCL LMP1 KO/rescue system.

We next tested the effects of LMP1 WT, TES1m, TES2m or DM rescue cDNA expression on
GM12878 proliferation. Whereas signaling from both TES1 and TES2 are required for EBV-
driven primary human B-cell growth transformation, we unexpectedly found that LCLs require
signaling only from TES1 for growth and survival: LMP1 KO LCLs with WT vs TES2m rescue
cDNA proliferated indistinguishably. By contrast, LCLs with TES1m or DM rescue cDNA
expression failed to proliferate (**Fig. 2C**). To characterize this unexpected result further, we next
measured effects of endogenous LMP1 KO and rescue LMP1 cDNA expression on LCL
survival. Consistent with our growth curve analysis, LMP1 KO LCLs with TES1m or DM rescue
cDNA exhibited widespread cell death, as judged by uptake of the vital dye 7aminoactinomycin
D (7-AAD) by FACS analysis (**Fig. 2D-E**). Consistent with apoptosis as the cell death pathway
triggered by loss of TES1 signaling in LCLs, levels of Annexin V and executioner caspase 3/7
activity were significantly higher in LMP1 KO GM12878 with TES1m or DM than with WT or
TES2m rescue cDNA expression (**Fig. 2F-G and S2**). Taken together, these data newly
suggest that TES1 signaling is necessary for LCL growth and survival in a manner that is not
redundant with TES2, and that cannot be rescued by TES2 signaling alone.

Identification of TES1 versus TES2 roles in LCL gene regulation

To gain insights into overlapping versus non-redundant TES1 and TES2 LCL roles, we
performed biological triplicate RNAseq analyses to cross-compare GM12878 transcriptomes at
day 6 post endogenous LMP1 KO and with doxycycline-induced WT, TES1m or TES2m rescue
cDNA expression. We selected this early timepoint as it is just prior to the divergence of the
growth curves (**Fig. 2C, Table S8-9**) and the onset of apoptosis. K-means analysis identified six
clusters in which host gene expression differed between LMP1 KO LCLs with WT, TES1m or
TES2m cDNA rescue. KEGG analysis highlighted pathways most highly enriched in each
cluster (**Fig. 3A**). Notably, apoptosis pathway genes were the most highly enriched in cluster 4
genes, which were expressed at lower levels in cells with TES1m than with WT or TES2m
rescue cDNA expression, suggesting that TES1 signaling may induce their expression.
Apoptosis genes were also enriched amongst cluster 5 genes, where levels were lower in cells
with TES2m expression, suggesting that TES2 signaling may induce their expression (**Fig. 3A**).

Given this apoptosis signal, we next analyzed the KEGG apoptosis gene set responses to
WT, TES1m or TES2m cDNA rescue (**Fig. 3B**). Interestingly, a cluster of genes were more
highly expressed in cells with WT and TES2m than with TES1m expression, including the anti-
apoptotic genes CFLAR, BCL2 and BIRC3, which encodes the cIAP2 ubiquitin ligase, which
counteracts TNF-driven cell death. BCL2 and BIRC3 were not defined as LCL dependency
factors by genome-wide CRISPR analysis, whereas CLFAR was (25). Intriguingly, while a small
number of the 87 LCL dependency factors (25) were more highly downmodulated in LMP1 KO
LCLs with TES1 rescue than with TES2 rescue, CFLAR was the LCL dependency factor most
highly depleted from with TES1m rescue (nearly 8-fold) (**Fig. 3C, S3A**). By contrast, CFLAR

was only mildly depleted (<0.5 fold) in cells with TES2m rescue (**Fig. 3C**). We validated that
CFLAR-encoded c-FLIP was highly downmodulated on the protein level in LMP1 KO LCLs with
TES1m or DM LMP1 rescue, but not in LCLs with WT or TES2m rescue at an early timepoint
prior to cell death (**Fig. 3D**). The LCL dependency factors NFKB2 and CCND2 were also more
highly downmodulated in LMP1 KO cells with TES1 than TES2 cDNA rescue but not to the
same extent as CFLAR (**Fig. 3C**), suggesting that their loss may not be responsible for
apoptosis in the absence of TES1 signaling. Collectively, these analysis underscore distinct
TES1 versus TES2 signaling roles in control of LCL apoptosis pathway gene expression.

To gain further insights into potential TES1 vs TES2 roles in regulation of genes with
relevance to Hodgkin Reed-Sternberg cells, we analyzed KEGG Hodgkin lymphoma pathway
gene expression in LMP1 KO GM12878 rescued with WT, TES1m or TES2m LMP1. This
analysis highlighted that many KEGG Hodgkin lymphoma pathway genes are jointly induced by
TES1 and TES2 signaling in LCLs, including CCL22, BCL3, cRel, IRF4, STAT3, STAT6 and
CD70, each of which have prominent roles in Hodgkin lymphoma pathogenesis (**Fig. S3B**). We
also confirmed the transcript levels of CCL22, EBI3 and IRF4 through qRT-PCR and the results
were concordant with our RNA-seq analysis where TES1 and TES2 are jointly responsible in
the induction of the expression of these genes (**Fig. S3 C-D**). Interestingly, the Hodgkin
lymphoma therapeutic target CD27 was expressed at lower level with TES1m rescue but more
highly expressed in cells with TES2m rescue, suggesting that TES1 and TES2 signaling may
jointly balance its expression.

On the transcriptome-wide level, multiple well characterized LMP1 target genes were more
highly expressed in LCLs rescued with WT LMP1 than with TES1m cDNA, establishing these as
key TES1 LCL target genes. These included CD40, TRAF1, EBI3 and ICAM1 (**Fig. 4A**), which
was previously established as a TES1 target genes in studies of cell lines with LMP1 over-
expression, including BL-41 Burkitt and BJAB diffuse large B-cell lymphoma models(55).
Notably, this approach also newly suggests a large number of B-cell targets whose upregulation
or downregulation is dependent on TES1 signaling. These include CFLAR and TLR6 (which
encodes Toll-like receptor 6), which were significantly more highly expressed in LMP1 KO LCLs
with WT LMP1 cDNA rescue. By contrast, the mRNA encoding the histone loader DAXX, which
can serve as an epigenetic suppressor of EBV gene expression (60, 61) and DNA damage
pathway TP53 (which encodes p53), were expressed at considerably higher in LMP1 KO LCLs
with TES1m than WT LMP1 rescue cDNA (**Fig. 4A**). This result suggests that TES1 may
repress their expression. Enrichr analysis of genes more highly expressed with WT LMP1
rescue highlighted TNF and NF-κB signaling as enriched KEGG pathways, whereas p53
signaling and apoptosis were amongst the pathways most highly enriched in genes more highly
expressed with TES1m rescue (**Fig. 4B**).

Volcano plot and KEGG pathway analysis highlighted LCL genes differentially expressed in
LMP1 KO LCLs with WT vs TES2m rescue (**Fig. 4C-D**). KEGG pathways enriched amongst
genes more highly expressed with WT LMP1 rescue again included antigen presentation and
cytokine/receptor interaction, but also included systemic lupus erythematosus (SLE). Pathways
enriched amongst genes more highly expressed with TES2m rescue instead included cyclic
GMP protein kinase G (cGMP-PKG) signaling and phosphatidylinositol signaling (**Fig. 4D**).
Notably, TP53 (which encodes p53) was also more highly expressed in LMP1 KO LCLs with
TES2m than WT LMP1 rescue, suggesting that both TES1 and 2 signaling regulate its

expression. By comparison, CFLAR expression was similar in LMP1 KO LCLs with WT and
TES2m rescue, further establishing it as a TES1 target in LCLs (**Fig. 4C**).

Direct cross-comparison of genes in LCLs with TES1m vs TES2m rescue further identified
roles of TES1 versus TES2 signaling on LCL target gene expression. The oncogenic kinase
CLK2, which has roles in splicing regulation, was the host gene most highly expressed in LMP1
KO with TES2m rescue than in cells with TES1m rescue, newly indicating that it is strongly
induced by TES1 or strongly inhibited by TES1 (**Fig. 4E**). Enrichr analysis indicated that multiple
KEGG metabolism pathways were the most highly enriched in cells with TES2m rescue,
including fructose/mannose metabolism, HIF1 signaling and AMPK signaling (**Fig. 4E-F**). In
support, the glycolytic enzyme PFKFB4 and the kinase PDK1, which regulates flux of glycolytic
products to mitochondrial metabolism pathways at the level of pyruvate, were more highly
expressed with TES2m rescue, suggesting that they are either driven by TES1 or repressed by
TES2 signaling (**Fig. 4F**). Cell cycle regulation was the KEGG pathway most enriched amongst
genes more highly expressed with TES1m rescue. The cyclin dependent kinase (CDK)
substrate and cytokinesis regulator PRC1, as well as the CDK1 kinase and mitosis regulator
PKMYT1, were amongst the genes most highly differentially expressed in TES1m rescue (Fig.
4E-F), suggesting that TES2 drives or that TES1 instead inhibits their expression.

B-cell genes induced by conditional expression LMP1 in EBV-negative Burkitt cells

As a complementary approach to our loss-of-function CRISPR KO LCL analyses, we next
profiled B-cell responses to conditional LMP1 expression. A goal of this approach was to identify
LMP1-specific effects on host gene expression, since LMP1 KO significantly altered expression
of several EBV latency III genes. Furthermore, EBNA and LMP latency III oncoproteins often
jointly target host genes. Therefore, to study LMP1-specific effects in isolation of other latency III
genes, we engineered EBV-negative Akata and BL-41 Burkitt B-cell lines with doxycycline-
inducible WT, TES1m, TES2m or DM LMP1 alleles. The Akata cell line was originally
established from a human EBV+ Burkitt tumor (62), but an EBV-negative subclone that
spontaneously lost the viral genome was isolated shortly thereafter (63), which we used for
these studies. Similarly, the BL-41 cell line was established from an EBV-negative human
Burkitt lymphoma tumor (64). BL-41 were used for early microarray analysis of latency III or
LMP1 effects on a subset of human genes (50). We validated that WT and point mutant LMP1
were expressed to similar extents across the panel. As expected, TES1m and DM exhibited
impaired non-canonical NF-κB pathway activation, as judged by p100:p52 processing (**Fig S4A-**
B). LMP1 signaling was also validated by FACS analysis of ICAM-1 and Fas upregulation.
Consistent with a published study (55), TES1 signaling more strongly induced ICAM-1 and Fas
in both Burkitt cell lines, even though BL-41 had somewhat higher basal NF-κB activity than
Akata, as judged by Fas and ICAM-1 levels in uninduced cells (**Fig. S4C-J**).

We then profiled effects of conditional WT, TES1m, TES2m or DM expression for 24 hours
on the Akata transcriptome using biological triplicate RNAseq datasets. K-means heatmap
analysis with n=6 clusters revealed strikingly distinct patterns of host gene responses to WT,
TES1m, TES2m and DM LMP1 signaling (**Fig. 5A, Table S1-3**). Cluster 1 genes were highly
upregulated by WT LMP1, to a lesser extent by TES2m (in which only TES1 signals), and more
modestly by TES1m (in which only TES2 signals). This result suggests that TES1 signaling
contributes more strongly than TES2 to their expression. Notably, CFLAR was a cluster I gene

target, consistent with our finding that TES1 drives CFLAR expression in LCLs, as was the
interferon stimulated gene IFIT1 (**Fig. 5B**). KEGG pathways enriched amongst Cluster 1 genes
included TLR signaling, chemokine signaling, IFN signaling and NLR signaling (**Fig. 5C**).
Cluster 1 also contained well described LMP1 target genes, including TRAF1, which we
validated by immunoblot (**Fig. S4A**), consistent with a prior study(55). Notably, MAP3K7, which
encodes the kinase TAK1 is also a Cluster 1 gene. Since TAK1 is critical for TES2/canonical
NF-κB and MAP kinase signaling (26), this result suggests an important mechanism of cross-
talk between TES1 and TES2. Likewise, the Cluster I gene IRF7 binds to and is activated by
TES2 (65-68), again suggesting cross-talk between LMP1 pathways. STAT1 and STAT3 are
also Cluster 1 genes, raising the question of whether these STATs may drive interferon
stimulated gene induction downstream of LMP1.

Cluster 2 genes were induced by WT, TES1m or TES2m LMP1 to a similar extent (**Fig. 5C**),
suggesting that they redundantly respond to TES1 or TES2 signaling. GO analysis highlighted
enrichment of NF-κB signaling in this cluster, and which included mRNAs encoding four NF-κB
transcription factor subunits, as well as the NF-κB induced inhibitors IκBα, IκBζ and IκBε.
mRNA fold changes for *NFBK2*, which encodes non-canonical pathway NF-κB p52 transcription
factor are shown in **Fig. 5C** and are consistent with our LCL rescue analysis, which identified
similarly important roles for both TES1 and TES2 in support of NFKB2 expression (**Fig. 3C**).
Consistent with a prior study (55), Fas and ICAM-1 are Cluster 2 genes similarly induced on the
mRNA level by TES1m and TES2m, though interestingly, plasma membrane ICAM-1 levels
were lower in cells expressing TES1m (**Fig. S4C-F**). This result raises the possibility that TES1
signaling may play a role in ICAM-1 post-transcriptional regulation and/or trafficking.

Cluster 3 genes were expressed at lower levels in cells expressing TES1m, even as
compared with cells expressing LMP1 DM, suggesting that unopposed TES2 signaling results in
their downregulation (**Fig. 5B and S4A**). Cluster 3 genes were enriched for multiple KEGG
metabolism pathways, including oxidative phosphorylation (**Fig. S5A**). Cluster 4 contained a
smaller subset of host genes, downregulated by TES1 signaling, including in the WT LMP1
context. This gene set was enriched for SNARE interactions in vesicular transport and
sphingolipid metabolism (**Fig. S5B**). Cluster 5 mRNAs were instead upregulated by unopposed
TES2 signaling (**Fig. 5B, S5C**) and enriched for the KEGG ubiquitin mediated proteolysis
pathway. Finally, Cluster 6 genes were repressed by TES1 and TES2 signaling in an additive
manner (**Fig. 5D**). This gene set was enriched for mismatch and base excision repair,
nucleotide metabolism, cystine and methionine metabolism. TES1 and TES2 signaling may
additively recruit the same repressors or may instead recruit co-repressors to these sites. We
also confirmed the concordance of our RNAseq through qRT-PCR of the transcript levels of
CCL22, EBI3 and IRF4 in WT, TES1m, TES2m or DM induced expression (**Fig. S5 D-F**). As
expected TES1m and TES2m significantly abrogate the expression of the genes while we
barely detect anything for the DM.

We next used RNAseq to profile BL-41 Burkitt cells. RNAseq was performed at 24 hours
post-expression of WT, TES1m, TES2m or DM LMP1. K-means analysis with n=6 clusters
again revealed categories of genes that respond differently to LMP1 alleles (**Fig. S6A, Table
S4-6**). As observed in Akata, Cluster 1 genes were most highly induced by WT, and to a lesser
extent by TES1m or TES2m, suggesting that TES1/2 additively or synergistically induce their
expression. Cluster 1 genes contained multiple pro-inflammatory factors, including chemokines
and the interferon pathway transcription factors STAT1, IRF4, IRF5 and IRF9 (**Fig. S6A-B**). As

observed in Akata, cluster 2 genes were induced more strongly by TES2m than by TES1m, and
to a somewhat higher level by WT LMP1, suggesting that these are predominantly TES1 target
genes (**Fig. S6C**). Consistent with our LCL and Akata cell analyses, CFLAR was a Cluster 2
gene more highly induced by LMP1 alleles with TES1 signaling, further underscoring it as a key
TES1 target gene (**Fig. S6C**). By contrast, BL-41 Cluster 4 genes were instead suppressed by
unopposed TES1 signaling even relative to levels observed in cells with LMP1 DM expression,
suggesting that TES2 may block TES1 repressive effects on these host targets (**Fig. S6D**).
Cluster 6 genes were enriched for the antigen presentation pathway and were most highly
induced by TES2m, suggesting positive TES1 and potentially also negative TES2 roles in their
induction (**Fig. S6E**). While concordant to a large degree, we speculate that observed
differences between LMP1 effects on Akata versus BL-41 host gene expression may likely
reflect the somewhat higher basal NF-κB levels observed in BL-41, and perhaps also
differences in driver mutations pathways frequently found in EBV+ vs EBV- Burkitt lymphomas
(69, 70). Nonetheless, both models highlight distinct clusters of host B-cell target genes that
differ in responses to TES1, TES2 or combined TES1/2 signaling.

LMP1 WT, TES1, TES2 and DM target genes

We next cross-compared the most highly differentially-expressed genes across the LMP1
conditions. At a fold-change >2 and adjusted p value <0.05 cutoff, WT LMP1 highly upregulated
1021 and downregulated 518 Akata genes, respectively. The most highly upregulated genes
included multiple interferon stimulated genes, TRAF1, FAS and CFLAR (**Fig. 6A**). Interestingly,
WT LMP1 decreased expression of the recombinase RAG1 and RAG2 mRNAs, as well as
MME, which encodes CD10, a plasma membrane protein that we and others have found is
downmodulated by EBV latency III (71-73). Enrichr analysis identified that EBV infection was
the KEGG pathway most highly upregulated by WT LMP1 (**Fig. 6B**), reflecting the major LMP1
contribution to latency III datasets used in KEGG. Likewise, NF-κB and TLR signaling were also
highly enriched, whereas downmodulated genes were enriched in the term primary
immunodeficiency. Highly concordant effects were observed in the BL-41 cell context, where the
same KEGG pathways were the most enriched amongst LMP1 upregulated genes (**Fig. S7A-**
C). Cross-comparison of expression patterns in Akata and BL-41 with WT vs DM LMP1
expression again revealed highly concordant results (**Fig. 6C-D and S7D-F**), further validating a
range of host genes as targets of TES1 and 2 signaling.

To gain insights into how TES2 signaling shapes LMP1 genome-wide targets, we next
cross-compared transcriptomes from Akata expressing WT vs TES1 mutant LMP1. At a fold-
change >2 and adjusted p-value <0.05 cutoff, 561 genes were more highly expressed in WT
than LMP1 TES1m, whereas 201 were less highly expressed. Interestingly, multiple interferon
stimulated genes, including IFIT1, IFI6, STAT1 and IFI44 were amongst the most highly
upregulated in WT LMP1 expressing cells (**Fig. 6E**). Enrichr analysis identified TNF, Nod-like
receptor (NLR) and JAK/STAT signaling to be the most highly enriched KEGG pathways
amongst genes more highly expressed in WT LMP1+ cells, whereas oxidative phosphorylation
was the most highly enriched KEGG pathway amongst genes more highly expressed in cells
expressing the TES1 mutant (**Fig. 6F**). Similar analyses on BL-41 cell datasets again revealed
large numbers of differentially expressed genes in WT vs TES1m LMP1 expressing cells (**Fig.**
S8A-B and **Table S5**).

To then gain insights into how TES1 signaling shapes LMP1 genome-wide target gene
effects, we cross-compared Akata differentially expressed genes at 24 hours post expression of
WT vs TES2 mutant LMP1. At a fold-change >2 and adjusted p value <0.05 cutoff, 275 genes
were more highly expressed in Akata with WT than TES2 mutant LMP1, whereas 118 were less
highly expressed. Once again, multiple interferon stimulated genes (ISG), including STAT1, IFI6
and OAS1 were more highly expressed in WT cells. Enrichr analysis identified sphingolipid
signaling and metabolism to be most highly enriched KEGG pathways amongst genes
upregulated genes, whereas TCA cycle was the most significant KEGG pathway amongst
genes more highly expressed with TES2 mutant LMP1 expression (**Fig. 6G-H**). Similar numbers
of genes were differentially regulated between WT and TES2m expressing cells in the BL-41
context, where Toll-like receptor signaling was the most highly enriched term amongst genes
more highly expressed in WT LMP1+ cells (**Fig S8C-D** and **Table S6**). These analyses are
consistent with a model in which TES1 and TES2 signaling additively or synergistically
upregulate ISGs. Direct cross-comparison of TES1 vs TES2 signaling in Akata and BL41 further
revealed pathways selectively targeted by either (**Fig. S8E-H**). In the Akata environment, cells
expressing TES2m more highly induced ISGs, including IFIT1, IFI6, OAS, IFI44 and DDX58
(**Fig. S8E**). Enrichr analysis indicated that TES1 signaling most strongly induced the Nod-like
receptor (NLR), necroptosis and chemokine signaling KEGG pathways. By contrast, TES2
signaling (from the TES1 mutant) most highly induced growth hormone and multiple amino acid
metabolism KEGG pathways (**Fig. S8F**). In BL-41 cells, interferon stimulated genes were not as
highly induced by TES2m (**Fig. S8G**). Since non-canonical NF-κB activity can strongly impact B
cell type I interferon pathways (74), we suspect that differences in basal NF-κB activity in BL-41
may compensate to some extent to reduce this phenotype. Instead, cell adhesion molecules
and TNF signaling were most highly enriched. For instance, CFLAR was significantly more
highly induced by TES1 signaling, as was OTULIN, a deubiquitinating enzyme that controls
TNF/NF-κB canonical pathway. FoxO and Toll-like receptor signaling were the most highly
enriched KEGG pathways induced by TES2 signaling (by the TES1 mutant) in BL-41, with
FOXO signaling the most selectively induced by TES1 mutant LMP1 (**Fig. S8H**).

We next directly cross-compared results from our LCL and Burkitt systems. Volcano plot
analysis identified host cell genes whose expression was induced by Akata WT LMP1
expression but decreased by LCL LMP1 KO, suggesting that they are bone fide LMP1 targets
(**Fig. S9A,** blue circles and **Table S1, S7**). This gene set included CFLAR, TRAF1, EBI3, CCL2,
CD40, consistent with prior studies (27, 50, 55, 75-78). Similarly, genes whose expression was
suppressed by Akata WT LMP1 expression but induced by LCL LMP1 KO were identified as
LMP1-repressed host targets (**Fig. S9A**, red circles and **Table S1, S7**). We similarly cross-
compared data from our Akata LMP1 expression and LCL LMP1 rescue datasets. Key targets of
TES1 signaling, whose expression was significantly lower in Akata with TES1 mutant than WT
LMP1 and also in LCLs rescued by TES1 mutant versus WT LMP1, included CFLAR, TRAF1,
NFKB2 and CCL22 (**Fig. S9B** and **Table S2, S7**). Likewise, key TES2 targets more highly
induced by WT than by TES2 mutant in both contexts included CCL22 and EBI3, whereas CR2,
which encodes the EBV B-cell receptor complement receptor 2, was instead more highly
expressed in cells with TES2m than WT LMP1 expression, suggesting it is repressed by TES2
signaling (**Fig. S9C** and **Table S3, S7**). Taken together, these findings serve to validate a class
of host genes as LMP1 targets in the LCL context, although we cannot exclude that they are
regulated through secondary effects.

LMP1 TES1 and TES2 roles in LCL Dependency Factor BATF and IRF4 Expression

We next characterized LMP1 pathways important for BATF and IRF4 induction, given their
key LCL but not Burkitt B-cell dependency factor roles (25, 79, 80). Notably, BATF and Jun
family members bind cooperatively with IRF transcription factors to AP1-IRF composite DNA
elements (AICE) (81), and JunB is the Jun family member predominantly expressed in LCLs
(**Fig. 7A**). WT and TES2m LMP1 upregulated IRF4 mRNA abundance to a similar extent in
Akata, whereas TES1m did so to a somewhat lesser extent. By contrast, TES1m and TES2m
each upregulated BATF, but not quite as strongly as WT LMP1 (**Fig. 7B**). Taken together with
the LCL LMP1 knockout data, these results suggest that LMP1 TES1 and TES2 signaling each
support expression of the host transcription factors that bind to AICE.

We next examined the contribution of LMP1 NF-κB pathways to B cell IRF4 and BATF
expression. LMP1 CRISPR knockout in GM12878 LCLs or in latency III Jijoye Burkitt cells
significantly reduced BATF and to IRF4 expression at the protein level, though effects on IRF4
were more subtle at this early timepoint (**Fig. S10A**). LMP1 induction of IRF4 and BATF was
strongly impaired by induction of either TES1m or TES2m in Akata or in LCLs, relative to levels
in cells with WT LMP1 (**Fig. 7B and S10B**). To test canonical NF-κB pathway roles in IRF4 and
BATF induction, we then induced LMP1 in the absence or presence of a small molecule
antagonist of the kinase IKKβ, which is critical for canonical NF-κB pathway signaling. IKKβ
inhibition blocked their residual induction by TES1m and TES2m (**Fig. 7C-D**). The IKKβ inhibitor
also reduced BATF and IRF4 expression in GM12878 (**Fig. S10C**). Similar results were
obtained in Akata cells that co-induced LMP1 with an IκBα super-repressor (IκBα-SR), in which
IκBα serine 32 and 36 to alanine point mutations prevent its canonical pathway phosphorylation
and proteasomal degradation (**Fig. S10D**). Furthermore, CRISPR KO of the canonical NF-κB
pathway kinase TAK1 significantly impaired IRF4 induction by WT and also TES2m LMP1 (**Fig.**
7E). Taken together, these results suggests that canonical NF-κB pathways driven by both
TES1 and TES2 signaling are each important for BATF and IRF4 expression (**Fig. 7F**).

LMP1 TES1 and TES2 roles in EBV Super-Enhancer target induction

The five LMP1-activated NF-κB transcription factor subunits and four EBNAs target a set of
LCL host genome enhancers termed EBV super-enhancers (EBV SE). EBV SE are
characterized by occupancy by all five NF-κB subunits, EBNAs 2, LP, 3A and 3C and markedly
higher and broader histone H3K27ac ChIP-seq signals than at typical LCL enhancers (**Fig. 8A**).
SE are critical for cell identity and oncogenic states (82), and EBV SE are important for LCL
growth and survival (83-85). However, little has remained known about the extent to which
LMP1 TES1 versus TES2 signaling contribute to EBV SE. To gain insights, we therefore
interrogated EBV SE gene target responses, we first plotted EBV SE gene target responses to
GM12878 LMP1 CRISPR KO. At the early timepoint of 48 hours after LMP1 editing, expression
of SE targets TRAF1 is significantly decreased while expression of PRDM1 (which encodes the
transcription repressor BLIMP1) and GPR15 (which encodes a G-protein coupled chemokine
receptor) each is significantly increased, though most other EBV SE gene targets did not
significantly change at this early timepoint (**Fig. 8B,** red circles and **Table S7**). This data
suggests that EBV SE are robust to short-term perturbations of LMP1 expression, perhaps
given persistence of the established epigenetic landscape built at these key sites.

To then identify the extent to which LMP1 is sufficient to alter expression of these LCL EBV
SE targets, we next visualized effects of WT, TES1m or TES2m rescue of LMP1 KO LCLs.
Potentially because this analysis could be done at a later timepoint post LMP1 CRISPR editing
(day 6 post LMP1 KO), we observed stronger effects on EBV SE target gene expression.
Rescue with TES1 mutant LMP1 caused significant loss of CFLAR, BCL2 and TRAF1 relative to
levels with WT rescue, whereas rescue with either TES mutant significantly lowered levels of
CD86 and BIRC3 messages from levels observed with WT rescue (**Fig. 8C-D** and **Table S8-9**).
As a complementary approach, we also analyzed effects of conditional LMP1 induction in our
Burkitt models on EBV SE target gene expression. In Akata B cells, WT LMP1 induction was
sufficient to significantly upregulate the abundance of the majority of EBV SE target gene
mRNAs (**Fig. S11A** and **Table S1**). This result suggests that while EBNA-2, LP, 3A and 3C also
target these sites in LCLs, LMP1 can independently alter expression of most EBV SE targets,
albeit not necessarily to the same extent as latency III. By contrast, Akata TES1m or TES2m
induction less strongly induced most EBV SE targets (**Fig. S11B-C**). A similar pattern was
observed in BL-41 cells (**Fig. S11D-F** and **Table S4**). Taken together, our results indicate that
TES1 and TES2 likely play key joint roles in the induction of EBV SE target genes.

Discussion:

Why the LMP1 C-terminal tail TES1 and TES2 domains are each necessary for lymphoblastoid B-cell immortalization, and whether each are necessary for LCL survival have remained longstanding questions. To gain insights into key LMP1 B-cell roles, we used a novel LCL LMP1 KO with conditional LMP1 rescue system to identify that signaling by TES1, but not TES2, is required for LCL survival. We performed systematic B-cell transcriptome-wide analyses to identify effects of LMP1 knockout in the absence or presence of rescue by WT, TES1 mutant or TES2 mutant LMP1, at early timepoints where cells remained viable. These highlighted key LMP1 TES1 and TES2 roles in support of LCL dependency factor and EBV super-enhancer target gene expression. As a complimentary approach to identify host B-cell genome-wide targets of LMP1 signaling, we also profiled EBV-negative Burkitt B-cell responses to conditional expression of wildtype LMP1, or LMP1 point mutants abrogated for signaling by TES1 or TES2. As has previously been described in comparisons of LMP1 vs LMP2A expression(86), TES1 and TES2 signaling effects were not simply additive, but yielded distinct effects on host target genes, with either TES domain more strongly inducing or repressing genes at particular sites, but opposing one another at other sites to fine tune target gene expression (**Fig. 9**).

LMP1 KO rapidly altered expression of ~3400 LCL genes, with roughly similar numbers of host genes being downregulated as upregulated. This result is consistent with prior microarray analyses of LMP1 targets in 293 cells induced for TES2 signaling and in Burkitt-cells induced for LMP1 (27, 50), as well as in microarray analysis of EBV-infected B-cells at timepoints where LMP1 expression increases(87), further suggesting that LMP1 strongly remodels the transcriptome by pleotropic effects on host gene expression. Similar numbers of LMP1 targets were also found in microarray profiling of B-cells with transgenic LMP1 expression (77). While TES1 and TES2 induce host genes through activation of NF-κB, MAP kinase, PI3K and interferon regulatory factor pathways (3-5, 11-20), comparatively little is known about how LMP1 downmodulates target gene expression. However, one mechanism by which LMP1 may repress target genes could be through NF-κB complexes, including p50 or p52 homodimers, or p50:52 heterodimers, potentially together with BCL3 (88, 89), as these NF-κB complexes lack transactivation domains. We do not suspect that these changes were secondary to cell death, as we performed profiling on viable cells at an early timepoint post-CRISPR editing. However, the result that LMP1 is critical for LCL survival builds on prior analyses, which showed that blockade of LMP1/NF-κB signaling triggers LCL apoptosis (50, 87, 90, 91).

Conditional expression of WT LMP1 rescued LCL survival, confirming on-target CRISPR effects on EBV genomic LMP1. Our rescue approach identified that loss of TES1, but not TES2 signaling, triggered LCL apoptosis, as judged by upregulation of caspase 3 and 7 activity and by FACS analysis for plasma membrane Annexin V. Disruption of cell death signaling is a hallmark of cancer (92), and enrichment analysis identified that the KEGG apoptosis pathway was highly altered by loss of TES1. Notably, LMP1 has thus far remained an undruggable target. Therefore, these results suggest that small molecule or peptide inhibitors that block TES1 signaling may have therapeutic benefit, even in the absence of effects on TES2, for instance in the setting of EBV-driven post-transplant and central nervous system lymphomas, which frequently express the latency III program and which are modeled by LCLs. It will be of interest to determine whether TES1 signaling has similarly important roles in apoptosis blockade in other EBV-infected tumor contexts, including in Hodgkin lymphoma Reed-Sternberg tumor cells and in nasopharyngeal carcinoma, where little is presently known about TES1 vs TES2 roles.

The LCL dependency factor CFLAR, which encodes the extrinsic apoptosis pathway inhibitor c-
FLIP, was highly downmodulated upon loss of TES1 signaling, to a significantly greater extent
than upon loss of TES2 signaling. This is consistent with prior microarray analysis identified that
identified cFLIP as an LMP1 target (50), which we now mostly induced by TES1 signaling. We
previously identified that c-FLIP is required for LCL survival and is required to block the extrinsic
apoptosis pathway that is otherwise triggered by TNFα signaling, likely in response to EBV
oncogenic stress (25). Therefore, our data newly identified CFLAR as a key TES1 target gene
and suggest that TES1 signaling is required for LCL survival, at least in part due to obligatory
roles in cFLIP induction. Our data raises the interesting question of why CFLAR expression is
particularly dependent on TES1 signaling, to a much greater extent than TES2. It is plausible
that a TES1-driven non-canonical NF-κB pathway is particularly important for cFLIP
transcription. However, TES1 signaling also strongly activates canonical NF-κB pathways (13,
15, 89), which may be critical for CFLAR induction. Alternatively, MAP kinases or PI3K activated
by TES1 (4, 36, 79, 88, 93-95) may also support CFLAR expression.

LMP1 expression both activates and blocks apoptosis (42, 43, 96), and our data suggests that
TES1 induction of CFLAR is central to this balance, perhaps together with BCL2 family
members such as BFL1 (96, 97). However, in addition to targeting CFLAR, apoptosis pathways
were enriched amongst LMP1 target genes. It has also been reported that the six LMP1
transmembrane domains induce apoptosis through activation of an unfolded protein response,
while LMP1 C-terminal domain signaling counteracts this (96). Similarly, LMP1 induces *c-jun*,
junB and *junD* (98), which may play roles in balancing proliferation and apoptosis responses
(99). LMP1 also closely regulates the expression of pro-apoptotic and anti-apoptotic genes to
allow for cell proliferation (98). We also observed downregulation of the p53 antagonist MDM2
and upregulation of p53 upon LMP1 KO. Both TES1 and TES2 had important roles in regulation
of MDM2 expression. Thus, taken together with prior studies, our data sheds light into the
balance of cell death and survival signals triggered by LMP1 signaling.

Our data further highlight LMP1 TES1 and TES2 roles in support of additional LCL dependency
factors, in particular BATF and IRF4. In contrast to CFLAR, TES1 and TES2 signaling were
each found to be important for BATF and IRF4 expression, both in LCL and Burkitt cell models.
TES1 and TES2 driven canonical NF-κB signaling supported BATF induction in EBV-negative
Burkitt cells, where EBNA2 is not expressed. BATF and IRF4 expression rapidly decreased
upon LMP1 KO in LCLs, even upon rescue by LMP1 signaling from only one TES domain.
Thus, in LCLs, EBNA2 and LMP1-stimulated canonical NF-κB jointly induce this critical LCL
dependency factor. LMP1 canonical NF-κB pathways were also critical for inducing IRF4, which
binds with BATF to composite AICE DNA sites. As EBNA2 also supports BATF (100) and
EBNA3C also supports IRF4 expression (101), our results further highlight BATF and IRF4 as
major hubs of EBV oncoprotein cross-talk.

We previously used ChIP-seq to characterize the LCL NF-κB genomic binding landscape (32).
Rather than identifying readily recognizable LMP1 canonical versus non-canonical NF-κB target
genes, this study identified complex patterns of occupancy by the five NF-κB transcription factor
subunits at LCL enhancers and promoters. However, LCL enhancers often target multiple
genes, often from long distances (83), complicating cross-comparison with this study.
Furthermore, concurrent LMP1 TES1 and TES2 signaling yields up to 13 distinct NF-κB
transcription factor dimers in LCL nuclei (32), including dimers such as cRel:p52 that are under
control of both NF-κB pathways. Conditional expression of TES1m or TES2m should yield a

considerably less complex NF-κB. Therefore, a future objective will therefore be to perform NF-
κB ChIP-seq in using the conditional TES1m and TES2m conditional Burkitt models reported
here, as these may yield less complex patterns of NF-κB occupancy.

Our analyses highlight independent, shared and antagonistic LMP1 TES1 and TES2 roles in B-
cell genome-wide target gene regulation (**Fig. 9**). How independent or combined TES1 and
TES2 signaling have different effects on clusters of target genes will be important to define. Our
results suggest multiple testable models. For instance, with regards to genes which were
induced weakly by TES2 signaling, somewhat more by TES1 signaling but more highly by WT
LMP1, we speculate that TES1 and TES2 may cross-talk at the epigenetic level. For example,
our results are consistent with a model in which TES1 signaling increases chromatin
accessibility at these sites, including the genes encoding IFIT1 and CXCL9. Once accessible,
both TES1 and TES2 signal-dependent pathways may then additively or perhaps synergistically
upregulate these sites. Alternatively, TES1 signaling could be needed to dismiss a repressor,
such that these sites can then be stimulated by both LMP1 domains. TES1 signaling may also
activate a key positive regulator such as BCL3 (88), which then functions together with
transcription factors activated by TES1 and TES2 pathways. By contrast, a large number of
genes appeared to be additively repressed by TES1 and TES2 signaling. TES1 and TES2 may
recruit co-repressors to these sites, may additively recruit the same repressor or may reduce
chromatin accessibility.

We also identified a cluster of LMP1 response genes, in which unopposed TES2 signaling
caused target gene downregulation (**Fig. 9**). In Akata cells, this cluster (Cluster 3) was enriched
for metabolism genes, raising the possibility that another key TES1 signaling role is to support
metabolic pathway remodeling by EBV, such as glutathione metabolism or OXPHOS (86, 102-
104). It is possible that TES2 induces a repressor that targets these sites, but that TES1
signaling serves to blunt its induction. Alternatively, TES1 signaling may induce an activator that
counter-balances TES2-driven repressor activity. Or, TES2 signaling may reduce chromatin
accessibility at these sites in the absence of TES1. By contrast, genes in Akata Cluster 5 were
upregulated by unopposed TES2 signaling, but not by WT LMP1 or unopposed TES1 signaling
(**Fig. 9**). TES1 signaling may instead recruit repressors or alter chromatin accessibility at these
sites. Epigenetic analyses of histone repressive marks, such as ChIP studies of H3K9me3 and
H3K27me3, as well as ATAC-seq studies of DNA packaging, should help to differentiate
between these and other possibilities.

Our studies provide insights into TES1 and TES2 roles in regulation of B-cell EBV SE targets.
Although EBV SE are highly co-occupied by all five LMP1-activated NF-κB subunits, individual
TES1 and TES2 roles in EBV SE target gene regulation has remained unstudied. Interestingly,
while either TES1 or TES2 signaling was sufficient to induce many EBV SE targets, TES1
induced a larger number, perhaps because it highly induces both canonical and non-canonical
pathways and therefore activates all 5 NF-κB subunits. By contrast, LMP1 KO perturbed
expression of only a small number of EBV SE targets in LCLs, likely because of the early
timepoint profiled prior to cell death, which left little time for epigenetic remodeling of these sites.

WT or DM LMP1 expression caused highly concordant changes in host gene expression in
Akata and BL-41 Burkitt models. We speculate that differences in response to signaling by
TES1 or TES2 alone may have instead arisen from distinct host genome mutation landscapes
between these two human tumor derived models, which alter the basal NF-κB level.

Nonetheless, since EBV can infect a wide range of B-cells, including of distinct differentiation or
activation states that alter NF-κB states, differences between Akata and BL-41 provide insights
into how LMP1 may function in differing human B-cell contexts, and suggest that LMP1 may
have evolved signaling by both TES domains to increase robustness across the spectrum of
infected B-cell states.

LMP1 polymorphisms have been observed across EBV strains, in particular between type I and
II EBV (105). In addition, LMP1 C-terminal tail polymorphisms have been observed in several
analyses of EBV genomes isolated from Hodgkin lymphoma and nasopharyngeal carcinoma
tumor cells, though the roles of these variant LMP1 sequences remain controversial. Amongst
the best studied is a 30 base pair deletion present in the EBV CAO and 1510 strains isolated
from Asian NPC tumors, which causes loss of LMP1 residues 343-352 (106, 107). This 30bp
deletion has also been reported as enriched in EBV genomes isolated from Hodgkin tumor
samples (107-110). A meta-analysis of 31 observational studies suggested a possible
association between this LMP1 C-terminal tail deletion and nasopharyngeal carcinoma
susceptibility, but was limited by small sample size and considerable variation between studies
(111). Deletion of these residues, which comprise the last eight residues between the
TES1/CTAR1 and the first two residues of TES2/CTAR2, enhance rodent fibroblast
transformation by LMP1 (112) and may reduce immunogenicity (113), but were not found to
enhance LMP1-mediated NF-κB activation (114). These strains also have multiple additional
LMP1 amino acid polymorphisms, which are instead implicated in enhanced NF-κB activation,
and were mapped to the LMP1 transmembrane domains (114). Little information is presently
available about how these polymorphisms alter LMP1 target gene expression. It will therefore
be of interest to use the approaches presented here to characterize how LMP1 polymorphisms
present in tumor-derived EBV strains may alter transcriptome responses to TES1 and TES2
signaling.

In summary, we identified LMP1 genome-wide B-cell targets and characterized their responses
to signaling by TES1 and/or TES2. Signaling by TES1, but not TES2 was identified to be critical
for blockade of LCL apoptosis, and CFLAR was identified as the LCL dependency factor most
strongly impacted by shutoff of TES1 signaling as opposed to TES2. CRISPR KO approaches
highlighted LCL genes that are highly sensitive to perturbation of TES1 and/or TES2 signaling.
K-means analysis highlighted gene clusters with distinct expression responses to signaling by
one or both LMP1 transformation essential domains in the latency III LCL versus EBV-negative
Burkitt B-cell contexts. These studies highlight multiple levels by which TES1 and TES2
signaling alter LMP1 target gene expression, including by additive vs opposing roles.
Collectively, these studies provide new insights into key non-redundant versus joint TES1 and
TES2 roles in B-cell target gene regulation and highlight TES1 signaling as a key
lymphoblastoid B-cell therapeutic target.

Acknowledgements:

This work was supported by NIH R01 CA228700 and AI164709, U01 CA275301, P01 CA269043 and a Burroughs Wellcome Career Award in Medical Sciences to B.E.G, T32 T32AI007245 and an American Cancer Society Post-doctoral Fellowship to E.M.B, T32AI007061 to N.B., T32 T32AI007245 and F32 AI172329 to L.A.M.N and NIH K99 1K99DE031016 to R.G. All RNAseq and proteomics datasets will be published in the GEO omnibus GSE228158, GSE228167, GSE228178 and GSE240732 upon manuscript publication.

AUTHOR CONTRIBUTIONS

B.M and E.M.B performed the experiments. B.M, N.R.B., R.G, E.M.B and L.A.M.N performed bioinformatics analyses. B.M and B.E.G designed the experiments. B.M and B.E.G wrote the manuscript. All authors analyzed the results, read and approved the manuscript**.**

Figure Legends:

Fig. 1. Characterization of LMP1 KO effects on GM12878 LCL target gene regulation

(A) Immunoblot analysis of whole cell lysates (WCL) from GM12878 LCLs transduced with lentiviruses that express control or LMP1 targeting single guide RNAs (sgRNAs). Transduced cells were puromycin selected for 0 vs 2 days, as indicated. Blots for LMP1, for LMP1 target genes TRAF1 and IRF4, and for LMP1-driven non-canonical NF-κB pathway p100/p52 processing are shown. Blots are representative of n=3 experiments.

(B) K-means heatmap analysis of GM12878 LCLs transduced as in (A) with lentivirus expressing control or LMP1 sgRNA and puromycin selected for 48 hours. The heatmap depicts relative Z-scores in each row from n=3 independent RNAseq replicates, divided into two clusters. The Z-score scale is shown at bottom, where blue and red colors indicate lower versus higher relative expression, respectively. Two-way ANOVA P-value cutoff of <0.01 and >2-fold gene expression cutoffs were used.

(C) Enrichr analysis of KEGG pathways most highly changed in GM12878 expressing control versus LMP1 sgRNA (LMP1 KO), as in (A). The X-axis depicts the -Log10 adjusted p-value (adj p-value) scale. The top three most enriched KEGG pathways are shown.

(D) Abundances of two representative Cluster 1 genes from n=3 RNAseq analyses in cells with control vs LMP1 sgRNA. p-values were determined by one-sided Fisher's exact test. *p<0.05, **p<0.01.

(E) Abundances of two representative Cluster 2 genes from n=3 RNAseq analyses in cells with control vs LMP1 sgRNA. p-values were determined by one-sided Fisher's exact test. ***p<0.001.

(F) Volcano plot analysis of host transcriptome-wide GM12878 genes differentially expressed in cells with control versus LMP1 sgRNA expression, as in (B), using data from n=3 RNAseq datasets.

(G) Scatter plot cross comparison of log2 transformed fold change mRNA abundances in GM12878 expressing LMP1 vs. control sgRNA (Y-axis) versus log2 transformed fold change abundances of sgRNAs at Day 21 versus Day 1 post-transduction of GM12878 LCLs in a genome-wide CRISPR screen (25) (X-axis).

(H) String analysis of genes shown in (G). Pathway identifiers for each gene and interaction are colored coded.

(I) Volcano plot analysis of EBV mRNA values in GM12878 expressing LMP1 vs control sgRNAs, as in (B). p-value <0.05 and >2-fold change mRNA abundance cutoffs were used.

Fig. S1. Characterization of LMP1 KO effects on GM12878 LCL target gene regulation.

(A) Relative mean + standard deviation (SD) live cell numbers from CellTitreGlo analysis of n=3 replicates of Cas9+ GM12878 LCLs, transduced with lentiviruses that expressed control or LMP1 sgRNAs and puromycin selected, for 2 versus 4 days.

(B) RT-PCR analysis of CCL22 mRNA abundance in Cas9+ GM12878 post-transduction with lentiviruses that expressed control or LMP1 sgRNA and puromycin selected for 2 days, as in **Fig. 1B**. The same RNA used for RNA-seq (**Fig. 1B**) was used for these qPCR experiments.

Values from cells with control sgRNA were set to 1, and mean fold-change of CCL22 mRNA abundance + SD in cells with LMP1 sgRNA are shown. p-values were determined by one-sided Fisher's exact test from two independent experiments, each with two technical replicates. ***p<0.001.

(C) RT-PCR analysis of EBI3 mRNA abundance in Cas9+ GM12878 post-transduction with lentiviruses that expressed control or LMP1 sgRNA and puromycin selected for 2 days, as in **Fig. 1B**. The same RNA used for RNA-seq (**Fig. 1B**) was used for these qPCR experiments. Values from cells with control sgRNA were set to 1, and mean fold-change of EBI3 mRNA abundance + SD in cells with LMP1 sgRNA are shown. p-values were determined by one-sided Fisher's exact test from two independent experiments, each with two technical replicates. ***p<0.001.

(D) RT-PCR analysis of IRF4 mRNA abundance in Cas9+ GM12878 post-transduction with lentiviruses that expressed control or LMP1 sgRNA and puromycin selected for 2 days, as in Fig. 1B. The same RNA used for RNA-seq (**Fig. 1B**) was used for these qPCR experiments. Values from cells with control sgRNA were set to 1, and mean fold-change of IRF4 mRNA abundance + SD in cells with LMP1 sgRNA are shown. p-values were determined by one-sided Fisher's exact test from two independent experiments, each with two technical replicates. ***p<0.001.

(E) Scatter plot analysis cross-comparing the significance of changes in LCL dependency factor expression upon GM127878 LMP1 KO versus the CRISPR screen significance score for selection against sgRNAs in LCL vs Burkitt dependency factor analysis (25). Shown on the Y-axis are -log10 transformed P-values from RNAseq analysis of GM12878 LCLs transduced with lentiviruses expressing LMP1 versus control sgRNA (as in **Fig. 1F**), versus -log10 transformed P-values from CRISPR LCL vs Burkitt cell dependency factor analysis (25). Higher Y-axis scores indicate more significant differences in expression for the indicated genes in GM12878 with LMP1 vs control sgRNA. Higher X-axis scores indicate a stronger selection against sgRNA targeting the indicated genes in GM12878 LCLs versus P3HR1 Burkitt cells over 21 days of cell culture. Shown are genes with p<0.05 in both analyses.

(F) Volcano plot analysis visualizing KEGG Hodkin lymphoma pathway gene -Log10 (P-value) on the y-axis versus log2 transformed fold change in mRNA abundances on the x-axis of GM12878 genes in cells expressing LMP1 versus control sgRNA (as in **Fig. 1F**). P-value <0.05 and >2-fold change mRNA abundance cutoffs were used.

Fig. 2. **Loss of TES1 but not TES2 signaling triggers LCL apoptosis.**

(A) Schematic diagram of LMP1 WT with TES1 and TES2 domains highlighted. Wildtype (WT) or point mutants abrogated for signaling from TES1 (TES1m), TES2 (TES2m) or double TES1/TES2 mutant (DM) are shown.

(B) Immunoblot analysis of WCL from GM12878 LCLs that expressed control or LMP1 sgRNAs and puromycin selected for 3 days, then induced for expression with the indicated LMP1 rescue cDNA construct for 6 days. Blots are representative of n = 3 experiments.

(C) Growth curve analysis of GM12878 LCLs at the indicated day post expression of control or LMP1 sgRNAs and the indicated LMP1 WT, TES1m, TES2m or DM rescue cDNA. Shown are mean ± SD from n=3 independent experiments. **p<0.01.

(D) FACS analysis of 7-AAD vital dye uptake in GM12878 on day 7 post- expression of LMP1 sgRNAs and the indicated LMP1 rescue cDNA. Shown are percentages of 7-AAD+ cells within the indicated gates. Representative of n=3 experiments.

(E) Mean ± SD of fold change 7-AAD values from n=3 independent experiments of GM12878 with the indicated control or LMP1 sgRNA and rescue cDNA expression, as in (D). Values in GM12878 with control sgRNA and no LMP1 rescue cDNA were set to 1.

(F) FACS analysis of plasma membrane annexin V abundance in GM12878 on day 7 post-expression of control or LMP1 sgRNAs and the indicated LMP1 rescue cDNA. Shown are percentages of 7-AAD+ cells within the indicated gates. Representative of n=3 experiments.

(G) Mean ± SD of fold-change caspase 3/7 activity levels, as determined by caspase 3/7 Glo assay, from n=3 independent experiments of GM12878 with the indicated control or LMP1 sgRNA and rescue cDNA expression. Values in GM12878 with control sgRNA and no LMP1 rescue cDNA were set to 1.

Fig. S2. Loss of TES1 but not TES2 signaling triggers LCL apoptosis.

Mean ± SD of fold change plasma membrane Annexin V values from n=3 independent experiments, using GM12878 with the indicated control or LMP1 sgRNA and rescue cDNA expression. Values in GM12878 with control sgRNA and no LMP1 rescue cDNA were set to 1.

Fig. 3. Characterization of host genome-wide TES1 vs TES2 LCL target genes

(A) RNAseq K-means heatmap analysis of GM12878 LCLs transduced with lentivirus expressing LMP1 sgRNA and induced for WT, TES1m or TES2m rescue cDNA expression for 6 days. The heatmap depicts relative Z-scores in each row from n=3 independent RNAseq datasets, divided into six clusters. The Z-score scale is shown at bottom, where blue and red colors indicate lower versus higher relative expression, respectively. Two-way ANOVA P-value cutoff of <0.05 and >2-fold gene expression cutoffs were used. The top three most highly enriched KEGG pathways amongst genes within each cluster are shown at right.

(B) Heatmap analysis of KEGG apoptosis pathway gene relative row Z-scores from RNAseq analysis as in (A). The Z-score scale is shown at bottom, where blue and red colors indicate lower versus higher relative expression, respectively. Two-way ANOVA P-value cutoff of <0.05 and >2-fold gene expression cutoffs were used.

(C) Scatter plot analysis cross comparing log2 transformed fold change of LCL dependency factor mRNA abundances in GM12878 expressing LMP1 sgRNA together with TES2 mutant versus wildtype cDNA rescue (Y-axis) and TES1 mutant versus wildtype cDNA rescue (X-axis) from triplicate RNAseq datasets, as in (A). This analysis highlighted that CFLAR and to a lesser extent NFKB2 and CCND2 mRNAs were more highly downmodulated by TES1m than TES2m rescue, relative to levels in cells with WT LMP1 rescue. Shown are genes differentially regulated by >2 fold with either TES1m or TES2m rescue, relative to levels with WT LMP1 rescue.

(D) Immunoblot analysis of c-FLIP and load control GAPDH expression in WCL from GM12878 LCLs with the indicated control or LMP1 sgRNA and LMP1 rescue cDNA expression. Representative of n=3 experiments.

Figure S3. Characterization of TES1 vs TES2 LCL dependency factor and Hodgkin lymphoma pathway targets.

(A) Heatmap analysis of CRISPR defined LCL dependency factor gene relative row Z-scores from RNAseq of GM12878 expressing LMP1 sgRNA and the indicated rescue cDNA, as in **Fig. 3**. The Z-score scale is shown at bottom, where blue and red colors indicate lower versus higher relative expression, respectively. Two-way ANOVA P-value cutoff of <0.05 and >2-fold gene expression cutoffs were used.

(B) Heatmap analysis of KEGG Hodgkin Lymphoma pathway gene relative row Z-scores from RNAseq of GM12878 expressing LMP1 sgRNA and the indicated rescue cDNA, as in **Fig. 3**. Two-way ANOVA P-value cutoff of <0.05 and >2-fold gene expression cutoffs were used.

(C) RT-PCR analysis of CCL22 mRNA abundance in GM12878 LCLs transduced with lentivirus expressing LMP1 sgRNA and induced for WT, TES1m or TES2m rescue cDNA expression for 6 days, as in **Fig. 3A**. The same RNA used for RNA-seq in **Fig. 3A** was used for these qPCR experiments. Values from cells with WT LMP1 rescue cDNA were set to 1, and mean fold-change of CCL22 mRNA abundance + SD in cells with TES1m or TES2m rescue cDNA are shown. p-values were determined by one-sided Fisher's exact test from two independent experiments, each with two technical replicates. ***p<0.001.

(D) RT-PCR analysis of EBI3 mRNA abundance in GM12878 LCLs transduced with lentivirus expressing LMP1 sgRNA and induced for WT, TES1m or TES2m rescue cDNA expression for 6 days, as in **Fig. 3A**. The same RNA used for RNA-seq in **Fig. 3A** was used for these qPCR experiments. Values from cells with WT LMP1 rescue cDNA were set to 1, and mean fold-change of EBI3 mRNA abundance + SD in cells with TES1m or TES2m rescue cDNA are shown. p-values were determined by one-sided Fisher's exact test from two independent experiments, each with two technical replicates. ***p<0.001.

(E) RT-PCR analysis of IRF4 mRNA abundance in GM12878 LCLs transduced with lentivirus expressing LMP1 sgRNA and induced for WT, TES1m or TES2m rescue cDNA expression for 6 days, as in **Fig. 3A**. The same RNA used for RNA-seq in **Fig. 3A** was used for these qPCR experiments. Values from cells with WT LMP1 rescue cDNA were set to 1, and mean fold-change of IRF4 mRNA abundance + SD in cells with TES1m or TES2m rescue cDNA are shown. p-values were determined by one-sided Fisher's exact test from two independent experiments, each with two technical replicates. ***p<0.001.

Fig. 4. Characterization of LCL pathways targeted by TES1 vs TES2 signaling

(A) Volcano plot analysis of host transcriptome-wide GM12878 genes differentially expressed in LMP1 KO GM12878 with WT vs TES1 mutant cDNA rescue. Higher X-axis fold changes indicate higher expression with WT LMP1 rescue, whereas lower X-axis fold changes indicate higher expression with TES1m rescue. Data are from n=3 RNAseq datasets, as in **Fig. 3**.

(B) Enrichr analysis of KEGG pathways most highly enriched in RNAseq data as in (A) amongst genes more highly expressed in LMP1 KO GM12878 with WT than TES1m rescue (red) vs amongst genes more highly expressed with TES1m than WT rescue (blue).

(C) Volcano plot analysis of host transcriptome-wide GM12878 genes differentially expressed in LMP1 KO GM12878 with WT vs TES2 mutant cDNA rescue. Higher X-axis fold changes indicate higher expression with WT LMP1 rescue, whereas lower X-axis fold changes indicate higher expression with TES2m rescue. Data are from n=3 RNAseq datasets, as in **Fig. 3**.

(D) Enrichr analysis of KEGG pathways most highly enriched in RNAseq data as in (C) amongst genes more highly expressed in LMP1 KO GM12878 with WT than TES2m rescue (red) vs amongst genes more highly expressed with TES2m than WT rescue (blue).

(E) Volcano plot analysis of host transcriptome-wide GM12878 genes differentially expressed in LMP1 KO GM12878 with TES1 vs TES2 mutant cDNA rescue. Higher X-axis fold changes indicate higher expression with TES1m rescue, whereas lower X-axis fold changes indicate higher expression with TES2m rescue. Data are from n=3 RNAseq datasets, as in **Fig. 3**.

(F) Enrichr analysis of KEGG pathways most highly enriched in RNAseq data as in (E) amongst genes more highly expressed in LMP1 KO GM12878 with TES1m than TES2m rescue (red) vs amongst genes more highly expressed with TES2m than TES1m rescue (blue).

Fig. 5. Characterization of host genome-wide Akata B-cell LMP1 target genes.

(A) K-means heatmap analysis of RNAseq datasets from n=3 replicates generated in EBV-Akata Burkitt cells with conditional LMP1 WT, TES1m, TES2m or DM expression induced by 250 ng/ml doxycycline for 24 hours. The heatmap visualizes host gene Log2 Fold change across the four conditions, divided into six clusters. A two-way ANOVA P value cutoff of <0.01 and >2-fold gene expression were used. # of genes in each cluster is indicated at right.

(B) Heatmaps of representative Cluster 1 differentially regulated genes (top), with column maximum (max) colored red and minimum (min) colored blue, as shown by the scalebar. Also shown are expression values of two representative Cluster 1 genes (lower left) and Enrichr analysis of KEGG pathways significantly enriched in Cluster 1 gene sets (lower right). p-values were determined by one-sided Fisher's exact test. ***p<0.001.

(C) Heatmaps of representative Cluster 2 differentially regulated genes (top), as in (B). Also shown are expression values of two representative Cluster 2 genes (lower left) and Enrichr analysis of KEGG pathways significantly enriched in Cluster 2 gene sets (lower right). p-values were determined by one-sided Fisher's exact test. *p<0.05, **p<0.01, ***p<0.001.

(D) Heatmaps of representative Cluster 6 differentially regulated genes (top), as in (B). Also shown are expression values of two representative Cluster 6 genes (lower left) and Enrichr analysis of KEGG pathways significantly enriched in Cluster 6 gene sets (lower right). p-values were determined by one-sided Fisher's exact test. *p<0.05, ***p<0.001.

Fig. S4. Validation of LMP1 WT, TES1m, TES2m and DM conditional expression system in EBV-negative Akata Burkitt B-cells.

(A) Immunoblot analysis of WCL from Akata cells induced for LMP1 WT, TES1m, TES2m or DM expression by addition of 250 ng/ml doxycycline (Dox) for 24 hours, as indicated. For cross-comparison, WCL from equal numbers of Mutu I Burkitt lymphoma (latency I, lacks LMP1 expression) and GM12878 were also included at right. Blots are representative of n = 3 experiments.

(B) Immunoblot analysis of WCL from BL-41 cells induced for WT, TES1m, TES2m or DM LMP1 by addition of 250 ng/ml Dox for 24 hours, as indicated. For cross-comparison, WCL from equal numbers of Mutu I Burkitt and GM12878 were also included at right. Blots are representative of n = 3 experiments.

(C) FACS analysis of plasma membrane (PM) ICAM-1 abundance in Akata cells induced for
LMP1 by 250 ng/ml Dox for 24 hours, as indicated. Y-axis are histogram cell counts, X-axis
represents PM ICAM-1 abundance. For comparison, levels in GM12878 LCLs or latency I Mutu
I Burkitt cells are shown.

(D) PM ICAM-1 Mean Fluorescence Intensity (MFI) + standard deviation (SD) from n=3
replicates in Akata cells with the indicated LMP1 expression, as in (C). p-values were
determined by one-sided Fisher's exact test. **p<0.001, ***p<0001.

(E) FACS analysis of PM Fas abundance in Akata cells induced for LMP1 by 250ng/ml of Dox
for 24 hours as indicated. For comparison, GM12878 LCLs or latency I Mutu I Burkitt cells were
also analyzed.

(F) PM Fas MFI + SD from n=3 replicates in Akata cells with the indicated LMP1 expression, as
in (E). P-values were determined by one-sided Fisher's exact test. **p<0.001, ***p<0.0001.

(G) FACS analysis of PM ICAM-1 in BL-41 cells induced for LMP1 expression by 250ng/ml Dox
for 24 hours, as indicated. For comparison, GM12878 and Mutu I were also analyzed. p-values
were determined by one-sided Fisher's exact test. **p<0.001, ***p<0.0001.

(H) PM ICAM-1 MFI + SD from n=3 replicates in BL-41 cells with the indicated LMP1
expression, as in (G). P-values were determined by one-sided Fisher's exact test. **p<0.001,
***p<0.0001.

(I) FACS analysis of PM Fas levels in BL-41 cells induced for LMP1 expression by 250 ng/ml
dox for 24 hours, as indicated. GM12878 and Mutu I were analyzed for cross-comparison. p-
values were determined by one-sided Fisher's exact test. **p<0.001, ***p<0.0001.

(J) PM Fas MFI + SD from n=3 replicates of BL-41-LMP1 with the indicated LMP1 expression,
as in (I). Mutu I and GM12878 were analyzed for comparison. p-values were determined by
one-sided Fisher's exact test. **p<0.001, ***p<0.0001.

Fig. S5. Characterization of host genome-wide Akata B-cell LMP1 target genes, related to
Figure 4.

(A) Heatmaps of representative **Figure 4** Cluster 3 differentially regulated genes (top), with
column maximum colored red and minimum colored blue, as shown by the scalebar. Also
shown are expression values of two representative Clusters 3 genes (lower left) and Enrichr
analysis of KEGG pathways most significantly enriched Cluster 3 gene sets (lower right). p-
values were determined by one-sided Fisher's exact test. **p<0.01.

(B) Heatmaps of representative **Figure 4** Cluster 4 differentially regulated genes (top), with
column maximum colored red and minimum colored blue, as shown by the scalebar. Also
shown are expression values of two representative Clusters 4 genes (lower left) and Enrichr
analysis of KEGG pathways most significantly enriched Cluster 4 gene sets (lower right). p-
values were determined by one-sided Fisher's exact test. *p<0.05, **p<0.01, ***p<0.001.

(C) Heatmaps of representative **Figure 4** Cluster 5 differentially regulated genes (top), with
column maximum colored red and minimum colored blue, as shown by the scalebar. Also
shown are expression values of two representative Clusters 5 genes (lower left) and Enrichr
analysis of KEGG pathways most significantly enriched Cluster 5 gene sets (lower right). p-
values were determined by one-sided Fisher's exact test. *p<0.05, ***p<0.001.

(D) RT-PCR analysis of CCL22 mRNA abundance in cells with conditional LMP1 WT, TES1m, TES2m or DM expression induced by 250 ng/ml doxycycline for 24 hours as in Fig. 5A. The same RNA used for RNA-seq in Fig. 5A was used for these qPCR experiments. Values from cells with conditional WT LMP1 expression were set to 1, and mean fold-change of CCL22 mRNA abundance + SD in cells with conditional TES1m,TES2m or DM expression are shown. p-values were determined by one-sided Fisher's exact test from two independent experiments, each with two technical replicates. **$p<0.01$.

(E) RT-PCR analysis of EBI3 mRNA abundance in cells with conditional LMP1 WT, TES1m, TES2m or DM expression induced by 250 ng/ml doxycycline for 24 hours as in Fig. 5A.The same RNA used for RNA-seq in Fig. 5A was used for these qPCR experiments. Values from cells with conditional WT LMP1 expression were set to 1, and mean fold-change of CCL22 mRNA abundance + SD in cells with conditional TES1m,TES2m or DM expression are shown. p-values were determined by one-sided Fisher's exact test from two independent experiments, each with two technical replicates. **$p<0.01$.

(F) RT-PCR analysis of IRF4 mRNA abundance in cells with conditional LMP1 WT, TES1m, TES2m or DM expression induced by 250 ng/ml doxycycline for 24 hours as in Fig. 5A. The same RNA used for RNA-seq in Fig. 5A was used for these qPCR experiments. Values from cells with conditional WT LMP1 expression were set to 1, and mean fold-change of CCL22 mRNA abundance + SD in cells with conditional TES1m,TES2m or DM expression are shown. p-values were determined by one-sided Fisher's exact test from two independent experiments, each with two technical replicates. **$p<0.01$.

Fig. S6. RNAseq analysis of BL-41 B-cell responses to WT, TES1, TES2 or DM LMP1.

(A) K-means heatmap analysis of RNAseq datasets from n=3 replicates generated in EBV- BL-41 Burkitt cells with conditional LMP1 WT, TES1m, TES2m or DM expression induced by 250 ng/ml doxycycline for 24 hours. The heatmap visualizes host gene Log2 Fold change across the four conditions, divided into six clusters. A two-way ANOVA P value cutoff of <0.01 and >2-fold gene expression were used. # of genes in each cluster is indicated at right.

(B) Heatmaps of representative Cluster 1 differentially regulated genes (top), with column maximum (max) colored red and minimum (min) colored blue, as shown by the scalebar. Also shown are expression values of two representative Cluster 1 genes (lower left) and Enrichr analysis of KEGG pathways significantly enriched in Cluster 1 gene sets (lower right). p-values were determined by one-sided Fisher's exact test. ***$p<0.001$.

(C) Heatmaps of representative Cluster 2 differentially regulated genes (top). Also shown are expression values of two representative Cluster 2 genes (lower left) and Enrichr analysis of KEGG pathways significantly enriched in Cluster 2 gene sets (lower right). p-values were determined by one-sided Fisher's exact test. ***$p<0.001$.

(D) Heatmaps of representative Cluster 4 differentially regulated genes (top). Also shown are expression values of two representative Cluster 4 genes (lower left) and Enrichr analysis of KEGG pathways significantly enriched in Cluster 4 gene sets (lower right). p-values were determined by one-sided Fisher's exact test. *$p<0.05$, ***$p<0.001$.

(E) Heatmaps of representative Cluster 6 differentially regulated genes (top). Also shown are expression values of two representative Cluster 6 genes (lower left) and Enrichr analysis of KEGG pathways significantly enriched in Cluster 6 gene sets (lower right). p-values were determined by one-sided Fisher's exact test.*$p<0.05$, **$p<0.01$, ***$p<0.001$.

Fig. 6. Characterization of Akata B-cell pathways targeted by TES1 vs TES2 signaling.

(A) Volcano plot analysis of host transcriptome-wide genes differentially expressed in Akata
cells conditionally induced for WT LMP1 expression for 24h by 250 ng/ml Dox versus in mock
induced cells. Higher X-axis fold changes indicate genes more highly expressed in cells with
WT LMP1 expression, whereas lower X-axis fold changes indicate higher expression in cells
mock induced for LMP1. Data are from n=3 RNAseq datasets, as in **Fig. 5**.

(B) Enrichr analysis of KEGG pathways most highly enriched in RNAseq data as in (A) amongst
genes more highly expressed in Akata with WT LMP1 (red) vs amongst genes more highly
expressed with mock LMP1 induction (blue).

(C) Volcano plot analysis of host transcriptome-wide genes differentially expressed in Akata
cells conditionally induced for WT vs DM LMP1 expression for 24h by 250 ng/ml Dox. Higher X-
axis fold changes indicate genes more highly expressed in cells with WT LMP1 expression,
whereas lower X-axis fold changes indicate higher expression in cells with DM LMP1. Data are
from n=3 RNAseq datasets, as in **Fig. 5**.

(D) Enrichr analysis of KEGG pathways most highly enriched in RNAseq data as in (C) amongst
genes more highly expressed in Akata with WT LMP1 (red) vs amongst genes more highly
expressed with DM LMP1 induction (blue).

(E) Volcano plot analysis of host transcriptome-wide genes differentially expressed in Akata
cells conditionally induced for WT vs TES1m LMP1 expression for 24h by 250 ng/ml Dox.
Higher X-axis fold changes indicate genes more highly expressed in cells with WT LMP1
expression, whereas lower X-axis fold changes indicate higher expression in cells with TES1m
LMP1. Data are from n=3 RNAseq datasets, as in **Fig. 5**.

(F) Enrichr analysis of KEGG pathways most highly enriched in RNAseq data as in (E) amongst
genes more highly expressed in Akata with WT LMP1 (red) vs amongst genes more highly
expressed with TES1m LMP1 induction (blue).

(G) Volcano plot analysis of host transcriptome-wide genes differentially expressed in Akata
cells conditionally induced for WT vs TES2m LMP1 expression for 24h by 250 ng/ml Dox.
Higher X-axis fold changes indicate genes more highly expressed in cells with WT LMP1
expression, whereas lower X-axis fold changes indicate higher expression in cells with TES2m
LMP1. Data are from n=3 RNAseq datasets, as in **Fig. 5**.

(H) Enrichr analysis of KEGG pathways most highly enriched in RNAseq data as in (G) amongst
genes more highly expressed in Akata with WT LMP1 (red) vs amongst genes more highly
expressed with TES2m LMP1 induction (blue).

1031

Fig. S7. Cross-comparison of WT and DM LMP1 effects on Akata vs BL-41
transcriptomes.

1034 (A) Volcano plot analysis of host transcriptome-wide genes differentially expressed in BL-41
1035 cells conditionally induced for WT LMP1 expression for 24h by 250 ng/ml Dox versus in mock
1036 induced cells. Higher X-axis fold changes indicate genes more highly expressed in cells with
1037 WT LMP1 expression, whereas lower X-axis fold changes indicate higher expression in cells
1038 mock induced for LMP1. Data are from n=3 RNAseq datasets.

(B) Enrichr analysis of KEGG pathways most highly enriched in RNAseq data as in (A) amongst genes more highly expressed in BL-41 with WT LMP1 (red) vs amongst genes more highly expressed with mock LMP1 induction (blue).

(C) Volcano plot cross-comparison of Log2 transformed fold change of host mRNA levels in BL-41 cells (X-axis) versus Akata cells (Y-axis) uninduced versus induced for WT LMP1 by 250 ng/ml Dox for 24 hours. Selected genes highly WT LMP1 induced in both Burkitt contexts are highlighted in red, whereas selected genes suppressed by LMP1 in both Burkitt contexts are highlighted in blue.

(D) Volcano plot analysis of host transcriptome-wide genes differentially expressed in BL-41 cells conditionally induced for DM versus WT LMP1 expression for 24h by 250 ng/ml Dox. Higher X-axis fold changes indicate genes more highly expressed in cells with WT LMP1 expression, whereas lower X-axis fold changes indicate higher expression in cells induced for DM LMP1. Data are from n=3 RNAseq datasets.

(E) Enrichr analysis of KEGG pathways most highly enriched in RNAseq data as in (D) amongst genes more highly expressed in BL-41 with WT LMP1 (red) vs amongst genes more highly expressed with DM LMP1 induction (blue).

(F) Volcano plot cross-comparison of Log2 transformed fold change of host mRNA levels in BL-41 cells (X-axis) versus Akata cells (Y-axis) induced for DM versus WT LMP1 by 250 ng/ml Dox for 24 hours. Selected genes highly WT LMP1 induced in both Burkitt contexts relative to levels in cells with DM LMP1 expression are highlighted in red, whereas selected genes suppressed by WT LMP1 in both Burkitt contexts are highlighted in blue.

Fig. S8. Cross-comparison of TES1 and TES2 LMP1 effects on Akata vs BL-41 transcriptomes.

A) Volcano plot analysis of host transcriptome-wide genes differentially expressed in BL-41 cells conditionally induced for TES1m vs WT LMP1 expression for 24h by 250 ng/ml Dox. Higher X-axis fold changes indicate genes more highly expressed in cells with WT LMP1 expression, whereas lower X-axis fold changes indicate higher expression induced for TES1m LMP1. Data are from n=3 RNAseq datasets.

(B) Enrichr analysis of KEGG pathways most highly enriched in RNAseq data as in (A) amongst genes more highly expressed in BL-41 with WT LMP1 (red) vs amongst genes more highly expressed with TES1m LMP1 induction (blue).

(C) Volcano plot analysis of host transcriptome-wide genes differentially expressed in BL-41 cells conditionally induced for TES2m vs WT LMP1 expression for 24h by 250 ng/ml Dox. Higher X-axis fold changes indicate genes more highly expressed in cells with WT LMP1 expression, whereas lower X-axis fold changes indicate higher expression induced for TES2m LMP1. Data are from n=3 RNAseq datasets.

(D) Enrichr analysis of KEGG pathways most highly enriched in RNAseq data as in (A) amongst genes more highly expressed in BL-41 with WT LMP1 (red) vs amongst genes more highly expressed with TES2m LMP1 induction (blue).

(E) Volcano plot analysis of host transcriptome-wide genes differentially expressed in Akata cells conditionally induced for TES1m vs TES2m LMP1 expression for 24h by 250 ng/ml Dox. Higher X-axis fold changes indicate genes more highly expressed in cells with TES1m LMP1

expression, whereas lower X-axis fold changes indicate higher expression induced for TES2m
LMP1. Data are from n=3 RNAseq datasets.

(F) Enrichr analysis of KEGG pathways most highly enriched in RNAseq data as in (E) amongst
genes more highly expressed in Akata with TES1m LMP1 (red) vs amongst genes more highly
expressed with TES2m LMP1 induction (blue).

(G) Volcano plot analysis of host transcriptome-wide genes differentially expressed in BL-41
cells conditionally induced for TES1m vs TES2m LMP1 expression for 24h by 250 ng/ml Dox.
Higher X-axis fold changes indicate genes more highly expressed in cells with TES1m LMP1
expression, whereas lower X-axis fold changes indicate higher expression induced for TES2m
LMP1. Data are from n=3 RNAseq datasets.

(H) Enrichr analysis of KEGG pathways most highly enriched in RNAseq data as in (G) amongst
genes more highly expressed in BL-41 with TES1m LMP1 (red) vs amongst genes more highly
expressed with TES2m LMP1 induction (blue).

1095

1096 **Figure S9. Cross-comparison of host genes differentially expressed upon perturbation of**
1097 **LCL LMP1 versus upon LMP1 induction in Akata cells.**

1098

(A) Volcano plot analysis of host genes differentially expressed upon WT LMP1 induction in
Akata (X-axis) versus upon LMP1 KO in GM12878 (Y-axis). Shown are Log2 transformed
mRNA fold change values for Akata cells mock induced versus induced for LMP1 WT
expression for 24 hours (X-axis) versus upon expression of LMP1 vs control sgRNA in
GM18278 for 48 hours. Genes more highly expressed in mock-induced Akata have higher x-
axis values, whereas genes more highly expressed in Akata induced for WT LMP1 have lower
x-axis values. Likewise, genes with higher expression with control sgRNA expression have
higher y-axis values, whereas genes with lower expression in GM12878 with LMP1 KO have
lower Y-axis values. P value <0.05 and >2-fold gene expression cutoffs were used.

(B) Volcano plot analysis of host genes differentially expressed upon TES1m vs WT LMP1
induction in Akata (X-axis) versus upon rescue of LMP1 KO GM12878 with TES1m versus WT
LMP1 (Y-axis). Shown are Log2 transformed mRNA fold change values for Akata cells induced
for TES1m versus WT LMP1 expression for 24 hours (X-axis) versus upon rescue of GM18278
LMP1 KO with TES1m vs WT LMP1 cDNA, as in **Fig 3**. Genes more highly expressed in Akata
with TES1m than WT LMP1 expression have higher x-axis values, whereas genes more highly
expressed in Akata induced for WT LMP1 than TES1m have lower x-axis values. Likewise,
GM12878 genes with higher expression with TES1m rescue have higher y-axis values, whereas
genes with lower expression in GM12878 with TES1m than WT rescue have lower Y-axis
values. P value <0.05 and >2-fold gene expression cutoffs were used.

(C) Volcano plot analysis of host genes differentially expressed upon TES2m vs WT LMP1
induction in Akata (X-axis) versus upon rescue of LMP1 KO GM12878 with TES2m versus WT
LMP1 (Y-axis). Shown are Log2 transformed mRNA fold change values for Akata cells induced
for TES2m versus WT LMP1 expression for 24 hours (X-axis) versus upon rescue of GM18278
LMP1 KO with TES2m vs WT LMP1 cDNA, as in **Fig 3**. Genes more highly expressed in Akata
with TES2m than WT LMP1 expression have higher x-axis values, whereas genes more highly
expressed in Akata induced for WT LMP1 than TES2m have lower x-axis values. Likewise,
GM12878 genes with higher expression with TES2m rescue have higher y-axis values, whereas

genes with lower expression in GM12878 with TES2m than WT rescue have lower Y-axis
values. P value <0.05 and >2-fold gene expression cutoffs were used.

1128

Fig. 7. Roles of TES1 and TES2 canonical NF-κB pathways in LCL dependency factor
BATF and IRF4 expression.

(A) Schematic diagram of JUNB, BATF and IRF4 at an AP-1/IRF composite DNA site.

(B) Mean + SD fold changes of IRF4, BATF and JUNB mRNA abundances from n=3 RNAseq
replicates of Akata cells expressing the indicated LMP1 cDNA for 24 hours, as in Fig. 5. p-
values were determined by one-sided Fisher's exact test. *p<0.05, ***p<0.001.

(C) Schematic diagram of LMP1 TES1 and TES2 NF-κB pathways. TES1 and TES2 each
activate canonical NF-κB pathways, whereas TES1 also activates non-canonical NF-κB.

(D) Immunoblot analysis of WCL from Akata cells induced for LMP1 expression by 250 ng/ml
Dox for 24 hours, either without or with 1 μM IKKβ inhibitor VIII. Shown below are relative fold
changes + SD from n=3 replicates of IRF4 or BATF vs GAPDH load control densitometry
values. Values in vehicle control treated WT LMP1 expressing cells were set to 1. P-values
were determined by one-sided Fisher's exact test. **p<0.001, ***p<0.0001.

(E) Immunoblot analysis of WCL from Cas9+ Akata cells expressing control or either of two
TAK1 targeting sgRNAs, induced for LMP1 expression by 250 ng/ml Dox for 24 hours. Shown
below are relative foldchanges + SD from n=3 replicates of IRF4 or BATF vs GAPDH load
control densitometry values. Levels in cells with control sgRNA (sgControl) and WT LMP1 were
set to 1. **p<0.001, ***p<0.0001.

(F) Model of additive TES1 and TES2 canonical NF-κB pathway effects on BATF and IRF4
induction.

1149

Fig. S10. Roles of TES1 and TES2 canonical NF-κB pathways in BATF and IRF4
induction.

(A) Immunoblot analysis of WCL from latency III Jijoye Burkitt cells or GM12878 LCL expressing
LMP1 targeting sgRNA, as indicated. Shown below are relative fold changes + SD from n=3
replicates of IRF4 or BATF vs GAPDH load control densitometry values, with values in sgRNA
control expressing cells set to 1. P-values were determined by one-sided Fisher's exact test.
**p<0.001, ***p<0.0001.

(B) Relative +SD BATF and IRF4 mRNA levels from n=3 RNAseq replicates from LMP1 KO
GM12878 with TES1m, TES2m or WT LMP1 rescue cDNA expression, as in **Fig 3-4**. BATF or
IRF4 levels in cells with WT LMP1 rescue were defined as 1. ***p<0.0001.

(C) Immunoblot analysis of WCL from GM12878 LCLs treated with vehicle control or 1 μM IKKβ
inhibitor VIII for 24 hours. Shown below are relative foldchanges + SD from n=3 replicates of
IRF4 or BATF vs GAPDH load control densitometry values. Levels in vehicle control treated WT
LMP1 expressing cells were set to 1. P-values were determined by one-sided Fisher's exact
test. **p<0.001, ***p<0.0001.

(D) Immunoblot analysis of WCL from Akata cells induced for the indicated LMP1 construct
expression, either without or together with an IκBα super-repressor (IκBα-S.R.) that blocks
canonical NF-κB signaling. Shown below are relative foldchanges + SD of IRF4 or BATF vs

GAPDH load control densitometry values from n=3 replicates, with values in cells expressing
WT LMP1 but not IκBα- set to 1. P-values were determined by one-sided Fisher's exact test.
**p<0.001, ***p<0.0001.

1171

Fig. 8. LMP1 TES1 and TES2 roles in EBV super-enhancer target gene regulation in
GM12878 LCLs.

(A) Schematic diagram of typical LCL enhancers vs super-enhancers. Super-enhancers have
significantly broader and taller histone 3 lysine 27 acetyl (H3K27Ac) peaks. EBV SE are host
genomic enhancer sites bound by all five LMP1-activated NF-κB transcription factor subunits,
EBNA-2, LP, 3A, and 3C.

(B) Volcano plot analysis of mRNA values in GM12878 expressing LMP1 vs control sgRNAs as
in **Fig. 1**. Genes targeted by EBV super-enhancers (SE) are highlighted by red circles, whereas
other LCL genes are indicated by blue circles. Genes more highly expressed with LMP1 KO
have higher x-axis values, whereas those downmodulated by LMP1 KO have lower values. P
value <0.05 and >2-fold gene expression cutoffs were used.

(C) Volcano plot analysis of mRNA values in GM12878 expressing LMP1 sgRNA with TES1m
versus WT LMP1 cDNA rescue, as in **Fig. 2-3**. Genes targeted by EBV super-enhancers (SE)
are highlighted by red circles, whereas other LCL genes are indicated by blue circles. Genes
more highly expressed with endogenous LMP1 KO and TES1m rescue have higher x-axis
values, whereas those more highly expressed with WT LMP1 rescue have lower values. P value
<0.05 and >2-fold gene expression cutoffs were used.

(D) Volcano plot analysis of mRNA values in GM12878 expressing LMP1 sgRNA with TES2m
versus WT LMP1 cDNA rescue, as in (C). Genes targeted by EBV super-enhancers (SE) are
highlighted by red circles, whereas other LCL genes are indicated by blue circles. Genes more
highly expressed with endogenous LMP1 KO and TES2m rescue have higher x-axis values,
whereas those more highly expressed with WT LMP1 rescue have lower values. P value <0.05
and >2-fold gene expression cutoffs were used.

1195

Fig. S11. LMP1 TES1 and TES2 roles in EBV super-enhancer target gene regulation in
Akata and BL-41 Burkitt B-cells.

(A) Volcano plot analysis of Akata RNA-seq, comparing mRNA values in cells induced for LMP1
WT vs. mock-induced for 24 hours. SE targeted genes highlighted by red circles and other B-
cell genes indicated by blue circles. P value <0.05 and >2-fold gene expression cutoffs were
used.

(B) Volcano plot analysis of Akata RNA-seq, comparing mRNA values in cells induced for LMP1
TES1m vs. mock-induced for 24 hours. SE targeted genes highlighted by red circles and other
B-cell genes indicated by blue circles. P value <0.05 and >2-fold gene expression cutoffs were
used.

(C) Volcano plot analysis of Akata RNA-seq, comparing mRNA values in cells induced for LMP1
TES2m vs. mock-induced for 24 hours. SE targeted genes highlighted by red circles and other
B-cell genes indicated by blue circles. P value <0.05 and >2-fold gene expression cutoffs were
used.

(D) Volcano plot analysis of BL-41 RNA-seq, comparing mRNA values in cells induced for LMP1
WT vs. mock-induced for 24 hours. SE targeted genes highlighted by red circles and other B-
cell genes indicated by blue circles. P value <0.05 and >2-fold gene expression cutoffs were
used.

(E) Volcano plot analysis of BL-41 RNA-seq, comparing mRNA values in cells induced for LMP1
TES1m vs. mock-induced for 24 hours. SE targeted genes highlighted by red circles and other
B-cell genes indicated by blue circles. P value <0.05 and >2-fold gene expression cutoffs were
used.

(F) Volcano plot analysis of BL-41 RNA-seq, comparing mRNA values in cells induced for LMP1
TES2m vs. mock-induced for 24 hours. SE targeted genes highlighted by red circles and other
B-cell genes indicated by blue circles. P value <0.05 and >2-fold gene expression cutoffs were
used.

1222

1223   **Fig. 9.** Model highlighting different modes of LMP1 TES1 and TES2 cross-talk in B-cell target
1224   gene regulation.

1225

Materials and Methods

Cell lines, culture, and vectors. HEK293T cells were purchased from ATCC and cultured in Dulbecco's modified Eagle medium (DMEM, Gibco) with 10% fetal bovine serum (FBS, Gibco). EBV-negative Akata and BL-41 cells were obtained from Elliott Kieff; GM12878 were purchased from Coriell. Mutu I was obtained from Jeff Sample and Jijoye was purchased from ATCC. All B-cell lines stably expressed *Streptococcus pyogenes* Cas9 and were grown in Roswell Park Memorial Institute (RPMI) 1640 (Life Technologies) with 10% fetal bovine serum (Gibco) and penicillin-streptomycin in a humidified chamber with 5% carbon dioxide. LMP1 wildtype, TES1 alanine point mutant 204PQQAT208 -> AQQAT, TES2 384YYD386->ID mutant and double mutant LMP1 with both AQQAT and ID mutations were cloned into the pLIX-402 vector. pLIX-402 uses a TET-On TRE promoter to drive transgene expression and a C-terminal HA-tag fusion. Lentivirus vectors were used to establish stable Cas9+/GM12878, Cas9+/EBV- Akata and Cas9+/EBV- BL-41 Burkitt cells. Cell lines were then maintained with 0.5 µg/mL puromycin or 25 µg/mL hygromycin. For LMP1 inducible expression studies, 0.5X10^6 cells/ml were plated on Day-1 in 2ml of fresh RPMI in a 12-well plate. Cell were treated with 250ng/ml (in Burkitt cell models) for 24 hours prior to sample collection for downstream analyses or with 400ng/ml doxycycline (in LCL LMP1 cDNA rescue model) (Sigma #D9891) to allow for LMP1 rescue upon CRISPR knockout of LMP1. The IκBα Super-repressor (SR) lacking residues 1-67 has previously been reported (27).

Antibodies and Reagents. Cell Signaling Technology (CST) TRAF1 (#4715, Rabbit mAb), p105/50 (#3035, Rabbit mAb), RelA (#8242, Rabbit mAb), phospho-RelA (Ser536) (3033, Rabbit mAb), RelB (#4922, Rabbit mAb), cRel (#4727, Rabbit mAb), IκBα (#9247, Mouse mAb), IRF4 (#4964, Rabbit mAb), TAK1 (#4505, Rabbit mAb), V5 (#13202, Rabbit mAb), HRP-linked anti-mouse IgG(7076), FLIP (#8510, Rabbit mAb), HRP-linked anti-rabbit IgG (#7074) were used in this study at 1:1000 dilution. p100/52 (EMD Millipore #05-361, Mouse mAb, 1:1000), GAPDH (EMD Millipore #MAB374, Mouse mAb, 1:500) was used. S12 mouse monoclonal antibody against LMP1 was purified from hybridoma supernatant (115). The IKKβ inhibitor IKK-2 inhibitor VIII (ApexBio, #A3485), puromycin dihydrochloride (Thermo Fisher #A1113803), and hygromycin B (Millipore #400052).

Growth Curve Analysis: For growth curve analysis, cells were counted and then normalized to the same starting concentration, using the CellTiterGlo (CTG) luciferase assay (Promega, Cat#G7570). Live cell numbers were quantitated at each timepoint by CTG measurements, and values were corrected for tissue culture passage. Fold change of live cell number at each timepoint was calculated as a ratio of the value divided by the input value. For the Caspase-Glo 3/6 Assay (Promega #G8092), Caspase-Glo 3/7 reagent was added to the cells, mixed and incubated for 30 minutes followed by recording of the luminescence. Readings were normalized to respective CTG values of the samples that were performed and collected concurrently.

CRISPR/Cas9 editing. B-cell lines with stable Cas9 expression were established as described previously (47). Briefly, HEK293T cells were plated at a density of 300,000 cells per well in 2 mL DMEM supplemented with 10% FBS on Day -1. The following day (Day 0) plated cells were transfected with the TransIT-LT1 Transfection Reagent (Mirus #2306), according to the manufacturer's protocol. Transfection media was replaced by RPMI 16 hours later (Day 1). B-cells were plated at 1.2X 10^6 density in a 6-well plate on Day 1. Lentivirus collected on Day 2 were added to the B-cells for spinoculation at 2000rpm for 2 hours at 37C and 4µg/ml of polybrene. Spinoculated cells were placed in in a humidified chamber with 5% carbon dioxide for 6 hours, then pelleted and resuspended in fresh RPMI/FBS. 48 hours post-transduction, transduced cells were selected by addition of puromycin 3 µg/ml or 200 µg/ml hygromycin. Broad Institute pXPR-515 control sgRNA (targets a non-coding intergenic region), Avana or

Brunello library sgRNAs, as listed in Table 1, were cloned into lentiGuide-Puro (Addgene,
catalog #52963) or pLenti SpBsmBI sgRNA Hygro (Addgene, catalog #62205).

RNAseq. Total RNA was isolated using RNeasy mini kit (Qiagen #74106) with in-column
genomic DNA digestion step (RNAse-free DAase set, Qiagen #79254) according to the
manufacturer's protocol. To construct indexed libraries, 1 µg of total RNA was used for polyA
mRNA selection using NEBNext Poly(A) mRNA Magnetic Isolation Module (Cat#E7490S), and
library preparation with NEBNext Ultra RNA Library Prep with Sample Purification Beads
(Cat#E7765S). Each experimental treatment was performed in biological triplicate. Libraries
were multi-indexed (NEB 7335L and E7500S), pooled and sequenced on an Illumina NextSeq
500 sequencer using single 75 bp read length. Adaptor-trimmed Illumina reads for each
individual library were mapped back to the human GRCh37.83 transcriptome assembly using
STAR2.5.2b (116). FeatureCounts was used to estimate the number of reads mapped to each
contig (117). Only transcripts with at least 5 cumulative mapping counts were used in this
analysis. DESeq2 was used to evaluate differential expression (DE) (118). DESeq2 uses a
negative binomial distribution to account for overdispersion in transcriptome data sets. It is
conservative and uses a heuristic approach to detect outliers while avoiding false positives.
Each DE analysis was composed of a pairwise comparison between experimental group and
the control group. Differentially expressed genes were identified after a correction for false
discovery rate (FDR). For more stringent analyses, we set the cutoff for truly differentially
expressed genes as adjusted p value (FDR corrected) < 0.05 and absolute fold change > 1.5.
DE genes meeting this cutoff were selected and subject to downstream bioinformatics and
functional analyses, including clustering, data visualization, GO annotation and pathway
analysis. DE genes were also subjected to Enrichr analysis (https://maayanlab.cloud/Enrichr/)
for pathway analysis. Heatmaps were generated by feeding the Variance-Stabilizing
Transformed values of selected DE genes from DESeq2 into Morpheus
(https://software.broadinstitute.org/morpheus/).

Quantitative real-time qRT-PCR Analysis. Total RNA was isolated using RNeasy mini kit
(Qiagen #74106) with in-column genomic DNA digestion step (RNAse-free DAase set, Qiagen
#79254) according to the manufacturer's protocol. Reverse transcription was performed with
400ng of total RNA using iScript Reverse Transcription supermix (Bio-Rad #1708841) in a 20ul
reaction. The cDNA mixture was diluted 1:20 and 4ul of the diluted cDNA was taken to perform
qPCR using the Power SYBR green PCR master mix (Fisher Scientific #4368708) in CFX96
Touch™ Real-Time PCR Detection System (Bio-Rad). Data was normalized to internal control
18s RNA levels. Relative expression was calculated using 2-ΔΔCt method. All samples were
run in technical triplicates and at least three independent experiments were performed. Primer
sequences are outlined in Table 2 below.

Immunoblot Analysis. Cells were lysed in Laemelli buffer (0.2M Tris-HCL, 0.4 M Dithiothreitol,
277mM SDS, 6mM Bromophenol blue and 10% glycerol v/v) and sonicated at 4℃ for five
seconds using a probe sonicator at 20% amplitude and boiled at 95℃ for eight minutes. The
whole cell lysates were resolved by 12% or 15% SDS-PAGE, transferred to nitrocellulose filters
at 100V at 4℃ for 1.5 hour, blocked with 5% non-fat dried milk in 1X TBST for 30 min at room-
temperature, and then probed with the indicated primary antibodies (diluted in 1x TBS-T with
0.02% sodium azide at recommended manufacturer concentrations) overnight at 4℃ on a
rotating platform. Blots were washed three times in TBST for ten minutes each, and then probed
with horse-radish peroxidase (HRP)-conjugated secondary antibodies, at a dilution of 1:3000 in
1X TBST with 5% non-fat dried milk. Blots were then washed three times in TBST for ten
minutes each, developed by ECL chemiluminescent substrate (Thermo Scientific, #34578) and
imaged on Li-COR Odyssey workstation.

Flow cytometry analysis. FACS was performed using a FACSCalibur instrument (BD). For
ICAM-1 and FAS detection, cells were washed in PBS w/2%FBS, then stained on ice for 30 min
with BioLegend PE-conjugated anti-CD54/ICAM-1 and APC-conjugated anti-CD95/Fas
antibody, washed three times with PBS with 2%FBS and analyzed by FACS. 7-AAD viability
assays were carried out suing 7-AAD (Thermo Fisher, #A1310) where cells were harvested and
washed twice in 1XPBS supplemented with 2% FBS (Gibco). Washed cells were incubated with
1ug/ml 7-AAD solution in 1X PBS/2%FBS buffer for five minutes at room temperature and
protected from light. Stained cells were analyzed via flow cytometry. Annexin V assay was
performed with harvesting 1 X 10^6 cells and washed with 1X PBS twice to remove excess
RPMI. 2 X10^5 cells were then resuspended to be stained with 5ul of Annexin V-FITC
(#640945, Biolegend) in 100ul of Annexin V binding buffer (10mM HEPES, 140mM NaCl and
2.5mM CaCl$_2$). Cells were incubated at room temperature for 15 minutes and protected from
light before analyzed by FACS.

Bioinformatic analysis and Software: All the growth curves and column charts were made
with GraphPad Prism v.9. FACS data was analyzed by FlowJo V10.

Table 1. sgRNAs used in this study.

Guide No.	Gene Target	sgRNA Sequence (5' -> 3')
#1	Control	TTGACCTTTACCGTCCCGCG
#1	LMP1	TCTATCTACAACAAAACTGG
#1	TAK1	CACCGGCTTACTGCTGGTTGCAGGG
#2	TAK1	CACCGCGCAATGAGTTGGTGTTTAC

1338

1339 **Table 2. RT-PCR primers used in this study.**

Gene Name	Primer Sequence(5' -> 3')	Primer Sequence (3' -> 5')
CCL22	CGCGTGGTGAAACACTTCTA	GGATCGGCACAGATCTCCT
EBI3	GATCCGTTACAAGCGTCAGG	ACGTAGTACCTGGCTCGGG
IRF4	ACAGCAGTTCTTGTCAGAG	GAGGTTCTACGTGAGCTG
18s	CCTGCGGCTTAATTTGACTC	AACCAGACAAATCGCTCCAC

1340

References

References

1. Gewurz BE, Longnecker, R & Cohen, JI. Epstein-Barr Virus. Fields Virology. 7 ed. pages 324-388.
Howley, PM and Knipe, DM, editors: Wolters Kluwer; 2021.

2. Stunz LL, Bishop GA. 2014. Latent membrane protein 1 and the B lymphocyte-a complex
relationship. Crit Rev Immunol 34:177-98.

3. Wang LW, Jiang S, Gewurz BE. 2017. Epstein-Barr Virus LMP1-Mediated Oncogenicity. J Virol 91.

4. Kieser A, Sterz KR. 2015. The Latent Membrane Protein 1 (LMP1). Curr Top Microbiol Immunol
391:119-49.

5. Wang L, Ning S. 2021. New Look of EBV LMP1 Signaling Landscape. Cancers (Basel) 13.

6. Fish K, Comoglio F, Shaffer AL, 3rd, Ji Y, Pan KT, Scheich S, Oellerich A, Doebele C, Ikeda M,
Schaller SJ, Nguyen H, Muppidi J, Wright GW, Urlaub H, Serve H, Staudt LM, Longnecker R,
Oellerich T. 2020. Rewiring of B cell receptor signaling by Epstein-Barr virus LMP2A. Proc Natl
Acad Sci U S A 117:26318-26327.

7. Weniger MA, Küppers R. 2021. Molecular biology of Hodgkin lymphoma. Leukemia 35:968-981.

8. Soni V, Cahir-McFarland E, Kieff E. 2007. LMP1 TRAFficking activates growth and survival
pathways. Adv Exp Med Biol 597:173-87.

9. Kaykas A, Worringer K, Sugden B. 2002. LMP-1's transmembrane domains encode multiple
functions required for LMP-1's efficient signaling. J Virol 76:11551-60.

10. Rastelli J, Hömig-Hölzel C, Seagal J, Müller W, Hermann AC, Rajewsky K, Zimber-Strobl U. 2008.
LMP1 signaling can replace CD40 signaling in B cells in vivo and has unique features of inducing
class-switch recombination to IgG1. Blood 111:1448-55.

11. Meckes DG, Jr., Menaker NF, Raab-Traub N. 2013. Epstein-Barr virus LMP1 modulates lipid raft
microdomains and the vimentin cytoskeleton for signal transduction and transformation. J Virol
87:1301-11.

12. Shair KH, Bendt KM, Edwards RH, Bedford EC, Nielsen JN, Raab-Traub N. 2007. EBV latent
membrane protein 1 activates Akt, NFkappaB, and Stat3 in B cell lymphomas. PLoS Pathog
3:e166.

13. Greenfeld H, Takasaki K, Walsh MJ, Ersing I, Bernhardt K, Ma Y, Fu B, Ashbaugh CW, Cabo J,
Mollo SB, Zhou H, Li S, Gewurz BE. 2015. TRAF1 Coordinates Polyubiquitin Signaling to Enhance
Epstein-Barr Virus LMP1-Mediated Growth and Survival Pathway Activation. PLoS Pathog
11:e1004890.

14. Kung CP, Meckes DG, Jr., Raab-Traub N. 2011. Epstein-Barr virus LMP1 activates EGFR, STAT3,
and ERK through effects on PKCdelta. J Virol 85:4399-408.

15. Mainou BA, Everly DN, Jr., Raab-Traub N. 2007. Unique signaling properties of CTAR1 in LMP1-
mediated transformation. J Virol 81:9680-92.

16. Huen DS, Henderson SA, Croom-Carter D, Rowe M. 1995. The Epstein-Barr virus latent
membrane protein-1 (LMP1) mediates activation of NF-kappa B and cell surface phenotype via
two effector regions in its carboxy-terminal cytoplasmic domain. Oncogene 10:549-60.

17. Zhang L, Wu L, Hong K, Pagano JS. 2001. Intracellular signaling molecules activated by Epstein-
Barr virus for induction of interferon regulatory factor 7. J Virol 75:12393-401.

18. Floettmann JE, Rowe M. 1997. Epstein-Barr virus latent membrane protein-1 (LMP1) C-terminus
activation region 2 (CTAR2) maps to the far C-terminus and requires oligomerisation for NF-
kappaB activation. Oncogene 15:1851-8.

19. Voigt S, Sterz KR, Giehler F, Mohr AW, Wilson JB, Moosmann A, Kieser A. 2020. A central role of
IKK2 and TPL2 in JNK activation and viral B-cell transformation. Nat Commun 11:685.

20.   Shkoda A, Town JA, Griese J, Romio M, Sarioglu H, Knöfel T, Giehler F, Kieser A. 2012. The
1387   germinal center kinase TNIK is required for canonical NF-κB and JNK signaling in B-cells by the
1388   EBV oncoprotein LMP1 and the CD40 receptor. PLoS Biol 10:e1001376.
1389   21.   Schultheiss U, Püschner S, Kremmer E, Mak TW, Engelmann H, Hammerschmidt W, Kieser A.
1390   2001. TRAF6 is a critical mediator of signal transduction by the viral oncogene latent membrane
1391   protein 1. Embo j 20:5678-91.
1392   22.   Arcipowski KM, Stunz LL, Bishop GA. 2014. TRAF6 is a critical regulator of LMP1 functions in vivo.
1393   Int Immunol 26:149-58.
1394   23.   Arcipowski KM, Stunz LL, Graham JP, Kraus ZJ, Vanden Bush TJ, Bishop GA. 2011. Molecular
1395   mechanisms of TNFR-associated factor 6 (TRAF6) utilization by the oncogenic viral mimic of
1396   CD40, latent membrane protein 1 (LMP1). J Biol Chem 286:9948-55.
1397   24.   Wang L, Howell MEA, Sparks-Wallace A, Zhao J, Hensley CR, Nicksic CA, Horne SR, Mohr KB,
1398   Moorman JP, Yao ZQ, Ning S. 2021. The Ubiquitin Sensor and Adaptor Protein p62 Mediates
1399   Signal Transduction of a Viral Oncogenic Pathway. mBio 12:e0109721.
1400   25.   Ma Y, Walsh MJ, Bernhardt K, Ashbaugh CW, Trudeau SJ, Ashbaugh IY, Jiang S, Jiang C, Zhao B,
1401   Root DE, Doench JG, Gewurz BE. 2017. CRISPR/Cas9 Screens Reveal Epstein-Barr Virus-
1402   Transformed B Cell Host Dependency Factors. Cell Host Microbe 21:580-591.e7.
1403   26.   Gewurz BE, Towfic F, Mar JC, Shinners NP, Takasaki K, Zhao B, Cahir-McFarland ED,
1404   Quackenbush J, Xavier RJ, Kieff E. 2012. Genome-wide siRNA screen for mediators of NF-κB
1405   activation. Proc Natl Acad Sci U S A 109:2467-72.
1406   27.   Gewurz BE, Mar JC, Padi M, Zhao B, Shinners NP, Takasaki K, Bedoya E, Zou JY, Cahir-McFarland
1407   E, Quackenbush J, Kieff E. 2011. Canonical NF-kappaB activation is essential for Epstein-Barr
1408   virus latent membrane protein 1 TES2/CTAR2 gene regulation. J Virol 85:6764-73.
1409   28.   Bentz GL, Moss CR, 2nd, Whitehurst CB, Moody CA, Pagano JS. 2015. LMP1-Induced
1410   Sumoylation Influences the Maintenance of Epstein-Barr Virus Latency through KAP1. J Virol
1411   89:7465-77.
1412   29.   Selby TL, Biel N, Varn M, Patel S, Patel A, Hilding L, Ray A, Ross T, Cramblet WT, Moss CR, Lowrey
1413   AJ, Bentz GL. 2019. The Epstein-Barr Virus Oncoprotein, LMP1, Regulates the Function of SENP2,
1414   a SUMO-protease. Sci Rep 9:9523.
1415   30.   Gires O, Kohlhuber F, Kilger E, Baumann M, Kieser A, Kaiser C, Zeidler R, Scheffer B, Ueffing M,
1416   Hammerschmidt W. 1999. Latent membrane protein 1 of Epstein-Barr virus interacts with JAK3
1417   and activates STAT proteins. Embo j 18:3064-73.
1418   31.   Izumi KM, Cahir McFarland ED, Riley EA, Rizzo D, Chen Y, Kieff E. 1999. The residues between the
1419   two transformation effector sites of Epstein-Barr virus latent membrane protein 1 are not
1420   critical for B-lymphocyte growth transformation. J Virol 73:9908-16.
1421   32.   Zhao B, Barrera LA, Ersing I, Willox B, Schmidt SC, Greenfeld H, Zhou H, Mollo SB, Shi TT,
1422   Takasaki K, Jiang S, Cahir-McFarland E, Kellis M, Bulyk ML, Kieff E, Gewurz BE. 2014. The NF-κB
1423   genomic landscape in lymphoblastoid B cells. Cell Rep 8:1595-606.
1424   33.   Wang D, Liebowitz D, Kieff E. 1985. An EBV membrane protein expressed in immortalized
1425   lymphocytes transforms established rodent cells. Cell 43:831-40.
1426   34.   Baichwal VR, Sugden B. 1988. Transformation of Balb 3T3 cells by the BNLF-1 gene of Epstein-
1427   Barr virus. Oncogene 2:461-7.
1428   35.   Moorthy RK, Thorley-Lawson DA. 1993. All three domains of the Epstein-Barr virus-encoded
1429   latent membrane protein LMP-1 are required for transformation of rat-1 fibroblasts. J Virol
1430   67:1638-46.
1431   36.   Mainou BA, Everly DN, Jr., Raab-Traub N. 2005. Epstein-Barr virus latent membrane protein 1
1432   CTAR1 mediates rodent and human fibroblast transformation through activation of PI3K.
1433   Oncogene 24:6917-24.

37. Minamitani T, Ma Y, Zhou H, Kida H, Tsai CY, Obana M, Okuzaki D, Fujio Y, Kumanogoh A, Zhao B,
Kikutani H, Kieff E, Gewurz BE, Yasui T. 2017. Mouse model of Epstein-Barr virus LMP1- and
LMP2A-driven germinal center B-cell lymphoproliferative disease. Proc Natl Acad Sci U S A
114:4751-4756.
38. Wirtz T, Weber T, Kracker S, Sommermann T, Rajewsky K, Yasuda T. 2016. Mouse model for
acute Epstein-Barr virus infection. Proc Natl Acad Sci U S A 113:13821-13826.
39. Zhang B, Kracker S, Yasuda T, Casola S, Vanneman M, Hömig-Hölzel C, Wang Z, Derudder E, Li S,
Chakraborty T, Cotter SE, Koyama S, Currie T, Freeman GJ, Kutok JL, Rodig SJ, Dranoff G,
Rajewsky K. 2012. Immune surveillance and therapy of lymphomas driven by Epstein-Barr virus
protein LMP1 in a mouse model. Cell 148:739-51.
40. Thornburg NJ, Kulwichit W, Edwards RH, Shair KH, Bendt KM, Raab-Traub N. 2006. LMP1
signaling and activation of NF-kappaB in LMP1 transgenic mice. Oncogene 25:288-97.
41. Pich D, Mrozek-Gorska P, Bouvet M, Sugimoto A, Akidil E, Grundhoff A, Hamperl S, Ling PD,
Hammerschmidt W. 2019. First Days in the Life of Naive Human B Lymphocytes Infected with
Epstein-Barr Virus. mBio 10.
42. Kaye KM, Izumi KM, Kieff E. 1993. Epstein-Barr virus latent membrane protein 1 is essential for
B-lymphocyte growth transformation. Proc Natl Acad Sci U S A 90:9150-4.
43. Dirmeier U, Neuhierl B, Kilger E, Reisbach G, Sandberg ML, Hammerschmidt W. 2003. Latent
membrane protein 1 is critical for efficient growth transformation of human B cells by epstein-
barr virus. Cancer Res 63:2982-9.
44. Cohen JI, Wang F, Kieff E. 1991. Epstein-Barr virus nuclear protein 2 mutations define essential
domains for transformation and transactivation. J Virol 65:2545-54.
45. Kaye KM, Izumi KM, Mosialos G, Kieff E. 1995. The Epstein-Barr virus LMP1 cytoplasmic carboxy
terminus is essential for B-lymphocyte transformation; fibroblast cocultivation complements a
critical function within the terminal 155 residues. J Virol 69:675-83.
46. Izumi KM, Kaye KM, Kieff ED. 1997. The Epstein-Barr virus LMP1 amino acid sequence that
engages tumor necrosis factor receptor associated factors is critical for primary B lymphocyte
growth transformation. Proc Natl Acad Sci U S A 94:1447-52.
47. Jiang S, Wang LW, Walsh MJ, Trudeau SJ, Gerdt C, Zhao B, Gewurz BE. 2018. CRISPR/Cas9-
Mediated Genome Editing in Epstein-Barr Virus-Transformed Lymphoblastoid B-Cell Lines. Curr
Protoc Mol Biol 121:31.12.1-31.12.23.
48. Kuleshov MV, Jones MR, Rouillard AD, Fernandez NF, Duan Q, Wang Z, Koplev S, Jenkins SL,
Jagodnik KM, Lachmann A, McDermott MG, Monteiro CD, Gundersen GW, Ma'ayan A. 2016.
Enrichr: a comprehensive gene set enrichment analysis web server 2016 update. Nucleic Acids
Res 44:W90-7.
49. Forte E, Luftig MA. 2009. MDM2-dependent inhibition of p53 is required for Epstein-Barr virus
B-cell growth transformation and infected-cell survival. J Virol 83:2491-9.
50. Cahir-McFarland ED, Carter K, Rosenwald A, Giltnane JM, Henrickson SE, Staudt LM, Kieff E.
2004. Role of NF-kappa B in cell survival and transcription of latent membrane protein 1-
expressing or Epstein-Barr virus latency III-infected cells. J Virol 78:4108-19.
51. Nakayama T, Hieshima K, Nagakubo D, Sato E, Nakayama M, Kawa K, Yoshie O. 2004. Selective
induction of Th2-attracting chemokines CCL17 and CCL22 in human B cells by latent membrane
protein 1 of Epstein-Barr virus. J Virol 78:1665-74.
52. Jorapur A, Marshall LA, Jacobson S, Xu M, Marubayashi S, Zibinsky M, Hu DX, Robles O, Jackson
JJ, Baloche V, Busson P, Wustrow D, Brockstedt DG, Talay O, Kassner PD, Cutler G. 2022. EBV+
tumors exploit tumor cell-intrinsic and -extrinsic mechanisms to produce regulatory T cell-
recruiting chemokines CCL17 and CCL22. PLoS Pathog 18:e1010200.

53.   Demetriades C, Mosialos G. 2009. The LMP1 promoter can be transactivated directly by NF-
kappaB. J Virol 83:5269-77.
54.   Johansson P, Jansson A, Rüetschi U, Rymo L. 2009. Nuclear factor-kappaB binds to the Epstein-
Barr Virus LMP1 promoter and upregulates its expression. J Virol 83:1393-401.
55.   Devergne O, Cahir McFarland ED, Mosialos G, Izumi KM, Ware CF, Kieff E. 1998. Role of the TRAF
binding site and NF-kappaB activation in Epstein-Barr virus latent membrane protein 1-induced
cell gene expression. J Virol 72:7900-8.
56.   Izumi KM, Kieff ED. 1997. The Epstein-Barr virus oncogene product latent membrane protein 1
engages the tumor necrosis factor receptor-associated death domain protein to mediate B
lymphocyte growth transformation and activate NF-kappaB. Proc Natl Acad Sci U S A 94:12592-
7.
57.   Lee DY, Sugden B. 2008. The LMP1 oncogene of EBV activates PERK and the unfolded protein
response to drive its own synthesis. Blood 111:2280-9.
58.   Bentz GL, Whitehurst CB, Pagano JS. 2011. Epstein-Barr virus latent membrane protein 1 (LMP1)
C-terminal-activating region 3 contributes to LMP1-mediated cellular migration via its
interaction with Ubc9. J Virol 85:10144-53.
59.   Luftig M, Yasui T, Soni V, Kang MS, Jacobson N, Cahir-McFarland E, Seed B, Kieff E. 2004. Epstein-
Barr virus latent infection membrane protein 1 TRAF-binding site induces NIK/IKK alpha-
dependent noncanonical NF-kappaB activation. Proc Natl Acad Sci U S A 101:141-6.
60.   Tsai K, Thikmyanova N, Wojcechowskyj JA, Delecluse HJ, Lieberman PM. 2011. EBV tegument
protein BNRF1 disrupts DAXX-ATRX to activate viral early gene transcription. PLoS Pathog
7:e1002376.
61.   Zhang Y, Jiang C, Trudeau SJ, Narita Y, Zhao B, Teng M, Guo R, Gewurz BE. 2020. Histone Loaders
CAF1 and HIRA Restrict Epstein-Barr Virus B-Cell Lytic Reactivation. mBio 11.
62.   Takada K, Horinouchi K, Ono Y, Aya T, Osato T, Takahashi M, Hayasaka S. 1991. An Epstein-Barr
virus-producer line Akata: establishment of the cell line and analysis of viral DNA. Virus Genes
5:147-56.
63.   Shimizu N, Tanabe-Tochikura A, Kuroiwa Y, Takada K. 1994. Isolation of Epstein-Barr virus (EBV)-
negative cell clones from the EBV-positive Burkitt's lymphoma (BL) line Akata: malignant
phenotypes of BL cells are dependent on EBV. J Virol 68:6069-73.
64.   Anonymous. 1985. Burkitt's lymphoma: a human cancer model. Proceedings of a symposium.
Lyon, 6-9 December 1983. IARC Sci Publ:1-484.
65.   Song YJ, Izumi KM, Shinners NP, Gewurz BE, Kieff E. 2008. IRF7 activation by Epstein-Barr virus
latent membrane protein 1 requires localization at activation sites and TRAF6, but not TRAF2 or
TRAF3. Proc Natl Acad Sci U S A 105:18448-53.
66.   Huye LE, Ning S, Kelliher M, Pagano JS. 2007. Interferon regulatory factor 7 is activated by a viral
oncoprotein through RIP-dependent ubiquitination. Mol Cell Biol 27:2910-8.
67.   Ning S, Hahn AM, Huye LE, Pagano JS. 2003. Interferon regulatory factor 7 regulates expression
of Epstein-Barr virus latent membrane protein 1: a regulatory circuit. J Virol 77:9359-68.
68.   Ning S, Pagano JS. 2010. The A20 deubiquitinase activity negatively regulates LMP1 activation of
IRF7. J Virol 84:6130-8.
69.   Thomas N, Dreval K, Gerhard DS, Hilton LK, Abramson JS, Ambinder RF, Barta S, Bartlett NL,
Bethony J, Bhatia K, Bowen J, Bryan AC, Cesarman E, Casper C, Chadburn A, Cruz M, Dittmer DP,
Dyer MA, Farinha P, Gastier-Foster JM, Gerrie AS, Grande BM, Greiner T, Griner NB, Gross TG,
Harris NL, Irvin JD, Jaffe ES, Henry D, Huppi R, Leal FE, Lee MS, Martin JP, Martin MR, Mbulaiteye
SM, Mitsuyasu R, Morris V, Mullighan CG, Mungall AJ, Mungall K, Mutyaba I, Nokta M,
Namirembe C, Noy A, Ogwang MD, Omoding A, Orem J, Ott G, Petrello H, Pittaluga S, et al. 2023.

Genetic subgroups inform on pathobiology in adult and pediatric Burkitt lymphoma. Blood
141:904-916.

70.   Grande BM, Gerhard DS, Jiang A, Griner NB, Abramson JS, Alexander TB, Allen H, Ayers LW,
Bethony JM, Bhatia K, Bowen J, Casper C, Choi JK, Culibrk L, Davidsen TM, Dyer MA, Gastier-
Foster JM, Gesuwan P, Greiner TC, Gross TG, Hanf B, Harris NL, He Y, Irvin JD, Jaffe ES, Jones
SJM, Kerchan P, Knoetze N, Leal FE, Lichtenberg TM, Ma Y, Martin JP, Martin MR, Mbulaiteye
SM, Mullighan CG, Mungall AJ, Namirembe C, Novik K, Noy A, Ogwang MD, Omoding A, Orem J,
Reynolds SJ, Rushton CK, Sandlund JT, Schmitz R, Taylor C, Wilson WH, Wright GW, Zhao EY, et
al. 2019. Genome-wide discovery of somatic coding and noncoding mutations in pediatric
endemic and sporadic Burkitt lymphoma. Blood 133:1313-1324.

71.   Guo R, Zhang Y, Teng M, Jiang C, Schineller M, Zhao B, Doench JG, O'Reilly RJ, Cesarman E,
Giulino-Roth L, Gewurz BE. 2020. DNA methylation enzymes and PRC1 restrict B-cell Epstein-
Barr virus oncoprotein expression. Nat Microbiol 5:1051-1063.

72.   Wang F, Gregory C, Sample C, Rowe M, Liebowitz D, Murray R, Rickinson A, Kieff E. 1990.
Epstein-Barr virus latent membrane protein (LMP1) and nuclear proteins 2 and 3C are effectors
of phenotypic changes in B lymphocytes: EBNA-2 and LMP1 cooperatively induce CD23. J Virol
64:2309-18.

73.   Gregory CD, Rowe M, Rickinson AB. 1990. Different Epstein-Barr virus-B cell interactions in
phenotypically distinct clones of a Burkitt's lymphoma cell line. J Gen Virol 71 (Pt 7):1481-95.

74.   Jin J, Hu H, Li HS, Yu J, Xiao Y, Brittain GC, Zou Q, Cheng X, Mallette FA, Watowich SS, Sun SC.
2014. Noncanonical NF-κB pathway controls the production of type I interferons in antiviral
innate immunity. Immunity 40:342-54.

75.   DeKroon RM, Gunawardena HP, Edwards R, Raab-Traub N. 2018. Global Proteomic Changes
Induced by the Epstein-Barr Virus Oncoproteins Latent Membrane Protein 1 and 2A. mBio 9.

76.   Devergne O, Hatzivassiliou E, Izumi KM, Kaye KM, Kleijnen MF, Kieff E, Mosialos G. 1996.
Association of TRAF1, TRAF2, and TRAF3 with an Epstein-Barr virus LMP1 domain important for
B-lymphocyte transformation: role in NF-kappaB activation. Mol Cell Biol 16:7098-108.

77.   Shair KH, Raab-Traub N. 2012. Transcriptome changes induced by Epstein-Barr virus LMP1 and
LMP2A in transgenic lymphocytes and lymphoma. mBio 3.

78.   Faumont N, Durand-Panteix S, Schlee M, Grömminger S, Schuhmacher M, Hölzel M, Laux G,
Mailhammer R, Rosenwald A, Staudt LM, Bornkamm GW, Feuillard J. 2009. c-Myc and Rel/NF-
kappaB are the two master transcriptional systems activated in the latency III program of
Epstein-Barr virus-immortalized B cells. J Virol 83:5014-27.

79.   Wang L, Ren J, Li G, Moorman JP, Yao ZQ, Ning S. 2017. LMP1 signaling pathway activates IRF4 in
latent EBV infection and a positive circuit between PI3K and Src is required. Oncogene 36:2265-
2274.

80.   Wang L, Ning S. 2013. Interferon regulatory factor 4 is activated through c-Src-mediated tyrosine
phosphorylation in virus-transformed cells. J Virol 87:9672-9.

81.   Glasmacher E, Agrawal S, Chang AB, Murphy TL, Zeng W, Vander Lugt B, Khan AA, Ciofani M,
Spooner CJ, Rutz S, Hackney J, Nurieva R, Escalante CR, Ouyang W, Littman DR, Murphy KM,
Singh H. 2012. A genomic regulatory element that directs assembly and function of immune-
specific AP-1-IRF complexes. Science 338:975-80.

82.   Hnisz D, Abraham BJ, Lee TI, Lau A, Saint-André V, Sigova AA, Hoke HA, Young RA. 2013. Super-
enhancers in the control of cell identity and disease. Cell 155:934-47.

83.   Jiang S, Zhou H, Liang J, Gerdt C, Wang C, Ke L, Schmidt SCS, Narita Y, Ma Y, Wang S, Colson T,
Gewurz B, Li G, Kieff E, Zhao B. 2017. The Epstein-Barr Virus Regulome in Lymphoblastoid Cells.
Cell Host Microbe 22:561-573.e4.

84.   Zhou H, Schmidt SC, Jiang S, Willox B, Bernhardt K, Liang J, Johannsen EC, Kharchenko P, Gewurz
1576   BE, Kieff E, Zhao B. 2015. Epstein-Barr virus oncoprotein super-enhancers control B cell growth.
1577   Cell Host Microbe 17:205-16.
1578   85.   Yan B, Wang C, Chakravorty S, Zhang Z, Kadadi SD, Zhuang Y, Sirit I, Hu Y, Jung M, Sahoo SS,
1579   Wang L, Shao K, Anderson NL, Trujillo-Ochoa JL, Briggs SD, Liu X, Olson MR, Afzali B, Zhao B,
1580   Kazemian M. 2023. A comprehensive single cell data analysis of lymphoblastoid cells reveals the
1581   role of super-enhancers in maintaining EBV latency. J Med Virol 95:e28362.
1582   86.   Burton EM, Voyer J, Gewurz BE. 2022. Epstein-Barr virus latency programs dynamically sensitize
1583   B cells to ferroptosis. Proc Natl Acad Sci U S A 119:e2118300119.
1584   87.   Price AM, Tourigny JP, Forte E, Salinas RE, Dave SS, Luftig MA. 2012. Analysis of Epstein-Barr
1585   virus-regulated host gene expression changes through primary B-cell outgrowth reveals delayed
1586   kinetics of latent membrane protein 1-mediated NF-κB activation. J Virol 86:11096-106.
1587   88.   Edwards RH, Marquitz AR, Raab-Traub N. 2015. Changes in expression induced by Epstein-Barr
1588   Virus LMP1-CTAR1: potential role of bcl3. mBio 6.
1589   89.   Kung CP, Raab-Traub N. 2010. Epstein-Barr virus latent membrane protein 1 modulates
1590   distinctive NF- kappaB pathways through C-terminus-activating region 1 to regulate epidermal
1591   growth factor receptor expression. J Virol 84:6605-14.
1592   90.   Schneider F, Neugebauer J, Griese J, Liefold N, Kutz H, Briseno C, Kieser A. 2008. The viral
1593   oncoprotein LMP1 exploits TRADD for signaling by masking its apoptotic activity. PLoS Biol 6:e8.
1594   91.   Le Clorennec C, Youlyouz-Marfak I, Adriaenssens E, Coll J, Bornkamm GW, Feuillard J. 2006. EBV
1595   latency III immortalization program sensitizes B cells to induction of CD95-mediated apoptosis
1596   via LMP1: role of NF-kappaB, STAT1, and p53. Blood 107:2070-8.
1597   92.   Hanahan D, Weinberg RA. 2000. The hallmarks of cancer. Cell 100:57-70.
1598   93.   Morris MA, Laverick L, Wei W, Davis AM, O'Neill S, Wood L, Wright J, Dawson CW, Young LS.
1599   2018. The EBV-Encoded Oncoprotein, LMP1, Induces an Epithelial-to-Mesenchymal Transition
1600   (EMT) via Its CTAR1 Domain through Integrin-Mediated ERK-MAPK Signalling. Cancers (Basel) 10.
1601   94.   Kutz H, Reisbach G, Schultheiss U, Kieser A. 2008. The c-Jun N-terminal kinase pathway is critical
1602   for cell transformation by the latent membrane protein 1 of Epstein-Barr virus. Virology
1603   371:246-56.
1604   95.   Lambert SL, Martinez OM. 2007. Latent membrane protein 1 of EBV activates
1605   phosphatidylinositol 3-kinase to induce production of IL-10. J Immunol 179:8225-34.
1606   96.   Pratt ZL, Zhang J, Sugden B. 2012. The latent membrane protein 1 (LMP1) oncogene of Epstein-
1607   Barr virus can simultaneously induce and inhibit apoptosis in B cells. J Virol 86:4380-93.
1608   97.   Price AM, Dai J, Bazot Q, Patel L, Nikitin PA, Djavadian R, Winter PS, Salinas CA, Barry AP, Wood
1609   KC, Johannsen EC, Letai A, Allday MJ, Luftig MA. 2017. Epstein-Barr virus ensures B cell survival
1610   by uniquely modulating apoptosis at early and late times after infection. Elife 6.
1611   98.   Dirmeier U, Hoffmann R, Kilger E, Schultheiss U, Briseno C, Gires O, Kieser A, Eick D, Sugden B,
1612   Hammerschmidt W. 2005. Latent membrane protein 1 of Epstein-Barr virus coordinately
1613   regulates proliferation with control of apoptosis. Oncogene 24:1711-7.
1614   99.   Shaulian E, Karin M. 2002. AP-1 as a regulator of cell life and death. Nat Cell Biol 4:E131-6.
1615   100.   Johansen LM, Deppmann CD, Erickson KD, Coffin WF, 3rd, Thornton TM, Humphrey SE, Martin
1616   JM, Taparowsky EJ. 2003. EBNA2 and activated Notch induce expression of BATF. J Virol
1617   77:6029-40.
1618   101.   Banerjee S, Lu J, Cai Q, Saha A, Jha HC, Dzeng RK, Robertson ES. 2013. The EBV Latent Antigen 3C
1619   Inhibits Apoptosis through Targeted Regulation of Interferon Regulatory Factors 4 and 8. PLoS
1620   Pathog 9:e1003314.
1621   102.   Wang LW, Shen H, Nobre L, Ersing I, Paulo JA, Trudeau S, Wang Z, Smith NA, Ma Y, Reinstadler B,
1622   Nomburg J, Sommermann T, Cahir-McFarland E, Gygi SP, Mootha VK, Weekes MP, Gewurz BE.

2019. Epstein-Barr-Virus-Induced One-Carbon Metabolism Drives B Cell Transformation. Cell
Metab 30:539-555.e11.

103.   Guo R, Liang JH, Zhang Y, Lutchenkov M, Li Z, Wang Y, Trujillo-Alonso V, Puri R, Giulino-Roth L,
Gewurz BE. 2022. Methionine metabolism controls the B cell EBV epigenome and viral latency.
Cell Metab 34:1280-1297.e9.

104.   Bonglack EN, Messinger JE, Cable JM, Ch'ng J, Parnell KM, Reinoso-Vizcaíno NM, Barry AP,
Russell VS, Dave SS, Christofk HR, Luftig MA. 2021. Monocarboxylate transporter antagonism
reveals metabolic vulnerabilities of viral-driven lymphomas. Proc Natl Acad Sci U S A 118.

105.   Correia S, Palser A, Elgueta Karstegl C, Middeldorp JM, Ramayanti O, Cohen JI, Hildesheim A,
Fellner MD, Wiels J, White RE, Kellam P, Farrell PJ. 2017. Natural Variation of Epstein-Barr Virus
Genes, Proteins, and Primary MicroRNA. J Virol 91.

106.   Hu LF, Zabarovsky ER, Chen F, Cao SL, Ernberg I, Klein G, Winberg G. 1991. Isolation and
sequencing of the Epstein-Barr virus BNLF-1 gene (LMP1) from a Chinese nasopharyngeal
carcinoma. J Gen Virol 72 (Pt 10):2399-409.

107.   Chen ML, Tsai CN, Liang CL, Shu CH, Huang CR, Sulitzeanu D, Liu ST, Chang YS. 1992. Cloning and
characterization of the latent membrane protein (LMP) of a specific Epstein-Barr virus variant
derived from the nasopharyngeal carcinoma in the Taiwanese population. Oncogene 7:2131-40.

108.   Knecht H, Bachmann E, Brousset P, Sandvej K, Nadal D, Bachmann F, Odermatt BF, Delsol G,
Pallesen G. 1993. Deletions within the LMP1 oncogene of Epstein-Barr virus are clustered in
Hodgkin's disease and identical to those observed in nasopharyngeal carcinoma. Blood 82:2937-
42.

109.   Kingma DW, Weiss WB, Jaffe ES, Kumar S, Frekko K, Raffeld M. 1996. Epstein-Barr virus latent
membrane protein-1 oncogene deletions: correlations with malignancy in Epstein-Barr virus--
associated lymphoproliferative disorders and malignant lymphomas. Blood 88:242-51.

110.   Knecht H, Bachmann E, Joske DJ, Sahli R, Eméry-Goodman A, Casanova JL, Zilić M, Bachmann F,
Odermatt BF. 1993. Molecular analysis of the LMP (latent membrane protein) oncogene in
Hodgkin's disease. Leukemia 7:580-5.

111.   da Costa VG, Marques-Silva AC, Moreli ML. 2015. The Epstein-Barr virus latent membrane
protein-1 (LMP1) 30-bp deletion and XhoI-polymorphism in nasopharyngeal carcinoma: a meta-
analysis of observational studies. Syst Rev 4:46.

112.   Li SN, Chang YS, Liu ST. 1996. Effect of a 10-amino acid deletion on the oncogenic activity of
latent membrane protein 1 of Epstein-Barr virus. Oncogene 12:2129-35.

113.   Trivedi P, Winberg G, Klein G. 1997. Differential immunogenicity of Epstein-Barr virus (EBV)
encoded growth transformation-associated antigens in a murine model system. Eur J Cancer
33:912-7.

114.   Miller WE, Cheshire JL, Baldwin AS, Jr., Raab-Traub N. 1998. The NPC derived C15 LMP1 protein
confers enhanced activation of NF-kappa B and induction of the EGFR in epithelial cells.
Oncogene 16:1869-77.

115.   Mann KP, Staunton D, Thorley-Lawson DA. 1985. Epstein-Barr virus-encoded protein found in
plasma membranes of transformed cells. J Virol 55:710-20.

116.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR.
2012. STAR: ultrafast universal RNA-seq aligner. Bioinformatics 29:15-21.

117.   Liao Y, Smyth GK, Shi W. 2013. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. Bioinformatics 30:923-930.

118.   Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. Genome Biol 15:550.

1669

Fig. 1. Characterization of LMP1 KO effects on GM12878 LCL target gene regulation

(A)Immunoblot analysis of whole cell lysates (WCL) from GM12878 LCLs transduced with lentiviruses that express control or LMP1 targeting single guide RNAs (sgRNAs). Transduced cells were puromycin selected for 0 vs 2 days, as indicated. Blots for LMP1, for LMP1 target genes TRAF1 and IRF4, and for LMP1-driven non-canonical NF-κB pathway p100/p52 processing are shown. Blots are representative of n=3 experiments.

(B) K-means heatmap analysis of GM12878 LCLs transduced as in (A) with lentivirus expressing control or LMP1 sgRNA and puromycin selected for 48 hours. The heatmap depicts relative Z-scores in each row from n=3 independent RNAseq replicates, divided into two clusters. The Z-score scale is shown at bottom, where blue and red colors indicate lower versus higher relative expression, respectively. Two-way ANOVA P-value cutoff of <0.01 and >2-fold gene expression cutoffs were used.

(C) Enrichr analysis of KEGG pathways most highly changed in GM12878 expressing control versus LMP1 sgRNA (LMP1 KO), as in (A). The X-axis depicts the -Log10 adjusted p-value (adj p-value) scale. The top three most enriched KEGG pathways are shown.

(D) Abundances of two representative Cluster 1 genes from n=3 RNAseq analyses in cells with control vs LMP1 sgRNA. p-values were determined by one-sided Fisher's exact test. *p<0.05, **p<0.01.

(E) Abundances of two representative Cluster 2 genes from n=3 RNAseq analyses in cells with control vs LMP1 sgRNA. p-values were determined by one-sided Fisher's exact test. ***p<0.001.

(F) Volcano plot analysis of host transcriptome-wide GM12878 genes differentially expressed in cells with control versus LMP1 sgRNA expression, as in (B), using data from n=3 RNAseq datasets.

(G) Scatter plot cross comparison of log2 transformed fold change mRNA abundances in GM12878 expressing LMP1 vs. control sgRNA (Y-axis) versus log2 transformed fold change abundances of sgRNAs at Day 21 versus Day 1 post-transduction of GM12878 LCLs in a genome-wide CRISPR screen (25) (X-axis).

(H) String analysis of genes shown in (G). Pathway identifiers for each gene and interaction are colored coded.

(I) Volcano plot analysis of EBV mRNA values in GM12878 expressing LMP1 vs control sgRNAs, as in (B). P-value <0.05 and >2-fold change mRNA abundance cutoffs were used.

A

B

C

D

E

F

Fig. S1. Characterization of LMP1 KO effects on GM12878 LCL target gene regulation.

(A) Relative mean + standard deviation (SD) live cell numbers from CellTitreGlo analysis of n=3 replicates of Cas9+ GM12878 LCLs, transduced with lentiviruses that expressed control or LMP1 sgRNAs and puromycin selected, for 2 versus 4 days.

(B) RT-PCR analysis of CCL22 mRNA abundance in Cas9+ GM12878 post-transduction with lentiviruses that expressed control or LMP1 sgRNA and puromycin selected for 2 days, as in **Fig. 1B**. Values from cells with control sgRNA were set to 1, and mean fold-change of CCL22 mRNA abundance + SD in cells with LMP1 sgRNA are shown. p-values were determined by one-sided Fisher's exact test from two independent experiments, each with two technical replicates. ***p<0.001. The same RNA used for RNA-seq (**Fig. 1B**) was used for these qPCR experiments.

(C) RT-PCR analysis of EBI3 mRNA abundance in Cas9+ GM12878 post-transduction with lentiviruses that expressed control or LMP1 sgRNA and puromycin selected for 2 days, as in **Fig. 1B**. Values from cells with control sgRNA were set to 1, and mean fold-change of EBI3 mRNA abundance + SD in cells with LMP1 sgRNA are shown. p-values were determined by one-sided Fisher's exact test from two independent experiments, each with two technical replicates. ***p<0.001. The same RNA used for RNA-seq (**Fig. 1B**) was used for these qPCR experiments.

(D) RT-PCR analysis of IRF4 mRNA abundance in Cas9+ GM12878 post-transduction with lentiviruses that expressed control or LMP1 sgRNA and puromycin selected for 2 days, as in **Fig. 1B**. Values from cells with control sgRNA were set to 1, and mean fold-change of IRF4 mRNA abundance + SD in cells with LMP1 sgRNA are shown. p-values were determined by one-sided Fisher's exact test from two independent experiments, each with two technical replicates. ***p<0.001. The same RNA used for RNA-seq (**Fig. 1B**) was used for these qPCR experiments.

(E) Scatter plot analysis cross-comparing the significance of changes in LCL dependency factor expression upon GM127878 LMP1 KO versus the CRISPR screen significance score for selection against sgRNAs in LCL vs Burkitt dependency factor analysis (25). Shown on the Y-axis are -log10 transformed P-values from RNAseq analysis of GM12878 LCLs transduced with lentiviruses expressing LMP1 versus control sgRNA (as in **Fig. 1F**), versus -log10 transformed P-values from CRISPR LCL vs Burkitt cell dependency factor analysis (25). Higher Y-axis scores indicate more significant differences in expression for the indicated genes in GM12878 with LMP1 vs control sgRNA. Higher X-axis scores indicate a stronger selection against sgRNA targeting the indicated genes in GM12878 LCLs versus P3HR1 Burkitt cells over 21 days of cell culture. Shown are genes with p<0.05 in both analyses.

(F) Volcano plot analysis visualizing KEGG Hodgkin lymphoma pathway gene -Log10 (P-value) on the y-axis versus log2 transformed fold change in mRNA abundances on the x-axis of GM12878 genes in cells expressing LMP1 versus control sgRNA (as in **Fig. 1F**). P-value <0.05 and >2-fold change mRNA abundance cutoffs were used.

Fig. 2. **Loss of TES1 but not TES2 signaling triggers LCL apoptosis.**

(A) Schematic diagram of LMP1 WT with TES1 and TES2 domains highlighted. Wildtype (WT) or point mutants abrogated for signaling from TES1 (TES1m), TES2 (TES2m) or double TES1/TES2 mutant (DM) are shown.

(B) Immunoblot analysis of WCL from GM12878 LCLs that expressed control or LMP1 sgRNAs and puromycin selected for 3 days, then induced for expression with the indicated LMP1 rescue cDNA construct for 6 days. Blots are representative of n = 3 experiments.

(C) Growth curve analysis of GM12878 LCLs at the indicated day post expression of control or LMP1 sgRNAs and the indicated LMP1 WT, TES1m, TES2m or DM rescue cDNA. Shown are mean ± SD from n=3 independent experiments. **P<0.01.

(D) FACS analysis of 7-AAD vital dye uptake in GM12878 on day 7 post- expression of LMP1 sgRNAs and the indicated LMP1 rescue cDNA. Shown are percentages of 7-AAD+ cells within the indicated gates. Representative of n=3 experiments.

(E) Mean ± SD of fold change 7-AAD values from n=3 independent experiments of GM12878 with the indicated control or LMP1 sgRNA and rescue cDNA expression, as in (D). Values in GM12878 with control sgRNA and no LMP1 rescue cDNA were set to 1.

(F) FACS analysis of plasma membrane annexin V abundance in GM12878 on day 7 post- expression of control or LMP1 sgRNAs and the indicated LMP1 rescue cDNA. Shown are percentages of 7-AAD+ cells within the indicated gates. Representative of n=3 experiments.

(G) Mean ± SD of fold-change caspase 3/7 activity levels, as determined by caspase 3/7 Glo assay, from n=3 independent experiments of GM12878 with the indicated control or LMP1 sgRNA and rescue cDNA expression. Values in GM12878 with control sgRNA and no LMP1 rescue cDNA were set to 1.

Fig. S2. Loss of TES1 but not TES2 signaling triggers LCL apoptosis.

Mean ± SD of fold change plasma membrane annexin V values from n=3 independent experiments, using GM12878 with the indicated control or LMP1 sgRNA and rescue cDNA expression. Values in GM12878 with control sgRNA and no LMP1 rescue cDNA were set to 1.

Figure 3

Fig. 3. Characterization of host genome-wide TES1 vs TES2 LCL target genes

(A) RNAseq K-means heatmap analysis of GM12878 LCLs transduced with lentivirus expressing LMP1 sgRNA and induced for WT, TES1m or TES2m rescue cDNA expression for 6 days. The heatmap depicts relative Z-scores in each row from n=3 independent RNAseq datasets, divided into six clusters. The Z-score scale is shown at bottom, where blue and red colors indicate lower versus higher relative expression, respectively. Two-way ANOVA P-value cutoff of <0.05 and >2-fold gene expression cutoffs were used. The top three most highly enriched KEGG pathways amongst genes within each cluster are shown at right.

(B) Heatmap analysis of KEGG apoptosis pathway gene relative row Z-scores from RNAseq analysis as in (A). The Z-score scale is shown at bottom, where blue and red colors indicate lower versus higher relative expression, respectively. Two-way ANOVA P-value cutoff of <0.05 and >2-fold gene expression cutoffs were used.

(C) Scatter plot analysis cross comparing log2 transformed fold change of LCL dependency factor mRNA abundances in GM12878 expressing LMP1 sgRNA together with TES2 mutant versus wildtype cDNA rescue (Y-axis) and TES1 mutant versus wildtype cDNA rescue (X-axis) from triplicate RNAseq datasets, as in (A). This analysis highlighted that CFLAR and to a lesser extent NFKB2 and CCND2 mRNAs were more highly downmodulated by TES1m than TES2m rescue, relative to levels in cells with WT LMP1 rescue. Shown are genes differentially regulated by >2 fold with either TES1m or TES2m rescue, relative to levels with WT LMP1 rescue.

(D) Immunoblot analysis of c-FLIP and load control GAPDH expression in WCL from GM12878 LCLs with the indicated control or LMP1 sgRNA and LMP1 rescue cDNA expression. Representative of n=3 experiments.

Figure S3

Figure S3. Characterization of TES1 vs TES2 LCL dependency factor and Hodgkin lymphoma pathway targets.

(A) Heatmap analysis of CRISPR defined LCL dependency factor gene relative row Z-scores from RNAseq of GM12878 expressing LMP1 sgRNA and the indicated rescue cDNA, as in **Fig. 3**. The Z-score scale is shown at bottom, where blue and red colors indicate lower versus higher relative expression, respectively. Two-way ANOVA P-value cutoff of <0.05 and >2-fold gene expression cutoffs were used.

(B) Heatmap analysis of KEGG Hodgkin Lymphoma pathway gene relative row Z-scores from RNAseq of GM12878 expressing LMP1 sgRNA and the indicated rescue cDNA, as in **Fig. 3**. Two-way ANOVA P-value cutoff of <0.05 and >2-fold gene expression cutoffs were used.

(C) RT-PCR analysis of CCL22 mRNA abundance in GM12878 LCLs transduced with lentivirus expressing LMP1 sgRNA and induced for WT, TES1m or TES2m rescue cDNA expression for 6 days, as in Fig. 3A. The same RNA used for RNA-seq in Fig. 3A was used for these qPCR experiments. Values from cells with WT LMP1 rescue cDNA were set to 1, and mean fold-change of CCL22 mRNA abundance + SD in cells with TES1m or TES2m rescue cDNA are shown. p-values were determined by one-sided Fisher's exact test from two independent experiments, each with two technical replicates. ***$p<0.001$.

(D) RT-PCR analysis of EBI3 mRNA abundance in GM12878 LCLs transduced with lentivirus expressing LMP1 sgRNA and induced for WT, TES1m or TES2m rescue cDNA expression for 6 days, as in Fig. 3A. The same RNA used for RNA-seq in Fig. 3A was used for these qPCR experiments. Values from cells with WT LMP1 rescue cDNA were set to 1, and mean fold-change of EBI3 mRNA abundance + SD in cells with TES1m or TES2m rescue cDNA are shown. p-values were determined by one-sided Fisher's exact test from two independent experiments, each with two technical replicates. ***$p<0.001$.

(E) RT-PCR analysis of IRF4 mRNA abundance in GM12878 LCLs transduced with lentivirus expressing LMP1 sgRNA and induced for WT, TES1m or TES2m rescue cDNA expression for 6 days, as in **Fig. 3A**. The same RNA used for RNA-seq in Fig. 3A was used for these qPCR experiments. Values from cells with WT LMP1 rescue cDNA were set to 1, and mean fold-change of IRF4 mRNA abundance + SD in cells with TES1m or TES2m rescue cDNA are shown. p-values were determined by one-sided Fisher's exact test from two independent experiments, each with two technical replicates. ***$p<0.001$.

Figure 4

Fig. 4. Characterization of LCL pathways targeted by TES1 vs TES2 signaling

(A) Volcano plot analysis of host transcriptome-wide GM12878 genes differentially expressed in LMP1 KO GM12878 with WT vs TES1 mutant cDNA rescue. Higher X-axis fold changes indicate higher expression with WT LMP1 rescue, whereas lower X-axis fold changes indicate higher expression with TES1m rescue. Data are from n=3 RNAseq datasets, as in **Fig. 3**.

(B) Enrichr analysis of KEGG pathways most highly enriched in RNAseq data as in (A) amongst genes more highly expressed in LMP1 KO GM12878 with WT than TES1m rescue (red) vs amongst genes more highly expressed with TES1m than WT rescue (blue).

(C) Volcano plot analysis of host transcriptome-wide GM12878 genes differentially expressed in LMP1 KO GM12878 with WT vs TES2 mutant cDNA rescue. Higher X-axis fold changes indicate higher expression with WT LMP1 rescue, whereas lower X-axis fold changes indicate higher expression with TES2m rescue. Data are from n=3 RNAseq datasets, as in **Fig. 3**.

(D) Enrichr analysis of KEGG pathways most highly enriched in RNAseq data as in (C) amongst genes more highly expressed in LMP1 KO GM12878 with WT than TES2m rescue (red) vs amongst genes more highly expressed with TES2m than WT rescue (blue).

(E) Volcano plot analysis of host transcriptome-wide GM12878 genes differentially expressed in LMP1 KO GM12878 with TES1 vs TES2 mutant cDNA rescue. Higher X-axis fold changes indicate higher expression with TES1m rescue, whereas lower X-axis fold changes indicate higher expression with TES2m rescue. Data are from n=3 RNAseq datasets, as in **Fig. 3**.

(F) Enrichr analysis of KEGG pathways most highly enriched in RNAseq data as in (E) amongst genes more highly expressed in LMP1 KO GM12878 with TES1m than TES2m rescue (red) vs amongst genes more highly expressed with TES2m than TES1m rescue (blue).

Fig. 5. Characterization of host genome-wide Akata B-cell LMP1 target genes.

(A) K-means heatmap analysis of RNAseq datasets from n=3 replicates generated in EBV- Akata Burkitt cells with conditional LMP1 WT, TES1m, TES2m or DM expression induced by 250 ng/ml doxycycline for 24 hours. The heatmap visualizes host gene Log2 Fold change across the four conditions, divided into six clusters. A two-way ANOVA P value cutoff of <0.01 and >2-fold gene expression were used. # of genes in each cluster is indicated at right.

(B) Heatmaps of representative Cluster 1 differentially regulated genes (top), with column maximum (max) colored red and minimum (min) colored blue, as shown by the scalebar. Also shown are expression values of two representative Cluster 1 genes (lower left) and Enrichr analysis of KEGG pathways significantly enriched in Cluster 1 gene sets (lower right). p-values were determined by one-sided Fisher's exact test. ***p<0.001.

(C) Heatmaps of representative Cluster 2 differentially regulated genes (top), as in (B). Also shown are expression values of two representative Cluster 2 genes (lower left) and Enrichr analysis of KEGG pathways significantly enriched in Cluster 2 gene sets (lower right). p-values were determined by one-sided Fisher's exact test. *p<0.05, **p<0.01, ***p<0.001.

(D) Heatmaps of representative Cluster 6 differentially regulated genes (top), as in (B). Also shown are expression values of two representative Cluster 6 genes (lower left) and Enrichr analysis of KEGG pathways significantly enriched in Cluster 6 gene sets (lower right). p-values were determined by one-sided Fisher's exact test. *p<0.05, ***p<0.001.

Fig. S4. Validation of LMP1 WT, TES1m, TES2m and DM conditional expression system in EBV-negative Akata Burkitt B-cells.

(A) Immunoblot analysis of WCL from Akata cells induced for LMP1 WT, TES1m, TES2m or DM expression by addition of 250 ng/ml doxycycline (Dox) for 24 hours, as indicated. For cross-comparison, WCL from equal numbers of Mutu I Burkitt lymphoma (latency I, lacks LMP1 expression) and GM12878 were also included at right. Blots are representative of n = 3 experiments.

(B) Immunoblot analysis of WCL from BL-41 cells induced for WT, TES1m, TES2m or DM LMP1 by addition of 250 ng/ml Dox for 24 hours, as indicated. For cross-comparison, WCL from equal numbers of Mutu I Burkitt and GM12878 were also included at right. Blots are representative of n = 3 experiments.

(C) FACS analysis of plasma membrane (PM) ICAM-1 abundance in Akata cells induced for LMP1 by 250 ng/ml Dox for 24 hours, as indicated. Y-axis are histogram cell counts, X-axis represents PM ICAM-1 abundance. For comparison, levels in GM12878 LCLs or latency I Mutu I Burkitt cells are shown.

(D) PM ICAM-1 Mean Fluorescence Intensity (MFI) + standard deviation (SD) from n=3 replicates in Akata cells with the indicated LMP1 expression, as in (C). p-values were determined by one-sided Fisher's exact test. **p<0.001, ***p<0001.

(E) FACS analysis of PM Fas abundance in Akata cells induced for LMP1 by 250ng/ml of Dox for 24 hours as indicated. For comparison, GM12878 LCLs or latency I Mutu I Burkitt cells were also analyzed.

(F) PM Fas MFI + SD from n=3 replicates in Akata cells with the indicated LMP1 expression, as in (E). P-values were determined by one-sided Fisher's exact test. **p<0.001, ***p<0.0001.

(G) FACS analysis of PM ICAM-1 in BL-41 cells induced for LMP1 expression by 250ng/ml Dox for 24 hours, as indicated. For comparison, GM12878 and Mutu I were also analyzed. p-values were determined by one-sided Fisher's exact test. **p<0.001, ***p<0.0001.

(H) PM ICAM-1 MFI + SD from n=3 replicates in Akata cells with the indicated LMP1 expression, as in (G). P-values were determined by one-sided Fisher's exact test. **p<0.001, ***p<0.0001.

(I) FACS analysis of PM Fas levels in BL-41 cells induced for LMP1 expression by 250 ng/ml dox for 24 hours, as indicated. GM12878 and Mutu I were analyzed for cross-comparison. p-values were determined by one-sided Fisher's exact test. **p<0.001, ***p<0.0001.

(J) PM Fas MFI + SD from n=3 replicates of BL-41-LMP1 with the indicated LMP1 expression, as in (I). Mutu I and GM12878 were analyzed for comparison. p-values were determined by one-sided Fisher's exact test. **p<0.001, ***p<0.0001.

Figure S5

Fig. S5. Characterization of host genome-wide Akata B-cell LMP1 target genes, related to Figure 4.

(A) Heatmaps of representative **Figure 4** Cluster 3 differentially regulated genes (top), with column maximum colored red and minimum colored blue, as shown by the scalebar. Also shown are expression values of two representative Clusters 3 genes (lower left) and Enrichr analysis of KEGG pathways most significantly enriched Cluster 3 gene sets (lower right). p-values were determined by one-sided Fisher's exact test. **p<0.01.

(B) Heatmaps of representative **Figure 4** Cluster 4 differentially regulated genes (top), with column maximum colored red and minimum colored blue, as shown by the scalebar. Also shown are expression values of two representative Clusters 4 genes (lower left) and Enrichr analysis of KEGG pathways most significantly enriched Cluster 4 gene sets (lower right). p-values were determined by one-sided Fisher's exact test. *p<0.05, **p<0.01, ***p<0.001.

(C) Heatmaps of representative **Figure 4** Cluster 5 differentially regulated genes (top), with column maximum colored red and minimum colored blue, as shown by the scalebar. Also shown are expression values of two representative Clusters 5 genes (lower left) and Enrichr analysis of KEGG pathways most significantly enriched Cluster 5 gene sets (lower right). p-values were determined by one-sided Fisher's exact test. *p<0.05, ***p<0.001.

(D) RT-PCR analysis of CCL22 mRNA abundance in cells with conditional LMP1 WT, TES1m, TES2m or DM expression induced by 250 ng/ml doxycycline for 24 hours as in Fig. 5A. The same RNA used for RNA-seq in Fig. 5A was used for these qPCR experiments. Values from cells with conditional WT LMP1 expression were set to 1, and mean fold-change of CCL22 mRNA abundance + SD in cells with conditional TES1m,TES2m or DM expression are shown. p-values were determined by one-sided Fisher's exact test from two independent experiments, each with two technical replicates. **p<0.01.

(E) RT-PCR analysis of EBI3 mRNA abundance in cells with conditional LMP1 WT, TES1m, TES2m or DM expression induced by 250 ng/ml doxycycline for 24 hours as in Fig. 5A. The same RNA used for RNA-seq in Fig. 5A was used for these qPCR experiments. Values from cells with conditional WT LMP1 expression were set to 1, and mean fold-change of CCL22 mRNA abundance + SD in cells with conditional TES1m,TES2m or DM expression are shown. p-values were determined by one-sided Fisher's exact test from two independent experiments, each with two technical replicates. **p<0.01.

(F) RT-PCR analysis of IRF4 mRNA abundance in cells with conditional LMP1 WT, TES1m, TES2m or DM expression induced by 250 ng/ml doxycycline for 24 hours as in Fig.5A. The same RNA used for RNA-seq in Fig. 5A was used for these qPCR experiments. Values from cells with conditional WT LMP1 expression were set to 1, and mean fold-change of CCL22 mRNA abundance + SD in cells with conditional TES1m,TES2m or DM expression are shown. p-values were determined by one-sided Fisher's exact test from two independent experiments, each with two technical replicates. **p<0.01.

Fig. S6. RNAseq analysis of BL-41 B-cell responses to WT, TES1, TES2 or DM LMP1.

(A) K-means heatmap analysis of RNAseq datasets from n=3 replicates generated in EBV- BL-41 Burkitt cells with conditional LMP1 WT, TES1m, TES2m or DM expression induced by 250 ng/ml doxycycline for 24 hours. The heatmap visualizes host gene Log2 Fold change across the four conditions, divided into six clusters. A two-way ANOVA P value cutoff of <0.01 and >2-fold gene expression were used. # of genes in each cluster is indicated at right.

(B) Heatmaps of representative Cluster 1 differentially regulated genes (top), with column maximum (max) colored red and minimum (min) colored blue, as shown by the scalebar. Also shown are expression values of two representative Cluster 1 genes (lower left) and Enrichr analysis of KEGG pathways significantly enriched in Cluster 1 gene sets (lower right). p-values were determined by one-sided Fisher's exact test. ***p<0.001.

(C) Heatmaps of representative Cluster 2 differentially regulated genes (top). Also shown are expression values of two representative Cluster 2 genes (lower left) and Enrichr analysis of KEGG pathways significantly enriched in Cluster 2 gene sets (lower right). p-values were determined by one-sided Fisher's exact test. ***p<0.001.

(D) Heatmaps of representative Cluster 4 differentially regulated genes (top). Also shown are expression values of two representative Cluster 4 genes (lower left) and Enrichr analysis of KEGG pathways significantly enriched in Cluster 4 gene sets (lower right). p-values were determined by one-sided Fisher's exact test. *p<0.05, ***p<0.001.

(E) Heatmaps of representative Cluster 6 differentially regulated genes (top). Also shown are expression values of two representative Cluster 6 genes (lower left) and Enrichr analysis of KEGG pathways significantly enriched in Cluster 6 gene sets (lower right). p-values were determined by one-sided Fisher's exact test.*p<0.05, **p<0.01, ***p<0.001.

Fig. 6. Characterization of Akata B-cell pathways targeted by TES1 vs TES2 signaling.

(A) Volcano plot analysis of host transcriptome-wide genes differentially expressed in Akata cells conditionally induced for WT LMP1 expression for 24h by 250 ng/ml Dox versus in mock induced cells. Higher X-axis fold changes indicate genes more highly expressed in cells with WT LMP1 expression, whereas lower X-axis fold changes indicate higher expression in cells mock induced for LMP1. Data are from n=3 RNAseq datasets, as in **Fig. 5**.

(B) Enrichr analysis of KEGG pathways most highly enriched in RNAseq data as in (A) amongst genes more highly expressed in Akata with WT LMP1 (red) vs amongst genes more highly expressed with mock LMP1 induction (blue).

(C) Volcano plot analysis of host transcriptome-wide genes differentially expressed in Akata cells conditionally induced for WT vs DM LMP1 expression for 24h by 250 ng/ml Dox. Higher X-axis fold changes indicate genes more highly expressed in cells with WT LMP1 expression, whereas lower X-axis fold changes indicate higher expression in cells with DM LMP1. Data are from n=3 RNAseq datasets, as in **Fig. 5**.

(D) Enrichr analysis of KEGG pathways most highly enriched in RNAseq data as in (C) amongst genes more highly expressed in Akata with WT LMP1 (red) vs amongst genes more highly expressed with DM LMP1 induction (blue).

(E) Volcano plot analysis of host transcriptome-wide genes differentially expressed in Akata cells conditionally induced for WT vs TES1m LMP1 expression for 24h by 250 ng/ml Dox. Higher X-axis fold changes indicate genes more highly expressed in cells with WT LMP1 expression, whereas lower X-axis fold changes indicate higher expression in cells with TES1m LMP1. Data are from n=3 RNAseq datasets, as in **Fig. 5**.

(F) Enrichr analysis of KEGG pathways most highly enriched in RNAseq data as in (E) amongst genes more highly expressed in Akata with WT LMP1 (red) vs amongst genes more highly expressed with TES1m LMP1 induction (blue).

(G) Volcano plot analysis of host transcriptome-wide genes differentially expressed in Akata cells conditionally induced for WT vs TES2m LMP1 expression for 24h by 250 ng/ml Dox. Higher X-axis fold changes indicate genes more highly expressed in cells with WT LMP1 expression, whereas lower X-axis fold changes indicate higher expression in cells with TES2m LMP1. Data are from n=3 RNAseq datasets, as in **Fig. 5**.

(H) Enrichr analysis of KEGG pathways most highly enriched in RNAseq data as in (G) amongst genes more highly expressed in Akata with WT LMP1 (red) vs amongst genes more highly expressed with TES2m LMP1 induction (blue).

Fig. S7. Cross-comparison of WT and DM LMP1 effects on Akata vs BL-41 transcriptomes.

(A) Volcano plot analysis of host transcriptome-wide genes differentially expressed in BL-41 cells conditionally induced for WT LMP1 expression for 24h by 250 ng/ml Dox versus in mock induced cells. Higher X-axis fold changes indicate genes more highly expressed in cells with WT LMP1 expression, whereas lower X-axis fold changes indicate higher expression in cells mock induced for LMP1. Data are from n=3 RNAseq datasets.

(B) Enrichr analysis of KEGG pathways most highly enriched in RNAseq data as in (A) amongst genes more highly expressed in BL-41 with WT LMP1 (red) vs amongst genes more highly expressed with mock LMP1 induction (blue).

(C) Volcano plot cross-comparison of Log2 transformed fold change of host mRNA levels in BL-41 cells (X-axis) versus Akata cells (Y-axis) uninduced versus induced for WT LMP1 by 250 ng/ml Dox for 24 hours. Selected genes highly WT LMP1 induced in both Burkitt contexts are highlighted in red, whereas selected genes suppressed by LMP1 in both Burkitt contexts are highlighted in blue.

(D) Volcano plot analysis of host transcriptome-wide genes differentially expressed in BL-41 cells conditionally induced for DM versus WT LMP1 expression for 24h by 250 ng/ml Dox. Higher X-axis fold changes indicate genes more highly expressed in cells with WT LMP1 expression, whereas lower X-axis fold changes indicate higher expression in cells induced for DM LMP1. Data are from n=3 RNAseq datasets.

(B) Enrichr analysis of KEGG pathways most highly enriched in RNAseq data as in (D) amongst genes more highly expressed in BL-41 with WT LMP1 (red) vs amongst genes more highly expressed with DM LMP1 induction (blue).

(C) Volcano plot cross-comparison of Log2 transformed fold change of host mRNA levels in BL-41 cells (X-axis) versus Akata cells (Y-axis) induced for DM versus WT LMP1 by 250 ng/ml Dox for 24 hours. Selected genes highly WT LMP1 induced in both Burkitt contexts relative to levels in cells with DM LMP1 expression are highlighted in red, whereas selected genes suppressed by WT LMP1 in both Burkitt contexts are highlighted in blue.

Fig. S8. Cross-comparison of TES1 and TES2 LMP1 effects on Akata vs BL-41 transcriptomes.

A) Volcano plot analysis of host transcriptome-wide genes differentially expressed in BL-41 cells conditionally induced for TES1m vs WT LMP1 expression for 24h by 250 ng/ml Dox. Higher X-axis fold changes indicate genes more highly expressed in cells with WT LMP1 expression, whereas lower X-axis fold changes indicate higher expression induced for TES1m LMP1. Data are from n=3 RNAseq datasets.

(B) Enrichr analysis of KEGG pathways most highly enriched in RNAseq data as in (A) amongst genes more highly expressed in BL-41 with WT LMP1 (red) vs amongst genes more highly expressed with TES1m LMP1 induction (blue).

(C) Volcano plot analysis of host transcriptome-wide genes differentially expressed in BL-41 cells conditionally induced for TES2m vs WT LMP1 expression for 24h by 250 ng/ml Dox. Higher X-axis fold changes indicate genes more highly expressed in cells with WT LMP1 expression, whereas lower X-axis fold changes indicate higher expression induced for TES2m LMP1. Data are from n=3 RNAseq datasets.

(D) Enrichr analysis of KEGG pathways most highly enriched in RNAseq data as in (A) amongst genes more highly expressed in BL-41 with WT LMP1 (red) vs amongst genes more highly expressed with TES2m LMP1 induction (blue).

(E) Volcano plot analysis of host transcriptome-wide genes differentially expressed in Akata cells conditionally induced for TES1m vs TES2m LMP1 expression for 24h by 250 ng/ml Dox. Higher X-axis fold changes indicate genes more highly expressed in cells with TES1m LMP1 expression, whereas lower X-axis fold changes indicate higher expression induced for TES2m LMP1. Data are from n=3 RNAseq datasets.

(F) Enrichr analysis of KEGG pathways most highly enriched in RNAseq data as in (E) amongst genes more highly expressed in Akata with TES1m LMP1 (red) vs amongst genes more highly expressed with TES2m LMP1 induction (blue).

(G) Volcano plot analysis of host transcriptome-wide genes differentially expressed in BL-41 cells conditionally induced for TES1m vs TES2m LMP1 expression for 24h by 250 ng/ml Dox. Higher X-axis fold changes indicate genes more highly expressed in cells with TES1m LMP1 expression, whereas lower X-axis fold changes indicate higher expression induced for TES2m LMP1. Data are from n=3 RNAseq datasets.

(B) Enrichr analysis of KEGG pathways most highly enriched in RNAseq data as in (G) amongst genes more highly expressed in BL-41 with TES1m LMP1 (red) vs amongst genes more highly expressed with TES2m LMP1 induction (blue).

A

C

Figure S9. Cross-comparison of host genes differentially expressed upon perturbation of LCL LMP1 versus upon LMP1 induction in Akata cells.

(A) Volcano plot analysis of host genes differentially expressed upon WT LMP1 induction in Akata (X-axis) versus upon LMP1 KO in GM12878 (Y-axis). Shown are Log2 transformed mRNA fold change values for Akata cells mock induced versus induced for LMP1 WT expression for 24 hours (X-axis) versus upon expression of LMP1 vs control sgRNA in GM18278 for 48 hours. Genes more highly expressed in mock-induced Akata have higher x-axis values, whereas genes more highly expressed in Akata induced for WT LMP1 have lower x-axis values. Likewise, genes with higher expression with control sgRNA expression have higher y-axis values, whereas genes with lower expression in GM12878 with LMP1 KO have lower Y-axis values. P value <0.05 and >2-fold gene expression cutoffs were used.

(B) Volcano plot analysis of host genes differentially expressed upon TES1m vs WT LMP1 induction in Akata (X-axis) versus upon rescue of LMP1 KO GM12878 with TES1m versus WT LMP1 (Y-axis). Shown are Log2 transformed mRNA fold change values for Akata cells induced for TES1m versus WT LMP1 expression for 24 hours (X-axis) versus upon rescue of GM18278 LMP1 KO with TES1m vs WT LMP1 cDNA, as in **Fig 3**. Genes more highly expressed in Akata with TES1m than WT LMP1 expression have higher x-axis values, whereas genes more highly expressed in Akata induced for WT LMP1 than TES1m have lower x-axis values. Likewise, GM12878 genes with higher expression with TES1m rescue have higher y-axis values, whereas genes with lower expression in GM12878 with TES1m than WT rescue have lower Y-axis values. P value <0.05 and >2-fold gene expression cutoffs were used.

(C) Volcano plot analysis of host genes differentially expressed upon TES2m vs WT LMP1 induction in Akata (X-axis) versus upon rescue of LMP1 KO GM12878 with TES2m versus WT LMP1 (Y-axis). Shown are Log2 transformed mRNA fold change values for Akata cells induced for TES2m versus WT LMP1 expression for 24 hours (X-axis) versus upon rescue of GM18278 LMP1 KO with TES2m vs WT LMP1 cDNA, as in **Fig 3**. Genes more highly expressed in Akata with TES2m than WT LMP1 expression have higher x-axis values, whereas genes more highly expressed in Akata induced for WT LMP1 than TES2m have lower x-axis values. Likewise, GM12878 genes with higher expression with TES2m rescue have higher y-axis values, whereas genes with lower expression in GM12878 with TES2m than WT rescue have lower Y-axis values. P value <0.05 and >2-fold gene expression cutoffs were used.

Figure 7

Fig. 7. Roles of TES1 and TES2 canonical NF-κB pathways in LCL dependency factor BATF and IRF4 expression.

(A) Schematic diagram of JUNB, BATF and IRF4 at an AP-1/IRF composite DNA site.

(B) Mean + SD log2 fold changes of IRF4 and BATF mRNA abundances from n=3 RNAseq replicates of Akata cells expressing the indicated LMP1 cDNA for 24 hours, as in Fig. 5. p-values were determined by one-sided Fisher's exact test. *p<0.05, ***p<0.001.

(C) Schematic diagram of LMP1 TES1 and TES2 NF-κB pathways. TES1 and TES2 each activate canonical NF-κB pathways, whereas TES1 also activates non-canonical NF-κB.

(D) Immunoblot analysis of WCL from Akata cells induced for LMP1 expression by 250 ng/ml Dox for 24 hours, either without or with 1 μM IKKβ inhibitor VIII. Shown below are relative fold changes + SD from n=3 replicates of IRF4 or BATF vs GAPDH load control densitometry values. Values in vehicle control treated WT LMP1 expressing cells were set to 1. P-values were determined by one-sided Fisher's exact test. **p<0.001, ***p<0.0001.

(E) Immunoblot analysis of WCL from Cas9+ Akata cells expressing control or either of two *TAK1* targeting sgRNAs, induced for LMP1 expression by 250 ng/ml Dox for 24 hours. Shown below are relative foldchanges + SD from n=3 replicates of IRF4 or BATF vs GAPDH load control densitometry values. Levels in cells with control sgRNA (sgControl) and WT LMP1 were set to 1. **p<0.001, ***p<0.0001.

(F) Model of additive TES1 and TES2 canonical NF-κB pathway effects on BATF and IRF4 induction.

A

B

C

D

Fig. S10. Roles of TES1 and TES2 canonical NF-κB pathways in BATF and IRF4 induction.

(A) Immunoblot analysis of WCL from latency III Jijoye Burkitt cells or GM12878 LCL expressing LMP1 targeting sgRNA, as indicated. Shown below are relative fold changes + SD from n=3 replicates of IRF4 or BATF vs GAPDH load control densitometry values, with values in sgRNA control expressing cells set to 1. P-values were determined by one-sided Fisher's exact test. **p<0.001, ***p<0.0001.

(B) Immunoblot analysis of WCL from GM12878 LCLs treated with vehicle control or 1 μM IKKβ inhibitor VIII for 24 hours. Shown below are relative foldchanges + SD from n=3 replicates of IRF4 or BATF vs GAPDH load control densitometry values. Levels in vehicle control treated WT LMP1 expressing cells were set to 1. P-values were determined by one-sided Fisher's exact test. **p<0.001, ***p<0.0001.

(C) Immunoblot analysis of WCL from Akata cells induced for the indicated LMP1 construct expression, either without or together with an IκBα super-repressor (IκBα-S.R.) that blocks canonical NF-κB signaling. Shown below are relative foldchanges + SD of IRF4 or BATF vs GAPDH load control densitometry values from n=3 replicates, with values in cells expressing WT LMP1 but not IκBα- set to 1. P-values were determined by one-sided Fisher's exact test. **p<0.001, ***p<0.0001.

(D) Mean + SD fold changes of IRF4 and BATF mRNA abundances from n=3 RNAseq replicates of GM12878 LCLs transduced with lentivirus expressing LMP1 sgRNA and induced for WT, TES1m or TES2m rescue cDNA expression for 6 days, as in Fig 3. P-values were determined by one-sided Fisher's exact test. ***p<0.001.

A Typical LCL Enhancer   LCL Super-Enhancer

B   ○ LCL SE Targets ○ Other LCL genes

C   ○ LCL SE Targets ○ Other LCL genes

D   ○ LCL SE Targets ○ Other LCL genes

Fig. 8. LMP1 TES1 and TES2 roles in EBV super-enhancer target gene regulation in GM12878 LCLs.

(A) Schematic diagram of typical LCL enhancers vs super-enhancers. Super-enhancers have significantly broader and taller histone 3 lysine 27 acetyl (H3K27Ac) peaks. EBV SE are host genomic enhancer sites bound by all five LMP1-activated NF-κB transcription factor subunits, EBNA-2, LP, 3A, and 3C.

(B) Volcano plot analysis of mRNA values in GM12878 expressing LMP1 vs control sgRNAs as in **Fig. 3**. Genes targeted by EBV super-enhancers (SE) are highlighted by red circles, whereas other LCL genes are indicated by blue circles. Genes more highly expressed with LMP1 KO have higher x-axis values, whereas those downmodulated by LMP1 KO have lower values. P value <0.05 and >2-fold gene expression cutoffs were used.

(C) Volcano plot analysis of mRNA values in GM12878 expressing LMP1 sgRNA with TES1m versus WT LMP1 cDNA rescue, as in **Fig. 2-3**. Genes targeted by EBV super-enhancers (SE) are highlighted by red circles, whereas other LCL genes are indicated by blue circles. Genes more highly expressed with endogenous LMP1 KO and TES1m rescue have higher x-axis values, whereas those more highly expressed with WT LMP1 rescue have lower values. P value <0.05 and >2-fold gene expression cutoffs were used.

(D) Volcano plot analysis of mRNA values in GM12878 expressing LMP1 sgRNA with TES2m versus WT LMP1 cDNA rescue, as in (C). Genes targeted by EBV super-enhancers (SE) are highlighted by red circles, whereas other LCL genes are indicated by blue circles. Genes more highly expressed with endogenous LMP1 KO and TES2m rescue have higher x-axis values, whereas those more highly expressed with WT LMP1 rescue have lower values. P value <0.05 and >2-fold gene expression cutoffs were used.

Figure S11. LMP1 TES1 and TES2 roles in EBV super-enhancer target gene regulation in Akata and BL-41 Burkitt B-cells.

(A) Volcano plot analysis of Akata RNA-seq, comparing mRNA values in cells induced for LMP1 WT vs. mock-induced for 24 hours. SE targeted genes highlighted by red circles and other B-cell genes indicated by blue circles. P value <0.05 and >2-fold gene expression cutoffs were used.

(B) Volcano plot analysis of Akata RNA-seq, comparing mRNA values in cells induced for LMP1 TES1m vs. mock-induced for 24 hours. SE targeted genes highlighted by red circles and other B-cell genes indicated by blue circles. P value <0.05 and >2-fold gene expression cutoffs were used.

(C) Volcano plot analysis of Akata RNA-seq, comparing mRNA values in cells induced for LMP1 TES2m vs. mock-induced for 24 hours. SE targeted genes highlighted by red circles and other B-cell genes indicated by blue circles. P value <0.05 and >2-fold gene expression cutoffs were used.

(D) Volcano plot analysis of BL-41 RNA-seq, comparing mRNA values in cells induced for LMP1 WT vs. mock-induced for 24 hours. SE targeted genes highlighted by red circles and other B-cell genes indicated by blue circles. P value <0.05 and >2-fold gene expression cutoffs were used.

(E) Volcano plot analysis of BL-41 RNA-seq, comparing mRNA values in cells induced for LMP1 TES1m vs. mock-induced for 24 hours. SE targeted genes highlighted by red circles and other B-cell genes indicated by blue circles. P value <0.05 and >2-fold gene expression cutoffs were used.

(F) Volcano plot analysis of BL-41 RNA-seq, comparing mRNA values in cells induced for LMP1 TES2m vs. mock-induced for 24 hours. SE targeted genes highlighted by red circles and other B-cell genes indicated by blue circles. P value <0.05 and >2-fold gene expression cutoffs were used.

Fig. S12. Model highlighting different modes of LMP1 TES1 and TES2 cross-talk in Burkitt B-cell target gene regulation. We identify 6 different modes of TES 1/2 cross-talk where (a) TES1 is required for the transcription of these subset of genes over TES2 (in limited amounts) or (b) TES1 or TES2 are individually sufficient for the expression of these subset of genes, or (c) TES1 is required to block the TES2-mediated suppression of this subset, or (d) TES2 is required to block the TES1-mediated suppression of this subset, or (e) TES1 is required to block the TES2-mediated upregulation of this subset, or (f) TES1 and TES2 are required for LMP1-mediated suppression of this gene subset.