Title: Sorting of persistent morphological polymorphisms links paleobiological pattern to population process

Author: C. Tomomi Parins-Fukuchi

Affiliation: Department of Ecology and Evolutionary Biology, University of Toronto, Toronto, Ontario, Canada M5S 3B2

Abstract

Biological variation fuels evolutionary change. Across longer timescales, however, polymorphisms at both the genomic and phenotypic levels often persists longer than would be expected under standard population genetic models such as positive selection or genetic drift. Explaining the maintenance of this variation within populations across long timespans via balancing selection has been a major triumph of theoretical population genetics and ecology. Although persistent polymorphisms can often be traced in fossil lineages over long periods through the rock record, paleobiology has had little to say about either the long-term maintenance of phenotypic variation or its macroevolutionary consequences. I explore the dynamics that occur when persistent polymorphisms maintained over long lineage durations are filtered into descendant lineages during budding speciation events. I evaluate these patterns in two lineages: *Ectocion*, a genus of Eocene mammals, and botryocrinids, a Mississippian echinoderm family. Following budding events, descendants are less variable than their ancestors. The patterns by which ancestral variation is sorted cannot be distinguished from drift. Maintained and accumulated polymorphisms in highly variable ancestral lineages such as *Barycrinus rhombiferus* may fuel radiations as character states are sorted into multiple descendant lineages. Interrogating the conditions under which trans-specific polymorphism is either maintained or lost during periods of demographic and ecological upheaval can explain how population-level processes contribute to the emergent macroevolutionary dynamics that shape the history of life as preserved in the fossil record.

Keywords: Phylogeny, budding speciation, crinoids, mammals, polymorphism

Non-technical abstract

Understanding how morphological variation changes within populations over relatively short timescales in response to environmental changes and ecology (i.e., thousands of years) is a major focus of paleontology and evolutionary biology. A distinct focus is in understanding the broad-scale patterns by which lineages have diversified into distinct environments over geologic time (i.e., millions of years). One major challenge has been reconciling how and whether processes acting over shorter timescales shape the patterns observed over long timescales. One way of examining morphological variation at the population level is by examining the distribution of polymorphic character states-- discrete anatomical features that vary within a population. Fossil species often maintain such polymorphisms for long periods of time, with such variation even sometimes inherited by new species from their ancestors. In this article, I suggest that examining

how these polymorphisms are distributed among incipient descendant lineages might help link
the ecological and evolutionary processes that act at the population level (e.g., natural selection,
genetic drift, competition, predation) to the paleobiological patterns that are often reconstructed
across many species and over long timescales. I explore these dynamics in two lineages:
Ectocion, a genus of Eocene mammals, and botryocrinids, a Mississippian echinoderm family. I
found that new lineages typically have fewer polymorphisms than their ancestors, suggesting that
ancestral variation is "sorted" into incipient lineages during speciation events. This variation
appears to be sorted randomly, which means that it is not possible to detect the influence of
natural selection in guiding the inheritance of ancestral morphologies. I suggest that the patterns
by which ancestral variation is sorted into new species may explain patterns of lineage
diversification over long timescales, highlighting how population processes can extend their
influence over longer timescales to shape large-scale evolutionary dynamics.

Introduction

Polymorphism is ubiquitous within natural populations. Spanning levels from the genome to the
phenotype, it is difficult to look far without observing variation. While variation within
populations is most often examined from the standpoint of microevolution, there is increasing
evidence that polymorphisms can persist over long time periods. Long-sustained intrapopulation
variation has been frequently observed at the protein and genetic level since the 1960's (e.g.,
Lewontin and Hubby 1966, Hughes and Nei 1988, Klein et al. 1998). Polymorphisms at the
genetic level can even be maintained for millions of years and across speciation events
(Muirhead et al. 2002). There is also growing evidence that polymorphisms in complex
phenotypic characters and gene networks may persist for very long time periods (Hittiger et al.
2010). Because it spans temporal and taxonomic scales, persistent polymorphic variation might
offer a window into understanding how the microevolutionary processes that promote long-term
polymorphism might shape macroevolutionary patterns and processes.

The evolution and maintenance of genetic and phenotypic diversity is a longstanding and well-
explored topic in population genetics. Explaining the high levels of variation maintained in
natural populations was an early challenge in post-synthesis evolutionary biology. Positive
selection, purifying selection, and genetic drift are all expected to remove variation from the
population, which has appeared at odds with the vast reservoirs of variation displayed by many
organisms. However, theoretical and empirical work has shown how environmental
heterogeneity (McDonald and Ayala 1974, Christiansen 1975, Chakraborty and Fry 2016, Gallet
et al. 2018) and negative frequency-dependent selection (NFDS) (Clark 1979, Charlesworth
2006, Fitzpatrick et al. 2007, Takahashi et al. 2010) protect such "balanced polymorphisms",
maintaining genetic and phenotypic variation over long timescales.

Despite its omnipresence in nature and frequent treatment in population genetics, the causes and
effects of intra-population polymorphism have been understudied at deeper timescales.
Characters employed in phylogenetic studies frequently display polymorphism across multiple
lineages but have frequently downplayed this variation. Nevertheless, they are sufficiently
common to substantially impact phylogenetic inferences when not explicitly accommodated in

analyses (Wiens 1995, 1999). Phenotypic polymorphisms also feature strongly in paleontological
systematics. Characters chosen for cladistic analysis between fossil lineages are frequently rife
with polymorphism, which can present challenges to existing algorithms (e.g., Trueman 2010,
Gilbert 2013, Whiting et al., 2016, plus many others). Relative to the goals of systematics, these
are often treated as nuisances. This is perhaps due to a historical gaps in the integration between
systematics, phylogenetic methods, and population genetics. Despite being frequently observed,
the apparent maintenance of polymorphism across populations separated by millions of years of
independent evolution remains understudied as a macroevolutionary phenomenon.

A small, but growing, body of work has contributed early inklings of the potential for sustained
variation to explain evolutionary dynamics across species boundaries and over increasingly deep
timescales. One increasingly important issue in evolutionary ecology concerns how phenotypic
and genetic variation evolves across speciation events (e.g., Thompson et al. 2019). Standing
genetic variation can facilitate rapid adaptation when populations are exposed to new ecological
pressures, such as might occur during ecological speciation (Barrett and Schluter 2008, Sicard et
al. 2016, Lai et al. 2019, McGee et al. 2020). There has also been an increasing focus in
population genomics upon understanding how (often balanced) polymorphisms in large ancestral
populations may be "sieved" by speciation events, becoming fixed among descendants either
randomly or by selection (Pease et al. 2016, Guerrero and Hahn 2017). "Deeply persistent"
polymorphisms can even be maintained by NFDS over millions of years, shaping
macroevolutionary dynamics over large clades (Igic et al. 2006, Goldberg et al. 2010).

Paleobiologists have spent less time seeking process-based explanations for the evolution of
phenotypic variation. However, the fossil record might be leveraged to offer unique insight into
this fundamental question. Fossilized populations often display maintain variation over long
intervals of time (e.g., Brothwell 1963, also see examples reviewed extensively by Van Valen
1969). Although the fossil record has been underutilized as a laboratory to investigate population
genetics, the marine invertebrate record has shown promise in its ability to illuminate the
processes governing the maintenance of balanced polymorphisms (Kermack 1954). Intraspecific
variation has also been hypothesized to have contributed to the explosive diversification of
trilobites during the Cambrian (Webster 2007). Fossil lineages also frequently display high
population-level variation, both in quantitative traits and discrete character states. This variation
can be maintained for long periods of time, across a range of environmental conditions
(including environmental stasis—see Schopf and Gooch 1972). Despite these substantial
advances, the full incorporation of persistent, trans-specific phenotypic polymorphisms in the
fossil record into paleobiological theory has remained incomplete. Nevertheless, the presence of
persistent intraspecific variation shared between fossilized lineages is an often-unrecognized
curiosity when viewed from the lens of population genetics that may contain hidden insights
when bridging micro- and macroevolutionary scales.

The evolution of phenotypic variation across speciation events has been a fundamental question
in paleobiology for decades (Simpson 1944, Eldredge and Gould 1972, Gingerich 1974, Polly
1997). Key neontological work has also suggested that examining the maintenance of ancestral
variation and its distribution among descendant populations may explain the formation and
adaptive divergence of new lineages (Wright 1982, Schluter 2000, Schluter and Conte 2009).

Although often discussed in different terms and using different sources of data than the modern
population ecology literature, paleobiologists' interests and goals are often fundamentally
overlapping. Paleobiology is often concerned with understanding the mode of speciation— for
example, in comparing splitting vs budding cladogenesis, and determining the long-term
consequences thereafter. Each of these has distinct demographic predictions and can stem from a
variety of different population processes, each leading to a different set of macroevolutionary
consequences.

Budding speciation, identified as asymmetric speciation between a large ancestral population,
and a smaller descendant population, has long served as paleobiological bread and butter (e.g.,
Rensch 1959, Raup and Gould 1974, Van Valen 1975, Raup 1985, Jackson and Cheetham 1994,
Wagner and Erwin 1995, Aze et al. 2011, Bapst and Hopkins 2017, Warnock et al. 2020, Parins-
Fukuchi 2021). One important characteristic of budding is the persistence of the ancestral lineage
alongside its newly formed descendant. There may be a geographic component, wherein the
ancestral lineage occupies a wide geographic range and the descendant lineage arises due to
peripheral isolation within a smaller area, but this is not essential. Dynamics in the distribution of
phenotypic changes and survivorship between coexisting ancestor and descendant species can
offer unique insights into large-scale evolutionary pattern and process (e.g., Pearson 1998). It
may also be sufficiently common to warrant explicit incorporation into phylogenetic comparative
methods (Caetano and Quental 2022).

Tracing the evolution of polymorphic characters across the repeated episodes of budding
speciation found in dense fossil records would provide a unique opportunity to test the
predictions from population genetics surrounding the inheritance of persistent, standing variation
into descendant populations and examine their effects over longer time scales. For example,
when populations bud off of an ancestral lineage, the resulting speciation event may be
"parallel", "divergent", or "random", with respect to the traits and alleles that become fixed
(Schluter and Nagel 1995, Thompson 2019). In the parallel case, new descendant lineages tend
to fix the same traits as they radiate into similar environments. In the divergent case, descendants
evolve in opposing phenotypic directions as they radiate into distinct environments. Lastly,
newly budded species may randomly sort variation present in the ancestor due to population
bottlenecks caused by the demographic asymmetry of the speciation event. Each of these leads to
different implications both within lineages and over deeper time.

In this paper, I attempt to explain the evolution of phenotypic variation across episodes of
budding speciation in two fossil lineages: *Ectocion*, a genus of Eocene mammals closely related
to perissodactyla; and botryocrinids, a family of mostly Mississippian crinoid echinoderms. My
first goal was to explain how an explicit treatment of persistent polymorphisms can clarify our
interpretation of biological patterns in the fossil record. I then sought to further examine the ways
by which persistent ancestral variation is sorted into descendant lineages. I sought to test whether
1) descendant species tended to be less variable than ancestral species, a key prediction of
budding speciation, and 2) whether ancestral variation tended to sort randomly, consistent with
the fixation of polymorphisms due to population bottlenecking, or non-randomly, consistent with
either convergent or divergent selection following budding events.

Materials and Methods

Data and code availability. All data and code used in the analyses presented here are available
on Figshare and associated with the digital object identifier 10.6084/m9.figshare.22584097.

Datasets. I harvested morphologic and stratigraphic datasets encompassing two lineages from the
literature: Ectocion (Thewissen 1992) and botryocrinidae (Gahn and Krammer 2002). These two
lineages offer several advantages. Both possess dense fossil records, simplifying the
identification of likely ancestor-descendant relationships. They have also benefited from careful
work delimiting species and tracing continuous lineages through stratigraphic zones in
particularly well-studied regional faunas. The OTUs represented within each dataset can
therefore likely be regarded as trustworthy. This careful work has also provided greater certainty
in trusting the persistence of polymorphisms observed in each dataset—while character state
frequencies among several polymorphic characters fluctuates throughout the range of each
lineage, overall, the polymorphic variation attributed to each lineage largely remained intact. The
maintenance of polymorphism throughout lineage durations is further supported by the
persistence of polymorphic characters across several speciation events. The repeated observation
of polymorphism across multiple lineages favors the persistence of polymorphisms within and
between lineages as the most parsimonious explanation.

The *Ectocion* dataset was entirely dental. After reconstructing the phylogeny, I mapped the
evolution of the P3 metacone, which was polymorphic across two lineages over long time spans.
While explicitly mapping such detailed dental morphology to dietary function is a very
challenging prospect, it is reasonable to imagine that the metacone, which is haphazardly
distributed across *Ectocion*, may have provided some dimension of dietary function. The
botryocrinid dataset sampled characters across the bauplan, spanning several anatomical regions,
including the rami, stalk, plates, and anus. Polymorphisms were observed across all regions.
Functional interpretations for each of these traits is very challenging. However, all, or nearly all,
of them do interact with the environment in some way, for example *Ectocion* premolar
morphology in food processing, or crinoid columnal morphology in hydrodynamics. As a result,
it is feasible that they could be under some form of selection, although closer biomechanical
analysis would be needed to better understand the functional context of each of these traits.

Several of the characters observed by Gahn and Kammer (2002) were not true discrete
characters, but rather, discretized descriptions of variation that is fundamentally continuous.
While this challenges interpretations of these particular characters within the framework of strict
allelic polymorphism, I believe they were appropriate to retain in the current study. This is
because my goal in analyzing the botyrocrinid dataset was in examining how ancestral variation
is sorted and maintained across budding speciation events. While coarse, even these discretized
treatments offer resolution into this topic, especially if the continuous variation represented
within is at least somewhat discontinuous in its frequency distribution. Such variation is also
likely to correspond to higher heterozygosity among underlying loci, supporting their use in
interpreting broad patterns underlying the evolution of variation.

Phylogenetic reconstruction. I implemented a very simple method for the reconstruction of
phylogeny, including ancestor-descendant relationships, for this study. It is based upon the

greedy, agglomorative algorithm employed by neighbor-joining (Saitou and Nei 1987). It only
differs by incorporating stratigraphic information into the distance matrix and allowing earlier-
occurring lineages to serve as direct ancestors of those that occur later. The approach is also very
similar to the greedy algorithm described by Alroy (1995), with some simplifications. It starts by
constructing a matrix of all of the morphological distances between each OTU, calculated as the
number of distinct character states between each possible OTU pair. In the case of
polymorphism, OTU pairs are assigned a distance of zero if the intersection between character
states displayed by each was non-empty. Stratigraphic distances are then added. These are
calculated as the number of discrete gaps separating non-overlapping lineages. Lineages with
overlapping or abutting ranges are assigned a stratigraphic distance of zero. The least dissimilar
OTU pair is then identified and joined. If one of the OTUs first occurs before the other, it is
assumed to be ancestor. If they start in the same time horizon, they are assigned a shared
hypothetical ancestor that is placed in the same time bin. The algorithm then proceeds iteratively,
adding OTU pairs until all pairs are joined. The tree is then rooted using either an outgroup or by
presuming the most stratigraphically basal OTU is the root. The resulting tree is one that
minimizes the amount of evolutionary change and stratigraphic gaps across lineages. It thus
bears similarities to both stratophenetics (Gingerich 1979) and stratocladistics (Fisher 2008).

The method used here is very simplistic. I do not endorse its use in large, complex datasets.
Nevertheless, I believe it is adequate for my purposes here. This is because the datasets
employed in this study are quite small and benefit from having very dense and well-characterized
stratigraphic and geographic ranges. In addition, the high degree of polymorphism displayed by
both lineages supported the use of the distance approach used here over existing implementations
in that it offered a simple solution to the treatment of polymorphism when calculating
evolutionary distances. This was important given the high degree of polymorphism in the
Barycrinus dataset and the inability of most existing parametric and parsimony approaches to
accommodate polymorphism explicitly. Several existing Bayesian approaches entertain ancestor-
descendant hypotheses, including budding speciation (Stadler et al. 2018). However, given the
extensive challenges associated with searching treespace to explicitly identify such hypotheses,
rather than simply integrating over them, I felt the simplicity of the method here was justified for
use on the particular datasets employed here, given the importance of putative budding events on
the final biological interpretations. As a final note, in the absence of contrary information, it is
generally more parsimonious to assume that taxa are related through ancestor-descendant
sequences rather than invoking hypothetical ancestors. Given that the density of the fossil
records displayed by the lineages examined here increases the odds of sampling ancestral taxa
(Foote 1996), this reasoning should lend some epistemological confidence in the method's
behavior of minimizing the number of hypothetical ancestors. Nevertheless, my approach to
phylogeny reconstruction, while likely appropriate for the densely sampled and carefully studied
lineages employed here, is propositional, and ultimately somewhat crude, in its nature. It should
therefore be accompanied by a more rigorous, evaluative, approach for future larger-scale studies
of budding dynamics in the fossil record, such as stratocladistics (e.g., Fisher 1991, 2008)
stratolikelihood (e.g., Wagner 1998, 2000), or full-Bayesian approaches (e.g., Wright et al.
2021). On larger datasets, the simple approach implemented here may remain useful in
generating a starting tree for more exhaustive heuristic searches.

I applied this algorithm to reconstruct relationships in the taxa *Ectocion* and botryocrinids.
Stratigraphic ranges were harvested from the same references from which I sourced the
morphologic information. One small adjustment was made in the botryocrinids. I allowed
Barycrinus rhombiferus to be ancestral to *Barycrinus spurius*, despite them having originated in
the same layer. This is because Gahn and Krammer (2002) made a compelling case for *B.
rhombiferus'* likely status as an ancestor to multiple other *Barycrinus* lineages. In addition, while
B. rhombiferus and *B. spurius* were both long-lasting and highly polymorphic lineages, *B.
rhombiferus* possess fewer derived character states than *B. spurius*. This makes a *B. rhombiferus*
-> *B. spurius* ancestor-descendant pair more parsimonious, according to stratocladistic criteria,
than the reverse. In addition, allowing the possibility also achieved a result with several fewer
hypothetical ancestors, making a strong appeal to model parsimony. It should be clarified that *B.
rhombiferus* was not, *a priori*, constrained or assumed to be ancestral to *B. spurius*. This
possibility was simply *allowed* by the analysis.

Random vs. non-random sorting of ancestral variation. I devised a simple permutation test to
identify whether ancestral variation in botryocrinids tended to sort randomly among descendant
lineages, which would be suggestive of drift driving fixation during speciation bottlenecks, or
whether descendant lineages tended to fix the same ancestral variants, which would be consistent
with selection during adaptation in similar environments. Using reconstructed budding events
across *Barycrinus*, I took the polymorphic character states displayed by the ancestral lineage. I
then randomly sampled a single character state from each ancestrally-polymorphic character to
generate a hypothetical character sequence of inherited polymorphisms for each descendant
lineage observed in the dataset. I then compared the pairwise Hamming distances between the
simulated character states displayed by each descendant. To calculate distances, polymorphisms
were treated as a unique character state. As a result, a comparison between a (01) polymorphism
and (1) monomorphism resulted in a distance of 1. This was done according to the reasoning
that, if incipient species were experiencing positive selection in new environments, they should

purge the same ancestral character states. My expectation was that descendants adapting to
similar environments would likely share more character states, and thus have fewer differences. I
repeated this sampling procedure 1000 times to generate a frequency distribution of pairwise
distances between the character states displayed by each descendant that were randomly sampled
from the polymorphic ancestor. I then compared the observed distances to the empirical
distribution to identify any descendant taxon pairs that were more similar (indicating parallel
adaptation) or different (indicating divergent adaptation) than expected under random sorting,
using the 2.5% quantile as a significance threshold.

Results

Ectocion phylogeny. Phylogenetic relationships reconstructed within *Ectocion* (Fig. 1) are largely
concordant with the interpretation that Thewissen (1992) derived from the same dataset based on
stratocladistic criteria. The main point of contention lies in the placement of *E. mediotuber* and
E. cedrus in relation to *E. collinus*. The original study suggested the three taxa to form a grade,
separated by hypothetical ancestors, whereas the current reconstruction posits that *E. cedrus* and

E. mediotuber independently budded from *E. collinus*. This is a minor difference that will be
discussed further below. Otherwise, the ancestor-descendant linking of *E. mediotuber* to *E.*
osbornianus and *E. parvus* corresponds perfectly to the original study. Thewissen was
conservative in his interpretation of *E. superstes*. However, the approach here suggested that it
may have speciated from *E. osbornianus*. It should be noted that, while the diagram presented
here gives the appearance of a direct ancestor-descendant relationship between the two, it is very
possible that their true genealogical relationship is separated by multiple additional speciation
events. This would still make *E. osbornianus* an indirect ancestor.

Evolution of the P3 metacone polymorphism. Mapping the evolution of the P3 metacone to the
Ectocion phylogeny illustrates the importance of considering both ancestor-descendant
relationships and polymorphism when reconstructing patterns in morphological evolution in the
fossil record. A standard cladogram would fail to capture how polymorphisms in *E. mediotuber*
and *E. osbornianus* persisted and sorted among the respective descendants of each, instead
representing the pattern of character changes through multiple character reversals. One slightly
confusing pattern in P3 metacone evolution is the apparent sudden emergence of a monomorphic
metacone in *E. cedrus*. A more parsimonious interpretation, based on P3 morphology, of the
sequence of lineage divergence than that achieved using the greedy method implemented here is
given in the inset to Figure 1, which implies that the metacone evolved one time and sorted
among descendant lineages while being maintained across multiple species. Alternatively, it is
possible that the metacone truly did evolve twice in *Ectocion*. A stronger understanding of the
developmental programs controlling dental variation would help to distinguish whether the
metacone is sufficiently evolvable to have emerged twice, but distinguishing between these two
possibilities falls outside the scope of this study. While the invocation of an additional,
unaccounted polymorphic hypothetical ancestor provides a better explanation for P3 morphology
evolution, detailed phylogenetic work that more thoroughly explores treespace using parsimony
or probabilistic criteria will be needed to confidently distinguish between these possibilities.

Distinguishing between inheritance of ancestral polymorphism and character reversal is
important for deriving meaningful biological interpretations. The pattern in P3 evolution
reconstructed here is consistent with a scenario in which balancing selection maintains variation
across a long-lived lineage, giving rise to descendant lineages that fix that ancestral variation in
different ways. If the maintenance of polymorphism by balancing selection is as common as
ecological theory might suggest, many character distributions among fossil taxa (which are often
interpreted as being highly homoplaseous) may simply be driven by the sorting of variation
among a small handful of variable ancestral lineages. In addition to potentially helping solve
problems in paleontological systematics, exploration of these dynamics may shed light on the
processes that drive the diversification of clades over paleobiological timescales.

The persistence of the P3 metacone polymorphism across *E. mediotuber* and *E. osbornianus* over
several million years is difficult to explain by genetic drift alone. While some fluctuation was
observed in the character state frequency throughout the range of *E. osbornianus*, the
polymorphism remained at reasonably intermediate frequency, with >30% and <60% of

individuals displaying the metacone throughout the duration of the lineage (Thewissen 1992).
NFDS provides one possible explanation for this maintenance. Competition for food resources
could provide negative, frequency-dependent feedback on the frequency of each P3 morph if
metacone presence yielded different fitness outcomes related to the processing of some limiting
food resource as a function of population density. Several modes of competition have long been
recognized as an important cause of NFDS (Antonovics and Kareiva 1988, Dijikstra and Border
2018). The most relevant mode for P3 morphology might be resource competition associated
with density-dependence. Alternatively, the persistence of the P3 polymorphism might instead
result from some form of bet-hedging or response to soft selection in the presence of
environmental heterogeneity over either space or time. Whatever maintained this polymorphism,
the results here demonstrate how sorting from ancestral stock shape the phenotypes displayed by
incipient descendant lineages. The fixation of monomorphic P3 forms in the descendant lineages
E. major, *E. parvus*, and *E. superstes* likely reflects either adaptation or drift from standing
variation present in the ancestral lineages. If some form of balancing selection maintained P3
polymorphism in *E. mediotuber* and *E. osbornianus*, it is possible that demographic bottlenecks
encountered during speciation of the three descendant lineages lead to random fixation. More
exploration into the overall levels of phenotypic diversity and geography of these descendant
lineages would help to examine the feasibility of this explanation.

Barycrinus phylogeny. *Barycrinus* displays several multiple-budding events throughout its
evolutionary history (Fig. 2). In particular, *B. rhombiferus* is identified as having been a highly
prolific lineage, with four immediate descendants. This is consistent with its status in Gahn and
Kammer's (2002) analysis, which found the placement of *B. rhombiferus* to be highly uncertain,
leading the authors to identify the lineage as a "rogue taxon". The authors interpreted this to
suggest that this uncertainty may have resulted from *B. rhombiferus* having itself given rise to
multiple later-occurring *Barycrinus* lineages. Based on patterns in the inheritance of ancestral
character states, Gahn and Kammer directly suggested that *B. rhombiferus* was likely the direct
ancestor of *B. magister*, *B. spectabilis*, and *B. scitulus*. They also suggested a possible close link
between *B. rhombiferus* and *B. spurius*, going on to speculate that *B. rhombiferus* may be
ancestral to most other *Barycrinus* lineages. Nevertheless, Gahn and Kammer's analysis, which
was relied only on traditional cladistic methods, was very poorly resolved. Given the pattern of
multiple buddings found here, this lack of resolution is not surprising. Nevertheless, the
congruence of our results with the original authors' qualitative interpretations is encouraging.

Barycrinus underwent a rapid radiation while drawing upon a stock of ancestral variation
maintained by *B. rhombiferus* and, to a lesser extent, *B. spurius*. This dynamic provides a small-
scale demonstration of how the filtering of phenotypic variation across budding species events
might shape patterns at deeper timescales. *B. rhombiferus* gave rise to four descendant lineages.
Two of these, *B. spectabilis* and *B. scitulus*, displayed a small fraction of the variation displayed
by their ancestor as well as significantly reduced durations. If each of these incipient lineages
occupied the same ecological landscape as *B. rhombiferus*, they may have been at a competitive
disadvantage (at the lineage level) which would explain their shorter durations. The high
phenotypic variability displayed by *B. rhombiferus* could stem from environmental heterogeneity

caused by broader niche occupancy. If so, the increased survivorship displayed by generalist
crinoids (Kammer et al. 1997) might be explained by increased bet-hedging capability displayed
by generalists that are able to maintain pools of phenotypic variation.

B. magister and *B. spurius* experienced longer durations than their siblings. This could be
explained in *B. magister* by its evolution of additional derived character states not present in *B.*
rhombiferus, which may be indicative of an escape to distinct ecological conditions. While
crinoids, as generalist suspension feeders, often occupy very similar ecological niches, a
spectrum of niche differentiation does exist (Ausich 1980). The enhanced duration and additional
speciations produced by *B. spurius* might be explained by its continued maintenance of much of
the variation possessed by *B. rhombiferus* as well as its derivation of several new characters.
Nevertheless, while interesting to speculate, more ecological information and careful testing of
extended phenotypic datasets is needed to more rigorously explain the distinct causes underlying
the patterns displayed by these lineages. The analysis here represent only a starting point from
which to draw deeper links between the evolution of phenotypic variation across species bounds
and the ecological strategies employed by crinoid species.

Random versus adaptive sorting of ancestral variation. Budding speciation events might often
yield population bottlenecks. If the bottleneck is strong enough, different character states would
be expected to fix when small sub-populations are drawn from polymorphic ancestral
populations over repeated trials (Fig. 3a). Alternatively, sufficiently strong positive selection
would fix the same character state over repeated trials if incipient lineages repeatedly move out
of the ancestral niche and into the same derived niche (Fig. 3b). I leveraged the repeated budding
events displayed across *Barycrinus* to identify whether ancestral variation tended to sort
randomly (Fig. 3a) or adaptively (Fig. 3b). Based on the permutation tests, it was not possible to
distinguish the pattern by which the descendants of *B. rhombiferus* and *B. spurius*, sorted and
fixed variation from their respective ancestors from the expectation under drift induced by
bottlenecking (Tables 1 and 2). None of the pairs of descendants of either lineage showed
statistically significant signs of parallel nor divergent speciation. Nevertheless, *B. spectabilis* and
B. magister were somewhat more similar than expected under random sorting, while *B.*
spectabilis and *B. spurius* were more divergent than expected. The lack of significance could be
a consequence of the small pool of traits included in this dataset and so further tests are needed to
see whether more comprehensive phenotypic sampling would distinguish the observed patterns
from drift.

The occurrence of polymorphisms in botyrocrinid traits that are likely functionally important,
such as rami, columnal, and plate morphology, might indicate the possibility of an ecological
role for maintained, trans-specific polymorphisms. If ecology played a role in the maintenance of
these polymorphisms, their maintenance and fixation in different descendant lineages may have
yielded a landscape of distinct ecological consequences. Throughout the radiation of *Barycrinus*,
variation maintained within ancestral lineages filtered down through a series of successively less
variable descendants (Fig. 2). The only exceptions lie in the emergence of *B. rhombiferus* from
C ibericus and *C. rex* from the hypothetical descendant of *C. ibericus*. The sudden and almost
coincident emergence of high phenotypic variability in *B. rhombiferus* and *C. rex* from the
highly monomorphic *C. ibericus* is mysterious and begs closer examination in future work. The

descendants of polymorphic ancestors are universally less variable than their ancestors. In all,
seven total lineages formed across *Barycrinus* by filtering variation maintained over millions of
years by *B. rhombiferus* and *B. spurius*. While some of the variation displayed by *B. rhombiferus*
may not have persisted over its entire range, the lineage did maintain high levels of
polymorphism throughout its existence. Each descendant lineage that budded from *B.*
rhombiferus and *B. spurius* might be characterized as a unique natural experiment, randomly
fixing ancestral variation and either persisting or perishing in its new habitat. More ecological
information would be needed to understand the specific phenotypic determinants of persistence
and extinction among the varied *Barycrinus* offspring.

Discussion

Budding, phylogenetic pattern, and paleobiological process. The analyses presented here show
how the traditional, bifurcating, cladograms imposed by pattern cladistics on paleontology are
woefully inadequate in their representation of evolutionary relationships among densely sampled
clades of fossil organisms. The multiple-budding events inferred in both *Ectocion* and
botyrocrinids cannot be represented meaningfully in a cladogram, resulting only in the
appearance of "rogue" taxa and polytomies. Biological reality demands the accommodation of
diverse speciation modes when reconstructing phylogeny in the fossil record. Moving forward, it
is important to consider how diverse biological processes such as multiple buddings, anagenesis,
and even hybridization (Ausich and Meyer 1992) might shape relationships among fossil
lineages. Although substantial advances have been made recently in the evaluation and
representation of a greater diversity of speciation modes (Bapst and Hopkins 2017, Stadler et al.
2018, Parins-Fukuchi et al. 2019, Wright et al. 2021, Wright et al. 2022), many of these still
operate from the premise that most relationships tend to be bifurcating and that hypothetical
ancestors predominately link fossil lineages. This is largely a matter of implementation,
stemming from the fact that most advances in phylogenetic tree searching over the past several
decades have focused on inferring trees between contemporaneous lineages and perhaps also the
pattern cladists' denial of the identifiability of ancestral taxa (Engelmann and Wiley 1977). It has
also been pointed out that it is more scientific to expose hypotheses of direct ancestry to
falsification than to "assume taxa are not ancestors until proven otherwise" (Polly 1997). Lastly,
in the absence of overturning information (which may also exist) it is simply more parsimonious
to initially assume hypotheses of direct ancestry, since these do not demand the *ad-hoc*
construction of unobserved "ghost" lineages to explain phylogeny. As a result, paleobiology
might benefit from algorithms that begin from the premise that taxa originating at different times
represent ancestor-descendent sequences *a priori*, and allow character data to overturn this
starting assumption when necessary.

Gahn and Kammer (2002) provided an excellent example of how a poorly resolved cladistic
analysis might be supplemented with extensive natural history knowledge to derive deep
evolutionary insights. However, these should ideally be supported by quantitative evidence and
thorough graphical explorations. Once patterns can be reconstructed with a reasonably high
degree of confidence, it becomes possible to narrow interpretations into a smaller range of

possible explanatory, process-driven, scenarios (Fisher 1981). Evidence supporting alternative
scenarios consistent with the reconstructed pattern might then be weighed using statistical
criteria. Nevertheless, the first step is to further develop theory and methods supporting the
reconstruction of ancestor-descendant relationships. While much important work has already
been done in this area (see references in methods section), the results here especially highlight
the importance of a continued rethinking of paleobiological phylogenetics and encourage a
further untethering of the field from the shackles of cladistic dogma.

Ancestral polymorphism, parallel speciation, and adaptive radiation. A major issue in the
biology of adaptive radiation has been understanding how ecology shapes the phenotypes
displayed by incipient species in the process and aftermath of adaptive radiation. Several authors
have highlighted the role of parallel evolution in shaping these episodes (Schluter and Nagel
1995). Under this scenario, rapidly forming species evolve similar traits as they radiate into
simlar ecological niches. This model of adaptive radiation has been thought to be incompatible
with the classic model of "peripatric" speciation (Mayr 1963), wherein species form rapidly
during founder events. Paleobiologists might be inclined to call these budding events. The
incompatibility between these two models presumably stems from the small population sizes
displayed by the descendants that result from these budding, founder events. This would
decrease the efficiency with which natural selection within each new population can fix similar
mutations in parallel among newly formed species.

While many examples of parallel speciation exist in the neontological literature, the
paleobiological literature remains rife with examples of budding, suggesting a role for peripatry
in the formation of new lineages. Careful examination of how variation is filtered and maintained
at different taxonomic levels across varying ecological contexts might provide the key to
reconciliation between these apparently conflicting views. Adaptation that occurs when
populations move into new habitats can be fuelled by large pools of standing ancestral variation
(e.g., McGee 2020). If selection is strong enough among the descendant lineages, advantageous
alleles may still overcome the effects of drift in small, peripherally isolated populations, allowing
ancestral variation to fix in beneficial ways (Fig. 3b). This demands that the ancestral and
descendant lineages exist in distinct ecological conditions, since the maintenance of variation in
the ancestral population would require a different, balancing, selective regime than the
positively-selective descendants. This type of dynamic may be shown between large, marine,
ancestral populations of sticklebacks and their smaller descendant freshwater populations (e.g.,
Schluter and Conte 2009). Species selection also may be able to generate a similar pattern when
the effect of drift is too great for selection to operate at the population level. Variation in
peripherally-isolated populations might become fixed randomly, with only populations that
stochastically fixed beneficial alleles able to persist. The *Barycrinus* analyses here are more
consistent with the latter scenario, which showed that the pattern of character state sorting in
descendant lineage cannot be distinguished from drift. Each of these alternatives would yield
similar patterns to parallel speciation, with the only difference being the level at which variation
filters out deleterious variation. While, in sticklebacks, population-level positive selection may
have driven the appearance of parallel speciation, the *Barycrinus* analysis illustrates the possible

role for species selection in generating similar patterns by differentially culling descendant
lineages that each inherited a random suite of characters from their ancestor.

Long term maintenance of polymorphisms. The trans-specific maintenance of polymorphic char-
acters creates a natural link between the population processes maintaining phenotypic and ge-
netic variation and the macroevolutionary processes shaping the evolution of lineages. Each of
the datasets examined here displays polymorphisms maintained over long timescales, both within
and between species. The polymorphisms present within lineages were confirmed or implied by
the original studies to have been present throughout the duration of each lineage, even though
trait frequencies may have oscillated over time in some cases. The true temporal persistence of
these polymorphisms was also supported by their frequent appearance across species boundaries.
While polymorphisms often sorted into monomorphic descendant lineages, they also persisted in
several cases, such as the transition from *Ectocion mediotuber* to *Ectocion osbornianus* or
Barycrinus rhombiferus and *Barycrinus spurius*. Patterns in the maintenance and sorting of poly-
morphic traits across species boundaries in *Barycrinus* highlight how variation at the population
level stochastically generates variation between species. The transformation of intrapopulation
into trans-specific variation paves the way for higher-level processes, such as species selection
(Stanley 1975), to operate. Because evolution is fundamentally rooted in the study of biological
variation, it makes sense that understanding its persistence across scales provides the indispens-
able link between population processes and paleobiological patterns. Further developing this
framework may provide the key to developing a true understanding of macroevolutionary
process and how it connects to the mechanisms of population genetics.

Linking population and macroevolutionary processes. When maintained over long periods
against a consistent backdrop of balancing selection, lineages that maintain polymorphic
variation will be better suited to persist compared to lineages that lose diversity to drift. It has
been argued that the basic prerequisite for selection to operate above the species level is that
phenotypic variation be fixed within lineages, but variable between lineages (Stearns 1986).
However, the persistence of polymorphisms maintained over long periods within and distributed
between fossil lineages adds additional complexity. When balancing selection is the maintaining
force, even lineages that display variation may be subject to species selection under certain
conditions. These polymorphic lineages display what might be viewed as a lineage-level
analogue to heterozygote advantage (left side of Fig. 4). Under this scenario, bet-hedging in the
face of spatial or temporal variation in environment and/or NFDS at the population level
maintains stability in trait or genotype frequencies that facilitates higher-level dynamics, such as
species selection, to operate simultaneously. In this case, the necessary condition for species
selection to operate would be ecologically-induced maintenance of polymorphism (potentially
both within and across lineages), rather than a total lack of intraspecific variation. On the other
hand, lineages that fix variation from polymorphic ancestors may achieve greater persistence by
escaping into a new ecological niche that does not impose balancing selection (Fig. 4). Overall, it
is important to consider both inter- and intraspecific variation and the ecological contexts within
which they are maintained and sorted among lineages when exploring higher-level evolutionary
dynamics.

While the level at which variation is displayed remains a critical point in identifying the
boundary conditions under which higher-level selection can operate, perhaps an even more
crucial point lies in considering the selective and ecological contexts occupied by different
lineages. Rather than demanding the absence of variation within species, the conditions allowing
species-level selection may instead be specified by comparing the distribution of variation within
species (poly- vs. monomorphic) to the ecological conditions occupied across species (e.g., does
competition or predation universally impose frequency-dependent fitness effects within
species?). The longer durations exhibited by *B. rhombiferus* and *B. spurius* may stem from
higher species-level fitness for polymorphic taxa, which could be driven by balancing forces
such as environmental variability or negative frequency-dependence resulting from competition
or predation. If intra-specific polymorphism is maintained by environmental heterogeneity
associated with more generalist ecologies, the new niches occupied by successful monomorphic
lineages should tend to be more specialized. This would yield distinct macroevolutionary
dynamics, with generalist polymorphic lineages displaying high lineage survivorship and
specialist monomorphic lineages displaying rapid species turnover. This dynamic would explain
the persistence of polymorphism in the fossil record by the ecological dynamics displayed by
Mississippian crinoids (Kammer et al. 1997)

It has long been hypothesized that biased patterns of extinction among new species might shape
macroevolutionary patterns observed in the fossil record by culling "ephemeral" species before
they are able to fossilize (Raup and Stanley 1978; pp. 105, Stanley 1979, Rosenblum et al. 2012,
Rabosky 2013). Such biased extinction patterns could arise when new lineages are unable to
adapt rapidly enough to new environmental conditions (De Lisle et al. 2021). Meanwhile,
paleontological work has suggested that high morphological variation within species might help
fuel lineage radiations (Webster 2007). If drawn from highly-diverse ancestral populations, the
variation displayed by rapidly radiating descendant lineages will become sorted through a mix of
stochasticity and adaptation, potentially occurring at multiple levels. Incipient species that are
able to draw upon a large well of ancestral variation may stand a greater chance of inheriting
"rescue" alleles that enable them to escape extinction long enough to avoid non-preservation in
the fossil record due to ephemerality.

The "natural experiments" displayed by descendant *Barycrinus* species illustrate how filtering of
ancestral variation, a process that occurs at the population level, might form the basic conditions
and raw material for macroevolutionary processes, such as species selection, to operate. Random
sorting of ancestral variation into descendant lineages is a pattern predicted by classic
evolutionary theory (e.g., Simpson 1944, Mayr 1963, Wright 1982). The manner by which this
occurs, when compared against the range of selective conditions inhabited by incipient lineages,
might explain how variation at the population level in highly-diverse ancestral populations might
filter into descendant lineages to shape macroevolutionary dynamics. Recent work has suggested
that gene tree discordance, which is at least partially caused by incomplete lineage sorting
(ILS)-- the random sorting of ancestral variation into descendant lineages-- is greater during
periods of rapid phenotypic innovation, such as during the early evolution of mammals, birds, or
angiosperms (Parins-Fukuchi et al. 2021). This link implies high genetic diversity (and thus
perhaps high phenotypic variation as well) within lineages during the early stages of rapid clade

diversification. Such variation will stand a good chance of sorting in ways that are discordant
with the order of speciation events, yielding two main effects: 1) high gene-tree discordance and
2) the apparent appearance of rapid morphological variation distributed haphazardly across
lineages. The resulting combination of stochastic sorting, parallel speciation, and species
selection may yield the extensive gene-tree discordance and hemiplasy (Avise 2008, Gurrero and
Hahn 2018) that often accompanies rapid radiations. Future work drawing more explicit links
between coalescent expectations under ILS and the sorting of phenotypic ancestral variation
across fossil speciation events may help shed light on how population processes provide the fuel
for large-scale macroevolutionary processes and patterns to operate.

Conclusion. Explaining the patterns observed by paleontologists in the fossil record in terms of
population processes has been a major goal of evolutionary paleobiology since the modern
synthesis (e.g., Simpson 1944, Kermack 1954, Van Valen 1963, Eldredge and Gould 1972,
Fisher 1985). The persistence of phenotypic variation displayed across fossil lineages, while
interesting from a population genetic perspective, may provide even more groundbreaking
insights when considered from a macroevolutionary perspective. The patterns reconstructed here
highlight how further advances in how we model 1) speciation dynamics, including ancestor-
descendant relationships and 2) the filtering of polymorphic phenotypic variation can vastly
increase the scope of evolutionary questions we are able to evaluate in the fossil record. More
work will also be needed at the population level to better understand how frequencies in
polymorphic traits evolve across the stratigraphic ranges of continuous populations. This will
contribute to a stronger quantiative understanding of the processes that maintain biological
variation among long-lived fossil lineages, such as *Ectocion osbornianus* or *Barycrinus*
rhombiferus. Scaling up, understanding how pools of maintained phenotypic variation segregate
during speciation events can provide a crucial link between the mechanisms explored by
population genetics and the patterns historically explored in evolutionary paleobiology.
Developing stronger links across these levels can help provide more cohesive and deeper
explanations of how population-level evolutionary change scales to the luxuriant diversity of
patterns observed throughout the history of life.

Acknowledgements

Several of the core themes present in this manuscript were sparked by a series of engaging
conversations with Locke Rowe. James Saulsbury offered much salient guidance on the
interpretation of the evolutionary morphology and stratigraphy of fossil crinoids as well as
excellent discussion around the macroevolutionary points of the manuscript. Any errors or
overwrought biological interpretations that remain lie solely on my own shoulders. Nathanael
Walker-Hale offered helpful criticism of the manuscript and useful discussion of the population
genetics of persistent polymorphisms. I thank Miriam Ahmad-Gawel for being a constant
sounding board and (sometimes highly) critical discussion partner for my more audacious
hypotheses of evolutionary process and for offering her keen and skilled eye for editing that was

indispensable while hammering the initial draft into a more coherent form. I was supported by
National Science Foundation grant DEB-2217117 during the course of this work.

Figures

Figure 1. Reconstructed ancestor-descendant relationships within *Ectocion*. Bars represent
stratigraphic ranges. Dotted lines indicate genealogical relationship between ancestral and
descendant lineages. Colors reflect presence or absence of the metacone on the P3. Inset
phylogeny represents hypothetical alternative reconstruction that better accommodates pattern in
the evolution of P3 polymorphism. Further evaluation would be needed to distinguish between
these two. Timescale reflects approximate correlation of the discrete stratigraphic subdivisions
with continuous absolute time.

Figure 2. Phylogeny of *Barycrinus*. Bars represent stratigraphic ranges. Dotted lines indicate
genealogical relationship between ancestral and descendant lineages. Width of stratigraphic lines
represents scaled number of polymorphisms—a measure of genetic variation within each lineage.
Shading represents the proportion of character states displayed by each lineage that were also
possessed by its ancestor. Timescale reflects approximate correlation of the discrete stratigraphic
subdivisions with continuous absolute time.

Figure 3. Distribution of simulated allele frequencies over 10 replicated populations while a
balanced ancestral polymorphism is filtered into a budding descendant under two different
adaptive scenarios: A) Polymorphism maintained by negative frequency dependent selection
(NFDS) in a large ancestral population that becomes randomly fixed in bottlenecking
encountered during a budding speciation event. B) Polymorphism maintained by NFDS that
becomes fixed due to positive selection (PS) in a budding descendant that has dispersed into a
new environment. Lineage widths in budding lineage diagrams represent effective population
size. Under scenario A, variation is filtered randomly by drift into the descendant lineages.
Under scenario B, the new regime rapidly fixes one allele/character state according to its new
selective landscape.

Figure 4. Species selection dynamics stemming from a persistently polymorphic ancestral
population. In this hypothetical scenario, negative frequency-dependent selection forms the
selective background in the ancestor. When descendant lineages randomly fix this ancestral
variation during budding speciation events, they demonstrate low survivorship in the ancestral
niche. Only lineages that can escape and radiate into a new niche while fixing ancestral variation
display high survivorship. Ecological opportunity afforded by the new niche may even facilitate
enhanced lineage survivorship and proliferation if the trait fixed in the descendant is congruent
with the selective demands of the new habitat.

Tables

	B. magister	*B. scitulus*	*B. spectabilis*
B. magister	-	-	-
B. scitulus	0.559	-	-
B. spectabilis	-1.512	0.583	-
B. spurius	0.289	0.306	1.285

741 *Table 1*. Pairwise phenotypic distances calculated among characters sorted from polymorphisms
742 present in the ancestor, *Barycrinus rhombiferus* between descendant lineages, relative to null

743 expectation generated under random sorting of ancestral polymorphisms. No pairs were
744 statistically significant at the 2.5% threshold.

	B. stellatus
B. stellatus	-
B. crassibrachiatus	-0.22

747 *Table 2*. Pairwise phenotypic distances calculated among characters sorted from polymorphisms
748 present in the ancestor, *Barycrinus spectabilis* between descendant lineages, relative to null
749 expectation generated under random sorting of ancestral polymorphisms. No pairs were
750 statistically significant at the 2.5% threshold.

References

Alroy, J., 1995. Continuous track analysis: a new phylogenetic and biogeographic method.
Systematic Biology, 44(2), pp.152-178.

Antonovics, J. and Kareiva, P., 1988. Frequency-dependent selection and competition: empirical
approaches. Philosophical Transactions of the Royal Society of London. B, Biological Sciences,
319(1196), pp.601-613.

Ausich, W.I., 1980. A model for niche differentiation in Lower Mississippian crinoid
communities. Journal of Paleontology, pp.273-288.

Ausich, W.I. and Meyer, D.L., 1994. Hybrid crinoids in the fossil record (Early Mississippian,
Phylum Echinodermata). Paleobiology, 20(3), pp.362-367.

Avise, J.C. and Robinson, T.J., 2008. Hemiplasy: a new term in the lexicon of phylogenetics.
Systematic biology, 57(3), pp.503-507.

Aze, T., Ezard, T.H., Purvis, A., Coxall, H.K., Stewart, D.R., Wade, B.S. and Pearson, P.N.,
2011. A phylogeny of Cenozoic macroperforate planktonic foraminifera from fossil data.
Biological Reviews, 86(4), pp.900-927.

Bapst, D.W. and Hopkins, M.J., 2017. Comparing cal3 and other a posteriori time-scaling
approaches in a case study with the pterocephaliid trilobites. Paleobiology, 43(1), pp.49-67.

Barrett, R.D. and Schluter, D., 2008. Adaptation from standing genetic variation. Trends in
ecology & evolution, 23(1), pp.38-44.

Benton, M.J. and Pearson, P.N., 2001. Speciation in the fossil record. Trends in Ecology &
Evolution, 16(7), pp.405-411.

Brothwell, D.R. ed., 2014. Dental Anthropology: Volume V: Society for the Study of Human
Biology. 5, pp.286. Elsevier.

Caetano, D.S. and Quental, T.B., 2022. How Important is Budding Speciation for Comparative
Studies?. bioRxiv, pp.2022-05.

Chakraborty, M. and Fry, J.D., 2016. Evidence that environmental heterogeneity maintains a
detoxifying enzyme polymorphism in Drosophila melanogaster. Current Biology, 26(2), pp.219-
223.

Charlesworth, D., 2006. Balancing selection and its effects on sequences in nearby genome
regions. PLoS Genetics, 2(4), p.e64.

Christiansen, F.B., 1975. Hard and soft selection in a subdivided population. The American
Naturalist, 109(965), pp.11-16.

Clarke, B.C., 1979. The evolution of genetic diversity. Proceedings of the Royal Society of
London. Series B. Biological Sciences, 205(1161), pp.453-474.

De Lisle, S.P., Punzalan, D., Rollinson, N. and Rowe, L., 2021. Extinction and the temporal
distribution of macroevolutionary bursts. Journal of Evolutionary Biology, 34(2), pp.380-390.

Dijkstra, P.D. and Border, S.E., 2018. How does male–male competition generate negative
frequency-dependent selection and disruptive selection during speciation?. Current Zoology,
64(1), pp.89-99.

Engelmann, G.F. and Wiley, E.O., 1977. The place of ancestor-descendant relationships in
phylogeny reconstruction. Systematic Biology, 26(1), pp.1-11.

Eldredge, N. and Gould, S.J., 1972. Punctuated equilibria: an alternative to phyletic gradualism.
Models in Paleobiology, 1972, pp.82-115.

Fisher, D.C., 1991. Phylogenetic analysis and its application in evolutionary paleobiology. Short
Courses in Paleontology, 4, pp.103-122.

Fisher, D.C., 2008. Stratocladistics: integrating temporal data and character data in phylogenetic
inference. Annual Review of Ecology, Evolution, and Systematics, 39, pp.365-385.

Fitzpatrick, M.J., Feder, E., Rowe, L. and Sokolowski, M.B., 2007. Maintaining a behaviour
polymorphism by frequency-dependent selection on a single gene. Nature, 447(7141), pp.210-
212.

Fisher, D.C., 1981. The role of functional analysis in phylogenetic inference: examples from the
history of the Xiphosura. American Zoologist, 21(1), pp.47-62.

Fisher, D.C., 1985. Evolutionary morphology: beyond the analogous, the anecdotal, and the ad
hoc. Paleobiology, 11(1), pp.120-138.

Foote, M., 1996. On the probability of ancestors in the fossil record. Paleobiology, 22(2),
pp.141-151.

Gallet, R., Froissart, R. and Ravigné, V., 2018. Experimental demonstration of the impact of
hard and soft selection regimes on polymorphism maintenance in spatially heterogeneous
environments. Evolution, 72(8), pp.1677-1688.

Gahn, F.J. and Kammer, T.W., 2002. The cladid crinoid Barycrinus from the Burlington
Limestone (early Osagean) and the phylogenetics of Mississippian botryocrinids. Journal of
Paleontology, 76(1), pp.123-133.

Gilbert, C.C., 2013. Cladistic analysis of extant and fossil African papionins using craniodental
data. Journal of human evolution, 64(5), pp.399-433.

Gingerich, P.D., 1979. The stratophenetic approach to phylogeny reconstruction in vertebrate
paleontology. In Phylogenetic analysis and paleontology (pp. 41-78). Columbia University Press.
New York.

Gingerich, P.D., 1974. Stratigraphic record of early Eocene Hyopsodus and the geometry of
mammalian phylogeny. Nature, 248(5444), pp.107-109.

Goldberg, E.E., Kohn, J.R., Lande, R., Robertson, K.A., Smith, S.A. and Igić, B., 2010. Species
selection maintains self-incompatibility. Science, 330(6003), pp.493-495.

Guerrero, R.F. and Hahn, M.W., 2017. Speciation as a sieve for ancestral polymorphism.
Molecular Ecology, 26(20), pp.5362-5368.

Guerrero, R.F. and Hahn, M.W., 2018. Quantifying the risk of hemiplasy in phylogenetic
inference. Proceedings of the National Academy of Sciences, 115(50), pp.12787-12792.

Hittinger, C.T., Gonçalves, P., Sampaio, J.P., Dover, J., Johnston, M. and Rokas, A., 2010.
Remarkably ancient balanced polymorphisms in a multi-locus gene network. Nature, 464(7285),
pp.54-58.

Hughes, A.L. and Nei, M., 1988. Pattern of nucleotide substitution at major histocompatibility
complex class I loci reveals overdominant selection. Nature, 335(6186), pp.167-170.

Igic, B., Bohs, L. and Kohn, J.R., 2006. Ancient polymorphism reveals unidirectional breeding
system shifts. Proceedings of the National Academy of Sciences, 103(5), pp.1359-1363.

Jackson, J.B. and Cheetham, A.H., 1994. Phylogeny reconstruction and the tempo of speciation
in cheilostome Bryozoa. Paleobiology, 20(4), pp.407-423.

Kammer, T.W., Baumiller, T.K. and Ausich, W.I., 1997. Species longevity as a function of niche
breadth: evidence from fossil crinoids. Geology, 25(3), pp.219-222.

Kermack, K.A., 1954. A biometrical study of Micraster coranguinum and M.(Isomicraster)
senonensis. Philosophical Transactions of the Royal Society of London. Series B, Biological
Sciences, pp.375-428.

Klein, J., Sato, A., Nagl, S. and O'hUigín, C., 1998. Molecular trans-species polymorphism.
Annual Review of Ecology and Systematics, 29(1), pp.1-21.

Lai, Y.T., Yeung, C.K., Omland, K.E., Pang, E.L., Hao, Y.U., Liao, B.Y., Cao, H.F., Zhang,
B.W., Yeh, C.F., Hung, C.M. and Hung, H.Y., 2019. Standing genetic variation as the
predominant source for adaptation of a songbird. Proceedings of the National Academy of
Sciences, 116(6), pp.2152-2157.

Lewontin, R.C. and Hubby, J.L., 1966. A molecular approach to the study of genic
heterozygosity in natural populations. II. Amount of variation and degree of heterozygosity in
natural populations of Drosophila pseudoobscura. Genetics, 54(2), p.595.

Mayr, E. 1963. Animal species and evolution. Harvard University Press, Cambridge.

McDonald, J.F. and Ayala, F.J., 1974. Genetic response to environmental heterogeneity. Nature,
250(5467), pp.572-574.

McGee, M.D., Borstein, S.R., Meier, J.I., Marques, D.A., Mwaiko, S., Taabu, A., Kishe, M.A.,
O'Meara, B., Bruggmann, R., Excoffier, L. and Seehausen, O., 2020. The ecological and
genomic basis of explosive adaptive radiation. Nature, 586(7827), pp.75-79.

Muirhead, C.A., Glass, N.L. and Slatkin, M., 2002. Multilocus self-recognition systems in fungi
as a cause of trans-species polymorphism. Genetics, 161(2), pp.633-641.

Parins-Fukuchi, C.T., Stull, G.W. and Smith, S.A., 2021. Phylogenomic conflict coincides with
rapid morphological innovation. Proceedings of the National Academy of Sciences, 118(19),
p.e2023058118.

Parins-Fukuchi, C.T., 2021. Morphological and phylogeographic evidence for budding
speciation: an example in hominins. Biology Letters, 17(1), p.20200754.

Pearson, P.N., 1998. Speciation and extinction asymmetries in paleontological phylogenies:
evidence for evolutionary progress?. Paleobiology, 24(3), pp.305-335.

Pease, J.B., Haak, D.C., Hahn, M.W. and Moyle, L.C., 2016. Phylogenomics reveals three
sources of adaptive variation during a rapid radiation. PLoS biology, 14(2), p.e1002379.

Polly, P.D., 1997. Ancestry and species definition in paleontology: a stratocladistic analysis of
Paleocene-Eocene Viverravidae (Mammalia, Carnivora) from Wyoming. Contributions from the
Museum of Paleontology. 30(1), pp.1-53.

Prout T., 2000. How well does opposing selection maintain variation? In: Singh RS, Krimbas
CB, editors. Evolutionary genetics: from molecules to morphology. Cambridge: Cambridge
University Press. pp.157–181.

Rabosky, D.L., 2013. Diversity-dependence, ecological speciation, and the role of competition in
macroevolution. Annual Review of Ecology, Evolution, and Systematics, 44, pp.481-502.

Raup, D.M. and Gould, S.J., 1974. Stochastic simulation and evolution of morphology-towards a
nomothetic paleontology. Systematic Biology, 23(3),pp.305–322.

Raup, D. and Stanley, S.M., 1978. Principles of Paleontology. Macmillan.

Raup, D.M., 1985. Mathematical models of cladogenesis. Paleobiology, 11(1), pp.42-52.

Rensch, B., 1959. 6. The Rules of Kladogenesis (Phylogenetic Branching). In Evolution Above
the Species Level (pp. 97-280). Columbia University Press.

Rosenblum, E.B., Sarver, B.A., Brown, J.W., Des Roches, S., Hardwick, K.M., Hether, T.D.,
Eastman, J.M., Pennell, M.W. and Harmon, L.J., 2012. Goldilocks meets Santa Rosalia: an
ephemeral speciation model explains patterns of diversification across time scales. Evolutionary
Biology, 39, pp.255-261.

Saitou, N. and Nei, M., 1987. The neighbor-joining method: a new method for reconstructing
phylogenetic trees. Molecular biology and evolution, 4(4), pp.406-425.

Schluter, D., 2000. Ecological character displacement in adaptive radiation. The American
Naturalist, 156(S4), pp.S4-S16.

Schluter, D. and Conte, G.L., 2009. Genetics and ecological speciation. Proceedings of the
National Academy of Sciences, 106(supp. 1), pp.9955-9962.

Schluter, D. and Nagel, L.M., 1995. Parallel speciation by natural selection. The American
Naturalist, 146(2), pp.292-301.

Schopf, T.J. and Gooch, J.L., 1972. A natural experiment to test the hypothesis that loss of
genetic variability was responsible for mass extinctions of the fossil record. The Journal of
Geology, 80(4), pp.481-483.

Sicard, A., Kappel, C., Lee, Y.W., Woźniak, N.J., Marona, C., Stinchcombe, J.R., Wright, S.I.
and Lenhard, M., 2016. Standing genetic variation in a tissue-specific enhancer underlies selfing-
syndrome evolution in Capsella. Proceedings of the National Academy of Sciences, 113(48),
pp.13911-13916.

Simpson, G.G., 1944. Tempo and Mode in Evolution. Columbia University Press. New York.

Stadler, T., Gavryushkina, A., Warnock, R.C., Drummond, A.J. and Heath, T.A., 2018. The
fossilized birth-death model for the analysis of stratigraphic range data under different speciation
modes. Journal of Theoretical Biology, 447, pp.41-55.

Stanley, S.M., 1975. A theory of evolution above the species level. Proceedings of the National
Academy of Sciences, 72(2), pp.646-650.

Stanley SM. 1979. Macroevolution: Pattern and Process. San Francisco, CA: Freeman

Stearns, S.C., 1986. Natural selection and fitness, adaptation and constraint. In: Raup D. M. and
Jablonski, D., eds., Patterns and Processes in the History of Life: Report of the Dahlem
Workshop on Patterns and Processes in the History of Life, pp. 23-44. Springer Berlin
Heidelberg.

Takahashi, Y., Yoshimura, J., Morita, S. and Watanabe, M., 2010. Negative frequency-
dependent selection in female color polymorphism of a damselfly. Evolution, 64(12), pp.3620-
3628.

Thewissen, J.G.M., 1992. Temporal data in phylogenetic systematics: an example from the
mammalian fossil record. Journal of Paleontology, 66(1), pp.1-8.

Thompson, K.A., Osmond, M.M. and Schluter, D., 2019. Parallel genetic evolution and
speciation from standing variation. Evolution Letters, 3(2), pp.129-141.

Trueman, J.W., 2010. A new cladistic analysis of Homo floresiensis. Journal of human
evolution, 59(2), pp.223-226.

Van Valen, L., 1969. Variation genetics of extinct animals. The American Naturalist, 103(931),
pp.193-224.

Van Valen, L., 1975. Group selection, sex, and fossils. Evolution, pp.87-94.

Van Valen, L., 1963. Selection in natural populations: *Merychippus primus*, a fossil horse.
Nature, 197, pp.1181-1183.

Wagner, P.J., 1998. A likelihood approach for evaluating estimates of phylogenetic relationships
among fossil taxa. Paleobiology, 24(4), pp.430-449.

Wagner, P.J., 2000. Phylogenetic analyses and the fossil record: tests and inferences, hypotheses
and models. Paleobiology, 26(S4), pp.341-371.

Wagner, P.J., Erwin, D.H., 1995. Phylogenetic patterns as tests of speciation models. In: Anstey,
R.L, editor. New approaches to speciation in the fossil record. Columbia University Press, New
York, pp.87-122.

Warnock, R.C., Heath, T.A. and Stadler, T., 2020. Assessing the impact of incomplete species
sampling on estimates of speciation and extinction rates. Paleobiology, 46(2), pp.137-157.

Wiens, J.J., 1995. Polymorphic characters in phylogenetic systematics. Systematic Biology,
44(4), pp.482-500.

Wiens, J.J., 1999. Polymorphism in systematics and comparative biology. Annual Review of
Ecology and Systematics, 30(1), pp.327-362.

Whiting, E.T., Steadman, D.W. and Vliet, K.A., 2016. Cranial polymorphism and systematics of
Miocene and living Alligator in North America. Journal of Herpetology, 50(2), pp.306-315.

Webster, M., 2007. A Cambrian peak in morphological variation within trilobite species.
Science, 317(5837), pp.499-502.

Wood, A.R., Zelditch, M.L., Rountrey, A.N., Eiting, T.P., Sheets, H.D. and Gingerich, P.D.,
2007. Multivariate stasis in the dental morphology of the Paleocene-Eocene condylarth Ectocion.
Paleobiology, 33(2), pp.248-260.

Wright, A.M., Bapst, D.W., Barido-Sottani, J. and Warnock, R.C., 2022. Integrating fossil
observations into phylogenetics using the fossilized birth–death model. Annual Review of
Ecology, Evolution, and Systematics, 53, pp.251-273.

Wright, A.M., Wagner, P.J. and Wright, D.F., 2021. Testing character evolution models in
phylogenetic paleobiology: a case study with Cambrian echinoderms. Cambridge University
Press. Cambridge.

Barycrinus stellatus

B. punctus

B. spurius

B. crassibrachiatus

B. magister

B. rhombiferus

B. scitulus

B. spectabilis

Costalocrinus ibericus

C. sentosus

C. rex

C. cornutus

Meniscocrinus magnitubus

Prop. anc.
char. states

1.0
0.9
0.8
0.7
0.6
0.5

360   340   320   300

ma