Myosin in autoinhibited *off* state(s), stabilized by mavacamten, can be recruited via inotropic effectors

Weikang Ma[1,#*], Carlos L. del Rio[2,#*], Lin Qi[3], Momcilo Prodanovic[4,5], Srboljub Mijailovich[5], Christopher Zambataro[2], Henry Gong[3], Rafael Shimkunas[2], Sampath Gollapudi[2], Suman Nag[2*], and Thomas C. Irving[1]

1. BioCAT, Department of Biology, Illinois Institute of Technology, Chicago, IL, USA
2. Cardiovascular Drug Discovery, Bristol Myers Squibb, Brisbane, CA 94005
3. Department of Biology, Illinois Institute of Technology, Chicago, IL, USA
4. Institute for Information Technologies, University of Kragujevac, Kragujevac, Serbia
5. FilamenTech, Inc., Newtown, MA 02458, USA

\# authors contributed equally

Short title: mavacamten-stabilized myosins are rectuitable

* Corresponding authors:

Suman Nag, sumannag@gmail.com, 3160 Porter Drive Palo Alto, CA 94304;
Carlos L. del Rio, cdelrio.heart@gmail.com, 200 Technology Square Cambridge, MA 02139;
Weikang Ma, wma6@iit.edu, 9700 S Cass Ave Argonne, IL 60439

Total word count: 10725

Abstract

Rationale: Mavacamten is a novel, FDA-approved, small molecule therapeutic designed to regulate cardiac function at the sarcomere level by selectively but reversibly inhibiting the enzymatic activity of myosin. It shifts myosin towards ordered *off* states close to the thick filament backbone.

Objective: The purpose of this study is to investigate whether mavacamten permanently sequester these myosin heads in the *off* state(s) and can these heads be recruited in response to physiological stimuli when required to boost cardiac output.

Methods and Results: We show that cardiac myosins stabilized in these *off* state(s) by mavacamten are recruitable by 1) Ca^{2+}, 2) increased chronotropy (heart rate), 3) stretch, and 4) β-adrenergic (β-AR) stimulation, all known physiological inotropic effectors. At the molecular level, we show that Ca^{2+} increases myosin ATPase activity by shifting mavacamten-stabilized myosin heads from the inactive super-relaxed (SRX) state to the active disordered relaxed (DRX) state. At the myofilament level, both Ca^{2+} and passive lengthening can shift mavacamten-ordered *off* myosin heads from positions close to the thick filament backbone to disordered *on* states closer to the thin filaments. In isolated rat cardiomyocytes, increased stimulation rates enhanced shortening fraction in mavacamten-treated cells. This observation was confirmed *in vivo* in telemetered rats, where left-ventricular dP/dt_{max}, an index of inotropy, increased with heart rate in mavacamten treated animals. Finally, we show that β-AR stimulation *in vivo* increases left-ventricular function and stroke volume in the setting of mavacamten.

Conclusions: Our data demonstrate that the mavacamten-promoted *off* states of myosin in the thick filament are activable, thus leading to the preservation of the cardiac reserve. These results provide a potential mechanistic explanation, beyond mere LV outflow tract obstruction removal, for the clinical observation of increased peak oxygen uptake (pVO_2) with exercise in HCM patients receiving mavacamten.

Introduction

The heart is the powerhouse of the circulatory system, and its stroke volume is precisely regulated by various inotropic effectors to fulfill the body's demands. Four foremost positive inotropic effectors exist in the heart: 1) increase in calcium (Ca^{2+}) concentration via the action of L-type Ca^{2+} channels[1]; or 2) via increased heart rate or frequency, known as the Bowditch effect or the Treppe phenomenon[2, 3]; 3) increased diastolic ventricular preload, also known as length-dependent activation (LDA)[4]; and 4) β-adrenergic stimulation that increases cyclic AMP, activating protein kinase A (PKA) which subsequently phosphorylates regulatory proteins in the heart muscle resulting in increased cardiac contractility and, consequently, cardiac output[5]. The ability to maintain these positive inotropic effects is vital for the heart to function properly in response to increased physiological demands, and deficits in the heart response to both inotropes and increased demand is a common feature of both cardiomyopathies[6] and/or acquired cardiovascular diseases[7].

Such is the case in hypertrophic cardiomyopathy (HCM), a myocardial disorder characterized by left ventricular (LV) hypertrophy, hyperdynamic contraction, and impaired relaxation[8]. The cardiac changes in HCM often manifest as hindered exercise capacity, as reflected by reduced peak oxygen uptake (pVO_2) and cardiac output recruitment. The root causes of the disease in a significant fraction of HCM patients are within the sarcomere, the basic contractile apparatus of the heart. Approximately 45% of affected individuals, and a significant portion of those with a family history of clinical disease, have a mutation(s) in one or more genes encoding sarcomere proteins[8]. It has been shown that these sarcomeric mutations lead to excessive actin-myosin interactions, or cross-bridges, arising from a pathogenic shift of myosin heads from reserve and energy-sparing *off* states towards *on* states[9, 10]. Other mutations outside of the sarcomere also contribute to HCM[11].

Mavacamten (commercially known as Camzyos) is an FDA approved small molecule developed to address this pathogenicity by selectively attenuating the ATPase activity of cardiac myosin[12]. This targeted myosin inhibition results in a novel pharmacological profile that is capable of normalizing contraction and relaxation properties, as demonstrated in both preclinical and phase-II and phase-III clinical studies, showing great promise for the treatment of HCM[13-15]. More importantly, despite reducing the intrinsic activity of myosin, the sarcomere motor of the heart, mavacamten has been shown to increase pVO_2 in patients with obstructive HCM[14, 16] and

in a subset of non-obstructive HCM patients[17]. Thus, as pVO$_2$ reflects the ability to recruit

forward flow and stroke volume during exercise, the mechanistic basis for these effects remains

to be elucidated and may lie within the molecular impacts of mavacamten in the sarcomere.

Although the atomic structure of myosin with mavacamten bound has yet to be published,

electron microscopy[13] and SAXS[18] studies show that mavacamten stabilizes a compact folded-

back structure of myosin[13] that is reminiscent of the interacting-heads motif (IHM) state of

myosin that is seen in relaxed skeletal, cardiac, and smooth muscle myosin from diverse

organisms[19]. Additionally, mavacamten stabilizes the biochemically energy-sparing auto-

inhibited super-relaxed (SRX) state(s) of myosin[13, 18, 20, 21]. Consistent with these *in vitro*

observations, *ex vivo* low-angle X-ray diffraction studies of porcine cardiac muscle fibers after

mavacamten treatment show more heads sequestered closer to the thick filaments, thereby

increasing the quasi-helical ordering of myosin heads along the thick filament backbone[13, 22, 23].

Whether the SRX state(s) is always structurally identical to the canonical IHM state of myosin[10,

19, 24], or whether the structurally defined sequestered heads are either in the biochemically

defined SRX state or necessarily adopt the IHM configuration remain open questions in the field

(see[25]). Regardless, altogether, this body of evidence suggests that mavacamten induces a

population of myosin heads to adopt sequestered *off* state(s) with low probability (availability) to

participate in contraction under normal physiological conditions. This raises the question,

however, whether these mavacamten-stabilized *off* state(s) can be recruited during increased

workload in order to increase forward flow and meet the physiological demands of mavacamten-

treated patients under physical stress.

Some hints of this potential recruitability can be found in the literature[12, 26], where

mavacamten has been shown to preserve, at least partially, responsiveness to exercise and β-AR

stimulation in *vivo*, as well as length-dependent activation (LDA) in *vitro*[22, 27, 28]. LDA is a key

feature of cardiac muscle, wherein the muscle responds with increased Ca^{2+} sensitivity and force

to increases in length (preload) and is the cellular basis of the Frank-Starling mechanism,

whereby the heart increases its cardiac output in response to an increase in preload[4].

Functionally, mavacamten is shown to preserve, at least partially, the systolic gains of LDA[22, 27,

28], suggesting that the myosin heads sequestered in the *off* SRX state(s) induced by mavacamten

are potentially available for recruitment. To test the this hypothesis, we designed experiments

ranging from molecular level studies on cardiac myosin synthetic thick filaments and cardiac

myofibrils using SRX/DRX and ATPase assays to sarcomere level studies on permeabilized

porcine myocardium using small-angle X-ray diffraction, computational simulations, and in *vivo*

experiments in conscious and anesthetized animals to demonstrate that the myosin sequestered

off states stabilized by mavacamten are controllable and can be recruited, at least partially, by 1)

direct increase in Ca^{2+} in *vitro* or 2) via increase in Ca^{2+} due to increased heart rate in *vivo*, 3)

sarcomere stretch as well as 4) β-AR stimulation.

Methods

β-cardiac full-length myosin protein purification. β-cardiac full-length myosin from the

bovine left ventricle is isolated following established methods described previously[29]. Following

this, full-length myosin is dialyzed in a buffer containing 10 mM PIPES (pH 6.8), 300 mM KCl,

0.5 mM $MgCl_2$, 0.5 mM EGTA, 1 mM $NaHCO_3$, and 1 mM DTT and stored at −80°C. Based on

densitometry analysis of SDS-PAGE, the purity of the myosin preparation varies between 90 and

95%, with negligible actin contamination. Also, the basal myosin ATPase from these preps

always turns out to be 0.03 ± 0.01 s^{-1}, suggesting negligible actin-activation. Bovine cardiac

myofibrils were prepared following the methods described previously[30] and were stored at

−80°C in a buffer containing 10 mM PIPES (pH 6.8), 30 mM KCl, 0.5 mM $MgCl_2$, 0.5 mM

EGTA, 1 mM $NaHCO_3$, and 1 mM DTT.

Reconstitution of myosin thick filaments. Full-length myosin remains fully soluble in a buffer

of high ionic strength (300 mM). However, many previous studies had previously shown that the

full-length myosin could spontaneously self-assemble into bipolar thick filaments when the ionic

strength of the buffer was reduced to 150 mM or below[31, 32]. This method was used to construct

thick filaments with some modifications in this study. Briefly, the ionic strength of the myosin

sample and the myosin concentration were first adjusted to the desired value by diluting it with a

buffer containing 20-mM Tris-HCl (pH 7.4), 0-mM KCl, 1-mM EGTA, 3-mM MgCl2, and 1-

mM DTT. After dilution, the myosin sample was incubated for 2 h on ice to form thick filaments

before using it for experiments. In this study, such reconstituted myosin filaments will be

referred to as STFs. Alternatively, STFs were also made by slowly dialyzing full-length myosin

into a low ionic (150 mM or below) strength buffer. Quantitatively, the measurements reported

in STFs were not different between these two methods.

Steady-state ATPase measurements. Measurements of basal myosin and myofibrillar ATPase activity as a function of Ca^{2+} were performed at 23 °C on a plate-based reader (SpectraMax 96-well) using an enzymatically coupled assay as described previously[33]. The buffer conditions used were 12 mM Pipes (pH 6.8), 2 mM MgCl2, 10 mM KCl, and 1 mM DTT. For the basal ATPase experiments with STFs, a final myosin concentration of 1 µM was used. When using myofibrils, approximately 40% of the total weight was assumed to be from myosin, and accordingly, the myofibril amount was loaded to attain a final myosin concentration of 1 µM. A concentrated stock (20 mM) of mavacamten and MYK-7660 (TF inhibitor) was first prepared using dimethyl sulfoxide (DMSO), and this was used to attain the desired concentration in the final buffer samples. A final DMSO concentration of 2% was achieved in all samples. Data recorded by the instrument-compatible software package, SoftMax Pro, was exported to GraphPad Prism and analyzed.

Biochemical SRX/DRX measurements in reconstituted myosin synthetic thick filament. β-cardiac full-length myosin from the bovine left ventricle was isolated following established methods described elsewhere[29]. Bipolar synthetic thick filaments (STF) from this full-length myosin were prepared (~ 0.4 µM) as described previously[21]. Estimations of the SRX population of myosin preparations were performed using a single ATP turnover kinetic protocol with a fluorescent 2′/3′-O-(N-Methylanthraniloyl) (mant)-ATP probe chased with excess unlabeled ATP[21]. The fluorescence decay profiles were fit to a bi-exponential function to estimate four parameters corresponding to fast and slow phases and the normalized % amplitude and the observed ATP turnover rate of each phase. The fast and the slow phases correspond to the myosin activity in the DRX and SRX states, respectively. Experiments in the both the control and mavacamten group contain 2% DMSO.

X-ray diffraction measurements on permeabilized porcine myocardium. Isolated hearts (N =2) from male Yucatan pig were provided by Exemplar Genetics Inc. Humane euthanasia and tissue collection procedures were approved by the Institutional Animal Care and Use Committees at Exemplar Genetics. The procedures were conducted according to the principles in the "Guide for the Care and Use of Laboratory Animals", Institute of Laboratory Animals Resources, Eighth Edition[34], the Animal Welfare Act as amended, and with accepted American Veterinary Medical Association (AVMA) guidelines at the time of the experiments[35]. Left

ventricular myocardium samples were prepared as described previously[36, 37]. Briefly, samples of

frozen left ventricular myocardium were permeabilized in relaxing solution (2.25 mM Na_2ATP,

3.56 mM $MgCl_2$, 7 mM EGTA, 15 mM sodium phosphocreatine, 91.2 mM potassium

propionate, 20 mM Imidazole, 0.165 mM $CaCl_2$, creatine phosphate kinase 15 U/ml) containing

15 mM 2,3-Butanedione 2-monoxime (BDM) and 1% Triton-X100 and 3% dextran at pH 7) at

room temperature for 2-3h). Muscles were then washed with fresh relaxing solution for ~ 10

minutes, repeated 3 times. The tissue was dissected into ~200 μm diameter fiber bundles and

clipped with aluminum T-clips. The preparations were then stored in cold (4°C) relaxing solution

with 3% dextran for the day's experiments. X-ray diffraction experiments were performed at the

BioCAT beamline 18ID at the Advanced Photon Source, Argonne National Laboratory[38]. The

muscles were incubated in a customized chamber, and experiments were performed at 28 °C to

30 °C. For the results presented in figure 2, the muscle was held at a sarcomere length of 2.3 µm

and the X-ray patterns were collected at seven different Ca^{2+} concentrations (pCa 8, pCa 6.4, pCa

6, pCa 5.8, pCa 5.6, pCa 5.3, and pCa 4.5) in the presence of 100 µM thin filament inhibitor

(MYK-7660) plus 2 µM of mavacamten (mavacamten group) to compare the results from

previous studies in the absence of these compounds[23]. For the results presented in figure 3, two

groups of the X-ray patterns were collected at a sarcomere length of 2.1 µm and then at a

sarcomere length of 2.3 µm in the presence and absence of 50 µM mavacamten respectively. The

X-ray patterns were collected on a MarCCD 165 detector (Rayonix Inc., Evanston IL) with a 1 s

exposure time. 2-3 patterns were collected under each condition. Reflection spacings and

intensities were extracted from these patterns using data reduction programs belonging to the

open-source MuscleX software package developed at BioCAT[39] and averaged. Briefly, the

equatorial reflections were measured by the "Equator" routine in MuscleX as described

previously[40]. The intensities[22] and spacings[41] of meridional and layer line reflections were

measured by the "Projection Traces" routine in MuscleX.

Isolated Myocytes (Bowditch). The rat experiments were done in compliance with protocols

approved by the Institutional Animal Care and Use Committees at Bristol-Myers Squibb

(formerly MyoKardia). Primary left-ventricular cardiomyocytes were isolated were from adult

Sprague Dawley rats via enzymatic digestion in a Langendorff apparatus as previously

described[12]. Subsequently, aliquots of cardiomyocyte suspensions in Tyrode's buffer solution

kept at room temperature (20-22 ℃) were placed into the perfusion chamber on an inverted

microscope, and cardiomyocytes were allowed to settle and adhere to the chamber glass coverslip for 5 minutes. Tyrode's solution containing DMSO (control, with 0.05% DMSO) was then superfused into the chamber at 1.5 mL/min and solution temperature gradually increased to and kept at 31-32 ℃ by in-line pre-heater temperature controller (Cell MicroControls, Norfolk, VA, USA). All measurements were performed at 31-32 ℃. Once at set temperature, cardiomyocyte pacing was started by electrical stimulations between two platinum electrodes at 1 Hz and 6 V for 2 ms (MyoPacer, IonOptix, Westwood, MA, USA). Cardiomyocytes were paced for at least 5 minutes to reach steady state contraction prior to experiments. Solutions containing either DMSO or mavacamten (0.1 μM) were superfused into the chamber for 5 minutes followed by progressive increases in pacing rate (to 2 Hz, 3 Hz, and 4 Hz). Measurements were taken from the last 30 seconds of each pacing/exposure period. Concurrent measurements of cell-length and cytosolic calcium fluorescence were acquired from single cardiomyocytes for each experiment recording.

For calcium fluorescence measurements, cardiomyocytes were incubated with 1 μM Fura-2-acetoxymethyl ester (Fura-2 AM, cell permeant, ThermoFisher) and 0.075% Pluronic F-127 (20% in DMSO, ThermoFisher) for 15 minutes at room temperature. Cardiomyocytes were then washed with fresh Tyrode's solution and incubated for 15 minutes at room temperature to allow for de-esterification. Calcium fluorescence was measured by excitation of the Fura-2 fluorophore by separate light-emitting diode (LED) light sources producing light in the UV spectra at 340 and 380 nm. Bandpass filters on each LED light restricted excitation light at 340 and 380 nm. Digital modulation generated alternating excitation wavelengths at 250 Hz, and the light was directed into the perfusion chamber. The framing aperture was centered around single cardiomyocytes to ensure that the fluorescence signal was acquired from only the selected cardiomyocyte. Emitted light at 510 nm passed through a dichroic mirror and directed to a photomultiplier tube (PMT) where the photon count was amplified and converted to a digital signal for fluorescence quantification. The output signal for the Fura-2 ratiometric method consisted of 250 ratios per second and recorded with IonWizard software (IonOptix). For contractility measurements, cardiomyocytes were imaged through a 40X objective in an inverted microscope. Cardiomyocyte contractility was measured by video-based CMOS sensor (MyoCam, IonOptix) quantification of changes in SL by fast Fourier transform (FFT) of a region on interest (ROI) selected on a longitudinal section of the cardiomyocyte with a clear striation

pattern. The ROI was kept in the same location on the cardiomyocyte for the duration of the

experiment. Video capture produced SL measurements at 500 Hz. SL signals were recorded with

IonWizard software (IonOptix) concurrent with calcium fluorescence. For analysis, a segment of

steady-state cardiomyocyte contraction was selected for each protocol period: 1) pre-treatment

(PRE), 2) either DMSO (Ctrl), or mavacamten (MAVA) at 1Hz, and 3) for stimulations at 2, 3,

and 4Hz. At least 10 calcium transients were included for each analyzed period.

In vivo **studies (Bowditch).** The rat experiments were done in compliance with protocols

approved by the Institutional Animal Care and Use Committees at Bristol-Myers Squibb

(formerly MyoKardia). Conscious cardiovascular responses to acute myosin-modulation via

mavacamten were studied in telemetered adult male Sprague Dawley rats instrumented with an

implantable radio-telemetry unit providing both systemic arterial (abdominal aorta, AoP) and

left-ventricular pressures (model HD-21; Data Sciences International, Inc.). These animals were

(orally) administered either DMSO (Ctrl, 0 mg/kg, n = 11) or mavacamten (MAVA: 1.0 mg/kg

PO, n = 6). In these experiments, LV hemodynamics were recorded continuously via telemetry in

conscious free-roaming animals both prior to (up 1 hour) and following (up to 24 hours) each

dosing. Heart rate (HR) as well as the peak rates of pressure rise (dP/dtmax) and decline

(dP/dtmin) were derived/measured from LV pressure signals, and are reported for the post-

dosing period.

In vivo **studies (β-adrenergic stimulation).** The pig experiments were done in compliance with

protocols approved by the Institutional Animal Care and Use Committees at Exemplar Genetics.

Four healthy Yucatan male pigs (58.0 ± 9.5 Kg), were chemically-restrained

(Telazol/Xylazine/Isoflurane) and underwent transthoracic echocardiographic examinations.

Left-ventricular end-systolic and end-diastolic dimensions were measured from 2D-guided M-

mode images in the parasternal long- and mid-papillary short-axis views; volumes and left-

ventricular ejection fraction (LVEF) were calculated from internal dimension via the Teichholz

method. End-diastolic volumes were indexed to body-surface area (EDVI = EDV/BSA) as

described before [42] to account for differences in body weight. These measurements were then

repeated after mavacamten administration (0.23 ± 0.03 mg/kg IV, at ~2mL/kg) (DOSE). These

measurements were then repeated after dobutamine administration (+DOB, 5 μg/kg/min) for 5

min to evaluate functional recruitment.

Computational simulations of thick filament regulation by calcium. For simulations of thick

filament regulation by calcium, we used a recently updated version of the MUSICO

computational platform[43-45] but with parameters modified from the previous studies to account

for differences between rodent and human myocardium and porcine myocardium. The full set of

parameters used in the model is given in Table S1. The model incorporated into MUSICO

accounts for all interactions between sarcomere proteins, e.g., myosin with actin, the troponin-

tropomyosin complex with the actin surface, and proteins with ions and small molecules such as

Ca^{2+} in the context of a spatially explicit, three-dimensional (3D) structure of the sarcomere[43-45].

The kinetic schemes describing these interactions include two major parts: (1) calcium binding to

troponin and the regulation of availability of actin sites for myosin binding; (2) kinetics of the

crossbridge cycle including interactions of myosin heads with actin and with the thick filament

backbone. The first process is defined by four states that describe the thin filament-based

regulation that binding of calcium to cTnC to form a Ca-cTnC complex that turn on thin

filament. The second process is defined by a six-state crossbridge cycle that includes five

biochemical states consistent with observed structural states: M.D.Pi, A.M.D.Pi, A.M.D, A.M,

M.T and an additional sixth state, the so-called "parked state" (PS), represents the interaction of

myosin heads with the thick filament backbone where they are unable to interact with actin. The

transition rate from the PS to M.D.Pi is assumed to be strongly dependent on $[Ca^{2+}]$, i.e.

$k_{+PS}(Ca) = k_{PS}^o + [(k_{PS}^{max} - k_{PS}^o)[Ca]^b]/([Ca]_{50}{}^b + [Ca]^b)$ where k_{PS}^o is baseline rate from

parked state, k_{PS}^{max} is the rate at high calcium, e.g. at pCa=4.5, b is Hill coefficient of the rate

sigmoidal rise and $[Ca]_{50}$ is calcium concentration when k_{+PS} is equal to $k_{PS}^{max}/2$ [43-45]. In

performing the simulations, the model incorporated known structural (lattice spacing, sarcomere

length) and functional parameters (force pCa curve). The simulations then varied the adjustable

parameters (k_{-PS}, k_{PS}^o , k_{PS}^{max}, Hill coefficient, b, and reverse rate, k_{-PS}) to provide good fits to

the data in control experiments. With mavacamten, only k_{-PS} needed to be adjusted to

quantitatively match the data. Since this is not a parametric model, there is no minimization

process, so normal goodness of fit statistics are not meaningful. See Supplemental Information

(SI) for further details.

Statistics. Statistical analyses were performed using GraphPad Prism 9 (Graphpad Software).

The relative changes versus pCa curves in figure 2 were fit to a four-parameter modified Hill

equation (minimum response + (maximum response - minimum response)/(1+10[h]. (pCa50-pCa)))

[46], where pCa_{50} is the calcium concentration yielding a response halfway between the minimum

and maximum values reported in the article. Mann-Whitney unpaired nonparametric t-test were

performed on the data in figure 2 between the thin filament inhibitor group and mavacamten

groups. Wilcoxon matched-pairs nonparametric t-test were performed on the data in figure 3

within control and mavacamten groups. Mann-Whitney unpaired nonparametric t-test were

performed on the data in figure 3 between control and mavacamten groups. The effects of

mavacamten in isolated myocytes and in vivo were evaluated via a one-factor analysis of

variance (ANOVA) with a repeated-measures design and set predetermined comparisons (e.g.,

DOSE vs. PRE and DOSE vs. +DOB). In addition, multiple linear regression analyses were used

to compare the HR-dependent effects in dP/dt_{max} and dP/dt_{min}. The significance level was set at

0.05 a priori. Symbols on figures: ns: not significant, *: $p<0.05$, **: $p<0.01$, ***: $p<0.001$,****:

$p<0.0001$. N and n indicate biological and technical repeats respectively

Results

Ca^{2+} destabilizes the SRX state(s) of myosin in thin-filament-free reconstituted synthetic

myosin thick filaments.

We have recently reported that Ca^{2+} binding to myosin filaments in reconstituted cardiac

synthetic thick filaments (STF) can directly destabilize myosin SRX state(s) towards the DRX

state(s)[23]. Here we studied Ca^{2+}-mediated destabilization in the presence of mavacamten, whose

primary mechanism of biochemical inhibition is by arresting myosin heads in the SRX state (Fig.

1a). The basal myosin SRX population in the STF system at pCa 10 is 10 ± 5 % (n=3), which

progressively increases with increasing concentration of mavacamten. The IC_{50} of this transition

is 1.2 ± 0.5 µM (Fig. 1a). Consistent with our earlier reports of a Ca^{2+}-mediated destabilization

mechanism[23], the myosin SRX population in the control sub-group progressively decreases with

an increase in Ca^{2+} with a pCa_{50} of 5.5 (5.3 to 5.6 for 95% CI). Interestingly, this mechanism of

Ca^{2+}-mediated destabilization of the SRX population holds at all mavacamten concentrations

tested (0.5, 1, and 2 µM) (Fig. 1b). These results suggest that while mavacamten puts more

myosin into the SRX state(s) (Fig. 1b, left, upwards arrow), this effect was blunted by increasing

Ca^{2+} concentrations (Fig. 1b, right, upwards arrow). The pCa_{50} of the Ca^{2+}-mediated

destabilization in the presence of various concentrations of mavacamten lies in the range of 5.2

to 5.6 (4.9 to 5.7 for 95% CI) (n=3), suggesting that the myosin heads held in the SRX

configuration by mavacamten can be similarly destabilized as in the absence of it. Taken together, these data signify that myosin heads are not permanently locked even at saturating doses of mavacamten (2 μM) but instead are recruitable as needed.

Consistent with this mechanism, the basal ATPase activity of myosin in STF progressively increases with an increase in Ca^{2+} with a pCa_{50} of 6.32 (6.28 to 6.37 for 95% CI) (n=8) in the control sub-group (Fig. 1c). This mechanism of Ca^{2+}-mediated activation of myosin ATPase also holds at 0.5, 1, and 2 μM mavacamten concentrations with pCa_{50} values of 6.31 (6.28 to 6.37 for 95% CI), 6.23 (6.18 to 6.28 for 95% CI), and 6.22 (6.14 to 6.30 for 95% CI) (Fig. 1c), respectively, suggesting that myosin heads inhibited by mavacamten can be re-activated with increasing Ca^{2+} concentrations. This premise is further tested in a more complex bovine cardiac myofibril system, encompassing both the thick and the thin filament components of the sarcomere. Increasing Ca^{2+} is expected to increase the myofibrillar ATPase via activating the thin-filament troponin-tropomyosin regulatory system (Fig. 1d, black; Control group; pCa_{50} of 6.36 and 6.32 to 6.40 for 95% CI). The Ca^{2+}-dependent direct thick filament activation mechanism as demonstrated in the STF system above can be isolated from myofibrillar system by using a 100 μM thin filament (TF) inhibitor (MYK-7660)[23] to inhibit acto-myosin interactions, thus abolish active contraction. Our data shows that Ca^{2+}-dependent thick filament activation, although blunted, was retained in the absence of active force (Fig 1d, red; TF inhibitor; pCa_{50} of 5.78 and 5.73 to 5.83 for 95% CI), suggesting that a part of the Ca^{2+}-mediated activation of the sarcomere lies in the thick filament, consistent with our previous report[23]. Increasing mavacamten concentrations in the presence of 100 μM MYK-7660, in general, progressively blunted the ATPase activity due to myosin inhibition, but, very interestingly, was partially rescued by increasing Ca^{2+} (Fig. 1d, traces in blue). The pCa_{50} values of ATPase in myofibrils with 100 μM MYK-7660 treated with 0.5, 1, and 2 μM concentrations of mavacamten lies in the range of 5.76 to 5.93 (5.62 to 6.04 for 95% CI) (n=3). We note that it is also possible that at this high concentration of the MYK-7660 and mavacamten, the sarcomere could be inhibited by other non-specific mechanisms, which would only lead to an underestimation of the degree of rescue by Ca^{2+}, further bolstering our hypothesis.

Figure 1. The biochemical SRX states of myosin stabilized by mavacamten are recruitable by Ca^{2+}. a, Absolute percentage (%) of myosin SRX population in synthetic thick filaments (STF) reconstituted from bovine cardiac full-length myosin. The IC$_{50}$ for this transition is 1.2 ± 0.5 μM. **b**, Absolute % myosin SRX population in STF in the absence (Control (2% DMSO), black) and the presence of increasing concentrations of mavacamten (0.5, 1 and 2 μM; increasing darker shades of blue) at different Ca^{2+} concentrations. Ca^{2+} destabilizes the myosin SRX state (s), otherwise stabilized by mavacamten. While mavacamten puts more myosins in the SRX state(s) at all concentrations (left, upwards arrow), this effect was blunted by increasing Ca^{2+} concentrations (right, upwards arrow). **c**, Absolute basal myosin ATPase in STF in the absence (Control (2% DMSO), black) and the presence of 0.5, 1, and 2 μM mavacamten concentrations (profiles in blue) at different Ca^{2+} concentrations. **d**, Absolute myosin ATPase in bovine cardiac myofibrils in the absence (Control (2% DMSO), black), and in the presence of 100 μM of thin filament inhibitor without (TF inhibitor, red) and with 0.5, 1, and 2 μM mavacamten concentrations (profiles in blue) at different Ca^{2+} concentrations.

Ca^{2+} mediated myosin filament structural changes are preserved in the presence of mavacamten.

We have previously shown Ca^{2+} can directly move myosin heads from ordered *off* states to disordered *on* states in the absence of active force[23]. Here we investigated how myosin thick filaments react to increasing Ca^{2+} concentration in the presence of mavacamten (blue curves in Fig. 2) and compared it to previously reported data[23] in the absence of mavacamten with (TF Inhibitor, red curve in Fig 2) and without TF inhibitor (Control, black curve in Fig 2). The

equatorial intensity ratio ($I_{1,1}/I_{1,0}$) is an indicator of the proximity of myosin heads to actin[25]. The

change in $I_{1,1}/I_{1,0}$ ($\Delta I_{1,1}/I_{1,0}$) vs. pCa curve shows a sigmoidal shape (Fig. 2b) with a pCa_{50} of 5.7

for the Mavacamten group (N =2 and n = 7), the same as in the TF inhibitor and Control groups

as previously reported[23]. The increase in $\Delta I_{1,1}/I_{1,0}$ with increasing Ca^{2+} concentration indicates

that myosin heads move radially away from the thick filament backbone even in the presence of

mavacamten. Relaxed myosin heads are quasi-helically ordered on the surface of the thick

filament producing the myosin-based layer line reflections[25]. The first-order myosin-based layer

line (MLL1) arising from the helically ordering of myosin heads is disrupted when myosin heads

are turned *on* to participate in contraction, leading to a decrease in the intensities of MLL1

(I_{MLL1}). Similarly, the intensity of the third-order myosin-based meridional reflection (I_{M3})

decreases when myosin heads become disordered[25, 47]. I_{M3} (Fig. 2c) and I_{MLL1} (Fig. 2d) decays as

a function of increasing Ca^{2+} concentration yields a pCa_{50} of 6.14 (5.93 to 6.38 for 95% CI) and

6.01 (5.84 to 6.20 for 95% CI) respectively in the presence of mavacamten. The spacing of the

sixth-order meridional reflection (S_{M6}) reports the periodicity of the thick filament backbone

with increases in S_{M6} indicating extension of the thick filament backbone[41]. The degree of thick

filament lengthening has been shown to correlate with the degree of myosin heads being in the

on states[48] and increases with increasing Ca^{2} concentration with a pCa_{50} of 6.35 (>5.97 for 95%

CI) for the mavacamten groups (Fig 2e).

These Ca^{2+}-mediated thick filament structural transitions in the presence of mavacamten,

(blue curves in Fig 2) have a pCa_{50} in the range of 5.9-6.4, essentially identical to the previously

reported range in the absence of mavacamten (Control and TF Inhibitor group in Fig 2).

Consistent with the biochemical studies in figure 1, the amplitudes of the Ca^{2+}-induced structural

changes are the greatest in the Control group and the least in the Mavacamten group, while the

TF inhibitor group is in between, indicating that even though mavacamten shifts more myosin

heads to *off* states, these *off* state myosin heads are still sensitive to Ca^{2+}.

Figure 2. Thick filament structure changes in the presence of Ca^{2+}. a, Representative X-ray diffraction patterns from permeabilized porcine myocardium under relaxing conditions (pCa 8, top panel) and activating conditions (pCa 4.5) with (bottom right) and without (bottom left) mavacamten in the absence of active force (treated with 100 μM MYK-7660). **b,** Change of equatorial intensity ratio ($\Delta I_{1,1}/I_{1,0}$) at different Ca^{2+} concentrations in the Control sub-group (black), thin filament (TF) inhibitor group (red), and the Mavacamten group (blue). The intensity of the third-order of myosin-based meridional reflection (**c**) and the intensity of the first-order myosin-based layer line (**d**) and in different concentrations of Ca^{2+} in the Control (black), TF inhibitor (red), and the Mavacamten (blue) groups. **e,** The spacing of the sixth-order myosin-based meridional reflection (S_{M6}) in different Ca^{2+} concentrations in the Control (black), TF inhibitor (red), and the Mavacamten (blue) groups. The Ca^{2+} induced structural changes in the thick filament were blunted but not eliminated by the presence of mavacamten. * $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$ between TF inhibitor and mavacamten groups. The results are given as mean ± SEM.

Mavacamten preserves positive force-frequency response in *vitro* and in *vivo*

As expected, and consistent with the Bowditch effect, increasing stimulation frequencies (from 1 to 4Hz) increases shortening fraction in isolated, membrane-intact, rat ventricular myocytes (Fig 3a & 3b, Table S2, n=12); this Treppe effect was driven by increases in Ca^{2+} transient amplitudes (Fig 3c, Table S2). Mavacamten (at 0.1 μM) preserved this Ca^{2+}-mediated functional recruitment, despite markedly inhibiting shortening (3.9 ± 0.5 ($p = 0.005$) vs. 6.1 ± 0.6% in control; ($p = 0.0002$)). Indeed, comparable gains in function with pacing (i.e., linear slopes) were noted between control and mavacamten treated cells (+12 ± 3%/Hz vs. +13 ±

5%/Hz in VEH, n.s.). Higher stimulation frequencies also elevated diastolic cytosolic Ca^{2+}

concentrations and resulted in progressive reductions in diastolic (resting) cell length (Fig 3b,

Table S2). Mavacamten preserved diastolic cell length (Fig 3a), blunting the diastolic contracture

(loss of diastolic length) with pacing (i.e., linear slope: -0.26 ± 0.04%/Hz (p = 0.00003) vs. -0.50

± 0.07%/Hz in Control, (p = 0.00004); Fig 3b) despite comparable increases in diastolic Ca^{2+}-

levels. This effect is likely mediated via the ability of mavacamten to facilitate mechanical re-

lengthening (RT75: 86 ± 3%, or 0.15 ± 0.01 (p = 0.00002) vs. 0.17 ± 0.01s pre-dose (p =

0.00007)).

These in *vitro* observations were also noted *in vivo* in conscious, telemetered rats, where

circadian changes in heart rate (over 24 h) are leveraged to establish the chronotropy-dependence

of both dP/dt$_{max}$ and dP/dt$_{min}$, inotropic and lusitropic indices derived from the left-ventricular

pressure (Fig 3d). Under control conditions (N = 11), both a negative (strong) heart rate (HR)-

dependence of dP/dt$_{min}$ (-18.2 ± 0.5 mm Hg/s/bpm in Ctrl, R^2=0.70, p < 0.0001) and a positive

HR-dependence of dP/dt$_{max}$ (20.0 ± 1.0 mm Hg/s/bpm in Ctrl, R^2=0.41, p < 0.0001) are noted

(black curve in Fig 3e & 3f). Mavacamten administration (1 mg/kg PO, N = 6) enhanced these

relationships as steeper dP/dt$_{min}$ (-21.1 ± 0.9 mm Hg/s/bpm) and dP/dt$_{max}$ (24.6 ± 1.8 mm

Hg/s/bpm) (p = 0.0077) vs. HR slopes were noted under treatment. Taken together, these data

demonstrates that mavacamten permits and/or enhances systolic and diastolic functional

recruitment as HR increases, consistent with the Bowditch effect.

446
Figure 3. In *vitro* and in *vivo* Bowditch effect in the presence of mavacamten. In isolated rat
ventricular myocytes, increases in stimulation frequency (from to 1 to 4 Hz) increasing
shortening fraction (**a**) and decreasing resting length (**b**) by increase peak $[Ca^{2+}]$ during systole
(**c**); In *vivo*, circadian changes in heart rate (HR) are accompanied by changes in the left-
ventricular pressure derived dP/dt_{max} (**e**, direct) and dP/dt_{min} (**f**, inverse), reflecting HR-dependent
function and mavacamten (1 mg/kg PO) preserved/enhanced these relationships. ******: $p < 0.01$,
***: $p < 0.001$, ********: $p < 0.0001$. The bar graphs are given as mean values. ******: $p < 0.01$, *******: p
< 0.001, ********: $p < 0.0001$, # $p < 0.05$ vs. Ctrl. Solid lines in panel **e** and **f** are simple linear
regression fits and the dashed lines are 95% confidence bands of the best-fit line.
456

Stretch-mediated myosin filament structural changes are preserved in the presence of mavacamten.

Systolic contractility of the heart is also regulated by the degree of diastolic filling. At the cellular level, passive stretch of porcine myocardium can recruit reserve myosin heads from *off* states to *on* states promoting cross-bridge formation during activation[22]. While it has been previously shown that mavacamten can stabilize the *off-state* of the thick filament[13, 22], it is unknown whether this stretch-mediated thick filament recruitment is preserved in the presence of mavacamten. Here we demonstrated that both $I_{1,1}/I_{1,0}$ and the radius of the center of mass of the cross-bridges (R_m), a direct measurement of the distance of helically ordered myosin heads from the thick filament backbone[49], increased significantly in both the control (N=2 and n=7; $p < 0.05$) and the mavacamten (N=2 and n=8; $p < 0.05$) group when muscle is stretched form SL of 2.1 µm to 2.3 µm (Fig 4 a & b, Table S3). Correspondingly, both I_{MLL1} and I_{M3} decreased significantly in both the control (N=2 and n=7; $p < 0.05$) and the mavacamten (N=2 and n=8; $p < 0.05$) groups when muscle is stretched from SL of 2.1 µm to 2.3 µm (Fig 4 c&d, Table S3). The $I_{1,1}/I_{1,0}$ and R_m data indicate that myosin heads move radially away from the thick filament backbone concomitantly with a loss of helical ordering, indicated by I_{MLL1} and I_{M3}, upon lengthening in the presence and absence of mavacamten. Remarkably, there are no significant differences ($p > 0.05$, Fig 4, Table S3) in all these changes induced by SL lengthening between the control and mavacamten group strongly indicating that length-mediated structural changes in the myosin filaments are preserved in the presence of mavacamten.

Figure 4. Length-dependent thick filament structural changes in the presence and absence of mavacamten. $I_{1,1}/I_{1,0}$ **(a)** and the radius of the average mass of myosin heads (R_m) **(b)** increases while the intensity of the first-order myosin-based layer line (I_{MLL1}) **(c)** and the intensity of the third-order myosin-based meridional reflection (I_{M3}) **(d)** decreases at longer sarcomere length indicate structural transitions of the thick filament towards the *on* states upon sarcomere lengthening in the presence (blue) and absence (grey) of mavacamten. *: $p < 0.05$, **: $p < 0.01$. The results are given as mean ± SEM.

β-adrenergic recruitment is preserved in the presence of mavacamten in *vivo*.

In addition, the effects of β-adrenergic stimulation in the presence of mavacamten were studied *in vivo*. As expected at the supra-therapeutic (i.e., producing functional depression greatly exceeding those observed in the clinic ($350 - 700$ng/mL[14])) dose (>1000ng/mL) used, mavacamten results in marked functional inhibition in healthy pigs (N = 4), decreasing left ventricular ejection fraction (LVEF) by $32 \pm 6\%$ (from 72 ± 5 to $40 \pm 5\%$, $p = 0.0437$, Fig 5a), resulting in increased end-systolic and end-diastolic ventricular dimensions. Notably, hyperdynamic contraction and decreased chamber size contribute to left-ventricular outflow tract obstruction in HCM. Mavacamten triggers significant increases in end-diastolic volume index (EDVI) from 41 ± 6 mL/m^2 to 59 ± 6 mL/m^2 ($p = 0.0263$, Fig 5b), which has been shown[26] to reflect increased ventricular distensibility, as suggested by a reduction in the number of residual

diastolic cross-bridges recruited by stretch. The functional effects of mavacamten are recruitable

with acute β-AR stimulation. Dobutamine restores LVEF towards baseline values (65 ± 4%, vs.

DOSE, $p = 0.0437$, Fig 5a), decreasing end-systolic volumes (40 ± 3 to 24 ± 3 mL, $p = 0.0302$,

Fig 5c) and increasing stroke-volume (27 ± 5 to 45 ± 8 mL, $p = 0.0471$, Fig 5d), suggesting an

increase in end-systolic elastance and forward flow as negligible effects in heart rate are

observed. β-AR stimulation fails to decrease diastolic volumes (59 ± 6 % to 59 ± 6%, n.s.),

suggesting that the diastolic effects of mavacamten remain unaltered. Taken together, these

observations indicate that despite mavacamten-mediated thick-filament stabilization, a

population of myosin-heads remains available to be recruited via β-adrenergic receptor

stimulation.

507
Figure 5. **In *vivo* cardiac effects of mavacamten, and overlapping dobutamine, in healthy**
pigs. **a,** mavacamten (DOSE) at supra-therapeutic doses results in a marked decrease in left
ventricular ejection fraction (LVEF), which are acutely recruitable by β-adrenergic receptor
stimulation with dobutamine (+DOB; 5 µg/kg/min for 5 min). **b,** end-diastolic volumes index
(EDVI) is increased by mavacamten and it is insensitive to β-AR stimulation. **c,** mavacamten
increases LV end-systolic volume (LVESV) but -AR stimulation is able to reverse it. **d**,
mavacamten does not significantly affect LV stroke volume (LVSV) but β-AR stimulation
significantly increases LVSV. ns: not significant, *: $p < 0.05$, **: $p < 0.01$. The results are given

as mean ± SEM.

Simulations of thick filament regulation by calcium.

We used the computational simulation program MUSICO (see methods) to explore the

effect of Ca^{2+} concentration on transitions in the population of myosin heads from sequestered

off states to *on* states[45, 50]. Simulations considered these transitions in the context of biochemical

transitions from a "parked state" (PS) to detached Myosin-ADP-phosphate (M.D.Pi) state and

then allowing myosin to bind to actin to form the attached Actin-Myosin-ADP-phosphate

(A.M.D.Pi) state. (Fig 6). The "parked state" is functionally defined as any heads unable to bind

to actin and would include any heads in the biochemically defined SRX state or the structurally

defined *off* states regardless of whether these are the same or different populations. From the

fraction of the myosin heads in the PS estimated from the X-ray diffraction data, assumed here to

be proportional to the square root of the number of ordered heads, we establish the Ca^{2+}-

dependent state transition rate from the PS using the data from experiments involving the thin

filament inhibitor MYK-7660 (TF inhibitor: Fig 6, red symbols). We achieved very good

matches to the data (Fig 6; red line) by setting the rate of Ca^{2+} binding to very small values, i.e.

mimicking inhibition of opening thin filament regulatory units, and setting parameters for the

Ca^{2+}-dependent state transition from the PS to M.D.Pi, $k_{+PS}(Ca^{2+})$, as: $k_{PS}^{o} = 27$ s^{-1}, $k_{PS}^{max} = 220$

s^{-1} and Hill coefficient, $b = 1.25$. The reverse rate constant from M.D.Pi to PS is taken to be k_{-PS}

$= 200$ s^{-1}. Using parameters for the cross-bridge cycle slightly modified from simulations for

human myocardium [44] and adjusted for temperature, species, and muscle preparation (i.e.

demembranated vs. intact muscles, following [50]) the predicted changes in the fraction of the

myosin heads in the PS as a function of Ca^{2+} concentration (pCa) (Fig 6; black line) match well

the observed values for *off* state heads estimated from the X-ray diffraction experiments (Fig 6;

black symbols). We also explored the effect of 2 μM mavacamten on permeabilized porcine

muscle in the absence of actin-bound cross-bridges (achieved with the TF inhibitor MYK-7660).

The predicted fraction of myosin heads in the PS (Fig 6, blue line) matches the observations (Fig

6, blue symbols), by changing only one rate constant k_{-PS} from 200 to 263 s^{-1}. This increase in

k_{-PS} reflects the activity of mavacamten in increasing the population of myosin heads in the PS

at all Ca^{2+} concentrations. Notably, even at low Ca^{2+} concentration (pCa 8), the population of PS

heads is increased (3.2%) with mavacamten.

547
Figure 6. Simulation of the Ca²⁺-dependent parked state. a, Parked state (PS) myosin heads
that are unavailable to form cross-bridges can be recruited by calcium to detached state(s) that
are ready to participate in contraction. **b,** The population of PS heads decreases as the calcium
concentration increases in both control muscle where active force is present (Exp.: experimental
data; Sim.: simulation data) and thin filament inhibitor (TF inhibitor) group where active force is
totally inhibited. In muscle where 2μM mavacamten is present, the population of the PS myosin
heads decreases in the same manner as the control and TF inhibitor group, albeit with a reduced
magnitude.
556

Discussion

Here, we show that cardiac myosins stabilized in these *off* state(s) by mavacamten are
recruitable by 1) Ca²⁺, 2) increased chronotropy (heart rate), 3) stretch, and 4) β-adrenergic (β-
AR) stimulation, all known physiological inotropic effectors. At the molecular level, we show
that Ca²⁺ increases myosin ATPase activity by shifting mavacamten-stabilized myosin heads
from the inactive super-relaxed (SRX) state to the active disordered relaxed (DRX) state. At the
myofilament level, both Ca²⁺ and passive lengthening can shift mavacamten-ordered *off* myosin
heads from positions close to the thick filament backbone to disordered *on* states closer to the
thin filaments. In isolated membrane intact rat cardiomyocytes, increased stimulation rates
enhanced shortening fraction in mavacamten-treated cells. This observation was confirmed *in*
vivo in telemetered rats, where left-ventricular dP/dt$_{max,}$ an index of inotropy, increased with
heart rate in mavacemten treated animals. Finally, we show that β-AR stimulation *in vivo*
increases left-ventricular function and stroke volume in the setting of mavacamten. In

conclusion, these data demonstrate that the mavacamten-promoted *off* states of myosin in the

thick filament are activable.

At the molecular level, excessive cross-bridge cycling due to mutations in sarcomeric

proteins have been widely recognized as the primary contributor to hypercontractility in HCM[9,

10]. Apart from the mutation-induced primary effect on myofilament function, other

pathophysiological insults such as high ADP:ATP ratio or increased Ca^{2+} may add to the

hypercontractile phenotype in HCM. Consistent with this hypothesis we have shown, in a recent

report, that independent of the thin filament-mediated activation, Ca^{2+} can directly destabilize the

parked *off* states of myosin[23], thereby increasing the probability of excess actin-myosin

interactions that could drive hypercontractility in HCM pathology, but also suggesting a

mechanistic link between thick-filament recruitment and inotropism.

Mavacamten is a small-molecule, FDA-approved, therapeutic for HCM that directly

addresses this causal biology by directly targeting and inhibiting cardiac myosin[12] thus correcting

the hypercontractility seen in HCM which can potentially prevent the HCM associated symptoms

and cardiac remodeling. It does so by sequestering myosin heads from the functionally more

active DRX state to a less active SRX state, presumed to be structurally similar to the closed

IHM-like state[13, 20, 24]. Myosin heads in this state are thought to lie close to the thick filament

backbones where they are quasi-helically ordered (the structurally defined *off* state) and away

from the thin filaments so that they are unavailable to interact with actin[10, 51]. Based on these two

major propositions, a bigger question emerges. Does mavacamten permanently park these

myosin heads in the *off* state(s)? Can these heads be recruited as and when required to boost the

output of the heart? The answer(s) to these questions may provide a mechanistic explanation for

the therapeutic benefits of mavacamten beyond mere obstruction relief, especially in the context

of increased cardiac demand in patients during exercise.

This report shows that, at the molecular level, mavacamten shifts the SRX/DRX

equilibrium towards the SRX state(s) and mitigates the deviation caused by the liberation of SRX

heads to the DRX state(s) seen in HCM as indicated by the STF SRX assays. Our computational

simulations predict the population of the myosin heads in the parked state (PS) in the presence of

mavacamten, and a reasonable fit is achieved by changing only one rate constant k_{-PS} from 200

to 263 s^{-1}. This result indicates that mavacamten promotes the transition of more myosin heads in

the detached state into the sequestered PS. However, our simulation studies show that binding of

mavacamten to myosin heads does not affect k_{+PS}, suggesting that Ca^{2+} dependent recruitment

of myosin heads from the PS is preserved in the presence of mavacamten. Additionally, the

magnitude of Ca^{2+}-mediated destabilization of SRX (Fig 1b) increases from ~12% in the absence

of mavacamten (from 15% to 3%) to ~21% in the presence of 2 μM mavacamten (from 60% to

38%), suggesting that the additional myosin heads that are parked in the SRX state(s) by

mavacamten are destabilized by Ca^{2+}, highlighting 1) the reversibility of such states, but also, 2)

the role of myosin-recruitment during contraction. These results also serve to support the *in vitro*

(Fig 3 a & b) and *in vivo* observations (Figure 3 d, e and f) that mavacamten preserves and/or

enhances the Bowditch effect, as stimulation-frequency and heart-rate dependent recruitment of

function was evident in the presence of mavacamten. It should also be noted that Matsubara and

colleagues showed that the Bowditch, or staircase phenomenon in cardiac muscle is mediated by

a frequency-dependent increase in the number of myosin projections transferred to the vicinity of

the thin filaments during diastole[52] and during systolic contraction[53]. Our studies provide a

molecular mechanism for these findings, by associating the increases in myosin-mass transfer

during Treppe with direct Ca^{2+}-dependent regulation of *off* to *on* transitions of myosin on the

thick filament.

617   In addition, the sarcomere-level observation that both Ca^{2+} and passive lengthening can

shift myosin heads from ordered *off* states close to the thick filament backbone to disordered *on*

states closer to the thin filaments in the presence of saturating mavacamten (Fig 4). Previous

mechanical studies showed that mavacamten blunts but does not eliminate the systolic gains of

LDA[22, 27, 28] supporting the notion that the myosin heads sequestered in the *off* SRX state(s)

induced by mavacamten are available for length dependent recruitment. All these lines of

evidence are supported by the observations from *in vivo* studies showing that mavacamten blunts

but does not eliminate β-adrenergic stimulation effects on the heart.

625   Our data not only demonstrate that myosins in mavacamten induced *off* states in the thick

filament are recruitable, thus leading to the preservation of the cardiac reserve, but also indicate

that mavacamten enhances cardiac output in HCM patients beyond mere LV outflow tract

obstruction removal in oHCM patients. This hypothesis is partially validated by the phase II

MAVERICK-HCM trial in patients without obstruction in which a high-risk subgroup with

elevated myocardial injury bio-markers (cardiac troponin I >99[th] percentile) or elevated diastolic

filling pressures (average E/e'>14 on echocardiogram), one-third of mavacamten-treated patients

(N =7 compared to zero in the placebo group, p = 0.03) met the composite functional endpoint—
defined as achieving: 1) an improvement of at least 1.5 ml/kg/min in pVO_2 with a reduction of
≥NYHA functional class; or 2) an improvement of 3.0 ml/kg/min or more in pVO_2 with no
worsening in NYHA functional class[17]. It is worth noting that the increase in pVO_2 was
insufficiently powered in the trial probably due to it was designed as a dose-finding study. We
anticipate this question will be resolved by the ongoing phase III ODYSSEY-HCM trial.

Our current studies are limited to the short term recruitability of mavacamten stabilized
myosin in healthy systems. We did not investigate how things might be in diseased myocardium.
In clinical trials where HCM patients were administered mavacamten showed improved exercise
capability beyond pVO_2 strongly suggesting that the myosin stabilized by mavacamten is
recruitable to accommodate increased physiological demands during exercise[54]. While we
showed β-adrenergic stimulation was able to reverse the effects of mavacamten, no specific
targets of β-adrenergic stimulation were identified in the current study. It has been shown that β-
adrenergic stimulation has multiple effects: 1) it prolongs L-type calcium channel opening,
effectively raising the Ca^{2+} concentration[55]; 2) it phosphorylates multiple sarcomeric proteins
including myosin-binding protein C [56], troponin I and C[57] and titin[58] all of which could modulate
heart contractility. We also did not investigate the recruitability of mavacamten-stabilized
myosin in the presence of other more commonly used drugs for HCM such as β blockers and
calcium blockers[54]. It has been shown that pVO_2 change was blunted in HCM patients on β
blockers compared with patients that were not in the EXPLORER-HCM trial[54]. We have shown
that β-AR blockade does not alter the diastolic and systolic effects of mavacamten[59], suggesting
that this clinical dichotomy could be attributed to the effects of β blockers on cardiac recruitment
such as blunted heart rate recruitment[60] rather than to mavacamten per se. Additionally, our
experiment did not allow us to investigate the side effects of either mavacamten or how it might
affect β-adrenergic stimulation long term. All of these questions, while beyond the scope of
current study, should be examined experimentally or clinically in the future.

The ultimate function of the heart is to maintain adequate blood supplies under various
conditions especially during increased demand. Here we provide direct evidence, ranging from in
silico, to in *vitro* and in *vivo*, that Ca^{2+}-mediated activation, chronotropic inotropism, sarcomere
stretch-mediated activation, and β-adrenergic receptor stimulation are all preserved with
mavacamten treatment, resulting in a likely maintenance (or in disease enhancement) of cardiac

reserve. These results not only provide a mechanistic explanation for the clinically observed increase in peak oxygen uptake (pVO_2) with exercise in patients with HCM receiving mavacamten, but also lay the groundwork for a potential methodology to investigate the sarcomere basis of chronotropic incompetence in disease states motivate new therapeutic interventions.

Acknowledgments

We thank Robert S. McDowell of Bristol Myers Squibb (formerly MyoKardia) for providing helpful comments to the manuscript. This research used resources of the Advanced Photon Source, a U.S. Department of Energy (DOE) Office of Science User Facility operated for the DOE Office of Science by Argonne National Laboratory under Contract No. DE-AC02-06CH11357. This project is supported by grant P30 GM138395 from the National Institute of General Medical Sciences of the National Institutes of Health and the Ministry of Education, Science and Technological Development of the Republic of Serbia through Contracts No. 451-03-68/2022-14/200378. The content is solely the authors' responsibility and does not necessarily reflect the official views of the National Institute of General Medical Sciences or the National Institutes of Health.

Author contributions

W.M, C.dR, S.N designed the experiments; W.M, H.G, C.dR, C.Z, R.S, S.G, S.N performed the experiments; W.M, L.Q, C.dD, C.Z. R.S, S.G, S.N. analyzed the data; M.P, S.M.M performed the simulations and interpreted the simulation results; W.M, S.M.M, C.dR, S.N, T.C.I. wrote the manuscript. All authors approved the final version of the manuscript.

Competing interests

C.Z and R.S are employees and own shares in Bristol Myers Squibb (formerly MyoKardia). S.N and S.G are former employees of Bristol Myers Squibb. C.dR. has a consulting relationship with Bristol Myers Squibb. S.M.M and M.P are partners in FilamenTech, Inc. and they provided the simulations at no cost. T.C.I. consults for Edgewise Therapeutics, but this activity has no relation to the current work.

Data Availability

694   The dataset generated or analyzed during this study are included in this article. The raw

695 data are available from the corresponding authors (Suman Nag: sumannag@gmail.com; Carlos

696 L. del Rio: cdelrio.heart@gmail.com; Weikang Ma: wma6@iit.edu) on reasonable request.

698 **Code Availability**

699   The X-ray dataset were analyzed using data reduction programs belonging to the open-

700 source MuscleX software package developed at BioCAT. Source codes are deposited at Github

701 (https://github.com/biocatiit/musclex).

References

1. Catterall WA. Voltage-gated calcium channels. *Cold Spring Harb Perspect Biol*. 2011;3:a003947

2. Usman A, Gandhi J, Gupta G. Physiology, bowditch effect. *Statpearls*. Treasure Island (FL); 2022.

3. Palomeque J, Vila Petroff MG, Mattiazzi A. Pacing staircase phenomenon in the heart: From bodwitch to the xxi century. *Heart Lung Circ*. 2004;13:410-420

4. de Tombe PP, Mateja RD, Tachampa K, Ait Mou Y, Farman GP, Irving TC. Myofilament length dependent activation. *Journal of molecular and cellular cardiology*. 2010;48:851-858

5. Xiang Y, Kobilka BK. Myocyte adrenoceptor signaling pathways. *Science*. 2003;300:1530-1532

6. Ahmad T, Miller PE, McCullough M, Desai NR, Riello R, Psotka M, Bohm M, Allen LA, Teerlink JR, Rosano GMC, Lindenfeld J. Why has positive inotropy failed in chronic heart failure? Lessons from prior inotrope trials. *Eur J Heart Fail*. 2019;21:1064-1078

7. Sarma S, Stoller D, Hendrix J, Howden E, Lawley J, Livingston S, Adams-Huet B, Holmes C, Goldstein DS, Levine BD. Mechanisms of chronotropic incompetence in heart failure with preserved ejection fraction. *Circ Heart Fail*. 2020;13:e006331

8. Marian AJ, Braunwald E. Hypertrophic cardiomyopathy: Genetics, pathogenesis, clinical manifestations, diagnosis, and therapy. *Circulation research*. 2017;121:749-770

9. Spudich JA. Three perspectives on the molecular basis of hypercontractility caused by hypertrophic cardiomyopathy mutations. *Pflugers Archiv : European journal of physiology*. 2019;471:701-717

10. Nag S, Trivedi DV. To lie or not to lie: Super-relaxing with myosins. *Elife*. 2021;10

11. Mazzarotto F, Olivotto I, Boschi B, Girolami F, Poggesi C, Barton PJR, Walsh R. Contemporary insights into the genetics of hypertrophic cardiomyopathy: Toward a new era in clinical testing? *J Am Heart Assoc*. 2020;9:e015473

12. Green EM, Wakimoto H, Anderson RL, Evanchik MJ, Gorham JM, Harrison BC, Henze M, Kawas R, Oslob JD, Rodriguez HM, Song Y, Wan W, Leinwand LA, Spudich JA, McDowell RS, Seidman JG, Seidman CE. A small-molecule inhibitor of sarcomere contractility suppresses hypertrophic cardiomyopathy in mice. *Science*. 2016; 351:617-621

13. Anderson RL, Trivedi DV, Sarkar SS, Henze M, Ma W, Gong H, Rogers CS, Gorham JM, Wong FL, Morck MM, Seidman JG, Ruppel KM, Irving TC, Cooke R, Green EM, Spudich JA. Deciphering the super relaxed state of human beta-cardiac myosin and the mode of action of mavacamten from myosin molecules to muscle fibers. *Proceedings of the National Academy of Sciences of the United States of America*. 2018;115:E8143-E8152

14. Olivotto I, Oreziak A, Barriales-Villa R, Abraham TP, Masri A, Garcia-Pavia P, Saberi S, Lakdawala NK, Wheeler MT, Owens A, Kubanek M, Wojakowski W, Jensen MK, Gimeno-Blanes J, Afshar K, Myers J, Hegde SM, Solomon SD, Sehnert AJ, Zhang D, Li W, Bhattacharya M, Edelberg JM, Waldman CB, Lester SJ, Wang A, Ho CY, Jacoby D, investigators E-Hs. Mavacamten for treatment of symptomatic obstructive hypertrophic cardiomyopathy (explorer-hcm): A randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet*. 2020;396:759-769

15. Spertus JA, Fine JT, Elliott P, Ho CY, Olivotto I, Saberi S, Li W, Dolan C, Reaney M, Sehnert AJ, Jacoby D. Mavacamten for treatment of symptomatic obstructive hypertrophic cardiomyopathy (explorer-hcm): Health status analysis of a randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet*. 2021;397:2467-2475

16. Heitner SB, Jacoby D, Lester SJ, Owens A, Wang A, Zhang D, Lambing J, Lee J, Semigran M, Sehnert AJ. Mavacamten treatment for obstructive hypertrophic cardiomyopathy: A clinical trial. *Ann Intern Med*. 2019;170:741-748

17. Ho CY, Mealiffe ME, Bach RG, Bhattacharya M, Choudhury L, Edelberg JM, Hegde SM, Jacoby D, Lakdawala NK, Lester SJ, Ma Y, Marian AJ, Nagueh SF, Owens A, Rader F, Saberi S, Sehnert AJ,

Sherrid MV, Solomon SD, Wang A, Wever-Pinzon O, Wong TC, Heitner SB. Evaluation of mavacamten in symptomatic patients with nonobstructive hypertrophic cardiomyopathy. *J Am Coll Cardiol*. 2020;75:2649-2660

18. Gollapudi SK, Ma W, Chakravarthy S, Combs AC, Sa N, Langer S, Irving TC, Nag S. Two classes of myosin inhibitors, blebbistatin, and mavacamten, stabilize beta-cardiac myosin in different structural and functional states. *Journal of molecular biology*. 2021:167295

19. Craig R, Padron R. Structural basis of the super- and hyper-relaxed states of myosin ii. *J Gen Physiol*. 2022;154

20. Rohde JA, Roopnarine O, Thomas DD, Muretta JM. Mavacamten stabilizes an autoinhibited state of two-headed cardiac myosin. *Proceedings of the National Academy of Sciences of the United States of America*. 2018;115:E7486-E7494

21. Gollapudi SK, Yu M, Gan QF, Nag S. Synthetic thick filaments: A new avenue for better understanding the myosin super-relaxed state in healthy, diseased, and mavacamten-treated cardiac systems. *The Journal of biological chemistry*. 2021;296:100114

22. Ma W, Henze M, Anderson RL, Gong H, Wong FL, Del Rio CL, Irving T. The super-relaxed state and length dependent activation in porcine myocardium. *Circulation research*. 2021;129:617-630

23. Ma W, Nag S, Gong H, Qi L, Irving T. Cardiac myosin filaments are directly regulated by calcium. *J Gen Physiol*. 2022;154 e202213213

24. Chu S, Muretta JM, Thomas DD. Direct detection of the myosin super-relaxed state and interacting-heads motif in solution. *The Journal of biological chemistry*. 2021;297:101157

25. Ma W, Irving TC. Small angle x-ray diffraction as a tool for structural characterization of muscle disease. *Int J Mol Sci*. 2022;23

26. del Rio CL, Yukie Ueyama, Dale C Baker, Shana R Dalton, Laurence Lucats, Philip Janiak, Bari Olivier, Joseph Lambing, Marc J Evanchik, Eric M Green. In vivo cardiac effects of mavacamten (myk-461): Evidence for negative inotropy and improved compliance. *Circulation*. 2017;136:A20593-A20593

27. Sewanan LR, Shen S, Campbell SG. Mavacamten preserves length-dependent contractility and improves diastolic function in human engineered heart tissue. *American journal of physiology. Heart and circulatory physiology*. 2021;320:H1112-H1123

28. Awinda PO, Bishaw Y, Watanabe M, Guglin MA, Campbell KS, Tanner BCW. Effects of mavacamten on ca(2+) sensitivity of contraction as sarcomere length varied in human myocardium. *Br J Pharmacol*. 2020;177:5609-5621

29. Margossian SS, Lowey S. Preparation of myosin and its subfragments from rabbit skeletal muscle. *Methods in enzymology*. 1982;85 Pt B:55-71

30. Solaro RJ, Pang DC, Briggs FN. The purification of cardiac myofibrils with triton x-100. *Biochimica et biophysica acta*. 1971;245:259-262

31. Huxley HE. Electron microscope studies on the structure of natural and synthetic protein filaments from striated muscle. *Journal of molecular biology*. 1963;7:281-308

32. Davis JS. Assembly processes in vertebrate skeletal thick filament formation. *Annu Rev Biophys Biophys Chem*. 1988;17:217-239

33. Kawas RF, Anderson RL, Ingle SRB, Song Y, Sran AS, Rodriguez HM. A small-molecule modulator of cardiac myosin acts on multiple stages of the myosin chemomechanical cycle. *The Journal of biological chemistry*. 2017;292:16571-16577

34. National Research Council (U.S.). Committee for the Update of the Guide for the Care and Use of Laboratory Animals., Institute for Laboratory Animal Research (U.S.), National Academies Press (U.S.). *Guide for the care and use of laboratory animals*. Washington, D.C.: National Academies Press; 2011.

797   35.   Leary S. *Avma guidelines for the euthanasia of animals: 2013 edition*. Journal of the American
798   Veterinary Medical Association: American Veterinary Medical Association 2013.

799   36.   Ma W, Gong H, Jani V, Lee KH, Landim-Vieira M, Papadaki M, Pinto JR, Aslam MI, Cammarato A,
800   Irving T. Myofibril orientation as a metric for characterizing heart disease. *Biophysical journal*.
801   2022;121:565-574

802   37.   Ma , W, McMillen T, Childers M, Gong H, Regnier M, Irving T. Structural off/on transitions of
803   myosin in relaxed porcine myocardium predict calcium activated force. *Proceedings of the*
804   *National Academy of Sciences of the United States of America*. 2023;120:e2207615120

805   38.   Fischetti R, Stepanov S, Rosenbaum G, Barrea R, Black E, Gore D, Heurich R, Kondrashkina E,
806   Kropf AJ, Wang S, Zhang K, Irving TC, Bunker GB. The biocat undulator beamline 18id: A facility
807   for biological non-crystalline diffraction and x-ray absorption spectroscopy at the advanced
808   photon source. *J. Synchrotron Radiat*. 2004;**11,**:399-405

809   39.   Jiratrakanvong J, Shao J, Menendez M, Li X, Li J, Ma W, Agam G, Irving T. Musclex: Software suite
810   for diffraction x-ray imaging v1.13.1. Doi:10.5281/zenodo.1195050. 2018

811   40.   Ma W, Gong H, Irving T. Myosin head configurations in resting and contracting murine skeletal
812   muscle. *Int J Mol Sci*. 2018;19

813   41.   Ma W, Gong H, Kiss B, Lee EJ, Granzier H, Irving T. Thick-filament extensibility in intact skeletal
814   muscle. *Biophys J*. 2018;115:1580-1588

815   42.   Itoh T, Kawabe M, Nagase T, Endo K, Miyoshi M, Miyahara K. Body surface area measurement in
816   laboratory miniature pigs using a computed tomography scanner. *J Toxicol Sci*. 2016;41:637-644

817   43.   Mijailovich SM, Prodanovic M, Poggesi, C., Geeves MA, Regnier M. Multiscale modeling of twitch
818   contractions in cardiac trabeculae. *Journal of General Physiology*. 2021;153 e202012604

819   44.   Prodanovic M, Geeves MA, Poggesi C, Regnier M, Mijailovich SM. Effect of myosin isoforms on
820   cardiac muscle twitch of mice, rats and humans. *Int J Mol Sci*. 2022;23

821   45.   Mijailovich SM, Prodanovic M, Poggesi C, Powers JD, Davis J, Geeves MA, Regnier M. The effect
822   of variable troponin c mutation thin filament incorporation on cardiac muscle twitch
823   contractions. *Journal of molecular and cellular cardiology*. 2021;155:112-124

824   46.   Walker JS, Li X, Buttrick PM. Analysing force-pca curves. *Journal of muscle research and cell*
825   *motility*. 2010;31:59-69

826   47.   Jani V, Qian W, Yuan S, Irving T, Ma W. Emd-57033 augments the contractility in porcine
827   myocardium by promoting the activation of myosin in thick filaments. *Int J Mol Sci*. 2022;23

828   48.   Linari M, Brunello E, Reconditi M, Fusi L, Caremani M, Narayanan T, Piazzesi G, Lombardi V,
829   Irving M. Force generation by skeletal muscle is controlled by mechanosensing in myosin
830   filaments. *Nature*. 2015;528:276-279

831   49.   Ma W, Childers M, Murray J, Moussavi-Harami F, Gong H, Weiss R, Daggett V, Irving T, Regnier
832   M. Myosin dynamics during relaxation in mouse soleus muscle and modulation by 2'-deoxy-atp.
833   *The Journal of physiology*. 2020;598:5165-5182

834   50.   Mijailovich SM, Prodanovic M, Poggesi C, Geeves MA, Regnier M. Multiscale modeling of twitch
835   contractions in cardiac trabeculae. *J Gen Physiol*. 2021;153

836   51.   Toepfer CN, Garfinkel AC, Venturini G, Wakimoto H, Repetti G, Alamo L, Sharma A, Agarwal R,
837   Ewoldt JF, Cloonan P, Letendre J, Lun M, Olivotto I, Colan S, Ashley E, Jacoby D, Michels M,
838   Redwood CS, Watkins HC, Day SM, Staples JF, Padron R, Chopra A, Ho CY, Chen CS, Pereira AC,
839   Seidman JG, Seidman CE. Myosin sequestration regulates sarcomere function, cardiomyocyte
840   energetics, and metabolism, informing the pathogenesis of hypertrophic cardiomyopathy.
841   *Circulation*. 2020;141:828-842

842   52.   Matsubara I, Suga H, Yagi N. An x-ray diffraction study of the cross-circulated canine heart. *The*
843   *Journal of physiology*. 1977;270:311-320

844   53.   Matsubara I, Yagi N, Endoh M. Behaviour of myosin projections during the staircase
845   phenomenon of heart muscle. *Nature*. 1978;273:67

846   54.   Wheeler MT, Olivotto I, Elliott PM, Saberi S, Owens AT, Maurer MS, Masri A, Sehnert AJ,
847   Edelberg JM, Chen YM, Florea V, Malhotra R, Wang A, Oreziak A, Myers J. Effects of
848   mavacamten on measures of cardiopulmonary exercise testing beyond peak oxygen
849   consumption: A secondary analysis of the explorer-hcm randomized trial. *JAMA Cardiol*.
850   2023;8:240-247

851   55.   Xiao RP. Beta-adrenergic signaling in the heart: Dual coupling of the beta2-adrenergic receptor
852   to g(s) and g(i) proteins. *Sci STKE*. 2001;2001:re15

853   56.   Gresham KS, Stelzer JE. The contributions of cardiac myosin binding protein c and troponin i
854   phosphorylation to beta-adrenergic enhancement of in vivo cardiac function. *The Journal of*
855   *physiology*. 2016;594:669-686

856   57.   Kuschel M, Zhou YY, Spurgeon HA, Bartel S, Karczewski P, Zhang SJ, Krause EG, Lakatta EG, Xiao
857   RP. Beta2-adrenergic camp signaling is uncoupled from phosphorylation of cytoplasmic proteins
858   in canine heart. *Circulation*. 1999;99:2458-2465

859   58.   Yamasaki R, Wu Y, McNabb M, Greaser M, Labeit S, Granzier H. Protein kinase a phosphorylates
860   titin's cardiac-specific n2b domain and reduces passive tension in rat cardiac myocytes.
861   *Circulation research*. 2002;90:1181-1188

862   59.   Sun S, Ferguson BS, Rio CLd. Abstract 14837: Pro-compliant effects of mavacamten are
863   preserved in the setting of β-adrenergic receptor blockade: In vivo and ex vivo evidence.
864   *Circulation*. 2019;140:A14837-A14837

865   60.   Jacoby DL, Wheeler MT, Elliott PM, Saberi S, Sehnert AJ, Edelberg JM, Li W, olivotto i. Abstract
866   10201: Efficacy of mavacamten in patients with symptomatic hypertrophic cardiomyopathy:
867   Sub-group analyses by background beta-blocker use from the explorer-hcm and mava-lte
868   studies. *Circulation*. 2021;144:A10201-A10201