Comprehensive analysis of nasal IgA antibodies induced by intranasal administration of the
SARS-CoV-2 spike protein.

Kentarou Waki [a] Hideki Tani [b], Eigo Kawahara [c,d] , Yumiko Saga [b], Takahisa Shimada [b], Emiko Yamazaki
[b], Seiichi Koike [e] , Yoshitomo Morinaga [c,d], Masaharu Isobe [d,e], Nobuyuki Kurosawa [d,e]

[a] Laboratory of Molecular and Cellular Biology, Graduate School of Science and Engineering for
Education, University of Toyama, Toyama, Japan

[b] Department of Virology, Toyama Institute of Health, Toyama, Japan

[c] Department of Microbiology, Toyama University Graduate School of Medicine and Pharmaceutical
Sciences, Toyama, Japan

[d] Center for Advanced Antibody Drug Development, University of Toyama, Toyama, Japan

[e] Laboratory of Molecular and Cellular Biology, Graduate School of Innovative Life Science, University
of Toyama, Toyama, Japan

Corresponding author: Nobuyuki Kurosawa

E mail: kurosawa@eng.u-toyama.ac.jp

Laboratory of Molecular and Cellular Biology, Graduate School of Innovative Life Science, University of
Toyama, 3190 Gofuku, Toyama-shi, Toyama, 930-8555, Japan

Abstract:

Intranasal vaccination is an attractive strategy for preventing COVID-19 disease as it stimulates the production of multimeric secretory immunoglobulin A (IgA), the predominant antibody isotype in the mucosal immune system, at the target site of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) entry. Currently, intranasal vaccine efficacy is evaluated based on the measurement of polyclonal antibody titers in nasal lavage fluid. However, how individual multimeric secretory IgA protects the mucosa from SARS-CoV-2 infection remains to be elucidated. To understand the precise contribution and molecular nature of multimeric secretory IgA induced by intranasal vaccines, we developed 99 monoclonal IgA clones from nasal mucosa and 114 monoclonal IgA or IgG clones from nonmucosal tissues of mice that were intranasally immunized with the SARS-CoV-2 spike protein. The nonmucosal IgA clones exhibited shared origins and common and unique somatic mutations with the related nasal IgA clones, indicating that the antigen-specific plasma cells in the nonmucosal tissues originated from B cells stimulated at the nasal mucosa. Comparing the spike protein binding reactivity, angiotensin-converting enzyme-2-blocking and *in vitro* SARS-CoV-2 virus neutralization of monomeric and multimeric secretory IgA pairs recognizing different epitopes showed that even non-neutralizing monomeric IgAs, which represents 70% of the nasal IgA repertoire, can protect against SARS-CoV-2 infection when expressed as multimeric secretory IgAs. We also demonstrated that the intranasal administration of multimeric secretory IgA delivered as prophylaxis in the hamster model reduced infection-induced weight loss. Our investigation is the first to demonstrate the function of nasal IgA at the monoclonal level, showing that nasal immunization can provide effective immunity against SARS-CoV-2 by inducing multimeric secretory IgAs at the target site of the virus infection.

Introduction

Immunoglobulin A (IgA) is differentially distributed between the systemic and mucosal immune systems (*Li et al. , 2020*). Monomeric IgA (M-IgA) is predominantly present in serum, whereas secretory IgA (S-IgA) is the most prevalent in mucous secretions. S-IgA comprises two IgA units and one J chain, which are synthesized and assembled in local plasma cells. Secretory components expressed on the basolateral surface of mucosal epithelial cells bind the IgA complex through the J chain and transport the molecule to the apical cell membrane (*Woof and Kerr, 2006*). Recent studies have shown that S-IgA in the nasal mucosa exists not only as dimers but also as trimers and tetramers in the human upper respiratory mucosa (*Saito et al. , 2019*). S-IgA plays an important role in the protection and homeostatic regulation of the airway, intestinal and vaginal epithelium through a process known as immune exclusion and immunosuppression (*Matsumoto, 2022*). Recent studies have demonstrated that multimeric S-IgA is more effective and provides greater cross-protection than IgG and M-IgA (*Okuya et al. , 2020b*) (*Asahi et al. , 2002*) (*Dhakal et al. , 2018*) (*Asahi-Ozaki et al. , 2004*) (*Wang et al. , 2021*).

COVID-19 is a disease caused by infection with severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) that causes mild upper respiratory symptoms in most cases, but some patients develop bilateral pneumonia with acute respiratory distress (*Budinger et al. , 2021*). The SARS-COV-2 spike protein expressed on the virus surface is a multidomain homotrimer protein composed of an S1 domain consisting of an N-terminal domain (NTD) and receptor binding domain (RBD) and an S2 domain that mediates fusion of the virus and host cell membrane. Viral infection is initiated through the interaction between the RBD and the host receptor angiotensin-converting enzyme-2 (ACE2) (*Jackson et al., 2022*). Thus, the spike protein is the main target for current vaccine development because antibodies against this protein can neutralize the infection. Currently approved intramuscular COVID-19 vaccines predominantly induce a systemic immune response by producing IgG in serum before they cause severe tissue damage, resulting in a high degree of efficacy (*Polack et al. , 2020*). However, these vaccines mainly induce IgG and M-IgA in serum; they do not induce S-IgA, which coats the upper respiratory tract mucosal surface (*DeFrancesco, 2020*). The emergence of highly infectious SARS-CoV-2 omicron variants may undermine the therapeutic efficacy of vaccines, requiring more effective vaccination to prevent SARS-CoV-2 infection. Since nose epithelial cells are a primary target for SARS-CoV-2, intranasal vaccinations that induce S-IgA in the upper respiratory tract are desirable for protection against the infection and transmission of the virus (*Sungnak et al. , 2020*). To date, some intranasal vaccines are under development and have shown robust mucosal and humoral immune responses in animal models (*Houston, 2023*) (*Chavda et al. , 2021*) (*Alu et al. , 2022*) (*Barrett et al. , 2021, Bricker et al. , 2021, D'Arco et al. , 2021, Kim et al. , 2021*) (*Vabret et al. , 2020*) (*Ohtsuka et al. , 2021*). However, there have been few published clinical trials of intranasal SARS-CoV-2 vaccines on human (*Ewer et al. , 2021, Zhu et al. , 2023*). To evaluate the response to intranasal vaccines, researchers have utilized nasal lavage fluids containing polyclonal S-IgA (*Maltseva et al. , 2022*) (*Gianchecchi et al. , 2019*) (*Wong et al. , 2022*) (*Afkhami et al. , 2022*) (*Sui et al. , 2021*) (*Azzi et al. , 2022*). However, obtaining information about the functions of individual antibodies in the polyclonal anti-viral mucosal repertoire is challenging due to the broad nature of such assays. It has also been shown that intranasal vaccination induces a systemic antibody response, but the origin of these antibodies has not previously been investigated at the molecular level (*Tumpey et al. , 2001*) (*Zhu et al. , 2023*). Thus, intranasal vaccines require novel approaches to evaluate the quality and quantity of IgA response.

To understand the precise contribution and molecular nature of S-IgA induced by intranasal vaccines in relation to its antiviral function, it is essential to develop monoclonal S-IgAs from plasma cells localized in the nasal mucosa and study their function. However, because of the difficulties in developing monoclonal IgA antibodies from nasal mucosal tissue, many studies have used artificial switching of IgG to IgA for recombinant production, and this approach has been used to study the protective properties of IgA against pathogens (*Ejemel et al. , 2020, Saito et al. , 2019*). Such technological limitations have

hampered the biochemical and clinical evaluation of intranasal vaccination at the molecular level.

Our group has established robust protocols for isolating single antigen-specific plasma cells from a

variety of animals and the high-throughput production of monoclonal antibodies (*Kurosawa et al. , 2012*).

Here, using this technology, we developed 213 antigen-specific monoclonal antibodies from plasma cells

derived from mice that were intranasally immunized with a stabilized SARS-CoV-2 spike trimer protein

(Spike) and demonstrated that intranasal immunization induced potentially nasal-derived antibodies in the

spleen, lung, and blood. Analysis of the properties of monoclonal IgAs recognizing different epitopes

revealed that multimerization of M-IgAs could induce neutralizing activity *in voitro* and *in vivo*. These

results are the first of their kind to be demonstrated by obtaining monoclonal S-IgAs from nasal mucosal

tissue. Our data provide insights into the nasal immunity that may help shape future intranasal vaccine

development.

Results

Development of anti-spike monoclonal M-IgAs from plasma cells localized in nasal mucosa.

To evaluate the immune response after the intranasal administration of the SARS-CoV-2 spike protein of

the Wuhan-Hu-1/D614G strain (Spike$_{Wuhan}$), nasal lavage fluids and serum were collected from the mice

one week after the last immunization. The intranasal administration of Spike$_{Wuhan}$ induced high levels of

antigen-specific IgA but not IgG in nasal lavage fluids compared with those of the phosphate-buffered

saline (PBS) control (Fig. 1A and Supplementary Fig.1).

Fig. 1. Isolation of antigen-specific IgA[+] plasma cells from mice that were intranasally immunized with Spike_{Wuhan}.

(A) Mice were inoculated intranasally with 10 μg of SpikeWuhan and 1 μg of Cholera toxin as an adjuvant in 10 μl of PBS, delivering the fluid dropwise into the nostril a total of three times at three-week intervals. Nasal lavage fluid and serum were collected from the mice one week after the last immunization, and antibody responses were evaluated using ELISA. The antibody titers are expressed as optical density (OD_{450}) value per total protein in nasal lavage fluids or serum.

(B) FACS gating strategy for the isolation of S1-specific plasma cells from mice. Plots represent the sequential gating strategy. Lymphocytes (R1 gate) were stained with anti-CD138 and ER-Tracker to enrich plasma cells (CD138[+] ER-Tracker[High] fraction, R2 gate). IgA[+] plasma cells gated in R3 were selected from the R2-gated plasma cell fraction by staining with anti-IgA antibody. The antigen-specific

plasma cells gated in R4 were further selected from the R3-gated Ig A$^+$ plasma cell fraction by staining

with S1 domain of Wuhan SARS-CoV-2 spike protein (S1). The numbers indicate the percentages of cells

in the gated area. A total of 100,000 events were recorded. Representative data from the No. 1 mouse are

shown.

Elevated levels of antigen-specific IgA and IgG responses were detected in serum from the immunized

mice, consistent with a prior study demonstrating that intranasal vaccinations induce both nasal and

serum IgA levels (*Sterlin et al. , 2021*) (*Maltseva et al. , 2022*). We selected three immunized mice with

high nasal IgA titers and isolated antigen-specific plasma cells from the nasal mucosa, spleen, lung and

blood. Cells isolated from these tissues were stained with fluorescently labeled anti-CD138, ER-Tracker

Blue-White DPX (ER-Tracker), anti-IgA and S1 domain of Spike$_{Wuhan}$ (S1), and then antigen-specific

plasma cells defined as anti-CD138$^+$, ER-tracker^{high}, anti-IgA$^+$ and S1$^+$ were isolated by

fluorescence-activated cell sorting (FACS). ER-Tracker is a fluorescent dye that is highly selective for the

endoplasmic reticulum of living cells. Because plasma cells have an expanded endoplasmic reticulum for

properly folding and secreting large quantities of antibodies, using ER-Tracker along with anti-CD138

facilitates the isolation of plasma cells from lymphocytes without the need for additional antibodies.

(Kurosawa et al., 2012). Consistent with the pattern of the IgA response observed in nasal lavage fluid,

the proportions of antigen-specific IgA$^+$ plasma cells in the nasal mucosa of immunized mice were

significantly higher than those of control mice (Fig. 1B and Supplementary file 1A).

Cognate pairs of immunoglobulin heavy chain variable (VH) and kappa light chain variable genes (VL)

were amplified by rapid amplification of 5' cDNA ends PCR from the single-sorted cells. After

constructing full-length immunoglobulin heavy and kappa light chain genes, antibodies were expressed

by DNA transfection into CHO-S cells. Antigen-specific antibody clones were identified by

enzyme-linked immunosorbent assay (ELISA) with immobilized Spike$_{Wuhan}$. After sequencing the entire

coding region of heavy and light chain genes, the antibodies sharing the same V-(D)-J genes were

grouped (Supplementary Table 1). Then, representative antibody clones from each group were analyzed

for their binding to the spike protein RBD of Wuhan, Beta, Kappa and Delta variants and an NTD of

Spike$_{Wuhan}$. The antibodies were also analyzed for their ability to block ACE2 binding to RBDs and to

neutralize the Wuhan pseudotyped virus. Based on their properties, each group of antibodies was

arbitrarily categorized into five types. Type 1: anti-RBD, *ACE2 blocking* neutralizing *antibody; Type 2:*

anti-RBD, *ACE2 blocking* non-neutralizing *antibody; Type 3:* anti-RBD, *non-ACE2 blocking*

non-neutralizing *antibody; Type 4:* anti-NTD, *non-ACE-2 blocking* non-neutralizing *antibody; and Type*

5: non-ACE-2 blocking non-neutralizing *antibody targeting* epitopes other than RBD and NTD.

Intranasal immunization induces functionally diverse IgA in the nasal mucosa and spleen.

We conducted a detailed characterization of antibodies obtained from the No. 1 mouse, as many

antigen-specific M-IgA clones were obtained. Of the 51 nasal M-IgA clones analyzed, they were

classified into 11 groups based on their V-(D)-J usage, with the majority (83%) clonally expanding into

four major clusters (G2, G3, G4 and G10) and the remainder (17%) scattered across branches (Fig. 2A

and B).

Fig. 2 Intranasal immunization induces functionally diverse antibodies in the nasal mucosa and

spleen.

(A) Characterization of S1-specific monoclonal antibodies obtained from No. 1 mouse. The heatmap

represents the relative intensity of antibody binding to RBDs and blocking of the RBD-ACE2 interaction.

Blue (0–25%), green (25–50), orange (50–75%) and red (>75%). NTD binding was considered positive

(+) when the OD at 405 nm was >0.3 after the background was subtracted. Neutralizing activity was

considered positive (+) when the antibody suppressed Wuhan pseudotyped virus infection by 50%. The

figure reports values from a single experiment. UN, antibody type not determined. ND, antibody activity not determined.

(B) Maximum-likelihood phylogenetic tree of the VH and VL chains of the S1-specific antibodies.

Different colored fonts indicate antibodies obtained from the nose (red), spleen (blue), and lung (green). Antibody groups are indicated by bands on the outer ring. The color of the ring indicates antibody types: Type 1 (red), Type 2 (orange), Type 3 (green), Type 4 (blue) and Type 5 (gray). The antibody group is defined as clones using the same V-(D)-J usage and having an overall sequence identity of at least 95% from the signal peptide to framework 4 (FR4). The prefixes N, S and L in the antibody clone numbers refer to antibodies derived from the nose, spleen and lung, respectively. The suffixes A, G and K in the antibody clone numbers refer to alpha, gamma and kappa chain, respectively.

(C) Nucleotide sequence arraignment of VH and VL genes in the G2 and G3 antibodies form No. 1 mouse. The VH and VL sequences from the beginning of the signal peptide through the end of FR4 are shown as horizontal lines. Nucleotide changes relative to S632A and N109A are depicted as vertical bars across the horizontal lines. Different colored fonts indicate antibodies derived from the nose (red) and spleen (blue). Antibody phylogenetic trees based on VH/VK paired sequences are depicted. Gray circles represent the hypothetical germline configuration. White circles represent hypothetical ancestors. Colors indicate nasal (red) and splenic (blue) antibodies. Circles and squares indicate IgA and IgG, respectively.

The antibodies in the G1-G4 clusters were categorized as Type 1. The G1 antibody appeared only once. This clone displayed potent binding to the Wuhan, Kappa and Delta RBD, and moderately to the Beta RBD. It also blocked ACE2 binding to all RBDs except the Beta RBD. The antibodies in G2-4 clusters bound to Wuhan and Delta RBD but not Beta and Kappa RBD. They were able to block ACE2 binding to the RBD of Wuhan and Delta but not to the RBD of Beta and Kappa. The antibodies in G5-G9 clusters were categorized as Type 2, bound to all four RBD variants, and uniformly blocked ACE2 binding to the RBDs of all four strains. G10 antibodies were classified as Type 3, accounting for 38% of the nasal IgA repertoire. They bound to the RBDs of all four strains but did not prevent ACE2 binding to the RBDs of any of the strains. The G11 antibody, classified as Type 4, was an anti-NTD antibody that did not prevent ACE2 binding to the RBDs of any strains.

Intranasal immunization induces potentially nasal-derived antibodies in the spleen, lung, and blood.

Previous reports suggest that class-switched, affinity-matured B cells selected by NALT migrate to the regional lymph node and then return to the nasal mucosa and non-mucosal tissues to differentiate into

plasma cells (*Shimoda et al. , 2001*). If this is the case, a fraction of S1-specific plasma cells differentiated from nose-originated B cells may reside in the spleen and produce antibodies. To directly evaluate the cellular origin of the anti-S1 antibodies in nonmucosal tissues, we analyzed the presence of S1-specific plasma cells in the spleen, lung, bone marrow and Peyer's patches. FACS analysis of splenocytes harvested from No. 1 mice showed the presence of S1-specific IgA$^+$ plasma cells, but the splenocytes harvested from control mice did not exhibit S1-specific IgA$^+$ plasma cells (Fig. 1B and Supplementary file 1B). Single-cell-based immunoglobulin gene cloning from the splenocytes of No.1 mouse resulted in the successful production of 57 S1-specific monoclonal IgAs. DNA sequence analysis of these clones revealed significant clonal overlap between the nose and spleen. (Fig. 2A and B). Clonal overlap was found in the G2-G3, G5, G6 and G10 clusters, in which G10 possessed the most expanded splenic clones, as in the case of nasal IgA (Fig. 2). Further investigation was conducted to determine if plasma cells expressing S1-specific IgG$^+$ antibody, which share V-(D)-J with nasal IgA, could be identified in the spleen. Although the number of antigen-specific IgG$^+$ plasma cells was limited, eight IgG clones specific for S1 were isolated, among which five were potentially nasal-derived. When we attempted to isolate antigen-specific IgA$^+$ plasma cells from the lung, nine S1-specific IgA clones were obtained, among which three were potentially nasal-derived. The gut mucosa and bone marrow possessed a large population of IgA$^+$ plasma cells. However, we could not detect S1-specific IgA$^+$ plasma cells in Peyer's patches and bone marrow (Supplementary file 1B).

Next, we focused on the expanded antibody groups G2, G3 and G10 to analyze the patterns of somatic hypermutation (SHM). We found complex patterns of shared and unique SHMs in antibodies obtained from the nose, spleen and lung. In G2, the VHs of the three splenic clones (S530A, S208G, and S619A) had the same sequence, except for framework 4 of an IgG clone (S208G) and had three shared SHMs with the nasal clones (N219A and N226A). In addition, the VKs of the three splenic clones and N226A had the same sequence. In Group 3, seven nasal clones (N109A, N112A, N135A, N139A, N244A, N245A and N715A) and two splenic clones (S545 and S607) were 100% identical in the nucleotide sequence, with N135A and N245 differing from the other clones by only one nucleotide in their VKs (Fig. 2C). Sequence analysis of the most expanded group, Group 10, also demonstrated clonal overlap in the nose, lung and spleen. For example, the VHs of a nasal clone (N221), spleen clones (S647, S503, S640 and S543) and a lung clone (L009) were 100% identical, with S647 differing from the other clones by only one nucleotide in VK (Supplementary file 2). Antibody lineage analysis of these clones suggests that each family of related antibodies originates from a common ancestor gene that was subjected to class switching and SHM. We also developed S1-specific Abs from No. 2 and No. 3 mice and analyzed their sequences (Fig. 3 and Supplementary file 3).

Fig. 3. Characterization of S1-specific monoclonal antibodies obtained from No. 2 mouse.

(A) A total of 51 S1-reactive antibodies were analyzed for their properties as in Fig 2A. UN, antibody type not determined. ND, antibody activity not determined.

(B) Maximum-likelihood phylogenetic tree of the VH and VL chains of the S1-specific antibodies.

Different colored fonts indicate antibodies obtained from the nose (red), spleen (blue), lung (green) and blood (magenta). Antibody groups are indicated by bands on the outer ring. The color of the band indicates antibody types: Type 1 (red), Type 2 (orange), Type 3 (green), Type 4 (blue) and Type 5 (gray). The prefixes N, S and L in the antibody clone numbers refer to antibodies derived from the nose, spleen and lung, respectively. The suffixes A, G and K in the antibody clone numbers refer to alpha, gamma and kappa chain, respectively.

(C) Nucleotide sequence arraignment of VH and VL genes in the G6 and G8 antibodies from No. 2 mouse. The. nucleotide changes relative to N5120A and N5105A are depicted as vertical bars across the horizontal lines. Different colored fonts indicate antibodies derived from the nose (red), spleen (blue), lung (green) and blood (magenta). Antibody phylogenetic trees based on VH/VK paired sequences are depicted. Gray circles represent the hypothetical germline configuration. White circles represent hypothetical ancestors. Colors indicate nasal (red), splenic (blue), lung (green) and blood (magenta) antibodies.

In No. 2 mice, we obtained 29 nasal IgAs, 15 splenic antibodies (13 IgAs and two IgGs), five lung IgAs and six blood IgAs. They were classified into 17 groups based on their V-(D)-J usage, and each group was categorized as either Type 1, 2, 3, 4 or 5 based on antibody properties. (Fig. 3). Potentially nasal-derived clones were found in the spleen, lung and blood (G5, 6, and 8). These clones also showed a complex pattern of shared and unique SHM with nasal clones throughout the full length of the VH and VL genes. In mouse No. 3, we cloned 19 nasal IgAs and 22 splenic antibodies (17 IgAs and five IgGs). They were classified into 10 groups, and each group was categorized as either Type 1, 3, 4 or 5. A high degree of clonal overlap between the nose and spleen was found, in which 13 out of 22 splenic Abs were potentially nasal-derived clones (Supplementary file. 3). Analysis of the VH and VL repertoires of the three mice revealed that the expanded clones constituted varying fractions of the antibody repertoire among different mice despite having received the same antigen, and no group of antibodies stood out across mice, suggesting that individual mice had immunologically distinct responses (Supplementary figure 4). In all mice, only a few bound to the NTD, and most of them were RBD-directed. Approximately 30% of M-IgAs categorized as Type 1 showed neutralizing activity (Supplementary figure 4). Taken together, regarding the mutation frequency and the pattern of SHM, it can be assumed that B cells stimulated by nasal challenges were the major precursor of antigen-specific plasma cells in the spleen, lung and blood, which may contribute to antibody production in the lower respiratory tract and systemic circulation.

Multimerization of M-IgA enhances antigen-binding activity.

To examine how IgA multimerization affects antigen-binding activity, four representative clones (N5203, N142, N114 and N217) were selected from each type of antibody, and S-IgAs were expressed by

cotransfecting alpha heavy chain, kappa light chain, J-chain and secretory component into CHO-S cells.

Analysis of the purified S-IgAs by polyacrylamide gel electrophoresis (SDS□PAGE) revealed a band

corresponding to the alpha heavy chain, kappa light chain, and secretory component. Native PAGE

analysis revealed that the antibodies comprised three quaternary structures, including a dimer (~400 kDa),

a trimer (~550 kDa), and a tetramer (~750 kDa) at a molar ratio of 5:1:3 (Fig. 4A).

Fig. 4 Comparison of reactivity between monomeric and multimeric IgAs.

(A) Production of recombinant M-IgAs and S-IgAs. Recombinant M-IgA and S-IgA purified from the

culture supernatant of CHO cells were subjected to SDS□PAGE and Blue native-PAGE analysis. Bands

corresponding to a monomer (M), dimer (Di), trimer (Ti) and tetramer (Te) are shown. H, α heavy chain;

L, light chain; J, J chain; SC, secretory component; M, M-IgA; S, S-IgA

(B) Binding dynamics of M-IgAs and S-IgAs to Wuhan, Delta or Omicron spike protein by SPR. The S-IgAs used are a mixture of dimers, trimers, and tetramers. The observed values refrect the average affinity of the S-IgAs. The curves shown are representative of two or three determinations. RU, resonance units. The table shows the association (ka) ($M^{-1}s^{-1}$), dissociation (kd) (s^{-1}) rate constants and apparent equilibrium dissociation constants (KD) expressed as the mean of two or three determinations (lower panel).

Then, we compared the binding kinetics of each pair of M-IgA/S-IgA by surface plasmon resonance (SPR) with immobilized $Spike_{Wuhan}$, $Spike_{Delta}$ or $Spike_{Omicron}$. As shown in Fig. 4B, N5203 M-IgA, which has a moderate binding affinity to $Spike_{Wuhan}$ (apparent equilibrium constant, K_D=5.2E-9), acquired dramatic enhancement of the binding activity to $Spike_{Wuhan}$ after multimerization (K_D=1.3E-13). However, multimerization did not enhance the binding activity to $Spike_{Delta}$ (K_D=2.1E-9). N142 M-IgA, which has a high affinity for $Spike_{Wuhan}$ (K_D=4.5E-11) and $Spike_{Delta}$ (K_D=7.8E-11), did not show enhanced binding activity to them after multimerization. Almost the same phenomenon was found for N114 M-IgA, which has a high affinity for $Spike_{Wuhan}$ (K_D=6.8E-11) and $Spike_{Delta}$ (K_D=9.7E-11). N142 M-IgA, which has the lowest affinity to $Spike_{Wuhan}$ (K_D=2.1E-8), also acquired dramatic enhancement of the binding activity to $Spike_{Wuhan}$ after multimerization (K_D=6.4E-11) but not to $Spike_{Delta}$. All antibodies showed little or marginal levels of binding to $Spike_{Omicron}$. These results suggest that the degree of avidity of S-IgAs depends on the affinity of the parent monomeric antibody: antibodies with low or intermediate affinity in the monomeric state (N5203 and N217) can increase their avidity by multimerization but not antibodies with high affinity in the monomeric state (N142 and N114). These results are consistent with recent work by Saito, S. et al., who examined the function of multimerized IgA against influenza viruses (*Saito et al. , 2019*).

Multimerization facilitates stronger neutralization activity in non-neutralizing M-IgA

Since ACE2-blocking activity is one indicator for evaluating the neutralizing activity of anti-SARS-CoV-2 antibodies, we next examined whether the multimerization of IgAs influences the RBD-ACE2 interaction by competitive ELISA (Fig. 5A).

Fig. 5 Multimerization facilitates the neutralization activity of nonneutralizing M-IgAs.

(A) Graphs of the competitive ELISA results showing the binding of biotinylated ACE2 to the
immobilized Wuhan, Delta or Omicron RBD in the presence of antibodies. The results are expressed as
the mean ± SD of three technical replicates. The IC_{50} values of the indicated antibodies that inhibit the
RBD-ACE2 interaction are shown in the diagrams.

(B) Comparison of neutralization activity between M-IgaA and S-IgA against SARS-CoV-2 pseudotyped
viruses. Neutralization curves of the indicated antibody against pseudotyped viruses bearing spike
proteins of Wuhan, Delta or Omicron are shown. Pseudotyped viruses preincubated with antibodies at the
indicated concentrations were used to infect VeroE6 cells, and luciferase activities in cell lysates were
measured at 20 h post-transduction to calculate infection (%) relative to nonantibody-treated controls. The
results are expressed as the mean ± SD of three technical replicates. The NT_{50} values of the indicated
antibodies are shown in the diagrams. Antibodies that did not reach >70% inhibition at the highest
concentration tested were listed as data not determined (ND).

(C) Comparison of neutralization potential between M-IgA and S-IgA against authentic SARS-CoV-2 BA.1. The neutralizing potential of the antibody was determined using an RT-PCR-based SARS-CoV-2 neutralization assay. VeroE6 cells preincubated with authentic SARS-CoV-2 BA.1 virus were incubated with the indicated antibodies at various concentrations. The virus in the cell culture medium was measured at 48 h post-transduction to calculate infection (%) relative to non-antibody-treated controls. The results are expressed as the mean ± SD of three technical replicates. The NT_{50} values of the indicated antibodies are shown in the diagrams. Antibodies that did not reach >50% inhibition at the highest concentration tested are listed as ND. **p<0.01

Although the multimerization of N5203 M-IgA led to the dramatic enhancement of $Spike_{Wuhan}$ binding, it only increased RBD_{Wuhan}-ACE2 blocking by 3.3-fold, and there were no differences in RBD_{Delta}-ACE2 blocking between M-IgA and S-IgA. In N142 M-IgA, multimerization slightly increased RBD_{Wuhan}-ACE2 blocking but not RBD_{Delta}-ACE2 blocking. Multimerization of N114 and N217 M-IgA did not enhance RBD_{Wuhan}-ACE2 and RBD_{Delta}-ACE2 blocking, as in the case of their monomeric forms. None of the antibodies showed $RBD_{Omicron}$-ACE2 blocking even at the highest antibody concentrations.

To analyze whether multimerization affects the functionality of M-IgAs, we tested all M-IgA/S-IgA pairs for their ability to neutralize pseudotyped lentiviruses bearing the Wuhan, Delta or Omicron spike protein (Fig. 5B). Both monomeric and multimeric forms of N5203 showed strong neutralizing activity against the Wuhan and Delta pseudotyped lentiviruses, but the concentrations required to achieve the 50% neutralizing titer (NT_{50}) were almost the same for both. Although both forms of N142 displayed RBD_{Wuhan}-ACE2 and RBD_{Delta}-ACE2 blocking, only the multimeric form showed neutralization activity against the Wuhan and Delta pseudotyped lentiviruses. Notably, N114 M-IgA and N217 M-IgA, which did not have RBD_{Wuhan}-ACE2 blocking activity, displayed neutralizing activity against the Wuhan but not the Delta pseudotyped lentivirus after multimerization. None of the antibodies exhibited neutralization activity against the Omicron pseudotyped lentivirus. We further tested the neutralization activity of these antibodies by using an authentic SARS-CoV-2 Wuhan strain (Fig. 5C). N5203 S-IgA showed twofold higher neutralization activity than its monomeric counterpart. As predicted from the results of the pseudotyped lentivirus assay, multimerization of N142, N114 and N217 M-IgA induced neutralization activity, while their monomeric counterparts failed to neutralize live virus at the highest antibody concentration. These results suggest that the neutralizing activity of S-IgAs is not solely attributed to the spike protein affinity and the RBD-ACE2 blocking activity of M-IgAs, and that the valence and binding mode between the epitope and paratope affect their function.

Next, we selected three types of S-IgAs (N5203, N142, and N217) and investigated whether nasally administered S-IgAs could protect against Wuhan SARS-CoV-2 infection in a Syrian hamster model as a prophylactic treatment (Fig.6). The animals in the control group showed significant and

progressive weight loss (20% by day 3) after the viral challenge. In contrast, animals treated with S-IgAs showed only a small weight loss (5% with N5203, 8% with N142, and 5% with N217). (Fig. 6A). Analysis of viral load at day 3 by qPCR for SARS-CoV-2 nucleoprotein (N2) RNA showed some hamsters treated with N5203 S-IgA or N217 S-IgA showed a trend of decreased virus production in nasal turbinate compared to the control, though this did not reach statistical significance. These antibodies did not reduce the viral load in the lungs. These results indicate that the prophylactic nasal administration of S-IgA confers a protective effect against SARS-CoV-2, but did not lead to the elimination of virus from the animals.

Fig. 6 Intranasal administration of S-IgA suppresses SARS-CoV-2 infection in Syrian hamsters.

(A) Experimental schedule. Three groups of hamsters received a single intranasal dose of 1.0 mg/kg of S-IgA three hours before infection (-3) for pre-exposure prophylaxis and at 24 hours (24) and 48 hours post-infection for early treatment, respectively. Control hamsters (n=3) received PBS at the same dose. On day 0, each hamster was intranasally challenged with the Wuhan SARS-CoV-2 virus (6×10^5 Median Tissue Culture Infectious Dose).

(B) Hamster body weights were recorded hourly (0, 24, 48 and 72 hours), and weight loss was defined as the percentage loss from 0 hours. Data represent the mean value $\pm$ SD at the indicated time points ($n = 3$) at the indicated time points and were analyzed using a Kruskal–Wallis one-way ANOVA.

(C) Animals were euthanized 72 hours post-infection, and RNA was extracted from nasal turbinates and lungs. The SARS-CoV-2 viral load was analyzed using qRT-PCR targeting the SARS-CoV-2 nucleoprotein. Assays were normalized relative to total RNA levels. Data represent the mean value (n = 3).

DISCUSSION

In this study, we generated hundreds of monoclonal antibodies from nasal mucosa and non-mucosal tissues, providing molecular evidence that intranasal immunization can induce both mucosal and systemic antibody responses. We also generated M-IgA and S-IgA with the same antigen recognition site for the Spike protein and compared their antiviral activities *in vitro*. SARS-CoV-2 research has mainly focused on anti-RBD neutralizing antibodies, while the role of non-neutralizing antibodies has not been adequately analyzed. We demonstrated that non-neutralizing M-IgAs, comprising 70% of the nasal IgA repertoire, showed strong neutralizing activity when expressed as S-IgAs. Nasal delivery of S-IgA offered potential prophylaxis against SARS-CoV-2 infection. The ability of nasal immunization to induce various neutralizing S-IgAs that recognize different epitopes may be a significant advantage not seen with injectable vaccines. Our data highlight the key role of S-IgA in the protective effect of mucosal immunity, which may be useful for better understanding how intranasal vaccines can help protect against SARS-CoV-2 infection.

It has been shown that the mucosal route of immunization elicits immune responses at local sites and systemic immune responses. (*Lapuente et al. , 2021*). Our analysis demonstrated that the spleen possessed a larger population of antigen-specific plasma cells that expressed potentially nasal-derived IgAs and IgGs. These results suggest that the B cells activated in the nasal immune system give rise to plasma cells that reside in the nasal mucosa and produce S-IgA, while the B cells also migrate to the spleen and produce M-IgA and IgG. The role of the spleen in the induction of potentially nasal-derived antibodies by intranasal immunization is yet to be determined. The antigen-specific plasma cells in the spleen may respond in cases of systemic infection by supplying M-IgAs and IgGs into the bloodstream or protecting the body against subsequent infection (*Afkhami et al. , 2022*) (*Sheikh-Mohamed et al. , 2022*). Although bone marrow and the intestines are reservoirs for IgA-producing B cells, we could not detect antigen-specific plasma cells in these tissues. The molecular mechanisms regulating potentially nasal-derived B-cell migration into the spleen but not into bone marrow and Peyer's patches remain to be determined. The extent of functional and reactive differences between M-IgA and IgG sharing the same V gene remains to be determined in this paper. Recent studies on replacing the Fc domain of IgG with IgA have shown conflicting data, with some antibodies being enhanced while others were weakened. This suggests that the degree of increase in antibody reactivity and functionality is not solely correlated with

the Fc domain but also depends on the selected epitopes and paratopes. (*Tudor et al. , 2012*) (*Muramatsu*

et al. , 2014) (*Ejemel et al. , 2020*) (*Saito et al. , 2019*).

As illustrated in Fig. 7, four types of antibodies exhibited different reactions between M-IgA and S-IgA

regarding spike protein binding, ACE2-blocking, and virus neutralization.

Fig.7 The affinity, ACE2 inhibitory activity, and the in vitro neutralizing activity of the indicated

antibodies are illustrated in a 3D scatter plot.

N5203 M-IgA, categorized as Type 1, binds to the RBD_{Wuhan} and RBD_{Delta}, inhibits ACE2-binding to

these RBDs, and effectively neutralizes Wuhan and Delta pseudotyped viruses. The multimeric form of

this antibody had a minimal effect on ACE-blocking and neutralization breadth despite the increased

affinity for $Spike_{Wuhan}$. This suggests that the increase in valency does not necessarily correlate with an

increase in functionality. N142 M-IgA, categorized as Type 2, binds to the RBD_{Wuhan} and RBD_{Delta},

inhibiting ACE2-binding to these RBDs, yet fails to neutralize the viruses. Interestingly, its S-IgA

exhibited neutralization activity against both pseudotyped viruses without enhancing spike binding or

ACE-blocking activity. These results suggest that the neutralizing activity of N142 S-IgA partially

depends on the ACE2-blocking activity of the corresponding M-IgA, but its mechanism differs from that

of N5203 S-IgA, whose monomeric form shows cross-neutralizing activity. Notably, two types of

non-ACE2-blocking, non-neutralizing M-Abs (N114 and N217), expressed as S-IgA, gained the ability to

neutralize the Wuhan but not the Delta pseudotyped viruses. These findings indicate that the two S-IgAs

exert their neutralizing activity through mechanisms apart from the general ACE2-RBD axis.

Recent reports show that antibodies that induce inter- or intraspike crosslinking can inhibit viral binding

or shedding by the host cell through steric hindrance or cause conformational changes in the spike

proteins (Galimidi et al., 2015) (Jackson et al., 2022) (Klein and Bjorkman, 2010). Such inter- and

intraspike cross-linking may be limited in M-IgAs and IgGs if the spike protein density on the virus

surface is low or if the epitopes are unfavorably located. We hypothesize that S-IgA could exert the above

effects through its multivalent arms, a mode of action that is hard to achieve with M-IgA, and factors such

as valency, epitope selection, antibody binding angle, and the bulkiness of the Fc are involved in this process. (Okuya et al. , 2020a) (Callegari et al. , 2022).

Intranasal delivery of monoclonal antibodies was shown to confer viral protection in a prophylactic setting in animal models (*Haga et al. , 2021*) (*Zhou et al. , 2023*) (*Marcotte et al. , 2024*). We have shown that hamsters receiving S-IgA intranasally are protected against weight loss by SARS-CoV-2 infection. However, this treatment did not lead to a decrease in viral load in the nasal turbinates and lungs. Since the S-IgAs used in this experiment represent the general antibodies found in the nasal mucosa rather than the potent antibodies selected through screening, they lacked sufficient capacity to prevent viral infections completely. However, when these antibodies exist as polyclonal antibodies on the nasal mucosa, they may be more effective in preventing infections.

Our study has limitations, including that the number of antigen-specific plasma cells isolated from the nasal mucosa of immunized mice was insufficient for a comprehensive analysis of systemic and mucosal immune responses. In this study, antibodies that did not share V-(D)-J with nasal clones were found in the spleen, lungs, and blood of immunized mice. This finding was probably due to the limited number of clones that were isolated from the nose. However, we cannot exclude the possibility that these clones differentiated from local naïve B cells. While S-IgA has many possible advantages in antibody therapy, S-IgA has the property of aggregating. Therefore, it encounters challenges in expression and stabilization. Addressing these issues is crucial for large-scale production for *in vivo* experiments.

Materials and methods

Study approval

All experiments were performed in accordance with relevant guidelines and regulations. The Committee on Animal Experimentation at the University of Toyama approved the animal experimental protocols, which were conducted using project license A2017eng-1. All experiments involving live SARS-CoV-2 followed the approved standard operating procedures in the Biosafety Level 3 facility at the University of Toyama and Toyama Institute of Health.

Materials

The materials used in this study can be found in the Supplementary Materials. VeroE6/TMPRSS2 cells were purchased from the Japanese Collection of Research Bioresources (JCRB) Cell Bank (the National Institute of Biomedical Innovation, Health, and Nutrition, Osaka, Japan) (JCRB1819). Female ICR mice and female Syrian golden hamsters were purchased from Japan SLC, Inc. (Tokyo, Japan). Other materials are listed in Supplementary Table 2.

Immunization

Anesthetized 6-week-old female ICR mice were intranasally immunized with 10 μg of Spike$_{Wuhan}$ or PBS

with 1 μg of cholera toxin in 10 μl of PBS by using a pipette to deliver the fluid dropwise into each nostril

a total of 3 times at 3-week intervals. One week after the final immunization, *mi*ce were sacrificed by CO_2

inhalation, and blood samples were collected from the inferior vena cava. To avoid contamination with

circulating blood lymphocytes, the mice were perfused with 2 ml of PBS via the heart. Then, the lungs,

the spleen, bone marrow and Peyer's patches were dissected, and single-cell suspensions were obtained

by mincing the tissue using a 100-μm nylon mesh. The mice were decapitated at the larynx. After the

removal of the facial skin from the head, the nose was separated from the rest of the head. A pipette tip

was inserted through the pharyngeal opening into the choana, and then two consecutive volumes of 250

μl of PBS were gently perfused, and the nostril fluid was collected in a tube. The nasal lavage fluids were

centrifuged, and supernatants were stored at −80°C until assayed. The isolated nose was cut into two

along the midline, the layer of epithelium was mechanically removed from the nasal septum by gently

rubbing the sample with a needle under a stereoscopic microscope, and the tissues were then

mechanically shredded. Single-cell suspensions containing lymphocytes were isolated by using

Lympholyte-M (Cedar Lane, Ontario, Canada).

ELISA

ELISA to evaluate antibody binding to the Spike-trimer, RBDs or NTD was performed by coating Nunc

MaxiSorp flat-bottom high-binding 96-half-well plates (ThermoFisher Scientific, MS, USA) with 50 μl

per well of a 0.1 μg ml^{-1} protein solution in PBS overnight at 4□°C. Plates were washed 3 times with

PBS and incubated with 170 μl per well Blocking One solution (Nakarai, Tokyo, Japan) for 1 h at room

temperature. Serially diluted antibody, serum or nasal lavage field was added to PBST (1× PBS with

0.1% Tween-20) and incubated for 1 h at room temperature. The plates were washed three times with

PBST and then incubated with anti-mouse IgG or IgA secondary antibody conjugated to horseradish

peroxidase (HRP) (Abcam, Cambridge, UK) in PBST for 1 h at room temperature. After washing with

PBST 3 times, the plates were developed by addition of the SureBlue/TMB peroxidase substrate and stop

solution (KPL, MS, USA), and absorbance was measured at 450 nm with an ELISA microplate reader.

Protein concentration in serum and nasal lavage field is determined by BCA Protein Assay Kit (TaKaRa,

Tokyo, Japan).

ACE2 blocking assay.

Nunc Maxisorp plates were coated with SARS-CoV-2 RBD at 50 ng per well and incubated overnight at

4°C. After blocking with Blocking One solution for 1 h at room temperature, serially diluted antibody

mixed with 5 ng of biotinylated ACE2 in 100 µL of PBST was transferred to the plate in triplicate. After incubation for 1 h at room temperature, the assay plate was washed with PBST three times, and 100 µL of streptavidin-HRP (Abcam, Cambridge, UK) diluted 1:5000 in PBST was transferred to each well and incubated for 30 min. After three washes, the plate was developed with streptavidin-HRP and SureBlue/TMB peroxidase substrate.

Plasmid construction

The DNA fragments encoding the human IgA2 constant region, human J-chain and extracellular domain of human pIgR were amplified by PCR. The IgA2 constant region was replaced with the IgG1 constant region of pJON-mIgG or pETmIgA to make pJON-hIgA and pET-hIgA, respectively (*Kurosawa et al. , 2012*). The human J chain was replaced with the DsRed2 gene of pDsRedN1 (Takara Bio, Shiga, Japan). The pIgR gene was inserted into the pEF-Myc-His vector (ThermoFisher Scientific, MS, USA).

Isolation of antigen-specific plasma cells

Isolation of antigen-specific plasma cells was performed as described previously with slight modifications (*Kurosawa et al. , 2012*). Cells were stained with PE-labeled anti-mouse CD138, DyLight-650-labeled S1 and DyLight488-labeled anti-mouse IgA or anti-mouse IgG at 4℃ for 30 min with gentle agitation. After washing with PBS, the cells were suspended in PBS containing ER-Tracker Blue-White DPX (ER-tracker) and subsequently analyzed by FACS. The forward-versus-side-scatter (FSC vs. SSC) lymphocyte gate (R1) was applied to exclude dead cells. The plasma cells (IgA+ or IgG+, CD138+, ER-Tracker[high], R2 gate) were further subdivided into fractions according to their binding of DyLight-650-labeled S1 domain of SARS-CoV-2 Spike protein to define antigen-specific plasma cells (IgA+, ER-Tracker [high], S1+). Single-cell sorting was performed using a JSAN Cell Sorter that was equipped with an automatic cell deposition unit (JSAN, Kobe, Japan) with DyLight488-labeled antibodies against IgA or IgA monitored in the FL-l channel, PE-labeled CD138 in the FL-2 channel, ER-tracker in the FL-7 channel and DyLight650-labeled S1 in the FL-6 channel.

Monoclonal antibody generation

Molecular cloning of VH and VL genes from single cells was performed by 5'-RACE PCR as previously described. For the first antibody screening, the PCR-amplified VHa and VL genes were joined to pJON-hIgA and pJON-mIgK to make full-length immunoglobulin alpha heavy and kappa light chain genes by TS-jPCR, respectively (*Yoshioka et al. , 2011*). M-IgA was expressed by transfecting a pair of alpha heavy and kappa light chain genes into FreeStyle CHO-S cells that were grown in a 24-well plate according to the manufacturer's protocol (CHOgro High Yield Expression System, Takara Bio, Shiga,

Japan). For large-scale antibody production, the PCR-amplified alpha heavy and kappa light chain genes

were inserted into pET-hIgA and pET-mIgK by TS-HR, respectively (*Kurosawa et al. , 2011*).

M-IgA was expressed by cotransfecting plasmids encoding the IgA gene (5 μg) and IgK gene (5 μg) into

FreeStyle™ CHO-S cells (2.0×10^5 cells/10 ml). S-IgA was expressed by cotransfecting plasmids

encoding the IgA gene (5 μg), IgK gene (5 μg), J-chain (1 μg) and pIgR gene (1 μg) into FreeStyle™

CHO-S cells (2.0×10^5 cells/10 ml). M-IgA was purified by using Peptide M Agarose (ThermoFisher

Scientific). S-IgA was purified by two-step chromatography using the Capturem His-Taged Purification

Kit (Takara Bio, Shiga, Japan) followed by size exclusion chromatography (Cytiva, Acta go, MS, USA).

The purified antibodies were analyzed by SDS□PAGE and native polyacrylamide gel electrophoresis on

NuPAGE 4–12% Bis-Tris gels (Thermo Fisher Scientific MS, USA).

Antibody binding kinetics by SPR

The binding kinetics and affinity of monoclonal antibodies to the SARS-CoV-2 spike trimer were

analyzed by SPR (Biacore T100, GE Healthcare). Specifically, a biotinylated spike trimer was covalently

immobilized to an SA Sensor Chip for a final RU of approximately 200 and interacted with S-IgA

containing a mixture of dimers, trimers and tetramers or M-IgA at various concentrations (0.3, 1.0, 3.0,

9.0 and 27 nM of each antibody). SPR assays were run at a flow rate of 30 µl/min in HEPES buffer. The

dissociation phase was monitored for 5 minutes. The sensograms were fit in a two-component model with

BIA Evaluation software (GE Healthcare).

Neutralization activity of monoclonal antibodies against pseudotyped SARS-CoV-2 virus

20 VeroE6/TMPRSS2 cells were incubated with serially diluted antibodies and pseudotyped virus possessing

21 the spike protein of the Wuhan, Delta, Omicron strains or vesicular stomatitis virus (VSV) and cultured

22 for 48 h at 37°C. After exposure to the virus–antibody mixture, the infectivity of the pseudotyped viruses

23 was determined by measuring the luciferase activities using a PicaGene Luminescence Kit (Fujifilm

24 Wako, Osaka, Japan) with a GloMax Navigator Microplate Luminometer (Promega, WI, USA).

Neutralization assay using an authentic virus strain.

The neutralizing activity of monoclonal antibodies against an authentic Wuhan SARS-CoV-2 strain was

determined by a neutralization test in a biosafety level 3 laboratory at the Toyama Institute of Health as

previously described (*Ozawa et al. , 2022*). VeroE6/TMPRSS2 cells plated at 2×10^4 cells in each well of

96-well plates were infected with the Wuhan SARS-CoV-2 strain at a multiplicity of infection of 0.001

per cell in the presence of serially 2-fold diluted monoclonal antibodies for 1 h. After discarding the

culture supernatants, cells were cultured for 24 h with DMEM containing 10% FBS in the presence of the

indicated concentration of monoclonal antibodies. The viral infectious dose was determined by the level of viral genomic RNA in the culture supernatant, which was measured using a real-time PCR assay with a SARS-CoV-2 direct detection RT‑qPCR kit (Takara Bio, Siga, Japan). The IC_{50} was calculated by IC50 Calculator (https://www.aatbio.com/tools/ic50-calculator) and represented the neutralization titer.

Hamster models

Six-week-old female Syrian hamsters were purchased from Japan SLC Inc. (Shizuoka, Japan). The hamsters were maintained under a 12-h light–dark cycle and had unrestricted access to food and water in the Division of Animal Resources and Development at the University of Toyama. All animal procedures were approved by the Animal Experiment Committee of the University of Toyama (protocol number: A2023MED-10). Hamsters were intranasally administered 1 mg/kg of IgA in 100 μL of PBS (50 μL to each nostril) under anesthesia three hours before intranasal inoculation of 6.0×10^5 TCID50 in 100 μL of PBS (50 μL to each nostril) of SARS-CoV-2 (2019-nCoV/Japan/TY/WK-521/2020) under anesthesia. At 24 and 48 hours after SARS-CoV-2 inoculation, 1 mg/kg of IgA was administered intranasally under anesthesia. Both SARS-CoV-2 inoculation and IgA administration were performed under anesthesia with a mixture of midazolam (2.4 mg/kg) (Maruishi Pharmaceutical Co.,Ltd. Osaka, Japan), butorphanol tartrate (3 mg/kg) (Meiji, Tokyo, Japan), and medetomidine hydrochloride (0.18 mg/kg) (ZENOAQ, Fukushima, Japan), and the hamsters were awakened from anesthesia with atipamezole hydrochloride (0.18 mg/kg) (ZENOAQ) after the inoculation or administration. The hamsters were anesthetized and euthanized 72 hours after inoculation with SARS-CoV-2.

RNA extraction and real-time reverse transcription polymerase chain reaction (RT-PCR)

The nasal turbinates and lungs were collected in ISOGEN (Nippon Gene, Tokyo, Japan) and homogenized with 1.4 mm Ceramic Beads (Thermo Fisher Scientific, Hampton, NH, USA) using BeadMill 24 (Thermo Fisher Scientific) at a speed of 4 m/s for 3 min and stored at −80 °C until subsequent use. Total RNA was extracted using ISOGEN according to the manufacturer's instructions. Real-time RT-PCR was performed by amplifying the SARS-CoV-2 N2 gene using QuantStudio 6 Pro Real-Time PCR System (Thermo Fisher Scientific) and THUNDERBIRD Probe One-step qRT-PCR Kit (TOYOBO, Osaka, Japan) with the N2 primer/probe set (Nihon Gene Research Laboratories, Miyagi, Japan). The absolute copy number of the SARS-CoV-2 N2 gene was determined by serially diluted RNA control (Nihon Gene Research Laboratories).

Gene family and phylogenetic analysis of monoclonal antibodies

The antibody sequences were annotated against the IMGT mouse heavy and light chain gene database

using NCBI IgBlast to determine IGHV, IGHD, IGLV, IGHJ, IGLV and IGLJ gene annotations. Antibody clones were assigned to clonal groups using Sequencher software. The heavy and light chain variable gene arrangement and phylogenetic analyses were performed using MAFFT, a multiple alignment program, in the GENETYX sequence analysis package (https://www.genetyx.co.jp). For somatic hypermutation, IGHV and IGLV sequences were aligned against representative clones of each group. Antibody clones consisting of pairs of heavy and light chain variable genes (signal sequence to FW4) were used to generate an antibody phylogenetic tree. Full-length germline sequences were reconstructed, with nucleotide additions/deletions in the junction between V-(D)-J adjusted to match the sequence of each antibody group. Within these groups, if the combined sequence of the heavy and light chains differed by more than five bases from each other, they were defined as separate clones. The sequence alignment tool Clustal Omega (https://www.ebi.ac.uk/Tools/msa/clustalo/) was used to identify deletions and insertions and to align the length of the sequence. The nucleotide lengths were aligned by adding "-" to the missing bases. The data file required by Alakazam was created in RStudio (version 2022.12.0+353), and a phylogenetic tree was created by Alakazam (https://www.rdocumentation.org/packages/alakazam/versions/1.2.1 and https://alakazam.readthedocs.io/en/stable/). The R script was as follows: https://www.rdocumentation.org/packages/alakazam/versions/1.2.1/topics/buil dPhylipLineage

Statistical analysis

All statistical analyses were performed using JMP statistical software (JMP Statistical Discovery, NC, USA). Unpaired Student's t tests were used to analyze each dataset. The threshold for statistical significance was set at $p < 0.01$ (**).

Data availability

The nucleotide sequence data were submitted to the DDBJ/EMBL/GenBank databases (Accession No. LC761310-LC761361, LC761441-LC761570, LC761965-LC762059, LC764607-LC764701, LC761389-LC761570). Other relevant data are available from the corresponding authors upon request.

Acknowledgments

We thank past and current members of our laboratory for fruitful discussions. We also thank M. Nozaki and K. Takai for technical support.

References

Afkhami S, D'Agostino MR, Zhang A, Stacey HD, Marzok A, Kang A, Singh R, Bavananthasivam J, Ye
G, Luo X, Wang F, Ang JC, Zganiacz A, Sankar U, Kazhdan N, Koenig JFE, Phelps A, Gameiro
SF, Tang S, Jordana M, Wan Y, Mossman KL, Jeyanathan M, Gillgrass A, Medina MFC, Smaill
F, Lichty BD, Miller MS, Xing Z. 2022. Respiratory mucosal delivery of next-generation
COVID-19 vaccine provides robust protection against both ancestral and variant strains of
SARS-CoV-2. *Cell* **185**:896-915 e819. DOI: 10.1016/j.cell.2022.02.005, PMID: PMC8825346

Alu A, Chen L, Lei H, Wei Y, Tian X, Wei X. 2022. Intranasal COVID-19 vaccines: From bench to bed.
EBioMedicine **76**:103841. DOI: 10.1016/j.ebiom.2022.103841, PMID: PMC8785603

Asahi Y, Yoshikawa T, Watanabe I, Iwasaki T, Hasegawa H, Sato Y, Shimada S, Nanno M, Matsuoka Y,
Ohwaki M, Iwakura Y, Suzuki Y, Aizawa C, Sata T, Kurata T, Tamura S. 2002. Protection
against influenza virus infection in polymeric Ig receptor knockout mice immunized intranasally
with   adjuvant-combined   vaccines.   *J   Immunol*   **168**:2930-2938.   DOI:
10.4049/jimmunol.168.6.2930

Asahi-Ozaki Y, Yoshikawa T, Iwakura Y, Suzuki Y, Tamura S, Kurata T, Sata T. 2004. Secretory IgA
antibodies provide cross-protection against infection with different strains of influenza B virus. *J
Med Virol* **74**:328-335. DOI: 10.1002/jmv.20173

Azzi L, Dalla Gasperina D, Veronesi G, Shallak M, Ietto G, Iovino D, Baj A, Gianfagna F, Maurino V,
Focosi D, Maggi F, Ferrario MM, Dentali F, Carcano G, Tagliabue A, Maffioli LS, Accolla RS,
Forlani G. 2022. Mucosal immune response in BNT162b2 COVID-19 vaccine recipients.
EBioMedicine **75**:103788. DOI: 10.1016/j.ebiom.2021.103788, PMID: PMC8718969

Barrett JR, Belij-Rammerstorfer S, Dold C, Ewer KJ, Folegatti PM, Gilbride C, Halkerston R, Hill J,
Jenkin D, Stockdale L, Verheul MK, Aley PK, Angus B, Bellamy D, Berrie E, Bibi S, Bittaye M,
Carroll MW, Cavell B, Clutterbuck EA, Edwards N, Flaxman A, Fuskova M, Gorringe A, Hallis
B, Kerridge S, Lawrie AM, Linder A, Liu X, Madhavan M, Makinson R, Mellors J, Minassian A,
Moore M, Mujadidi Y, Plested E, Poulton I, Ramasamy MN, Robinson H, Rollier CS, Song R,
Snape MD, Tarrant R, Taylor S, Thomas KM, Voysey M, Watson MEE, Wright D, Douglas AD,
Green CM, Hill AVS, Lambe T, Gilbert S, Pollard AJ, Oxford CVTG. 2021. Phase 1/2 trial of
SARS-CoV-2 vaccine ChAdOx1 nCoV-19 with a booster dose induces multifunctional antibody
responses. *Nat Med* **27**:279-288. DOI: 10.1038/s41591-020-01179-4

Bricker TL, Darling TL, Hassan AO, Harastani HH, Soung A, Jiang XP, Dai YN, Zhao HY, Adams LJ,
Holtzman MJ, Bailey AL, Case JB, Fremont DH, Klein R, Diamond MS, Boon ACM. 2021. A
single intranasal or intramuscular immunization with chimpanzee adenovirus-vectored
SARS-CoV-2 vaccine protects against pneumonia in hamsters. *Cell Reports* **36**. DOI: ARTN
109400

10.1016/j.celrep.2021.109400

Budinger GRS, Misharin AV, Ridge KM, Singer BD, Wunderink RG. 2021. Distinctive features of
severe SARS-CoV-2 pneumonia. *J Clin Invest* **131**. DOI: 10.1172/JCI149412, PMID:
PMC8279580

Callegari I, Schneider M, Berloffa G, Muhlethaler T, Holdermann S, Galli E, Roloff T, Boss R, Infanti L,
Khanna N, Egli A, Buser A, Zimmer G, Derfuss T, Sanderson NSR. 2022. Potent neutralization
by monoclonal human IgM against SARS-CoV-2 is impaired by class switch. *EMBO Rep*
23:e53956. DOI: 10.15252/embr.202153956, PMID: PMC9253785

Chavda VP, Vora LK, Pandya AK, Patravale VB. 2021. Intranasal vaccines for SARS-CoV-2: From
challenges to potential in COVID-19 management. *Drug Discov Today* **26**:2619-2636. DOI:
10.1016/j.drudis.2021.07.021, PMID: PMC8319039

D'Arco C, McCormick AA, Arnaboldi PM. 2021. Single-dose intranasal subunit vaccine rapidly clears
secondary sepsis in a high-dose pneumonic plague infection. *Vaccine* **39**:1435-1444. DOI:
10.1016/j.vaccine.2021.01.040

DeFrancesco L. 2020. COVID-19 antibodies on trial. *Nat Biotechnol* **38**:1242-1252. DOI:
10.1038/s41587-020-0732-8, PMID: PMC7576980

Dhakal S, Renu S, Ghimire S, Shaan Lakshmanappa Y, Hogshead BT, Feliciano-Ruiz N, Lu F,
HogenEsch H, Krakowka S, Lee CW, Renukaradhya GJ. 2018. Mucosal Immunity and
Protective Efficacy of Intranasal Inactivated Influenza Vaccine Is Improved by Chitosan
Nanoparticle Delivery in Pigs. *Front Immunol* **9**:934. DOI: 10.3389/fimmu.2018.00934, PMID:
PMC5940749

Ejemel M, Li Q, Hou S, Schiller ZA, Tree JA, Wallace A, Amcheslavsky A, Kurt Yilmaz N, Buttigieg
KR, Elmore MJ, Godwin K, Coombes N, Toomey JR, Schneider R, Ramchetty AS, Close BJ,
Chen DY, Conway HL, Saeed M, Ganesa C, Carroll MW, Cavacini LA, Klempner MS, Schiffer
CA, Wang Y. 2020. A cross-reactive human IgA monoclonal antibody blocks SARS-CoV-2

spike-ACE2 interaction. *Nat Commun* **11**:4198. DOI: 10.1038/s41467-020-18058-8, PMID: PMC7442812

Galimidi RP, Klein JS, Politzer MS, Bai S, Seaman MS, Nussenzweig MC, West AP, Jr., Bjorkman PJ. 2015. Intra-spike crosslinking overcomes antibody evasion by HIV-1. *Cell* **160**:433-446. DOI: 10.1016/j.cell.2015.01.016, PMID: PMC4401576

Gianchecchi E, Manenti A, Kistner O, Trombetta C, Manini I, Montomoli E. 2019. How to assess the effectiveness of nasal influenza vaccines? Role and measurement of sIgA in mucosal secretions. *Influenza and Other Respiratory Viruses* **13**:429-437. DOI: 10.1111/irv.12664

Houston S. 2023. SARS-CoV-2 mucosal vaccine. *Nat Immunol* **24**:1. DOI: 10.1038/s41590-022-01405-w

Jackson CB, Farzan M, Chen B, Choe H. 2022. Mechanisms of SARS-CoV-2 entry into cells. *Nat Rev Mol Cell Biol* **23**:3-20. DOI: 10.1038/s41580-021-00418-x, PMID: PMC8491763

Kim E, Weisel FJ, Balmert SC, Khan MS, Huang SH, Erdos G, Kenniston TW, Carey CD, Joachim SM, Conter LJ, Weisel NM, Okba NMA, Haagmans BL, Percivalle E, Cassaniti I, Baldanti F, Korkmaz E, Shlomchik MJ, Falo LD, Gambotto A. 2021. A single subcutaneous or intranasal immunization with adenovirus-based SARS-CoV-2 vaccine induces robust humoral and cellular immune responses in mice. *European Journal of Immunology* **51**:1774-1784. DOI: 10.1002/eji.202149167

Klein JS, Bjorkman PJ. 2010. Few and far between: how HIV may be evading antibody avidity. *PLoS Pathog* **6**:e1000908. DOI: 10.1371/journal.ppat.1000908, PMID: PMC2877745

Kurosawa N, Yoshioka M, Fujimoto R, Yamagishi F, Isobe M. 2012. Rapid production of antigen-specific monoclonal antibodies from a variety of animals. *Bmc Biology* **10**. DOI: Artn 80

10.1186/1741-7007-10-80

Kurosawa N, Yoshioka M, Isobe M. 2011. Target-selective homologous recombination cloning for high-throughput generation of monoclonal antibodies from single plasma cells. *Bmc Biotechnology* **11**. DOI: Artn 39

10.1186/1472-6750-11-39

Lapuente D, Fuchs J, Willar J, Vieira Antao A, Eberlein V, Uhlig N, Issmail L, Schmidt A, Oltmanns F, Peter AS, Mueller-Schmucker S, Irrgang P, Fraedrich K, Cara A, Hoffmann M, Pohlmann S,

1   Ensser A, Pertl C, Willert T, Thirion C, Grunwald T, Uberla K, Tenbusch M. 2021. Protective
2   mucosal immunity against SARS-CoV-2 after heterologous systemic prime-mucosal boost
3   immunization. *Nat Commun* **12**:6871. DOI: 10.1038/s41467-021-27063-4, PMID: PMC8626513

4   Li Y, Jin L, Chen T. 2020. The Effects of Secretory IgA in the Mucosal Immune System. *Biomed Res Int*
5   **2020**:2032057. DOI: 10.1155/2020/2032057, PMID: PMC6970489

6   Maltseva M, Galipeau Y, Renner TM, Deschatelets L, Durocher Y, Akache B, Langlois MA. 2022.
7   Characterization of Systemic and Mucosal Humoral Immune Responses to an Adjuvanted
8   Intranasal SARS-CoV-2 Protein Subunit Vaccine Candidate in Mice. *Vaccines (Basel)* **11**. DOI:
9   10.3390/vaccines11010030, PMID: PMC9865305

Matsumoto ML. 2022. Molecular Mechanisms of Multimeric Assembly of IgM and IgA. *Annu Rev*
Immunol. DOI: 10.1146/annurev-immunol-101320-123742

Nicasio M, Sautto G, Clementi N, Diotti RA, Criscuolo E, Castelli M, Solforosi L, Clementi M, Burioni
R. 2012. Neutralization interfering antibodies: a "novel" example of humoral immune
dysfunction facilitating viral escape? *Viruses* **4**:1731-1752. DOI: 10.3390/v4091731, PMID:
PMC3499828

Ohtsuka J, Imai M, Fukumura M, Maeda M, Eguchi A, Ono R, Maemura T, Ito M, Yamayoshi S,
Kataoka Y, Kawaoka Y, Nosaka T. 2021. Non-propagative human parainfluenza virus type 2
nasal vaccine robustly protects the upper and lower airways against SARS-CoV-2. *iScience*
24:103379. DOI: 10.1016/j.isci.2021.103379, PMID: PMC8596570

Okuya K, Eguchi N, Manzoor R, Yoshida R, Saito S, Suzuki T, Sasaki M, Saito T, Kida Y,
Mori-Kajihara A, Miyamoto H, Ichii O, Kajihara M, Higashi H, Takada A. 2020a. Comparative
Analyses of the Antiviral Activities of IgG and IgA Antibodies to Influenza A Virus M2 Protein.
Viruses **12**. DOI: 10.3390/v12070780, PMID: PMC7411592

Okuya K, Yoshida R, Manzoor R, Saito S, Suzuki T, Sasaki M, Saito T, Kida Y, Mori-Kajihara A,
Kondoh T, Sato M, Kajihara M, Miyamoto H, Ichii O, Higashi H, Takada A. 2020b. Potential
Role of Nonneutralizing IgA Antibodies in Cross-Protective Immunity against Influenza A
Viruses of Multiple Hemagglutinin Subtypes. *Journal of Virology* **94**. DOI: ARTN e00408-20

10.1128/JVI.00408-20

Ozawa T, Tani H, Anraku Y, Kita S, Igarashi E, Saga Y, Inasaki N, Kawasuji H, Yamada H, Sasaki SI,
Somekawa M, Sasaki J, Hayakawa Y, Yamamoto Y, Morinaga Y, Kurosawa N, Isobe M,
Fukuhara H, Maenaka K, Hashiguchi T, Kishi H, Kitajima I, Saito S, Niimi H. 2022. Novel
super-neutralizing antibody UT28K is capable of protecting against infection from a wide
variety of SARS-CoV-2 variants. *MAbs* **14**:2072455. DOI: 10.1080/19420862.2022.2072455,
PMID: PMC9103358

Polack FP, Thomas SJ, Kitchin N, Absalon J, Gurtman A, Lockhart S, Perez JL, Perez Marc G, Moreira
ED, Zerbini C, Bailey R, Swanson KA, Roychoudhury S, Koury K, Li P, Kalina WV, Cooper D,
Frenck RW, Jr., Hammitt LL, Tureci O, Nell H, Schaefer A, Unal S, Tresnan DB, Mather S,
Dormitzer PR, Sahin U, Jansen KU, Gruber WC, Group CCT. 2020. Safety and Efficacy of the
BNT162b2 mRNA Covid-19 Vaccine. *N Engl J Med* **383**:2603-2615. DOI:
10.1056/NEJMoa2034577, PMID: PMC7745181

Saito S, Sano K, Suzuki T, Ainai A, Taga Y, Ueno T, Tabata K, Saito K, Wada Y, Ohara Y, Takeyama H,
Odagiri T, Kageyama T, Ogawa-Goto K, Multihartina P, Setiawaty V, Pangesti KNA, Hasegawa
H. 2019. IgA tetramerization improves target breadth but not peak potency of functionality of
anti-influenza virus broadly neutralizing antibody. *PLoS Pathog* **15**:e1007427. DOI:
10.1371/journal.ppat.1007427, PMID: PMC6317788

Sheikh-Mohamed S, Isho B, Chao GYC, Zuo M, Cohen C, Lustig Y, Nahass GR, Salomon-Shulman RE,
Blacker G, Fazel-Zarandi M, Rathod B, Colwill K, Jamal A, Li Z, de Launay KQ, Takaoka A,
Garnham-Takaoka J, Patel A, Fahim C, Paterson A, Li AX, Haq N, Barati S, Gilbert L, Green K,
Mozafarihashjin M, Samaan P, Budylowski P, Siqueira WL, Mubareka S, Ostrowski M, Rini JM,
Rojas OL, Weissman IL, Tal MC, McGeer A, Regev-Yochay G, Straus S, Gingras AC,
Gommerman JL. 2022. Systemic and mucosal IgA responses are variably induced in response to
SARS-CoV-2 mRNA vaccination and are associated with protection against subsequent
infection. *Mucosal Immunol* **15**:799-808. DOI: 10.1038/s41385-022-00511-0, PMID:
PMC9037584

Shimoda M, Nakamura T, Takahashi Y, Asanuma H, Tamura S, Kurata T, Mizuochi T, Azuma N, Kanno
C, Takemori T. 2001. Isotype-specific selection of high affinity memory B cells in
nasal-associated lymphoid tissue. *Journal of Experimental Medicine* **194**:1597-1607. DOI: DOI
10.1084/jem.194.11.1597

Sterlin D, Malaussena A, Gorochov G. 2021. IgA dominates the early neutralizing antibody response to
SARS-CoV-2 virus. *M S-Medecine Sciences* **37**:968-970. DOI: 10.1051/medsci/2021154

Sui Y, Li J, Zhang R, Prabhu SK, Andersen H, Venzon D, Cook A, Brown R, Teow E, Velasco J, Greenhouse J, Putman-Taylor T, Campbell TA, Pessaint L, Moore IN, Lagenaur L, Talton J, Breed MW, Kramer J, Bock KW, Minai M, Nagata BM, Lewis MG, Wang LX, Berzofsky JA. 2021. Protection against SARS-CoV-2 infection by a mucosal vaccine in rhesus macaques. *JCI Insight* **6**. DOI: 10.1172/jci.insight.148494, PMID: PMC8262352

Sungnak W, Huang N, Becavin C, Berg M, Queen R, Litvinukova M, Talavera-Lopez C, Maatz H, Reichart D, Sampaziotis F, Worlock KB, Yoshida M, Barnes JL, Network HCALB. 2020. SARS-CoV-2 entry factors are highly expressed in nasal epithelial cells together with innate immune genes. *Nat Med* **26**:681-687. DOI: 10.1038/s41591-020-0868-6, PMID: PMC8637938

Tan GS, Leon PE, Albrecht RA, Margine I, Hirsh A, Bahl J, Krammer F. 2016. Broadly-Reactive Neutralizing and Non-neutralizing Antibodies Directed against the H7 Influenza Virus Hemagglutinin Reveal Divergent Mechanisms of Protection. *PLoS Pathog* **12**:e1005578. DOI: 10.1371/journal.ppat.1005578, PMID: PMC4833315

Tumpey TM, Renshaw M, Clements JD, Katz JM. 2001. Mucosal delivery of inactivated influenza vaccine induces B-cell-dependent heterosubtypic cross-protection against lethal influenza A H5N1 virus infection. *J Virol* **75**:5141-5150. DOI: 10.1128/JVI.75.11.5141-5150.2001, PMID: PMC114919

Vabret N, Britton GJ, Gruber C, Hegde S, Kim J, Kuksin M, Levantovsky R, Malle L, Moreira A, Park MD, Pia L, Risson E, Saffern M, Salome B, Esai Selvan M, Spindler MP, Tan J, van der Heide V, Gregory JK, Alexandropoulos K, Bhardwaj N, Brown BD, Greenbaum B, Gumus ZH, Homann D, Horowitz A, Kamphorst AO, Curotto de Lafaille MA, Mehandru S, Merad M, Samstein RM, Sinai Immunology Review P. 2020. Immunology of COVID-19: Current State of the Science. *Immunity* **52**:910-941. DOI: 10.1016/j.immuni.2020.05.002, PMID: PMC7200337

Wong TY, Lee KS, Russ BP, Horspool AM, Kang J, Winters MT, Allison Wolf M, Rader NA, Miller OA, Shiflett M, Izac J, Varisco D, Sen-Kilic E, Cunningham C, Cooper M, Cyphert HA, Barbier M, Martinez I, Bevere JR, Ernst RK, Damron FH. 2022. Intranasal administration of BReC-CoV-2 COVID-19 vaccine protects K18-hACE2 mice against lethal SARS-CoV-2 challenge. *NPJ Vaccines* **7**:36. DOI: 10.1038/s41541-022-00451-7

Woof JM, Kerr MA. 2006. The function of immunoglobulin A in immunity. *J Pathol* **208**:270-282. DOI: 10.1002/path.1877

Yoshioka M, Kurosawa N, Isobe M. 2011. Target-selective joint polymerase chain reaction: A robust and rapid method for high-throughput production of recombinant monoclonal antibodies from single cells. *Bmc Biotechnology* **11**. DOI: 10.1186/1472-6750-11-75

Ejemel M, Li Q, Hou S, Schiller ZA, Tree JA, Wallace A, Amcheslavsky A, Kurt Yilmaz N, Buttigieg KR, Elmore MJ, Godwin K, Coombes N, Toomey JR, Schneider R, Ramchetty AS, Close BJ, Chen DY, Conway HL, Saeed M, Ganesa C, Carroll MW, Cavacini LA, Klempner MS, Schiffer CA, Wang Y. 2020. A cross-reactive human IgA monoclonal antibody blocks SARS-CoV-2 spike-ACE2 interaction. *Nat Commun* **11**:4198. DOI: 10.1038/s41467-020-18058-8, PMID: PMC7442812

Ewer KJ, Barrett JR, Belij-Rammerstorfer S, Sharpe H, Makinson R, Morter R, Flaxman A, Wright D, Bellamy D, Bittaye M, Dold C, Provine NM, Aboagye J, Fowler J, Silk SE, Alderson J, Aley PK, Angus B, Berrie E, Bibi S, Cicconi P, Clutterbuck EA, Chelysheva I, Folegatti PM, Fuskova M, Green CM, Jenkin D, Kerridge S, Lawrie A, Minassian AM, Moore M, Mujadidi Y, Plested E, Poulton I, Ramasamy MN, Robinson H, Song R, Snape MD, Tarrant R, Voysey M, Watson MEE, Douglas AD, Hill AVS, Gilbert SC, Pollard AJ, Lambe T, Oxford CVTG. 2021. T cell and antibody responses induced by a single dose of ChAdOx1 nCoV-19 (AZD1222) vaccine in a phase 1/2 clinical trial. *Nat Med* **27**:270-278. DOI: 10.1038/s41591-020-01194-5

Haga K, Takai-Todaka R, Matsumura Y, Song C, Takano T, Tojo T, Nagami A, Ishida Y, Masaki H, Tsuchiya M, Ebisudani T, Sugimoto S, Sato T, Yasuda H, Fukunaga K, Sawada A, Nemoto N, Murata K, Morimoto T, Katayama K. 2021. Nasal delivery of single-domain antibody improves symptoms of SARS-CoV-2 infection in an animal model. *PLoS Pathog* **17**:e1009542. DOI: 10.1371/journal.ppat.1009542, PMID: PMC8516304

Marcotte H, Cao Y, Zuo F, Simonelli L, Sammartino JC, Pedotti M, Sun R, Cassaniti I, Hagbom M, Piralla A, Yang J, Du L, Percivalle E, Bertoglio F, Schubert M, Abolhassani H, Sherina N, Guerra C, Borte S, Rezaei N, Kumagai-Braesch M, Xue Y, Su C, Yan Q, He P, Gronwall C, Klareskog L, Calzolai L, Cavalli A, Wang Q, Robbiani DF, Hust M, Shi Z, Feng L, Svensson L, Chen L, Bao L, Baldanti F, Xiao J, Qin C, Hammarstrom L, Yang X, Varani L, Xie XS, Pan-Hammarstrom Q. 2024. Conversion of monoclonal IgG to dimeric and secretory IgA restores neutralizing ability and prevents infection of Omicron lineages. *Proc Natl Acad Sci U S A* **121**:e2315354120. DOI: 10.1073/pnas.2315354120, PMID: PMC10801922

Muramatsu M, Yoshida R, Yokoyama A, Miyamoto H, Kajihara M, Maruyama J, Nao N, Manzoor R,
Takada A. 2014. Comparison of antiviral activity between IgA and IgG specific to influenza
virus hemagglutinin: increased potential of IgA for heterosubtypic immunity. *PLoS One*
9:e85582. DOI: 10.1371/journal.pone.0085582, PMID: PMC3895000

Saito S, Sano K, Suzuki T, Ainai A, Taga Y, Ueno T, Tabata K, Saito K, Wada Y, Ohara Y, Takeyama H,
Odagiri T, Kageyama T, Ogawa-Goto K, Multihartina P, Setiawaty V, Pangesti KNA, Hasegawa
H. 2019. IgA tetramerization improves target breadth but not peak potency of functionality of
anti-influenza virus broadly neutralizing antibody. *PLoS Pathog* **15**:e1007427. DOI:
10.1371/journal.ppat.1007427, PMID: PMC6317788

Tudor D, Yu H, Maupetit J, Drillet AS, Bouceba T, Schwartz-Cornil I, Lopalco L, Tuffery P, Bomsel M.
2012. Isotype modulates epitope specificity, affinity, and antiviral activities of anti-HIV-1
human broadly neutralizing 2F5 antibody. *Proc Natl Acad Sci U S A* **109**:12680-12685. DOI:
10.1073/pnas.1200024109, PMID: PMC3412007

Wang Z, Lorenzi JCC, Muecksch F, Finkin S, Viant C, Gaebler C, Cipolla M, Hoffmann HH, Oliveira
TY, Oren DA, Ramos V, Nogueira L, Michailidis E, Robbiani DF, Gazumyan A, Rice CM,
Hatziioannou T, Bieniasz PD, Caskey M, Nussenzweig MC. 2021. Enhanced SARS-CoV-2
neutralization by dimeric IgA. *Sci Transl Med* **13**. DOI: 10.1126/scitranslmed.abf1555, PMID:
PMC7857415

Zhou B, Zhou R, Chan JF, Zeng J, Zhang Q, Yuan S, Liu L, Robinot R, Shan S, Liu N, Ge J, Kwong HY,
Zhou D, Xu H, Chan CC, Poon VK, Chu H, Yue M, Kwan KY, Chan CY, Chan CC, Chik KK,
Du Z, Au KK, Huang H, Man HO, Cao J, Li C, Wang Z, Zhou J, Song Y, Yeung ML, To KK,
Ho DD, Chakrabarti LA, Wang X, Zhang L, Yuen KY, Chen Z. 2023. SARS-CoV-2 hijacks
neutralizing dimeric IgA for nasal infection and injury in Syrian hamsters(1). *Emerg Microbes*
Infect **12**:2245921. DOI: 10.1080/22221751.2023.2245921, PMID: PMC10444022

Zhu F, Huang S, Liu X, Chen Q, Zhuang C, Zhao H, Han J, Jaen AM, Do TH, Peter JG, Dorado AG,
Tirador LS, Zabat GMA, Villalobos REM, Gueco GP, Botha LLG, Iglesias Pertuz SP, Tan J,
Zhu K, Quan J, Lin H, Huang Y, Jia J, Chu X, Chen J, Chen Y, Zhang T, Su Y, Li C, Ye X, Wu
T, Zhang J, Xia N, Team C-P-S. 2023. Safety and efficacy of the intranasal spray SARS-CoV-2
vaccine dNS1-RBD: a multicentre, randomised, double-blind, placebo-controlled, phase 3 trial.
Lancet Respir Med **11**:1075-1088. DOI: 10.1016/S2213-2600(23)00349-1, PMID:
PMC10682370