Comparative Clustering (CompaCt) of eukaryote complexomes identifies novel interactions and sheds light on protein complex evolution

Joeri van Strien[1*], Felix Evers[2], Madhurya Lutikurti[3], Stijn L. Berendsen[3], Alejandro Garanto[3,4,5], Geert-Jan van Gemert[2], Alfredo Cabrera-Orefice[1], Richard J. Rodenburg[5,6], Ulrich Brandt[3,5,7], Taco W.A. Kooij[2], Martijn A. Huynen[1*]

[1] Medical BioSciences Department, Radboud University Medical Center, Nijmegen, the Netherlands

[2] Medical Microbiology, Radboud Center for Infectious Diseases, Radboud University Medical Center, Nijmegen, The Netherlands.

[3] Department of Pediatrics, Amalia Children's Hospital, Radboud University Medical Center, Nijmegen, the Netherlands

[4] Department of Human Genetics, Radboud University Medical Center, Nijmegen, The Netherlands

[5] Radboud Center for Mitochondrial Medicine (RCMM), Radboud University Medical Center, Nijmegen, the Netherlands

[6] Department of Pediatrics, Translational Metabolic Laboratory, Radboud University Medical Center, Nijmegen, the Netherlands

[7] Cologne Excellence Cluster on Cellular Stress Responses in Aging-Associated Diseases (CECAD), University of Cologne, 50931 Cologne, Germany

* Corresponding Authors

E-mail: joeri.vanStrien@radboudumc.nl, Martijn.Huijnen@radboudumc.nl

Abstract

Complexome profiling allows large-scale, untargeted, and comprehensive characterization of protein complexes in a biological sample using a combined approach of separating intact protein complexes e.g., by native gel electrophoresis, followed by mass spectrometric analysis of the proteins in the resulting fractions. Over the last decade, its application has resulted in a large collection of complexome profiling datasets. While computational methods have been developed for the analysis of individual datasets, methods for large-scale comparative analysis of complexomes from multiple species are lacking. Here, we present Comparative Clustering (CompaCt), that performs fully automated integrative analysis of complexome profiling data from multiple species, enabling systematic characterization and comparison of complexomes. CompaCt implements a novel method for leveraging orthology in comparative analysis to allow systematic identification of conserved as well as taxon-specific elements of the analyzed complexomes. We applied this method to a collection of 53 complexome profiles spanning the major branches of the eukaryotes. We demonstrate the ability of CompaCt to robustly identify the composition of protein complexes, and show that integrated analysis of multiple datasets improves characterization of complexes from specific complexome profiles when compared to separate analysis. We identified novel candidate interactors and complexes in a number of species from previously analyzed datasets, like the emp24, the V-ATPase and mitochondrial ATP synthase complexes. Lastly, we demonstrate the utility of CompaCt for the automated large-scale characterization of the complexome of the mosquito *Anopheles stephensi* shedding light on the evolution of metazoan protein complexes. CompaCt is available from https://github.com/cmbi/compact-bio.

Author summary

Proteins carry out essential functions in the majority of processes in life, often by binding with other proteins to form multiprotein complexes. State of the art experimental techniques

such as complexome profiling enable large-scale identification of protein complexes in a biological sample. With the increase in use of this method in recent years these experiments have been performed on a variety of species, of which the results are publicly available. Combining the results from these experiments presents a computational challenge, but could identify novel protein complexes and provide insights into their evolution. Here, we introduce CompaCt as a method to integrate complexome profiles from multiple species enabling automatic large-scale characterization of protein complexes. It identifies commonalities as well as the differences between species. By applying CompaCt to a collection of complexome profiles, we identified candidate complexes and interacting proteins in a number of species that were not detected in previous separate analyses of these datasets. In doing so we shed light on the evolutionary origin of several protein complex members, pinpointed the function of biomedically relevant proteins, whose role was previously unknown, and performed the first investigation of the *Anopheles stephensi* complexome, a mosquito that transmits the malaria parasite.

Introduction

Most biological processes are mediated by proteins, which in many cases need to form multiprotein complexes to carry out their function. The complexome is the complete set of multiprotein complexes present in a biological system, be it an entire organism or a more limited subsystem such as a specific tissue, cell type, organelle or life stage. Complexome profiling, an 'omics approach aimed at large-scale identification of protein complexes in a single experiment (2) has in the last decade enabled untargeted and systematic analysis of complexomes, reviewed in (1). In a complexome profiling experiment, native protein complexes are systematically separated across a number of fractions that are then separately analyzed by tandem mass spectrometry to assess their content in quantitative manner. Proteins contained in the same complex or subassembly typically co-migrate and exhibit similar abundance profiles across all or a subset of the fractions. This approach

allows interrogation of the composition of a large number of protein complexes as well as of

their assembly in a given biological sample in an untargeted manner.

In recent years several computational approaches have been developed to infer protein

complexes from these data. Some of these rely solely on the complexome profiling data (3–

5), while others use a reference of known protein complexes (6–8) or integrate additional

types of interaction evidence to improve protein complex identification (7,9,10). Additionally,

several tools allow comparison of multiple complexome profiles from one species to examine

the effect of certain mutations or conditions on the composition or assembly of the

investigated complexome (8,11,12).

Since the initial application of the complexome profiling approach, an increasing number of

complexome profiling datasets have been generated from at least 21 different species,

covering all kingdoms of life (1), which are available from the CEDAR database of

complexome profiles (13). Leveraging these data in comparative analyses in principle has

the potential for more sensitive detection of novel interactors and to facilitate analyses of the

evolution of protein complexes. Nevertheless, comparative analysis of multiple complexome

profiles poses a methodological challenge. Firstly, available datasets have been generated

by different laboratories, using different experimental protocols and resolutions for the

separation of protein complexes. This renders direct comparison of the "raw" protein

migration profile data difficult. Secondly, while integrating complexome profiles from related

species has helped to identify evolutionarily conserved protein interactions consistently

present in multiple species (9,10), these approaches preclude characterization of species-

specific interactors. Large-scale identification of not just conserved, but also taxon-specific

complex members and interactions would result in a more complete picture of the

investigated complexomes, and would facilitate evolutionary analyses of protein complexes.

To comparatively analyze the protein complexes in multiple complexome profiling datasets, we developed Comparative Clustering (CompaCt), a computational approach that performs fully automated large-scale integration of protein interaction data from multiple species. Our method of utilizing orthology relations allows combined analysis of multiple species in a manner that enables identification of conserved as well as taxon-specific protein complexes and interactions in a single, uniform analysis platform. Additionally, by including multiple interaction datasets per analyzed species CompaCt is able to distinguish consistently co-migrating proteins representing true interactors from spuriously co-migrating proteins that do not interact. Therewith, it improves the reliability of the analysis of complexes from the individual species included in the comparative analysis.

We have applied our tool to a collection of 53 complexome profiling datasets from nine different species covering the major eukaryotic branches. We demonstrate that integrative analysis of these data with CompaCt is able to systematically recover known conserved and taxon-specific interactions and pinpoint novel interactors and complexes missed in previous analyses. Moreover, by presenting the first analysis of the complexome of the malaria transducing mosquito *A. stephensi*, we demonstrate the usefulness of CompaCt for performing large scale and automatic identification of protein complexes to increase our understanding of the evolution of eukaryotic protein complexes.

Results

CompaCt: Comparative Clustering of interaction datasets

Our comparative clustering (CompaCt) approach performs automated comparative analysis of protein interaction data representing a number of complexomes (i.e., the set of complexes present in a species, cell type, etc.), and enables systematic identification of not just conserved, but also of taxon-specific interactions. To this end, rather than asking whether

interactions between orthologous proteins are conserved, which precludes analysis of

proteins without known orthologs, we ask whether any two proteins from different species

interact with the same set of orthologous proteins. Thus, even though proteomes of different

species will only partly overlap, we are still able to estimate whether two proteins are part of

the same complex by identifying orthologs among their interactors. An overview of the

CompaCt workflow is shown in Figure 1.

correlation datasets   comparing interactor profiles   clustering combined network   processing cluster results

Figure 1: Overview of CompaCt analysis steps. A. Input correlation datasets from multiple biological

systems, with lines indicating correlations between proteins based on the similarity in their migration

profiles in the complexome profiling data and colors indicating proteins from different datasets. B.

Determining interactor profile similarity of two proteins, I and II, from different datasets, by calculating

the overlap between their interactors (A, B, C, etc.), using orthology (A-A', C-C') when they are from

different species, with the rank biased overlap metric (14). C. Clustering the combined network of all

proteins (nodes) with RBO scores (edges) into clusters with MCL (15). D. Processing of MCL clusters,

separating into subclusters per system (e.g., from the same tissue or species), while pooling

information of datasets representing the same system.

To estimate whether two proteins have common interactors, we compare their sets of

interactions that are ranked based on scores representing interaction likelihood inferred from

experimental interaction data. We refer to these ranked sets as their interactor profile. To be

able to quantify the similarity between interactor profiles, we use the rank biased overlap

(RBO) metric (14). This metric determines the similarity between ranked lists, while allowing

lists of different lengths with non-common elements. Applying this metric to compare

interactor profiles we can determine whether any two proteins from different proteomes have

common, orthologous interactors, independent of whether they themselves are orthologous

or not. To systematically compare interaction datasets, the interactor profile similarity is then

computed between all possible protein pairs from all included datasets. The complete set of

RBO similarity scores computed between protein pairs form the edges of a large

hypernetwork that connects proteins identified across all included datasets. Clusters of

connected proteins are extracted from this network using the Markov Cluster (MCL)

algorithm (15). We call the clusters resulting from the MCL analysis superclusters. They

contain a mix of proteins from multiple datasets that can originate from different

complexomes, e.g. if they come from different species, or a different subsystem such as a

tissue, cell type or life stage. The clustered proteins are then sorted into subclusters, each

corresponding to one of the included complexomes. To robustly identify interactors, multiple

interaction datasets can be included for each complexome. Exploiting the information in

these datasets for the same complexome, the clustered proteins are scored by determining

the fraction of datasets for which they are part of this cluster. This "fraction clustered" (FrC)

score enables prioritization of likely complex members. To prioritize clusters with high

probability to present actual protein complexes, the resulting superclusters are scored and

filtered based on the similarity between the sets of clustered proteins from different datasets,

as determined for all datasets represented in the cluster. The CompaCt software is available

as a user-friendly command line tool, as well as a flexible Python package, along with

detailed documentation and instructions, from github (https://github.com/cmbi/compact-bio),

the python package index (https://pypi.org/project/compact-bio) and dockerhub

(https://hub.docker.com/r/joerivanstrien/compact-bio).

Combined analysis of eukaryote complexome profiles with CompaCt improves recovery of known protein complexes

To comparatively explore the complexomes of eukaryotes, 53 complexome profiling datasets

were analyzed with CompaCt. These datasets represent a set of 12 complexomes from

different biological systems (e.g.: tissues, life stages, cell-types) in nine different eukaryotic

species. A detailed overview of the analyzed complexome profiling data is shown in Table 1.

Sensitive pairwise orthology predictions (16) were computed by including best bidirectional

hits at the sequence profile level. An overview of the resulting superclusters is presented in

Figure 2, showing the size and cluster coherence score of all clusters, including those that

did not pass CompaCt's filtering step. The cluster coherence score reflects the degree of

commonality between the clustered proteins originating from different datasets. A total of

726 superclusters were identified, each consisting of one or more subclusters that represent

one of the included complexomes. 332 of these passed CompaCt's filtering for coherence

across datasets. Out of these 332 superclusters, 254 were consistently represented in more

than one complexome. To automatically annotate the results, CompaCt identifies clusters

that overlap with a provided set of reference complexes: 36 of the 81 consistently

represented human subclusters overlapped with complexes listed in the CORUM database

(17) (i.e., contain more than half of the total number of subunits comprising the reference

complex). An overview of all clusters that passed filtering, with detailed information, scores,

their composition and automatic annotation is available in Supplemental File SF1.

Table 1. Overview of analyzed complexomes

Alias	Species	Sample	n	Proteins	Fractions	CEDAR study id(s)	Reference (s)
HUM	*H. sapiens*	Fibroblast	8	5630	60	CRX22 CRX17 CRX15 CRX9 CRX8	(18–22)
BOVIN	*B. taurus*	Heart	6	1180	60	CRX33	(23)
ANOST	*A. stephensi*	Salivary tissue	2	925	60	CRX41	This paper

YARLI	*Y lipolytica*		4	3515	60	CRX40	-
PF3_GAM	*P. falciparum*	Gametocyte	4	1672	60-61	CRX23	(24)
PF3_AS	*P. falciparum*	Asexual stage	4	1732	60-61	CRX23	(24)
PF3_SCH	*P. falciparum*	Blood-stage schizont	6	1816	48	CRX20	(9)
BER_SCH	*P. berghei*	Blood-stage schizont	6	1553	48	CRX20	(9)
KNO_SCH	*P. knowlesi*	Blood-stage schizont	6	1885	48	CRX20	(9)
TOX	*T. gondii*	Tachyzoite	1	839	61	CRX27	(25)
AT_LF	*A. thaliana*	Leaf	3	1518	30	CRX24	(26)
AT_SD	*A. thaliana*	Seedling	3	1957	30	CRX24	(26)

Overview of analyzed complexomes, with descriptions of the datasets included per complexome.

Note that a number of analyzed complexomes were specifically designed to focus on the

mitochondrial component. Despite this, most of these datasets also include proteins and complexes

from other cellular compartments. For more detailed descriptions of the specific datasets, please refer

to the methods, respective CEDAR entries and the relevant publications.

Figure 2: Overview of clusters resulting from CompaCt analysis of a collection of complexome

profiling datasets. Each dot corresponds to an MCL supercluster, with dot size corresponding to the

number of subclusters (e.g., species) consistently represented in this cluster. The x-axis represents

the number of proteins consistently represented in the cluster. The cluster coherence score displayed

on the y-axis is a measure for the degree of commonality between the clustered proteins per dataset.

Various clusters that we identified to correspond to known protein complexes have been annotated in

the figure as such. Note that the cluster consisting mainly of ras-related proteins is unlikely to

represent a large complex, as the comigration of multiple members of this protein family is most likely

caused by their similar mass.

CompaCt identifies complexome-specific clusters after first integrating data from multiple

complexomes into a single network. To determine whether the combined analysis of multiple

species with CompaCt improves recovery of protein complexes for a single species when

compared to independent, species-specific analyses, we computed the agreement of human

subclusters with CORUM (17) while progressively including additional complexomes in the

analysis (Figure 3). To compute the agreement of the human subclusters resulting from

CompaCt with the CORUM reference, we use the maximum matching ratio metric (MMR), a

score that computes overlap with a reference based on a mapping between the result

clusters and the reference complexes (27). Starting from separate analysis of just the

datasets from the human complexome, each collection of datasets representing one of the

other complexomes is added to the human data, computing the MMR resulting from its

inclusion. After then adding the complexome data that increased the MMR the most, this

process was repeated until finally data from all 12 complexomes were included. At each step

the complexome data that increased the MMR the most was added to demonstrate that even

after adding the most informative data in the early steps, the MMR increased further as the

remaining datasets were included. In most cases the inclusion of datasets from other

species resulted in a higher MMR, reflecting improved agreement of the human subclusters

with the CORUM reference complexes as compared to separate analysis (Figure 3). This

highlights the utility of including a diverse set of complexome data as opposed to separate

analysis even if the focus is on a specific system.

Figure 3: Effect of including data from additional complexomes to CompaCt analysis on the

agreement of *H. sapiens* cluster results with CORUM v4 (17). Starting with only datasets representing

the human complexome, datasets from additional complexomes were added to the analysis one by

one, computing the MMR after including each. The remaining datasets are then added in a

progressive manner, prioritizing those that resulted in the highest MMR increase. The y-axis displays

the maximum matching ratio (MMR) between the human subclusters and the CORUM reference set

of protein complexes, reflecting the overlap of the identified clusters with the reference. At each step

the MMR resulting from including data from each remaining complexome is shown.

253   CompaCt allows comparative analysis of complexes

254   through robust identification of conserved as well as

255   taxon-specific subunits

To demonstrate that CompaCt is able to accurately recover the composition of complexes

from the analysis of complexome profiling data, including taxon-specific subunits, we

focused on the five well studied and resolved oxidative phosphorylation (OXPHOS)

complexes in *H. sapiens*, *Y. lipolytica* and *A. thaliana*. The ability to include multiple datasets

representing the same complexome allows CompaCt to assign membership confidence

scores to cluster members using the FrC score. Therefore, we determined the recovery of

the OXPHOS complexes by their corresponding clusters while varying the FrC-based

inclusion criteria of cluster members (Figure 4). These results show that selecting proteins

based on a mimimum FrC greatly reduces inclusion of false positives, while losing a limited

number of known subunits. To quantify this, we computed the ratio of recovery of true

positives over false positive and missed scores (i.e., the Jaccard index) for each selection

criterion. Aside from providing the FrC scores for cluster members, CompaCt implements an

additional criterion to include likely relevant cluster members, termed the "best guess"

selection: in addition to proteins that themselves meet the FrC threshold, other clustered

proteins are also included when they have an ortholog or equivalent in one of the

subclusters corresponding to another complexome that does meet the FrC threshold. The

recovery of OXPHOS complexes using this criterion results in the highest Jaccard index in

all but one complexome (Figure 4). Thus, the ability to prioritize proteins based on consistent

comigration in datasets from its own species or one of the other species is helpful in

discerning true complex members from spuriously clustered proteins.

Figure 4: Recovery of the five oxidative phosphorylation complexes by the corresponding subclusters

in four complexomes. The x axis categories correspond to different protein inclusion criteria to the

subclusters. The fractions reflect the minimum fraction clustered (FrC) score for proteins to be

included: i.e., in how many datasets per complexome this protein is part of the cluster. The "bg" ("best

guess") selection criterion includes all proteins that have a fraction clustered of over 1/2. In addition to

those, it also includes proteins with lower FrC scores, but that cluster with an orthologous protein

scoring higher than 1/2. Bars above zero correspond to counts of all proteins included in the

corresponding clusters. Bars below zero correspond to known complex members that are present in

at least one dataset but are not part of the corresponding cluster (false negatives). The numeric

values above the bars represent Jaccard index values (true positives divided by true positive, false

positive and false negative) for each selection criterion. A) *H. sapiens*, B) *Y. lipolytica,* C) *A. thaliana*

seedling, D) *A. thaliana* leaf.

The recovery of complex I, the largest of the OXPHOS complexes, was analyzed in detail to

illustrate that CompaCt's output can be used to reconstruct the composition of a complex in

multiple species (Figure 5). Out of a total of 149 known subunits in these three species, 135

were detected in at least one of the analyzed datasets. Of these, 128 were part of the best

guess selection of the corresponding cluster, four were part of the cluster but fell below the

threshold, and three subunits were not part of the Complex I cluster. 11 proteins are part of

the best guess selection of the cluster while not being a known subunit (HS:0 YL:6, AT:5).

The two Arabidopsis proteins that do not cluster with the other complex I subunits,

At1g65290 and At3g08610, were both only detected in only one out of six Arabidopsis

datasets, and in these data show visibly different migration than other complex I subunits. *Y.*

lipolytica sulfur transferase (ST1) that has been observed in cryo-EM structures to be

associated with complex I (28) does not cluster with the other complex I subunits. Since ST1

was found only in a subset of particles and genetic deletion of its gene does not affect

complex I assembly and function (29) it is considered to be a sub-stoichiometric and non-

essential subunit of *Y. lipolytica* complex I. Like the two *A. thaliana* proteins, its migration in

the complexome profiling datasets differs from the other subunits. This is not surprising since

ST1 is known to dissociate from complex I during BN-PAGE (30). Aside from these three, we

were able to recover all other 132 detected complex I subunits across these three species,

including 13 taxon-specific subunits that are only present in one of the analyzed species.

Figure 5: recovery of the respiratory chain complex I in *H. sapiens, Y. lipolytica* and *A. thaliana* by the

corresponding CompaCt output cluster. Subunits in green are detected and are part of the cluster.

Subunits in yellow were not detected in any of the analyzed complexome profiles. Subunits in red

were detected in at least one complexome profiling dataset, but are not part of the cluster. Taxon-

specific elements of the protein complex are depicted against a colored background, the others

correspond to conserved elements. Proteins in columns are orthologs or best hit homologs.

Large-scale analysis of complexome profiles pinpoints

novel candidate complexes and interactors

All datasets used in this analysis, except those from *A. stephensi,* have been studied

separately before, often to answer a research question regarding a specific complex or set of

complexes. Despite the fact that most analyzed complexome profiles are generated from

samples enriched for mitochondria, they also consistently capture many complexes located

in other compartments of the cell. One of the aims of this study is to leverage the systematic

approach and increased sensitivity from integrative analysis of these data with CompaCt to

identify novel candidate interactors and protein complexes. To this end, we inspected a selection of 45 clusters resulting from the analysis that likely represent (partial) protein complexes (Table 2). The "best guess" selection members of these clusters, from seven species whose complexomes were previously studied, comprise a total of 1603 proteins. By evaluation of the UniProt records of these proteins we determined that for 1343 of these proteins there is already evidence associating them with the corresponding protein complexes, ranging from sequence based predictions to experimental evidence (31). The remaining 260 entries in UniProt contain no evidence of involvement with the respective complexes. Of these, 17 clustered proteins were identified as orthologs of known complex subunits in one of the other analyzed species, suggesting that their function has been conserved. A detailed overview of the identified members per cluster and species is available in Supplemental File SF2. We will discuss three of the analyzed clusters in more detail: the emp24 complex, the F_1F_o-ATP synthase complex and the vacuolar type ATPase complex.

Table 2. Overview of 45 inspected superclusters

Species	Clusters	Assoc. with complex	Not assoc. with complex	Not assoc., subunit ortholog	Total proteins
A. thaliana	23	136	83	2	221
B. taurus	24	169	21	0	190
H. sapiens	38	340	37	3	380
P. berghei	21	143	15	3	161
P. falciparum	28	209	39	3	251
P. knowlesi	24	165	14	5	184
Y. lipolytica	27	181	34	1	216
All species total	45	1343	243	17	1603

Overview of 45 inspected superclusters resulting from CompaCt analysis of eukaryote complexome profiling data, filtered using the "best guess" cluster member selection criterion. The second column shows how many of the 45 inspected clusters contain proteins from each analyzed species. Protein members are divided in three categories based on information available in their Uniprot (31) entries: proteins that have evidence associating them with the respective protein complex, proteins that have no evidence associating them with the complex, and proteins that have no evidence associating them

347   with the complex, but we predicted based on sequence-based homology to be orthologous with a

348   known subunit of the respective complex.

Evidence for an evolutionary conserved p24 complex

Members of the p24 protein family have been found to play a role as cargo receptors in

anterograde as well as retrograde vesicular ER-Golgi trafficking. In yeast it has been shown

that members of this protein family form a heteromeric protein complex, consisting of

emp24p, erv25p, and most likely erp1p and erp2p (32,33). These four proteins consistently

clustered together in *Y. lipolytica*, supporting the hypothesis that they form a heteromeric

complex (Figure 6A). The supercluster containing the *Y. lipolytica* p24 proteins also contains

H. sapiens and *P. falciparum* subclusters with members of the p24 protein family (Figure

6A). Phylogenetic analysis of all the members of this protein family (InterPro ID:

IPR015720)(34) in these three species shows that all but one (TMED6) of the human

representatives of this family were detected in the analyzed complexome profiles, and are

part of the same cluster. While interactions between some of the human p24 proteins have

previously been shown and are e.g., present in the protein-protein interaction database

STRING (35), the proteins have not yet been assigned to a multisubunit complex in *H.

sapiens*. The apparent mass of the detected protein complex estimated from the human

complexome profiles is ~110-120 kDa (supplementary figure S1), approximately matching

the mass of a putative heteromeric complex of the five clustered subunits (125 kDa).

TMED4, TMED5 and TMED7 result from recent gene duplications at the root of the

vertebrates (Figure 6B). They are part of the p24 cluster, but do not make the "best guess"

selection threshold because they do not cluster as consistently with the complex and are not

predicted as one-to-one orthologs with one of the other clustered proteins. Nevertheless,

their presence in the cluster and their homology suggests that they may sub-

stoichiometrically replace their respective in-paralogs in a subset of the human p24

complexes. The *P. falciparum* subcluster contains all four *P. falciparum* representatives of

the p24 protein family (Figure 6A), but no specific investigation of the p24 complex in malaria

parasites has been performed. Two of these (PF3D7_0526900, PF3D7_0422100) have

however been shown to interact in a pairwise assay and to be essential for liver-stage

viability or sporozoite infectivity (36). Notably, PF3D7_13333300 has been predicted to

interact with an apicoplast-resident protease, which seems inconsistent with a role in ER-

Golgi protein trafficking (37). To what degree the *Plasmodium* emp24 complex function is

conserved and whether there is a specific function in the liver or sporozoite stages remains

to be explored.

Figure 6: A) Overview of CompaCt output cluster containing proteins of the p24 family in *Y. lipolytica*,

H. sapiens and *P. falciparum*. Clustered proteins with a fraction clustered score of over 0.5 or best hit

homologs of those are displayed. Proteins in columns are orthologs or best hit homologs with two or

more proteins in the same block, reflecting likely gene duplications. B) Phylogenetic tree of all

representatives of the p24 protein family in the three aforementioned species. Species-specific

branches have been colored. Members of the protein family that were not detected in the data used in

this project are shown in gray. Sequences were aligned with ClustalOmega (38), and the phylogenetic

tree was reconstructed with PhyML (39). The dynamic evolution of the protein family whose members

are part of one complex are well captured by the CompaCt approach.

High confidence assignment of ATP synthase subunits

The F_1F_o-ATP synthase complex is a highly conserved complex central to energy conversion

in the mitochondria of eukaryotes and bacteria and has been well studied in all model

species. In multiple complexomes, many members of this complex form a supercluster

(Figure 7). Two human cluster members were not reported as ATP synthase subunits

previously, but one of them (C15orf61) was recently observed to comigrate with ATP

synthase subunits by Morgenstern *et al.* in a complexome profiling experiment (40). We

could confirm that C15orf61 consistently co-migrates with ATP synthase across multiple

replicates and studies, as it is the only non-ATP synthase subunit cluster member with an

FrC as high as 6/8. While it has been annotated as a secreted protein based on

experimental data, it is predicted to be mitochondrial protein in Mitocarta 3.0 based on

MS/MS and APEX labelling experiments and on the presence of a predicted mitochondrial

target sequence (41). To determine whether absence of C15orf61 affects functioning of the

ATP synthase complex, we generated a C15orf61 knock-out in human HEK cells. Enzymatic

activities of the five individual OXPHOS complexes were not significantly affected by ablating

C15orf61, although the activity of complex I tended to be somewhat lower (Figure S2).

IGF2BP2 (FrC: 5/8) is not predicted to be a mitochondrial protein, and has been ascribed a

function in mRNA binding and transport (42). The *P. knowlesi* ATP synthase subcluster

contains four proteins that have not been described previously as subunits of this complex.

Three of those subunits: PKNH_0117300, PKNH_0725400 and PKNH_1124100 are

orthologous to known *T. gondii* ATP synthase subunits, suggesting that their function as

complex V subunits is conserved among apicomplexa. Supporting this, PKNH_0725400 and

PKNH_1124100 are predicted to be mitochondrially located in PlasmoMitoCarta (43). Lastly,

the uncharacterized TGGT1_233890 protein, one of the *T. gondii* cluster members not

previously described as a subunit, has been assigned to the inner mitochondrial membrane

(44), consistent with our assignment.

Figure 7: Overview of CompaCt output cluster containing subunits from the mitochondrial ATP

Synthase complex in *H. sapiens, P. knowlesi and T. gondii complexomes*. Clustered *P. falciparum*

and *H. sapiens* proteins with a fraction clustered score of over 0.5 or best hit homologs of those are

displayed, as well as their clustered *Toxoplasma* orthologs. Human proteins labeled "predicted

mitochondrial" are listed in Mitocarta 3.0 (41). Plasmodium and Toxoplasma proteins with this label

are predicted as such by PlasmoMitoCarta (43) and a HyperLOPIT study (44), respectively.

The V-ATPase complex outside the Metazoa

Vacuolar type ATPase (V-ATPase) is a highly conserved eukaryotic proton pump that is

responsible for acidification of a variety of intracellular compartments or whole cells. The

complex is subdivided into two modules, the membrane bound V_0 module and the

hydrophilic V_1 module. The V_0 module of this complex is represented in a separate

supercluster (cluster 7) from the V_1 module supercluster (cluster 13, Supplemental File SF1),

caused by the differing migration patterns of the two modules, which in turn is likely due to

dissociation during the experimental procedure. Of the V_0 module the *H. sapiens*, *Y.*

lipolytica, *A. thaliana* and *P. falciparum* subclusters are shown in Figure 8. We present the

first characterization of the *P. falciparum* V-ATPase complex based on experimental data.

While some of the detected subunit identities are clear from sequence-based homology, our

cluster contains several potential novel subunits of this complex that are either taxon-specific

or whose orthology could not be established through standard homology detection methods.

Figure 8: Overview of CompaCt output cluster containing subunits from the membrane component

(V_0) of the Vacuolar ATPase complex (V-ATPase) in *H. sapiens, Y. lipolytica, A. thaliana* and *P.*

falciparum complexomes. Clustered proteins with a fraction clustered score of over 0.5 or best hit

homologs with those are displayed. Proteins in columns are orthologs or best hit homologs, and are

labeled depending on whether they have previously been associated with this protein complex. Two

or more proteins in the same block indicate likely gene duplications. For proteins labeled as

"associated with the complex" there is some previous evidence associating them with the V-ATPase

complex, while for the other proteins there is not.

Although two members of this cluster were annotated as proteins of unknown function,

profile-based homology searches identified them as orthologs of the human V-ATPase

subunit ATP6AP2: at3g24160 (*A. thaliana*) and PF3D7_0613100 (*P. falciparum*). Orthologs

of ATP6AP2 were not identified previously as subunits of V-ATPase outside of the Metazoa.

Another protein that is part of the *P. falciparum* subcluster is PF3D7_0713700, which we

identify as a likely ortholog of human V-ATPase subunit ATP6AP1, even though it was not

predicted to have a human ortholog as the hhsearch result was above the threshold (E-

value=1.6). It should be noted that ATP6AP1 evolves at a very high rate, as also the

sequence identity between the human and the *Saccharomyces cerevisiae* ATP6AP1

proteins is very low (45). Additionally, we predict A0A182XX75 as the *A. stephensi* ortholog

of ATP6AP1, which is also part of the cluster corresponding to the V_0 component of V-

ATPase. Notably, the C-terminal transmembrane (TM) region is the only part that is

conserved in this protein family, whose lack of sequence conservation has been established

before (45,46). We now extend ATP6AP1 to *Anopheles* and *Plasmodium*, of which the

Plasmodium ATP6AP1 is only 209 amino acids long with two predicted TM helices

(positions: 5-25, 168-190), compared to 239 amino acids in human, with a single TM region

after cleavage of the original protein at the furin cleavage site (46). An alignment of the C-

terminal TM region of ATP6AP1 and its orthologs in a number of species in shown in

Supplemental Figure S3. The *Y. lipolytica* subcluster contains YALI0_F09405g, a protein

that is predicted to contain a "V-ATPase proteolipid subunit" domain (IPR035921) in the

InterPro database (34), but whose complex membership had not yet been established.

Additionally, several proteins that have no orthologs in one of the other analyzed species are

part of this cluster. The human MCUR1 protein has a known function as a regulator of the

mitochondrial calcium uniporter and is localized to mitochondria (47), so it is not likely to be

part of this complex. The *Y. lipolytica* protein YALI0_E16126g has no known function, but is

co-expressed with known subunits as indicated in the STRING database (35) and thus a

potential interactor of the *Y. lipolytica* V-ATPase complex. Finally, several V-ATPase

candidates are included in one of the two *P. falciparum* complexomes, of which only

PF3D7_1112000 has a high fraction clustered score (6/8) in both. PF3D7_1112000 is a

small (72 amino acid) protein of unknown function that is predicted to contain two

transmembrane helices that take up the majority of the protein, suggesting a possible role in

the V_0 module of the V-ATPase complex.

CompaCt analysis of novel complexome sheds light on respiratory chain complex evolution

To demonstrate the suitability of CompaCt for the characterization of an unstudied complexome, we included a previously unpublished dataset from the mosquito *A. stephensi.* This is the first large-scale analysis of complexome profiling data from a species from the Protostomes, one of the two major taxa of the Bilateria. We provide experimental evidence for the origin of the constituents of multiprotein complexes that hereto were mainly based on their orthology to experimentally characterized proteins from the Deuterostomes, the other major taxon that contains the vertebrates. We generated two datasets as byproducts from isolation of *P. falciparum* sporozoites from *A. stephensi* salivary glands and detected 925 proteins of the mosquito host of the parasite. Supplementary Table S1 provides an overview of putative *A. stephensi* complex subunits identified through analysis with CompaCt. From the superclusters corresponding to known complexes in one of the well-studied species like *H. sapiens* and *A. thaliana*, we identified fifteen clusters that contain *A. stephensi* proteins homologous to known cluster members in one of the other species. Across these clusters we provide experimental support for 87 proteins whose association with the respective complexes were hereto solely based on sequence similarity. Additionally, we identify five proteins of unknown function that both cluster with the complex and are orthologous to a known subunit in one of the other analyzed species. The complete list of identified proteins per cluster is available in supplemental file SF2.

One of the identified clusters corresponds to the mitochondrial respiratory chain complex cytochrome *c* oxidase (complex IV). This cluster contains four proteins predicted to be associated with this protein complex based on sequence similarity: A0A182XZQ2, A0A182YLP4, A0A182YLP7 and A0A182Y5T1. Additionally, this cluster contains *Anopheles* protein A0A182Y1F7, the ortholog of the *Drosophila CG7630* protein, recently identified as an ortholog of mammalian cytochrome *c* oxidase subunit COX7B, which was previously

believed to be vertebrate specific (48). The alignment of COX7B, the *Anopheles* A0A182Y1F7 protein and the *Drosophila CG7630* protein is shown in Supplemental Figure S4. The *A. stephensi* ATP synthase cluster contains two proteins (A0A182YCB1, A0A182YSV3) that are best hit homologs with the human beta subunit, and two proteins (A0A182Y5P7, A0A182YCU3) that are best hit homologs with the epsilon subunit, indicating that these two genes have been duplicated and that all four have retained their role as subunits of complex V.

Discussion

The increasing availability of complexome profiling data from various species in the public CEDAR database has opened the door for large scale and automated analysis of these data (13). However, methods that enable automated large-scale comparative analysis of protein interaction data to exploit this wealth of information were lacking. In this work, we demonstrate the potential of large-scale integration of protein interaction data from multiple species with CompaCt. Our approach of integrating interaction data has several benefits over existing methods. Firstly, comparing datasets using interactor profiles avoids direct comparison of migration patterns, circumventing difficulties caused by batch effects introduced through differences in protocols and complex separation resolution hampering direct comparisons. Secondly, indirect usage of orthology to determine similarity between two proteins' set of interactors rather than directly connecting orthologous proteins enables incorporation of proteins with no known orthologs and therewith the detection of taxon-specific interaction members or the proteins that have diverged beyond what can be detected by sequence-based homology detection methods. An example of the former is the emp24 complex supercluster, where we determined the putative composition of this complex in *H. sapiens*, *P. falciparum* and *Y. lipolytica*, even though it varies between these species and contains taxon-specific elements. This demonstrates that the dynamic evolution of the protein family whose members are part of one complex are well captured by our approach.

An additional benefit of this approach is that it exhibits robustness to incomplete orthologies

or imperfect correspondence of protein identifiers between datasets. We found that in many

such cases CompaCt is still able to recover these proteins and interactions based on

orthology between the interactors. For example, a number of mitochondrially encoded

complex I subunits were correctly clustered, even though their orthology was missed

because of mismatched identifiers between the various complexome data and the

proteomes used for orthology predictions (Figure 5, Supplemental File SF1).

One of the main challenges in the identification of complexes from complexome profiling

data is the occurrence of spuriously comigrating proteins or protein complexes, as

thousands of identified proteins are separated into typically just 60 fractions. We

demonstrate that by systematically prioritizing cluster members based on consistency of

interactions, enabled through inclusion of multiple interaction datasets from the same

species, we were able to greatly simplify identification of novel potential interactors that were

missed in previous individual analyses of the datasets used. For example, we identified a

number of novel candidate interactors in the V-ATPase complex, of which PF3D7_0613100

and AT3G24160 are predicted as orthologs of human V-ATPase subunit ATP6AP2 through

profile-based homology (Figure 8). Similarly, PF3D7_0208800 and PF3D7_0505900

consistently cluster with the ER membrane and the prefoldin complexes respectively, and

are predicted as orthologs of human subunits of these complexes (EMC10, PFDN6) using

profile-based homology (Supplemental file SF1).

CompaCt integrates data from multiple complexomes in a single network and then clusters

them together, after which complexome-specific clusters are extracted. We show that this

has two major advantages over separate analyses in which proteins are assigned to

complexes based on data from a single complexome. Firstly, conserved protein complexes

present in multiple species will generally be grouped together as subclusters of the same

cross-species supercluster, foregoing the need to a posteriori match or align clusters from

separate analyses. More importantly, we demonstrate that our method of combined

clustering of multiple complexomes results in overall better recovery of protein complex

composition, by showing the agreement with a known reference improves when including

additional complexomes in the analysis. E.g., protein complexes that are only partially

detected in a dataset will often be difficult to detect, but can still contain novel interactors

missed in other experiments. An example of this are the three candidate *Plasmodium* ATP

synthase subunits identified in this study, that were missed in the initial publications despite

being orthologous to known *T. gondii* ATP synthase subunits. Evers et al. analyzed the

mitochondrial complexome of gametocyte and asexual blood-stage parasites, where they

described the composition of ATP synthase but failed to detect these proteins (24). Hillier et

al performed large-scale clustering analysis of whole-cell schizont complexome profiles,

where they did not identify a well-defined ATP synthase cluster, as the complex was well

represented in only one of their replicates (9). In our combined analysis, ATP synthase was

represented in multiple complexomes, resulting in a well-defined ATP synthase cluster,

which included the three subunits detected in the data from Hillier et al, that remained

undetected in the complexome profiles of Evers et al.

By inherently allowing transfer of information between species in an unsupervised manner,

CompaCt simplifies large-scale characterization of previously unstudied complexomes. This

is demonstrated by the identification of a number of complexes and interactors in the *A.*

stephensi complexome. Automated large-scale characterization of complexes in previously

unstudied species can shed light on complex evolution. While the evolutionary origin of

subunits can be inferred based solely on the distribution of orthologs (49), this approach has

two pitfalls. Firstly, it may not be able to detect the homolog through sequence-based

methods. An example of this is A0A182XX58, an *A. stephensi* protein that clusters with

complex I; its closest human homolog is the complex I subunit NDUFA3, but it was not

identified as orthologous in our analysis because the E-value resulting from HHsearch did

not meet the significance threshold (hhsearch E-value = 0.056). While drafting this

manuscript, the *D. melanogaster* ortholog of this protein was identified as an ortholog of

NDUFA3 based on Cryo-EM data (50), supporting this finding. Secondly, the presence of an

ortholog in a species does not necessarily indicate that it is part of the same complex. As an

example, the *P. falciparum* protein ApiCOX13 (PF3D7_1022900) has recently been

identified as a subunit of complex IV, and is part of our complex IV cluster. However, using

sequence-based homology this protein can be identified as orthologous to human protein

CISD3, which is not part of human complex IV. The *Anopheles* subcluster corresponding to

respiratory chain complex I contains A0A182YAZ9, the *A. stephensi* ortholog of the

mammalian complex I subunit NDUFA10. While this protein has orthologs throughout the

Metazoa, it is unclear when this protein became part of complex I, as evidence suggests a

nucleoside kinase function outside of its role as complex I subunit and the protein binds

dGTP (51). Notably, this protein is homologous to deoxyribonucleoside kinases, and its

active sites have been conserved (52). Furthermore, its migration in human and bovine

complexome profiles (Supplemental Figure S5) suggest that it is, besides being part of

complex I, also present as a monomer, pointing towards a potential dual role as monomeric

enzyme and subunit of complex I. Our results indicate that this protein is part of complex I in

Protostomes, suggesting it was already a subunit of complex I at the root of the Bilateria.

The migration of this protein in *A. stephensi* does not clearly indicate it is present as a

monomer (Figure S5).

In this study, to score interaction likelihood between protein pairs within complexome profiles

we used the Pearson correlation. This metric is most commonly used and generally performs

very well to determine interaction partners from complexome profiles. However, recent work

has suggested and demonstrated the effectiveness of several other metrics to determine

interactions (7,8,11,53,54). Here we refrained from using those, as our focus was on the

method of integrating interaction datasets rather than identifying the best within-dataset

interaction metric. However, as other metrics might be better suited in certain situations to

determine interaction likelihood, it is important to note that the CompaCt software is able to

analyse interaction matrices previously scored with any similarity metric.

In conclusion, the approach of comparing protein interaction data between species

presented here enables more complete comparative analyses of multiple complexomes than

existing methods. While CompaCt was used to analyze multiprotein complexes and their

interactors here, it could also be used for the comparative analysis of other types of

interaction data like gene co-expression data or to identify modules of coregulated genes.

Methods

CompaCt software

Input data

CompaCt performs integrative cluster analysis on multiple protein interaction datasets

representing one or more biological systems (e.g., species, tissue, cell-type etc.).

An interaction dataset consists of a set of elements that represent proteins, with real number

scores between each pair of elements representing interaction likelihood, which can be

represented as a symmetric matrix of correlation or interaction scores. The provided

datasets are subdivided into "collections", where each collection should represent a distinct

biological system in which we expect the protein interactions to be identical to each other

(e.g., human fibroblast, *Arabidopsis* leaf, *Plasmodium* mitochondria, etc.). Inclusion of

multiple datasets representing the same collection can be included to more robustly identify

true interactors, by prioritizing consistently clustering proteins. Multiple datasets representing

the same species are expected to use the same protein identifiers, but are allowed to

contain different sets of elements of varying sizes. In order to be able to compare and

identify common elements between datasets from different species, pairwise, "one-to-one"

orthologies between these species need to be provided. A pairwise orthology between

species is represented as a set of identifier pairs, one from each species, that represent

orthologous proteins. It is not required that every protein present in the data of one of the

species has an ortholog in any of the other species.

Comparing interactor profiles with rank biased overlap

For each element in each dataset, a ranked list of all other elements within this dataset,

ordered from highest interaction score to lowest, is determined. This ranked list is referred to

as the molecule's interactor profile. To be able to determine for two proteins from different

interaction datasets to which degree they have a common set of interacting proteins, we

determine the similarity between their interactor profiles, using the non-extrapolated rank

biased overlap (RBO) metric (14) (github.com/changyaochen/rbo). RBO is a set-based

overlap metric, that assigns a value between 0 and 1 representing the degree of overlap

between two ranked lists.

The RBO metric has two properties that make it suitable for the comparison of interactor

profiles. First, the ranked lists can contain elements not common in both lists and allowed to

have different lengths. This is required for the comparison of interactor profiles, as often a

different set of proteins is identified in different datasets, and when comparing between

species it is likely that part of the proteins will not have an ortholog in the other species.

Secondly, RBO is a "top-heavy" score, assigning more weight to the top of the list when

determining overlap, with a tunable parameter that determines the extent to which the metric

is "top-heavy". A focus on the high-scoring interactors makes biological sense, as only the

top scoring proteins are likely to be true interactors of the given protein. This RBO top-

heaviness parameter is a tunable parameter in the CompaCt tool. To determine a suitable

default value for this parameter for the analysis of complexome profiling data, we determined

the effect of changing this parameter on recovering pairs of proteins belonging to the same

protein complex (Supplemental Figure S6), showing that a parameter value of 0.9 results in

the highest median rank of all protein pairs part of the same complex, and that the rank

biased overlap is able to prioritize protein pairs that are part of the same complex over other

protein pairs.

To limit computational cost, the search depth when computing RBO scores, i.e., the length of

the ranked lists, is limited so that lower ranks that have very little influence on the RBO score

are not considered. By default, the contribution of each rank is computed, and the ranks that

cumulatively contribute 99% of the RBO score are used. To compute rank biased overlap

between interactor profiles from different species, pairwise orthologies are used to determine

the common elements between the ranked lists. Proteins from different species that are

orthologous are considered common elements when computing RBO scores, while the

others are considered unique to that list.

Reciprocal top hits
687

Pairwise all-to-all comparisons between 2 datasets result in a large number of RBO scores,

of which the majority will not represent proteins with actual shared interactors. To reduce the

amount of data and CPU time that is considered in the next steps of the CompaCt workflow

we want to exclude uninformative similarity scores. However, absolute RBO score values

representing a true set of shared interactors vary greatly, because proteins from a small

protein complex will result in much lower RBO scores than proteins that are part of a large

complex, due to the difference in the number of shared interactors. Therefore, to select

relevant RBO scores when comparing two datasets we determine whether for both proteins

the RBO score between them is in the top 1% (adjustable parameter in CompaCt) of all

scores for the protein in question. If this is the case, we refer to it as a "reciprocal top hit",

and the edge between this pair is included in the hypernetwork used for cluster analysis.

Combined network clustering
700

The approach described above is used to systematically determine interactor profile

similarity of all proteins between two datasets. This is then performed in a pairwise manner

comparing each dataset to all others regardless of which collection (i.e., a number of

datasets representing the same species/biological system) they are a part of. To perform

combined clustering of all datasets, similarity scores originating from all pairwise dataset

comparisons should be comparable. However, as different datasets likely have a different

number of common proteins, the similarity scores will likely vary between the pairs of

datasets compared. Therefore, before these scores are combined and then clustered, they

are normalized such that they have the same mean. After computation of similarity scores,

selection of reciprocal top hits and normalization, the resulting hypernetwork connecting

proteins from all included datasets is clustered using the MCL cluster algorithm version 14-

137 (15) using the default parameters.

Processing clusters: prioritizing proteins

The resulting clusters, extracted from the hypernetwork by MCL, can contain elements

originating from multiple datasets and collections. These super-clusters are first split into

subclusters separating elements by input collection. The subclusters, containing elements

from datasets belonging to the same collection, are then further processed to assign a single

"fraction clustered" (FrC) score to each unique protein that represents the consistency with

which it is part of a cluster. Per protein and subcluster, the number of elements from different

datasets representing this protein are counted, and divided by the total number of datasets

in the corresponding collection (e.g.; if 4 datasets are provided for a given collection, and the

current subcluster contains elements corresponding to this protein for 3 out of 4 of those

datasets, the fraction clustered score for this protein in this cluster equals ¾). Note that to

compute the FrC, the total number of datasets from that complexome is used as the

denominator instead of the number of datasets that this protein is detected in, as we

consider a protein not being detected lack of evidence of interaction.

Processing clusters: prioritizing clusters

Additionally, to determine the coherence score of a supercluster, the fraction of possible matches is computed, reflecting the degree to which the same or orthologous proteins from different datasets are clustered together. A match is defined as: two "equivalent" elements from different datasets are part of the same supercluster. Equivalent elements are either two elements representing the same protein in two datasets from the same species, or two orthologous proteins from different species. The total number of matches found within a supercluster is then divided by the number of possible matches given the cluster's composition, to get the fraction of possible matches. To prioritize likely biologically relevant clusters, CompaCt performs a filtering step, retaining only clusters with a minimum of 2 matches as well as having at least one protein with a fraction clustered of at least 0.5.

Annotation of clusters with reference complexes

To simplify identification of protein complexes from the output clusters, CompaCt implements automatic annotation of clusters using a provided reference. For one of the input collections a collection of reference complexes can be provided. For each provided reference complex, the cluster that contains the most members of this complex is identified. If a large enough fraction of the complex is present in the cluster, for which a threshold can be provided, it will be annotated with the name of the reference complex.

Analysis of eukaryote complexomes

Complexome profiling

Salivary glands containing *P. falciparum* sporozoites were collected from female *A. stephensi* mosquitoes (Sind-Kasur Nijmegen strain) and consequently homogenized (55,56). The homogenate was then either separated into gland material and *P. falciparum* sporozoite material or kept unseparated. The resulting protein samples were solubilized and

subsequently separated using BN-PAGE (57). A complete description of this process is

available in the supplemental methods. In-gel trypsin digestion followed by mass

spectrometric analysis was performed as described in (24).

The remaining complexome profiles used in this paper are reused from published studies

and have been made publicly available on the CEDAR database, from which they were

retrieved. A detailed overview of the used datasets including CEDAR accessions is available

from Supplementary Table S2. The protein abundances were used as is, no normalization or

preprocessing was performed before analysis with CompaCt. Identifiers were converted to

match those used in orthology predictions.

Cell culture conditions

765 HEK293T cells (ATTCC, 293T-CRL-3216) were cultured in High glucose DMEM medium

766 supplemented with 10% fetal calf serum, 1% Penicillin/Streptomycin and 1% Sodium

767 pyruvate. Cells were cultured at 37 °C and 5% CO_2.

Generation of KO HEK293T cells

770 C15orf61 was knocked-out in HEK293T using CRISPR/Cas9 system, followed by clonal

771 expansion and characterization. The characteristics of the resulting clones used in this work

772 is shown in Table 3. A complete description of the methods is available in the supplementary

773 material.

774 **Table 3. Characteristics of the different c15orf61 HEK293T clones used in this work**

Clone	Alias	Passage	Allele 1	Allele 2	gRNA used
HEK293T not transfected	FM-WT HEK		WT	WT	None
Clone 2E7	FM-2-E7	29	WT	WT	gRNA1

Clone 4E7	FM-4-E7	29	WT	WT	gRNA1+2
Clone 2D4	FM-2-D4	29	c.-8_132delins83 p.Met1?	c.-101_95delins45 p.Met1?	gRNA1
Clone 2E11	FM-2-E11	29	c.-149_103delins149 p.Met1?	c.-149_103delins149 p.Met1?	gRNA1
Clone 3C10	FM-3-C10	29	c.26_*243del p.(Ala11Cysfs*21)	c.26_*243del p.(Ala11Cysfs*21)	gRNA1+2
Clone 4F5	FM-4-F5	29	c.-152_*267del p.Met1?	c.-152_*267del p.Met1?	gRNA1+2

Enzyme activity measurements

777 Enzyme activity measurements of respiratory chain complexes in clonal cell-lines were

778 performed as previously described (58). Presented activity values are relative to the activity

779 of respiratory chain complex II.

Orthology prediction

Pairwise orthologies can be provided as input to CompaCt, alongside interaction datasets, to

allow comparison between interaction data from different species. To sensitively determine

pairwise orthology between all species included in this project, we determined the

bidirectional best hits between their respective proteomes. The proteomes used in the

homology searches are listed in Supplementary Table S3. To determine bidirectional best

hits between two proteomes, protein blast searches were performed in both "directions". Any

proteins that did not have a significant (Evalue < 0.001) hit, were subsequently searched

using hhsearch from the hhsuite toolkit using default parameters (59). The set of best hits

from blast was complemented with any significant best hit hhsearch results. This combined

set of best hits was used to determine bidirectional best hits between each pair of

proteomes. The blast software version 2.9.0 was used. Custom blast databases were

generated from the aforementioned proteomes. All-to-all blast searches were run against

these databases using blastp with default settings. The hhsuite software version 3.0 was

used. Custom profile databases were generated from the proteomes used in each species,

following the protocol available from (https://github.com/soedinglab/hh-suite/wiki#building-

customized-databases), skipping the secondary structure annotation as recommended.

Profile based homology searches were run with hhsearch using default settings.

Annotation and evaluation of results using CORUM reference

For automatic annotation of CompaCt's output clusters, a reference set of known protein

complexes can be provided. For this we used the complete human complexes from CORUM

v4 (17). To evaluate the agreement of CompaCt's resulting clusters with a set of human

reference complexes, we performed some further processing on the human CORUM

reference. Firstly, any duplicate complexes have been removed, so that each complex

occurs once. Additionally, any complexes that are a subset of other complexes, generally

corresponding to sub-assemblies, were removed. To enable fair assessment of the

clustering approach, rather than of the completeness of the underlying data, any proteins not

detected in at least two of the eight human datasets were removed from the reference.

Lastly, any protein complexes containing two or fewer proteins (after the previous steps)

were removed from the reference, to focus on multiprotein complexes that are well

represented in the underlying data. To quantitatively assess the agreement of CompaCt's

output clusters with the aforementioned reference, we used the maximum matching ratio, as

described in (27).

Analysis of eukaryote complexome profiles with CompaCt

The complexome profiles described earlier were analyzed together with the CompaCt

command line tool, using default settings. The aforementioned pairwise orthologies were

included to allow comparison between species, and the CORUM reference was included for

automatic annotation of the resulting clusters. In order to assess the effect of including data

from additional complexomes on the human cluster results, several additional CompaCt

analyses were run using various combinations of complexome datasets. First, a single

collection of eight human complexome profiling datasets corresponding to the human

complexome were analyzed. Then, in separate runs, the human datasets were analyzed

together with each of the remaining complexomes. The combination that resulted in the best

agreement of human subclusters with the CORUM reference was selected for further

analysis. This process was then repeated progressively, until all complexomes were

included in the final analysis.

Annotation of cluster members in inspected clusters

To determine whether protein members of the analyzed result clusters have previously been

associated with the respective protein complexes, their UniProt entries were inspected for

any evidence associating them with the protein complexes represented by the respective

clusters (last accessed on Nov 3, 2022) (31). Any type of evidence suggesting involvement

with this protein complex was deemed sufficient, be it experimental evidence or automated

predictions. Additionally, some proteins that lacked evidence of involvement on UniProt but

for which we were aware of recent evidence showing association with the corresponding

complex were categorized as such.

Acknowledgements

840
841   We would like to thank Marga van de Vegte-Bolmer, Rianne Stoter and Wiebe Kooijman for

842   generating the *P. falciparum* parasites and the mosquito infections. We would like to

843   acknowledge Astrid Pouwelsen, Saskia Mulder, Jolanda Klaassen, Laura Pelser-Posthumus

844   and Jacqueline Kuhnen for breeding of the mosquitoes and for all the salivary gland

845   dissections to obtain the sporozoites.

References

1. Cabrera-Orefice A, Potter A, Evers F, Hevler JF, Guerrero-Castillo S. Complexome Profiling—Exploring Mitochondrial Protein Complexes in Health and Disease. Front Cell Dev Biol. 2022 Jan 12;9:796128.

2. Heide H, Bleier L, Steger M, Ackermann J, Dröse S, Schwamb B, et al. Complexome Profiling Identifies TMEM126B as a Component of the Mitochondrial Complex I Assembly Complex. Cell Metab. 2012 Oct;16(4):538–49.

3. Giese H, Ackermann J, Heide H, Bleier L, Drose S, Wittig I, et al. NOVA: a software to analyze complexome profiling data. Bioinformatics. 2015 Feb 1;31(3):440–1.

4. Páleníková P, Harbour ME, Ding S, Fearnley IM, Van Haute L, Rorbach J, et al. Quantitative density gradient analysis by mass spectrometry (qDGMS) and complexome profiling analysis (ComPrAn) R package for the study of macromolecular complexes. Biochim Biophys Acta BBA - Bioenerg. 2021 Jun;1862(6):148399.

5. Schulte U, den Brave F, Haupt A, Gupta A, Song J, Müller CS, et al. Mitochondrial complexome reveals quality-control pathways of protein import. Nature. 2023 Feb 2;614(7946):153–9.

6. Bludau I, Heusel M, Frank M, Rosenberger G, Hafen R, Banaei-Esfahani A, et al. Complex-centric proteome profiling by SEC-SWATH-MS for the parallel detection of hundreds of protein complexes. Nat Protoc. 2020 Aug;15(8):2341–86.

7. Hu LZ, Goebels F, Tan JH, Wolf E, Kuzmanov U, Wan C, et al. EPIC: software toolkit for elution profile-based inference of protein complexes. Nat Methods. 2019 Aug;16(8):737–42.

869 8. Stacey RG, Skinnider MA, Scott NE, Foster LJ. A rapid and accurate approach for

870   prediction of interactomes from co-elution data (PrInCE). BMC Bioinformatics. 2017

871   Dec;18(1):457.

872 9. Hillier C, Pardo M, Yu L, Bushell E, Sanderson T, Metcalf T, et al. Landscape of the

873   Plasmodium Interactome Reveals Both Conserved and Species-Specific Functionality.

874   Cell Rep. 2019 Aug;28(6):1635-1647.e5.

875 10. Wan C, Borgeson B, Phanse S, Tu F, Drew K, Clark G, et al. Panorama of ancient

876   metazoan macromolecular complexes. Nature. 2015 Sep;525(7569):339–44.

877 11. Nolte H, Langer T. ComplexFinder: A software package for the analysis of native protein

878   complex fractionation experiments. Biochim Biophys Acta BBA - Bioenerg. 2021

879   Aug;1862(8):148444.

880 12. Van Strien J, Guerrero-Castillo S, Chatzispyrou IA, Houtkooper RH, Brandt U, Huynen

881   MA. COmplexome Profiling ALignment (COPAL) reveals remodeling of mitochondrial

882   protein complexes in Barth syndrome. Valencia A, editor. Bioinformatics. 2019 Sep

883   1;35(17):3083–91.

884 13. van Strien J, Haupt A, Schulte U, Braun HP, Cabrera-Orefice A, Choudhary JS, et al.

885   CEDAR, an online resource for the reporting and exploration of complexome profiling

886   data. Biochim Biophys Acta BBA - Bioenerg. 2021 Jul;1862(7):148411.

887 14. Webber W, Moffat A, Zobel J. A similarity measure for indefinite rankings. ACM Trans Inf

888   Syst. 2010 Nov;28(4):1–38.

889 15. van Dongen SM. Graph clustering by flow simulation. 2000.

890 16. Szklarczyk R, Wanschers BF, Cuypers TD, Esseling JJ, Riemersma M, van den Brand

891   MA, et al. Iterative orthology prediction uncovers new mitochondrial proteins and

identifies C12orf62 as the human ortholog of COX14, a protein involved in the assembly

of cytochrome c oxidase. Genome Biol. 2012 Feb 22;13(2):R12.

17. Giurgiu M, Reinhard J, Brauner B, Dunger-Kaltenbach I, Fobo G, Frishman G, et al.

CORUM: the comprehensive resource of mammalian protein complexes—2019. Nucleic

Acids Res. 2019 Jan 8;47(D1):D559–63.

18. Gardeitchik T, Mohamed M, Ruzzenente B, Karall D, Guerrero-Castillo S, Dalloyaux D,

et al. Bi-allelic Mutations in the Mitochondrial Ribosomal Protein MRPS2 Cause

Sensorineural Hearing Loss, Hypoglycemia, and Multiple OXPHOS Complex

Deficiencies. Am J Hum Genet. 2018 Apr 5;102(4):685–95.

19. Lobo-Jarne T, Pérez-Pérez R, Fontanesi F, Timón-Gómez A, Wittig I, Peñas A, et al.

Multiple pathways coordinate assembly of human mitochondrial complex IV and

stabilization of respiratory supercomplexes. EMBO J. 2020 Jul 15;39(14):e103912.

20. Alston CL, Veling MT, Heidler J, Taylor LS, Alaimo JT, Sung AY, et al. Pathogenic Bi-

allelic Mutations in NDUFAF8 Cause Leigh Syndrome with an Isolated Complex I

Deficiency. Am J Hum Genet. 2020 Jan 2;106(1):92–101.

21. Huynen MA, Mühlmeister M, Gotthardt K, Guerrero-Castillo S, Brandt U. Evolution and

structural organization of the mitochondrial contact site (MICOS) complex and the

mitochondrial intermembrane space bridging (MIB) complex. Biochim Biophys Acta.

2016 Jan;1863(1):91–101.

22. Chatzispyrou IA, Guerrero-Castillo S, Held NM, Ruiter JPN, Denis SW, IJlst L, et al.

Barth syndrome cells display widespread remodeling of mitochondrial complexes without

affecting metabolic flux distribution. Biochim Biophys Acta Mol Basis Dis. 2018

Nov;1864(11):3650–8.

23. Hevler JF, Zenezeni Chiozzi R, Cabrera-Orefice A, Brandt U, Arnold S, Heck AJR.

Molecular characterization of a complex of apoptosis-inducing factor 1 with cytochrome c

oxidase of the mitochondrial respiratory chain. Proc Natl Acad Sci U S A. 2021 Sep

28;118(39):e2106950118.

24. Evers F, Cabrera-Orefice A, Elurbe DM, Kea-te Lindert M, Boltryk SD, Voss TS, et al.

Composition and stage dynamics of mitochondrial complexes in Plasmodium falciparum.

Nat Commun. 2021 Dec;12(1):3820.

25. Maclean AE, Bridges HR, Silva MF, Ding S, Ovciarikova J, Hirst J, et al. Complexome

profile of Toxoplasma gondii mitochondria identifies divergent subunits of respiratory

chain complexes including new subunits of cytochrome bc1 complex. PLoS Pathog.

2021 Mar;17(3):e1009301.

26. Ligas J, Pineau E, Bock R, Huynen MA, Meyer EH. The assembly pathway of complex I

in Arabidopsis thaliana. Plant J Cell Mol Biol. 2019 Feb;97(3):447–59.

27. Nepusz T, Yu H, Paccanaro A. Detecting overlapping protein complexes in protein-

protein interaction networks. Nat Methods. 2012 May;9(5):471–2.

28. D'Imprima E, Mills DJ, Parey K, Brandt U, Kühlbrandt W, Zickermann V, et al. Cryo-EM

structure of respiratory complex I reveals a link to mitochondrial sulfur metabolism.

Biochim Biophys Acta. 2016 Dec;1857(12):1935–42.

29. Abdrakhmanova A, Dobrynin K, Zwicker K, Kerscher S, Brandt U. Functional

sulfurtransferase is associated with mitochondrial complex I from *Yarrowia lipolytica* , but

is not required for assembly of its iron-sulfur clusters. FEBS Lett. 2005 Dec

19;579(30):6781–5.

30. Morgner N, Zickermann V, Kerscher S, Wittig I, Abdrakhmanova A, Barth HD, et al.

Subunit mass fingerprinting of mitochondrial complex I. Biochim Biophys Acta BBA -

Bioenerg. 2008 Oct;1777(10):1384–91.

31. The UniProt Consortium, Bateman A, Martin MJ, Orchard S, Magrane M, Agivetova R, et

al. UniProt: the universal protein knowledgebase in 2021. Nucleic Acids Res. 2021 Jan

8;49(D1):D480–9.

32. Belden WJ, Barlowe C. Distinct Roles for the Cytoplasmic Tail Sequences of Emp24p

and Erv25p in Transport between the Endoplasmic Reticulum and Golgi Complex. J Biol

Chem. 2001 Nov;276(46):43040–8.

33. Marzioch M, Henthorn DC, Herrmann JM, Wilson R, Thomas DY, Bergeron JJM, et al.

Erp1p and Erp2p, Partners for Emp24p and Erv25p in a Yeast p24 Complex. Kaiser C,

editor. Mol Biol Cell. 1999 Jun;10(6):1923–38.

34. Paysan-Lafosse T, Blum M, Chuguransky S, Grego T, Pinto BL, Salazar GA, et al.

InterPro in 2022. Nucleic Acids Res. 2023 Jan 6;51(D1):D418–27.

35. Szklarczyk D, Gable AL, Nastou KC, Lyon D, Kirsch R, Pyysalo S, et al. The STRING

database in 2021: customizable protein–protein networks, and functional

characterization of user-uploaded gene/measurement sets. Nucleic Acids Res. 2021 Jan

8;49(D1):D605–12.

36. Knöckel J, Dundas K, Yang ASP, Galaway F, Metcalf T, Gemert GJ van, et al.

Systematic Identification of Plasmodium Falciparum Sporozoite Membrane Protein

Interactions Reveals an Essential Role for the p24 Complex in Host Infection. Mol Cell

Proteomics. 2021;20:100038.

37. Florentin A, Stephens DR, Brooks CF, Baptista RP, Muralidharan V. Plastid biogenesis
in malaria parasites requires the interactions and catalytic activity of the Clp proteolytic
system. Proc Natl Acad Sci. 2020 Jun 16;117(24):13719–29.

38. Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, et al. Fast, scalable
generation of high-quality protein multiple sequence alignments using Clustal Omega.
Mol Syst Biol. 2011 Jan;7(1):539.

39. Guindon S, Dufayard JF, Lefort V, Anisimova M, Hordijk W, Gascuel O. New Algorithms
and Methods to Estimate Maximum-Likelihood Phylogenies: Assessing the Performance
of PhyML 3.0. Syst Biol. 2010 Mar 29;59(3):307–21.

40. Morgenstern M, Peikert CD, Lübbert P, Suppanz I, Klemm C, Alka O, et al. Quantitative
high-confidence human mitochondrial proteome and its dynamics in cellular context. Cell
Metab. 2021 Dec;33(12):2464-2483.e18.

41. Rath S, Sharma R, Gupta R, Ast T, Chan C, Durham TJ, et al. MitoCarta3.0: an updated
mitochondrial proteome now with sub-organelle localization and pathway annotations.
Nucleic Acids Res. 2021 Jan 8;49(D1):D1541–7.

42. Christiansen J, Kolte AM, Hansen T v O, Nielsen FC. IGF2 mRNA-binding protein 2:
biological function and putative role in type 2 diabetes. J Mol Endocrinol. 2009
Nov;43(5):187–95.

43. van Esveld SL, Meerstein-Kessel L, Boshoven C, Baaij JF, Barylyuk K, Coolen JPM, et
al. A Prioritized and Validated Resource of Mitochondrial Proteins in *Plasmodium*
Identifies Unique Biology. Blader IJ, editor. mSphere. 2021 Oct 27;6(5):e00614-21.

44. Barylyuk K, Koreny L, Ke H, Butterworth S, Crook OM, Lassadi I, et al. A
Comprehensive Subcellular Atlas of the Toxoplasma Proteome via hyperLOPIT Provides
Spatial Context for Protein Functions. Cell Host Microbe. 2020 Nov;28(5):752-766.e9.

45. Jansen EJR, Timal S, Ryan M, Ashikov A, van Scherpenzeel M, Graham LA, et al.

ATP6AP1 deficiency causes an immunodeficiency with hepatopathy, cognitive

impairment and abnormal protein glycosylation. Nat Commun. 2016 May 27;7:11600.

46. Wang L, Wu D, Robinson CV, Wu H, Fu TM. Structures of a Complete Human V-

ATPase Reveal Mechanisms of Its Assembly. Mol Cell. 2020 Nov 5;80(3):501-511.e3.

47. Mallilankaraman K, Cárdenas C, Doonan PJ, Chandramoorthy HC, Irrinki KM, Golenár

T, et al. MCUR1 is an essential component of mitochondrial Ca2+ uptake that regulates

cellular metabolism. Nat Cell Biol. 2012 Dec;14(12):1336–43.

48. Brischigliaro M, Cabrera-Orefice A, Sturlese M, Elurbe DM, Frigo E, Fernandez-Vizarra

E, et al. CG7630 is the *Drosophila melanogaster* homolog of the cytochrome *c* oxidase

subunit COX7B. EMBO Rep [Internet]. 2022 Aug 3 [cited 2022 Oct 26];23(8). Available

from: https://onlinelibrary.wiley.com/doi/10.15252/embr.202254825

49. Gabaldón T, Rainey D, Huynen MA. Tracing the Evolution of a Large Protein Complex in

the Eukaryotes, NADH:Ubiquinone Oxidoreductase (Complex I). J Mol Biol. 2005

May;348(4):857–70.

50. Agip ANA, Chung I, Sanchez-Martinez A, Whitworth AJ, Hirst J. Cryo-EM structures of

mitochondrial respiratory complex I from Drosophila melanogaster. eLife. 2023 Jan

9;12:e84424.

51. Molina-Granada D, González-Vioque E, Dibley MG, Cabrera-Pérez R, Vallbona-Garcia

A, Torres-Torronteras J, et al. Most mitochondrial dGTP is tightly bound to respiratory

complex I through the NDUFA10 subunit. Commun Biol. 2022 Jun 23;5(1):620.

52. Elurbe DM, Huynen MA. The origin of the supernumerary subunits and assembly factors

of complex I: A treasure trove of pathway evolution. Biochim Biophys Acta BBA -

Bioenerg. 2016 Jul;1857(7):971–9.

53. Crozier TWM, Tinti M, Larance M, Lamond AI, Ferguson MAJ. Prediction of Protein

Complexes in Trypanosoma brucei by Protein Correlation Profiling Mass Spectrometry

and Machine Learning. Mol Cell Proteomics. 2017 Dec;16(12):2254–67.

54. Havugimana PC, Hart GT, Nepusz T, Yang H, Turinsky AL, Li Z, et al. A Census of

Human Soluble Protein Complexes. Cell. 2012 Aug;150(5):1068–81.

55. Feldmann AM, Ponnudurai T. Selection of Anopheles stephensi for refractoriness and

susceptibility to Plasmodium falciparum. Med Vet Entomol. 1989 Jan;3(1):41–52.

56. Ponnudurai T, Lensen AHW, Van Gemert GJA, Bensink MPE, Bolmer M, Meuwissen

JHETh. Infectivity of cultured *Plasmodium falciparum* gametocytes to mosquitoes.

Parasitology. 1989 Apr;98(2):165–73.

57. Wittig I, Braun HP, Schägger H. Blue native PAGE. Nat Protoc. 2006 Jun;1(1):418–28.

58. Rodenburg RJT. Biochemical diagnosis of mitochondrial disorders. J Inherit Metab Dis.

2011 Apr;34(2):283–92.

59. Steinegger M, Meier M, Mirdita M, Vöhringer H, Haunsberger SJ, Söding J. HH-suite3

for fast remote homology detection and deep protein annotation. BMC Bioinformatics.

2019 Dec;20(1):473.

60. Amos B, Aurrecoechea C, Barba M, Barreto A, Basenko EY, Bażant W, et al.

VEuPathDB: the eukaryotic pathogen, vector and host bioinformatics resource center.

Nucleic Acids Res. 2022 Jan 7;50(D1):D898–911.

61. Berardini TZ, Reiser L, Li D, Mezheritsky Y, Muller R, Strait E, et al. The arabidopsis

information resource: Making and mining the "gold standard" annotated reference plant

genome: Tair: Making and Mining the "Gold Standard" Plant Genome. genesis. 2015

Aug;53(8):474–85.

Supporting information

1030

Figure S1. Migration of clustered *H.sapiens* EMP24 proteins in three human fibroblast

complexome profiles.

Figure S2. Enzyme activities in C15orf61-KO and WT HEK cell lines.

Figure S3. Alignment of the conserved C-terminus of human ATP6AP12 and its predicted

orthologs

Figure S4. Alignment of COX7B with recently identified *D. melanogaster* and ortholog and a

clustered *A. stephensi* protein.

Figure S5. Migration of four complex I proteins in various complexome profiles.

Figure S6. Identification performance of protein complex members with rank biased overlap

for a range of "p" parameter values.

Table S1. Overview of 15 inspected *A. stephensi* subclusters resulting from CompaCt

analysis.

Table S2. Overview of complexome profiles used in main analysis

Table S3. Overview of proteomes used for sequence-based homology analysis

Supplemental methods. Complexome profiling of anopheles material and generation of

KO HEK293T cells.

Supplemental File S1. Output clusters from CompaCt analysis of eukaryote complexome

profiles.

Supplemental File S2. Annotation of inspected clusters and cluster members.

44