1
2   Title:
3   A Gene-expression Module Identifies Circulating Immune Cells with Enhanced Recruitment to Sites of
4   Inflammation
5
6   Debajyoti Sinha*[1], Thomas Laurent*[1], Alexis Broquet[1], Cynthia Fourgeux[1], Thibault Letellier[2], Gaelle
7   Tilly[1], Sarah Bruneau[1], Simon Ville[1,2], Laurence Bouchet-Delbos[1], Julien Brancherau[1,2], Clarisse
8   Kerleau[2], Sophie Brouard[1], Gilles Blancho[1,2], Magali Giral[1,2], Regis Josien[1,2], Richard Danger[1], Antoine
9   Roquilly[1,3], Nicolas Degauque[1], Jeremie Poschmann[1,4]
10
11
[1]Nantes Université, CHU Nantes, INSERM, Center for Research in Transplantation and Translational
Immunology, UMR 1064, ITUN; F-44000, Nantes, France.
[2] CHU Nantes, Nantes Université, Institut de Transplantation Urologie Néphrologie (ITUN), Service de
Néphrologie et Immunologie Clinique, 44000 Nantes, France.
[3] Nantes Université, CHU Nantes, INSERM, Anesthesie Réanimation, CIC 1413, F-44000 Nantes,
France
[4] Corresponding author (orcid : 0000-0002-9613-5297), 30 Boulevard Jean-Monnet, CR2TI, 44093
Nantes, France, jeremie.poschmann@univ-nantes.fr
* contributed equally

21
22
23
Abstract:

Circulating immune cells are critical mediators of inflammation upon recruitment to tissues, yet how
their gene expression state influences this recruitment is not well understood. Here, we report
longitudinal single-cell transcriptome profiling of peripheral blood mononuclear cells in patients
undergoing kidney transplantation rejection. We identify a novel gene expression module, termed
ALARM (early activation transcription factor module), associated with transcriptional regulation,
homing, and immune activation across multiple immune cell types. Circulating cells expressing this
module are significantly reduced in patients experiencing graft rejection, a finding confirmed in a pig
model of acute kidney transplantation rejection. Correspondingly, module expression is markedly
increased in kidney grafts undergoing rejection, indicating preferential recruitment of ALARM-
expressing cells to the inflamed tissue.
Within this module, we identify the receptor CXCR4 and its ligand CXCL12, expressed in the graft, as a
likely mechanism for recruitment. In vitro transwell assays combined with scRNA-seq reveal that this
CXCR4-CXCL12 interaction is critical for T cell migration and upregulation of CD69, an early activation
marker, and is accompanied by a metabolic switch towards glycolysis. Further exploration of publicly
available transcriptomic data demonstrates that this module is generally expressed in healthy
individuals and is strongly associated with responses to infection, including SARS-CoV-2 infection. This
finding is further supported by experiments in a pneumonia mouse model, which confirm the
recruitment of CXCR4-expressing T cells during lung infection. Moreover, we find that module
expression is predictive of immune-mediated diseases.
In summary, we have identified a key gene expression module in circulating immune cells that
orchestrates their preferential recruitment to inflamed tissues, metabolic reprogramming, promoting
tissue residency and effector functions. These insights advance our understanding of immune cell
recruitment and activation mechanisms in transplant rejection and infectious diseases, with potential
implications for therapeutic interventions.

49
50
51   **Introduction:**

Circulating immune cells are critical to be recruited to the site of inflammation, infection, and cancer. This compartment and particularly the peripheral blood mononuclear cells (PBMCs) thus offer an attractive resource for precision medicine as in a single experiment, diverse cell types, including the CD4+ T cells, CD8+ T cells, B cells, NK cells and monocytes are profiled. Especially since the introduction of single cell RNA-seq (scRNA-seq), profiling PBMCs has been highly successful in identifying gene expression signatures and cell-types associated with immune-related diseases[1–4]. For example, a monocyte signature associated with sepsis was discovered in circulating cells [1]. Recently, a study on systemic lupus erythematosus revealed gene expression changes with disease state and genetic variation [2]. In addition, distinct Covid-19 studies revealed signatures associated to infection and disease severity in blood [5,6].

Transcriptomics profiling enables the characterization of genes groups (i.e., modules) which perform critical cellular functions such as maintaining a cell identity, homeostasis & metabolism and respond to external signals. Notably, in circulating immune cells we have previously shown that monocytes express a gene module associated with Herpes Simplex virus reactivation after traumatic brain injury [7]. While gene expression programs of circulating immune cells are likely to be distinct from the same cells which migrated into the tissue, identifying modules in circulating cells may reveal early immune activation programs or modules associated with homing and migration. For example, in a previous study we identified large gene regulatory and gene expression alterations in circulating monocytes during active Tuberculosis which improved pathogen clearance for these cells [8].

Leveraging on scRNA-seq and the availability of recent module identification approaches tailored for single cell transcriptomics [9] we aimed to identify gene expression programs associated with kidney transplantation rejection. Currently, rejection status is monitored in clinical practice by analyzing metabolites in blood and urine, such as creatinine, to assess renal function. The diagnosis is then confirmed through pathologic examination of kidney biopsies[10]. However, metabolite monitoring is not specific to rejection and can be approximative. Pathologic analysis of kidney biopsies remains the primary diagnostic tool, offering reliable results. However, despite minimal risk to patients, it is an invasive procedure that cannot be performed regularly. Thus identifying gene expression signatures in circulating cells may improve precision medicine diagnostics of kidney rejection [11]. In addition, acute and chronic rejection are characterized by the infiltration of immune cells into the graft, which then mediate an inflammatory response in the tissue ultimately leading to the rejection of the graft. Blood thus constitutes an easily accessible compartment to identify gene expression modules associated with homing and early activation. Two archetypes of rejection are prominent according to the Banff classification[10], the antibody mediated rejection (ABMR) and T-cell mediated rejection (TCMR), which can also arise in a mixed form. In both cases, immune cell infiltration into the graft occurs via the bloodstream through either donor specific antibodies binding to the graft endothelium in ABMR, or cytokine and homing signals in TCMR[12].

In this study, our aim was to identify putative modules in circulating cells which may be associated to kidney transplantation rejection. We profiled a longitudinal patient cohort consisting of 3 stable patients, 3 TCMR and 3 ABMR patients at 0-month, 3 month and 12 months after transplantation or when rejection occurred. The PBMCs were collected at the same time of graft biopsy, allowing us to characterize relationship of graft rejection status with gene expression modules of peripheral immune cell-types. We identified a module associated with transcriptional regulation and early activation in the blood and used a pig-transplantation model to validate its association with rejection status. Further characterization of this module was carried out in transcriptomics data from over 1500 kidney biopsies revealing a cytokine-receptor interaction between the graft and circulating cells, respectively. Finally, we demonstrate that this module is not specific to graft rejection but is implicated and predictive of a variety of immuno-pathologies.

Results:
Single cell transcriptome analysis of circulating immune cells in a longitudinal kidney
transplantation cohort
In order to identify gene expression changes in circulating immune cells during kidney transplantation
rejection, we used scRNA-seq on PBMC isolated from 3 patients with stable allograft function (STA) for
which no sign of rejection was observed clinically after more than a year of follow-up, 3 antibody
mediated rejection (ABMR) patients and 3 T-cell mediated rejection (TCMR) patients (Figure 1A, Table
1). The patients were selected based on their treatment, age, sex, and collection time. (See Table 1).
112

Table 1: Clinical summary of the cohort composition.

	Name	Age	Sex	Treatment *	Rejection time (in days)	HLA ** mismatches	Induction Therapy
Batch 1	STA1	35	Male	Tacrolimus, MMF, Corticoids	—	5	Depleting
	TCMR1	70	Male	Tacrolimus, MPA, Corticoids	244	5	Depleting
	ABMR1	51	Female	Tacrolimus, MMF, Corticoids	374	2	Depleting
	TOL1	75	Male		—	4	Depleting
Batch 2	STA2	69	Male	Tacrolimus, MMF, Corticoids	—	5	Non − depleting
	TCMR2	35	Male	Tacrolimus, MPA, Corticoids	369	3	Depleting
	ABMR2	56	Male	Tacrolimus, MPA, Corticoids	427	4	Depleting
	TOL2	45	Male		—	4	NA
Batch 3	STA3	25	Female	Tacrolimus, MPA, Corticoids	—	5	Depleting
	TCMR3	24	Male	Tacrolimus, MMF, Corticoids	213	3	Non − depleting
	ABMR3	61	Female	Tacrolimus, MPA, Corticoids	365	2	Depleting
	TOL3	39	Female		—	0	NA

Mean Age	Sex Ratio
48.75	0.66

* Antiproliferative treatments: MMF=Mycophenolate mofetil, MPA=Mycophenolic Acid
** HLA mismatches: number of mismatches on the A, B and DR loci (0-6)

113
114
115
116
For each patient, three time points were profiled; T0, at the kidney transplantation, T1 at 90-150 days
after the transplantation and T2, which was sampled at the time of rejection for ABMR and TCMR or
at 1 year for STA patients after transplantation (Figure 1A). In addition, PBMCs from 3 kidney
transplantation patients who maintained allograft function in the absence of immunosuppression (i.e.,
operational tolerant (TOL)) were included in the cohort (Table 1).
Rejection status was defined by clinical pathology assessment of kidney biopsies performed at T1 and
T2 for all patients. ABMR and TCMR were defined by pathology biopsy assessment at time T2 whereas
patients were determined as STA when they which no sign of rejection in biopsies at T1 and T2. To
minimize scRNA-seq-related experimental variation, we performed CITE-seq (Cell Hashing)[13] using
hashtag oligo-conjugated antibodies (HTO) to label each patient and time point separately and then
pooled 10 samples (1 ABMR, 1 TCMR and 1 STA patient across T0, T1 & T2 and 1 TOL patient) into a
single experiment. We thus generated the complete transcriptomic data in three balanced batches
(Figure 1B). After removing doublets, data cleaning, normalization, and batch correction, we obtained

50,507 cells across the three batches (see Methods). Cells were automatically annotated using
Azimuth[14,15] and manually validated for cell-type specific markers (Figure 1C and Supplemental Figure
S1A). Cell-type proportions varied minimally between conditions when compared to PBMCs from two
separate cohorts of healthy volunteers (HV) from publicly available scRNA-seq data [1,5] (Supplemental
Figure S1B). For example, NK cells were significantly decreased in the stable and rejection conditions
as compared to HV. However, we note that there were also significant differences in the HV (e.g., CD14
and CD16 monocytes) from the two distinct control cohorts indicating that this may be due to
individual variation. We then inspected the distribution of cells across patients and by time points
(Supplemental Figure S1C) and across disease states (Figure 1D). Neither of these variables formed
unique clusters suggesting that the clusters were driven by cell-type specific expression rather than by
condition or cellular states as was also observed in other PBMC studies in patient cohorts [1,2,5]. We then
explored whether the clusters were affected by merging the three experimental batches (Figure 1E).
As no batch effects were apparent through visual inspection, we used the KBET metric to quantitatively
assess potential batch effects. KBET evaluates whether cells from different batches are clustering
together in shared neighborhoods (i.e., clusters) (Supplemental Figure S1D)[16]. The acceptance rate of
the KBET for the complete data set was 0.969, indicating that batch integration was successful. In
summary, the pooling strategy and subsequent bioinformatics analysis resulted in a robust dataset of
50,507 cells to be analyzed for time and disease state specific gene expression.
148
Gene co-expression analysis identifies a module associated to rejection state
To identify modules, ie. co-expressed groups of genes, we used gene co-expression analysis across all
three batches independently (Figure 2A). This approach was chosen to avoid potential signal
alterations induced during the batch correction step. We applied consensus non-negative matrix
factorization (cNMF)[9] to identify gene expression programs which may either be associated to cell-
type specific gene expression programs or to cellular activity (see methods). Nine overlapping modules
were identified which were evenly distributed across the three batches (Figure 2B, Supplemental
Figure 2A). These modules mostly revealed cell-type specific expressions, notably three of these
modules were associated to monocytes (Mod_1-Mod_3), and two modules were mostly expressed in
a specific cell type such as B cells and pDC cells (Mod_4 and Mod_5 respectively) (Figure 2C). Three
modules were enriched for the CD4 & CD8 T lymphocytes (Mod_6) and/or NK cells (Mod_7, Mod_8,
Figure 2C). Interestingly, the Mod_9 was expressed in all cell types, but with notable higher expression
in B cells, T cells, pDCs and NK cells as revealed by its module score (Figure 2C, Figure 2D).
Next, we aimed to independently validate the co-expression of the 9 modules (Supplemental file 1).
To achieve this, we quantified the gene co-expression using the pair-wise Pearson correlation
coefficient R. As the module was identified using cNMF, Pearson correlation thus acts as an
independent evaluation of gene co-expression. We note that this approach is inherent to weighted
gene co-expression analysis (WGCNA), a prominent method to identify modules in bulk and single cell
transcriptomics[17]. We compared the average Pearson correlation per cell for each module to the same
number of randomly picked genes in the same cells. The eight cell type specific modules showed robust
and significant correlation between module genes, and this correlation was strongest in the cell-types
to which they were associated (Supplemental Figure S2B). The cell-activity module Mod_9 was also
significantly correlated in all cell types (Supplemental Figure S2C). Of note, randomly chosen genes
picked 1000 times revealed a Pearson correlation of 0 in all cell-types indicating that unrelated genes
typically do not correlate with each other (Supplemental Figure S2B, S2C). Therefore, the modules
identified above are robustly co-expressed, as inferred by the two most prominent module detection
methods.
To test whether this cell-activity module or any of the cell-specific modules were associated to disease
state, (i.e., rejection or stable) and if it would vary throughout time, we estimated the module score
for each cell type and calculated the trend of the module score longitudinally in each cell-type (Figure
2E). A positive or negative slope thus indicates whether a module changes over time across the stable,
humoral & cellular rejection. Indeed, Mod1 and Mod2 showed a significant positive trend in both

rejection conditions (ABMR and TCMR) but not in the STA condition. Interestingly, the activity module
(Mod_9) showed a significant negative trend (regression β value) in the rejection conditions but not in
the stable condition in multiple cell types (B cells, CD4 and CD8 T cells and NK cells). Further inspection
of this negative trend was carried out by displaying the module score of each patient separately in the
form of a combined violin plot (Figure 2F). These module scores show that there was indeed a
reduction in ABMR patients, while in TCMR patients the module score followed a U-shape, reduced
more during T1 and increased again at T2. Interestingly, the 3 stable patient's module scores remained
consistent across time in NK, CD4 and CD8 and B cells. These results signify that Mod_9 expression is
associated to rejection state in a time dependent manner.
190
Discovery of the early activation, transcription factor module (ALARM)
To explore the function of the 61 genes found in Mod_9 we first investigated whether it was enriched
for ribosomal, proliferation or cell cycle genes using SEURAT-based list [18]. We did not observe any
genes involved in these cellular processes (data not shown). We thus explored the genes within this
module by performing gene ontology analysis. Enrichment of the module genes was quantified for
molecular function (MF) and biological processes (BP) compared to the combined set of 4000 most
variable genes from the three batches (Figure 2G). MF could be associated to 24 genes and was
significantly enriched (FDR <0.05) for helicase activity, ubiquitin-like protein binding, RNA catalytic
activity, transcription activator activity and ribonucleoprotein binding. The BP (20 genes) was
associated to NF-κB signaling, response to peptide and oxidative stress, regulation of RNA metabolic
processes and viral gene expression. Of the 61 genes in the module, 56 were annotated in the GSEA
database and 30 of these genes were linked to gene ontology enrichment. This indicates that Mod_9
is likely to be involved in multiple molecular functions associated with transcription, mRNA process
and ubiquitination. The BP suggested involvement in response to immune conditions (i.e., viral gene
expression, NF-κB signaling, oxidative stress). This result was further supported by the 5-fold
enrichment of Transcription factor genes in this module (OR: 4.9; Fisher-Test P-value $2.5e^{-5}$), such as
the AP-1 complex (*JUN, JUND, FOS*), *REL* (NF-κB subunit), *MAFF* and *NR4A2* (see methods,
Supplemental file 1). Further manual examination exposed the early activation marker CD69, a cell
surface type II lectin. This receptor was described to be rapidly expressed at the membrane in T cells
upon TCR activation[19]. In addition, CD69 has been described as a marker of tissue retention of T cells
[20–22]. Interestingly, *CD69* gene promoter is controlled by AP-1 TF complex and NF-κB, both of which are
also members of this module[23,24]. We also found the cell surface marker CXCR4 in this module, a
chemokine receptor known to play a role in recruiting CXCR4 positive cells to the kidney after an
ischemic injury via the chemokine CXCL12[25]. The role of the CXCR4/CXCL12 axis in kidney rejection is
still unclear[26] but an elevated expression of CXCL12 has been described in chronic kidney rejection[27]
suggesting it may act as a chemotactic signal to recruit immune cells in the inflamed tissues. In
summary, the Mod_9 module comprises genes implicated in the response to stress, mRNA processing,
early activation, and tissue-homing. For clarity, we named this module ALARM, which stands for for
eArLy activation trAnscription factoR Module.
220
Circulating ALARM cells are recruited to the graft during acute graft rejection in a pig kidney
transplantation model
We found that circulating cells expressing ALARM are altered in ABMR and TCMR in a timely fashion
(Figure 2). To independently validate this observation, i.e., whether ALARM high expressing cells are
depleted in the circulation during kidney transplantation rejection, we took advantage of an acute
rejection pig kidney transplantation model (Figure 3A). We decided to use this pig kidney
transplantation model as they share anatomical, physiological and genetic similarities to human and
offer the advantage to have well defined swine leukocyte antigen (SLA) genotypes [28]. Two SLA-
mismatched pigs were subjected to kidney transplantation, keeping one of their own kidneys (see
methods). This model typically results in an acute TCMR within a few days after transplantation, as no
immunosuppressive treatment is given (Figure 3A). Kidney biopsies and PBMCs were collected daily
before and after the transplantation. Microscopical analysis of the biopsies at D2, D4 and D6 indicated

a time-dependent infiltration of immune cells, culminating at day 6 (Figure 3B). This infiltration was
quantified in three areas (excluding glomeruli) of each biopsy time point (see methods, Supplemental
Figure S3A). Cell counts drastically increased from day 2 to D6, indicating a continuous immune cell
accumulation over time (Figure 3C). We noted that the second pig did not display any signs of rejection,
possibly due to early arterial ischemia of the transplanted kidney and it was thus discarded from the
subsequent analysis. PBMCs collected at D0, D2, D4 and D6, were pooled in a single scRNA-seq
experiment (see methods), resulting in a total of 4,411 annotated cells across cell-types and time-
points (Figure 3D and Supplemental Figure S3B). We found that cellular proportions within the PBMC
compartment drastically changed from D0 to D2, characterized by a drastic increase in monocyte
proportions concomitant with a reduction of B, CD4+ and CD8+ T cells (Figure 3E). This suggests that
the lymphocytes are rapidly recruited to the kidney graft and accumulate there, as demonstrated by
the cellular invasion observed in the biopsies at the same time (Figure 3C). To test whether the decline
of blood lymphocytes is accompanied by a reduction of ALARM high expressing cells, we quantified
ALARM expression across the time-points in CD4+, CD8+, B cells and NK cells (Figure 3F). Interestingly,
as soon as D2 the levels of ALARM expression drastically decreased and remained low in the blood
until sacrifice of the animal (D6). Taken together, this controlled time-course experiment reveals a
drastic immune cell infiltration in the graft associated with the depletion of ALARM high expressing
cells in the circulation. This result mirrors the reduction of circulating ALARM cells observed during the
kidney transplantation rejection in the human cohort (Figure 2). It is thus probable that ALARM high
expressing cells are readily circulating in healthy condition. Upon the kidney graft transplantation
ALARM high expressing cells are then preferentially recruited to the organ to mediate the immune
response.
255
ALARM gene expression increases in kidney grafts undergoing rejection
To further support the hypothesis that ALARM high-expressing cells are recruited to the graft during
rejection from the bloodstream, we evaluated the expression of ALARM genes in graft biopsies from
kidney transplantation patients. For this, we used a previously published Canadian transcriptomics
analysis of 569 transplant biopsies collected from 13 clinical sites and with a patient classification of
STA, TCMR, ABMR and mixed rejection (TCMR and ABMR)[29]. In parallel, we exploited a second similar
Belgian transcriptomics study performed on kidney biopsies in 224 patients who were either stable
(168 patients) or undergoing ABMR[30]. After precleaning and QC controls of the available microarray
data (Methods), we quantified the ALARM gene expression in each cohort separately (Figure 4A). The
ALARM score was consistently and significantly elevated in all three rejection cases compared to stable
biopsies. Similarly, in the second study, ABMR samples showed a significant increase in ALARM
expression, regardless of the presence of donor-specific antibodies. Analysis of the ALARM genes
ranked by expression, further revealed that this score is driven by the upregulation of a large fraction
of the ALARM genes, including *CD69*, *CXCR4*, *JUN* and *IRF1* in both cohorts (Figure 4B). Quantification
of the ALARM module across rejection and stable revealed a significant upregulation of ALARM
expression in both data sets (Figure 4C). These results indicate that ALARM genes are significantly
increased during graft rejection across over 793 biopsies in two distinct studies. Given that graft
rejection is defined by immune cell infiltration and that circulating ALARM high cells are depleted at
the same time, it is possible that ALARM expressing cells are preferentially recruited to the graft to
mediate the rejection. To investigate how circulating cells could be preferentially recruited to the graft,
we explored the cytokine expression in the graft and receptor expression in the circulating cell subsets.
We first identified all possible ligand-receptor pairs and then tested whether these pairs were
differentially expressed between rejection and stable status. We found 10 differentially expressed
cytokines in the graft pairing with 7 receptors upregulated in circulating immune cells in both cohorts
independently (Figure 4D). The most prominent receptor was CXCR4, expressed in CD4, NK, CD8 and
B cells. As mentioned above, CXCR4 is a member of ALARM genes, indicating a likely mechanism of
signaling from the graft via CXCL12 and recruitment of ALARM cells expressing *CXCR4*. This cytokine
receptor pair has been previously described as a homing mechanism in various distinct tissues, such as
bone marrow[31,32] and in cancer[33]. To test the relationship of this interaction, we quantified *CXCL12*

expression in the graft biopsies stratified by ALARM expression (low (<25%), mid (25-75%), high
(>75%)) and found that there was a significant increase in CXCL12 expression in the ALARM high group
in both cohorts (Figure 4E). In summary, these results show that ALARM is increased during rejection
in the graft, which strengthens the notion that circulating cells with high ALARM expression are
preferentially recruited to the kidney graft via the homing signaling axis of CXCL12 and CXCR4.
290
Analysis of ALARM Module Expression in an In Vitro Transwell Assay
To experimentally verify the observed CXCL12-mediated recruitment, we employed an *in vitro*
transwell assay using a cytokine gradient of CXCL12. This assay consists of a membrane covered with
primary microvascular endothelial cells (HDMEC), allowing for the comparison of unstimulated cells,
cells in direct contact with CXCL12, and those migrating based on the CXCL12 gradient (Figure 5A).
Flow cytometry quantifications of T cells indicate that CXCL12 significantly induces the recruitment and
migration of T cells, observed 4 hours post-deposition (Figure 5A; p=0.0286). The composition of naïve,
central memory (CM), effector memory expressing CD45RA (TEMRA), and effector memory (EM) T cells
remained similar after migration and in response to CXCL12 alone (Figure 5B, Supplemental Figure
S4A). This suggests that all T cell subsets are equally attracted and migrate in response to a CXCL12
gradient. We further analyzed the expression of several T cell surface markers, focusing on CD69 as a
component of the ALARM module, CD25 (an activation marker), CD49a (a tissue residency marker),
and CD95 (an apoptosis marker) (Figure 5C). Interestingly, CD69 increased slightly with CXCL12
addition but was even more highly upregulated in migrated cells, while CD25 expression did not
change, decoupling the role of CD69 in early activation from its role in tissue residency[34]. CD49a and
CD95 levels remained unchanged, indicating that CXCL12 in combination with migration specifically
induced the extracellular display of CD69. It is noteworthy that CD69 expression depends on both
CXCL12 and the direct contact with HDMEC cells (Supplemental Figure S4B).
To assess whether CXCL12-induced migration altered the T cells' response to immune stimuli, CXCL12
exposed or transmigrated cells were purified and restimulated polyclonally for an additional 24 hours
(Figure 5D). Migrated cells showed a significantly increased expression of IFN-γ in both CD4+ and CD8+
T cells in contrast to TNF-α, granzyme B (GZMB) or granulysin (GNLY). The increased expression of IFN-
γ suggests that migration via the CXCL12 gradient may enhance the effector functions of T cells but
that other signals are needed to induce cytotoxic mechanisms in this model. The observed increase in
CD69 membrane display indicates a functional role for the ALARM module, which enables T cells to
acquire "new functions" in the tissue i.e. to establish residency via CD69 expression and increased IFN-
γ expression.
To further investigate the impact of CXCL12 signaling on T cell migration and the role of ALARM module
expression, we performed scRNA-seq using the transwell assay under three distinct conditions (Figure
6A). In the first condition (CXCL12-), T cells were placed below the transwell membrane without any
exposure to CXCL12. In the second condition (CXCL12+), T cells were placed below the transwell
membrane in direct contact with CXCL12. The third condition, (transmigrated), involved placing T cells
above the membrane, which were then collected from below the membrane after migrating in the
presence of CXCL12. This setup allowed us to assess the transcriptional changes associated with T cell
migration in response to CXCL12 (Figure 6A). Notably, all T cell subtypes identified by gene expression
were found in the three conditions (Supplemental Figure S4C).
To more accurately quantify gene expression changes and minimize the impact of subtype differences,
we focused on the most abundant subsets, namely naïve CD4 and CD8 T cells. These naïve T cells
clustered into four distinct groups, primarily due to differences in ribosomal gene expression and the
annotation of CD4 versus CD8 cells (Figure 6B). Additionally, there was a clear separation between
CXCL12- T cells and those exposed to CXCL12 or that had transmigrated, indicating significant
underlying gene expression differences. This separation was further evidenced by distinct changes in
the expression of ALARM module genes, with the CXCL12- condition showing markedly lower
expression patterns compared to the CXCL12+ and transmigrated groups (Figure 6B). Next, we
investigated specific gene expression alterations which would increase or decrease from CXCL12-,
CXCL12+ to the transmigrated condition (see methods).

In addition to ALARM module genes (Supplemental Figure S4D), several others exhibited notable
changes in expression across the different conditions (Figure 6C, Supplemental file 1). HSP90AB1, a
member of the HSP90 family of chaperone proteins, which is crucial for stabilizing proteins involved in
cell survival and esponses[35] was significantly upregulated in both CD4 and CD8 T cells. This suggests
that HSP90AB1 may play an important role in enhancing the functional stability of proteins required
for T cell migration and adaptation during CXCL12 stimulation. Interestingly, two genes with roles in
cell migration, VIM[36] and STK38[37] (serine/threonine kinase 38), also showed differential expressions in
both cell types. VIM, a key regulator of cytoskeletal organization that promotes cell motility[36], was
upregulated in migrated cells, aligning with its role in facilitating the cytoskeletal rearrangements
necessary for migration. On the other hand, STK38 was upregulated predominantly in the CXCL12-
condition. To better understand the roles of all differentially expressed genes, we grouped them by
function (Supplemental file S1, Figure 6D). We found gene sets involved in immune response,
migration, cytoskeleton, adhesion, stress response and metabolism were gradually upregulated in the
CXCL12+ and migrated conditions in both CD4 and CD8 T cells. Conversely, gene sets associated with
structural organization, gene regulation, and membrane transport were downregulated. These
findings suggest that CXCL12 signaling, and migration induce profound metabolic and functional
changes in T cells, preparing them for new roles that require increased energy and biosynthetic
demands. Given the prominent upregulation of metabolic pathways, we further investigated the
metabolic reprogramming that accompanies T cell migration and activation in response to CXCL12. To
achieve this, we performed a comprehensive metabolic pathway analysis on T cells using Compass, an
algorithm designed to characterize the metabolic state of cells by integrating single-cell RNA-Seq data
with flux balance analysis[38]. This in silico approach allows us to infer the metabolic status of individual
cells based solely on transcriptomic data, providing insights at single-cell resolution. The analysis
revealed significant upregulation in several metabolic pathways, notably glycolysis/gluconeogenesis,
phosphatidylinositol signaling, and amino acid metabolism in response to CXCL12 and migration
(Figure 7A). To visualize the overall metabolic differences between the conditions, we performed PCA
on the Compass score matrix, which quantifies the metabolic state in each cell (see methods)[38]. The
PCA results showed clear clustering of samples according to their condition, with non-migrated T cells
forming a distinct cluster separate from CXCL12+ and migrated groups (Figure 7B). This separation
underscores the significant impact of CXCL12-induced migration on the metabolic state of T cells.
Because the metabolic shift towards glycolysis is an essential hallmark of T cell activation[39] we focused
on the glycolysis and gluconeogenesis pathways to investigate the gene expression changes involved
in these processes (Figure 7C). Key glycolytic enzymes, such as glucose-6-phosphate isomerase and
pyruvate dehydrogenase, were significantly upregulated in migrated T cells compared to controls. To
validate the transcriptomic findings experimentally, we performed flow cytometry analyses to assess
glucose uptake and the expression of glycolytic enzymes. We measured protein levels of glucose
transporter 1 (GLUT1) and lactate dehydrogenase A (LDHA), a key glycolytic enzyme that converts
pyruvate to lactate for rapid ATP production under anaerobic conditions. Although the increase in
GLUT1-positive cells post-transmigration was not statistically significant (p = 0.68) (Figure 7D), LDHA
expression showed a significant increase in migrated T cells compared to controls (p = 0.019) (Figure
7D). Additionally, uptake of the fluorescent glucose analog 2-NBDG was significantly elevated in
migrated cells (p = 0.031) (Figure 7D), indicating enhanced glucose metabolism. These results confirm
an increased glycolytic activity observed in migrated T cells, consistent with the metabolic
reprogramming identified in our pathway analysis. In summary, our results demonstrate that T cells
migrating in response to CXCL12 undergo functional reorganization, enabling their transition from
circulating cells to active immune responders at sites of inflammation or tissue injury.
383
ALARM is expressed in healthy individuals and variation is associated with infectious disease

Up to this point, we have examined the role of ALARM cells primarily in the context of kidney rejection
and stable kidney transplantation patients. However, the findings from the Transwell assay, using
healthy donor cells in an in vitro model, suggest that ALARM cells may play a broader role beyond
kidney-specific contexts. Therefore, we investigated whether the ALARM module is expressed across
a healthy population to assess its broader function. For this, we explored a publicly available scRNA-
seq data set of ~25,000 PBMC from 45 healthy volunteers (HV) for ALARM expression [40]. The data was
generated from the *LifeLines DEEP* cohort in Netherland, ranges in age from 20 to 79 and contained
46,6% female individuals and was described to be healthy time of collection as estimated by two
general practitioner visits [41]. The ALARM gene expression was prominent in all cell types as shown by
its module score suggesting that ALARM is generally expressed in HV (Supplemental Figure S5A). We
note that age and sex did not result a significantly different expression of the ALARM (Supplemental
Figure S5B). Since this module was generally expressed in HV, we next asked whether it may be
involved in other disease conditions than transplantation rejection, but which implicate the
recruitment of circulating immune cells to specific tissues. To test this hypothesis, we exploited a
publicly available scRNA-seq data on PBMCs in which healthy individuals were intravenously injected
with the endotoxin lipopolysaccharide (LPS), a component of the cell wall of Gram-negative bacteria [5].
LPS in the bloodstream causes an immediate systemic release of a variety of inflammatory mediators,
a fever and a rapid but transient leukopenia [42,43]. This experiment is thought to mimic an acute systemic
inflammatory response (SIRS) [5], and thus provides an ideal proxy of how ALARM expressing cells are
responding to LPS-induced SIRS. We used the preprocessed available scRNA-seq data which contained
the 0-time point (10 HV), 90 minutes (6 HV) and 10h after the LPS injection (6 HV) and first evaluated
how the cellular proportions changed over time. We measured the ALARM score in the three
conditions across the CD4, CD8, NK and B cells (Figure 8A). There was rapid and significant reduction
of ALARM high expressing cells as soon as 90 minutes and which further decreased until 10h after the
LPS injection (Figure 8A). This decrease was continuous within three individuals for which both
timepoints were available, indicating that ALARM cells are reduced in a time dependent manner
(Figure 8B). This drastic change of ALARM expression in such a short time frame suggests that cells
which highly express this module rapidly egress from circulation, directly contributing to the transient
leukopenia observed upon LPS injection. This is also consistent with the transient leukopenia
associated with LPS i.v. injection. Furthermore, this response to LPS which is thought to be initiated
via TLR4 receptor signaling expressed on circulating monocytes [44,45] signifies that ALARM is not solely
implicated in transplantation rejection or kidney immune cell invasion but appears to also be involved
in the inflammatory response to endotoxin.
Next, we investigated how ALARM may regulate when the site of inflammation is localized to a single
organ as in urinary tract infection. For this we explored a publicly available PBMC scRNA-seq dataset
which contained patients with leukocyte infiltrating urinary tract infection (UTI). We chose this
condition as the data was generated on patients which presented a localized infection with infiltrating
leukocytes. The study also provided results of HV and sepsis patients, notably patients with
bacteremia, i.e. bacterial presence in the blood. [1] (Figure 8C). The bacteremia patients were thus also
used for comparison since this condition reflects a generalized or systemic infection which is distinct
from a localized infection such as UTI (Figure 8C). Interestingly, we found that ALARM expression was
reduced in circulating cells in UTI patients (except B cells), consistent with the recruitment of
leukocytes to the tissue. In contrast, in patients with bacteremia, the ALARM cells accumulated in the
circulation, indicating that under this condition ALARM expressing cells may not be recruited to a
specific tissue. This contrasted with the response to LPS iv which is also thought to engender a systemic
response (see discussion).
To further explore the role of ALARM in response to localized infection, we analyzed two scRNA-seq
datasets generated from Covid-19 patients [5,46]. The aim was to explore the dynamics of ALARM cells in
the circulation in comparison to the lung. For this, we compared ALARM expression in PBMCs with

ALARM expression in broncho-alveolar lavage fluid (BALF). The two separate original studies stratified
the patients by healthy, moderate, and severe Covid-19 disease and we used this stratification to
compare the ALARM module expression in the blood (PBMC) and in the lung (BALF) (upper panel figure
8D). The ALARM cells diminished according to disease severity (T, B, Mono, but not NK) in the blood
stream. This reduction was concomitant with an increase of ALARM high cells in the lung suggesting
that ALARM cells are migrating to the site of infection. Interestingly, these changes were cell-type
specific, notably while CD4+ T cells increased it was not the case for CD8+T cells. We note that the cell-
type specific modules did not change between disease state and between blood and lung (lower panel,
figure 8D) indicating that the similar cell types were analyzed and that the cell-type specific modules
were not related to disease state.
Collectively these results show that ALARM displays a normal distribution of expression across healthy
individuals and changes in response to distinct disease states (bacterial, viral, kidney rejection). It is
noteworthy that expression alterations of this module are apparent in distinct cell types depending on
the disease conditions.
448
Recruitment of CXCR4+ T Cells During Lung Infection In Vivo
To independently validate the rapid recruitment of immune cells during lung infection, as exemplified
in the COVID-19 results above, we used a well-established pneumonia mouse model[47,48] to study T cell
recruitment. The infection was induced with *E. coli* and resolved after 7 days, with the peak of infection
occurring between day 1 and day 3. To specifically evaluate the recruitment of T cells from the blood,
we employed CD45-PE mediated immune staining of blood immune cells before and during infection[48]
(Figure 9A). This method allowed us to precisely quantify the recruitment of cells from the blood during
the infection. We observed that T cells were recruited from the blood as early as day 1, with a
significant peak at day 3, indicating a rapid response to the lung infection (Figure 9B). Upon analyzing
the membrane expression of CXCR4 on these cells, we found that 15% to 20% were CXCR4+ T cells, a
key component of the ALARM module, suggesting that a diverse set of T cells is recruited to the lung
(Figure 9B, Pie Charts). Next, we investigated the membrane display of CD69 and found that most
CXCR4+ CD69+ T cells were predominantly CD45+ cells (60-80% at day 1 and day 3), indicating their
blood-derived origin (Figure 9C). This phenomenon was particularly evident at day 1 and day 3,
corresponding with the peak of infection. Additionally, stratification of the CXCR4+ T cells by CD4+ and
CD8+ subsets showed that CD4+ T cell recruitment was much more abundant than CD8+ T cells,
mirroring observations from COVID-19 lung infection studies (Figure 9D).
These findings demonstrate that CXCR4+ T cells, particularly CD4+ T cells, are rapidly recruited from
the blood to the lung in response to infection, with peak recruitment occurring between day 1 and day
3. The presence of CD69+ cells among the recruited CXCR4+ T cells suggests that these cells are not
only migrating but are also likely being activated or display a residency program[34]. The observation of
the concomitant *protein* expression and membrane display of CXCR4 and CD69 is characteristic of the
ALARM module, which is defined by the *gene* expression of both CXCR4 and CD69. The results
underscore the crucial role of the ALARM module in orchestrating an effective immune response
during the peak of lung infection, highlighting its broader relevance beyond kidney-specific contexts.
474
475
476   **Disease classification of ALARM in immune mediated diseases**
477
The above results suggest that ALARM expression changes are associated to immune diseases, likely
via the recruitment of circulating immune cells to the site of inflammation. It may thus be possible that
the ALARM genes could be used as predictors for immune disease classification. The rationale is that
if ALARM genes are relevant for a precise disease state, they would be strong predictors to classify
healthy from disease [49]. To investigate this hypothesis, we analyzed 10 immune-mediated diseases
with distinct tissue tropisms (Supplemental Figure S8A). Out of these 10 diseases, systemic lupus
erythematosus (SLE), Sjogren syndrome and Anti-neutrophil cytoplasmic antibody-associated
vasculitis (ANCA) are known to favor the kidney among other organs. In contrast, mixed connective

tissue disease and systemic sclerosis rather favor the connective tissues and rheumatoid arthritis
targets the joints. Thus, we aimed to test whether ALARM genes are good classifiers of disease vs
healthy condition and whether the classification of disease state varies depending on the tissue
tropism. This was possible thanks to a large bulk RNA-seq dataset comprising 28 circulating immune
cell-types which included 337 patients across 10 immune mediated diseases and 79 healthy controls
[50] (Supplemental Figure S8A). We evaluated total gene expression using UMAP of this data set (Figure
10A) and found that ALARM module expression was present in all major cell types (Figure 10B). To test
the discriminative ability of the ALARM module, we then devised a classification pipeline comparing
disease state (all 10 immune diseases) vs healthy using logistic regression (Figure 10C). As in the data
set there were transcriptomics data from 28 cell types available, we focused on B cells, T cells, NK and
Monocytes by regrouping their respective sub cell types together (see Figure 10A).
We then generated an ALARM gene classification model compared it to two other models. The first
one using the most variable genes (coefficient of variation (CV^2), see methods) and the second model
was based on genes in cell type identity modules (see figure 2). The rationale for using CV^2 gene
selection was to use an independent gene selection process which is more predictive than random
gene selection. We compared the prediction performance of the three models using the F-1 score as
it gives equal weight to precision and recall (Figure 10D, F1 score). Interestingly, the ALARM genes
were the best predictors for CD8, Monocytes and NK cells and were similar in performance to the cell
type specific modules of CD4 and B cells. To account for possible imbalances in the numbers of disease
and control samples we also computed the Cohen's Kappa score (Supplemental Figure S8B). The results
were consistent with the F-scores. Of note, the CV^2 gene selection approach was less predictive in all
cell types (Figure 8D and Supplemental S8B).
To evaluate whether the ALARM module was prominent for any specific cell sub types within the major
cell types (e.g., CD4 T helper cells vs CD4 naïve cells) we estimated disease classification performance
of the ALARM module separately for each sub type. Predictability as estimated by F1 score remained
robust when each subtype was analyzed separately in comparison to the CV^2 method (Figure 10E). We
note that in certain subtypes there were too few samples to compute an accurate Kappa score
(Supplemental Figure S8C). Nevertheless, this result indicates that ALARM was found to be relevant in
all the subtypes analyzed.
Next, we evaluated whether each of the 10 immune mediated diseases could be individually classified
from healthy (Figure 10F). The best classification ability of ALARM was found for Mixed connective
tissue disease, rheumatoid arthritis, Sjogren's syndrome, SLE and Takayasu's Arteritis. This indicates
that ALARM genes are not specifically predictive for tissue tropism but appear to be relevant
independently of the targeted tissue. In most cases ALARM outperformed the CV^2 feature selection
suggesting that ALARM genes are likely to be implicated in their disease etiology. In summary, this
comprehensive classification analysis indicates that ALARM genes are strong predictors of disease
state across the majority of circulating immune cells and within the 10 immune related diseases.
523
ALARM is enriched for genetic disease associations
Since the ALARM genes are strong predictors of immune mediated disease and its general role within
multiple cell-types and across multiple infections, immune related and autoimmune diseases, it is likely
that ALARM is enriched for genes known to be associated to diseases. To test this, we exploited the
DisGeNET database[51] which provides a comprehensive compilation of published and curated human
gene disease associations (GDAs) from repositories including Mendelian, complex and environmental
diseases and enables enrichment analysis of such GDAs. Notably, we found that 24 out of the 61
ALARM genes (39%) were associated with a disease. To estimate the probability of this occurring by
chance, we compared this to random sampling of 61 genes and their quantification of GDAs
(Supplemental Figure S8D permutation). The probability of reaching 40 % of genes or more was below
> 0.001, indicating that ALARM genes are highly enriched for GDAs.
Next, we tested whether the ALARM genes were enriched for diseases associations (Figure 10G). The
top three disease categories enriched for GDAs were Juvenile arthritis, Hodgkin disease and
polyarthritis comprising by themselves distinct 13 genes. This enrichment analysis indicates that

ALARM is also genetically connected to disease state. In summary, ALARM is a strong classifier and
genetically linked to immune disease and thus is likely to play a general role in multiple immune
mediated diseases.
541
Discussion:
In this study, we gathered a cohort of matching ABMR, TCMR and stable patients and generated a
comprehensive scRNA-seq atlas of circulating immune cells across time and conditions. We then
identified multiple gene co-expression modules. Five out of nine modules were related to a single cell-
type while three were expressed in closely related cells (CD4+, CD8+ and NK cells) and only the ALARM
module was prominent in multiple cell-types. The observation that single cell transcriptomes mostly
reveal cell-type specific modules was also described by Kotliar et al., [9] in which they distinguish
between identity (i.e., cell-type specific) gene expression programs (GEP) and activity GEPs. It is
possible that cell-type specific co-expressed genes are better detected as they show a greater
coherence within a well-defined group of cells. This is also notable in the presented data as the gene
expression scores of cell-type specific modules show less variance than the ALARM module.
Nevertheless, cell-type specific gene expression may not necessarily imply that it remains constant
across conditions. For example, we noted that some cell type specific modules were associated to
disease state (Figure 2E), notably the NK cell and monocyte specific modules were increased during
rejection. It is also possible that because of the relatively low number of genes per cell detected when
compared to bulk transcriptomics, cell identity programs are preferentially detected, and more subtle
condition specific modules are not robustly detected. Indeed, while cNMF revealed additional modules
in the separate batches, only the ALARM module was consistently identified across the three batches.
The ALARM genes were found to be highly enriched for transcription factors and gene ontology
pathways associated with the gene expression machinery including transcription, mRNA processing
and ubiquitination. Prominent transcription factors included the AP-1 complex and the NFKB subunit
REL which are both associated with stress responses and inflammation. The membership of CD69 in
the ALARM module also suggests a role of stress response. CD69 is a classical early activation marker
of lymphocytes, as demonstrated by its rapid display on the surface of T cells after TCR stimulation[34].
In addition, CD69 is also known to be a tissue retention marker as it is expressed on resident memory
T cells in distinct tissues. In blood, this gene has been associated with chronic inflammation in various
diseases including rheumatoid arthritis [52] and systemic lupus erythematosus [53]. Concomitant with this,
CD69 protein expression is increased on infiltrated immune cells at the site of inflammation in immune
mediated diseases including systemic sclerosis, rheumatoid arthritis, systemic lupus erythematosus[34].
The membership of CD69 in this module thus indicates that this module role could be to prepare
circulating cells T cells for TCR stimulation and for tissue retention once moved into a tissue, i.e., to
become T resident effector cells. This notion is also consistent with the increased expression of the
ALARM module in the kidney biopsy transcriptomics data (Figure 4).
The cytokine receptor CXCR4 was also identified in this module. CXCR4 is predominantly expressed by
lymphocytes as well as monocytes and through which the CXCL12 ligand promotes chemotaxis to
tissues via a concentration gradient [54]. CXCL12 is expressed in multiple tissues including the kidney and
is altered during pathophysiological responses including immune diseases. Indeed, an alteration of
CXCL12 expression was observed in the kidney transplantation biopsies and this increase was
associated to elevated ALARM gene expression in the tissue (Figure 4E). It is possible that the CXCR4-
CXCL12 axis contributes to the recruitment of ALARM expressing cells in the case of kidney
transplantation rejection and other immune diseases. This is also consistent with the observation that
cells expressing the ALARM module decrease in the circulation during kidney graft rejection. This
observation was confirmed by both transcriptomics and histological studies of pigs as well as
transcriptomics in human kidney biopsies, thus, via CXCR4-CXCL12 leading ALARM cells to infiltrate the
tissues during rejection. This mechanism was further supported by an in vitro trans well assay, where
CXCL12 was shown to induce T cell migration. More importantly, it was found that both CXCL12
presence and migration significantly increased CD69 protein expression at the cell membrane.
Specifically, the combination of HMEC contact and CXCL12 presence was necessary for the highest

expression of CD69, similar to that observed during transmigration. Single cell transcriptomic analysis
of migrated and non-migrated cells further revealed that certain gene groups, particularly those
involved in cytoskeleton organization, migration, and immune response, were upregulated, while
others were downregulated, indicating a shift in cellular state to adapt to new functions after
migration. Furthermore,we found that migration of T cells in response to CXCL12 is accompanied by
significant metabolic reprogramming. We observed upregulation of glycolytic pathways, increased
expression of LDHA at protein levels, and enhanced glucose uptake in migrated T cells. Metabolic
reprogramming towards increased glycolysis is a hallmark of activated T cells and is essential for their
effector functions during immune responses [39]. These findings suggest that ALARM module expression
not only primes T cells for migration and tissue retention but also prepares them metabolically for the
demands of their new functional roles at sites of inflammation.
This CXCR4-CXCL12 axis also highlights the notion that the ALARM module is not necessarily specific to
transplantation rejection or the kidney. Indeed, reanalysis of circulating immune cells from publicly
available scRNA-seq data showed that the ALARM module was expressed in 45 unrelated healthy
individuals [40] and showed significant alteration between distinct pathological conditions (Figure 5).
First, the ALARM response to LPS iv injection in healthy individuals revealed that it is time-dependent,
illustrated by a gradual decrease of ALARM cells within the same individuals over time. Second, there
was a significant difference between ALARM cells depending on the location of the pathological
condition. ALARM cells were shown to be decreased in response to kidney rejection, in response to
leukocyte infiltrating UTI and Covid-19 infection of the lung highlighting the role of ALARM in the
recruitment of cells to the site of inflammation and infection. This was further supported by
bacteremia sepsis, a state of systemic inflammation in which ALARM cells were increased in the blood.
While LPS iv injection and bacteremia induce both a systemic immune response, the former is
associated is essentially an endotoxemia response associated to a transient leukopenia[42,43] while the
latter is a complex and heterogeneous condition that involves multiple factors beyond LPS, such as
pathogen virulence, host susceptibility, and coexisting medical conditions. It thus makes sense that
leukopenia is associated to the decrease of ALARM, while in bacteremia ALARM expression is
increased. Third, ALARM cells showed a gradually measurable response to disease severity. This notion
was observed by combining two distinct and complementary Covid-19 datasets one of which was
collected on BALF, and which had stratified their patients according to disease severity. ALARM cells
decreased in response to severity in the blood with a corresponding increase in the lung.
Fourth, we collected several lines of evidence suggesting that ALARM cells are indeed recruited to the
site of inflammation and/or infection. During acute rejection induced in the pig model there was a
rapid infiltration of leukocytes concomitant with the reduction of ALARM cells in the blood. The
analysis of kidney biopsies revealed an increase of ALARM gene expression during kidney transplant
rejection. Similarly, the recruitment of ALARM cells to the lung was observed during Covid-19 lung
infection. Finally, an in vivo mouse model of *E. coli* pneumonia demonstrated that CXCR4+ and CD69+
T cells are rapidly recruited from the blood to the lung during the peak of infection further supporting
the role of the ALARM module in mediating immune cell recruitment to sites of inflammation.
We thus propose a model in which ALARM expression priorities the infiltration capacity of each
circulating cell (see figure 11 model). This model has wide ranging consequences in precision medicine
as blocking of ALARM cells to the site of inflammation in the case of kidney rejection may prevent
further organ damage or attenuate the immune response in the case of Covid-19 lung infection. It may
also be useful to predict disease state as we have shown in figure 8. ALARM was a strong classifier of
immune disease when compared to healthy individuals. The importance of ALARM was independently
demonstrated by its enrichment for genes mutated in notably juvenile and poly- arthritis. Further
investigation is however required to test the specificity of disease detection or whether ALARM is
merely a response to inflammatory state in the circulation.
There are several limitations of this study, first it is based on gene transcription and thus remains to
be explored for protein expression, however this is difficult to achieve at single cell resolution and we
are not aware of any gene co-expression modules estimated at the protein level. Nevertheless, the
transwell assay and in vivo mouse model experiments indicate a connection between the CXCL12-

CXCR4 axis and CD69 and their display during and after migration. While direct evidence of the
recruitment of these cells to the tissue has been experimentally confirmed the module expression was
shown to be low in the blood and high in the kidney tissue during rejection which may also be caused
be lack of the source of these cells, or a slowing in cellular maturation before expressing ALARM genes
an avenue that should be explored in subsequent studies. A further limitation is that as of now we do
not have a protein surface marker panel that could be associated to cells with high or low ALARM
expression. Such markers would enable the purification of ALARM cells enabling further molecular and
cellular characterization. Markers would also allow the targeted modulation of the recruitment of
ALARM cells to the graft or during Covid19 as well as other immune mediated diseases may thus impact
disease severity. Nevertheless, our study remains important in terms of precision medicine,
highlighting the discovery of ALARM, which expression enables cells to be preferentially recruited to
the inflamed tissue. This notion is likely to open novel strategies of disease monitoring and disease
intervention.

655
656
657
658
659 **Methods:**

660
661
Study of kidney transplant patients
Kidney transplantation patients
The PBMC samples used in this study (see table 1) were obtained from the DIVAT biocollection (CNIL
agreement n°891735, Réseau DIVAT: 10.16.618). Every patient included in the study was enrolled in
the DIVAT biocollection following their informed consent. The PBMC from patients were isolated from
kidney transplantation biopsies, frozen with DMSO 10% and stored in liquid nitrogen at the Centre de
Ressources Biologiques (CRB, CHU Nantes, France).
Cell preparation
Frozen PBMC samples were rapidly thawed and resuspended in complete Roswell Park Memorial
Institute (RPMI) 1640 media (Invitrogen, Carlsbad, CA) with 5% FBS, pre-heated at 37°C.
Following washing steps in PBS+0.04%BSA, cell pellets were resuspended in 200µL FACS buffer (1X PBS
supplemented with 2mM EDTA, 2% FBS) in which dead cells were labelled by adding 0.1 µg/mL DAPI
(Invitrogen, Carlsbad, CA). Cells were filtered on 70 µm cell strainer and living cells were then sorted
using a Fluorescence-activated cell sorting (FACS) Aria II cell sorter (BD Biosciences, Mountain View,
CA).We used the same method as previously described for single cell RNAseq[55,56]. One million cells
were kept for each sample and resuspended in 100µL of staining buffer (PBS,2%BSA,0.01% Tween)
according to the cell hashing protocol[13] recommendations. Cells were incubated for 10 min with 10µL
of human Fc blocking reagent. Each sample was then mixed with 1uL of a specific TotalSeq-A hashtag
antibody (BioLegend, San Diego, CA) and incubated on ice for 30 min. Following 3 washing steps with
the staining buffer, cells were counted and their viability measured using an ADAM-MC automatic cell
counter (NanoEntek, Seoul,South Korea) to ensure a viability above the recommended 70%. All the
samples were pooled at an equal cell concentration in a single vial, centrifuged and resuspended in
PBS to obtain a concentration of 700 cells/µL, to match the targeted cell recovery of 32,000 cells.
Encapsulation of single cells was performed on a 10XChromium (10X Genomics, San Francisco, CA)
with the Chromium Single Cell 3' Library and Next GEM reagent kit v3. The libraries were sequenced
twice for each of the three experiments on the NovaSeq 6000 (Illumina, San Diego, CA) with S1 flow
cells. The sequenced libraries were aligned to the GRCh38-2020-A reference genome with CellRanger
v5.0.0 (10X Genomics, San Francisco, CA). The scRNA-seq was performed in 3 different experiments
following the same protocol. Each experiment included longitudinal samples from three patients (one
stable patient, one humoral rejection and one cellular rejection) as well as one late sample of a tolerant
patient.
Method demultiplexing and Seurat analysis

The count matrices were analyzed in R 4.0.3 using the Seurat R package (v4.0.2, Satija Lab[14]). Each
experiment was first processed separately, with the same workflow. First, following the standard
workflow recommendations, cells with less than 200 unique feature counts were removed (potential
empty droplets). Cells with a percentage of mitochondrial genes greater than 15% were excluded as it
results from mitochondria degradation from dead or dying cells. The hashing antibody sequences were
then collected to demultiplex and assign each cell to its sample using the MULTIseqDemux function.
Cells with too little labels information were called "Negative" while cells with a high count of two or
more different oligo-conjugated antibody sequences were called "Doublets". Only cells with a unique
HTO were kept for downstream analysis. Singlet cells were annotated automatically with the Azimuth
workflow within Seurat, by mapping the query cells on an annotated reference of 162,000 PBMC
measured with 228 antibodies[15].
All the runs were then merged in a single Seurat object. Doublets and contaminant cells to exclude
were selected by identifying cells co-expressing marker genes from distinct cell types. After
normalization of the global object, the 2,000 most variable genes in the data were selected to compute
the correction using the reciprocal principal component analysis (RPCA). The final annotated and
corrected object gathering the 12 patients was composed of 50,507 cells.

<u>Gene Module Identification</u>

Consensus Non-Negative Matrix Factorization (cNMF)[9] was used to decompose the cell vs gene
expression matrix into cell vs module and usage vs gene low-rank matrices. Non-Negative Matrix
Factorization is a stochastic method and therefore it was run with 200 NMF replicates to find a
consensus robust factorization. For each batch, the top 2000 over dispersed genes were selected as
input to the cNMF run. Different K values (7 to 14) were explored to determine the optimal number of
modules. For each K, the *stability* and *error* metrics were examined. The best K was chosen such that
the error was minimum, and the stability was maximum. Each batch was independently processed to
mitigate batch effects from the three different runs. For the three batches, the optimal number of
modules (K) were 14, 11 and 13.

<u>Module filtering:</u> All modules from the three batches were collated and hierarchical clustering was
performed to identify matching modules. Jaccard similarity score was used to define the similarity
between two modules. Finally, only those clusters were retained that could represent all three
batches. In this way, nine consistent modules were identified. A unique geneset was determined for
each consistent cluster by intersecting the top-ranking 200 genes from the modules. The threshold
was achieved by observing a scree plot of input number of genes vs the number of genes after
intersection.

<u>Module association:</u> The genes in the module were examined for a module to be associated with a
known cell type. The modules containing marker genes were associated with their respective cell type;
for example, the module with MS4A1 and CD79A genes was associated with B cells. However, one of
the modules (Module 9, later named as the ALARM module) could not be associated with a known cell
type as it did not contain cell type specific markers and was well expressed in multiple cell types.

<u>UCell module score</u>

The enrichment of a particular set of genes in an individual cell was measured with the UCell[58] module
score. The score is calculated using the Mann-Whitney U statistic, which compares the expression
levels of the module genes relative to the total gene expression of the cell. The U-statistic outcome is
then normalized between 0 to 1 to produce the UCell score.

<u>Regression analysis</u>

Change in the module score (μ) along time (τ) for each condition was modelled with linear regression
method. The 'lm()' function from R *stats* package was used to fit a distinct model $\mu \sim \tau$, per celltype
within the group of Stable, ABMR and TCMR individuals. A positive slope indicated that the module
score increased with time.

<u>Gene Ontology analysis</u>

The gene ontology (GO) functional enrichment of the ALARM module gene list was performed using
the R package WebGestaltR (v0.4.4) for Biological Process (BP) and Molecular function (MF)
annotation. P-value are obtained with the hypergeometric test for ORA (Over-Representation

Analysis). As background dataset for enrichment the top 2000 variable genes from each run were
merged and used.
Transcription factor enrichment analysis
To test for enrichment of transcription factors in the ALARM module, GSEA molecular signature
database was used to count the transcription factors gene family in the module and in the background
dataset used for gene ontology analysis. Fisher exact test was used to calculate the P-value and
enrichment of transcription factors in the ALARM module.
753
Allogeneic kidney rejection model in pig
Animal model
The study protocol was approved by the French Ministry of Higher Education, Research and Innovation
(APAFiS #30136). The experiment was performed on 60 to 80kg male pigs (*Sus scrofa*). Test card with
pre-applied antibodies from Serafol (Berlin, Germany) were used to identify the pigs' ABO blood
groups. The alloreactivity was performed by mixed lymphocyte reaction assay between donor and
recipient. The donor pig was selected from a different breed as inbred pigs might escape rejection.
Donor and recipients blood groups were ensured to be compatible to avoid hyper-acute rejection, and
mixed lymphocyte reaction assays were positive thus proving their alloreactivity.
Allogeneic transplantation
Unilateral nephrectomies were performed on two recipient pigs under general anesthesia with a
premedication by Zolazepam/Tiletamine (Zoletil ® Virbac, Carros, France) 15 mg/kg IM, before
intubation and a maintained ventilation with a mixture of 49% oxygen, 49% nitrous oxide and 2%
isoflurane. The two kidneys from a third donor pig were harvested in the same operating time. The
two recipient pigs received one collected kidney each for an orthotopic transplantation. During
surgery, a central venous catheter (CVC) was inserted into the internal jugular vein for hydration and
medication. Post-operative analgesia was performed every day with intravenous injections of
Nalbuphine (Nubain ®, Mylan, Canonsburg, Pennsylvania) and Paracetamol at a dose of 25 mg/kg.
Prophylactic antimicrobial therapy was conducted with Cefazolin 1 g (Cefovet ®, Dopharma, Ancenis,
France).
PBMC collection in Pig model
Blood samples were collected daily through the CVC and frozen in a CoolCell® container (Corning ®,
Corning, NY, USA) at -80°C following the PBMC isolation. Kidney transplant biopsies were collected
daily using automated biopsy needles of 16 gauges under ultrasound guidance while pigs were sedated
by Zolazepam/Tiletamine and locally anesthetized with Lidocaine. Kidney samples were then placed in
cryovials with 1mL fetal bovine serum (FBS) and 10% dimethylsulfoxyde (DMSO) for gradual cooling in
a CoolCell chamber.
Single cell preparation for pig model
Blood samples were processed as described previously (see Methods 1.2). After filtering, cells were
centrifuged at 300g for 10 minutes at 4°C and resuspended in staining buffer for the HTO antibody
labeling (see Methods 1.3). The sequenced libraries were aligned to the Sscrofa 11.1 (February 2017
release) reference genome with CellRanger v5.0.0.
786
Pig biopsy immunostaining analysis
Kidney biopsies fixed Carnoy's solution for 30 minutes followed by a fixation in formaldehyde for 24h
for optical microscopy purpose. A second batch of kidney biopsies was prepared for immunostaining
purpose: biopsies were placed in cryomold, covered with optimal cutting temperature (OCT)
compound and immersed in cold isopentane. Following their solidification, cryomolds were stored in
liquid nitrogen. Cryosectioning was performed and the resulting slides of kidney biopsies were stained
with periodic acid-Schiff (PAS) and Masson's trichrome stains (TM).
The cellular infiltration was counted using ImageJ[59] software on the PAS-stained kidney biopsies. Areas
of interest were selected to exclude areas with glomeruli. Pictures were first converted to 8-bit
grayscale, and the threshold of detection was set to capture only the stained cells.

<u>Transmigration Model</u>

HDMECs (10 × 10^4 cells) activated with TNF-α (100 U/ml) for 24 hours were seeded O/N onto 1%
gelatin-coated Transwell membrane inserts (24-well, 3-μm-pore polycarbonate membrane; Corning
Life Science) in endothelial cell growth medium at 37°C. On the day of the assay, purified CD3 T-cell
subsets (4×10^5) were added to the upper transwell migration chamber, and the chemokine CXCL12 (50
ng/ml) was added to the lower transwell migration chamber. Migration was assessed after 4 hours by
quantifying the number and phenotype of migrated cells in the lower chamber using 123count eBeads
counting beads and a Cytek AURORA flow cytometer (5 lasers). Migrated CD3 were surface stained
with specific antibodies to characterize phenotype CD3, CD8, CD4, CD45RA, CCR7 and activation
molecules CD69, CD25, CD127, CD95, CD103 and CD49. The antibodies used for the cytometric
analyses are listed below<u>Expression of Cytotoxic Molecules by Human transmigrated CD3 T Cell</u>
<u>Subsets</u>

To define the expression of cytotoxic molecules, transmigrated CD3 were restimulated with PMA
(50ng/mL), ionomycin (500ng/mL) and BFA (5ug/mL). Transmigrated CD3 were surface stained with
specific antibodies for phenotypic characterization of CD3, CD8, CD4, CD45RA, CCR7, and after fixation
and permeabilization (BD Cytofix/Cytoperm), intracellular staining was performed using antibodies
against granzyme B (GZMB) and perforin-1 (PERF-1), granulysin, and TNFa. The antibodies used for
cytometric analyses are listed below.

Metabolic characterization of Human transmigrated CD3 T Cell Subsets

Transmigrated T cells were stimulated over-night with plate bound anti-CD3 (1ug/mL) and anti-CD28
(2ug/mL) mAb. Cells were washed, surface stained with anti-CD3, CD4 and CD8 mAbs and cultured for
30' at 37°C 5% CO2 in glucose-free medium containing 50 µM 2-NBDG. Alternatively, cells were surface
stained with anti-CD3, CD4, CD8 and GLUT1 mAbs and after fixation and permeabilization (BD
Cytofix/Cytoperm), intracellular stained with anti-LDHA mAb. Data were acquired using a 5 lasers Cytek
AURORA flow cytometer and analyzed using OMIQ.

Antibodies used:

Target	Clone	Fluorochrome	Provider
CD3	OKT3	Purified	In house production
CD28	CD28.2	Purified	In house production
CD3	SK7	PE Cy7	BD
CD4	SK3 R7	CF568	Cytek
CD8	SK1	Sparkblue 550	Biolegend
CD69	FN50	APC Cy7	Biolegend
GLUT1	202915	FITC	R&D Systems
LDHA	APC	ab310866	Abcam
CD3	UCHT1	PE-Cy7	Biolegend
CD8	SK1	Spark Blue 550	Biolegend
CD4	SK3	cL-Fluor YG584	Cytek
CD45RA	HI100	BUV496, BV786	BD
CCR7	G043H7	APC Fire810	Biolegend
CD69	FN50	APC-Cy7	Biolegend
CD25	M-A251	PE-Fire700	Biolegend
CD127	HIL-7R-M21	APC R700	BD Biosciences
CD95	DX2	PE-Cy5	BD Biosciences
CD103	Ber-Act8	BV605	BD Biosciences

CD49a	SR84	BUV661	BD Biosciences
GZMb	QA16A02	PE-Cy5	Biolegend
IFNg	B27	APC	BD Biosciences
GNLY	RB1	AF488	BD Biosciences
TNFa	Mab11	BUV737	ThermoFisher

scRNA-seq of Transmigration model

T cells were collected as indicated in figure 5A and then processed for chromium loading. The three
conditions were processed at the same time using CITE-seq approach (see above).

Primary Analysis: Fastq files were generated from Illumina bcl files using Bcl2fastq version 2.2.
Cellranger v7.2 was employed to create a filtered scRNA gene expression matrix from the fastq files,
utilizing default parameters and the human genome reference version GRCh38-2020.

Seurat version 5.1 was used for subsequent quality control and preprocessing. HTODemux, with a
positive quality threshold of 0.95, was applied to demultiplex cells, identifying singlets and associating
each singlet with the corresponding condition. Cells with an RNA count exceeding 30,000 or exhibiting
mitochondrial gene expression above 20% were excluded. Cell type annotation was performed using
Celltypist annotation tool[60].

Secondary Analysis: For each CD8 and CD4 cell type, genes exhibiting significant gradients across the
three conditions were identified using a linear model. The dependent variable in the model
represented the condition, with control CXCL12-, control CXCL12+, and migrated groups assigned
values of 0, 1, and 2, respectively. To simulate multiple individuals, cells were randomly grouped into
10 groups. These groups served as pseudo-individuals, each containing at least 50 cells, created using
the `createfolds` function from the caret package. The final linear model was formulated as ~
condition + (1 | individual). Top genes were identified based on significant p-values (< 0.05), and the
direction of their gradients across the conditions was noted.

Among the top gene modules, several were found to participate in known pathways. The average gene
expression for these modules was calculated using the `AddModuleScore` function of Seurat.

Metabolome Pathway Analysis: The Compass algorithm[38] was used to characterize the metabolic
states of CD4+ T cells across three different conditions: Migrated, CXCL12-positive (CXCL12+), and
CXCL12-negative (CXCL12-). The algorithm designed to infer the metabolic state of cells from scRNA-
Seq data through flux balance analysis. It addresses the limitations of traditional metabolic assays in
assessing metabolic states at the single-cell level, leveraging transcriptome data to predict metabolic
activities. For reference to metabolic reactions and pathways, the RECON2 database was utilized for
this analysis. The scRNA data were first micropooled, resulting in 20 pseudobulk samples for each
condition in order to compare the same number of samples between conditions. The reaction
penalties were estimated for various metabolic pathways based on gene expression levels in each
pseudobulk sample. Reaction penalties were then converted to negative log scores, with higher values
indicating greater predicted activity. Significant active reactions were identified using the Wilcoxon
rank-sum test, comparing the Migrated samples to the CXCL12- samples. Reactions with an adjusted
p-value of less than 0.1 were considered significant.

Analysis of the ALARM module in kidney biopsies

Two separate studies were analyzed here. The Reeve *et al.* Affymetrix Microarray data in RAW CEL
format was downloaded from Gene Expression Omnibus (GEO) website with accession number
GSE98320. The samples were pre-processed using robust multi-chip averaging (RMA) implemented in
Bioconductor. The patient condition was obtained from the 'd96' metadata column as designated in
the corresponding Series Matrix file. The patients strictly defined as either TCMR (n=76), ABMR
(n=197), Mixed (n=39) or no major abnormality (STA, n=257) as a stable condition, were retained for

downstream analysis. The Callemeyn *et al.* dataset was downloaded from GEO with accession number
GSE147089. The CEL files were similarly pre-processed using the RMA method. The labels for each
sample were obtained from the Series Matrix file and the phenotypes are defined as biopsies without
ABMR (n=168), DSA negative ABMR (n=26) and DSA positive ABMR (n=30).
The score used to stratify patients was computed by averaging the z-scores of the ALARM module
genes.

<u>ALARM mean z-score distribution across conditions</u>
The Kolmogorov-Smirnov (KS) test was then used to compare the distribution of the ALARM module
score between stable and rejection conditions. This non-parametric statistical test compares the
cumulative distribution functions of the mean ALARM z-scores in both groups. The KS test statistic (D)
is the maximum vertical distance between the two distributions. The p-value of the test is the
probability of obtaining a test statistic as extreme as D or more extreme, assuming that the null
hypothesis is true. The null hypothesis is that the two samples are drawn from the same distribution.

<u>Receptor-Ligand analysis</u>
The 'iTalk' R package was used for the receptor-ligand (RL) analysis. The receptors were gene
candidates in single-cell kidney transplant stable and rejection patients. The *rawParse()* function with
`stats='mean'` was used to identify the candidate genes. For each celltype, the genes were ordered
by their average count expression. Only the top 50 percent of these genes were selected for the
subsequent RL analysis. The same criteria were adopted for selecting the ligand candidates from the
bulk RNAseq biopsy data. The significant interaction pairs were discovered from the iTalk database
restricted on the cytokine interactions only.
888

<u>Analysis of publicly available scRNA-seq datasets</u>

PBMCs of 45 healthy Volunteers (Van Der Wijst MG et al.):
Processed (de-anonymized) single-cell RNA-seq data and its relevant meta data was obtained from the
European Genome-phenome Archive (EGA) accession number EGAS00001002560. The available data
was merged to build a single Seurat object for downstream analysis. Azimuth reference for PBMCs was
used to annotate the cells.

LPS and Covid PBMC dataset (Stephenson et al.):
The processed data was downloaded from Array Express under accession number E-MTAB-10026. Only
individuals from the same batch containing the LPS-treated volunteers were selected to mitigate batch
effects. Fig 5B shows only those individuals where the major cell types were available. The Covid
patients originally annotated as Mild/Moderate was included as 'Moderate' in Fig 5D.

Sepsis-PBMC Dataset (Reyes et al.):
The pre-processed scRNA-seq data was downloaded through the Broad Institute Single Cell Portal
(SCP548). The data was further analyzed with Seurat to obtain the Module score based on UCell.

Covid – BALF Dataset (Liao et al.):
The data was accessed from GSE145926. The filtered cell matrix was processed with Seurat with the
code as provided by the authors of the original article.

<u>In vivo Mouse model experiments</u>
For induction of pneumonia, E. coli (DH5α strain), OVA-coated E. coli, grown for 18 h in Luria broth
medium at 37 °C, were washed twice (1,000g, 10 min, 37 °C), diluted in sterile isotonic saline and
calibrated by nephelometry. Bacteria (75 μl, OD600 = 0.6–0.7) were injected i.t. in anesthetized mice
to induce nonlethal acute pneumonia. Infected mice were intravenously (i.v.) injected with 10 μg of

CD45-PE on days 0, 1, 3 and 5 to evaluate T cell trafficking towards the lung. Five minutes before sample collection on each of these days, 10 µg of CD45-BV480 was i.v. injected to evaluate blood contamination during lung excision. To specifically assess the expression of membrane markers CD3, CD4, CD8 CXCR4 and CD69 conjugated monoclonal antibodies were used on cell suspensions: CD3-bv711 (145-2C11, 7311597, BD Biosciences, 1:200 dilution); CD4-buv395 (GK1.5, 1097734, BD Biosciences, 1:200 dilution); CD8-AlexaFluor700 (RPA-T8, 9025745, BD Biosciences, 1:200 dilution); anti-CD69-APC (H1.2F3, 9204727, BD Biosciences). Two independent experiments with each 3-4 mice were carried out. Anova test were used to evaluate for statistical significance across time points.

Analysis of publicly available data set of 10 immune mediated diseases

Bulk RNAseq of 28 pure immune cell types from 339 individuals divided into 10 immune-mediated diseases and 92 healthy controls was obtained from the National Bioscience Database Center (NBDC) Human Database with the accession number E-GEAD-397. The dataset was assembled as a large matrix with genes listed in rows and columns are individuals with homogeneous cell types. Functions from the Seurat pipeline were used to compute the module scores and generate the population's UMAP embedding.

Disease Classification

Combined diseases vs healthy approach was deployed for each major cell type and by cell subtype at the primary level. The Logistic Regression classifier from the R package 'caret' and the repeated cross-validation strategy for model evaluation were used.

In the next phase, for each major cell type, classification was evaluated for Healthy vs One Disease. An identical classification model was built in this phase as well. The coefficient of variation (CV^2)[61] method produced an unsupervised set of highly variable genes as a control for the ALARM module genes. The mean vs (variance/means^2) was modelled with *glmgam.fit* from the *statmod* R package for the variance estimate of every gene. The genes were ranked by the significance of deviation from the fit. The same number of variable genes was then used in the modules.

Gene association to disease terms

The association between the ALARM genes and immune disease terms was performed using the disgenet2r (v0.99.2) R package [51]. The ratio of genes associated with immune diseases to the total number of genes within the modules was compared to the corresponding ratio obtained from 1000 randomly selected sets of 61 genes.

Methodological Clarifications and Editing

Parts of this manuscript were edited and refined with the assistance of ChatGPT, the AI language model developed by OpenAI. This tool was used to improve the clarity and coherence of the text without altering the scientific content.

Acknowledgements:

We are most grateful to the Genomics Core Facility **GenoA**, member of Biogenouest and France Genomique and to the Bioinformatics Core Facility **BiRD**, member of Biogenouest and Institut Français de Bioinformatique (IFB) (ANR-11-INBS-0013) for the use of their resources and their technical support. DS was supported by Progreffe foundation. The project was supported by Region Pays de Loire "Etoile Montantes" grant for JP. We would like to thank the members of the DIVAT consortium for their involvement in the study, the physicians who helped recruit patients, and all patients who participated in this study. We also thank the clinical research associates who participated in the data collection. Data were collected from the French DIVAT multicentric prospective cohort of kidney and/or pancreatic transplant recipients (www.divat.fr, N° CNIL 914184, ClinicalTrials.gov recording: NCT02900040). We also thank the biological resource centre for biobanking (CHU Nantes, Nantes

958 Université, Centre de ressources biologiques (BB-0033-00040), F-44000 Nantes, France). The authors
959 have no conflicting financial interests.
960
Contributions:
TL, DS & JP analyzed the data and wrote the manuscript. CF and TL performed scRNA-seq experiments.
TL and SV, JB and GB performed pig transplantation model experiments. LBD performed cell sorting
experiments. CK, RD, MG and SB collected and provided patient data and kidney transplantation PBMC
samples. GT & ND performed and analyzed all transwell experiments. AB & AR performed and analyzed
mouse pneumonia experiments and critically revised the manuscript. RJ, SB provided samples,
contributed to study design, result interpretation and critically revised the manuscript. All authors
contributed to the draft of the manuscript. JP conceived the study and oversaw experiments and
analyses.
970
971 **Data availability**
972 The data underlying Figures 1 and 2 will be openly available in GEO and embargo will be lifted upon
973 acceptance.
974
975
976 Figure legends:
977
978
Figure 1. Comprehensive longitudinal single-cell RNA-sequencing of circulating immune cells in a
cohort of kidney allograft recipient.
A) Timeline of the blood sampling points post-transplantation for the patients followed longitudinally.
STA=Stable patient (n=3), TCMR=T cell mediated rejection (n=3), ABMR=Antibody-mediated rejection
(n=3). Tolerant patients are not shown.
B) Schematic diagram of the scRNA-seq preparation workflow using cell hashing. Peripheral blood
mononuclear cells (PBMCs) were collected from stable, ABMR and TCMR patients and then stained
with one different oligo-conjugated antibody before being pooled and processed using microfluidic
encapsulation.
C) UMAP dimensional-reduction embedding of the integrated samples (n=30). Each colour represents
a different cell subpopulation, adapted and manually curated from the automatic Azimuth annotation.
D) UMAP projection showing the disease state distribution, TOL=Tolerant patients.
E) UMAP projection coloured according to experimental batch of origin
992
Figure 2. ALARM Module identification
A) Schematic workflow of the gene co-expression analysis using consensus nonnegative matrix
factorisation (cNMF) module detection separate in each batch. The module selection was then refined
based on overlapping genes between the three batches.
B) Clustering of 9 modules across the 3 batches, using the Jaccard distance.
C) Heatmap showing the combined gene expression of each module, i.e. module scores of the 9
modules summarized for each distinct cell-type.
D) UMAP projection of the expression of Module 9 across all cells.
E) Regression analysis of the 9 modules by cell types. The outcome variable (y) was time (T0, T1 and
T2) and the independent variable (x) was the module score. Heatmap shows the beta values (trend)
for each cell-type and disease state. Negative values correspond to a decrease of the module score
across the three time points, positive values to an increase.
F) Super violin plots[62] showing the longitudinal trend of ALARM module in NK, CD4 T cells (top), CD8 T
cells and B cells (bottom), stratified by individuals from each batch.
G) Gene Ontology analysis of biological process (BP) and molecular function (MF) identified by
comparing the ALARM gene enrichment using the 4000 most variable genes as background (See
Supplemental file 1).

1010

Figure 3. Pig model scRNA-seq analysis

A) Schematic diagram of the acute kidney allograft rejection in pig model. Recipient pig with a unilateral nephrectomy received a kidney graft from a second healthy pig. Kidney biopsies and PBMCs were collected daily and observed in immunohistochemistry. PBMCs were prepared for a scRNA-seq analysis.

B) Immunohistochemistry of pig biopsies stained with Periodic Aid Schiff (PAS), to stain polysaccharides, glycoproteins and glycolipids at day 0, day 2, day 4 and day 6

C) Quantification of cell populations in the kidney graft at given days using Image J cell counting software (See supplemental figure 3 for details)

D) UMAP dimensional-reduction projection of the circulating immune cell types (PBMCs) scRNA-seq after PBMC isolation at D0, D2, D4 and D6.

E) Proportions of circulating immune cells (PBMC) across the different time points (D0-D6) in the recipient transplanted pig.

F) Violin plot of the ALARM module score in B cells, monocytes, CD4 T cells, CD8 T cells and NK cells throughout acute rejection in the kidney tissues (Day 0, Day2, Day 4 Day6). *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ****$P < 0.0001$.

Figure 4. ALARM gene expression increases in kidney tissues during rejection

Left panels from Reeve et al., 2017 and right panels from Callemeyn et al 2020. STA =Stable patient, ABMR=Antibody-mediated rejection, TCMR=T cell mediated rejection, Mixed=Graft undergoing ABMR and TCMR, DSA- = Donor specific antibody negative and DSA+= Donor specific antibodies positive.

A) Violin plot of the sum of z-scores of ALARM genes across conditions. Wilcoxon P-values are shown in panel comparing Stable (STA) to rejection status (ABMR, Mixed or TCMR and DSA- and DSA+ ABMR). The mean comparison p-values were computed using the Wilcoxon rank-sum test. B) Heatmap showing z-scores of ALARM genes (one gene per row) in all graft biopsies. Phenotype denotes the disease states. Patients are sorted on the mean of the module gene z-scores.

C) Cumulative distribution of the mean of z-scores of ALARM genes comparing stable vs the combined rejection conditions. K-S = Kolmogorov -Smirnov P-values and distance.

D) Ligand-Receptor analysis between receptors genes identified in circulating immune cells and cytokines genes expressed in the allograft kidney tissue under no rejection and rejection condition. Width of the arrow line is proportional to the expression of the ligands and the width of the arrowhead is proportional to the receptor. Only the top 5 associations from each cell type of differentially expressed receptors (PBMC scRNA-seq) and cytokines (Biopsy microarray) are shown.

E) Boxplots showing CXCL12 expression in biopsies from patients with high ALARM module expression (>75%), medium (25-75%) and low (<25%). Wilcoxon rank-sum test was used to calculate P-values shown above.

Figure 5: Analysis of ALARM Module Expression and T Cell Behavior in an In Vitro Transwell Assay

A) Schematic representation of the in vitro transwell assay used to study CXCL12-mediated recruitment of T cells. The assay involved a membrane covered with human dermal microvascular endothelial cells (HDMEC), allowing for the comparison between unstimulated cells, cells in direct contact with CXCL12, and those migrating across the membrane in response to a CXCL12 gradient.

B) Quantification of T cell subsets following migration. The composition of naïve, central memory (CM), effector memory RA (TEMRA), and effector memory (EM) T cells remained similar post-migration and in response to CXCL12 alone, indicating that all T cell subsets are attracted to and migrate in response to the CXCL12 gradient.

C) Analysis of T cell surface marker expression. CD69, a component of the ALARM module, showed a slight increase in response to CXCL12, with a more significant upregulation in migrated cells. CD25, CD49A, and CD95 levels remained unchanged.

D) Functional analysis of migrated T cells. CXCL12-exposed and transmigrated cells were stimulated with PMA/ionomycin after 24 hours. Migrated cells showed significantly increased expression of IFN-

1062 γ in both CD4+ and CD8+ T cells, while TNF-α, granzyme B (GZMB), and granulysin (GNLY) expression
1063 remained constant.
1064 Statistical significance was determined using one-way ANOVA, followed by post-hoc tests where
1065 appropriate. Significance levels are indicated as follows: *< 0.05, ** < 0.01, and *** < 0.001.
1066
Figure 6: Single Cell Characterization of T Cell Behavior in an In Vitro Transwell Assay
A) Schematic representation of the transwell assay used to investigate CXCL12-mediated signaling and
its effect on T cell migration. Three distinct conditions were assessed: (1) CXCL12-: T cells were placed
below the transwell membrane without exposure to CXCL12, (2) CXCL12+: T cells were placed below
the transwell membrane in direct contact with CXCL12, and (3) Migrated: T cells were placed above
the membrane and collected from below after migrating in the presence of CXCL12.
B) UMAP visualizations displaying the clustering of naïve CD4 and CD8 T cells based on gene expression.
Clusters were identified based on differences in ribosomal gene expression and CD4 versus CD8
annotation (top left & right). Annotation according to condition (bottom left) shows a notable
separation between CXCL12- T cells and those exposed to CXCL12 or that had transmigrated. The
ALARM module shows lower expression in the CXCL12- condition compared to the CXCL12+ and
Migrated groups (bottom right).
C) Dot plot showing the expression levels of specific genes across the different conditions (Migrated,
CXCL12+, CXCL12-).
D) Violin plots depicting the expression distribution of gene groups in pooled CD4 and CD8 T cells across
the three conditions. The plots demonstrate changes in gene expression associated with immune
response, migration, cytoskeleton organization, and stress response, as well as the downregulation of
genes related to structural organization and gene regulation.
1085
Figure 7. ALARM Module associates with a shift of immune cells metabolic fonctions
A) Metabolome pathway analysis of CD4+ T cells, using 20 pseudobulk sample for each experimental
condition (Migrated, CXCL12-positive, and CXCL12-negative). The differential activity of metabolic
reactions was evaluated by comparing the mean of Migrated samples and CXCL12- samples
B) PCA visualization of the pseudobulk samples by condition based on the metabolome pathway
analysis scores
C) Volcano plot illustrating glycolytic enzyme changes in migrated T cells compared to controls. The x-
axis shows Cohen's d effect sizes, and the y-axis indicates the statistical significance (−log10 p-values).
Key enzymes in the glycolysis/gluconeogenesis pathway are labeled.
D) Flow cytometry analysis and quantification of GLUT1 and 2-NBDG uptake in T cells before and after
transmigration. Left panels show representative flow cytometry plots, and right panels show paired
comparisons for each parameter. LDHA expression levels (measured by MFI) are also shown before
and after transmigration. Statistical significance is indicated, with p-values provided for each
comparison.
1100
Figure 8. ALARM gene expression is altered in distinct immune conditions
A) Outline of study on Lipopolysaccharide (LPS) intravenous injection (iv) scRNAseq experiment
performed on healthy volunteers obtained from Stephenson et al. The annotated expression data was
used to compute the ALARM module score across time points after LPS injection in healthy volunteers.
Violin plots show the module score across cell types and condition. Total number of cells are
mentioned below the plot. P-values are shown above the violin plots and were calculated using
Wilcoxon rank sum test comparing each cell type between healthy and LPS conditions.
B) Violin plots of ALARM module score for three individuals which had matching timepoints across all
cell types in the condition LPS 90 min and LPS 10h. Lines connect the median ALARM module score
across time points for each individual separately. P-values are shown above the violin plots and were
calculated using Wilcoxon rank sum test comparing the two distinct time points.
C) Outline of patients with sepsis and urinary tract infection obtained from Reyes et al,. Annotated
data was used to compute ALARM module score across the different cell-types in urinary tract infection

(UTI) patients and bacteraemia sepsis patients. Number of cells and individuals used are shown below
the violin plots. P-values are shown above the violin plots and were calculated using Wilcoxon rank
sum test comparing each cell type between healthy and LPS conditions.
D) Outline of PBMC scRNA-seq generated on healthy individuals, moderate covid-19 and severe covid-
19 patients (Stephenson et al.,) and Bronchoalveolar lavage fluid (BALF) from a distinct cohort (Liao et
al.). Upper panel shows median ALARM module score for each cell-type for PBMC (blue) and BALF
(red). Lower panel shows for each cell-type the module score for each corresponding cell-type specific
module.
1122
Figure 9 Recruitment and Characterization of CXCR4+ T Cells During Lung Infection in Vivo
A) Schematic representation of the experimental setup used to study T cell recruitment during *E. coli*
pneumonia in a mouse model. The infection was induced on day 0, with peak infection occurring
between days 1 and 3, and resolution by day 7. CD45-PE was intravenously injected at different time
points (days 0, 1, 3, and 5) to stain and track blood-derived immune cells migrating into the lungs
during infection.
B) Quantification of CD45+ T cells in the lungs at various time points post-infection. The left plot shows
the number of CD45+ T cells at each time point. The accompanying pie charts depict the proportion of
CXCR4+ and CXCR4- T cells over the course of the infection.
C) Percentage of CD45+ cells within the CXCR4+ CD69+ T cell population over time.
D) Comparative analysis of CD45+ CXCR4+ CD4+ and CD8+ T cell counts at different time points.
Independent experiments with 3-4 mice per condition were carried out, and statistical significance was
determined using one-way ANOVA, followed by post-hoc tests where appropriate. Significance levels
are indicated as follows: *< 0.05, ** < 0.01, and *** < 0.001.
1137
Figure 10 ALARM genes are implicated in and predictive of immune mediated diseases
A) UMAP of bulk transcriptomic study (Ota et al., 2020). Colours represent major cell types and each
single point represents a bulk transcriptomic dataset. Cell sub types are annotated in the plot.
B) The same UMAP showing expression of ALARM module expression across samples.
C) Outline of classification approach used to test disease prediction
D) Barchart showing the F-1 score of disease classification for ALARM, cell-type specific modules and
coefficient of variation (CV^2) selected genes in each major cell type separately.
E) Barchart of F1-scores computed for disease classification in each sub cell type separately for ALARM
and CV^2 genes
F) Radar chart showing the F1-score in each cell type for classification between each disease and
healthy separately using ALARM genes in each major cell type.
G) Barchart showing the top three enriched diseases in the ALARM genes using the DisGeNET curated
database.
1151
Figure S1 ScRNA-seq integration of a cohort of kidney allograft recipients
A) Expression profiles of cell-specific markers distinguish the PBMC populations. Average expression is
the log-normalized expression average of the cells by cell type, size of the dots is associated to the
fraction of cells of the cluster in which the gene is detected.
B) Cell type proportion by cluster in the overall PBMC population. Comparison include Stable patients,
Rejection (ABMR+TCMR) and two public datasets: healthy volunteers from Stephenson et al, 2021, and
healthy volunteers from Reyes et al, 2020. One-way ANOVA with Tukey's multiple comparisons post
hoc test was performed, *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.
C) UMAP projection showing the sample distribution, ABMR=Humoral rejection, TCMR=Cellular
rejection, STA=Stable, TOL=Tolerant. Second UMAP shows the timepoint distribution, T0=Time point
0 (Graft), T1=Time point 1, T2=Time point 2.
D) K-bet acceptance rate by cell types following the CCA batch integration. Complete k-bet acceptance
rate was computed on the overall PBMC population.
1165

Figure S2 Assessing module distribution
A) Alluvial plot showing the shared origin of each module across batches. Each cell is assigned to the most enriched module they express.
B) Boxplots showing the distribution of Pearson correlation coefficient *R* of pairwise gene expression correlation by cell type across modules. A module of random genes was generated to compare with the nine other modules. Multiple gene pairs from each module are selected to compute the rho values using a subset of common cells within the cell type. 1000 repetitions of 50 pairs each yielded 50000 *R* values, as represented in every box. The subset of common cells changes with every repetition.
C) Boxplots showing the distribution of Pearson correlation coefficient *R* of pairwise gene expression correlation by cell type for Module 9.

Figure S3 Cellular characterization of pig PBMC by scRNA-seq analysis
A) Image J analysis of immuno-stained slices to quantify cell infiltration. Red boxes show counting areas which excludes glomeruli. Right panel shows the cells which were then counted using Image J software.
B) Pig data were annotated using the Sscrofa reference genome and associated to their human homologue. The expression profiles of cell type-specific markers is shown in the dotplot of relative expression by cell type. Size of the dots are associated to the fraction of cells of the cluster in which the gene is detected. The UMAP embeds the 4 samples (D0, D2, D4, D6) with their corresponding cell type annotation.

Fig S4 Transwell assay experiments
A) Gating strategy used in the transwell assay for cell type annotation. Lymphocytes were gated based on forward scatter (FSC) and side scatter (SSC), followed by singlet selection and the identification of CD3+ T cells. CD4+ and CD8+ T cells were further delineated based on CD45RA and CCR7 expression to distinguish between naïve (Tn), central memory (Tcm), and effector memory RA (TemRA) subsets. Additionally, CD4 T regulatory (Treg) cells were identified by CD25 and CD127 expression to differentiate between activated (aTreg) and non-Treg populations.
B) Percentage of CD4+ and CD8+ T cells expressing CD69 under five different experimental conditions: before the transwell assay (baseline), CXCL12 only, HDMEC alone, CXCL12 with HDMEC, and after migration through HDMEC with CXCL12.
C) UMAP visualization of scRNA-seq data from the transwell experiment showing all cell populations. The top plot displays T cell subtypes identified by their transcriptional profiles, including CD4+ and CD8+ T naïve cells, CD8+ T effector memory RA (TEMRA) cells, mucosal-associated invariant T (MAIT) cells, and CD4+ and CD8+ T effector memory (TEM) cells annotated using Celltypist. The bottom plot shows the distribution of cells based on their experimental condition: control which is CXCL12-, CXCL12+ and transmigrated. The clustering of cells in the UMAP space indicates differences in transcriptional states based on both cell type and condition.
D) ALARM gene expression by condition. Dot plot showing the relative expression levels of ALARM-associated genes in CD4+ and CD8+ T cells across the different conditions of the transwell assay (Migrated, CXCL12+, and CXCL12-). The size of the dots represents the percentage of cells expressing the gene, and the color intensity represents the level of expression, with darker shades indicating higher expression.

Fig 5S. Overview of the cohort distribution
A) ALARM module score distribution by cell type based on scRNA-seq analysis of PBMC from 45 unrelated healthy individuals[40].
B) Left: ALARM module score distribution across sex. No significance was found for the sex parameter with a t-test. Right: ALARM module score distribution across age categories. No significance was found for the age parameter with a one-way ANOVA test.

Fig 6S. ALARM genes can classify immune-mediated diseases

A) Distribution of the bulk RNA-seq samples across the 28 circulating immune cell types for the 337 patients distributed across 10 immune mediated diseases and the 79 healthy controls.

B) Area Under Curve (AUC) and Cohen's Kappa score across cell types in disease and healthy patients.

C) AUC and Cohen's Kappa score across cell subtypes in disease and healthy patients

D) Distribution of the ratio of genes associated to Gene-Disease Association (GDA) terms in 1000 modules of 61 randomly selected genes. The ALARM module has a ratio of 0.39 genes associated to GDA terms (red line).

References:

1. Reyes, M. *et al.* An immune cell signature of bacterial sepsis. *Nat. Med.* **26**, 333–340 (2020).

2. Perez, R. K. *et al.* Single-cell RNA-seq reveals cell type-specific molecular and genetic associations to lupus. *Science* **376**, eabf1970 (2022).

3. Kazer, S. W. *et al.* Integrated single-cell analysis of multicellular immune dynamics during hyperacute HIV-1 infection. *Nat. Med.* **26**, 511–518 (2020).

4. Zhang, F. *et al.* Defining inflammatory cell states in rheumatoid arthritis joint synovial tissues by integrating single-cell transcriptomics and mass cytometry. *Nat. Immunol.* **20**, 928–942 (2019).

5. Stephenson, E. *et al.* Single-cell multi-omics analysis of the immune response in COVID-19. *Nat. Med.* **27**, 904–916 (2021).

6. Bernardes, J. P. *et al.* Longitudinal Multi-omics Analyses Identify Responses of Megakaryocytes, Erythroid Cells, and Plasmablasts as Hallmarks of Severe COVID-19. *Immunity* **53**, 1296-1314.e9 (2020).

7. Chaumette, T. *et al.* Monocyte Signature Associated with Herpes Simplex Virus Reactivation and Neurological Recovery After Brain Injury. *Am. J. Respir. Crit. Care Med.* (2022) doi:10.1164/rccm.202110-2324OC.

8. del Rosario, R. C. H. *et al.* Histone acetylome-wide associations in immune cells from individuals with active Mycobacterium tuberculosis infection. *Nat. Microbiol.* **7**, 312–326 (2022).

9. Kotliar, D. *et al.* Identifying gene expression programs of cell-type identity and cellular activity with single-cell RNA-Seq. *eLife* **8**, e43803 (2019).

10.   Loupy, A., Mengel, M. & Haas, M. Thirty years of the International Banff Classification for

Allograft Pathology: the past, present, and future of kidney transplant diagnostics. *Kidney Int.* **101**,

678–691 (2022).

11.   Sirota, M. & Sarwal, M. M. Transplantomics: Toward Precision Medicine in Transplantation

Research. *Transplantation* **101**, 1777–1782 (2017).

12.   Moreau, A., Varey, E., Anegon, I. & Cuturi, M.-C. Effector Mechanisms of Rejection. *Cold*

Spring Harb. Perspect. Med. **3**, a015461 (2013).

13.   Stoeckius, M. *et al.* Cell Hashing with barcoded antibodies enables multiplexing and doublet

detection for single cell genomics. *Genome Biol.* **19**, 224 (2018).

14.   Stuart, T. *et al.* Comprehensive integration of single-cell data. *Cell* **177**, 1888-1902.e21

(2019).

15.   Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587.e29

(2021).

16.   Büttner, M., Miao, Z., Wolf, F. A., Teichmann, S. A. & Theis, F. J. A test metric for assessing

single-cell RNA-seq batch correction. *Nat. Methods* **16**, 43–49 (2019).

17.   Langfelder, P. & Horvath, S. WGCNA: an R package for weighted correlation network analysis.

BMC Bioinformatics **9**, 559 (2008).

18.   Nestorowa, S. *et al.* A single-cell resolution map of mouse hematopoietic stem and

progenitor cell differentiation. *Blood* **128**, e20–e31 (2016).

19.   Yamashita, I., Nagata, T., Tada, T. & Nakayama, T. CD69 cell surface expression identifies

developing thymocytes which audition for T cell antigen receptor-mediated positive selection. *Int.*

Immunol. **5**, 1139–1150 (1993).

20.   Ziegler, S. F., Ramsdell, F. & Alderson, M. R. The activation antigen CD69. *Stem Cells* **12**, 456–

465 (1994).

21.   Werfel, T., Boeker, M. & Kapp, A. Rapid expression of the CD69 antigen on T cells and natural

killer cells upon antigenic stimulation of peripheral blood mononuclear cell suspensions. *Allergy*

52, 465–469 (1997).

22.   Turner, J.-E., Becker, M., Mittrücker, H.-W. & Panzer, U. Tissue-Resident Lymphocytes in the

Kidney. *J. Am. Soc. Nephrol. JASN* **29**, 389–399 (2018).

23.   López-Cabrera, M. *et al.* Transcriptional Regulation of the Gene Encoding the Human C-type

Lectin Leukocyte Receptor AIM/CD69 and Functional Characterization of Its Tumor Necrosis

Factor-α-responsive Elements (∗). *J. Biol. Chem.* **270**, 21545–21551 (1995).

24.   Castellanos, M. C. *et al.* Expression of the leukocyte early activation antigen CD69 is

regulated by the transcription factor AP-1. *J. Immunol. Baltim. Md 1950* **159**, 5463–5473 (1997).

25.   Tögel, F., Isaac, J., Hu, Z., Weiss, K. & Westenfelder, C. Renal SDF-1 signals mobilization and

homing of CXCR4-positive cells to the kidney after ischemic injury. *Kidney Int.* **67**, 1772–1784

(2005).

26.   Song, A., Jiang, A., Xiong, W. & Zhang, C. The Role of CXCL12 in Kidney Diseases: A Friend or

Foe? *Kidney Dis.* **7**, 167–176 (2021).

27.   Hoffmann, U. *et al.* SDF-1 expression is elevated in chronic human renal allograft rejection.

Clin. Transplant. **20**, 712–718 (2006).

28.   Pabst, R. The pig as a model for immunology research. *Cell Tissue Res.* **380**, 287–304 (2020).

29.   Reeve, J. *et al.* Assessing rejection-related disease in kidney transplant biopsies based on

archetypal analysis of molecular phenotypes. *JCI Insight* **2**, 94197 (2017).

30.   Callemeyn, J. *et al.* Transcriptional Changes in Kidney Allografts with Histology of Antibody-

Mediated Rejection without Anti-HLA Donor-Specific Antibodies. *J. Am. Soc. Nephrol.* **31**, 2168–

2183 (2020).

31.   Moll, N. M. & Ransohoff, R. M. CXCL12 and CXCR4 in bone marrow physiology. *Expert Rev.*

Hematol. **3**, 315–322 (2010).

32.   Yellowley, C. CXCL12/CXCR4 signaling and other recruitment and homing pathways in

1298   fracture repair. *BoneKEy Rep.* **2**, 300 (2013).

1299   33.   Burger, J. A. & Kipps, T. J. CXCR4: a key receptor in the crosstalk between tumor cells and

1300   their microenvironment. *Blood* **107**, 1761–1767 (2006).

1301   34.   Cibrián, D. & Sánchez-Madrid, F. CD69: from activation marker to metabolic gatekeeper. *Eur.*

1302   *J. Immunol.* **47**, 946–953 (2017).

1303   35.   Scarneo, S. A. *et al.* Expression of membrane Hsp90 is a molecular signature of T cell

1304   activation. *Sci. Rep.* **12**, 18091 (2022).

1305   36.   Nieminen, M. *et al.* Vimentin function in lymphocyte adhesion and transcellular migration.

1306   *Nat. Cell Biol.* **8**, 156–162 (2006).

1307   37.   Liu, Y. *et al.* Prognostic and Immunological Role of STK38 across Cancers: Friend or Foe? *Int. J.*

1308   *Mol. Sci.* **23**, 11590 (2022).

1309   38.   Wagner, A. *et al.* Metabolic modeling of single Th17 cells reveals regulators of autoimmunity.

1310   *Cell* **184**, 4168-4185.e21 (2021).

1311   39.   MacIver, N. J., Michalek, R. D. & Rathmell, J. C. Metabolic regulation of T lymphocytes. *Annu.*

1312   *Rev. Immunol.* **31**, 259–283 (2013).

1313   40.   van der Wijst, M. G. P. *et al.* Single-cell RNA sequencing identifies celltype-specific cis-eQTLs

1314   and co-expression QTLs. *Nat. Genet.* **50**, 493–497 (2018).

1315   41.   Tigchelaar, E. F. *et al.* Cohort profile: LifeLines DEEP, a prospective, general population cohort

1316   study in the northern Netherlands: study design and baseline characteristics. *BMJ Open* **5**,

1317   e006772 (2015).

1318   42.   Vos, A. F. de *et al.* In Vivo Lipopolysaccharide Exposure of Human Blood Leukocytes Induces

1319   Cross-Tolerance to Multiple TLR Ligands. *J. Immunol.* **183**, 533–542 (2009).

1320   43.   Andonegui, G., Goyert, S. M. & Kubes, P. Lipopolysaccharide-Induced Leukocyte-Endothelial

1321   Cell Interactions: A Role for CD14 Versus Toll-Like Receptor 4 Within Microvessels1. *J. Immunol.*

1322   **169**, 2111–2119 (2002).

44.   Beutler, B. TLR4 as the mammalian endotoxin sensor. *Curr. Top. Microbiol. Immunol.* **270**,

109–120 (2002).

45.   Beutler, B., Du, X. & Poltorak, A. Identification of Toll-like receptor 4 (Tlr4) as the sole conduit

for LPS signal transduction: genetic and evolutionary studies. *J. Endotoxin Res.* **7**, 277–280 (2001).

46.   Liao, M. *et al.* Single-cell landscape of bronchoalveolar immune cells in patients with COVID-

19. *Nat. Med.* **26**, 842–844 (2020).

47.   Roquilly, A. *et al.* Alveolar macrophages are epigenetically altered after inflammation, leading

to long-term lung immunoparalysis. *Nat. Immunol.* **21**, 636–648 (2020).

48.   Broquet, A. *et al.* Sepsis-trained macrophages promote antitumoral tissue-resident T cells.

Nat. Immunol. 1–18 (2024) doi:10.1038/s41590-024-01819-8.

49.   Mi, Z., Guo, B., Yin, Z., Li, J. & Zheng, Z. Disease classification via gene network integrating

modules and pathways. *R. Soc. Open Sci.* **6**, 190214 (2019).

50.   Ota, M. *et al.* Dynamic landscape of immune cell-specific gene regulation in immune-

mediated diseases. *Cell* **184**, 3006-3021.e17 (2021).

51.   Piñero, J. *et al.* The DisGeNET knowledge platform for disease genomics: 2019 update.

Nucleic Acids Res. **48**, D845–D855 (2020).

52.   Afeltra, A. *et al.* Expression of CD69 antigen on synovial fluid T cells in patients with

rheumatoid arthritis and other chronic synovitis. *Ann. Rheum. Dis.* **52**, 457–460 (1993).

53.   Su, C. C., Shau, W. Y., Wang, C. R., Chuang, C. Y. & Chen, C. Y. CD69 to CD3 ratio of peripheral

blood mononuclear cells as a marker to monitor systemic lupus erythematosus disease activity.

Lupus **6**, 449–454 (1997).

54.   Polyfunctionality of the CXCR4/CXCL12 axis in health and disease: Implications for

therapeutic interventions in cancer and immune-mediated diseases. doi:10.1096/fj.202001273R.

55.   Abidi, A. *et al.* Characterization of Rat ILCs Reveals ILC2 as the Dominant Intestinal Subset.

Front. Immunol. **11**, (2020).

56. Drouin, M. *et al.* CLEC-1 is a death sensor that limits antigen cross-presentation by dendritic

cells and represents a target for cancer immunotherapy. *Sci. Adv.* **8**, eabo7621 (2022).

57. Feregrino, C. & Tschopp, P. Assessing evolutionary and developmental transcriptome

dynamics in homologous cell types. *Dev. Dyn.* **251**, 1472–1489 (2022).

58. Andreatta, M. & Carmona, S. J. UCell: Robust and scalable single-cell gene signature scoring.

Comput. Struct. Biotechnol. J. **19**, 3796–3798 (2021).

59. Schneider, C. A., Rasband, W. S. & Eliceiri, K. W. NIH Image to ImageJ: 25 years of image

analysis. *Nat. Methods* **9**, 671–675 (2012).

60. Xu, C. *et al.* Automatic cell-type harmonization and integration across Human Cell Atlas

datasets. *Cell* **186**, 5876-5891.e20 (2023).

61. Brennecke, P. *et al.* Accounting for technical noise in single-cell RNA-seq experiments. *Nat.*

Methods **10**, 1093–1095 (2013).

62. Kenny, M. & Schoen, I. Violin SuperPlots: visualizing replicate heterogeneity in large data

sets. *Mol. Biol. Cell* **32**, 1333–1334 (2021).

1362

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Figure 9

Figure 10

Figure 11

circulating immune cells

ALARM
expression
"high"

ALARM

ALARM
expression
"low"

Early Activation Transcription
Factor Module
(61 genes)

- Peripheral blood mononuclear cells express ALARM at different levels
- ALARM cells are depleted in the blood in response to tissue infection & inflammation
- ALARM cells increase at site of infection and inflammation
- Recruitment of ALARM is associated to CXCL12 expression in the tissue
- ALARM cells are implicated across a wide variety of immune diseases
- ALARM genes are strong predictors of immune mediated diseases

Tissue effector activity

CXCL12

Supp. Figure 6

A

B

Classification: Combined Diseases vs. Healthy

C

Classification: Combined Diseases vs. Healthy

D

Ratio of unique genes in GDA terms passing FDR<0.05

Supp. Figure 1

A

B

C

D

Supp. Figure 2

Supp. Figure 3

A

B

Supp. Figure 4

Supp. Figure 5

A Healthy Individuals (Van der Wijst et al., dataset)

B