<u>Title</u>

Syncytiotrophoblast 5'-tRNA fragments are placental endocrine signals contributing

to sterile inflammation in preeclampsia

<u>Authors</u>

William Robert Cooke[1]*

Peiyong Jiang[2, 3]

Lu Ji[2, 3]

Jinyue Bai[2, 3]

Gabriel Davis Jones[1]

Y. M. Dennis Lo[2, 3]

Christopher Redman[1]

Manu Vatish[1]

*Corresponding author

<u>Affiliations</u>

1) Nuffield Department of Women's and Reproductive Health, University of Oxford,

Oxford, UK

2) Centre for Novostics, Hong Kong Science Park, Pak Shek Kok, New Territories,

Hong Kong SAR, China

3) Department of Chemical Pathology, The Chinese University of Hong Kong, Prince

of Wales Hospital, Shatin, New Territories, Hong Kong SAR, China

<u>Abstract</u>

The relationship between placental pathology and the maternal syndrome of

preeclampsia is incompletely characterised. Mismatch between placental nutrient

supply and fetal demands induces stress in the syncytiotrophoblast, the layer of

placenta in direct contact with maternal blood. Such stress alters the content and

increases the release of extracellular vesicles into the maternal circulation. 5'-tRNA

fragments (5'-tRF) constitute the majority of short RNA in these vesicles. Here we

characterise 5'-tRF profiles in syncytiotrophoblast vesicles in preeclampsia, reporting

differential expression of >900 fragments. We detected preeclampsia-dysregulated

5'-tRF in peripheral plasma, where we identified a placentally-derived load. We

demonstrated these 5'-tRF induce sterile inflammation in macrophages and

monocytes. Our findings suggest syncytiotrophoblast-derived 5'-tRF may contribute

to distant vascular inflammation, previously described in preeclampsia. This is the

first demonstration of endocrine actions of vesicle-bound 5'-tRF, which may have

relevance in other disciplines studying their functions, such as immunology and

cancer biology.

Introduction

Preeclampsia is a complex placental syndrome, with multiple causes and a variable phenotype. Clinically it is characterised by maternal sterile inflammation and endothelial dysfunction[1,2]. A point of convergence in the disorder is stress in the syncytiotrophoblast, the interface between maternal and fetal circulations[3]. The link between syncytiotrophoblast stress and maternal symptoms, likely through blood-borne factors, is incompletely characterised[4]. Angiogenic proteins (soluble fms-like tyrosine kinase [sFlt-1] and placental growth factor [PlGF]) are important syncytiotrophoblast stress signals which contribute to the preeclamptic syndrome[5]. Excess sFlt-1 sensitises endothelial cells to pro-inflammatory cytokines[6]. These biomarkers have also been successfully applied to the clinical diagnosis of preeclampsia[7]. Altered extracellular vesicle [EV] release from the syncytiotrophoblast, whilst also a key contributor in the pathophysiology of preeclampsia, remains underexplored[8].

The healthy syncytiotrophoblast releases EV into the maternal circulation in increasing quantities with advancing gestation[9]. These lipid bilayer-bound particles are decorated with surface proteins and shuttle their contents to distant cells[10,11]. Cellular stress increases EV release; this is reflected in preeclampsia where circulating placental EV are more abundant[12,13]. Placental EV cargoes also change in preeclampsia; for example nitric oxide synthase is reduced and neprilysin (a metalloprotease causing hypertension) is increased[14,15].

We recently reported 5'-transfer RNA fragments [5'-tRF] as the predominant short RNA species in healthy syncytiotrophoblast EV[16]. 5'-tRF form when mature tRNA

molecules are cleaved by a number of stress-induced ribonucleases including

angiogenin[17,18]. They can be exported as EV cargo[19]. 5'-tRF expression profiles can

be complex; at least 417 tRNA genes can be cleaved at multiple loci[20]. 5'-tRF are

also multifaceted signalling molecules, regulating transcription, translation and

epigenetic inheritance[21–24]. 5'-tRF have mostly been investigated in cancer biology

and immunology where they are described as intracellular, autocrine and paracrine

signals[19,25,26].

5'-tRF are stress signals; syncytiotrophoblast stress is a key feature of preeclampsia.

We hypothesised that syncytiotrophoblast 5'-tRF release may change in

preeclampsia. We used placental perfusion as a source of syncytiotrophoblast EV to

show that 5'-tRF expression in preeclampsia differs from normal pregnancy. Our in-

vivo work demonstrates circulating placentally-derived 5'-tRF. Cell culture studies

determined preeclampsia-dysregulated 5'-tRF trigger sterile inflammation in

macrophages. Together these findings suggest 5'-tRF are placental endocrine

signals, linking syncytiotrophoblast stress with maternal inflammation in

preeclampsia.

Results

The preeclamptic syncytiotrophoblast exports 5'-tRF in EV

We isolated small extracellular vesicles (SEV) and medium-large EV (MLEV),

alongside placental biopsies from women with early-onset preeclampsia (n=8) and

matched healthy controls (n=6, Supplementary Table 1) using dual-lobe placental

perfusion[16]. We performed single-end short RNA sequencing (size selecting <50

nucleotides) using the Illumina HiSeq 4000 platform[27]. Sequence length distribution

plots confirmed the majority of reads in MLEV and SEV from both preeclampsia and

normal placentas were 30-34 nucleotides long (Figure 1A). In contrast a peak at 22

nucleotides was greater in the placental samples from both groups. The majority of

fragments in MLEV and SEV mapped to tRNA species, rather than to the microRNA

and ribosomal RNA species seen in placental tissues (Figure 1B). Coverage plots

demonstrated that 5'-tRF constitute almost all tRNA reads in EV in preeclampsia as

well as normal pregnancies (Figure 1C). Enrichment of 5'-tRF (but not 3' fragments)

in EV compared to cells of origin is consistent with studies in immune cells[19,26].

5'-tRF are differentially expressed in preeclamptic syncytiotrophoblast EV

We used a bespoke bioinformatics pipeline to investigate differential expression of

tRNA fragments. To minimise data loss, we assigned each unique RNA fragment an

identifier, then annotated fragments after differential expression analysis (see details

in Methods). We identified 983 differentially expressed short RNA fragments in

preeclampsia MLEV compared to controls; 182 mapped to 5'-tRF using GtRNAdb

2.0[28]. No fragments were found to be differentially expressed in preeclampsia SEV.

This is likely consistent with the different biogenesis of smaller EV, which are

released constitutively via the endosomal pathway[29].

The 12 most abundant differentially expressed fragments are shown in Table 1 and

Figure 2A. Using the sum of preeclampsia MLEV normalised counts as a

denominator, these 12 fragments account for 64% of the differentially expressed

counts. 5% of these counts were accounted for by the 626 least abundant fragments.

Thus, a small number of abundant fragments represented the majority of the signal

in an otherwise complex dysregulated short RNA profile in preeclampsia MLEV.

Comparing to human small RNA databases, among the 12 most abundant

differentially expressed fragments, 3 were identified as 5'-tRF and 4 as piwi-

interacting RNA. The remaining 5 had no directly matched reference sequences in

human small RNA databases and are referred to as "unmatched". Review of

fragment sequences demonstrated overlap; indeed all 12 showed only one or two

nucleotide insertions or deletions differentiating them from known 5'-tRF (differences

highlighted in Table 1). A complex profile of differential expression in preeclampsia

MLEV was thus dominated by minor variants of a handful of abundant 5'-tRNA

fragments.

We sought target 5'-tRF which were differentially expressed in preeclampsia EV to

validate our findings *ex-vivo* (in placental perfusion samples) and *in-vivo* (in plasma).

5'-tRF are known to be expressed in other circulating EV, most notably from immune

cells. The most abundant EV in plasma are blood-cell derived[30]. Hence, we used

publicly available blood-cell datasets to quantify possible target 5'-tRF likely to be

within contaminating EV in plasma (Supplementary Table 2). Using these data we

selected three targets for validation which were: abundant in syncytiotrophoblast EV

(above 90th centile PE expression amongst 983 differentially expressed fragments),

upregulated in preeclampsia and of low relative abundance in potentially

contaminating EV source cells. The relative abundance of these three targets in

STBEV using RNAseq is shown in Figure 2B (5'-tRF targets named A, B and C for

brevity but full sequences and tRNA derivations shown in Supplementary Table 2).

We used a custom small RNA assay with stem-looped reverse transcription primers

and a specific minor-groove binding TaqMan™ probe to compare relative 5'-tRF

expression in placental EV obtained by perfusion using quantitative real-time

polymerase chain reaction. Results validated target 5'-tRF upregulation and were

consistent with RNAseq findings, showing a median of 1.47 fold upregulation in

preeclampsia (Figure 2C).

Plasma EV 5'-tRF are upregulated in pregnancy and placentally-derived

Total EV RNA were isolated from maternal plasma using membrane-affinity columns

(EV size profiles for plasma compared to perfusion MLEV are shown in

Supplementary Figure 1; extensive characterisation of plasma membrane-affinity EV

has previously been published[31]). Target 5'-tRF were significantly more abundant in

pregnant peripheral plasma EV than in non-pregnant matched control samples

(median 2.8 fold higher in pregnancy, $p<0.05$ for all three) (Figure 3A). The

difference between these sample groups suggested a pregnancy-specific load of 5'-

tRF in plasma EV. The high abundance of these 5'-tRF in non-pregnant samples

confirms they were not unique to pregnancy (median Cq values in non-pregnant

samples: A 24.0, B 27.9, C 25.9). We then acquired paired plasma samples

simultaneously from the uterine and peripheral veins of women without preeclampsia

undergoing elective caesarean section for an indication unrelated to preeclampsia

(e.g. breech presentation). The uterine vein directly receives blood from the

placenta, thus placentally-derived molecules are typically more abundant in these

samples (Figure 3B)[32,33]. All three 5'-tRF were more abundant in uterine vein

samples (median 1.4 fold, $p<0.05$ for all three) (Figure 3C). These data support a

placentally-derived load of 5'-tRF in circulating plasma EV, consistent with our

hypothesis that syncytiotrophoblast 5'-tRF are endocrine signals.

EV-bound preeclampsia-upregulated 5'-tRF promote macrophage, but not

monocyte activation

5'-tRF can influence cellular function through a wide-range of regulatory mechanisms

at the level of transcription, translation and the epigenome[24]. Vascular inflammation

is a key feature of preeclampsia[34]. Tissue-resident macrophages are known to be

activated in preeclampsia[35]; recent data suggest syncytiotrophoblast EV contribute

to this[36,37]. We treated monocytes and macrophages with the most abundant 5'-tRF

upregulated in preeclampsia (labelled A – full sequence in Supplementary Table 2),

contained within otherwise undecorated lipid vesicles (in order to distinguish the

effect of one tRF from the accompanying RNA, lipids and proteins in perfusion-

derived EV). We used a scrambled version of the most abundant STBEV 5'-tRF

sequence as a negative control, following validation against untreated cells

(Supplementary Figure 2). Macrophages were activated to a type 1 phenotype after

12 hours treatment with tRF-A, increasing expression of pro-inflammatory cytokines

(Figure 4A). This pro-inflammatory 5'-tRF action was confined to macrophages, with

no changes observed when the same experiment was repeated in undifferentiated

monocytes (Figure 4B).

EV-bound 5'-tRF may indirectly activate endothelial cells, reducing expression

of endothelial nitric oxide synthase

Endothelial damage is considered a common factor in the wide-ranging maternal

organ dysfunction of preeclampsia[1,2]. Vascular macrophages are known to modulate

adjacent endothelial cell function[38,39]. We treated human umbilical vein endothelial

cells with vesicle-bound tRF-A and with the supernatants from activated

macrophages (cells which lie adjacent to endothelial cells in the maternal vessel

wall). tRF-A activated endothelial cells (quantified through increased expression of

adhesion molecules) indirectly through macrophage stimulation, but not directly

(Figure 4C).

The primary presenting feature of preeclampsia is high blood pressure. Immediate

blood pressure regulation involves constitutive release of nitric oxide from endothelial

cells, which maintains a state of relative vasodilation. In preeclampsia, circulating

nitrite levels are reduced, though the mechanism for this remains unclear[40]. tRF-A

reduces expression of endothelial nitric oxide synthase in endothelial cells, indirectly

via macrophage activation (Figure 4C). Thus aberrant placental EV 5'-tRF release in

preeclampsia may contribute to the maternal endothelial dysfunction and

hypertension which characterise the disease.

Discussion

5'-tRF have gained significant interest as signalling molecules in diverse disciplines: from human biology to drosophila, yeasts and plants[41–43]. Most studies have used cell culture as a model system. Work in breast cancer has identified intracellular 5'-tRF release promoting metastasis[25]. Mycobacterium infection in human macrophages has been shown to trigger local 5'-tRF release in EV, resulting in Toll-like receptor 7 activation in neighbouring cells[26]. T cell activation can be inhibited by blocking released 5'-tRF, suggesting autocrine and paracrine actions[19]. Pregnancy provides a unique opportunity to investigate 5'-tRF at a whole-organ level. The placenta is expelled with the fetus during parturition and can be studied intact *ex-vivo*. Both our sequencing and qPCR data demonstrate 5'-tRF release from the syncytiotrophoblast in extracellular vesicles, in both healthy and preeclamptic pregnancies. Moreover, routine opening of the abdomen for caesarean section offers additional *in-vivo* evidence to further assess discoveries from placental perfusion. By sampling uterine and peripheral veins simultaneously, we have conclusively demonstrated a 5'-tRF load in maternal plasma which is placentally-derived. Together with functional data in macrophages which are found distant to the syncytiotrophoblast, this provides compelling evidence for placental endocrine signalling with 5'-tRF. This principle might be conserved across other organ systems and species, with far-reaching implications for short RNA biology.

Mature tRNA molecules are cleaved by a number of ribonucleases, most notably angiogenin, in response to various cellular stressors[18]. Preeclampsia is a heterogeneous syndrome, unified by stress in the syncytiotrophoblast[4]. A change to the 5'-tRF expression profile in syncytiotrophoblast EV is thus consistent with

existing models for preeclampsia pathophysiology. We note reports that angiogenin-

cleaved 5'-tRF have a 2-3 cyclic phosphate residue at their 3' end, meaning they are

not captured by conventional RNA sequencing[44]. Our approaches would only detect

3'-hydroxyl groups, suggesting syncytiotrophoblast 5'-tRF may be a product of other

ribonucleases.

There is evidence for a variety of regulatory functions for 5'-tRF[24]. This includes

binding argonaute proteins to regulate mRNA stability and affecting components of

the ribosomal machinery to impact translation. We found 5'-tRF-A directly activated

macrophages, but not monocytes or umbilical vein endothelial cells. We propose this

difference may be accounted for by phagocytosis of EV by macrophages, trafficking

to the endosomal compartment (usually free of nucleic acids) and activation of Toll-

like receptor 7 or 8[45]. This hypothesis is supported by published work demonstrated

a lack of macrophage response to unencapsulated 5'-tRF and warrants further

investigation[26]. Macrophages are not typically in direct contact with blood; however

in preeclampsia the endothelial barrier is disrupted[1]. Syncytiotrophoblast vesicles

contribute to this endothelial damage[46]. We suggest that circulating EV-bound 5'-tRF

might directly reach macrophages in the vessel walls in preeclampsia, where their

functional effect could contribute to the well-described sterile inflammation of the

disease. The close relationship between vascular macrophages and neighbouring

endothelial cells may indirectly implicate 5'-tRF in potentiation of the hypertension

and diffuse organ dysfunction which typify preeclampsia[38]. We found no direct effect

of the most abundant preeclampsia-dysregulated 5'-tRF on human umbilical vein

endothelial cells, yet syncytiotrophoblast EV are known to change endothelial cell

function[46]. This discrepancy may be attributed to the absence of other EV RNA and

proteins which are necessary to trigger or potentiate this effect; it may reflect our

decision to use umbilical vein endothelium as a model. We speculate that the direct

endocrine role for 5'-tRF in preeclampsia may extend beyond macrophages other to

cells in direct contact with maternal blood that are involved in maternal disease, such

as liver sinusoidal endothelium and pericytes. Investigating these alternative targets

is the topic of ongoing work in our laboratory.

High-profile studies have recently highlighted the importance of circulating RNA in

preeclampsia, reporting both transcriptomic and miRNA dysregulation[47–49]. The

placenta is both necessary and sufficient to cause preeclampsia, yet the majority of

cell-free plasma/serum RNA is non-placental[50]. By using placental perfusion as our

experimental model, we eliminated the confounding non-placental RNA in blood,

finding the majority of short RNA released into the maternal circulation are 5'-tRNA

fragments. Previous studies investigating syncytiotrophoblast EV short RNA have

used lower fidelity models to source EV (e.g. explants) and taken bioinformatic

approaches which disregard 5'-tRF data, despite reporting their presence[13,51]. Our

ongoing work also focuses on the optimisation of techniques to isolate placental EV

5'-tRF from complex biofluids such as plasma, with attention to their clinical

relevance. Other endocrine products of syncytiotrophoblast stress such as sFlt-1 and

PlGF have proved highly valuable as diagnostic markers in preeclampsia,

transforming the diagnostic approach at the bedside[7,52]. However, the relatively poor

positive predictive value of angiogenic markers makes the search for other

biomarkers of the stressed syncytiotrophoblast clinically important.

In summary, we demonstrate differential release of syncytiotrophoblast 5'-tRF

fragments into the maternal circulation in preeclampsia can trigger sterile

inflammation in potentially distant macrophages. We propose that 5'-tRF form

another piece of the jigsaw, contributing to an explanation for the transmission of

preeclampsia from a single organ (the placenta) to multiple distant diseased cells in

the unwell mother.

Methods

1. Sample collection and storage

This project was approved by the Central Oxfordshire Research Ethics Committee

(07/H0607/74 and 07/H0606/148). Participants provided informed written consent.

Preeclampsia was defined using the ISSHP classification. Placentas were obtained

at the time of caesarean section and perfused within 10 minutes of delivery. Uterine

vein samples were taken during caesarean section, just prior to uterine incision,

using a 21-gauge needle and syringe. Peripheral blood samples were taken from the

antecubital fossa using 21-gauge needle with vacutainer system into 4.5 ml sodium

citrate vacutainer tubes (BD Diagnostics, UK). Non-pregnant samples were from

females of reproductive age. Plasma was obtained by centrifugation at 1500 g for 15

minutes. Samples were processed within 30 minutes of collection, aliquoted and

stored immediately at -80°C.

2. Placental perfusion

Placentas from 8 women with early-onset preeclampsia and 6 normal pregnancies

were perfused using a well-established dual-lobe perfusion technique[53]. Maternal

perfusate was centrifuged at 10 000 g for 30 minutes to isolate medium-large

extracellular vesicles (MLEV). The supernatant was centrifuged at 150 000 g for 2

hours to isolate small extracellular vesicles (SEV). Biopsies of placental tissue were

taken from the maternal surface of a non-perfused lobe. EV were characterised

using Nanoparticle Tracking Analysis, transmission electron microscopy and

Western blotting as described previously[53].

3. RNA sequencing

RNA was isolated from MLEV, SEV and placental tissue using Total RNA

Purification Plus Kit (Norgen Biotek Corporation, Canada). After confirming RNA

quantity and integrity using Bioanalyser (Agilent Technologies, Germany), the same

amount of input RNA was loaded for library preparation using the NEBNext Multiplex

Small RNA library preparation kit (New England Biolabs, USA). Libraries were size-

selected for fragments 15-50 bp by gel electrophoresis; fragment size and

concentration was confirmed using High Sensitivity D1000 ScreenTape (Agilent,

UK). Single-end sequencing by synthesis was undertaken using an Illumina HiSeq

4000 machine (Illumina, USA).

Sequence reads were analyzed using sRNAnalyzer[54]. Briefly, sequencing adaptors

and low-quality reads were removed using Cutadapt[55]. All identical reads in

sequence reads were collapsed, thus generating a set of unique reads (referred to

as fragment ID in this study). The number of sequence reads attributed to a fragment

ID were defined as the raw expression level of such a fragment ID. The fragment IDs

were mapped to the human small RNA databases allowing 2 mismatches. The

human small RNA databases comprised miRNA, piRNA, snoRNA, rRNA, and tRNA.

The miRNA, piRNA, and snoRNA reference sequences were obtained from the

study by Wu et al[54]. The rRNA and tRNA reference sequences were obtained from

SILVA database[56] and GtRNAdb 2.0[28], respectively. The normalised expression

level for each fragment ID within a class of RNA was used for downstream

differential expression analyses. The normalised expression level for a fragment ID

in a RNA class was calculated by dividing the raw expression level by a per million

scaling factor of total reads of that RNA class, expressed as reads per million (RPM).

Compared to normal pregnancy placentas, differentially expressed fragment IDs in

the MLEV of preeclampsia placentas were determined using the following criteria: i)

At least one MLEV sample had expression level of greater than 100 RPM, ii) P-value

was required to be less than 0.05 based on Mann Whitney *U* test after Benjamini-

Hochberg correction. The in-house bioinformatics pipeline was written in Perl and R

languages for counting sequence reads, tag annotations, and differential expression

analysis. Sequence read archive (SRA) data for blood cell datasets were

downloaded from NCBI, with SRA identifiers shown in Supplementary Table 3[57].

4. Plasma EV RNA isolation

Plasma aliquots were thawed at 37°C and centrifuged at 3000 *g* for 5 minutes to

remove cryoprecipitates. For membrane-based affinity isolation of total vesicular

RNA, 500 µL plasma was loaded onto exoRNeasy™ Midi columns and

manufacturer's instructions followed (Qiagen, Hilden, Germany). RNA was stored in

aliquots at -80°C.

5. RT-qPCR detection

Stem-loop primers were designed for specific short RNA target sequences identified

from RNA-sequencing analysis. Reverse transcription reactions and qPCR detection

were undertaken using Custom TaqMan™ Small RNA Assays (Applied Biosystems,

USA) following manufacturer's protocols. Custom assay linearity and specificity were

verified.

Two qPCR instruments were used: QuantStudio™ 7 for plasma and QuantStudio™

6 for cell culture assays (Applied Biosystems, USA). Quantification cycles (Cq) were

determined automatically by instrument-associated software using standard settings.

Relative expression was determined by following the PFAFFL approach.

6. Cell culture

THP-1 cells (ATCC, Manassas, VA) were seeded onto 24-well Nunc plates at 50 000

cells per well in RPMI-1640 medium supplemented with 10% Fetal Calf Serum. Cells

were grown with 6.2 ng/mL phorbol 12-myristate-13-acetate (PMA) for 24 hours to

differentiate into macrophages. Transfection experiments were performed using RNA

oligonucleotides (IDT, San Diego, CA) with a 5'-P modification; the sequence for

tRF-A is shown in Supplementary Table 2. The scramble RNA control was the most

abundant STBEV tRF without 5'-P modification and U nucleotides replaced with A

(sequence 5'-GCAAAGGAGGAACAGAGGAAGAAAACACGCCA-3'). 9.2uM RNA

was packaged into lipid vesicles using N-[1-(2,3-Dioleoyloxy)propyl]-N,N,N-

trimethylammonium methylsulfate (DOTAP), following manufacturer's instructions

(Roche, Basel, Switzerland). Human Umbilical Vein Endothelial Cells were

purchased and seeded onto 24-well Nunc plates at 25 000 cells per well in

Endothelial Cell Growth Medium 2 (PromoCell, Heidelberg, Germany). Macrophage

supernatant experiments were conducted by preparing 2X Growth Medium and

mixing 50:50 with THP-1 supernatant. Cellular RNA was isolated using RNeasy™

Plus Mini Kit according to manufacturer's instructions (Qiagen, Hilden, Germany).

Target mRNA expression was quantified using the following TaqMan™ Assays

(Applied Biosystems, USA): Hs02758991_g1, Hs00164932_m1, Hs01003372_m1,

Hs01574665_m1, Hs00174128_m1, Hs01555410_m1, Hs01011518_m1.

7. Data presentation

All data within this manuscript are derived from distinct samples. Figure 3B was

created using BioRender.com (Toronto, Canada). Statistical analyses and figures

were generated using RStudio (RStudio, USA) and Prism9 (GraphPad, USA).

Normality of data was assessed and two-tailed Mann-Whitney U or t-tests used

accordingly in Figures 3A and 4. One-tailed Wilcoxon matched pairs signed rank

tests were used in Figure 3C given paired samples were from the same patient, and

only a reduction in peripheral venous samples was anticipated.

Data availability

Raw data and bioinformatic pipelines will be made available upon reasonable

request to the corresponding author.

<u>References</u>

1. Roberts, J. M. *et al.* Preeclampsia: An endothelial cell disorder. *Am. J. Obstet.*

Gynecol. **161**, 1200–1204 (1989).

2. Powe, C. E., Levine, R. J. & Karumanchi, S. A. Preeclampsia, a disease of the

maternal endothelium: the role of antiangiogenic factors and implications for

later cardiovascular disease. *Circulation* **123**, 2856–2869 (2011).

3. Magee, L. A., Nicolaides, K. H. & von Dadelszen, P. Preeclampsia. *N. Engl. J.*

Med. **386**, 1817–1832 (2022).

4. Redman, C. W. G., Staff, A. C. & Roberts, J. M. Syncytiotrophoblast stress in

preeclampsia: the convergence point for multiple pathways. *American Journal*

of Obstetrics and Gynecology vol. 226 S907–S927 (2022).

5. Levine, R. J. *et al.* Circulating Angiogenic Factors and the Risk of

Preeclampsia. *N. Engl. J. Med.* **350**, 672–683 (2004).

6. Cindrova-Davies, T., Sanders, D. A., Burton, G. J. & Charnock-Jones, D. S.

Soluble FLT1 sensitizes endothelial cells to inflammatory cytokines by

antagonizing VEGF receptor-mediated signalling. *Cardiovasc. Res.* **89**, 671–

679 (2011).

7. Zeisler, H. *et al.* Predictive Value of the sFlt-1:PlGF Ratio in Women with

Suspected Preeclampsia. *N. Engl. J. Med.* **374**, 13–22 (2016).

8. Cooke, W. R., Jones, G. D., Redman, C. W. G. & Vatish, M.

Syncytiotrophoblast Derived Extracellular Vesicles in Relation to Pre-

eclampsia. *Matern. Med.* (2021).

9. Salomon, C. *et al.* A gestational profile of placental exosomes in maternal

plasma and their effects on endothelial cell migration. *PLoS One* **9**, (2014).

10. Théry, C. *et al.* Minimal information for studies of extracellular vesicles 2018

(MISEV2018): a position statement of the International Society for Extracellular

Vesicles and update of the MISEV2014 guidelines. *J. Extracell. Vesicles* **7**,

(2018).

11. Cronqvist, T., Erlandsson, L., Tannetta, D. & Hansson, S. R. Placental

syncytiotrophoblast extracellular vesicles enter primary endothelial cells

through clathrin-mediated endocytosis. *Placenta* **100**, 133–141 (2020).

12. Knight, M., Redman, C. W. G., Linton, E. A. & Sargent, I. L. Shedding of

syncytiotrophoblast microvilli into the maternal circulation in pre-eclamptic

pregnancies. *BJOG An Int. J. Obstet. Gynaecol.* **105**, 632–640 (1998).

13. Salomon, C. *et al.* Placental Exosomes as Early Biomarker of Preeclampsia:

Potential Role of Exosomal MicroRNAs Across Gestation. *J. Clin. Endocrinol.*

Metab. **102**, 3182–3194 (2017).

14. Motta-Mejia, C. *et al.* Placental Vesicles Carry Active Endothelial Nitric Oxide

Synthase and Their Activity is Reduced in Preeclampsia. *Hypertension* **70**,

372–381 (2017).

15. Gill, M. *et al.* Placental Syncytiotrophoblast-Derived Extracellular Vesicles

Carry Active NEP (Neprilysin) and Are Increased in Preeclampsia.

Hypertension **73**, 1112–1119 (2019).

16. Cooke, W. R. *et al.* Maternal circulating syncytiotrophoblast-derived

extracellular vesicles contain biologically active 5'-tRNA halves. *Biochem.*

Biophys. Res. Commun. **518**, 107–113 (2019).

17. Fu, H. *et al.* Stress induces tRNA cleavage by angiogenin in mammalian cells.

FEBS Lett. **583**, 437–442 (2009).

18. Li, G. *et al.* Distinct Stress-Dependent Signatures of Cellular and Extracellular

tRNA-Derived Small RNAs. *Adv. Sci.* **9**, (2022).

19.   Chiou, N.-T., Kageyama, R. & Ansel, K. M. Selective Export into Extracellular

Vesicles and Function of tRNA Fragments during T Cell Activation. *Cell Rep.*

25, 3356-3370.e4 (2018).

20.   Lyons, S. M., Fay, M. M. & Ivanov, P. The role of RNA modifications in the

regulation of tRNA cleavage. *FEBS Letters* vol. 592 2828–2844 (2018).

21.   Sharma, U. *et al.* Biogenesis and function of tRNA fragments during sperm

maturation and fertilization in mammals. *Science (80-.).* **351**, 391–396 (2016).

22.   Ivanov, P., Emara, M. M., Villen, J., Gygi, S. P. & Anderson, P. Angiogenin-

Induced tRNA Fragments Inhibit Translation Initiation. *Mol. Cell* **43**, 613–623

(2011).

23.   Chen, Q. *et al.* Sperm tsRNAs contribute to intergenerational inheritance of an

acquired metabolic disorder. *Science* **351**, 397–400 (2016).

24.   Di Fazio, A. & Gullerova, M. An old friend with a new face: tRNA-derived small

RNAs with big regulatory potential in cancer biology. *Br. J. Cancer 2023* 1–11

(2023) doi:10.1038/s41416-023-02191-4.

25.   Liu, X. *et al.* A pro-metastatic tRNA fragment drives Nucleolin oligomerization

and stabilization of its bound metabolic mRNAs. *Mol. Cell* **82**, 2604-2617.e8

(2022).

26.   Pawar, K., Shigematsu, M., Sharbati, S. & Kirino, Y. Infection-induced 50-half

molecules of tRNAHisGUG activate Toll-like receptor 7. *PLoS Biol.* **18**, (2020).

27.   HiSeq 4000 - Illumina, Inc (San Diego, CA, USA).

https://emea.support.illumina.com/sequencing/sequencing_instruments/hiseq-

4000.html.

28.   Chan, P. P. & Lowe, T. M. GtRNAdb 2.0: An expanded database of transfer

RNA genes identified in complete and draft genomes. *Nucleic Acids Res.* **44**,

D184–D189 (2016).

29.   Teng, F. & Fussenegger, M. Shedding Light on Extracellular Vesicle

Biogenesis and Bioengineering. *Adv. Sci.* **8**, 2003505 (2021).

30.   Auber, M. & Svenningsen, P. An estimate of extracellular vesicle secretion

rates of human blood cells. *J. Extracell. Biol.* **1**, (2022).

31.   Stranska, R. *et al.* Comparison of membrane affinity-based method with size-

exclusion chromatography for isolation of exosome-like vesicles from human

plasma. *J. Transl. Med.* **16**, 1–9 (2018).

32.   Cerdeira, A. S. *et al.* Circulating soluble fms-like tyrosine kinase-1 is

placentally derived in normal pregnancy: First in vivo evidence. *Pregnancy*

Hypertens. **16**, 145–147 (2019).

33.   Cerdeira, A. S. *et al.* In vivo evidence of significant placental growth factor

release by normal pregnancy placentas. *Sci. Rep.* **10**, 1–5 (2020).

34.   Redman, C. W. G., Sacks, G. P. & Sargent, I. L. Preeclampsia: An excessive

maternal inflammatory response to pregnancy. *Am. J. Obstet. Gynecol.* **180**,

499–506 (1999).

35.   Huda, S. S. *et al.* Visceral adipose tissue activated macrophage content and

inflammatory adipokine secretion is higher in pre-eclampsia than in healthy

pregnancys. *Clin. Sci. (Lond).* **131**, 1529 (2017).

36.   Awoyemi, T. *et al.* Syncytiotrophoblast Extracellular Vesicles From Late-Onset

Preeclampsia Placentae Suppress Pro-Inflammatory Immune Response in

THP-1 Macrophages. *Front. Immunol.* **12**, (2021).

37.   Liu, X. *et al.* Trophoblast-Derived Extracellular Vesicles Promote Preeclampsia

by Regulating Macrophage Polarization. *Hypertens. (Dallas, Tex. 1979)* **79**,

2274–2287 (2022).

38.   Hernandez, G. E. & Iruela-Arispe, M. L. The many flavors of monocyte /

macrophage-endothelial cell interactions. *Curr. Opin. Hematol.* **27**, 181 (2020).

39.   Baer, C., Squadrito, M. L., Iruela-Arispe, M. L. & De Palma, M. Reciprocal

interactions between endothelial cells and macrophages in angiogenic

vascular niches. *Exp. Cell Res.* **319**, 1626–1634 (2013).

40.   Sandrim, V. C. *et al.* Nitric oxide formation is inversely related to serum levels

of antiangiogenic factors soluble fms-like tyrosine kinase-1 and soluble

endogline in preeclampsia. *Hypertens. (Dallas, Tex. 1979)* **52**, 402–407

(2008).

41.   Karaiskos, S., Naqvi, A. S., Swanson, K. E. & Grigoriev, A. Age-driven

modulation of tRNA-derived fragments in Drosophila and their potential

targets. *Biol. Direct* **10**, 51 (2015).

42.   Alves, C. S. & Nogueira, F. T. S. Plant Small RNA World Growing Bigger:

tRNA-Derived Fragments, Longstanding Players in Regulatory Processes.

Front. Mol. Biosci. **8**, 529 (2021).

43.   Tyczewska, A. & Grzywacz, K. tRNA-derived fragments as new players in

regulatory processes in yeast. *Yeast* (2022) doi:10.1002/YEA.3829.

44.   Shigematsu, M., Kawamura, T. & Kirino, Y. Generation of 2′,3′-Cyclic

Phosphate-Containing RNAs as a Hidden Layer of the Transcriptome. *Front.*

Genet. **9**, (2018).

45.   Lee, B. L. & Barton, G. M. Trafficking of endosomal Toll-like receptors. *Trends*

Cell Biol. **24**, 360 (2014).

46.   Gupta, A. K. *et al.* A comparative study of the effect of three different

syncytiotrophoblast micro-particles preparations on endothelial cells. *Placenta*

26, 59–66 (2005).

47.   Rasmussen, M. *et al.* RNA profiles reveal signatures of future health and

disease in pregnancy. *Nat. 2022 6017893* **601**, 422–427 (2022).

48.   Moufarrej, M. N. *et al.* Early prediction of preeclampsia in pregnancy with cell-

free RNA. *Nat. 2022 6027898* **602**, 689–694 (2022).

49.   Srinivasan, S. *et al.* Discovery and Verification of Extracellular miRNA

Biomarkers for Non-invasive Prediction of Pre-eclampsia in Asymptomatic

Women. *Cell Reports Med.* **1**, 100013 (2020).

50.   Roberts, J. M. & Escudero, C. The placenta in preeclampsia. *Pregnancy*

Hypertens. **2**, 72 (2012).

51.   Wei, J. *et al.* Placental trophoblast debris mediated feto-maternal signalling via

small RNA delivery: Implications for preeclampsia. *Sci. Rep.* **7**, 1–12 (2017).

52.   Maynard, S. E. *et al.* Excess placental soluble fms-like tyrosine kinase 1 (sFlt1)

may contribute to endothelial dysfunction, hypertension, and proteinuria in

preeclampsia. *J. Clin. Invest.* **111**, 649–58 (2003).

53.   Dragovic, R. A. *et al.* Isolation of syncytiotrophoblast microvesicles and

exosomes and their characterisation by multicolour flow cytometry and

fluorescence Nanoparticle Tracking Analysis. *Methods* **87**, 64–74 (2015).

54.   Wu, X. *et al.* sRNAnalyzer-a flexible and customizable small RNA sequencing

data analysis pipeline. *Nucleic Acids Res.* **45**, 12140–12151 (2017).

55.   Martin, M. Cutadapt removes adapter sequences from high-throughput

sequencing reads. *EMBnet.journal* **17**, 10 (2011).

56.   Quast, C. *et al.* The SILVA ribosomal RNA gene database project: improved

data processing and web-based tools. *Nucleic Acids Res.* **41**, (2013).

57.   Home - SRA - NCBI. https://www.ncbi.nlm.nih.gov/sra.

Acknowledgements

We wish to thank Fenella Roseman, research midwife, for her assistance in

collecting clinical samples.

Author contributions

Study was conceived and designed by WRC, CR and MV. Bioinformatic analyses

were performed by PJ, LJ, JB, WRC and GDJ. Supervision was provided by PJ,

YMDL, CR and MV. Experimental work was performed and manuscript written by

WRC. All authors edited and approved the final manuscript.

Competing interests

YMDL holds equity in DRA, Insighta Grail/Illumina and Take2. PJ holds equity in

Illumina. PJ is a consultant to Take2. PJ is a Director of DRA and KingMed Future.

YMDL, PJ, and LJ receives royalties from Illumina, LabCorp, Grail, DRA, Xcelom

and Take2.

Sources of Funding

WRC receives a clinical doctoral training fellowship from the Wellcome Trust, which

funded this research.

1   <u>Figures and Tables</u>

Figure 1

A

B

C

Figure 1

Assessment of reads from medium-large extracellular vesicles (MLEV), small

extracellular vesicles (SEV) and placental samples obtained from placentas in

normal pregnancy (Normal) and early-onset preeclampsia (PE).

A: Sequence length distribution plots showing size in base pairs (bp) for short RNA

fragments after removal of adaptors and low-quality reads. B: Mapping of short RNA

fragments. C: Coverage plots showing percentage of per-base coverage for mapped

tRNA fragments.

Figure 2

Figure 2

Expression in early-onset preeclampsia and normal pregnancy syncytiotrophoblast

EV obtained by placental perfusion:

A: twelve most abundant differentially expressed short RNA fragments quantified

using RNAseq (n=6 normal, n=8 preeclampsia).

B: three selected target tRF quantified using RNAseq (labelled A, B, C for brevity; full

sequences in Supplementary Table 2; n=6 normal, n=8 preeclampsia).

C: three target tRF quantified using stem-loop qPCR (normalised to reference gene

TBP, n=4 per group).

Boxes show median/interquartile range; whiskers show max/min.

Figure 3

Figure 3

Detection of target tRF in plasma EV

A: Expression of three target tRF using qPCR in total EV isolated from

pregnant and non-pregnant peripheral venous plasma (labelled A, B, C for brevity;

full sequences in Supplementary Table 2); n=7 pregnant, n=6 non-pregnant; boxes

show median/interquartile range; whiskers show max/min.

B: Diagram demonstrating sampling rationale for uterine and peripheral venous

blood

C: Expression of three target tRF using qPCR in total EV isolated from uterine and

peripheral venous plasma; n=5 paired samples from the same women taken

simultaneously.

Figure 4

Figure 4

tRF actions relevant to preeclampsia.

A: Cytokine expression by THP-1 derived macrophages after treatment with vesicle-

encapsulated RNA after 12 hours (labelled tRF-A and Scramble for brevity; full

sequences in Supplementary Table 2 and Methods).

B: Cytokine expression by THP-1 monocytes (undifferentiated) after treatment with

vesicle-encapsulated RNA after 12 hours.

C: mRNA expression in human umbilical vein endothelial cells treated with vesicle-

encapsulated RNA (Direct), or the cell culture medium from macrophages pre-

treated with vesicle-encapsulated RNA (Indirect) after 6 hours.

All data represent n=4 biological repeats per group.

Table 1

ID	Sequence	Length	GtRNAdb mapping	Median PE count	Median N count	Log2Fold Change	Adjusted p value	% PE DE reads
T1	GCATTGGTGGTTCAGTGGTAGAATTCTCGCCT	32	tRF-Gly-GCC	276174	307639	0.16	0.0335	24.5
T2	GCATTG_TGGTTCAGTGGTAGAATTCTCGCCT	31	Unmatch	104785	152208	0.66	0.0058	9.2
T3	TCCCT_GTGGTCTAGTGGTTAGGATTCGGCGCT	32	Unmatch	91141	42460	-1.10	0.0058	8.2
T4	GCATTGGTGGTTCAGTGGTAGAATTCTCGCCC	32	tRF-Gly-GCC	65470	76778	0.32	0.0150	5.6
T5	TCCCTGGTGGTCTAGTGGTTAGGATTCGGCGCT	33	tRF-Glu-CTC	44786	19826	-1.50	0.0054	4.7
T6	TCCCTGGTGGTCTAGTGGTTAGGATTCGGCGC_	32	piR-hsa-5938	28094	11533	-1.29	0.0085	2.5
T7	GCATTGTGGTTCAGTGGTAGAATTCTCGCCC_	31	Unmatch	26253	38285	0.80	0.0058	2.2
T8	TCCCTG_TGGTCTAGTGGTTAGGATTCGGCGCC	32	Unmatch	21370	11323	-0.88	0.0058	1.9
T9	GCATTGTGGTTCAGTGGTAGAATTCTCGCC__	30	Unmatch	20168	31843	0.65	0.0305	1.9
T10	TCCCTGGTGGTCTAGTGGTTAGGATTCGGCG__	31	piR-hsa-5938	13644	3284	-2.19	0.0085	1.3
T11	GCCCGGCTAGCTCAGTCGGTAGAGCATGAGAC	32	piR-hsa-27622	12516	20917	0.73	0.0220	1.1
T12	GTTTCCGTAGTGTAGTGGTTATCACGTTCGC__	31	piR-hsa-28877	12154	20820	0.87	0.0128	1.1

49   **Table 1**

50   Descriptors of twelve most abundant differentially expressed short RNA fragments in

51   preeclampsia EV. Normal pregnancy (N), early-onset preeclampsia (PE),

52   differentially expressed (DE). Nucleotide differences from GtRNAdb annotated tRNA

53   fragments are highlighted in grey.