<u>Title</u>

Differential Expression of 5'-tRNA Fragments in Circulating Preeclampsia

Syncytiotrophoblast Vesicles Drives Macrophage Inflammation

<u>Authors</u>

William Robert Cooke[1]*

Peiyong Jiang[2, 3]

Lu Ji[2, 3]

Jinyue Bai[2, 3]

Gabriel Davis Jones[1]

Y. M. Dennis Lo[2, 3]

Christopher Redman[1]

Manu Vatish[1]

*Corresponding author: william.cooke@wrh.ox.ac.uk

Dr William Cooke

Nuffield Department of Women's and Reproductive Health, University of Oxford

Level 3, Women's Centre, John Radcliffe Hospital. Oxford, OX3 9DU, UK

<u>Affiliations</u>

1) Nuffield Department of Women's and Reproductive Health, University of Oxford,

Oxford, UK

2) Centre for Novostics, Hong Kong Science Park, Pak Shek Kok, New Territories,

Hong Kong SAR, China

3) Department of Chemical Pathology, The Chinese University of Hong Kong, Prince

of Wales Hospital, Shatin, New Territories, Hong Kong SAR, China

<u>Short title</u>

5'-tRNA Fragments in Preeclampsia Vesicles

<u>Total word count</u>

5931

Abstract

Background: The relationship between placental pathology and the maternal

syndrome of preeclampsia is incompletely characterised. Mismatch between

placental nutrient supply and fetal demands induces stress in the

syncytiotrophoblast, the layer of placenta in direct contact with maternal blood. Such

stress alters the content and increases the release of extracellular vesicles (STB-

EVs) into the maternal circulation. We have previously shown 5'-tRNA fragments (5'-

tRFs) constitute the majority of small RNA in STB-EVs in healthy pregnancy. 5'-tRFs

are produced in response to stress. We hypothesised STB-EV 5'-tRF release might

change in preeclampsia.

Methods: We perfused placentas from eight women with early-onset preeclampsia

and six controls, comparing small RNA expression in STB-EVs. We used

membrane-affinity columns to isolate maternal plasma vesicles and investigate

placental 5'-tRFs *in-vivo*. We quantified 5'-tRFs from circulating STB-EVs using a

placental alkaline phosphatase immunoassay. 5'-tRFs and scrambled RNA controls

were added to monocyte, macrophage and endothelial cells in culture to investigate

transcriptional responses.

Results: 5'-tRFs constitute the majority of small RNA in STB-EVs from both

preeclampsia and normal pregnancies. >900 small RNA fragments are differentially

expressed in preeclampsia STB-EVs. Preeclampsia-dysregulated 5'-tRFs are

detectable in maternal plasma, where we identified a placentally-derived load. 5'-

tRF-Glu-CTC, the most abundant preeclampsia-upregulated 5'-tRF in perfusion STB-

EVs, is also increased in preeclampsia STB-EVs from maternal plasma. 5'-tRF-Glu-

CTC induced inflammation in macrophages but not monocytes. The conditioned

media from 5''-tRF-Glu-CTC-activated macrophages reduced eNOS expression in

endothelial cells.

Conclusions: Increased release of syncytiotrophoblast-derived vesicle-bound 5'-tRF-

Glu-CTC contributes to preeclampsia pathophysiology.

<u>Abbreviations</u>

DOTAP   N-[1-(2,3-Dioleoyloxy)propyl]-N,N,N- trimethylammonium methylsulfate

EV   Extracellular vesicle

MLEV   Medium-large extracellular vesicle

PlGF   Placental growth factor

SEV   Small extracellular vesicle

STB-EV   Syncytiotrophoblast extracellular vesicle

sFlt-1   Soluble fms-like tyrosine kinase-1

tRF   tRNA fragment

Introduction

Preeclampsia is a complex placental syndrome, with multiple causes and a variable

phenotype. Clinical features are characterised by maternal sterile inflammation and

endothelial dysfunction[1]. A point of convergence in the disorder is stress in the

syncytiotrophoblast, the interface between maternal and fetal circulations[2]. The link

between syncytiotrophoblast stress and maternal symptoms, likely through blood-

borne factors, is incompletely characterised[3]. Angiogenic proteins (soluble fms-like

tyrosine kinase [sFlt-1] and placental growth factor [PlGF]) are important

syncytiotrophoblast stress signals which contribute to the preeclamptic syndrome[4].

Excess sFlt-1 sensitises endothelial cells to pro-inflammatory cytokines[5]. These

molecules have been successfully applied to the clinical diagnosis of preeclampsia[6].

Altered extracellular vesicle [EV] release from the syncytiotrophoblast, whilst also a

key contributor in the pathophysiology of preeclampsia, remains underexplored[7].

The healthy syncytiotrophoblast releases EVs [STB-EVs] directly into the maternal

circulation[7]. These lipid bilayer-bound particles are decorated with surface proteins

and shuttle their contents to distant cells. Cellular stress increases EV release; this is

reflected in preeclampsia where circulating STB-EVs are more abundant[8]. STB-EV

cargoes also change in preeclampsia; for example nitric oxide synthase expression

is reduced and neprilysin (a metalloprotease causing hypertension) is increased[9,10].

We recently reported 5'-transfer RNA fragments [5'-tRFs] as the predominant small

RNA species in healthy STB-EVs[11]. 5'-tRFs form when mature tRNA molecules are

cleaved by a number of stress-induced ribonucleases including angiogenin[12]. They

can be exported as EV cargo[13]. 5'-tRF expression profiles can be complex; at least

417 tRNA genes can be cleaved at multiple loci[14]. 5'-tRFs are also multifaceted

signalling molecules, regulating transcription, translation and epigenetic

inheritance[15]. 5'-tRFs have mostly been investigated in cancer biology and

immunology where they are described as intracellular, autocrine and paracrine

signals[13,16,17].

5'-tRFs are stress signals; syncytiotrophoblast stress is a key feature of

preeclampsia. We hypothesised that syncytiotrophoblast 5'-tRF release may change

in preeclampsia. We used placental perfusion as a source of STB-EVs to show that

5'-tRF expression in preeclampsia differed from healthy pregnancy. Our *in-vivo* work

demonstrated a placentally-derived load of circulating 5'-tRFs. 5'-tRF-Glu-CTC (the

most abundant preeclampsia-upregulated 5'-tRF) was increased in preeclampsia

plasma STB-EVs. Cell culture studies found 5'-tRF-Glu-CTC triggered sterile

inflammation in macrophages. Together these findings suggest 5'-tRFs may link

syncytiotrophoblast stress with maternal inflammation in preeclampsia.

Methods

1. Sample collection and storage

This project was approved by the Central Oxfordshire Research Ethics Committee

(07/H0607/74 and 07/H0606/148). All participants provided informed written consent.

Preeclampsia was defined using the ISSHP classification[18]. Placentas were obtained

at the time of caesarean section and perfused within 10 minutes of delivery. Uterine

vein samples were taken during caesarean section, just prior to uterine incision,

ipsilateral to the placental site. Peripheral blood samples were taken from the

antecubital fossa. 21-gauge needles and 4.5 ml sodium citrate vacutainers were

used for venepuncture (BD Diagnostics, UK). Non-pregnant samples were from

female volunteers of reproductive age. Plasma was obtained by centrifugation at

1500 g for 15 minutes. Samples were processed within 30 minutes of collection,

aliquoted and stored at -80°C.

2. Placental perfusion

Placentas from eight women with early-onset preeclampsia and six normotensive

pregnancies were perfused using a well-established dual-lobe perfusion technique[19].

Maternal perfusate was centrifuged at 10 000 g for 30 minutes to isolate medium-

large extracellular vesicles [MLEVs]. The supernatant was centrifuged at 150 000 g

for 2 hours to isolate small extracellular vesicles [SEVs]. Biopsies of placental tissue

were taken from the maternal surface of a non-perfused lobe. EVs were

characterised using Nanoparticle Tracking Analysis, transmission electron

microscopy and Western blotting as described previously[19].

3. RNA sequencing

RNA was isolated from MLEVs, SEVs and placental tissue using Total RNA

Purification Plus Kit (Norgen Biotek Corporation, Canada). After confirming RNA

quantity and integrity using Bioanalyser (Agilent Technologies, Germany), the same

amount of input RNA was loaded for library preparation using the NEBNext Multiplex

Small RNA library preparation kit (New England Biolabs, USA). Libraries were size-

selected for fragments 15-50 bp by gel electrophoresis; fragment size and

concentration was confirmed using High Sensitivity D1000 ScreenTape (Agilent,

UK). Single-end sequencing by synthesis was undertaken using an Illumina HiSeq

2500 machine (Illumina, USA). One preeclampsia sample was removed from the

SEV/placenta groups due to a technical issue.

Sequence reads were analysed using sRNAnalyzer[20]. Briefly, sequencing adaptors

and low-quality reads were removed using Cutadapt[21]. All identical reads in

sequence reads were collapsed, thus generating a set of unique reads (referred to

as fragment ID in this study). The number of sequence reads attributed to a fragment

ID were defined as the raw expression level of such a fragment ID. The fragment IDs

were mapped to the human small RNA databases allowing 2 mismatches. The

human small RNA databases comprised miRNA, piRNA, snoRNA, rRNA, and

tRNA[20,22,23]. The normalised expression level for each fragment ID within a class of

RNA was used for downstream differential expression analyses. The normalised

expression level for a fragment ID in a RNA class was calculated by dividing the raw

expression level by a per million scaling factor of total reads of that RNA class,

expressed as reads per million (RPM). Compared to healthy placentas, differentially

expressed fragment IDs in the MLEVs of preeclampsia placentas were determined

using the following criteria: i) At least one MLEV sample had expression level of

greater than 100 RPM, ii) P-value was required to be less than 0.05 based on Mann

Whitney test after Benjamini-Hochberg correction. The in-house bioinformatics

pipeline was written in Perl and R languages for counting sequence reads, tag

annotations, and differential expression analysis. Sequence read archive data for

blood cell datasets were downloaded from NCBI, with identifiers shown in

Supplementary Table 3.

4. Plasma EV RNA isolation

Plasma aliquots were thawed at 37°C and centrifuged at 3000 g for 5 minutes to

remove cryoprecipitates. For membrane-based affinity isolation of total vesicular

RNA, 500 µL plasma was loaded onto exoRNeasy™ Midi columns (Qiagen,

Germany). For magnetic-bead isolation of STB-EV RNA, 500 µL plasma was

centrifuged at 10 000 g for 30 minutes and the total EV pellet washed once before

resuspension with biotin-saturated MojoSort streptavidin magnetic nanobeads

(Biolegend, USA) to deplete non-specific binding. The supernatant was resuspended

with nanobeads pre-bound with biotinylated in-house placental alkaline phosphatase

(PLAP) antibody, known as NDOG2[24]. Bead-STB-EV complexes were washed four

times before EV RNA isolation using Trizol LS (Invitrogen, USA). RNA was stored in

aliquots at -80°C.

5. RT-qPCR detection

Custom Taqman™ stem-loop assays were designed for specific small RNA target

sequences identified from RNA sequencing analysis (Applied Biosystems, USA).

Assay linearity and specificity were verified. Perfusion samples were normalised to

TBP (confirmed empirically to be a suitable reference). Plasma samples were

normalised to *C. Elegans* miR-39 spike-in.

qPCR assays are documented in Supplementary Table 4. QuantStudio™ qPCR

instruments (Applied Biosystems, USA) automatically determined quantification

cycles (Cq) using standard settings; Cq >35 was considered undetectable. Relative

expression was determined by following the PFAFFL approach, normalising to

median expression in the control group.

6. Cell culture

THP-1 cells (ATCC, USA) were seeded onto 24-well Nunc plates at 50 000 cells per

well in RPMI-1640 medium supplemented with 10% Fetal Calf Serum. Cells were

grown with 6.2 ng/mL phorbol 12-myristate-13-acetate (PMA) for 24 hours to

differentiate into macrophages. Transfection experiments were performed using RNA

oligonucleotides (IDT, USA) with a 5'-P modification; the sequence for tRF-A is

shown in Supplementary Table 2. The scramble RNA control was the most abundant

STB-EV tRF without 5'-P modification and U nucleotides replaced with A (sequence

5'-GCAAAGGAGGAACAGAGGAAGAAAACACGCCA-3'). 9.2uM RNA was

packaged into lipid vesicles using N-[1-(2,3-Dioleoyloxy)propyl]-N,N,N-

trimethylammonium methylsulfate [DOTAP], following manufacturer's instructions

(Roche, Switzerland) and in line with a prior publication[17]. Human Umbilical Vein

Endothelial Cells were purchased and seeded onto 24-well Nunc plates at 25 000

cells per well in Endothelial Cell Growth Medium 2 (PromoCell, Germany).

Macrophage supernatant experiments were conducted by preparing 2X Growth

Medium and mixing 50:50 with THP-1 supernatant. Cellular RNA was isolated using

RNeasy™ Plus Mini Kit (Qiagen, Germany). Target mRNA expression was

quantified and normalised to *GAPDH.* Pre-amplification was used to detect IL-12B

using TaqMan™ master mix #4391128 with 10 cycles.

7. Data presentation

All data within this manuscript are derived from distinct samples. Figures 3B and 4a

were created using BioRender.com (Toronto, Canada). Statistical analyses and

figures were generated using RStudio (RStudio, USA) and Prism9 (GraphPad, USA).

Suspected outliers were excluded if over the 85[th] centile. Unpaired two-tailed Mann

Whitney tests were used throughout unless specifically stated within the figure

legend.

8. Data availability

Data supporting findings of this study and bioinformatic pipelines are available

through collaboration upon reasonable request to the corresponding author.

<u>Results</u>

The preeclamptic syncytiotrophoblast exports 5'-tRFs in EVs, mirroring

healthy pregnancy

We isolated SEVs and MLEVs, alongside placental biopsies, from eight placentas

with early-onset preeclampsia and six normotensive controls using dual-lobe

placental perfusion. Pregnancy characteristics are shown in Supplementary Table 1.

We performed single-end small RNA sequencing (size selecting <50 nucleotides).

Sequence length distribution plots confirmed the majority of reads in MLEVs and

SEVs from both preeclampsia and normal placentas were 30-34 nucleotides long

(Figure 1A). In contrast a peak at 22 nucleotides was greater in the placental

samples from both groups. The majority of fragments in MLEVs and SEVs mapped

to tRNA species, rather than to the ribosomal RNA and micro-RNA species seen in

placental tissues (Figure 1B). Coverage plots demonstrated that 5'-tRFs (but not 3'-

tRFs) constitute almost all tRNA reads in EVs in preeclampsia as well as normal

pregnancies (Figure 1C).

5'-tRFs are differentially expressed in STB-EVs from preeclampsia compared

to normotensive placentas

We used a bespoke bioinformatics pipeline to investigate differential expression of

tRFs. To minimise data loss, we assigned each unique fragment an identifier, then

annotated fragments after differential expression analysis (Methods). We identified

983 differentially expressed small RNA fragments in preeclampsia MLEVs compared

to controls; 182 mapped to 5'-tRFs using GtRNAdb 2.0[23]. No fragments were found

to be differentially expressed in preeclampsia SEVs.

The 12 most abundant differentially expressed fragments are shown in Table 1.

Using the sum of preeclampsia MLEV normalised counts as a denominator, these 12

fragments account for 64% of the differentially expressed counts. 5% of these counts

were accounted for by the 626 least abundant fragments. Thus, a small number of

abundant fragments represented the majority of the signal in an otherwise complex

dysregulated small RNA profile in preeclampsia MLEVs.

Comparing to human small RNA databases using conventional mapping, among the

12 most abundant differentially expressed fragments, 3 were identified as 5'-tRFs

and 4 as piwi-interacting RNAs. The remaining 5 had no directly matched reference

sequences. Review of fragment sequences demonstrated substantial overlap;

indeed all 12 showed only one or two nucleotide insertions or deletions differentiating

them from known 5'-tRFs (Table 1). By adapting the post-hoc mapping strategy to

incorporate up to two insertions or deletions, it was evident that the majority of

differentially expressed small RNA in preeclampsia are 5'-tRFs. A complex profile of

differential expression in preeclampsia MLEVs was thus dominated by minor variants

of a handful of abundant 5'-tRFs.

We sought three target 5'-tRFs which were differentially expressed in preeclampsia

STB-EVs to validate our findings *ex-vivo* (in placental perfusion samples) and *in-vivo*

(in plasma). 5'-tRFs are known to be expressed in other circulating EVs, most

notably from immune cells. The most abundant EVs in plasma are blood-cell derived.

Hence, we used publicly available blood-cell datasets to quantify possible target 5'-

tRFs likely to be within contaminating EVs in plasma. Using these data we selected

three tRFs for validation which were: abundant in STB-EVs (above 90[th] centile PE

expression amongst 983 differentially expressed fragments), upregulated in

preeclampsia, and of low relative abundance in potentially contaminating EV source

cells (Supplementary Table 2). The relative abundance of these three targets in

STB-EVs using RNA sequencing is shown in Figure 2A (5'-tRF targets named A, B

and C for brevity but full sequences and tRNA derivations shown in Supplementary

Table 2).

We used custom small RNA assays with stem-looped reverse transcription primers

and specific minor-groove binding TaqMan™ probes to compare relative 5'-tRF

expression in STB-EVs obtained by perfusion using quantitative real-time

polymerase chain reaction. Results validated target 5'-tRF upregulation and effect

sizes were consistent with RNA sequencing findings (median 1.4-fold upregulated in

preeclampsia, Figure 2B).

A proportion of 5'-tRFs are pregnancy-specific and placentally-derived in the

maternal circulation

Total EV RNA was isolated from maternal plasma using membrane-affinity columns.

EV size profiles for plasma were comparable to perfusion MLEVs (Supplementary

Figure 1). Extensive characterisation of plasma membrane-affinity EVs has

previously been published[25]. Target 5'-tRFs were significantly more abundant in

pregnant peripheral plasma EVs than in non-pregnant matched control samples

(median 3.0-fold higher in pregnancy, p<0.05 for all three) (Figure 3A). The

difference between these sample groups suggested a pregnancy-specific load of 5'-

tRFs in plasma EVs. The high abundance of these 5'-tRFs in non-pregnant samples

confirms they were not unique to pregnancy (median Cq values in non-pregnant

samples: A 24.0, B 27.9, C 25.9). We then acquired paired plasma samples

simultaneously from the uterine and peripheral veins of women without preeclampsia

undergoing elective caesarean section for an indication unrelated to preeclampsia

(e.g. breech presentation) prior to delivery of the feto-placental unit. The uterine vein

directly receives blood from the placenta, thus placentally-derived molecules are

more abundant in these samples (Figure 3B)[26]. All three 5'-tRFs were more

abundant in uterine vein samples (median 1.3-fold, $p < 0.05$ for all three) (Figure 3C).

These data support a placentally-derived load of 5'-tRFs in circulating plasma EVs.

STB-EV 5'-tRF-Glu-CTC, the most abundant preeclampsia-upregulated 5'-tRF,

is increased in preeclampsia maternal plasma

A technique was optimised (Figure 4A) to isolate STB-EV RNA from maternal

plasma, targeting the syncytiotrophoblast marker protein placental alkaline

phosphatase (PLAP). Streptavidin nanobeads (130 nm diameter) were incubated

with a highly specific biotinylated anti-PLAP antibody (NDOG2, in-house). PLAP+

plasma MLEVs were separated from soluble PLAP by centrifugation and NDOG2-

nanobeads used to pull STB-EVs from total plasma EVs. Expression of miR518

(from the placental chromosome 19 microRNA cluster) was used to demonstrate

assay sensitivity (Figure 4B). Perfusion STB-EVs were spiked into non-pregnant

plasma as a positive control, achieving around 2000-fold greater miR518 expression

than pregnant plasma. Nanobeads without antibody (saturated with free biotin) were

added to pregnant plasma as a negative control; no miR518 expression was

detected.

This technique was used to quantify the abundance of 5'-tRF-Glu-CTC (tRF-A) in

peripheral plasma from 14 women with early-onset preeclampsia and 12 gestation-

matched normotensive controls (Supplementary Table 5). 5'-tRF-Glu-CTC was

upregulated (median 1.4-fold, p=0.017) in preeclampsia plasma STB-EVs (Figure

4C). The effect size was comparable to preeclampsia perfusion STB-EVs (median

1.4-fold upregulated).

EV-bound 5'-tRF-Glu-CTC promotes macrophage, but not monocyte activation

Inflammation is a key feature of preeclampsia[27]. Tissue-resident macrophages are

known to be activated in preeclampsia[28]. Recent studies suggest STB-EVs may

underlie this activation[29]. We treated macrophages and monocytes in culture with the

most abundant 5'-tRF upregulated in preeclampsia (5'-tRF-Glu-CTC, labelled A for

brevity; full sequence in Supplementary Table 2). RNA was packaged within

otherwise undecorated lipid vesicles, in order to distinguish the effect of one tRF

from the accompanying RNA, lipids and proteins in perfusion-derived STB-EVs. We

used a scrambled version of the most abundant STB-EV 5'-tRF sequence as a

negative control, following validation against untreated cells (Supplementary Figure

2). Macrophages were activated to a type 1 phenotype after 12 hours treatment with

tRF-A, increasing expression of pro-inflammatory cytokines (Figure 5A). This pro-

inflammatory 5'-tRF action was confined to macrophages, with no changes observed

when the same experiment was repeated in undifferentiated monocytes (Figure 5B).

EV-bound 5'-tRF-Glu-CTC indirectly activates endothelial cell adhesion

molecules and reduces expression of endothelial nitric oxide synthase

Endothelial damage is considered a common factor in the wide-ranging maternal

organ dysfunction of preeclampsia[1]. Vascular macrophages are known to modulate

adjacent endothelial cell function[30]. We treated human umbilical vein endothelial

cells with vesicle-bound tRF-A (direct) and with the supernatants from tRF-A-

activated macrophages (indirect) alongside corresponding scramble controls. tRF-A

activated endothelial cells (quantified through increased expression of adhesion

molecules) indirectly through macrophage stimulation, but not directly (Figure 5C).

The primary presenting feature of preeclampsia is high blood pressure. Immediate

blood pressure regulation involves constitutive release of nitric oxide from endothelial

cells, which maintains a state of relative vasodilation. In preeclampsia, circulating

nitrite levels are reduced, which may contribute to hypertension[31]. tRF-A reduced

expression of endothelial nitric oxide synthase in endothelial cells, indirectly via

macrophage activation (Figure 5C).

Discussion

We are the first to report 5'-tRFs as the predominant species of small RNA in

preeclampsia STB-EVs. These data are consistent with our prior finding in healthy

pregnancy STB-EVs[11]. Enrichment of 5'- (but not 3'-) tRFs in EVs is also reported in

immune cells and suggests a specific export process[13,17]. We demonstrate

differential expression of over 900 STB-EV small RNA fragments in early-onset

preeclampsia compared to normal pregnancy. Different preeclampsia phenotypes

are unified by stress in the syncytiotrophoblast[3]; tRFs are produced by stress-

dependent ribonucleases[12]. Thus, a change to STB-EV 5'-tRF expression fits with

our existing understanding of preeclampsia. Differential expression in preeclampsia

was identified in MLEVs but not SEVs, which is consistent with different EV

biogenesis: SEVs are released constitutively via the endosomal pathway whereas

MLEVs form by budding in response to stress[32]. The effect size between

preeclampsia and normal is consistent with other studies of differential 5'-tRF

expression in disease[33]. We corroborated discoveries in perfusion data by finding

increased STB-EV 5'-tRF-Glu-CTC in preeclampsia plasma compared to

normotensive controls.

One of the most studied tRNA ribonucleases is angiogenin, which is reported to

generate 2-3 phosphate residues at the 3' end of the 5'-tRF. Our sequencing

approach detected 5'-tRF with hydroxyl groups, but not 2-3 cyclic phosphate

residues, suggesting STB-EV 5'-tRFs were generated by ribonucleases other than

angiogenin[34].

5'-tRFs are known to influence cellular function through a variety of regulatory

mechanisms at the level of the transcriptome, translatome and the epigenome[15]. We

found 5'-tRF-Glu-CTC directly activated macrophages, but not monocytes or

umbilical vein endothelial cells. We speculate this difference may be accounted for

by phagocytosis of EVs by macrophages, trafficking 5'-tRFs to the endosomal

compartment (usually free of nucleic acids) where they could encounter Toll-like

receptor 7. This hypothesis is founded in published work demonstrating a lack of

macrophage response to unencapsulated 5'-tRFs, or 5'-tRFs with Toll-like receptor 7

antagonists and warrants further investigation[17]. Macrophages are not typically in

direct contact with blood; however in preeclampsia the endothelial barrier is

significantly disrupted[1]. We suggest that circulating EV-bound 5'-tRFs would directly

reach macrophages in the vessel walls in preeclampsia, where their functional effect

could contribute to the well-described sterile inflammation of the disease[27]. Our

findings of increased pro-inflammatory cytokine expression in response to tRF-Glu-

CTC correlate with plasma cytokine concentrations in preeclampsia[35].

STB-EVs are known to directly damage the endothelium, yet we found no direct

effect of 5'-tRF-Glu-CTC on human umbilical vein endothelial cells[36]. This

discrepancy may be attributed to the absence of other EV RNA and proteins which

could be necessary to trigger endothelial damage. Our data show 5'-tRF-Glu-CTC

macrophage activation indirectly activates the endothelium.

Prior studies of 5'-tRFs have used cell culture as a model system. Work in breast

cancer reported intracellular 5'-tRF expression promoted metastasis[16].

Mycobacterium infection in human macrophages triggered 5'-tRF release in EVs,

activating neighbouring cells[17]. Here we have investigated 5'-tRFs at a whole-organ

level: the placenta is expelled with the fetus during parturition and can be studied

intact *ex-vivo*. Sampling of the uterine vein during caesarean section has offered *in-*

vivo evidence for a placental 5'-tRF load in maternal plasma. An immuno-assay has

corroborated STB-EV 5'-tRF-Glu-CTC upregulation in preeclampsia plasma.

Together with functional data showing 5'-tRF-Glu-CTC macrophage activation, we

propose a possible endocrine signalling function for 5'-tRFs, contributing to

preeclampsia pathogenesis.

Our study's strength lies in the unique integration of cutting-edge techniques and

distinctive samples. Previous studies investigating STB-EV small RNA have used

lower fidelity models to source EVs (e.g. explants) and taken bioinformatic

approaches which disregard 5'-tRF data, despite noting their presence[37,38].

We have corroborated perfusion-based RNA sequencing discoveries *in vivo* using

qPCR in plasma. We overcame confounding in high-profile studies of total cell-free

RNA in preeclampsia by focussing our attention exclusively on placental RNA[39,40].

The smaller size of our discovery cohort could be considered a weakness in a

heterogeneous disease; we consider that by confining ourselves to a common step

(syncytiotrophoblast stress) in early-onset disease, as well as confirming our data *in*

vivo and *in vitro*, our findings are pertinent.

Perspectives

Preeclampsia is a multifactorial condition, with diverse clinical features.

Syncytiotrophoblast stress is common to all cases, but remains poorly understood.

Here we present 5'-tRFs, a novel and highly abundant class of RNA differentially

released by the preeclamptic syncytiotrophoblast. We demonstrate placentally-

derived 5'-tRFs in the maternal circulation. The most abundant preeclampsia-

upregulated STB-EV 5'-tRF was also increased in preeclampsia plasma. We find

pro-inflammatory effects of this tRF on macrophages. Together these data suggest

5'-tRFs may play a role as transducers of an inflammatory signal from placenta to

periphery in preeclampsia. Our findings offer a novel category of signalling molecule

released by the placenta, warranting further investigation. We speculate that 5'-tRFs

may dysregulate other cells in preeclampsia. Future studies will consider: other

putative 5'-tRF targets in preeclampsia, such as liver sinusoids and pericytes; the

actions of additional 5'-tRFs which are differentially expressed in preeclampsia;

whether STB-EV 5'-tRFs play a role in other pregnancy-related diseases. Our

ongoing work also focuses on the optimisation of techniques to isolate low-

abundance placental EV 5'-tRF signal from complex biofluids such as plasma, with

attention to their clinical relevance. 5'-tRFs may join other better-studied stress

markers such as sFlt-1 and PlGF in explaining the pathogenesis of preeclampsia.

Novelty and Relevance

1. What is New?

   5'-tRFs are the most abundant small RNA species within preeclampsia STB-EVs

   and are differentially expressed compared to normal pregnancy. A proportion of

   5'-tRFs in the maternal circulation are placentally-derived. The most abundant

upregulated 5'-tRF in preeclampsia STBEVs is 5'-tRF-Glu-CTC; this has been

discovered by placental perfusion and corroborated in plasma. 5'-tRF-Glu-CTC

has pro-inflammatory effects on macrophages.

2. What is Relevant?

Preeclampsia is a placentally-derived hypertensive disorder, which can result in

multi-organ failure. Recent studies report STB-EVs could promote preeclampsia

through macrophage activation. Our findings suggest 5'-tRFs may underlie some

pro-inflammatory actions of STB-EVs in preeclampsia.

3. Clinical/Pathophysiological Implications?

STB-EV 5'-tRFs represent a feto-maternal signal with sufficient complexity to

contribute to the varied clinical features of preeclampsia. 5'-tRFs may represent a

biomarker or therapeutic target in this syndrome.

<u>Acknowledgements</u>

We wish to thank Fenella Roseman, research midwife, for assistance with collecting

clinical samples.

<u>Supplemental Material</u>

Figures S1-S2

Tables S1-S5

<u>Sources of Funding</u>

WRC receives a clinical doctoral training fellowship from the Wellcome Trust, which

funded this study.

<u>Disclosures</u>

YMDL holds equity in DRA, Insighta Grail/Illumina and Take2. PJ holds equity in

Illumina. PJ is a consultant to Take2. PJ is a Director of DRA and KingMed Future.

YMDL, PJ, and LJ receives royalties from Illumina, LabCorp, Grail, DRA, Xcelom

and Take2.

<u>Author contributions</u>

Study was conceived and designed by WRC, CR and MV. Bioinformatic analyses

were performed by PJ, LJ, JB, WRC and GDJ. Supervision was provided by PJ,

YMDL, CR and MV. Experimental work was performed and manuscript written by

WRC. All authors edited and approved the final manuscript.

References

1.   Powe CE, Levine RJ, Karumanchi SA. Preeclampsia, a disease of the

489   maternal endothelium: the role of antiangiogenic factors and implications for

490   later cardiovascular disease. *Circulation*. 2011;123:2856–2869.

491   2.   Magee LA, Nicolaides KH, von Dadelszen P. Preeclampsia. *N. Engl. J. Med.*

492   2022;386:1817–1832.

493   3.   Redman CWG, Staff AC, Roberts JM. Syncytiotrophoblast stress in

494   preeclampsia: the convergence point for multiple pathways. Am. J. Obstet.

495   Gynecol. 2022;226:S907–S927.

496   4.   Levine RJ, Maynard SE, Qian C, Lim K-H, England LJ, Yu KF, Schisterman

497   EF, Thadhani R, Sachs BP, Epstein FH, et al. Circulating Angiogenic Factors

498   and the Risk of Preeclampsia. *N. Engl. J. Med.* 2004;350:672–683.

499   5.   Cindrova-Davies T, Sanders DA, Burton GJ, Charnock-Jones DS. Soluble

500   FLT1 sensitizes endothelial cells to inflammatory cytokines by antagonizing

501   VEGF receptor-mediated signalling. *Cardiovasc. Res.* 2011;89:671–679.

502   6.   Zeisler H, Llurba E, Chantraine F, Vatish M, Staff AC, Sennström M, Olovsson

503   M, Brennecke SP, Stepan H, Allegranza D, et al. Predictive Value of the sFlt-

504   1:PlGF Ratio in Women with Suspected Preeclampsia. *N. Engl. J. Med.*

505   2016;374:13–22.

506   7.   Cooke WR, Jones GD, Redman CWG, Vatish M. Syncytiotrophoblast Derived

507   Extracellular Vesicles in Relation to Pre-eclampsia. *Matern. Med.* 2021;

508   8.   Knight M, Redman CWG, Linton EA, Sargent IL. Shedding of

509   syncytiotrophoblast microvilli into the maternal circulation in pre-eclamptic

510   pregnancies. *BJOG*. 1998;105:632–640.

9.   Motta-Mejia C, Kandzija N, Zhang W, Mhlomi V, Cerdeira AS, Burdujan A,

Tannetta D, Dragovic R, Sargent IL, Redman CW, et al. Placental Vesicles

Carry Active Endothelial Nitric Oxide Synthase and Their Activity is Reduced in

Preeclampsia. *Hypertension*. 2017;70:372–381.

10. Gill M, Motta-Mejia C, Kandzija N, Cooke W, Zhang W, Cerdeira AS, Bastie C,

Redman C, Vatish M. Placental Syncytiotrophoblast-Derived Extracellular

Vesicles Carry Active NEP (Neprilysin) and Are Increased in Preeclampsia.

Hypertension. 2019;73:1112–1119.

11. Cooke WR, Cribbs A, Zhang W, Kandzija N, Motta-Mejia C, Dombi E, Ri R,

Cerdeira AS, Redman C, Vatish M. Maternal circulating syncytiotrophoblast-

derived extracellular vesicles contain biologically active 5'-tRNA halves.

Biochem. Biophys. Res. Commun. 2019;518:107–113.

12. Li G, Manning AC, Bagi A, Yang X, Gokulnath P, Spanos M, Howard J, Chan

PP, Sweeney T, Kitchen R, et al. Distinct Stress-Dependent Signatures of

Cellular and Extracellular tRNA-Derived Small RNAs. *Adv. Sci.* 2022;9.

13. Chiou N-T, Kageyama R, Ansel KM. Selective Export into Extracellular

Vesicles and Function of tRNA Fragments during T Cell Activation. *Cell Rep.*

2018;25:3356-3370.e4.

14. Lyons SM, Fay MM, Ivanov P. The role of RNA modifications in the regulation

of tRNA cleavage. FEBS Lett. 2018;592:2828–2844.

15. Di Fazio A, Gullerova M. An old friend with a new face: tRNA-derived small

RNAs with big regulatory potential in cancer biology. *Br. J. Cancer.* 2023;1–11.

16. Liu X, Mei W, Padmanaban V, Alwaseem H, Molina H, Passarelli MC, Tavora

B, Tavazoie SF. A pro-metastatic tRNA fragment drives Nucleolin

oligomerization and stabilization of its bound metabolic mRNAs. *Mol. Cell.*

2022;82:2604-2617.e8.

17.   Pawar K, Shigematsu M, Sharbati S, Kirino Y. Infection-induced 5'-half

molecules of tRNAHisGUG activate Toll-like receptor 7. *PLoS Biol.* 2020;18.

18.   Brown MA, Magee LA, Kenny LC, Karumanchi SA, McCarthy FP, Saito S, Hall

DR, Warren CE, Adoyi G, Ishaku S. Hypertensive disorders of pregnancy:

ISSHP classification, diagnosis, and management recommendations for

international practice. *Hypertension.* 2018;72:24–43.

19.   Dragovic RA, Collett GP, Hole P, Ferguson DJP, Redman CW, Sargent IL,

Tannetta DS. Isolation of syncytiotrophoblast microvesicles and exosomes and

their characterisation by multicolour flow cytometry and fluorescence

Nanoparticle Tracking Analysis. *Methods.* 2015;87:64–74.

20.   Wu X, Kim TK, Baxter D, Scherler K, Gordon A, Fong O, Etheridge A, Galas

DJ, Wang K. sRNAnalyzer-a flexible and customizable small RNA sequencing

data analysis pipeline. *Nucleic Acids Res.* 2017;45:12140–12151.

21.   Martin M. Cutadapt removes adapter sequences from high-throughput

sequencing reads. *EMBnet.journal.* 2011;17:10.

22.   Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J,

Glöckner FO. The SILVA ribosomal RNA gene database project: improved

data processing and web-based tools. *Nucleic Acids Res.* 2013;41.

23.   Chan PP, Lowe TM. GtRNAdb 2.0: An expanded database of transfer RNA

genes identified in complete and draft genomes. *Nucleic Acids Res.*

2016;44:D184–D189.

24.   Sunderland CA, Davies JO, Stirrat GM. Immunohistology of normal and

ovarian cancer tissue with a monoclonal antibody to placental alkaline

phosphatase. *Cancer Res.* 1984;44:4496–4502.

25.   Stranska R, Gysbrechts L, Wouters J, Vermeersch P, Bloch K, Dierickx D,

Andrei G, Snoeck R. Comparison of membrane affinity-based method with size-exclusion chromatography for isolation of exosome-like vesicles from human plasma. *J. Transl. Med.* 2018;16:1–9.

26. Cerdeira AS, Kandzija N, Pargmae P, Cooke W, James T, Redman C, Vatish M. Circulating soluble fms-like tyrosine kinase-1 is placentally derived in normal pregnancy: First in vivo evidence. *Pregnancy Hypertens.* 2019;16:145–147.

27. Redman CWG, Sacks GP, Sargent IL. Preeclampsia: An excessive maternal inflammatory response to pregnancy. *Am. J. Obstet. Gynecol.* 1999;180:499–506.

28. Huda SS, Jordan F, Bray J, Love G, Payne R, Sattar N, Freeman DJ. Visceral adipose tissue activated macrophage content and inflammatory adipokine secretion is higher in pre-eclampsia than in healthy pregnancys. *Clin. Sci.* 2017;131:1529.

29. Liu X, Fei H, Yang C, Wang J, Zhu X, Yang A, Shi Z, Jin X, Yang F, Wu D, et al. Trophoblast-Derived Extracellular Vesicles Promote Preeclampsia by Regulating Macrophage Polarization. *Hypertension.* 2022;79:2274–2287.

30. Hernandez GE, Iruela-Arispe ML. The many flavors of monocyte / macrophage-endothelial cell interactions. *Curr. Opin. Hematol.* 2020;27:181.

31. Sandrim VC, Palei ACT, Metzger IF, Gomes VA, Cavalli RC, Tanus-Santos JE. Nitric oxide formation is inversely related to serum levels of antiangiogenic factors soluble fms-like tyrosine kinase-1 and soluble endoglin in preeclampsia. *Hypertension.* 2008;52:402–407.

32. Teng F, Fussenegger M. Shedding Light on Extracellular Vesicle Biogenesis and Bioengineering. *Adv. Sci.* 2021;8:2003505.

33. Meng L, Jiang L, Chen J, Ren H, Gao Z, Wu F, Wen Y, Yang L. Transfer RNA-

derived fragment tRF-28-QSZ34KRQ590K in plasma exosomes may be a

potential biomarker for atopic dermatitis in pediatric patients. *Exp. Ther. Med.*

2021;21:489.

34. Shigematsu M, Kawamura T, Kirino Y. Generation of 2′,3′-Cyclic Phosphate-

Containing RNAs as a Hidden Layer of the Transcriptome. *Front. Genet.*

2018;9.

35. Kupferminc MJ, Peaceman AM, Aderka D, Wallach D, Socol ML. Soluble

tumor necrosis factor receptors and interleukin-6 levels in patients with severe

preeclampsia. *Obstet. Gynecol.* 1996;88:420–427.

36. Gupta AK, Rusterholz C, Huppertz B, Malek A, Schneider H, Holzgreve W,

Hahn S. A comparative study of the effect of three different syncytiotrophoblast

micro-particles preparations on endothelial cells. *Placenta.* 2005;26:59–66.

37. Salomon C, Guanzon D, Scholz-Romero K, Longo S, Correa P, Illanes SE,

Rice GE. Placental Exosomes as Early Biomarker of Preeclampsia: Potential

Role of Exosomal MicroRNAs Across Gestation. *J. Clin. Endocrinol. Metab.*

2017;102:3182–3194.

38. Wei J, Blenkiron C, Tsai P, James JL, Chen Q, Stone PR, Chamley LW.

Placental trophoblast debris mediated feto-maternal signalling via small RNA

delivery: Implications for preeclampsia. *Sci. Rep.* 2017;7:1–12.

39. Rasmussen M, Reddy M, Nolan R, Camunas-Soler J, Khodursky A, Scheller

NM, Cantonwine DE, Engelbrechtsen L, Mi JD, Dutta A, et al. RNA profiles

reveal signatures of future health and disease in pregnancy. *Nat. 2022*

6017893. 2022;601:422–427.

40. Moufarrej MN, Vorperian SK, Wong RJ, Campos AA, Quaintance CC, Sit R V.,

Tan M, Detweiler AM, Mekonen H, Neff NF, et al. Early prediction of

preeclampsia in pregnancy with cell-free RNA. *Nat. 2022 6027898*.

2022;602:689–694.

Tables

Table 1

Rank	Sequence	Length	Mapping no indel	Mapping ≤2 indel	Mean PE count	SD PE count	Mean N count	SD N count	Log2Fold Change	Adjusted p value	% PE DE reads
1	GCATTGGTGGTTCAGTGGTAGAATTCTCGCCT	32	tRF-Gly-GCC	tRF-Gly-GCC	273954	18135	305199	17604	-0.16	0.0335	24.5
2	GCATTGTGGTTCAGTGGTAGAATTCTCGCCT	31	Unmatch	tRF-Gly-GCC	102499	14060	161979	28840	-0.66	0.0058	9.2
3	TCCCTGTGGTCTAGTGGTTAGGATTCGGCGCT	32	Unmatch	tRF-Glu-CTC	91226	23407	42503	8037	1.10	0.0058	8.2
4	GCATTGGTGGTTCAGTGGTAGAATTCTCGCCC	32	tRF-Gly-GCC	tRF-Gly-GCC	62934	10260	78646	6927	-0.32	0.0150	5.6
5	TCCCTGGTGGTCTAGTGGTTAGGATTCGGCGCT	33	tRF-Glu-CTC	tRF-Glu-CTC	52051	18436	18405	5423	1.50	0.0054	4.7
6	TCCCTGGTGGTCTAGTGGTTAGGATTCGGCGC	32	piR-hsa-5938	tRF-Glu-CTC	27597	6781	11286	2564	1.29	0.0085	2.5
7	GCATTGTGGTTCAGTGGTAGAATTCTCGCCC	31	Unmatch	tRF-Gly-GCC	24069	6717	41872	9820	-0.80	0.0058	2.2
8	TCCCTGTGGTCTAGTGGTTAGGATTCGGCGCC	32	Unmatch	tRF-Glu-CTC	20922	2939	11389	1846	0.88	0.0058	1.9
9	GCATTGTGGTTCAGTGGTAGAATTCTCGCC	30	Unmatch	tRF-Gly-GCC	20992	4826	32864	7785	-0.65	0.0305	1.9
10	TCCCTGGTGGTCTAGTGGTTAGGATTCGGCG	31	piR-hsa-5938	tRF-Glu-CTC	15003	6543	3282	922	2.19	0.0085	1.3
11	GCCCGGCTAGCTCAGTCGGTAGAGCATGAGAC	32	piR-hsa-27622	tRF-Lys-CTT	12391	2710	20600	4403	-0.73	0.0220	1.1
12	GTTTCCGTAGTGTAGTGGTTATCACGTTCGC	31	piR-hsa-28877	tRF-Val-AAC	11802	2537	21618	6164	-0.87	0.0128	1.1

Descriptors of twelve most abundant differentially expressed small RNA fragments in

early-onset preeclampsia (PE) perfusion STB-EVs compared to normotensive (N)

perfusion STB-EVs, ranked by abundance. Rows showing targets downregulated in

preeclampsia are shaded grey. Mapping is shown with no insertions or deletions

(indel) or permitting up to two. Standard deviation (SD), differentially expressed (DE).

Figures with Figure Legends

Figure 1

Assessment of reads <50 base pairs from medium-large extracellular vesicles

(MLEV), small extracellular vesicles (SEV) and placental samples obtained from

placentas in normotensive pregnancy (Normal, n=6) and early-onset preeclampsia

(PE, n=8).

A: Sequence length distribution plots showing size in base pairs (bp) for small RNA

fragments after removal of adaptors and low-quality reads. B: Mapping of small RNA

fragments. C: Coverage plots showing percentage of per-base coverage for mapped

tRNA fragments.

Figure 2

Expression of three target small RNA fragments in early-onset preeclampsia and

normotensive pregnancy STB-EVs obtained by placental perfusion.

A: Quantified using RNA sequencing (n=6 normotensive, n=8 preeclampsia; Mann

Whitney tests after Benjamini-Hochberg correction displayed).

B: Quantified using stem-loop qPCR (normalised to reference gene TBP, n=4 per

group).

Boxes show median/interquartile range; whiskers show max/min. Fragments labelled

A, B, C for brevity; full sequences in Supplementary Table 2.

Figure 3

Detection of three target small RNA fragments in plasma using qPCR

A: Expression in total EVs isolated from peripheral venous plasma from seven

healthy third trimester pregnancies and six female volunteers of reproductive age

(boxes show median/interquartile range; whiskers show max/min).

B: Diagram demonstrating sampling rationale for uterine and peripheral venous

blood

C: Expression in total EVs isolated from uterine and peripheral venous plasma taken

simultaneously from five healthy term pregnancies (paired one-tailed Wilcoxon tests

displayed).

Fragments labelled A, B, C for brevity; full sequences in Supplementary Table 2.

Figure 4

Isolation of STB-EVs from venous plasma

A: Diagram summarising protocol for STB-EV isolation: plasma preparation, 10 000

g centrifugation to isolate MLEVs, immuno-isolation using magnetic beads coated

with antibody to placental alkaline phosphatase (NDOG2).

B: Relative expression of miR518 in EVs isolated from three samples using above

technique (from left to right: pregnant plasma using beads coated with NDOG2; non-

pregnant plasma spiked with perfusion STB-EVs using beads coated with NDOG2;

pregnant plasma using beads without NDOG2 coating). Bars represent median, error

bars represent interquartile range.

C: Relative expression of 5'-tRF-Glu-CTC (tRF-A) in perfusion-derived STB-EVs (not

gestation-matched, reproduced from Figure 2B for comparison) and gestation-

matched peripheral venous plasma STB-EVs from women with early-onset

preeclampsia (n=14) and normotensive pregnancies (n=12).

Figure 5

5'-tRF-Glu-CTC (tRF-A) actions in immune and endothelial cells.

A: Cytokine expression by THP-1 derived macrophages after treatment with vesicle-

encapsulated RNA for 12 hours.

B: Cytokine expression by THP-1 monocytes (undifferentiated) after treatment with

vesicle-encapsulated RNA for 12 hours.

C: Expression of cell adhesion molecule and endothelial nitric oxide synthase mRNA

65   in human umbilical vein endothelial cells treated with vesicle-encapsulated RNA

66   (Direct), or the cell culture medium from macrophages pre-treated with vesicle-

67   encapsulated RNA (Indirect) for 6 hours.

68   Fragments labelled tRF-A and Scramble for brevity; full sequences in Supplementary

69   Table 2 and Methods; n=4 per condition.