Physcomitrium patens SMXL homologs are PpMAX2-dependent negative regulators of growth

Ambre Guillory[1,2,*], Mauricio Lopez-Obando[1,3], Khalissa Bouchenine[1], Philippe Le Bris[1], Alain Lécureuil[1], Jean-Paul Pillot[1], Vincent Steinmetz[4], François-Didier Boyer[4], Catherine Rameau[1], Alexandre de Saint Germain[1], Sandrine Bonhomme[1,*]

[1]Université Paris-Saclay, INRAE, AgroParisTech, Institut Jean-Pierre Bourgin (IJPB), 78000, Versailles, France.

[2]LIPME, Université de Toulouse, INRAE, CNRS, Castanet-Tolosan 31326, France.

[3]Institut de biologie moléculaire des plantes (IBMP), CNRS, University of Strasbourg,12 rue du Général Zimmer, Strasbourg, France.

[4]Université Paris-Saclay, CNRS, Institut de Chimie des Substances Naturelles, UPR 2301, 91198, Gif-sur-Yvette, France.

[*]Corresponding authors: sandrine.bonhomme@inrae.fr and ambre.guillory@inrae.fr

Short title

SMXLs are negative regulators of the MAX2-dependent pathway in *Physcomitrium patens*

Abstract

SMXL proteins are a plant-specific clade of type I HSP100/Clp-ATPases. *SMXL* genes are found in virtually all land plants' genomes. However, they have mainly been studied in angiosperms. In *Arabidopsis thaliana*, three SMXL functional subclades have been identified: SMAX1/SMXL2, SMXL345 and SMXL678. Out of these, two subclades ensure transduction on endogenous hormone signals: SMAX1/SMXL2 are involved in KAI2-ligand (KL) signaling, while SMXL678 are involved in strigolactones (SLs) signaling. Many questions remain regarding the mode of action of these proteins, as well as their ancestral role. In light of recent discoveries in the liverwort *Marchantia polymorpha,* we addressed this second question by investigating the function of the four *SMXL* genes of the moss *Physcomitrium patens*. We demonstrate that PpSMXL proteins are negative regulators of growth, involved in the likely conserved ancestral MAX2-dependent KL signaling pathway. However, PpSMXL proteins expressed in *Arabidopsis thaliana* unexpectedly cannot replace SMAX1/SMXL2 function in KL signaling, whereas they can functionally replace SMXL4/5 and restore root growth. Therefore, the molecular function of SMXL could be conserved, but not their interaction network. Moreover, one PpSMXL clade also positively regulates transduction of the SL signal in *P. patens*, this function most probably having an independent evolutionary origin to angiosperms SMXL678.

Introduction

Strigolactones (SLs) are butenolide compounds with an early origin in land plant evolution (Delaux et al. 2012; Kyozuka et al. 2022; Bonhomme and Guillory 2022). In angiosperms, these molecules have first been identified as rhizospheric signals with both negative and positive outcomes for the producing plant: SLs stimulate seed germination of root parasitic plants (Cook et al. 1966) but also promote Arbuscular Mycorrhizal (AM) symbiosis (Akiyama et al. 2005) by boosting AM fungi mitochondrial metabolism and thus hyphae growth (Besserer et al. 2006; Besserer et al. 2008). SLs are also employed as phytohormones in angiosperms, where they play diverse roles, in particular regulating plant architecture (for review see (Machin et al. 2020)). Notably, SLs have a largely documented ability to repress axillary branching, by inhibiting axillary bud activity (Gomez-Roldan et al. 2008; Umehara et al. 2008; Jiang et al. 2013; Kerr et al. 2021).

Phylogenetic studies suggest that the SL biosynthesis pathway is ancient, genes encoding most SL biosynthesis hormones being found both in land plants and some algae (Delaux et al. 2012; Walker et al. 2019). The CAROTENOID CLEAVAGE DIOXYGENASE 8 (CCD8) enzyme is essential for SL biosynthesis, since its product, carlactone (CL), is considered to be a key precursor for SLs (Alder et al. 2012). Genetic studies have demonstrated that CCD8 function, first established in angiosperms models, is probably conserved in bryophytes, as this function was demonstrated at least in the moss *Physcomitrium patens* (*P. patens*) (Proust et al. 2011; Decker et al. 2017) and in the liverwort *Marchantia paleacea* (Radhakrishnan et al. 2020; Kodama et al. 2022). However, this conservation has not been proven yet for algae CCD8 homologs (Walker et al. 2019) and so far, SL synthesis has only been described in land plants. In *P. patens*, SLs have a hormonal function, since PpCCD8-derived compounds repress filament branching and elongation (Proust et al. 2011; Hoffmann et al. 2014), repress gametophore (leafy shoot) basal branching (Coudert et al. 2015) and enhance resistance to phytopathogenic fungi (Decker et al. 2017). Still, the exact molecule with a hormonal role has not been identified yet (Lopez-Obando et al. 2021). In contrast, in *M. paleacea*, the recently characterized SL Bryosymbiol (BSB) does not function as a hormone but is an essential rhizospheric signal needed for AM symbiosis establishment (Kodama et al. 2022).

In angiosperms, SL perception is achieved by the receptor DWARF14 (D14). SL perception triggers D14's binding to the F-box protein MORE AXILLARY GROWTH 2 (MAX2), and subsequent targeting of repressor proteins (Bennett et al. 2016). These repressors are encoded by a small gene family, and called SUPPRESSOR OF MAX2 (SMAX)1-LIKE (SMXL) in Arabidopsis (Stanga et al. 2013; Soundappan et al. 2015; Wang et al. 2015), and DWARF53 (D53) in rice (Jiang et al. 2013). SMXL6, 7, and 8 are key repressors of the SL pathway for branching control (Soundappan et al. 2015; Wang et al. 2015). SMAX1 and SMXL2 are repressors in a parallel signaling pathway controlling seed germination and seedling development (Stanga et al. 2016). This latter pathway also involves the MAX2

F-box protein, and the KARRIKIN INSENSITIVE 2 (KAI2) receptor (Conn and Nelson 2016), an ancestral paralog of D14. While the existence of an endogenous hormonal KAI2-ligand (KL) is expected (Waters et al. 2015; Conn and Nelson 2016; Sun et al. 2016), the identity of molecule(s) activating the so-called KL pathway is still unknown.

As described in angiosperms, response to KL or SL requires the degradation of respective SMXL repressors (SMAX1 or SMXL/D53) by the 26S proteasome, via the action of an SCF complex containing MAX2. SMAX1 and SMXL/D53 are large proteins with several conserved structural domains: a double Clp-N domain in N terminus and two ATPase domains (D1 and D2), separated by a middle region (M). The D2 domain is split into D2a and D2b subdomains by an ethylene-responsive element binding factor amphiphilic repression (EAR) motif (for review see (Temmerman et al. 2022)). In the D2a subdomain lies a specific degron motif (RGKT or P loop), necessary for KL/SL-triggered SMXL degradation (Zhou et al. 2013; Khosla et al. 2020). Their activity as repressors partly relies on the EAR motif mediating transcriptional repression (Liang et al. 2016; Ma et al. 2017). Angiosperms possess a third subclade of SMXL proteins, SMXL345, involved in cell differentiation, in a MAX2-independent manner (Wallner et al. 2017; Moturu et al. 2018; Wallner et al. 2020).

Members of gene families associated with SL or KL signaling are found in all plants (Walker et al. 2019), with the exception of the *D14* SL receptor gene, which likely emerged in the flowering plant lineage. Outside seed plants, putative receptors are homologs to KAI2 (Bythell-Douglas et al. 2017). Thanks to reverse genetics, SL and KL signaling pathways have been partially described in bryophyte models (Proust et al. 2011; Decker et al. 2017; Mizuno et al. 2021; Kodama et al. 2022; Kyozuka et al. 2022; Bonhomme and Guillory 2022). In *Marchantia polymorpha*, a KAI2 and MAX2-dependent pathway was recently identified, that regulates thallus growth, and where the unique MpSMXL homolog acts as a repressor (Mizuno et al. 2021). In the moss *P. patens*, we have previously shown that the homolog of MAX2 is not necessary for response to SLs but is involved in the response to red light and possibly in KL signaling (Lopez-Obando et al. 2018; Lopez-Obando et al. 2021). The *KAI2* gene family is extended in *P. patens*, comprising candidate receptors for SLs (GJM clade), and candidate receptors for the KL (euKAI2 clade). Thus, both SL (MAX2-independent) and KL (MAX2-dependent) pathways are present in *P. patens*, in contrast to *M. polymorpha* (Lopez-Obando et al. 2021). Also contrasting with the single SMXL copy of *M. polymorpha,* there are four *PpSMXL* genes. Hence, we have sought to determine whether SMXL proteins' role in the ancestral KL signaling pathway is conserved in *P. patens,* and whether the seemingly moss-specific SL signaling pathway also relies on SMXL proteins.

Here, we show that the four PpSMXL proteins all display the ancestral SMXL domain organization, as well as motifs matching the respective EAR and RGKT consensus (Walker et al. 2019). All four homologs appear to play a negative role downstream of PpMAX2 in the putative KL signaling pathway of *P. patens*. Introduction of moss *PpSMXL* genes in Arabidopsis failed to induce functional

complementation of the *Atsmax1* mutant, suggesting PpSMXL are not able to form a functional signaling complex with AtKAI2 and AtMAX2. Our results nonetheless support the hypothesis of the ancestral role of SMXL proteins being KL response. Moreover, two PpSMXL proteins could act as positive actors of SL signaling and thus constitute a level of crosstalk between the SL and the PpMAX2-dependent KL pathway. The finding that none of these SMXL homologs are repressors in the SL signaling pathway supports the hypothesis that the mechanism triggering SL signal transduction diverged between *P. patens* and seed plants, even though the same family of receptors was recruited in both lineages (Bythell-Douglas et al. 2017; Lopez-Obando et al. 2021). Furthermore, our work supports the hypothesis that the use of SLs as hormones is not an ancestral land plant trait and might not be shared by seedless plants outside of bryopsid mosses (Kodama et al. 2022).

Results

Phylogeny and expression analysis reveal two clades of *SMXL* genes in *P. patens*

We recovered four SMXL genes from *Physcomitrium patens* (formerly *Physcomitrella patens*) genome on Phytozome (V3.6) as the first four results of a BLAST against *P. patens* proteome, using the full protein sequence of AtSMAX1 (encoded by *AT5G57710.1*). The predicted PpSMXL proteins contain the four regions characteristic of the SMXL family: a double ClpN N-terminal domain (N) and two ATPase domains (D1 and D2) separated by a long middle domain (M) ((Temmerman et al. 2022) and Supplemental Figure S1). As per a previous extensive analysis of SMXL proteins phylogeny (Walker et al. 2019), proteins encoded by *Pp3c2_14220* and *Pp3c1_23530* (here respectively named as *PpSMXLA* and *PpSMXLB*) correspond to a SMXL clade that is specific to bryopsid mosses, while those encoded by *Pp3c9_16100* and *Pp3c15_16120* (*PpSMXLC* and *PpSMXLD* herein) correspond to a SMXL clade that is common to all mosses, including the sphagnopsida (Supplemental Figure S2). At the amino acid level, PpSMXLC and PpSMXLD show 72% identity and PpSMXLA and PpSMXLB 61%, whereas comparisons between proteins of the two different clades give 27-29% identity. This suggests these two PpSMXL clades might have been evolving separately since the emergence of the bryopsid lineage.

To further explore the putative functional divergence between these two clades, we checked for differences in the expression of the four *PpSMXL* genes. For each *PpSMXL* gene, the region preceding the start codon was cloned upstream of the GUS coding sequence and this construct was introduced into wild-type (WT) moss plants. By staining two-week-old plants from *proPpSMXL:GUS* lines, we found that *PpSMXLA* promoter is hardly active at this age, as staining was only observed at the tip of phyllids (Figure 1A). *PpSMXLC* and *PpSMXLD* promoters seem to be much more active since GUS staining was noted in most of the tissues (Figure 1A). Unfortunately no *proPpSMXLB:GUS* line was obtained. We

then investigated these genes' expression along vegetative development, using RT qPCR. With this method, *PpSMXLB* transcripts were detected and we could confirm that gene expression is generally lower in the *PpSMXLA/B* clade than in the *PpSMXLC/D* clade (Figure 1B). Indeed, compared to the mean expression level of *P. patens* genes which is around 16 reads per Kilobase of transcript per million mapped reads (RPKM), *PpSMXLC/D* levels are approximately 18 RPKM and *PpSMXLA/B* levels are around 2 and 6, respectively (Perroud et al. 2018). We could not distinguish any significant changes in transcript levels across *protonema* development, but *PpSMXL* transcript levels all tended to be higher in mature gametophores and/or rhizoids at later developmental stages (Figure 1B).

Ppsmxl loss-of-function mutants do not display a constitutive SL response phenotype

To investigate the function of the four *PpSMXL* genes in *P. patens* and their potential involvement in the SL/PpKAI2L-GJM or KL/PpMAX2 pathway, we employed two CRISPR-Cas9-based mutagenesis strategies. In the first one, we used only one guide RNA targeted against the CDS region to obtain *Ppsmxl* mutants with frameshift or small insertions/deletions (Lopez-Obando et al. 2016b). We ensured obtaining knock-out mutants by devising a second strategy using two guide RNAs, with one targeting each untranslated region (UTR) (Supplemental Figure S1; Supplemental Table S1). Mutations giving rise to a complete deletion of the CDS are noted as *Ppsmxl*Δ (Supplemental Figure S3 and S4). Reasoning that the four PpSMXL proteins grouping into two clades could reflect different functions (Supplemental Figure S2 and (Walker et al. 2019)), we first sought double mutants for each clade, e.g. *Ppsmxl(Δ)ab* and *Ppsmxl(Δ)cd* mutants. Single and double mutants in the *PpSMXLA/B* clade were highly similar to the WT when grown under white light (Figure 2, A and B and Supplemental Figure S5). On the other hand, both *Ppsmxlcd* and *Ppsmxl*Δ*cd* double mutants showed markedly enhanced protonema extension, resulting in very large plants (Figure 2, A and B), even surpassing the *Ppccd8* mutant in some experiments (Supplemental Figure S5). Single *Ppsmxl*Δ*c* and *Ppsmxld* mutants also tended to be larger than WT, although not to the same level as double mutants (Figure 2, A and B). Thus, it appears *PpSMXLC/D* loss of function results in a protonema extension phenotype opposite to the *Ppmax2-1* mutant, which would be compatible with PpSMXLC/D proteins acting as repressors downstream of PpMAX2.

We assayed the phenotypes of the *Ppsmxl* mutants in the dark and continuous red-light, in conditions that allowed us to clearly distinguish mutants affected in the SL pathway from those affected in the PpMAX2-dependent KL pathway (Lopez-Obando et al. 2018; Lopez-Obando et al. 2021). Compared to WT, SL deficient (*Ppccd8*) and SL insensitive (*Ppkai2Lgjm*) mutants elongate more caulonema filaments in the dark. In contrast, KL mutants (*Ppmax2* and *eu-KAI2* clade) elongate fewer filaments than WT in the dark, and develop taller gametophores than WT as a result of an altered photomorphogenesis response to red light (Lopez-Obando et al. 2021).

When grown vertically in the dark, *PpSMXLA/B* clade mutants tended to grow as many caulonema filaments as WT, of similar length (Supplemental Figure S6). On the contrary, *PpSMXLC/D* clade mutants developed more and longer caulonema filaments than WT, similarly to *Ppccd8* (Supplemental Figure S6). Additionally, when grown under continuous red light, *PpSMXLA/B* clade mutants showed a similar phenotype to WT, while *PpSMXLC/D* mutants had smaller and more stunted gametophores, again opposite to the *Ppmax2-1* mutant (Figure 2, C to F). Counting phyllid number allowed estimating phytomere length, which was shorter for PpSMXLC/D clade mutants than for any other genotype (including *Ppccd8*) (Supplemental Figure S7), suggesting constitutive photomorphogenesis. Overall, *Ppsmxl* mutants' phenotypic characterization suggests that (1) *PpSMXLC* and *PpSMXLD* act redundantly to limit protonema growth, at least partly by limiting caulonema filaments number and length; (2) *PpSMXLC* and *PpSMXLD* activate gametophore photomorphogenesis under red light; (3) *PpSMXLA* and *PpSMXLB* play a very modest role (if any) in regulation of protonema and gametophore growth in various light conditions. Taken together, the results indicate that *Ppsmxlcd* mutants display largely opposite phenotypes to the *Ppmax2-1* mutant, suggesting that PpSMXLC/D would act redundantly as repressors of the PpMAX2-dependent KL signaling pathway.

Our first mutagenesis strategy also granted us few higher order *Ppsmxl* mutants, of which we characterized one triple mutant (*Ppsmxla1b1d1*) and one quadruple mutant (*Ppsmxla2b2c1d2*). It should be noted that the *Ppsmxlc1* mutation in the quadruple mutant might not cause a loss of function of *PpSMXLC,* as it removed four residues of the protein (not in a predicted functional motif) and did not induce a frameshift (Supplemental Figure S4). Both of these mutants displayed highly restrained growth and browning of the protonema, and only rare stunted gametophores were produced (Supplemental Figure S5A). Moreover, these higher order mutants grew few agravitropic curled caulonema filaments in the dark (Supplemental Figure S6).

PpSMXL overexpressing lines display phenotypes alike *Ppmax2-1*

To further explore the putative role of PpSMXL as repressors of the PpMAX2-dependent pathway, we examined the phenotype of stable *P. patens* transgenic lines overexpressing *PpSMXLA* and *PpSMXLC* with a N-terminal GFP tag under the maize ubiquitin promoter (*pZmUbi:GFP-PpSMXL* lines). We checked the *GFP-PpSMXL* fusion transcript overexpression by semi-quantitative RT-PCR (Supplemental Figure S8A). For clarity, we simplified the notation of these lines as OE-SMXLA and OE-SMXLC, respectively. When grown under standard conditions in white light, these lines were significantly less radially extended than the WT (Figure 3, A and B). OE-SMXLA and OE-SMXLC displayed a growth phenotype similar to that of *Ppmax2-1,* with less (but bigger) gametophores than in WT (Figure 3A and Supplemental Figure S8B). Furthermore, when we examined the phenotype of these lines under red-light continuous illumination, we likewise found that these OE lines behaved very

similarly to *Ppmax2-1* (Figure 3, C-D) with very elongated gametophores. Thus, in red light conditions,
OE-PpSMXL lines show an opposite phenotype to that of *Ppsmxl* mutants, at least of the *PpSMXLC/D*
clade (comparing Figure 2, C-F to Figure 3, C-D). These features point to *PpSMXLC* but also *PpSMXLA*
acting downstream of *PpMAX2* and playing an opposite role to *PpMAX2* in development.

PpSMXL loss-of-function does not restore WT growth in the Ppccd8 background

To test a putative role of PpSMXL genes in the SL pathway, we combined *Ppccd8* and *Ppsmxl*
mutations. Replicating our first CRISPR mutagenesis strategy in the original *Ppccd8* SL deficient
background, we obtained *Ppccd8 Ppsmxlab*, *Ppccd8 Ppsmxlcd*, and *Ppccd8 Ppsmxlabcd* mutants.
Following the growth of these *Ppccd8 Ppsmxl* mutants (Figure 4; Supplemental Figure S9A), we noticed
that mutation of neither SMXL clade restored protonema extension to WT levels. Strikingly, the
mutation of all four *PpSMXL* genes, which had a dramatic negative effect on growth in the WT
background (Supplemental Figure S5 and S6), was completely circumvented by the *Ppccd8* mutation,
and the *Ppccd8 Ppsmxlabcd* quintuple mutant showed a phenotype similar to *Ppccd8*. (Figure 4;
Supplemental Figure S9B). It is however worthy of note that the *Ppsmxlc9* mutation in the *Ppccd8*
Ppsmxlabcd mutant does not lead to a frameshift, similarly to the *Ppsmxlc1* mutation (Supplemental
Figure S4), therefore the PpSMXLC protein is likely still functional in this mutant. These observations
indicate that PpSMXL proteins are not acting as repressors in the SL pathway. In addition, the lack of
functional PpSMXLA, B and D proteins appears to be detrimental only when endogenous SL are
present.

PpSMXL mutations partially restore Ppmax2-1 mutant phenotypes

To confirm the genetic relationship between *PpSMXL* and *PpMAX2*, we tested whether *Ppsmxl*
loss of function could restore a WT phenotype in a *Ppmax2* background. *Ppsmxlab* mutants in the
Ppmax2-1 background were obtained, but no *Ppsmxlcd* mutants. Therefore, we instead mutated
PpMAX2 in one of the *PpsmxlΔcd* mutants, using the CRISPR-Cas9 system, with five guide RNAs
targeting *PpMAX2* (Supplemental Figure S10 Supplemental Table S1). Using this approach, we
successfully obtained a new *Ppmax2* mutant allele (*Ppmax2-16*) in the *PpsmxlΔcd* mutant background.
When we examined the protonema extension of the *Ppmax2 Ppsmxl* mutants, we found that *Ppsmxlab*
mutations had no effect on *Ppmax2* mutant phenotype while *Ppsmxlcd* mutations could partially
suppress the dramatic growth decrease caused by *PpMAX2* loss of function (Figure 5, A-B). By contrast,
under red light, both *Ppsmxlab* and *Ppsmxlcd* mutations could partially restore the excessive elongation
of gametophores caused by *PpMAX2* loss of function (Figure 5, C-D).

Altogether, phenotype analysis led us to hypothesize that all four PpSMXL proteins could act

as repressors of the PpMAX2-dependent KL pathway, with both *PpSMXL* clades acting redundantly in

the regulation of gametophore elongation in red light, while *PpSMXLC/D* would play a predominant

role under standard light conditions.

Loss of PpSMXL C/D function dampens the response to SL mimic but not to KL mimic

We then explored whether *Ppsmxl(Δ)* mutants have a modified phenotypic response to GR24

enantiomers. To investigate this response, we relied on the test of filament growth in the dark, which

consists in counting the caulonema filaments after enantiomer application (Hoffmann et al. 2014; Lopez-

Obando et al. 2021). When treated with 1 µM (+)-GR24, WT and *Ppccd8* plants developed significantly

less caulonema filaments, which is the typical phenotypic response to SL treatment ((Lopez-Obando et

al. 2021) and Figure 6A). When investigating the response of *PpsmxlA/B* clade mutants to (+)-GR24, it

appeared that all *Ppsmxl* single mutants and *Ppsmxlab* double mutants could respond to this SL mimic,

apparently as much as the WT (Figure 6A; Supplemental Figure S11A and S11B). The response of

Ppsmxlcd double mutants to (+)-GR24 was less pronounced than that of *Ppsmxlab*, considering several

independent assays (Figure 6A; Supplemental Figure S11C). *PpSMXLC* and *PpSMXLD* play redundant

roles in responding to (+)-GR24, as response was not disturbed in single *Ppsmxlc* and *Ppsmxld* mutants

(Supplemental Figure S11B). This suggested that PpSMXLC/D could mediate at least part of the SL

response. In addition, single *Ppsmxlc* mutants developed slightly but significantly more filaments than

WT in control conditions (Supplemental Figure S11B), possibly indicating that *PpSMXLC* plays a

predominant role over *PpSMXLD* in the regulation of filament number.

In our previous assays, the use of (−)-GR24 or KAR$_2$ treatment to trigger the KL pathway gave

non-reproducible results, though sometimes tending to increase the filament number (Lopez-Obando et

al. 2021). Therefore, we instead chose to evaluate the effect of (−)-desmethyl-GR24, described as a

better mimic for KL in both Arabidopsis and *Marchantia* (Yao et al. 2021). In similar assays of filament

growth in the dark, increasing doses of (−)-desmethyl-GR24 (0.01 to 10 µM) significantly increased the

number of filaments in WT and *Ppccd8* mutant, but not in *Ppmax2-1* mutant (Supplemental Figure S12A

and S12B). Unexpectedly, the eu-KAI2 clade mutant, affected in *PpKAI2L-A* to *E* genes encoding

putative KL receptors, was responsive to the (−)-desmethyl-GR24 treatment, however not in all assays

(Supplemental Figure S12 A and S12B). We still considered (−)-desmethyl-GR24 our best mimic for

the KL in *P. patens*. We tested the effect of (−)-desmethyl-GR24 on *Ppsmxl* mutants and observed that

both clade A/B and clade C/D *Ppsmxl* double mutants were able to respond to this compound (here

tested at 1 and 10 µM, Figure 6B). However, considering several independent assays with 1 µM (−)-

desmethyl-GR24, the increase of the filament number was lower for the *Ppsmxlcd* double mutant than

for the *Ppsmxlab* double mutant (Figure 6B, Supplemental Figure S12C). This observation leads to the

hypothesis that all four *PpSMXL* genes may play a redundant role in response to (−)-desmethyl-GR24,

at least in this experimental setup (in the dark).

278   *PpSMXL* transcript levels are downregulated in response to activation of the SL signaling pathway

We wondered if a differential transcriptional regulation of all four *PpSMXL* genes could explain

the different responses of moss mutants to SL and KL mimics. We therefore compared the transcript

levels of *PpSMXL* genes in WT to that in the *Ppccd8* mutant and in the *Ppmax2-1* mutant, and assayed

the effect of (+)-GR24 application. Because we previously found that the transcriptional response of

SL-related genes was strongly affected by light and enhanced in the dark (Lopez-Obando et al. 2016a;

Lopez-Obando et al. 2018), we assessed this response in both dark and light conditions.

We observed that all four *PpSMXL* genes have much higher transcript levels in the dark than in

the light, in WT, in the *Ppccd8* SL synthesis mutant, and in the *Ppmax2-1* mutant (Supplemental Figure

S13, comparing the y-axis in left *versus* right panels). In control conditions, all four *PpSMXL* genes

were downregulated in the *Ppmax2-1* mutant compared to WT in the dark (Supplemental Figure S13 A-

D, left panels), and also in the light for *PpSMXLC* and *PpSMXLD* (Supplemental Figure S13 A-D, right

panels). This suggests that the PpMAX2-dependent KL pathway upregulates the expression of *PpSMXL*

genes. In light conditions, *PpSMXLC* and *D* transcript levels in control conditions were higher in *Ppccd8*

compared to WT, indicating that endogenous SLs repress the expression of these two genes.

Accordingly, the (+)-GR24 treatment down-regulated *PpSMXLC* and *D* transcript levels, in both WT

and *Ppccd8* mutant (Supplemental Figure S13, C and D, right panels). Surprisingly, in the *Ppmax2-1*

mutant, transcript levels of *PpSMXLC* and *D* were upregulated by the (+)-GR24 treatment in light

conditions (Supplemental Figure S13, C and D). Considered together, these results show that *PpSMXL*

genes' transcript levels are mainly down-regulated by light and that PpMAX2 function leads to enhanced

PpSMXL transcription. Hence, expression of *PpSMXL* genes could be induced by activation of the KL

pathway. In addition, the down-regulation of *PpSMXLC* and *PpSMXLD* transcripts by SLs further

suggests possible crosstalk with the SL pathway.

302   PpSMXL proteins are localized mainly in the nucleus, and not degraded following SL application

303   Functional nuclear localization signals (NLS) were found in a part of the N-terminal region of

304   AtSMXL7 (Liang et al., 2016) AtSMAX1 (Khosla et al. 2020) and OsSMAX1 proteins (Choi et al.

305   2020). This sequence enriched in residues with basic sidechains is broadly conserved in the four

306   PpSMXL proteins, suggesting they are also located in the nucleus (Supplemental Figure S14).

To determine the subcellular localization of PpSMXL proteins, we transiently overexpressed RFP-PpSMXL fusion proteins under the 35S promoter in *Nicotiana benthamiana* epidermal cells stably expressing CFP tagged H2b histone (thus all nuclei were marked by CFP, Supplemental Figure S15). A clear nuclear signal was observed for all four PpSMXL (Supplemental Figure S15A, S15C, S15E, S15G), but some cytoplasmic signal was also observed for RFP fusions with PpSMXLA, PpSMXLB and PpSMXLC. Thus, either the PpSMXL are actually nucleo-cytosolic, or high expression of RFP-PpSMXL under p35S led to proteins in excess leaking out of the nuclear compartment.

In angiosperms, the RGKT degron motif is responsible for SMXL proteasomal degradation in presence of SL (Zhou et al. 2013). Since the RGKT motif was found in PpSMXLC and PpSMXLD proteins (RGRT in PpSMXLA and PpSMXLB, Supplemental Figure S1; Supplemental Figure S15I), we obtained constructs where this motif was deleted. In *N. benthamiana* leaves transiently expressing *p35S:RFP-PpSMXL* constructs, we found similarly high levels of RFP-PpSMXL fusion proteins whether the degron motif was present or deleted (ΔRGRT and ΔRGKT lines, Supplemental Figure S15B, S15D, S15F, S15H). Nonetheless, the additional cytosolic RFP signal seemed increased in leaves expressing the ΔRGKT versions of PpSMXL proteins, especially for PpSMXLB and PpSMXLD.

Using our stable *P. patens* transgenic lines overexpressing *PpSMXL* (*pZmUbi:GFP-PpSMXL* lines), we could confirm the nuclear localization of PpSMXLA, PpSMXLC and PpSMXLD, even though the GFP signal was not restricted to this compartment for PpSMXLC and PpSMXLD (Figure 7). The GFP-PpSMXLB fusion led to a faint and more diffuse cytoplasmic signal. We then tested whether the SL treatment had an effect on GFP-PpSMXL protein stability and/or localization in *P. patens*. We could not see any effect of a 20 minute-long 1 µM (+)-GR24 treatment on the signal observed in the *pZmUbi:GFP-PpSMXLA* line (Supplemental Figure S16A). When introduced into the *Ppmax2-1* mutant background, the GFP-PpSMXLA fusion also led to fluorescent nuclei, that were easier to spot (Supplemental Figure S16B-C). In this background as well, the 1 µM (+)-GR24 treatment had no effect on the fluorescent signal (Supplemental Figure S16C). These results support the hypothesis that at least PpSMXLA is not a negative regulator of the SL pathway.

PpSMXL proteins interact with players of the PpMAX2-dependent pathway

Since we hypothesized the PpSMXL proteins to act in the PpMAX2-dependent KL pathway, we tested putative interactions of these proteins with PpMAX2. Bimolecular fluorescence complementation (BiFC) assays in *P. patens* protoplasts were unsuccessful. However, when overexpressed in *N. benthamiana* cells, each of the four PpSMXL proteins interacted with PpMAX2 when respectively fused to complementary parts of the eYFP fluorescent protein (Figure 8, A-D; Supplemental Figure S17). The eYFP signal was lower for PpSMXLA and PpSMXLC, than for PpSMXLB and PpSMXLD, but still indicated a nuclear interaction with PpMAX2 (Figure 8, A, C).

Interestingly, we also found that PpSMXLB and PpSMXLD could potentially form homo-oligomers (Figure 8, E, F).

Using the same technique, we investigated the interactions of PpSMXL proteins with PpKAI2L proteins, among which the previously reported putative receptors for SL and KL pathways (Lopez-Obando et al. 2021). Among the PpKAI2L proteins we tested in BiFC (at least one from each clade), only PpKAI2L-C from eu-KAI2 clade could interact with PpSMXL proteins, moreover the four of them (Figure 8, G-J). This is consistent with our previous demonstration that PpSMXL proteins act in the PpMAX2-dependent pathway since eu-KAI2 genes are also involved in this pathway (Lopez-Obando et al. 2021). To conclude, BiFC assays revealed interactions involving PpSMXL, PpMAX2 and eu-KAI2 putative KL receptors. It is also interesting to note that, while PpSMXL proteins could also be present in the nucleosol in this experimental system, the BiFC signal was strictly restricted to the nuclear compartment.

Altogether, our data in *P. patens* allow us to propose a model for PpSMXL role in moss development, highlighting a possible crosstalk between SL and KL-MAX2 pathway (Figure 9 and see discussion).

Expression of PpSMXL proteins rescues SMAX4/5 function, but not SMAX1 or SMXL6/7/8 function in *Arabidopsis*

In order to investigate the putative conservation of SMXL proteins function along land plant evolution, we conducted functional complementation assays in *Arabidopsis thaliana.* To this end, one PpSMXL protein from each clade (PpSMXLB and PpSMXLC) was expressed in Arabidopsis *smxl* mutants, under the control of appropriate native promoters from AtSMAX1, AtSMXL5, and AtSMXL6 for *Atsmax1, Atsmxl4,5* double and *Atsmxl678* triple mutant complementation, respectively. Furthermore, as positive controls for complementation, the Arabidopsis *SMAX1, SMXL5 and SMXL6* genes, driven by their promoter, were introduced into *Atsmxl* mutants.

First, complementation of AtSMXL4/5 function was tested by complementation assays under AtSMXL5 promoter, by measuring the primary root length of young seedlings (Figure 10). We found that all three PpSMXLB and three out of four PpSMXLC expressing lines (Supplemental Figure S18A) had similar root length as WT Arabidopsis plants and as the mutant plants complemented with native *AtSMXL5*. This surprising result suggests that both SMXL clades of *P. patens* have retained the same KL and SL-signaling independent function as Arabidopsis SMXL homologs.

Then, we explored the ability of PpSMXL proteins to functionally replace AtSMAX1, that is to say their capacity to integrate into the KL signaling pathway of Arabidopsis. Complementation of AtSMAX1 function was assayed by measuring hypocotyl length of *Atsmax1-2* mutant seedlings expressing PpSMXLB or PpSMXLC (Supplemental Figure S18B) and grown in low light for 11 days (Supplemental Figure S19A). Here we could only achieve partial complementation of *Atsmax1-2*

decreased hypocotyl length by reintroducing *pSMAX1:SMAX1*, furthermore only in two out of our three
independent transgenic lines (*pSMAX1:SMAX1* #20.6 and #38.1). However, the case of plants
expressing *PpSMXLB/C* was still different, as none of the independent lines was even partly restored.
Some of them even appeared to develop shorter hypocotyls (*e. g. pSMAX1:PpSMXLB # 5.4*). Thus,
although the used *pSMAX1* promoter sequence may not be enough efficient, it seems reasonable to
conclude that PpSMXL proteins are not able to replace SMAX1 in Arabidopsis KL signaling.

Finally, we explored whether PpSMXL proteins could functionally replace AtSMXL6, by
comparing the shoot phenotype of mature transgenic *Atsmxl678* plants expressing PpSMXLB or
PpSMXLC (Supplemental Figure S18C) to that of the untransformed *Atsmxl678* mutant (Supplemental
Figure S19, B-D). Several shoot phenotypes were assessed, such as plant height, caulinary secondary
branch (C2) number, and leaf shape (width). One transformed line out of three independent lines
expressing PpSMXLB (#13.2) showed partial complementation of both plant height and C2 number
phenotype (Supplemental Figure S19, B-C). However, this line showed no complementation of the leaf
width. Leaf width was partially complemented in another transformed line expressing PpSMXLB (#7.7).
None of the three independent lines expressing PpSMXLC showed any complementation of shoot
phenotype. Hence, it appears PpSMXL proteins are not able to replace SMXL6 in Arabidopsis SL
signaling either.

Discussion

395   SL involvement in mycorrhization and plant architecture implies that these molecules played
396 major roles in land plant evolution, for adaptation to land conditions and then for the emergence of seed
397 plants. The knowledge of SL roles and associated cellular pathways in non-vascular plants is necessary
398 to understand how these compounds could have contributed to the process of terrestrialization.

Phylogeny and expression profiles suggest a differential regulation of clade A/B and clade C/D
PpSMXL

It has already been shown that bryopsid mosses have an additional SMXL clade compared to
other non-seed plants (Walker et al. 2019). We report evidence of this in *P. patens*, in which
PpSMXLA/B belong to this divergent clade, which could potentially be neofunctional, as we have shown
earlier for some PpKAI2-L homologs (Lopez-Obando et al. 2021). The split between *PpSMXLA* and
PpSMXLB, and between *PpSMXLC* and *PpSMXLD* on the other hand, is probably quite recent given the
high similarity of encoded proteins in each given clade (Supplemental Figure S2). PpSMXLA/B
moreover have shorter D1 and M domains compared to PpSMXLC/D (Supplemental Figure S1) and
lack the typical degron motif (Supplemental Figure S1 and S15). The consensus RGKT sequence is also
modified in some flowering plants SMXL where the motif is nonetheless functional (OsSMAX1 (Zheng

et al. 2020)). However, among the sequences examined, PpSMXLA/B are the only one lacking the K
residue, which is functionally relevant in Walker A motifs as it is necessary for phosphate binding
(Bianchi et al. 2012). Inversely, the EAR motif is conserved in all four PpSMXL proteins, thus
transcriptional regulation could be a uniting feature between the two clades.

Expression level of *PpSMXLA/B* is lower than that of *PpSMXLC/D* in the protonema (Figure 1).
This could indicate that only the C/D clade has a relevant function in the protonema, which is supported
by phenotypic analysis of *Ppsmxl* mutants. All four *PpSMXL* genes are more highly expressed in older
gametophores and/or rhizoids than in filaments, implying that PpSMXL function becomes more
important as *P. patens* switches from early vegetative development to mature vegetative development.
Furthermore, available RNAseq data (Ortiz-Ramirez et al. 2016), indicate that *PpSMXL* transcripts are
even more abundant in spores.

***PpSMXL* genes are mainly regulated by light and the PpMAX2-dependent pathway**

PpSMXL gene expression is mainly downregulated by light, and only modestly sensitive to (+)-
GR24 treatments (Supplemental Figure S13). In the light, all four genes tend to be downregulated by
(+)-GR24 in WT plants. This trend becomes more evident in the SL deficient *Ppccd8* background, but
solely for *PpSMXLC* and *PpSMXLD,* which accordingly have elevated transcript levels in control-
treated *Ppccd8* compared to WT. Thus, it appears that only *PpSMXLC* and *PpSMXLD* are significantly
repressed by (+)-GR24 and endogenous SLs alike, in the light. The moderate effects of (+)-GR24
enantiomer on *PpSMXLC* and *PpSMXLD* expression are of similar amplitude as those reported for
angiosperms *SMXL* genes (Stanga et al. 2013; Kerr et al. 2021). However, the feedback regulation
exerted by SL on *PpSMXL* transcript levels is opposite to that in angiosperms, since we observed a
downregulation of these genes. We relied on the *Ppmax2* mutant as a proxy to explore the effect of the
KL pathway on *PpSMXL* gene expression (Supplemental Figure S13). We found that all four *PpSMXL*
genes are downregulated in the *Ppmax2* mutant compared to WT in the dark, suggesting that KL
signaling would have a positive effect on the transcription of these genes. In the light, only *PpSMXLC*
and *PpSMXLD* showed lower transcript levels in the *Ppmax2* mutant compared to WT. This observation
is comparable to what was reported for *M. polymorpha*, the unique *MpSMXL* gene being downregulated
in *Mpmax2* background (Mizuno et al. 2021). We could hypothesize that PpSMXL proteins bind to the
promoters of the *PpSMXL* genes to repress their transcription, as previously described for angiosperm
SMXL proteins (Wang et al. 2020). In *Ppmax2*, PpSMXL protein levels are likely very high (from our
results, since OE-PpSMXL lines mimic *Ppmax2* phenotype), which would lead to low transcript levels
in this mutant. Finally, it was puzzling to observe that (+)-GR24 treatment increased *PpSMXLC* and
PpSMXLD expression in the *Ppmax2-1* mutant, when the opposite was seen in *Ppccd8*. Thus, SL could

somehow prevent the KL-triggered induction of *PpSMXL* transcript levels, perhaps by stabilizing
PpSMXL proteins (Figure 9 and see below).

PpSMXL proteins take part in nuclear signaling complexes

The presence of predicted NLS sequences, as well as an EAR motif in the four PpSMXL
proteins hint at a nuclear localization (Supplemental Figure S1, Supplemental Figure S14). We have
shown that all four PpSMXL are indeed located in the nucleus, albeit not exclusively (Figure 7,
Supplemental Figures S15). For the moment, we cannot exclude that, as many proteins carrying an NLS,
PpSMXL proteins could actually shuttle between nucleus and cytosol, perhaps until a specific signal
(KL?) induces their retention into the nuclear compartment. Still, this partial nuclear localization is
enough to enable the interaction of PpSMXLs with proteins in the nucleus (namely PpMAX2 and
PpKAI2L-C, Figure 8). It would be interesting in the future to explore whether the EAR motif grants
PpSMXL proteins a function in transcriptional regulation. Notably, we could determine if they can
interact directly with DNA, like has been demonstrated for AtSMXL6 (Wang et al. 2020), and which
loci are targeted. In particular, it would be interesting to see whether PpSMXL are able to directly
downregulate their own expression, as is the case for AtSMXL6 that binds to the ATAACAA motif in
the promoter region of *AtSMXL6* (Wang et al. 2020).

PpSMXL proteins are not highly sensitive to SL levels and their sensitivity to KL probably
depends on light

We report herein that none of the four PpSMXL proteins are rapidly degraded in response to
(+)-GR24, not even PpSMXLA, which transcripts are increased by treatment with this enantiomer in
the dark (Supplemental Figure S13 and Supplemental Figure S16). Furthermore, when put in the
Ppmax2 background, GFP-PpSMXLA fusion protein was also found insensitive to the (+)-GR24
treatment (Supplemental Figure S16C). The absence of (+)-GR24 triggered degradation is probably not
the result of an insufficient amount of (+)-GR24 and/or treatment duration, as (1) AtSMXL6 is almost
completely degraded after 20 minutes of (±)-GR24 2μM in p35S:AtSMXL6-GFP seedling (Wang et al.
2015); (2) AtSMXL7 is significantly decreased after only 10 minutes of a 1 μM (±)-GR24 treatment in
p35S:AtSMXL7-YFP Arabidopsis roots (Soundappan et al. 2015); (3) as little as 12 minutes are
necessary for pOsAct:D53-GFP degradation in rice roots (Zhou et al. 2013). Thus, PpSMXL proteins
do not behave similarly to AtSMXL6/7 or D53 proteins and they seem insensitive to both the SL mimic
(+)-GR24 and endogenous SL from both *P. patens* (GFP lines, Figure 7 and Supplemental Figure S16)
and *N. benthamiana* (RFP transient expression lines, Supplemental Figure S15). While their stability in
N. benthamiana could be explained by an incompatibility with the angiosperm MAX2, PpSMXLs are

also stable in *P. patens.* This indicates that PpSMXL proteins are likely not degraded in the context of

SL signaling. Likewise, deletion of the degron motif (or degron-like for PpSMXLA/B) has no effect

since these proteins are stable (Supplemental Figure S15).

The stability of PpSMXLA and PpSMXLC in *P. patens* lines, in the WT background where

PpMAX2 is functional, suggests that the whole pool of PpSMXL proteins is not degraded in response

to the endogenous compound(s) perceived via PpMAX2 either (putative KL). However, it is important

to note that observation of GFP fluorescence was carried out on *P. patens* lines that were previously

incubated in low-light conditions or even in the dark for a few hours (Figure 7, Supplemental Figure

S16). Since we observed that the intensity of the GFP-PpSMXL signal tended to be lower or even

undetectable in high light conditions, it is tempting to hypothesize an activation of the PpMAX2-

dependent KL signaling pathway by light, which would lead to PpSMXL degradation. Hence, reported

enhanced expression of *PpKAI2-Like* (Lopez-Obando et al. 2016a)*, PpMAX2* (Lopez-Obando et al.

2018), and *PpSMXL* genes in the dark (Supplemental Figure S13), would be a mean to keep KL signaling

active in the absence of light.

There might be another link between PpSMXL proteins and SL response. Indeed, we have

observed that triple/quadruple *Ppsmxl* mutants display a very striking phenotype seemingly resulting

from protonema growth limitation coupled to early senescence. However, this phenotype was fully

alleviated by the *Ppccd8* mutation (Figure 4), suggesting that endogenous SLs produced via PpCCD8

activity are responsible. We previously showed that SLs have toxic effects at high concentrations on

moss growth (Lopez-Obando et al. 2018). Since the *Ppccd8* single mutant does not display this

phenotype, we can also infer that PpSMXL proteins would have a protective effect against SL toxicity.

500   *Ppsmxl* loss of function mutants contradict PpSMXL acting as repressors of SL response

At first glance, it is evident that clade C/D *Ppsmxl* mutants do not display a constitutive SL

response phenotype, instead they are alike the SL deficient *Ppccd8* mutant (Figure 2; Supplemental

Figure S7). This could be indicative of either a positive role of PpSMXLC/D in SL signaling or a

negative role of these proteins in PpMAX2-dependent signaling. The phenotype of clade A/B mutants

is more comparable to WT, which does not rule out the possibility of PpSMXLA/B acting as repressors

of SL signaling (Figure 2). However, both A/B clade and C/D clade loss-of-function in the *Ppccd8*

background result in mutants that are like *Ppccd8* (Figure 4; Supplemental Figure S9)*.* If PpSMXL

proteins were repressors of SL signaling, the loss of repression should circumvent the absence of

endogenous SL and we could expect restoration to a WT-like phenotype. Therefore, PpSMXL are most

likely not involved in the repression of the SL response. From our various assays, it appears instead that

PpSMXLC and PpSMXLD act redundantly to enable the response to (+)-GR24 in dark-grown

caulonema (Figure 6; Supplemental Figure S11), but are not the sole positive regulators of this response.

This hypothesis would be in line with the observed repression of *PpSMXLC/D* gene expression by SL

(Supplemental Figure S13). Indeed, the SL signal would be transduced through PpSMXLC/D proteins,

which would inhibit the expression of their genes, permitting a negative feedback regulation of SL

response (Figure 9).

Ppsmxl and *Ppsmxl*Δ mutations in the same genes have comparable effects on phenotype. Thus,

shortened proteins produced from *Ppsmxl* alleles are not functional, even when most of the D1 domain,

thought to mediate interaction with receptors (Khosla et al. 2020), is (putatively) still present for the

PpsmxlA, PpsmxlB and PpsmxlC mutant proteins (Supplemental Figures S4).

PpSMXL are negative actors of the PpMAX2-dependent pathway

Two major points of evidence show that PpSMXL proteins are collectively involved in the

PpMAX2-dependent signaling pathway and exert a negative role in this pathway: (1) *P. patens* lines

overexpressing *PpSMXLA* or *PpSMXLC* are phenotypically similar to the *Ppmax2-1* mutant (Figure 3);

(2) Loss of function of either *PpSMXL* clade partially restores developmental disturbances caused by

loss of *PpMAX2* function (Figure 5). Although we did not obtain triple or quadruple *Ppsmxl* mutants in

the *Ppmax2-1* background, we hypothesize that the four PpSMXL proteins act in an additive fashion to

repress PpMAX2-dependent signaling. (3) PpSMXL proteins can indeed interact with components of

the PpMAX2-dependent pathway: PpMAX2 itself, but also PpKAI2L-C (Figure 8).

We also found that both clades *Ppsmxl* double mutants are able to respond to the KL mimic (−)-

desmethyl-GR24, although this response seemed reduced in *Ppsmxlcd* mutants (Figure 6, Supplemental

Figure S12C). Thus, the two *PpSMXL* clades likely have a redundant function in the context of

PpMAX2-dependent signaling, with PpSMXLC/D playing a prevalent role in the growth conditions we

examined. This hypothesis needs to be confirmed by testing the ability of higher order mutants to

respond to (−)-desmethyl-GR24, but the dramatic phenotype of these mutants makes them challenging

to work with.

PpMAX2 is involved in the induction of *PpSMXL* expression, while PpSMXL proteins are

likely targets of PpMAX2-dependent degradation. Therefore, PpSMXL could constitute a level of

negative feedback regulation in the PpMAX2-dependent pathway. Since this induction by PpMAX2 is

especially prevalent in the dark, negative feedback might be more active in the absence of light.

Nevertheless, the PpMAX2-dependent regulation of *PpSMXL* expression needs to be further clarified.

PpSMXL could be a bridge linking SL signaling and PpMAX2-dependent signaling

It is interesting to note that "de-repression" of PpMAX2-dependent signaling (*Ppsmxlcd*
mutants) and "silencing" of SL signaling (*Ppccd8* mutant) have very similar effects on developmental
pattern of the protonema (more extended) and gametophores (smaller, especially in red light) (Figure
2). This, together with the opposite effects of (+)-GR24 (SL mimic) and (−)-desmethyl-GR24 (KL
mimic) on dark-grown caulonema in WT, indicates that these two pathways likely regulate the same
processes in opposite manners. In addition, we have previously shown that the *Ppmax2* mutation is
epistatic to *Ppccd8* (Lopez-Obando et al. 2018). As suggested in this article, the phenotype of *Ppmax2*
would phenocopy a constitutive response to SL, which, according to our present data, would result from
an overaccumulation of PpSMXL proteins. This is compatible with a model where SL and KL signaling
converge on PpSMXL proteins and regulate their stability/function in opposite manners (Figure 9).

One of the processes seemingly regulated by both SL and PpMAX2-dependent pathways, is
probably cell division, as demonstrated for the *Ppccd8* mutant (Hoffmann et al., 2014). Increased cell
division and filaments branching still must be confirmed in *Ppsmxl* mutants, however, it is supported by
the observation that double *Ppsmxl* mutants of both clades (tend to) develop more caulonema filaments
than WT in the dark (Figure 6; Supplemental Figure S6). Furthermore, the observation that phenotypic
response to (+)-GR24 is not seen in the absence of PpSMXLC/D in some experiments suggests that SL
act at least partially via the PpMAX2-PpSMXL pathway. However, as we have seen herein as well as
previously (Lopez-Obando et al. 2021), loss of *PpMAX2* function or loss of function in all euKAI2clade
does not abolish response to (+)-GR24. Hence, SL signaling does not solely rely on the PpMAX2-
dependent pathway, and we can hypothesize that perception of SL by PpKAI2-L proteins of the (GJM)
clade does not act via a regulation of the PpMAX2-dependent pathway but nonetheless requires
functional PpSMXLC/D. A tempting explanation would be that PpKAI2-L proteins of the GJM clade
could interact with and stabilize PpSMXLC/D proteins, thus acting downstream of PpMAX2 and
leading to opposite effects on phenotype (Figure 9). While we did not find any direct interaction between
PpKAI2-LG/J and PpSMXLC/D using BiFC, the hypothesis that an unknown interactor is needed to
permit this interaction is still open.

Does PpSMXLC have a specific role among PpSMXL proteins?

We noted that higher-order *Ppsmxl* mutants display highly disturbed growth of gametophores
and rapid browning of the protonema, suggesting that extreme activation of the PpMAX2-dependent
pathway is very detrimental to viability. The observation that the *Ppccd8* mutation can circumvent the
dramatic effect of *Ppsmxlabcd* loss of function (PpSMXLC is probably still functional in the presented
mutant) (Figure 4; Supplemental Figure S9), together with our previous hypothesis of SL repressing
PpMAX2-dependent signaling, implies that accumulation of PpSMXLC in the absence of other
PpSMXL is detrimental to development. Under this hypothesis, in the *Ppccd8* mutant, inhibition of the

PpMAX2-pathway by endogenous SL would be lifted and PpSMXLC would be degraded in a
PpMAX2-dependent manner.

SMXL proteins likely retained a common molecular function along evolution

We have shown through our complementation assays in Arabidopsis that PpSMXLB/C
expression in the *Atsmxl45* mutant was enough to restore primary root length to a WT level.
Interestingly, it has been reported before that, in Arabidopsis itself, SMXL proteins from different clades
can ensure equivalent function if expressed with the same spatio-temporal profile. Indeed,
pAtSMXL5:AtSMAX1 expression in *smxl45* could also complement root growth (Wallner et al. 2017).
Hence, all SMXL might have actually kept the same ancestral molecular "growth regulating" function
that has been described for SMXL4/5. In angiosperms, this function would have (1) become KL/SL
regulated for SMAX1 and SMXL678 clades through the evolutive gain of interaction with KAI2/KAI2L
receptors and with MAX2, (2) become differentially regulated in various types of tissues and
developmental contexts through major changes in regulatory sequences.

On the other hand, the observation that PpSMXL expression does not complement the *Atsmax1*
mutant is not surprising, given that our previous attempt to complement the *Atmax2* mutant with
PpMAX2 was also unsuccessful (Lopez-Obando et al. 2018). Hence, we could think that, despite MAX2
and SMAX1/PpSMXL acting in a conserved KL signaling pathway in both species, the interface for
MAX2/SMXL interaction is simply not compatible between Arabidopsis and *P. patens* proteins. The
same could be concluded from the absence of complementation in the *smxl678* mutant.

The MpKAI2a/MpMAX2 pathway, regulating thallus and gemmae growth in *M. polymorpha*
through the degradation of the unique MpSMXL repressor (Mizuno et al. 2021) is so far the "simplest"
form described of the above-described ancestral growth regulating function. It is interesting to note that
in *M. polymorpha*, like in *P. patens*, this pathway is regulated by light. *M. polymorpha* does not
synthesize SLs (no CCD8), and therefore the crosstalk between SL and KL pathways that we highlight
in *P. patens* cannot be observed. This crosstalk is also apparently absent in *M. paleacea* though it
synthesizes BSB, likely because of the absence of the required receptor for SL as hormones (Kodama
et al. 2022). Therefore, *P. patens* would represent a specific case among bryophytes where the SL and
KL pathway are interconnected. The identity of both SL and KL signals remains to be discovered.

Methods

Cultivation of *Physcomitrium patens* Gransden

Unless otherwise explicitly stated in legends, experiments were always carried out on PpNO$_3$ medium

(corresponds to the minimal medium described by Ashton et al., 1979), in the following control

conditions: 25°C during daytime and 23°C at night, 50% humidity, long days conditions with 16 hours

of day and 8 hours of night (quantum irradiance of ~80 μmol/m^2/s). Multiplication of tissues from young

protonema fragments prior to every experiment is done in the same conditions but using medium with

higher nitrogen content (PpNH$_4$ medium, PpNO$_3$ medium supplemented with 2.7 mM NH$_4$ tartrate). For

red light experiments, plants were grown on PpNO$_3$ medium in Magenta pots at 25°C, in continuous

red-light (~45 μmol μmol/m^2/s). Cellophanes of appropriate sizes were used for monitoring of

protonema extension kinetics, as well as for the cultures launched in 6-well plates for gene expression

studies (see a detailed protocol in our methods chapter titled "Methods for medium-scale study of the

biological effects of strigolactone-like molecules on the moss *Physcomitrella patens*"). Analysis of

caulonema growth in the dark was performed in 24-well plates, with ~2 weeks of growth in control

conditions before incubation (± treatment) in the dark and placed vertically for ~10 days (see the same

method chapter).

P. patens gene expression analyses by qPCR

Total *P. patens* RNA was extracted and rid of contaminant genomic DNA using RNeasy Plant Mini Kit

and on-column DNAse I treatment (Qiagen), following supplier's indications. cDNA was obtained using

the Maxima^{TM} H Minus retrotranscriptase (ThermoFisher), from 50-250 ng of total RNA. cDNA extracts

were diluted at 1/5-1/8 before use. RT qPCR was performed in a 384-well thermocycler

(QuantStudioTM^5, ThermoFisher), using SsoAdvanced Universal SYBR Green Supermix (BioRad) and

appropriate primers. The thermocycler was programmed to run for 3 min at 95°C, followed by 40-45

cycles of 10 sec at 95°C and 30 sec at 60°C. Each biological replicate was run twice to assess technical

variation. Expression of genes of interest was normalized by two reference genes among *PpElig2*

(*Pp3c14_21480*), *PpAPT* (*Pp3c8_16590*) and *PpACT3* (*Pp3c10_17080*) (all three are expressed at

similar levels (Le Bail et al. 2013). Relative expression was calculated as RE = 2-CT$_{gene}$/2-CT$_{ref}$ where

CT$_{ref}$ is the mean value of the two reference genes. For the study of *PpSMXL* genes' expression across

development (figure 2), WT *P. patens* was cultivated in petri dishes from fragmented tissues, or in

Magenta pots (for 35 days). The following tissues were collected: protonema at 6 days (mostly

chloronema in our culture conditions), at 10 days (mix of chloronema and caulonema), and at 14 days

(mix of chloronema, caulonema and buds), and mature gametophores and rhizoids at 35 days. Four

biological replicates were used for each timepoint. For the "response to GR24" experiment, plants were

cultivated from fragmented *protonema* in 6-well plates for 2 weeks in control conditions, then

transferred in the dark for one week, and treated with 1 μM (+)-GR24, or 0.01% DMSO in the dark for

6 hours. Six biological repeats were used for each genotype and treatment. For the "response to light"

experiment, 2-week-old WT, *Ppccd8* and *Ppmax2-1* were similarly transferred in the dark for 5 days,

and then either kept in the dark for 24 hours, kept in control white light conditions for 24 hours, or placed

under constant red-light for 1, 6 or 24 hours. Six biological repeats were used for each genotype and
treatment.

Cloning of *PpSMXL* CDS and promoters

Coding sequence of each *PpSMXL* gene was amplified on WT *P. patens* Gransden cDNA, using Phusion
DNA polymerase (ThermoFisher), following provided instructions and using primers with attB1 and
attB2 extensions (respectively on the forward and reverse primer, see Supplemental Table 1). A similar
strategy was used to amplify promoter sequences (full 5'UTR and 1000 bp upstream, from Phytozome
V3.1). Both CDS and promoters were then integrated into the pDONR207 plasmid using BP clonase II
mix (Thermofisher). pDONR207 plasmids containing *PpSMXL* CDS were submitted to PCR-mediated
mutagenesis to obtain ΔRGK/RT versions.

CRISPR-Cas9 mediated mutagenesis

Coding sequences of *PpSMXL* and *PpMAX2* were used to search for CRISPR RNA (crRNA) contiguous
to a PAM motif recognized by *Streptococcus pyogenes* Cas9 (NGG), using the webtool CRISPOR V4
against *P. patens* genome Phytozome V9 (http://crispor.tefor.net/). crRNAs located in the first third of
the coding sequence, with highest possible specificity score, and fewest possible predicted off-targets,
were selected. Small constructs containing each crRNA fused to either the proU6 or the proU3 snRNA
promoter in 5' U3 or U6 promoter (Collonnier et al. 2017), and to the tracrRNA in 3', encased between
attB1/attB2 GateWay recombination sequences, were synthesized by Twist Biosciences. These inserts
were then cloned into pDONR207 vectors. Polyethylene glycol–mediated protoplast transformation was
performed with multiple pDONR207-sgRNA according to the protocol described by Lopez-Obando et
al., 2016. Mutations of the *PpSMXL* genes were confirmed by PCR amplification of *PpSMXL* loci
around the recognition sequence of each guide RNA and sequencing of the PCR products. Alternatively,
for *PpSMXL* genes, a second strategy was employed where crRNAs were designed in the 5' and 3'UTR
sequences, to completely remove the coding sequence of *PpSMXL* genes from the genome when used
together. Mutants obtained from this second strategy were genotyped by monitoring the size and
sequence of amplicons spanning from the 5'UTR to the 3'UTR.

Generation of *proPpSMXL:GUS*, *proZmUbi:GFP-PpSMXL* and control *proZmUbi:flag-GFP* lines

proPpSMXL:GUS constructs were obtained by LR recombination of pDONR207 plasmids containing
PpSMXL promoters with the pMP1301 destination vector previously described (Lopez-Obando et al.,
2018). This method could not be employed for the promoter of *PpSMXLA*, which had to be amplified
with a NotI forward primer and a AscI reverse primer and was subcloned into the pTOPO-blunt II vector
(ThermoFisher), and then directly inserted into a NotI-AscI digested pMP1301. *proZmUbi:GFP-*
PpSMXL constructs were obtained by LR recombination of pDONR207 plasmids containing *PpSMXL*
coding sequences with the pMP1335 destination vector

(http://labs.biology.ucsd.edu/estelle/Moss_files/pK108N+Ubi-mGFP6-GW.gb). Similarly, the

proZmUbi:flag-GFP construct was obtained using the pMP1382 destination vector. These plasmids

were used independently to transform WT *P. patens* Gransden, together with pDONR207 containing

sgRNA recognizing Pp108 homology sequences contained in the three pMP vectors and appropriate

Cas9 and selection plasmids (Lopez-Obando et al., 2016b). Obtained G418 resistant lines were screened

for insertion using PCR (with proPpSMXL forward and GUS reverse, GFP forward and PpSMXL

reverse, or proZmUbi forward and GFP reverse primers, respectively).

GUS staining

690 Two to six independent G418 resistant lines with verified *GUS* insertion into the genome were obtained

691 for each *PpSMXL* genes except *PpSMXLB* and used for histochemical analyses. GUS staining of two-

692 week-old *P. patens* plants was carried out following the protocol detailed by Yuji Hiwatashi on the

693 NIBB PHYSCObase website (http://moss.nibb.ac.jp/protocol.html).

Generation of BiFC constructs

Gateway cloned inserts of genes of interest were integrated into pbiFP vectors using LR Clonase II mix

(ThermoFisher). Inserts containing a STOP codon were cloned in pbiFP2 and pbiFP3, those not

containing a STOP were cloned in pbiFP1 and pbiFP4 (when possible, all four vectors were obtained

for a given gene). Resulting vectors were electroporated into *Escherichia coli* DH10B cells and clones

were selected on spectinomycin. In phase integration of the coding sequence relative to the half eYFP

tag was checked by sequencing insert's ends. The previously published GLOBOSA/DEFICIENS

interaction (Azimzadeh et al. 2008) was consistently used a positive control of eYFP reconstruction.

Agroinfiltration of *Nicotiana benthamiana* leaves

pbiFP plasmids containing the genes of interest were electroporated into *Agrobacterium tumefaciens*

strain C58C1. Agrobacteria were incubated for 18 hours at 28°C under constant agitation and then

pelleted, washed twice, and resuspended in infiltration buffer (13 g/L S-medium (Duchefa Biochemie)

and 40 g/L sucrose, pH 5.7 with KOH) to attain an OD600 value of 0.5. To enhance transient expression

of RFP-PpSMXL and BiFC fusion proteins, the P19 viral suppressor of gene silencing from tomato

bushy stunt virus was co-expressed. Equal volumes of needed bacterial cultures were mixed and

infiltrated into the abaxial epidermis of 4–5-week-old *Nicotiana benthamiana* leaves. After incubation

at 25 °C (16 h light/8 h dark) for 4 days, leaves were harvested on wet paper and kept in similar

temperature and hygrometry conditions for short-term preservation until observation.

Confocal microscopy observations

713 Fragments of *P. patens proZmUBI:GFP-PpSMXL* plants and infiltrated parts of *Nicotiana benthamiana*

714 leaves were both observed on a TCS SP5 inverted or on a TCS SP8 Upright confocal microscopy system

(Leica), with a 20X objective. GFP fluorescence was acquired in the 495nm-520nm λ range, eYFP in the 525nm-540nm range, RFP in the 570nm-610nm range and CFP in the 465nm-505nm range. Signals in the 700nm-750nm range were attributed to chlorophyll autofluorescence. Lasers used for excitation have a peak wavelength of 488nm (GFP), 514nm (YFP), 458nm (CFP) and 561nm (RFP).

Constructs and generation of Arabidopsis transgenic lines

The expression vectors for transgenic Arabidopsis were constructed by MultiSite Gateway Three-Fragment Vector Construction kit (Invitrogen). AtSMAX/SMXL and PpSMXL constructs were tagged with m-Citrine protein at their C-terminus. Lines were resistant to hygromycin. For the Arabidopsis *SMAX1* (AT5G57710), *SMXL5* (A5G57130) and *SMXL6* (AT1G07200) promotors cloning, a 3128-bp fragment upstream from SMAX1 start codon, a 3028-bp fragment upstream from SMXL5 start codon and a 3058-bp fragment upstream from SMXL6 start codon were amplified from Col-0 genomic DNA using the primers described in Supplemental Table S1, and were cloned into the pDONR-P4P1R vector, using Gateway recombination (Invitrogen) as described in (Lopez-Obando et al. 2021). The m-Citrine tag was cloned into pDONR-P2RP3 (Invitrogen) as described in (de Saint Germain et al. 2016). AtSMAX1, AtSMXL5 and AtSMXL6 CDS were PCR amplified from Arabidopsis Col-0 cDNA and with the primers specified in Supplemental Table S1 and then recombined into the pDONR221 vector (Invitrogen). PpSMXLB and PpSMXLC CDS were obtained as described above. The suitable combinations of AtSMXL native promoters, AtSMXL or PpSMXL CDS, and m-Citrine tag were cloned into the pH7m34GW final destination vectors by using the three fragment recombination system (Karimi et al. 2007) and were thusly named *pSMAX1:SMAX1, pSMAX1:PpSMXLB, pSMAX1:PpSMXLC, pSMXL5:SMXL5, pSMXL5:PpSMXLB, pSMXL5:PpSMXLC, pSMXL6:SMXL6, pSMXL6:PpSMXLB and pSMXL6:PpSMXLC*. Transformation of Arabidopsis *smax1-1*, *smxl4-1 smxl5-1* and *smxl6/7/8* mutants was performed according to the conventional floral dipping method (Clough and Bent 1998), with Agrobacterium strain GV3101. For each construct, at least 12 independent T1 lines were isolated, and then 2 to 4 lines were selected in T2 for showing a 3 :1 segregation (single T-DNA insertion). Phenotypic analysis shown in Figure 10 and Supplemental Figures S18 and S19 were performed on the T3 homozygous lines.

Plant materials, growth conditions, and phenotypic assays

Arabidopsis thaliana plants used in this study originated from the Columbia (Col-0) ecotype background. *smax1-2* and *smxl678* Arabidopsis mutants are gift Dave Nelson (University of California, Riverside), and the Arabidopsis *smxl4-5* mutant gift from Thomas Greb (Heidelberg University, Heidelberg). The transgenic lines are described above. For *Arabidopsis thaliana* shoot branching assays, the plants were grown in greenhouse. Experiments were carried out in summer, under long photoperiods (15-16 h per day); daily temperatures fluctuated between 18 °C and 25 °C. Peak levels of PAR were between 700 and 1000 µmol m^{-2} s^{-1}. Plants were watered twice a week with tap water. The number of

caulinary branch and plant height was scored when the plants were 35 days old. Leaf Morphology Assay was performed as described in (Soundappan et al. 2015) on the 5th leaf of each plant marked with indelible marker at 3 weeks post-germination. Plants were cultivated as for branching assay. 12 plants were then used for Kruskal Wallis Dunn *post-hoc* test VS Col0. Hypocotyl elongation assays were performed as described in (Guercio et al. 2022) with the following modification: 11-day-old seedlings were photographed and hypocotyl lengths were quantified using ImageJ. 2 plates of 20 to 24 seeds were sown for each genotype. The data from the 45 seedlings were then used for a one-way ANOVA de Welch, Dunnett T3 *posthoc* VS Col0 ($n = 45-129$, $p<5\%$). For lateral Root Assay seedlings were grown on 3 different plates in the same media and condition as for hypocotyl assay. Roots were analyzed at 15 days post-germination (dpg). Primary root length was then measured using ImageJ. 24 seedlings per genotype were used for Kruskal Wallis followed by a Dunn *post-hoc* test.

Chemicals

GR24 enantiomers and (−)-desmethyl-GR24 were produced by V Steinmetz and F-D Boyer using organic synthesis and chiral separation as described in (de Saint Germain et al. 2021). Chemicals were diluted in DMSO.

RT-PCR analysis in Arabidopsis

Semi-quantitative RT-PCR analyses of Arabidopsis transformants were performed from leaf or seedling RNAs extracted and rid of contaminant genomic DNA using RNeasy Plant Mini Kit and on-column DNAse I treatment (Qiagen), following supplier's indications. cDNA was obtained using the H Minus retrotranscriptase (ThermoFisher), from 250 ng of total RNA. cDNA extracts were diluted twice, and 1μL was used for each PCR reaction. The thermocycler was programmed to run for 3 min at 95 °C, followed by 28 cycles of 30 sec at 95 °C, 30 sec at 56 °C and 40 sec at 72 °C.

Statistical analysis of results

Kruskal-Wallis, Mann-Whitney and *post-hoc* Dunn, Dunnett or Tukey multiple comparisons tests (details in figures legends) were carried out either in R 3.6.3 or in GraphPad Prism 8.4.2. Tests employed were mostly non-parametric as normality of distributions and/or homoscedasticity among groups could not be confirmed in most experiments (Kruskal-Wallis tests for multiple comparisons and Mann-Whitney for single comparisons, unless otherwise stated in legends). For some gene expression experiments, data points were excluded based on an outliers' search (Grubb's, $\alpha=0.05$) on in GraphPad Prism 8.4.2. Unless otherwise defined, used statistical significance scores are as follow: # $0.05 \leq p<0.1$, * $0.01 \leq p<0.05$, ** $0.001 \leq p<0.01$, *** $p<0.001$. Same letters scores indicate that $p \geq 0.05$ (non-significant differences).

Accession Numbers

Moss sequences used in the present article can be found on Phytozome (*P. patens* Gransden genome, V3.1 version). *PpSMXLA* is *Pp3c2_14220*, *PpSMXLB* is *Pp3c1_23530*, *PpSMXLC* is *Pp3c9_16100* and *PpSMXLD* is *Pp3c15_16120*. *PpMAX2* corresponds to *Pp3c17_1180*, *PpCCD8 to Pp3c6_21520*, *PpAPT* to *Pp3c8_16590*, *PpACT3* to *Pp3c10_17080,* and *PpElig2* corresponds to *Pp3c14_21480*.

Authors contributions

A.G and S.B are the main authors of the manuscript and A.G performed most of the experiments. Some steps in genotyping mutant and transgenic lines and support with molecular cloning was given by P.L.B, K.B, A.L and S.B helped for moss mutant phenotyping and J-P. P for transgenic Arabidopsis phenotyping. V.S and F-D. B furnished SL and KL analogs. Plasmids used for *Ppsmxl* mutant generation in the first strategy, as well as higher order *Ppsmxl* mutants were obtained by M.L-O. A.G, S.B, A.S.G, and C.R all contributed to experimental design and to the argumentation developed herein.

Acknowledgements

We thank Dave Nelson (University of California, Riverside) for the generous gift of Arabidopsis *smax1-2* and *smxl678* mutants, as well as for the SMXL6p:SMXL6-GFP construct. We thank Thomas Greb (Heidelberg University, Heidelberg) for the kind gift of Arabidopsis *smxl4-5* mutant. We thank Fabien Nogué (IJPB, Versailles) for his helpful advice and for giving us guide RNAs targeting the Pp108 non-coding locus. We thank Martine Pastuglia (IJPB, Versailles) for the gift of BiFC control constructs and Michael J. Prigge (University of California San Diego, La Jolla) for nicely sending us the pMP destination vectors. This work has benefited from the support of IJPB's Plant Observatory technological platforms.

Fundings

The IJPB benefits from the support of Saclay Plant Sciences-SPS (ANR-17-EUR-0007). This work has benefited from a French State grant (Saclay Plant Sciences, reference n° ANR-17-EUR-0007, EUR SPS-GSR) managed by the French National Research Agency under an Investments for the Future program integrated into France 2030 (reference n° ANR-11-IDEX-0003-02).

References

Akiyama K, Matsuzaki K, Hayashi H (2005) Plant sesquiterpenes induce hyphal branching in arbuscular mycorrhizal fungi. Nature 435 (7043):824-827. doi:10.1038/nature03608

Alder A, Jamil M, Marzorati M, Bruno M, Vermathen M, Bigler P, Ghisla S, Bouwmeester H, Beyer P,
Al-Babili S (2012) The path from beta-carotene to carlactone, a strigolactone-like plant
hormone. Science 335 (6074):1348-1351. doi:10.1126/science.1218094
Azimzadeh J, Nacry P, Christodoulidou A, Drevensek S, Camilleri C, Amiour N, Parcy F, Pastuglia M,
Bouchez D (2008) Arabidopsis TONNEAU1 proteins are essential for preprophase band
formation and interact with centrin. Plant Cell 20 (8):2146-2159. doi:tpc.107.056812 [pii]

10.1105/tpc.107.056812
Bennett T, Liang Y, Seale M, Ward S, Muller D, Leyser O (2016) Strigolactone regulates shoot
development through a core signalling pathway. Biol Open 5 (12):1806-1820.
doi:10.1242/bio.021402
Besserer A, Becard G, Jauneau A, Roux C, Sejalon-Delmas N (2008) GR24, a synthetic analog of
strigolactones, stimulates the mitosis and growth of the arbuscular mycorrhizal fungus
Gigaspora rosea by boosting its energy metabolism. Plant Physiol 148 (1):402-413
Besserer A, Puech-Pages V, Kiefer P, Gomez-Roldan V, Jauneau A, Roy S, Portais JC, Roux C, Becard
G, Sejalon-Delmas N (2006) Strigolactones stimulate arbuscular mycorrhizal fungi by
activating mitochondria. PLoS Biol 4 (7):e226
Bianchi A, Giorgi C, Ruzza P, Toniolo C, Milner-White EJ (2012) A synthetic hexapeptide designed to
resemble a proteinaceous P-loop nest is shown to bind inorganic phosphate. Proteins 80
(5):1418-1424. doi:10.1002/prot.24038
Bonhomme S, Guillory A (2022) Synthesis and Signalling of Strigolactone and KAI2-Ligand signals in
bryophytes. J Exp Bot. doi:10.1093/jxb/erac186
Bythell-Douglas R, Rothfels CJ, Stevenson DWD, Graham SW, Wong GK, Nelson DC, Bennett T
(2017) Evolution of strigolactone receptors by gradual neo-functionalization of KAI2
paralogues. BMC biology 15 (1):52. doi:10.1186/s12915-017-0397-z
Choi J, Lee T, Cho J, Servante EK, Pucker B, Summers W, Bowden S, Rahimi M, An K, An G,
Bouwmeester HJ, Wallington EJ, Oldroyd G, Paszkowski U (2020) The negative regulator
SMAX1 controls mycorrhizal symbiosis and strigolactone biosynthesis in rice. Nat Commun
11 (1):2114. doi:10.1038/s41467-020-16021-1
Clough SJ, Bent AF (1998) Floral dip: a simplified method for Agrobacterium-mediated transformation
of Arabidopsis thaliana. Plant J 16 (6):735-743. doi:10.1046/j.1365-313x.1998.00343.x
Collonnier C, Guyon-Debast A, Maclot F, Mara K, Charlot F, Nogué F (2017) Towards mastering
CRISPR-induced gene knock-in in plants: Survey of key features and focus on the model
Physcomitrella patens. Methods 121-122:103-117. doi:10.1016/j.ymeth.2017.04.024
Conn CE, Nelson DC (2016) Evidence that KARRIKIN-INSENSITIVE2 (KAI2) Receptors may
Perceive an Unknown Signal that is not Karrikin or Strigolactone. Front Plant Sci 6:1219.
doi:10.3389/fpls.2015.01219
Cook CE, Whichard LP, Turner B, Wall ME, Egley GH (1966) Germination of Witchweed (Striga lutea
Lour.): Isolation and Properties of a Potent Stimulant. Science 154 (3753):1189-1190
Coudert Y, Palubicki W, Ljung K, Novak O, Leyser O, Harrison CJ (2015) Three ancient hormonal cues
co-ordinate shoot branching in a moss. Elife 4. doi:10.7554/eLife.06808
de Saint Germain A, Clave G, Badet-Denisot MA, Pillot JP, Cornu D, Le Caer JP, Burger M, Pelissier
F, Retailleau P, Turnbull C, Bonhomme S, Chory J, Rameau C, Boyer FD (2016) An histidine
covalent receptor and butenolide complex mediates strigolactone perception. Nat Chem Biol 12
(10):787-794. doi:10.1038/nchembio.2147
de Saint Germain A, Jacobs A, Brun G, Pouvreau JB, Braem L, Cornu D, Clavé G, Baudu E, Steinmetz
V, Servajean V, Wicke S, Gevaert K, Simier P, Goormachtig S, Delavault P, Boyer FD (2021)
A Phelipanche ramosa KAI2 protein perceives strigolactones and isothiocyanates
enzymatically. Plant Commun 2 (5):100166. doi:10.1016/j.xplc.2021.100166
Decker EL, Alder A, Hunn S, Ferguson J, Lehtonen MT, Scheler B, Kerres KL, Wiedemann G, Safavi-
Rizi V, Nordzieke S, Balakrishna A, Baz L, Avalos J, Valkonen JPT, Reski R (2017)
Strigolactone biosynthesis is evolutionarily conserved, regulated by phosphate starvation and
contributes to resistance against phytopathogenic fungi in a moss, Physcomitrella patens. New
Phytol 216 (2):455-468. doi:10.1111/nph.14506

Delaux PM, Xie X, Timme RE, Puech-Pages V, Dunand C, Lecompte E, Delwiche CF, Yoneyama K, Becard G, Sejalon-Delmas N (2012) Origin of strigolactones in the green lineage. New Phytol 195 (4):857-871. doi:10.1111/j.1469-8137.2012.04209.x

Gomez-Roldan V, Fermas S, Brewer PB, Puech-Pages V, Dun EA, Pillot JP, Letisse F, Matusova R, Danoun S, Portais JC, Bouwmeester H, Becard G, Beveridge CA, Rameau C, Rochange SF (2008) Strigolactone inhibition of shoot branching. Nature 455 (7210):189-194. doi:10.1038/nature07271

Guercio AM, Torabi S, Cornu D, Dalmais M, Bendahmane A, Le Signor C, Pillot JP, Le Bris P, Boyer FD, Rameau C, Gutjahr C, de Saint Germain A, Shabek N (2022) Structural and functional analyses explain Pea KAI2 receptor diversity and reveal stereoselective catalysis during signal perception. Commun Biol 5 (1):126. doi:10.1038/s42003-022-03085-6

Hoffmann B, Proust H, Belcram K, Labrune C, Boyer FD, Rameau C, Bonhomme S (2014) Strigolactones inhibit caulonema elongation and cell division in the moss Physcomitrella patens. PLoS One 9 (6):e99206. doi:10.1371/journal.pone.0099206

Jiang L, Liu X, Xiong G, Liu H, Chen F, Wang L, Meng X, Liu G, Yu H, Yuan Y, Yi W, Zhao L, Ma H, He Y, Wu Z, Melcher K, Qian Q, Xu HE, Wang Y, Li J (2013) DWARF 53 acts as a repressor of strigolactone signalling in rice. Nature 504 (7480):401-405. doi:10.1038/nature12870

Jones D.T., Taylor W.R., and Thornton J.M. (1992) The rapid generation of mutation data matrices from protein sequences. *Computer Applications in the Biosciences* **8**: 275-282.

Karimi M, Bleys A, Vanderhaeghen R, Hilson P (2007) Building blocks for plant gene assembly. Plant Physiol 145 (4):1183-1191. doi:10.1104/pp.107.110411

Kerr SC, Patil SB, de Saint Germain A, Pillot JP, Saffar J, Ligerot Y, Aubert G, Citerne S, Bellec Y, Dun EA, Beveridge CA, Rameau C (2021) Integration of the SMXL/D53 strigolactone signalling repressors in the model of shoot branching regulation in Pisum sativum. Plant J 107 (6):1756-1770. doi:10.1111/tpj.15415

Khosla A, Morffy N, Li Q, Faure L, Chang SH, Yao J, Zheng J, Cai ML, Stanga J, Flematti GR, Waters MT, Nelson DC (2020) Structure-Function Analysis of SMAX1 Reveals Domains That Mediate Its Karrikin-Induced Proteolysis and Interaction with the Receptor KAI2. Plant Cell 32 (8):2639-2659. doi:10.1105/tpc.19.00752

Kodama K, Rich MK, Yoda A, Shimazaki S, Xie X, Akiyama K, Mizuno Y, Komatsu A, Luo Y, Suzuki H, Kameoka H, Libourel C, Keller J, Sakakibara K, Nishiyama T, Nakagawa T, Mashiguchi K, Uchida K, Yoneyama K, Tanaka Y, Yamaguchi S, Shimamura M, Delaux PM, Nomura T, Kyozuka J (2022) An ancestral function of strigolactones as symbiotic rhizosphere signals. Nat Commun 13 (1):3974. doi:10.1038/s41467-022-31708-3

Kumar S., Stecher G., Li M., Knyaz C., and Tamura K. (2018) MEGA X: Molecular Evolutionary Genetics Analysis across computing platforms. *Molecular Biology and Evolution* **35**:1547-1549.

Kyozuka J, Nomura T, Shimamura M (2022) Origins and evolution of the dual functions of strigolactones as rhizosphere signaling molecules and plant hormones. Curr Opin Plant Biol 65:102154. doi:10.1016/j.pbi.2021.102154

Le Bail A, Scholz S, Kost B (2013) Evaluation of reference genes for RT qPCR analyses of structure-specific and hormone regulated gene expression in Physcomitrella patens gametophytes. PLoS One 8 (8):e70998. doi:10.1371/journal.pone.0070998

Liang Y, Ward S, Li P, Bennett T, Leyser O (2016) SMAX1-LIKE7 Signals from the Nucleus to Regulate Shoot Development in Arabidopsis via Partially EAR Motif-Independent Mechanisms. Plant Cell 28 (7):1581-1601. doi:10.1105/tpc.16.00286

Lopez-Obando M, Conn CE, Hoffmann B, Bythell-Douglas R, Nelson DC, Rameau C, Bonhomme S (2016a) Structural modelling and transcriptional responses highlight a clade of PpKAI2-LIKE genes as candidate receptors for strigolactones in Physcomitrella patens. Planta 243 (6):1441-1453. doi:10.1007/s00425-016-2481-y

Lopez-Obando M, de Villiers R, Hoffmann B, Ma L, de Saint Germain A, Kossmann J, Coudert Y, Harrison CJ, Rameau C, Hills P, Bonhomme S (2018) Physcomitrella patens MAX2 characterization suggests an ancient role for this F-box protein in photomorphogenesis rather than strigolactone signalling. New Phytol 219 (2):743-756. doi:10.1111/nph.15214

Lopez-Obando M, Guillory A, Boyer FD, Cornu D, Hoffmann B, Le Bris P, Pouvreau JB, Delavault P,
Rameau C, de Saint Germain A, Bonhomme S (2021) The Physcomitrium (Physcomitrella)
patens PpKAI2L receptors for strigolactones and related compounds function via MAX2-
dependent and -independent pathways. Plant Cell 33 (11):3487-3512.
doi:10.1093/plcell/koab217

Lopez-Obando M, Hoffmann B, Géry C, Guyon-Debast A, Téoulé E, Rameau C, Bonhomme S, Nogué
F (2016b) Simple and Efficient Targeting of Multiple Genes Through CRISPR-Cas9 in
Physcomitrella patens. G3 (Bethesda) 6 (11):3647-3653. doi:10.1534/g3.116.033266

Ma H, Duan J, Ke J, He Y, Gu X, Xu TH, Yu H, Wang Y, Brunzelle JS, Jiang Y, Rothbart SB, Xu HE,
Li J, Melcher K (2017) A D53 repression motif induces oligomerization of TOPLESS
corepressors and promotes assembly of a corepressor-nucleosome complex. Science advances
3 (6):e1601217. doi:10.1126/sciadv.1601217

Machin DC, Hamon-Josse M, Bennett T (2020) Fellowship of the rings: a saga of strigolactones and
other small signals. New Phytol 225 (2):621-636. doi:10.1111/nph.16135

Mizuno Y, Komatsu A, Shimazaki S, Naramoto S, Inoue K, Xie X, Ishizaki K, Kohchi T, Kyozuka J
(2021) Major components of the KARRIKIN INSENSITIVE2-dependent signaling pathway are
conserved in the liverwort Marchantia polymorpha. Plant Cell 33 (7):2395-2411.
doi:10.1093/plcell/koab106

Moturu TR, Thula S, Singh RK, Nodzynski T, Vareková RS, Friml J, Simon S (2018) Molecular
evolution and diversification of the SMXL gene family. J Exp Bot 69 (9):2367-2378.
doi:10.1093/jxb/ery097

Ortiz-Ramirez C, Hernandez-Coronado M, Thamm A, Catarino B, Wang M, Dolan L, Feijo JA, Becker
JD (2016) A Transcriptome Atlas of Physcomitrella patens Provides Insights into the Evolution
and Development of Land Plants. Mol Plant 9 (2):205-220. doi:10.1016/j.molp.2015.12.002

Perroud PF, Haas FB, Hiss M, Ullrich KK, Alboresi A, Amirebrahimi M, Barry K, Bassi R, Bonhomme
S, Chen H, Coates JC, Fujita T, Guyon-Debast A, Lang D, Lin J, Lipzen A, Nogué F, Oliver
MJ, Ponce de León I, Quatrano RS, Rameau C, Reiss B, Reski R, Ricca M, Saidi Y, Sun N,
Szövényi P, Sreedasyam A, Grimwood J, Stacey G, Schmutz J, Rensing SA (2018) The
Physcomitrella patens gene atlas project: large-scale RNA-seq based expression data. Plant J 95
(1):168-182. doi:10.1111/tpj.13940

Proust H, Hoffmann B, Xie X, Yoneyama K, Schaefer DG, Nogue F, Rameau C (2011) Strigolactones
regulate protonema branching and act as a quorum sensing-like signal in the moss
Physcomitrella patens. Development 138 (8):1531-1539. doi:10.1242/dev.058495

Radhakrishnan GV, Keller J, Rich MK, Vernié T, Mbadinga Mbadinga DL, Vigneron N, Cottret L,
Clemente HS, Libourel C, Cheema J, Linde AM, Eklund DM, Cheng S, Wong GKS,
Lagercrantz U, Li FW, Oldroyd GED, Delaux PM (2020) An ancestral signalling pathway is
conserved in intracellular symbioses-forming plant lineages. Nat Plants 6 (3):280-289.
doi:10.1038/s41477-020-0613-7

Soundappan I, Bennett T, Morffy N, Liang Y, Stanga JP, Abbas A, Leyser O, Nelson D (2015) SMAX1-
LIKE/D53 Family Members Enable Distinct MAX2-Dependent Responses to Strigolactones
and Karrikins in Arabidopsis. Plant Cell 27 (11):3143-3159. doi:10.1105/tpc.15.00562

Stanga JP, Morffy N, Nelson DC (2016) Functional redundancy in the control of seedling growth by the
karrikin signaling pathway. Planta 243 (6):1397-1406. doi:10.1007/s00425-015-2458-2

Stanga JP, Smith SM, Briggs WR, Nelson DC (2013) SUPPRESSOR OF MORE AXILLARY
GROWTH2 1 controls seed germination and seedling development in Arabidopsis. Plant
Physiol 163 (1):318-330. doi:10.1104/pp.113.221259

Sun YK, Flematti GR, Smith SM, Waters MT (2016) Reporter Gene-Facilitated Detection of
Compounds in Arabidopsis Leaf Extracts that Activate the Karrikin Signaling Pathway. Front
Plant Sci 7:1799. doi:10.3389/fpls.2016.01799

Temmerman A, Guillory A, Bonhomme S, Goormachtig S, Struk S (2022) Masks Start to Drop:
Suppressor of MAX2 1-Like Proteins Reveal Their Many Faces. Front Plant Sci 13:887232.
doi:10.3389/fpls.2022.887232

Umehara M, Hanada A, Yoshida S, Akiyama K, Arite T, Takeda-Kamiya N, Magome H, Kamiya Y,
Shirasu K, Yoneyama K, Kyozuka J, Yamaguchi S (2008) Inhibition of shoot branching by new
terpenoid plant hormones. Nature 455 (7210):195-200. doi:10.1038/nature07272

Walker CH, Siu-Ting K, Taylor A, O'Connell MJ, Bennett T (2019) Strigolactone synthesis is ancestral in land plants, but canonical strigolactone signalling is a flowering plant innovation. BMC biology 17 (1):70. doi:10.1186/s12915-019-0689-6

Wallner ES, López-Salmerón V, Belevich I, Poschet G, Jung I, Grünwald K, Sevilem I, Jokitalo E, Hell R, Helariutta Y, Agustí J, Lebovka I, Greb T (2017) Strigolactone- and Karrikin-Independent SMXL Proteins Are Central Regulators of Phloem Formation. Curr Biol 27 (8):1241-1247. doi:10.1016/j.cub.2017.03.014

Wallner ES, Tonn N, Shi D, Jouannet V, Greb T (2020) SUPPRESSOR OF MAX2 1-LIKE 5 promotes secondary phloem formation during radial stem growth. Plant J 102 (5):903-915. doi:10.1111/tpj.14670

Wang L, Wang B, Jiang L, Liu X, Li X, Lu Z, Meng X, Wang Y, Smith SM, Li J (2015) Strigolactone Signaling in Arabidopsis Regulates Shoot Development by Targeting D53-Like SMXL Repressor Proteins for Ubiquitination and Degradation. Plant Cell 27 (11):3128-3142. doi:10.1105/tpc.15.00605

Wang L, Wang B, Yu H, Guo H, Lin T, Kou L, Wang A, Shao N, Ma H, Xiong G, Li X, Yang J, Chu J, Li J (2020) Transcriptional regulation of strigolactone signalling in Arabidopsis. Nature 583 (7815):277-281. doi:10.1038/s41586-020-2382-x

Waters MT, Scaffidi A, Moulin SL, Sun YK, Flematti GR, Smith SM (2015) A Selaginella moellendorffii Ortholog of KARRIKIN INSENSITIVE2 Functions in Arabidopsis Development but Cannot Mediate Responses to Karrikins or Strigolactones. Plant Cell 27 (7):1925-1944. doi:10.1105/tpc.15.00146

Yao J, Scaffidi A, Meng Y, Melville KT, Komatsu A, Khosla A, Nelson DC, Kyozuka J, Flematti GR, Waters MT (2021) Desmethyl butenolides are optimal ligands for karrikin receptor proteins. New Phytol 230 (3):1003-1016. doi:10.1111/nph.17224

Zheng J, Hong K, Zeng L, Wang L, Kang S, Qu M, Dai J, Zou L, Zhu L, Tang Z, Meng X, Wang B, Hu J, Zeng D, Zhao Y, Cui P, Wang Q, Qian Q, Wang Y, Li J, Xiong G (2020) Karrikin Signaling Acts Parallel to and Additively with Strigolactone Signaling to Regulate Rice Mesocotyl Elongation in Darkness. Plant Cell 32 (9):2780-2805. doi:10.1105/tpc.20.00123

Zhou F, Lin Q, Zhu L, Ren Y, Zhou K, Shabek N, Wu F, Mao H, Dong W, Gan L, Ma W, Gao H, Chen J, Yang C, Wang D, Tan J, Zhang X, Guo X, Wang J, Jiang L, Liu X, Chen W, Chu J, Yan C, Ueno K, Ito S, Asami T, Cheng Z, Lei C, Zhai H, Wu C, Wang H, Zheng N, Wan J (2013) D14-SCF(D3)-dependent degradation of D53 regulates strigolactone signalling. Nature 504 (7480):406-410. doi:10.1038/nature12878

Figure 1

Figure 1 Expression of *PpSMXL* genes along *P. patens* vegetative development. A, GUS staining of 2-week old *proPpSMXL:GUS* plants. Scale bars = 500 μm. B, Transcript levels of the four *PpSMXL* genes, relative to the two reference genes *PpACT3* (*Pp3c10_17080*) and *PpAPT* (*Pp3c8_16590*). RT-qPCR data used for the analysis was extracted from 4 biological replicates, each with 2 technical repeats. Each point represents the mean of two technical repeats. Some points were excluded from analysis following an outliers identification test carried out in GraphPad Prism (version 8.4.2). Statistical significance scores of comparisons among tissues are indicated as bold letters (Kruskal-Wallis test followed by a Dunn *post-hoc* test, $p<5\%$). Note the difference in expression values on y axes.

Figure 2

Figure 2 Phenotypes of the *Ppsmxl* mutants. *Ppsmxl* mutants were grown in white light long day conditions (A-B) , or under continuous red light (C-F).

A, Three-week-old plants grown in the light on low nitrogen content medium, without underlying cellophane. Bar = 5 mm. B, Diameters of six-week-old plants; n=35-49 plants of each genotype grown on 7 individual plates. Statistical significance of comparisons between all genotypes are indicated by bold symbols; left panel: Kruskal Wallis followed by a Dunn *post-hoc* test, n=42-49, p<5%; right panel: standard ANOVA followed by a Tukey *post-hoc* test, n=20-35, p<5%). C, E, Examples of gametophores from two-month-old plants grown in red light. Bar = 3 mm D, F, Gametophore height measured for 30 two-month-old gametophores of each genotype. Statistical significance of comparisons between all genotypes are indicated by bold symbols, Kruskal Wallis followed by a Dunn *post-hoc* test, n⩾30, p<5%.

Figure 3 Growth of *proZmUbi:GFP-PpSMXL* lines. (A-B): Plant extension in control conditions. A, Representative individuals after 5 weeks of growth. Scale bar = 1 cm. B, Diameters were measured from 6 week-old plants, n = 21-35, grown on at least 3 different plates. Statistical significance of comparisons between all genotypes are indicated as bold symbols (Welch ANOVA followed by a Dunnett post-hoc test, p<5%). (C-D) Gametophore height in red light. C, Representative individuals after two months culture under continuous red light. Scale bar = 1 mm. D, Gametophore height was measured for n = 28-30 two-month-old gametophores of each genotype. Statistical significance of comparisons between all genotypes are indicated by bold symbols (standard ANOVA followed by a Tukey *post-hoc* test, p<5%).

Figure 4 Genetic analysis of *PpSMXL* relationship with *PpCCD8*. A, Phenotype of three-week-old plants on low nitrogen content medium (without underlying cellophane). Scale bar = 5 mm. B, Extension phenotype of some of these mutants on low nitrogen content medium (with underlying cellophane), after 5 weeks growth. Statistical significance of comparisons relative to WT at the last time point is shown as bold black letters (Kruskal Wallis followed by a Dunn *post-hoc* test, n = 35-40 , p<5%).

Figure 5 Genetic analysis of *PpSMXL* relationship with *PpMAX2*. (A-B): Plant extension in control conditions. A, Representative individuals after 5 weeks of growth. Bar = 1 cm. B, Plant diameters of the mutants grown on low nitrogen content medium with underlying cellophane for 5 weeks. N ≥ 35 individuals grown on 7 different plates. (C-D) Gametophore height in red light. C, Representative gametophores after two months culture under constant red light. Scale bar = 3 mm. D, Gametophore height was measured for 30 two-month-old gametophores of each genotype. Each point represents a measurement. N ≥ 43 gametophores from 3 different Magenta boxes. (B,D): Statistical significance of comparisons between all genotypes are indicated by bold letters (Kruskal Wallis followed by a Dunn *post-hoc* test, p<0.05).

Figure 6 *Ppsmxl* mutant response to (±)-GR24 enantiomers. A, Response to (+)-GR24: n = 24 two-week-old plants of each genotype were mock treated with 0,01% DMSO (grey) or with 1 µM of (+)-GR24 (blue). Plants were incubated vertically in the dark for ten days. Negatively gravitropic caulonema filaments were enumerated for each plant. Statistical significance of comparisons of control groups relative to WT is shown as black symbols (Kruskal Wallis followed by a Dunn *post-hoc* test). Statistical significance of comparisons between control and treated for each genotype is shown as blue symbols (Mann-Whitney tests).

B, Response to (−)-desmethyl-GR24: n = 24 two-week-old plants of each genotype were mock treated with 0,01% DMSO (grey), 1 µM of (−)-desmethyl-GR24 (orange) or 10 µM of (−)-desmethyl-GR24 (brown). Statistical significance of comparisons of control groups relative to WT is shown as black symbols (Kruskal Wallis followed by a Dunn *post-hoc* test). Statistical significance of comparisons between control and treated for each genotype is shown as orange symbols (Kruskal Wallis tests followed by Tukey *post-hoc* tests).

A and B. For all statistical analyses, p-values are reported as * $0.01 \leq p < 0.05$, ** $0.001 \leq p < 0.01$, *** $p < 0.001$.

	GFP	Autofluorescence	Merge

GFP-PpSMXLA

GFP-PpSMXLB

GFP-PpSMXLC

GFP-PpSMXLD

Figure 7 Subcellular localization of GFP-PpSMXL fusion proteins in *P. patens*. Confocal images of protonema filaments of transgenic *P. patens* lines expressing *proZmUbi:GFP-PpSMXL* fusions as indicated on the left. Right images are overlays of autofluorescence (red) and GFP (green) channels. Bar = 50 µm. Arrow points to a nucleus.

Figure 8 PpSMXL protein interaction assays in *Nicotiana benthamiana*. All four PpSMXL can interact with PpMAX2 (A-D). PpSMXLB and PpSMXLD can form homo-oligomers (E,F). All four PpSMXL can interact with PpKAI2L-C (G-J)**.** Below each tryptic of images, the first indicated protein is fused to the N-terminal part of eYFP, while the second protein is fused to the C-terminal part (both tags are fused at the N-terminal end of *P. patens* proteins). Colocalization of CFP-H2b and eYFP biFC signals are pointed at with white arrows on. Merge are overlays of CFP (blue), YFP (green) and bright field images. Bar = 50 µm. Tested interactions with DEFICIENS and GLOBOSA proteins from *Antirrhinum major* (negative controls) are shown in Supplemental Figure S17. The previously published GLOBOSA/DEFICIENS interaction (Azimzadeh et al., 2008) was consistently used a positive control of eYFP reconstruction.

Figure 9 Current model of KL and SL signaling in *P. patens*. SLs and putative KL signal are mimicked by (+)-GR24 and (−)-desmethyl-GR24 respectively. The two pathways have opposite effects on protonema extension and gametophore development. In contrast to angiosperms, only the KL pathway is PpMAX2 dependent. All four PpSMXL are likely suppressors of the KL pathway, though their PpMAX2-dependent degradation following KL/(−)-desmethyl-GR24 perception has not been demonstrated (question mark). Green dotted arrow indicates that PpSMXL could be stabilized by SLs. A further level of regulation through possible transcriptional regulation of *PpSMXL* gene transcription by PpSMXL proteins has not been represented.

Figure 10 *Atsmxl4/5* **mutant root phenotype is complemented by PpSMXLB and PpSMXLC**. A, Root length was measured from 15 day-old seedlings grown vertically on 0.5 x MS with 1% sucrose. 20 to 40 seedlings per genotype, grown on 3 different plates. Statistical significance of comparisons relative to WT is shown as black symbols and statistical significance of comparisons relative to the double mutant *smxl4,5* is shown as red symbols (Kruskal Wallis followed by a Dunn *post-hoc* test, p<5%). B, Representative seedlings from each genotype. Scale bar = 1 cm