1 **Legacy effects of precipitation and land use impact maize growth and microbiome assembly**
2 **under drought stress**
3

4 Authors
5 Joel F. Swift[1,2*], Matthew R. Kolp[1,2,4], Amanda Carmichael[1,4], Natalie E. Ford[1], Paige M.
6 Hansen[2,4], Benjamin A. Sikes[1,2], Manuel Kleiner[3], Maggie R. Wagner[1,2*]
7

8 Affiliations
9 [1] Department of Ecology and Evolutionary Biology, University of Kansas, Lawrence, KS 66045
10

11 [2] Kansas Biological Survey & Center for Ecological Research, University of Kansas, Lawrence,
12 KS 66045
13

14 [3] Department of Plant and Microbial Biology, North Carolina State University, Raleigh, NC,
15 27695
16

17 [4] Current Address: Center for Animal and Human Health in Appalachia, College of Veterinary
18 Medicine, Lincoln Memorial University, Harrogate, TN 37752 (M.R.K.); Department of
19 Conservation and Research, Southeastern Center for Conservation, Atlanta Botanical Garden,
20 Atlanta, GA, 30309 (A.C.); Department of Soil and Crop Science, Colorado State University, Fort
21 Collins, CO 80521 (P.M.H.)
22

23 *Correspondence: joel.swift@ku.edu and maggie.r.wagner@ku.edu
24

Abstract

As the climate changes, plants and their associated microbiomes face greater water limitation and

increased frequency of drought. Historical precipitation patterns can leave a legacy effect on soil

and root-associated microbiomes, but the impact of this conditioning on future drought

performance is poorly understood. Moreover, agricultural practices, like tilling and fertilization,

may override precipitation legacies in natural systems. Precipitation gradients provide a means to

assess these legacy effects. We collected six soil microbiomes across a Kansas precipitation

gradient from both agricultural fields and native prairies. Seedlings of two *Zea mays* genotypes

were inoculated with each soil microbiome in a factorial drought experiment. Droughted plants

exhibited decreased shoot mass accumulation rates and greater root mass relative to shoot mass.

Restructuring of the bacterial root-associated microbiome was apparent, with depletion observed

in Pseudomonadota and enrichment in Actinomycetota, while the fungal microbiome was largely

unaffected by drought. A historical precipitation legacy effect on soil microbiomes interacted with

plants during drought treatment, but only among prairie soils. Prairie soils from historically wetter

locations increased maize shoot biomass under drought more so than agricultural or historically

drier prairie soils. We demonstrate links between legacy effects and drought performance,

suggesting that future drying climates may condition soils to negatively impact plant performance.

Introduction

Increasing drought frequency due to global climate change will impact both soil and plant-

associated microbiomes [1, 2]. Drought directly limits microbial mobility and access to resources,

but also alters plant host traits (e.g., root physiology, morphology, and exudate composition) that

comprise the habitat available to root-associated microorganisms [3]. Historical precipitation

patterns and drought can leave lasting impacts, or legacy effects, on soil and root-associated

microbiomes. Naturally-occurring precipitation gradients—which shape the biogeography of both

plants [4, 5] and soil microorganisms [6, 7]—provide opportunities to study legacy effects of

precipitation on plant-microbe interactions. Historical patterns of water availability across

gradients constrain the functional potential of soil microbiomes [8–10]; however, the implications

for host plants, including agriculturally critical species, are not well understood.

As plants recruit root microbes primarily from the soil [11], we would expect drought-

induced changes in the soil microbiome to be reflected in root-associated microbiomes. However,

because many root-associated microbes are also sensitive to plant root traits, their responses to

drought may also reflect drought-induced changes in the host phenotype. Both soil- and root-

associated microbiomes under drought exhibit consistent enrichment of gram-positive bacteria and

depletion of gram-negative bacteria [12–14]. Within those categories, however, some bacterial

phyla respond differently to drought in soil versus roots [3]. For example, enrichment of Bacillota

(formerly Firmicutes) and Actinomycetota (formerly Actinobacteria) during drought are less

pronounced in soil than in root-associated microbiomes, suggesting that these phyla respond to

drought-induced changes in root physiology [14–16]. Thus, drought impacts both soil- and root-

associated microbiomes, but in different ways, owing to the influence of the plant phenotype.

Changes in soil- and root-associated microbiomes due to historical precipitation may affect

plant fitness under drought. Drought-induced restructuring of root-associated microbiomes is

hypothesized to be a conserved mechanism [14, 17] that supports survival of both the plant host

and its symbionts [18]. In one study, plant fitness peaked when the soil microbiome's historic

moisture conditions matched contemporary experimental conditions [19]. Additional research

suggests that exposure to short-term drought creates a microbial legacy effect that moderates

drought stress in plants [20, 21] and strengthens positive plant-soil feedbacks, although many such

patterns were species-specific [22]. Thus, how historical precipitation patterns condition soil- and

root-associated microbiomes and their relevance to plant fitness under drought stress is an open

question.

Another open question is how historical sympatry between a plant species and the soil

microbiome influences their interactions during drought. For example, drought-stressed

Arabidopsis thaliana performed better in soil collected from a habitat where the plant had

naturalized, compared to soils from a maize field and a forest stand of *Pinus ponderosa* [23].

Conversely, historical sympatry can be detrimental in agricultural systems: soil-dwelling plant

pathogens tend to accumulate over time in monocultures [24–26], leading to negative plant-soil

feedbacks [27, 28] that drive lower plant performance in sympatric soils. Agricultural practices

such as tillage and irrigation, however, may obscure some of these patterns as has been observed

previously [29–31]. Thus, we designed an experiment to assess plant fitness under a contemporary

drought when grown in sympatric and non-sympatric conditioned soils from different land use

types and historic precipitation regimes.

Our experiment tested how soil microbiomes with different precipitation and land use

legacies impacted *Zea mays* L. (maize) under drought stress. We collected six soils across an east-

west precipitation gradient in Kansas (Figure 1), representing two land use regimes with differing

degrees of sympatry with maize: native prairies and agricultural fields. We inoculated maize

seedlings with these soil communities and measured plant growth during our experimental

drought. Specifically, we: **1**) characterized plant phenotype and microbiome responses to drought,

2) tested how soil microbiome histories (precipitation and land use type) affected maize growth,

and **3**) identified microbial taxa that were associated with plant growth. A more nuanced

understanding of drought-induced phenotypic and root microbiome changes will help stimulate

advancements towards alleviating crop losses due to drought.

Material and Methods

Soil inoculum sources and preparation

Soil samples were collected in May 2019 from 4 prairie sites and 2 agricultural sites across

a steep precipitation gradient in Kansas, USA (Figure 1). From each site, ~60 cm deep samples

were collected from each corner and the center. The first 2-3 cm of topsoil were removed. Soil

samples were homogenized, sieved, and stored in plastic bags at 4ºC until use (~4 months after

collection).

Inocula were prepared by mixing 20 g of each soil into 100 mL 1x phosphate-buffered

saline (PBS) and 0.0001% Triton X-100. After settling, the mixtures were filtered through

Miracloth (22-25 µm pore size; Calbiochem, San Diego, CA, USA) and then centrifuged for 30

minutes (3000 x g). After discarding the supernatant, the microbial pellets were resuspended in 20

mL 1x PBS; these slurries were kept shaking for aeration until use, later that day. Each final

inoculum consisted of 10 mL soil slurry in 1 L 0.5X Murashige and Skoog basal salt mixture (MS;

Sigma-Aldrich, Darmstadt, Germany). A control treatment was made by adding 10 mL sterile 1x

PBS into 1 L 0.5X MS.

Planting, inoculation, and greenhouse setup

To increase the generalizability of our results, two genotypes of *Zea mays* L. (maize) were

used: B73 and Mo17 [32]. In a laminar flow hood, seeds were surface sterilized by submerging in

70% ethanol, then 5% NaClO, followed by three rinses with sterile water. To avoid confounding

effects of soil physicochemical properties, autoclaved calcined clay ("Pro's Choice Rapid Dry";

Oil-Dri Corporation, Chicago, IL) was used as the growth medium. Seeds were planted one inch

deep in 107 mL "Cone-tainers" (Stuewe & Sons Inc., Tangent, OR, USA). Cone-tainers were

assigned randomly to treatments, inoculated with 25 mL of one of the six soil inocula (or the sterile

control), and placed two inches apart in randomized complete blocks (Figure S1A) containing the

24 treatment combinations and one of four controls. Plants were arranged under growth lights (250

W Sun Systems Compact Cool fluorescent lamps) to supplement natural light for 13 hours/day

during the experiment (January-March 2020).

Monitoring and harvesting plants

All plants received 30 mL of reverse osmosis water twice weekly until emergence. After

emergence, "well-watered" plants continued twice-weekly watering while "droughted" plants

were watered every 8-12 days. Plant height was measured weekly to the nearest 0.25 cm on the

tallest fully expanded leaf held parallel to the stem. Plants were sequentially harvested to measure

biomass at one of three time points during the experiment: 100 plants on day 24 and day 39, and

the remainder (n=344) on day 50 (Figure S1B). Calcined clay was removed from roots by hand.

Shoot and root fractions were separated and oven-dried at 65 ℃ to a constant weight. Before

drying, one nodal root (from the uppermost soil-borne root whorl) per plant was collected and

stored at -80 ℃ until DNA extraction.

Phenotypic analysis

All statistical analysis for phenotypic and microbiome data (see below) was conducted in

R v3.6 [33]. Differences among group means were assessed using analysis of variance (ANOVA)

with Type III sums of squares in the car v3.0.13 [34] or lmertest v3.1.3 packages [35] with post-

hoc comparisons made using the emmeans package v1.7.4.1 [36]. Assumptions of all models were

checked by inspecting model residuals [37]. Dry shoot and root mass were converted to

accumulation rates (g/day) to account for the variable number of growing days plants experienced

post-emergence. The lme4 package v1.1.29 [38] was used to fit a generalized linear mixed model

for emergence (binary) and linear mixed models for other growth responses (continuous). Models

used the following formula (*growth response~ Inoculum × Treatment × Genotype*) with all

possible interactions included. Models were also fit using '*Longitude*' instead of '*Inoculum*' to test

the linear relationship between plant growth and the longitude of the soil microbiome origin sites

(a proxy for mean annual precipitation; Figure 1). Experimental block was treated as a random

effect with time point included as an additional random effect in the root/shoot ratio model.

Root/shoot ratio was log-transformed and both shoot and root mass rate were square-root-

transformed to meet the assumptions of ANOVA.

DNA extraction and library preparation

Nodal roots were freeze-dried for 48 hours (FreeZone lyophilizer; Labconco, Kansas City,

MO, USA) and ground to powder with a HT Lysing Homogenizer (OHAUS, Parsippany, NJ,

USA). Each ground root sample was mixed into 800 µL lysis buffer (1M Tris pH=8.0; 0.5 mM

NaCl; 10 mM EDTA) and transferred to 96-well plates containing 500 µL 1 mm garnet rocks per

well (Biospec Products, Bartlesville, OK, USA), after which DNA purification proceeded via

bead-beating and chemical lysis as previously described [39]. DNA from soil and freshly generated

inocula samples was extracted in December 2022 using the same protocol but omitting the

lyophilization step and placing 250 mg soil or 800 µL inoculum directly into a garnet plate.

Amplicon sequencing libraries were prepared using established protocols [39]. Briefly, we

PCR-amplified the V4 region of the 16S rRNA gene and ITS1 region for bacteria and fungi,

respectively. Amplicons were sequenced (2x250 bp) on a NovaSeq 6000 SP flow cell (Illumina,

San Diego, CA, USA). Each "batch" of 96 samples remained together throughout DNA extraction,

amplification, and sequencing.

Bioinformatic processing

Sequence processing and quality filtering was conducted using an established workflow

[39]. Briefly, Cutadapt v2.3 [40] was used to remove primers and demultiplex sequences into 16S

and ITS sequence files. DADA2 v1.14.1 [41] was used to denoise reads, merge paired-end

sequences, dereplicate, and remove chimeric sequences. Taxonomy was assigned to the resulting

amplicon sequence variants (ASVs) by a naïve Bayesian classifier [42] trained on the 11.5 release

of the Ribosomal Database Project [43] for bacteria and the 01.12.2017 release of the UNITE

database [44] for fungi. Phyloseq v1.40.0 [45] was used to remove host contamination and ASVs

unassigned to a kingdom. Chao1 richness and diversity indices (Shannon and Inverse Simpson)

were calculated before the removal of ASVs not observed $\geq$25 times in $\geq$5 samples and samples

with <400 reads. Finally, ASV counts were centered-log-ratio (CLR) transformed using the

ALDEx2 package v1.28.1 [46, 47].

Microbiome Analysis

To test the similarity of inocula to their soils of origin, a linear model was fit for each alpha

diversity metric with collection site, substrate (soil *vs.* inoculum), and sequencing depth (Z-

transformed) as main effects. Community composition was assessed by permutational multivariate

analysis of variance (PERMANOVA) and constrained partial redundancy analysis (RDA). For

PERMANOVA, the *adonis2()* function from the vegan package v2.6.2 [48] was used to model the

matrix of between-sample Aitchison distances [47] with collection site and substrate as main

effects. The RDA was fit with the same model, removing variance attributed to log(Sequencing

Depth), and visualized using an ordination plot.

To investigate root microbiome responses, a model was fit for each alpha diversity metric

(*Metric ~ Soil Inoculum × Drought Treatment + Genotype*), with *Block* and *Batch* as random

effects. Genotype was treated as a fixed effect without interactions to generalize results across

genotypes and increase power to detect other main effects. Each phylum's CLR-transformed

abundance was modeled to test for responses to experimental factors. Across all models and where

appropriate, p-values were adjusted to correct for multiple comparisons using a Benjamini-

Hochberg correction [49]. To explore changes in community composition, a PERMANOVA

model was fit (*Aitchison matrix ~ Soil Inoculum × Drought Treatment + Genotype*) with

permutations constrained by *Batch.* Main effects were tested for homogeneity of dispersions using

the vegan package. Pairwise dispersions among factor levels were assessed using a permutation

test with 999 permutations constrained by *Batch.* Constrained partial redundancy analysis of the

Aitchison distance matrix was conducted to remove variance attributed to *Batch* and

log(Sequencing Depth) and to visualize the variance in microbial composition explained by

experimental factors.

The machine-learning package caret v6.0.92 [50] with ranger's v0.14.1 [51]

implementation of the random forest algorithm was used to assess how accurately roots could be

classified as droughted or well-watered based on microbiome community data (bacterial and

fungal, separately). Samples were randomly split into a training set (80%) and a test set (20%),

with the test set withheld from the model, and used to determine the final model's performance.

Hyperparameters were optimized using a grid search over minimum node size (1, 5, 10) and the

number of features available at each node (10-100% of the ASVs). For each combination in the

grid, classifier accuracy was used to assess performance on out-of-bag samples with 10-fold cross

validation. Final predictions were made with the trained model on the withheld test set and

visualized using confusion matrices. The contribution of each ASV to classification accuracy was

assessed using permuted importance: mean decrease in accuracy (MDA) values were calculated

for each ASV across the 10-fold cross validation. The benefits of this strategy are twofold, we can

assess if drought induced a predictable shift in the microbiome as a whole and identify individual

ASVs with the greatest differential abundance as a result of drought.

Connecting plant growth responses to microbial abundances

Linear models were used to test associations between taxon abundances and plant growth.

To focus on the phenotypic variation that could not be explained by Genotype, Inoculum, Block

and Batch effects, we extracted residuals from a model that contained these terms and used them

as the response variable for a second model (*Growth Response$_{residual}$ ~ Taxon abundance ×*

Drought Treatment). *Taxon abundance* described the association of the taxon (CLR-transformed

counts of bacterial families and fungal orders) with plant growth rate; the interaction term

described whether the taxon's effect on plant growth differed under droughted vs. well-watered

conditions.

Results

Soil and inocula community composition differs across collection sites

We assessed the microbial community composition of soils and their derived inocula from

the six collection sites (n = 8 per site). Bacterial and fungal alpha diversity of soils differed across

collection sites (Tables S2 and S3); bacterial diversity was highest in the westernmost prairie soil

whereas fungal diversity was highest in the easternmost prairie soil (Figures S2 and S3).

Additionally, community composition of both bacterial and fungal microbiomes differed among

sites (Table S3; R^2=0.22-0.23; *p*-value<0.01). Relative abundances of bacterial phyla and fungal

classes further showed site-specific patterns (Figures S4 and S5), however, few were correlated

with longitude.

Relative to the original soils, the inocula had similar bacterial alpha diversity but lower

fungal richness (Figure S3C-D; Tables S2 and S3). The composition of both bacterial and fungal

communities also shifted between soils and their derived inocula (Table S4); however, both soil

and inocula samples clustered in ordination space by collection site (Figure 2). Thus, the original

soil microbial communities were largely retained in the derived inocula.

Plant responses to soil microbiomes vary across genotype and drought treatment

We found no evidence that soil inocula impacted emergence rates (*p*-value = 0.37,

Pearson's χ^2 test; Table S1), however, maize genotype B73 (82.5%) emerged more successfully

than Mo17 (62.0%; *p*-value=0.05, Pearson's χ^2 test). Post-emergence, both drought treatment and

maize genotype affected plant growth responses (Table 1). Drought decreased both shoot and root

mass accumulation rates (Figure 3A-B; p-value=<0.01) and increased root/shoot ratio (Figure 3C;

p-value<0.01). These traits also differed between genotypes, with Mo17 investing more into root

growth than B73 (Figure S6; p-value=<0.01). Shoot mass rate was the only trait that responded to

soil inoculum (Table 1; p-value=<0.01); this effect was driven by the 'TLI$_P$' inoculum, which

slowed the accumulation of root mass.

We observed no evidence of interactions between drought treatment and soil inoculum on

plant growth responses. However, inoculum interacted with genotype to influence root/shoot ratio

(Table 1; p-value=0.02). Mo17 displayed greater root/shoot ratios than B73 across all inocula,

although the magnitude of this difference varied (Δ_{emmean} 0.176-0.493; p-value=0.02-<0.01). B73

accumulated shoot mass more quickly than Mo17 when inoculated with three of the four prairie

soil microbiomes ('SVR$_P$' p-value=<0.01, 'HAY$_P$' p-value=<0.01, 'TLI$_P$' p-value=0.03, 'KNZ$_P$'

p-value=0.27) but not with agricultural soil microbiomes ('SVR$_{Ag}$' p-value=0.33, 'TLI$_{Ag}$' p-

value=0.12). Thus, the effect of soil microbial communities on maize growth generally depended

on plant genotype, but not on the drought treatment.

Root microbiomes and community responses to drought modulated by soil inoculum

Drought treatment and soil inocula collection site impacted the richness and diversity of

root-associated bacterial and fungal communities (Tables S5 and S6). Drought increased Chao1

richness of bacterial communities (Tables S5 and S6; p-value=0.01) but affected neither Shannon

nor Inverse Simpson diversity. In contrast, Shannon and Inverse Simpson fungal diversity

increased under drought conditions, particularly for plants that received the 'TLI$_P$' inoculum

(Figure 1A; Table S6; p-value=0.01–0.02).

Bacterial community composition depended on the soil inoculum (Table 2; R^2=0.05, p-

value=<0.01, PERMANOVA) and drought treatment (R^2=0.01, p-value=<0.01, PERMANOVA).

We found no evidence that bacterial community dispersion differed among treatments, genotypes,

or inocula (p-value=>0.05, Multivariate Levene's test). Redundancy analysis aligned with

PERMANOVA results: soil inoculum and drought treatment explained the largest proportions of

variance (6.09% and 1.57%; Figure S7 and 4B, respectively), while genotype explained >1% of

variance (Figure S8). We attributed 7.1% of the variance in Deinococcota (formerly Deinococcus-

Thermus) to the interaction between soil inoculum and drought (q=0.03), with enrichment

observed in droughted plants (39.6% increase in relative abundance). In addition, 3.7% of the

variance in Pseudomonadota was attributable to our drought treatment (q=0.04); relative

abundance of this phylum decreased 4% in droughted plants.

In contrast, drought treatment did not alter relative abundances of any fungal classes, nor

overall root fungal community profiles (Figure 5A). Variance in fungal community composition

was mostly due to initial soil inoculum source (Figure S9; Table 3; R^2=0.09, p-value=<0.01). As

observed for bacterial communities, neither drought treatment, maize genotype, nor soil inoculum

source affected fungal community dispersion (p-value=>0.05, Multivariate Levene's test).

Drought and maize genotype each explained <1% of the variance in fungal community

composition (Figure 5B and S10, respectively). Thus, while root-associated bacterial community

composition responded weakly to drought treatment, we observed no trend for fungal community

composition.

Identification of bacterial and fungal ASVs responding to drought

Our random forest models showed greater accuracy in predicting the drought treatment of

a given sample when trained on bacterial vs fungal communities (Figure 4C&5C). Two bacterial

ASVs from the genus *Sphingomonas* (Pseudomonadota) were the top two features contributing to

prediction accuracy of the random forest classifier when attempting to predict drought treatment,

with a mean decrease in accuracy (MDA) of 2.88% and 0.59% when withheld from the model

(Figure 4C and S11-12). In other words, these two bacterial ASVs contributed to ~3.5% increase

of model accuracy in predicting drought and were positively correlated with one another (*p*-

value=<0.01; r=0.521 Pearson test). These *Sphingomonas* ASVs were depleted in droughted plants

relative to well-watered plants (10.8% decrease in relative abundance). In contrast, we did not

detect any fungal ASVs that were differentially abundant in response to drought treatment, nor any

that contributed to random forest classifier accuracy with MDA value >0.5% (Figure 5C and S13-

14).

Connecting plant growth responses to bacterial and fungal communities

Soil inocula derived from collection sites across Kansas, spanning an east-west

precipitation gradient had only a small impact on plant growth responses (Table 1; Figure 6A and

S15). These patterns were not correlated with the longitude of soil inocula collection sites (a proxy

for precipitation) when including both agricultural sites and native prairies (Figure S16; *p*-

value=0.08-0.70 and 0.17-0.93; ANOVA for *'Longitude'* and *'Longitude × Treatment'*,

respectively). However, when restricting the analysis to prairie soil communities, we detected an

interaction between longitude and drought treatment (Figure 6B; *p*-value=0.03). The shoot mass

rate of well-watered plants was negatively associated with longitude (slope=-3.29e^{-4}

g/day/°longitude) whereas droughted plants displayed a weak positive association (slope=3.04e^{-}

^{5}). Put another way, plants that experienced drought benefited from microbiomes from historically

wetter soils, while the opposite was true for well-watered plants in the experiment. Notably, B73

and Mo17 exhibited contrasting associations with longitude and soils across the Kansas

precipitation gradient (Figure 6C; p-value=0.03; Slope=$-3.07e^{-4}$ and $9.12e^{-6}$, respectively).

We identified bacterial and fungal taxa correlated with promotion or repression of plant

growth. Overall, 13 bacterial families showed a significant positive (6/13) or negative (7/13)

growth promotion effect on shoot mass (Table 4). These represented only a few phyla (6

Actinomycetota, 5 Pseudomonadota, and 2 Bacillota) and were predominantly gram positive

(8/13). Two of the identified families (Corynebacteriaceae and Streptococcaceae) interacted with

drought treatment, indicating that their effect on shoot mass rate was modulated by the amount of

water the plant received. The abundances of Corynebacteriaceae and Streptococcaceae were

negatively correlated with shoot mass rate in well-watered plants but not in droughted plants

(Figure S17). Using the same analysis framework, root mass rate was not associated with any

bacterial families, possibly because root mass rate variance decreased over time as plant roots

expanded to fill the entire container (Figure S18). In addition, bacterial families that correlated

with root/shoot ratio were simply a subset of the bacterial families identified for shoot mass rate.

For fungi, only shoot mass rate showed significant correlations with the abundance of fungal orders

(Table 5). All three significant fungal orders (Hypocreales, Malasseziales, and Sporidiobolales)

were positively correlated with shoot growth regardless of the water regime applied.

Discussion

Understanding plant-microbiome interactions under drought stress is critical to safeguard

agricultural systems in the face of global climate change. In this study, we measured the growth of

maize seedlings under experimental drought after inoculation with soil microbiomes from across

a natural precipitation gradient in Kansas. Under drought stress, we observed phenotypic

differences in maize resource allocation and the restructuring of bacterial, but not fungal, root

microbiome communities. Following the stress gradient hypothesis [52] we expected drought-

stressed maize seedlings to perform better in the presence of microorganisms from historically dry

soils, where chronic water stress would have fostered positive interactions with neighboring plants.

We detected an effect of historical precipitation exposure on soil microbiomes' interactions with

plant hosts during drought treatment, but only for the prairie soils. Notably, the direction of this

interaction was opposite of our prediction: prairie soils from historically wetter locations increased

maize biomass under experimental drought more than soils from historically dry locations. Deeper

analysis revealed several bacterial and fungal taxa that were associated with plant growth

promotion or repression and that were modulated by the experimental drought.

Plant and microbiome responses to drought

In our experiment, maize displayed strong phenotypic tradeoffs in response to drought.

Drought-treated plants increased root growth relative to shoot growth, resulting in shorter plants

over the course of the experiment. This supports our understanding of plant responses to drought,

as plants prioritize root over shoot growth under drought stress [53, 54], favoring deeper and often

denser root systems in order to obtain sufficient water [55–57]. These phenotypic changes in root

growth, along with direct effects of water limitation in the soil, contributed to the responses of the

root microbiomes in our study.

Drought altered bacterial communities of maize roots, although the magnitude of

compositional change within our study was small ($R^2=0.01$) relative to other studies ($R^2=0.04$-

0.10) [14–16, 18]. Similar to previous studies [17], we observed a depletion of gram-negative and

an enrichment for gram-positive bacteria in response to drought, likely relating to the desiccation

resistance provided by the single thick peptidoglycan cell wall of gram-positive bacteria [58].

Enrichment of Actinomycetota during drought tends to be more pronounced in root-associated

microbiomes than in soil, suggesting that shifts in plant root physiology also play a role [3, 17];

however, the root traits driving this effect are unknown. During drought, root exudate quantity is

generally maintained or increased [59–61], but the stronger changes in exudate composition [62–

64] may contribute to the root microbiome drought response [65]. In *Sorghum*, for instance,

drought-induced changes in root exudate composition and iron homeostasis contribute to

Actinomycetota enrichment in the rhizosphere [18, 66].

Fungal communities of maize roots were less sensitive to drought relative to their bacterial

neighbors. While drought increased the diversity of the plants inoculated with the TLI prairie

fungal community, it did not significantly affect overall fungal community composition. Machine

learning classification by drought treatment further illustrated this difference: when trained on

bacteria, we achieved 72.5% balanced accuracy, while for fungi, we achieved only 50%. This

cross-kingdom difference agrees with previous reports, as both free living and root-associated

fungal communities tend to tolerate desiccation [67, 68]. The observed difference in survival

between bacteria and fungi under drought may relate to both the vegetative and dormant growth

forms of fungi: hyphae comprise most vegetative fungal growth and can cross air-gaps within dry

soil in search of resources [58, 69], and dormant spores are generally desiccation-resistant [70].

Historical precipitation, soil community legacies, and their impact on plant drought tolerance

Our hypothesis that historical precipitation patterns would impart legacy effects on soil

microbiomes that influence plant growth under drought stress was partially supported. Bacterial

and fungal compositional differences were apparent among the soils and inocula from six sites

across a steep precipitation gradient in Kansas (Figure 1&2). In the four soils from prairie sites,

we found a significant interaction for shoot mass rate between longitude of soil inoculum source

and drought treatment. The fact that the two agricultural soils weakened this pattern provides

support for our hypothesis: the periodic irrigation of the agricultural sites may have obscured the

legacy effects of historic differences in annual rainfall across the gradient. Alternatively (or in

addition), intensive agricultural practices such as tillage and fertilization may have altered these

soil microbiomes in a way that disrupted their responses to drought and/or their interactions with

the host plant.

Interestingly, we found plants performed better when the historic conditions were opposite

the contemporary conditions. The reason for this surprising result is not immediately clear, but

there are several possible explanations. Contemporary well-watered conditions may have relaxed

the fitness consequences of expression of microbial traits that are critical under drier environments.

For instance, microorganisms counteract the physiologic stress of drought by accumulating

solutes, which reduces their internal water potential as a mechanism to avoid desiccation [71]. The

accumulation of solutes requires considerable energy expenditure and constitutes a large sink of C

and N resources [72–74]. The contemporary well-watered conditions relax these costly solute

needs, perhaps allowing resources to be reallocated to functions that are beneficial to plants.

Another possibility is that soils from historically wetter sites generally possess greater microbial

biomass and diversity [75–77], which is predicted to promote positive interactions and improved

community function [78]. A third potential explanation involves the balance of beneficial and

pathogenic microbes in the soil community. Pathogen richness in soil, root, and shoot microbiomes

is often positively associated with mean annual temperature and precipitation [79–81]. Thus, soil

communities from historically dry sites may have relatively fewer plant pathogens, resulting in

greater plant performance under contemporary well-watered conditions. We cannot distinguish

between these possible explanations using the data in hand; however, a future experiment that

leverages rain exclusion structures could effectively explore changes in the balance of beneficial,

neutral, and pathogenic microbes.

The degree of sympatry between the plant species and the soil microbiome may affect

performance under drought conditions for the host plant [20, 23]. Positive plant-soil feedbacks are

generally expected to strengthen under drought conditions [82, 83], but this effect can vary

strongly by plant species and functional group [84, 85]. Irrigation of agricultural sites would have

weakened historical drought stress, and in turn reduced positive interactions [52]. Furthermore,

repeated growth of a crop in one location, even in rotation with another species, frequently results

in negative feedbacks driven by soil microbial communities [86–88]. Therefore, we expected that

maize inoculated with agricultural soil microbiomes (*i.e.,* from grain fields where maize has long

been a dominant plant species) would perform worse under drought than plants inoculated with

prairie soils that were naïve to maize. Consistent with our expectation, inoculation with prairie-

derived communities increased shoot mass rate and root/shoot ratio in the maize genotype B73,

relative to farm-derived communities. This finding agrees with past work showing that frequent

cultivation of maize in monoculture can lead to a buildup of pathogenic microbes in the soil and

stronger negative plant-soil feedbacks [24–26, 89]. The prairie soils, naïve to maize, would have

had no such buildup. Our results support the need for additional research to tease apart the ability

of dominant plant species to condition soil microbiomes in either a beneficial or detrimental

manner, and to characterize the environmental dependency of such conditioning.

Microbial taxa associated with plant growth

Although only a small portion of plant growth variance was explained by soil inoculum

source and historical precipitation, we identified multiple bacterial and fungal taxa that correlated

with plant growth responses to drought. Two bacterial families, Corynebacteriaceae and

Streptococcaceae, repressed growth of well-watered but not droughted plants. Corynebacteriaceae

has been recovered from roots of multiple plant species [90, 91] and been identified as a keystone

species of the endosphere in two Antarctic endemic plants [92]. This family contains multiple

phytopathogenic species [93] with some strains specific to maize [94]. Additionally, during a

multi-generation experimental evolution study, Corynebacteriaceae (among other bacterial

families) was consistently associated with delayed flowering in multiple plant species [95]. Less

is known about the relevance of Streptococcaceae to plant growth. In *Zingiber officinale*, this

family was significantly enriched in rhizomes with advanced stages of soft rot [96]. However,

Streptococcaceae is prevalent in soil and plants [97, 98], typically at higher abundance in aerial

plant compartments than in the rhizosphere [99, 100]. Both of these families represent targets for

direct manipulation and *in-planta* testing to better understand the mechanism responsible for the

growth repression we observed under well-watered conditions.

Acknowledgements

We thank members of the Wagner lab for helpful comments in preparing this article. In addition,

we thank Zachary Harris and Nichole Ginnan for helpful discussions. This work was funded by

the United States Department of Agriculture National Institute of Food and Agriculture (USDA-

NIFA) grant #2022-67013-36672 to MK and MRW, National Science Foundation grant #IOS-

2016351 to MRW, Novo Nordisk Foundation INTERACT project under award number

NNF19SA0059360 to MK, and seed funding from the NSF under award No. OIA-1656006 to Dr.

Kristin Bowman-James and matching support from the State of Kansas through the Kansas Board

of Regents.

Competing Interests

The authors declare that the research was conducted in the absence of any commercial or financial

relationships that could be construed as a potential conflict of interest. Any opinion, findings, and

conclusions or recommendations expressed in this material are those of the authors and do not

necessarily reflect the views of the United States Department of Agriculture or the National

Science Foundation.

Data Availability Statement

Raw sequencing data is available on NCBI SRA under BioProject ID PRJNA913622 and SRA

accessions SAMN32302538-SAMN32303072 for root-associated samples and SAMN33863891-

SAMN33864082 for soil and inoculum samples. Code to reproduce the analysis and figures are

available on GitHub at https://github.com/Kenizzer/Maize_Drought_Microbiome.

Author Contributions

BAS and PMH collected soil samples across Kansas. MRK, MK, and MRW designed the

experiment and collection methods. MRK performed sample collection and processing. AC and

NEF performed DNA extractions and library preparations. NEF and JFS performed the data

analysis. JFS wrote the first draft of the manuscript. All authors contributed to data interpretation,

the writing and reviewing of the manuscript, and approved the final draft.

References

1. Balting DF, AghaKouchak A, Lohmann G, Ionita M. Northern Hemisphere drought risk in a warming climate. *npj Clim Atmos Sci* 2021; **4**: 61.

2. Chiang F, Mazdiyasni O, AghaKouchak A. Evidence of anthropogenic impacts on global drought frequency, duration, and intensity. *Nat Commun* 2021; **12**: 2754.

3. Naylor D, Coleman-Derr D. Drought Stress and Root-Associated Bacterial Communities. *Front Plant Sci* 2018; **8**: 1–16.

4. Korell L, Auge H, Chase JM, Harpole WS, Knight TM. Responses of plant diversity to precipitation change are strongest at local spatial scales and in drylands. *Nat Commun* 2021; **12**: 2489.

5. Field R, Hawkins BA, Cornell H V., Currie DJ, Diniz-Filho JAF, Guégan J-F, et al. Spatial species-richness gradients across scales: a meta-analysis. *J Biogeogr* 2009; **36**: 132–147.

6. Bachar A, Al-Ashhab A, Soares MIM, Sklarz MY, Angel R, Ungar ED, et al. Soil Microbial Abundance and Diversity Along a Low Precipitation Gradient. *Microb Ecol* 2010; **60**: 453–461.

7. Hawkes C V., Kivlin SN, Rocca JD, Huguet V, Thomsen MA, Suttle KB. Fungal community responses to precipitation. *Glob Chang Biol* 2011; **17**: 1637–1645.

8. Hawkes C V., Keitt TH. Resilience vs. historical contingency in microbial responses to environmental change. *Ecol Lett* 2015; **18**: 612–625.

9. Averill C, Waring BG, Hawkes C V. Historical precipitation predictably alters the shape and magnitude of microbial functional response to soil moisture. *Glob Chang Biol* 2016; **22**: 1957–1964.

10. Canarini A, Schmidt H, Fuchslueger L, Martin V, Herbold CW, Zezula D, et al. Ecological memory of recurrent drought modifies soil processes via changes in soil microbial community. *Nat Commun* 2021; **12**: 5308.

11. Zarraonaindia I, Owens SM, Weisenhorn P, West K, Hampton-Marcell J, Lax S, et al. The Soil Microbiome Influences Grapevine-Associated Microbiota. *MBio* 2015; **6**: e02527-14.

12. Acosta-Martínez V, Cotton J, Gardner T, Moore-Kucera J, Zak J, Wester D, et al. Predominant bacterial and fungal assemblages in agricultural soils during a record drought/heat wave and linkages to enzyme activities of biogeochemical cycling. *Appl Soil Ecol* 2014; **84**: 69–82.

13. Fuchslueger L, Bahn M, Hasibeder R, Kienzl S, Fritz K, Schmitt M, et al. Drought history affects grassland plant and microbial carbon turnover during and after a subsequent drought event. *J Ecol* 2016; **104**: 1453–1465.

497   14.   Naylor D, DeGraaf S, Purdom E, Coleman-Derr D. Drought and host selection influence
498   bacterial community dynamics in the grass root microbiome. *ISME J* 2017; **11**: 2691–2704.

499   15.   Santos-Medellín C, Edwards J, Liechty Z, Nguyen B, Sundaresan V. Drought Stress Results
500   in a Compartment-Specific Restructuring of the Rice Root-Associated Microbiomes. *MBio*
501   2017; **8**: 1–15.

502   16.   Fitzpatrick CR, Copeland J, Wang PW, Guttman DS, Kotanen PM, Johnson MTJ. Assembly
503   and ecological function of the root microbiome across angiosperm plant species. *Proc Natl*
504   *Acad Sci* 2018; **115**: E1157–E1165.

505   17.   Xu L, Coleman-Derr D. Causes and consequences of a conserved bacterial root microbiome
506   response to drought stress. *Curr Opin Microbiol* 2019; **49**: 1–6.

507   18.   Xu L, Naylor D, Dong Z, Simmons T, Pierroz G, Hixson KK, et al. Drought delays
508   development of the sorghum root microbiome and enriches for monoderm bacteria. *Proc*
509   *Natl Acad Sci* 2018; **115**: E4284–E4293.

510   19.   Lau JA, Lennon JT. Rapid responses of soil microorganisms improve plant fitness in novel
511   environments. *Proc Natl Acad Sci* 2012; **109**: 14058–14062.

512   20.   Kaisermann A, de Vries FT, Griffiths RI, Bardgett RD. Legacy effects of drought on plant–
513   soil feedbacks and plant–plant interactions. *New Phytol* 2017; **215**: 1413–1424.

514   21.   De Long JR, Semchenko M, Pritchard WJ, Cordero I, Fry EL, Jackson BG, et al. Drought
515   soil legacy overrides maternal effects on plant growth. *Funct Ecol* 2019; **33**: 1400–1410.

516   22.   Buchenau N, van Kleunen M, Wilschut RA. Direct and legacy-mediated drought effects on
517   plant performance are species-specific and depend on soil community composition. *Oikos*
518   2022; **2022**: 1–14.

519   23.   Zolla G, Badri D V., Bakker MG, Manter DK, Vivanco JM. Soil microbiomes vary in their
520   ability to confer drought tolerance to Arabidopsis. *Appl Soil Ecol* 2013; **68**: 1–9.

521   24.   Strom N, Hu W, Haarith D, Chen S, Bushley K. Interactions between soil properties, fungal
522   communities, the soybean cyst nematode, and crop yield under continuous corn and soybean
523   monoculture. *Appl Soil Ecol* 2020; **147**: 103388.

524   25.   Sumner DR, Gascho GJ, Johnson AW, Hook JE, Threadgill ED. Root Diseases, Populations
525   of Soil Fungi, and Yield Decline in Continuous Double-Crop Corn. *Plant Dis* 1990; **74**:
526   704.

527   26.   Wachowska U, Rychcik B. Plants control the structure of mycorrhizal and pathogenic
528   fungal communities in soil in a 50-year maize monoculture experiment. *Plant Soil* 2022.

529   27.   Cesarano G, Zotti M, Antignani V, Marra R, Scala F, Bonanomi G. Soil sickness and
530   negative plant-soil feedback: A reappraisal of hypotheses. *J Plant Pathol* 2017; **99**: 545–
531   570.

28. Jing J, Cong WF, Bezemer TM. Legacies at work: plant–soil–microbiome interactions underpinning agricultural sustainability. *Trends Plant Sci* 2022; **27**: 781–792.

29. Delavaux CS, Schemanski JL, House GL, Tipton AG, Sikes B, Bever JD. Root pathogen diversity and composition varies with climate in undisturbed grasslands, but less so in anthropogenically disturbed grasslands. *ISME J* 2021; **15**: 304–317.

30. Putten WH, Bradford MA, Pernilla Brinkman E, Voorde TFJ, Veen GF. Where, when and how plant–soil feedback matters in a changing world. *Funct Ecol* 2016; **30**: 1109–1121.

31. Raaijmakers JM, Paulitz TC, Steinberg C, Alabouvette C, Moënne-Loccoz Y. The rhizosphere: a playground and battlefield for soilborne pathogens and beneficial microorganisms. *Plant Soil* 2009; **321**: 341–361.

32. Schultz CR, Brantley KM, Wallace JG. The role of genetic variation in Zea mays response to beneficial endophytes. *Plant Growth Regul* 2022; **98**: 167–177.

33. R Core Team. R: A Language and Environment for Statistical Computing. 2021. Vienna, Austria.

34. Fox J, Weisberg S. An R Companion to Applied Regression, 3rd ed. 2019. Sage, Thousands Oaks, CA.

35. Kuznetsova A, Brockhoff PB, Christensen RHB. lmerTest Package: Tests in Linear Mixed Effects Models. *J Stat Softw* 2017; **82**.

36. Lenth R, Love J, Herve M. emmeans: Estimated Marginal Means, aka Least-Squares Means. 2020. CRAN. , https://cran.r-project.org/package=emmeans

37. Harrison XA, Donaldson L, Correa-Cano ME, Evans J, Fisher DN, Goodwin CED, et al. A brief introduction to mixed effects modelling and multi-model inference in ecology. *PeerJ* 2018; **6**: e4794.

38. Bates D, Mächler M, Bolker B, Walker S. Fitting Linear Mixed-Effects Models Using lme4. *J Stat Softw* 2015; **67**.

39. Wagner MR, Roberts JH, Balint-Kurti P, Holland JB. Heterosis of leaf and rhizosphere microbiomes in field-grown maize. *New Phytol* 2020; **228**: 1055–1069.

40. Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.journal* 2011; **17**: 10.

41. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2: High-resolution sample inference from Illumina amplicon data. *Nat Methods* 2016; **13**: 581–583.

42. Wang Q, Garrity GM, Tiedje JM, Cole JR. Naïve Bayesian Classifier for Rapid Assignment of rRNA Sequences into the New Bacterial Taxonomy. *Appl Environ Microbiol* 2007; **73**: 5261–5267.

43. Callahan BJ. RDP taxonomic training data formatted for DADA2 (RDP trainset 16/release 11.5). 2017. Zenodo.

44. UNITE Community. UNITE general FASTA release. 2017.

45. McMurdie PJ, Holmes S. phyloseq: An R Package for Reproducible Interactive Analysis and Graphics of Microbiome Census Data. *PLoS One* 2013; **8**: e61217.

46. Fernandes AD, Reid JNS, Macklaim JM, McMurrough TA, Edgell DR, Gloor GB. Unifying the analysis of high-throughput sequencing datasets: characterizing RNA-seq, 16S rRNA gene sequencing and selective growth experiments by compositional data analysis. *Microbiome* 2014; **2**: 15.

47. Gloor GB, Macklaim JM, Pawlowsky-Glahn V, Egozcue JJ. Microbiome Datasets Are Compositional: And This Is Not Optional. *Front Microbiol* 2017; **8**: 1–6.

48. Oksanen J, Guillaume Blanchet F, Friendly M, Kindt R, Legendre P, McGlinn D, et al. vegan: Community Ecology Package. 2019.

49. Benjamini Y, Hochberg Y. Controlling the False Discovery Rate : A Practical and Powerful Approach to Multiple Testing. *J R Stat Soc* 1995; **57**: 289–300.

50. Kuhn M. Building Predictive Models in R Using the caret Package. *J Stat Softw* 2008; **28**: 159–160.

51. Wright MN, Ziegler A. ranger : A Fast Implementation of Random Forests for High Dimensional Data in C++ and R. *J Stat Softw* 2017; **77**.

52. Bertness MD, Callaway R. Positive interactions in communities. *Trends Ecol Evol* 1994; **9**: 191–193.

53. Sharp RE, Davies WJ. Solute regulation and growth by roots and shoots of water-stressed maize plants. *Planta* 1979; **147**: 43–49.

54. Lynch JP, Chimungu JG, Brown KM. Root anatomical phenes associated with water acquisition from drying soil: Targets for crop improvement. *J Exp Bot* 2014; **65**: 6155–6166.

55. Gao Y, Lynch JP. Reduced crown root number improves water acquisition under water deficit stress in maize (Zea mays L.). *J Exp Bot* 2016; **67**: 4545–4557.

56. Zhan A, Schneider H, Lynch JP. Reduced Lateral Root Branching Density Improves Drought Tolerance in Maize. *Plant Physiol* 2015; **168**: 1603–1615.

57. Sebastian J, Yee M-C, Goudinho Viana W, Rellán-Álvarez R, Feldman M, Priest HD, et al. Grasses suppress shoot-borne roots to conserve water during drought. *Proc Natl Acad Sci* 2016; **113**: 8861–8866.

58. Manzoni S, Schimel JP, Porporato A. Responses of soil microbial communities to water stress: results from a meta-analysis. *Ecology* 2012; **93**: 930–938.

59. Henry A, Doucette W, Norton J, Bugbee B. Changes in Crested Wheatgrass Root Exudation Caused by Flood, Drought, and Nutrient Stress. *J Environ Qual* 2007; **36**: 904–912.

60. Preece C, Peñuelas J. Rhizodeposition under drought and consequences for soil communities and ecosystem resilience. *Plant Soil* 2016; **409**: 1–17.

61. Karlowsky S, Augusti A, Ingrisch J, Hasibeder R, Lange M, Lavorel S, et al. Land use in mountain grasslands alters drought response and recovery of carbon allocation and plant-microbial interactions. *J Ecol* 2018; **106**: 1230–1243.

62. Svenningsson H, Sundin P, Liljenberg C. Lipids, carbohydrates and amino acids exuded from the axenic roots of rape seedlings exposed to water-deficit stress. *Plant, Cell Environ* 1990; **13**: 155–162.

63. Canarini A, Merchant A, Dijkstra FA. Drought effects on Helianthus annuus and Glycine max metabolites: from phloem to root exudates. *Rhizosphere* 2016; **2**: 85–97.

64. Calvo OC, Franzaring J, Schmid I, Müller M, Brohon N, Fangmeier A. Atmospheric CO 2 enrichment and drought stress modify root exudation of barley. *Glob Chang Biol* 2017; **23**: 1292–1304.

65. Zhalnina K, Louie KB, Hao Z, Mansoori N, da Rocha UN, Shi S, et al. Dynamic root exudate chemistry and microbial substrate preferences drive patterns in rhizosphere microbial community assembly. *Nat Microbiol* 2018; **3**: 470–480.

66. Xu L, Dong Z, Chiniquy D, Pierroz G, Deng S, Gao C, et al. Genome-resolved metagenomics reveals role of iron metabolism in drought-induced rhizosphere microbiome dynamics. *Nat Commun* 2021; **12**: 3209.

67. Furze JR, Martin AR, Nasielski J, Thevathasan N V., Gordon AM, Isaac ME. Resistance and resilience of root fungal communities to water limitation in a temperate agroecosystem. *Ecol Evol* 2017; **7**: 3443–3454.

68. de Vries FT, Liiri ME, Bjørnlund L, Bowker MA, Christensen S, Setälä HM, et al. Land use alters the resistance and resilience of soil food webs to drought. *Nat Clim Chang* 2012; **2**: 276–280.

69. Griffin DM. A comparison of the roles of bacteria and fungi, Volume 1: *Bacteria in Nature* . 1985. Springer US, Boston, MA.

70. Wyatt TT, Wösten HAB, Dijksterhuis J. Fungal Spores for Dispersion in Space and Time. *Advances in Applied Microbiology*. 2013. pp 43–91.

71. Harris RF. Effect of Water Potential on Microbial Growth and Activity. In: Parr JF, Gardner WR, Elliott LF (eds). *Water Potential Relations in Soil Microbiology*, 9th ed. 1981. Soil

634   Science Society of America, Madison W.I., pp 23–95.

635   72.   Schimel J, Balser TC, Wallenstein M. Microbial stress-response physiology and its
636   implications for ecosystem function. *Ecology* 2007; **88**: 1386–1394.

637   73.   Schimel JP, Scott WJ, Killham K. Changes in Cytoplasmic Carbon and Nitrogen Pools in a
638   Soil Bacterium and a Fungus in Response to Salt Stress. *Appl Environ Microbiol* 1989; **55**:
639   1635–1637.

640   74.   Killham K, Firestone MK. Salt stress control of intracellular solutes in streptomycetes
641   indigenous to saline soils. *Appl Environ Microbiol* 1984; **47**: 301–306.

642   75.   Neilson JW, Califf K, Cardona C, Copeland A, van Treuren W, Josephson KL, et al.
643   Significant Impacts of Increasing Aridity on the Arid Soil Microbiome. *mSystems* 2017; **2**.

644   76.   Araya JP, González M, Cardinale M, Schnell S, Stoll A. Microbiome Dynamics Associated
645   With the Atacama Flowering Desert. *Front Microbiol* 2020; **10**: 1–13.

646   77.   Wan X, Chen X, Huang Z, Chen HYH. Global soil microbial biomass decreases with aridity
647   and land-use intensification. *Glob Ecol Biogeogr* 2021; **30**: 1056–1069.

648   78.   Hooper DU, Chapin FS, Ewel JJ, Hector A, Inchausti P, Lavorel S, et al. Effects of
649   biodiversity on ecosystem functioning: A consensus of current knowledge. *Ecol Monogr*
650   2005; **75**: 3–35.

651   79.   Delgado-Baquerizo M, Guerra CA, Cano-Díaz C, Egidi E, Wang JT, Eisenhauer N, et al.
652   The proportion of soil-borne pathogens increases with warming at the global scale. *Nat Clim*
653   *Chang* 2020; **10**: 550–554.

654   80.   Větrovský T, Kohout P, Kopecký M, Machac A, Man M, Bahnmann BD, et al. A meta-
655   analysis of global fungal distribution reveals climate-driven patterns. *Nat Commun* 2019;
656   **10**: 1–9.

657   81.   Chen QL, Hu HW, Yan ZZ, Li CY, Nguyen BAT, Zhu YG, et al. Precipitation increases
658   the abundance of fungal plant pathogens in Eucalyptus phyllosphere. *Environ Microbiol*
659   2021; **23**: 7688–7700.

660   82.   Beals KK, Moore JAM, Kivlin SN, Bayliss SLJ, Lumibao CY, Moorhead LC, et al.
661   Predicting plant-soil feedback in the field: Meta-analysis reveals that competition and
662   environmental stress differentially influence psf. *Front Ecol Evol* 2020; **8**.

663   83.   Veresoglou SD, Li GC, Chen J, Johnson D. Direction of plant–soil feedback determines
664   plant responses to drought. *Glob Chang Biol* 2022; **28**: 3995–3997.

665   84.   Hassan K, Carrillo Y, Nielsen UN. The effect of prolonged drought legacies on plant–soil
666   feedbacks. *J Veg Sci* 2021; **32**: 1–17.

85.   Lozano YM, Aguilar-Trigueros CA, Ospina JM, Rillig MC. Drought legacy effects on root

morphological traits and plant biomass via soil biota feedback. *New Phytol* 2022; **236**: 222–
234.

86.   Bennett AJ, Bending GD, Chandler D, Hilton S, Mills P. Meeting the demand for crop
production: the challenge of yield decline in crops grown in short rotations. *Biol Rev* 2012;
87: 52–71.

87.   Wang G, Bei S, Li J, Bao X, Zhang J, Schultz PA, et al. Soil microbial legacy drives crop
diversity advantage: Linking ecological plant–soil feedback with agricultural intercropping.
J Appl Ecol 2021; **58**: 496–506.

88.   Peralta AL, Sun Y, McDaniel MD, Lennon JT. Crop rotational diversity increases disease
suppressive capacity of soil microbiomes. *Ecosphere* 2018; **9**.

89.   Zhang BQ, Chen WD, Yang XB. Occurrence of Pythium species in long-term maize and
soybean monoculture and maize/soybean rotation. *Mycol Res* 1998; **102**: 1450–1452.

90.   Vik U, Logares R, Blaalid R, Halvorsen R, Carlsen T, Bakke I, et al. Different bacterial
communities in ectomycorrhizae and surrounding soil. *Sci Rep* 2013; **3**: 3471.

91.   Stafford WHL, Baker GC, Brown SA, Burton SG, Cowan DA. Bacterial diversity in the
rhizosphere of Proteaceae species. *Environ Microbiol* 2005; **7**: 1755–1768.

92.   Zhang Q, Acuña JJ, Inostroza NG, Duran P, Mora ML, Sadowsky MJ, et al. Niche
Differentiation in the Composition, Predicted Function, and Co-occurrence Networks in
Bacterial Communities Associated With Antarctic Vascular Plants. *Front Microbiol* 2020;
11: 1–13.

93.   Dye DW, Kemp WJ. A taxonomic study of plant pathogenic Corynebacterium species. *New
Zeal J Agric Res* 1977; **20**: 563–582.

94.   Vidaver AK, Mandel M. Corynebacterium nebraskense, a New, Orange-Pigmented
Phytopathogenic Species. *Int J Syst Bacteriol* 1974; **24**: 482–485.

95.   Panke-Buisse K, Poole AC, Goodrich JK, Ley RE, Kao-Kniffin J. Selection on soil
microbiomes reveals reproducible impacts on plant function. *ISME J* 2015; **9**: 980–989.

96.   Huang K, Sun X, Zou Y, Li H, Xu P, Zhang W, et al. Comparison of the Endophytic
Bacterial Microbiota of Asymptomatic and Symptomatic Ginger Rhizomes During the
Activation of Adventitious Bud Development. *Plant Dis* 2022; **106**: 2470–2479.

97.   Wang X, Shi W, Wang J, Gao Z, Li S, Wang N, et al. Control of Southern Root-knot
Nematodes on Tomato and Regulation of Soil Bacterial Community by Biofumigation with
Zanthoxylum bungeanum Seed. *Hortic Plant J* 2021; **7**: 49–58.

98.   Chuah L-O, Yap K-P, Thong KL, Liong MT, Ahmad R, Shamila-Syuhada AK, et al.
Genome Sequence of "Anthococcus," a Novel Genus of the Family Streptococcaceae
Isolated from Flowers. *Genome Announc* 2016; **4**.

99.   Yu AO, Leveau JHJ, Marco ML. Abundance, diversity and plant-specific adaptations of
704   plant-associated lactic acid bacteria. *Environ Microbiol Rep* 2020; **12**: 16–29.

705   100.   Minervini F, Celano G, Lattanzi A, Tedone L, De Mastro G, Gobbetti M, et al. Lactic Acid
706   Bacteria in Durum Wheat Flour Are Endophytic Components of the Plant during Its Entire
707   Life Cycle. *Appl Environ Microbiol* 2015; **81**: 6736–6748.

Figures

Figure 1. Kansas precipitation gradient. Soil samples were collected from either agricultural
grain fields (Ag) or native prairies (P) (land use types) at four locations in Kansas along an
increasing precipitation gradient (Smoky Valley Ranch [SVR] prairie [latitude 38.8665, longitude
-100.9951], SVR grain field [38.8791, -100.9828], Hayes [HAY] prairie [38.8355, -99.3033], The
Land Institute [TLI] grain field [38.6943, -97.5912], TLI prairie [38.9698, -97.4690], and Konza
[KNZ] native prairie [39.1056, -96.6099]).

718
719 **Figure 2. Soil community composition is strongly differentiated across the soil sources and**
720 **substrates.** Redundancy analysis for **A**) bacterial and **B**) fungal communities from soil sources
721 used to derive soil inocula. Soil source and substrate are represented by differing colors or shapes,
722 respectively.

Figure 3. Shoot and root growth vary across drought treatment. The effect of drought treatment on **A)** shoot mass rate, **B)** root mass rate, and **C)** root/shoot ratio. Boxplot hinges represent the 1st and 3rd quartiles; whiskers represent 1.5 times the interquartile range; letters above box plots denote significant differences in estimated marginal means (*p*-value=<0.01; see Table 1 for full ANOVA table).

Figure 4. Bacterial community composition across drought treatment. A) Bacterial and taxonomic barplots for drought (D) and well-watered (W) plants. Each sample is represented by a single line with the relative abundance (non CLR transformed data) of each phylum displayed by color. **B)** Redundancy analysis on bacterial communities fit by drought treatment. Redundancy analysis was conditioning for sequencing batch and log transformed usable reads per sample. **C)** Confusion matrix depicting the classification accuracy of a random forest classifier predicting drought treatment trained on bacterial communities. Above the confusion matrix balanced accuracy is shown, for additional classifier statistics see Table S8

Figure 5. Fungal community composition across drought treatment. A) Fungal taxonomic barplots for drought (D) and well-watered (W) plants. Each sample is represented by a single line with the relative abundance (non CLR transformed data) of each class displayed by color. **B)** Redundancy analysis on fungal communities fit by drought treatment. Redundancy analysis was conditioning for sequencing batch and log transformed usable reads per sample. **C)** Confusion matrix depicting the classification accuracy of random forest a classifier predicting drought treatment trained on fungal communities. Above the confusion matrix balanced accuracy is shown, for additional classifier statistics see Table S9.

749
Figure 6. Shoot mass accumulation rate by soil inoculum. Boxplots are colored by soil inoculum for **A)** shoot mass rate. Separate
facets represent drought (D) and well-watered plants (W). Boxplot hinges represent the 1st and 3rd quartiles; whiskers represent 1.5 times
the interquartile range. **B-C)** trend lines for shoot mass rate of significant interactions of longitude (Long) with drought treatment and
genotype, respectively, for plants inoculated with prairie soils.

Tables

Table 1. Type three analysis of variance (ANOVA) for plant growth responses.

Factor	DF	sqrt(Shoot mass rate)		sqrt(Root Mass Rate)		log(Root/shoot Ratio)	
		F-value	p-value	F-value	p-value	F-value	p-value
Drought Treatment (DT)	1	471.39	**<0.01**	31.09	**<0.01**	216.09	**<0.01**
Genotype (G)	1	32.57	**<0.01**	30.34	**<0.01**	111.01	**<0.01**
Soil Inoculum (SI)	5	3.42	**<0.01**	0.4	0.85	2.14	0.06
DT x G	1	0.11	0.74	2.36	0.12	0.15	0.7
DT x SI	5	1.3	0.26	1.18	0.32	0.99	0.42
G x SI	5	1.95	0.09	0.64	0.67	2.65	**0.02**
DT x G x SI	5	0.77	0.57	0.22	0.95	0.12	0.99

757 Bold values denote significant effects, p-value<0.05

759 **Table 2.** Permutational Multivariate ANOVA (PERMANOVA) table bacterial community

760 composition of maize nodal roots.

Factor	DF	SS	R^2	F-value	p-value
Genotype (G)	1	4109	0.01	1.23	0.21
Soil Inoculum (SI)	5	34055	0.05	2.04	**<0.01**
Drought Treatment (DT)	1	7483	0.01	2.24	**<0.01**
SI x DT	5	16011	0.02	0.96	0.52
Residual	185	617812	0.91		

761 Bold values denote significant effects, p-value<0.05

763 **Table 3.** PERMANOVA table for fungal community composition of maize nodal roots.

Factor	DF	SS	R^2	F-value	p-value
Genotype (G)	1	774	0.01	1.01	0.28
Soil Inoculum (SI)	5	9152	0.09	2.40	**<0.01**
Drought Treatment (DT)	1	693	0.01	0.91	0.42
SI x DT	5	4198	0.04	1.10	0.22
Residual	116	88478	0.86		

764 Bold values denote significant effects, p-value<0.05

765 **Table 4.** Bacterial families associated with square root transformed shoot mass rate.

Taxon	Model Term	R	X	PVE$_{Res}$	Rel. Abund.	q
Lactobacillaceae	Abundance	3.92E-04	-1.52E-02	3.6	0.14	3.78E-04
Nocardiaceae	Abundance	2.84E-04	-1.43E-02	3.63	0.02	4.78E-04
Microbacteriaceae	Abundance	1.11E-04	-1.51E-02	5.14	1.17	3.04E-05
Sinobacteraceae	Abundance	1.03E-04	-1.46E-02	2.31	0.05	4.81E-03
Micrococcaceae	Abundance	2.58E-05	-1.48E-02	2.77	8.06	2.21E-03
Alcaligenaceae	Abundance	1.58E-06	-1.42E-02	3.04	0.05	1.39E-03
Corynebacteriaceae*	Abundance	-1.19E-04	-1.40E-02	6.38	0.01	7.67E-07
Sphingomonadaceae	Abundance	-1.21E-04	-1.25E-02	5.49	1.18	1.98E-05
Streptococcaceae*	Abundance	-1.61E-04	-1.39E-02	5.52	0.02	7.59E-06
Rhodospirillaceae	Abundance	-4.46E-04	-1.19E-02	5.54	0.08	1.75E-05
Dietziaceae	Abundance	-6.54E-04	-1.38E-02	4.97	<0.01	2.43E-05
Beijerinckiaceae	Abundance	-8.49E-04	-1.46E-02	2.76	<0.01	2.19E-03
Actinomycetaceae	Abundance	-1.85E-03	-1.47E-02	2.18	<0.01	5.79E-03

766 *Corynebacteriaceae and Streptococcaceae both show significant interactions between the family
767 abundance and drought treatment, these are presented in Figure S11.
768
769 **Table 5.** Fungal orders associated with square root transformed shoot mass rate.

Taxon	Trait	Model Term	R	X	PVE$_{Res}$	Relative Abundance	q
Hypocreales	SMR	Abundance	4.75E-05	-1.42E-02	3.28	19.23	0.01
Malasseziales	SMR	Abundance	9.97E-05	-1.42E-02	2.92	1.22	0.01
Sporidiobolales	SMR	Abundance	6.36E-04	-1.34E-02	4.87	0.66	0.00

771 **Supplemental Figures**

773
774

Figure S1. Experimental design. A) A complete randomized block design (blocks=27) was used.
Each pot was filled with sterile calcined clay soil mix and received a seed of either *Zea mays*
genotype B73 or Mo17, was subject to either well-watered or drought conditions, and was
inoculated with one of the six inocula. One uninoculated control was used per randomized block
(denoted with white stripe). **B)** Plant biomass was collected in a time series: for time point one
(day 25) and two (day 39), 100 plants each were destructively harvested, then dried for root and
shoot biomass. For the third time point (day 50), all remaining plants (n=529 – non-germinates)
were harvested. Prior to drying for biomass measures, one nodal root from each plant was collected
for amplicon sequencing for bacterial and fungal microbiome composition.

784
785 **Figure S2.** Bacterial alpha diversity values for **A)** Chao1 **B)** Shannon Diversity and **C)** Inverse
786 Simpson's Diversity. Black squares represent estimated marginal means (EMM) for each
787 collection site and colored bars show the 95% confidence intervals for the EMMs.

789
790 **Figure S3.** Fungal alpha diversity values for **A/C)** Chao1 and **B/D)** Inverse Simpson's Diversity.
791 Black squares represent estimated marginal means (EMM) for each collection site or substrate and
792 colored bars show the 95% confidence intervals for the EMMs.

793
794 **Figure S4.** Taxonomic bar plots for bacterial phyla for **A)** soil and **B)** inocula samples. Historical
795 annual precipitation for each collection site (SVR, HAY, TLI, KNZ) increases from left to right.
796 Each sample is represented by a single line with the relative abundance (non CLR transformed
797 data) of each phylum displayed by color.

798
Figure S5. Taxonomic bar plots for fungal classes for **A)** soil and **B)** inocula samples. Historical
annual precipitation for each collection site (SVR, HAY, TLI, KNZ) increases from left to right.
Each sample is represented by a single line with the relative abundance (non CLR transformed
data) of each class displayed by color.

Figure S6. Maize genotypes growth differs across **A)** shoot mass rate, **B)** root mass rate, and **C)**

root/shoot ratio. Boxplot hinges represent the 1st and 3rd quartiles; whiskers represent 1.5 times the

interquartile range; letters above box plots denote significant differences in estimated marginal

means (*p*-value=<0.01; see Table 1 for full ANOVA table).

808
809 **Figure S7.** Redundancy analysis for root bacterial community composition. Points are colored by
810 soil inoculum. Normal data ellipses are colored by soil inoculum assuming a multivariate normal
811 distribution.

Figure S8. Redundancy analysis for root bacterial community composition. Points are colored by
maize genotype.

815
816 **Figure S9**. Redundancy analysis for root fungal community composition. Points are colored by
817 soil inoculum. Normal data ellipses are colored by soil inoculum assuming a multivariate normal
818 distribution.

820 **Figure S10.** Redundancy analysis for fungal community composition. Points are colored by maize

821 genotype.

822
Figure S11. Confusion matrices depicting the classification accuracy of random forest classifiers
trained on bacterial data predicting **A)** Soil Inoculum and **B)** Genotype. The models were trained
using an 80:20 data split (80% train, 20% test) with 10-fold cross validation, shading represents
the proportion of predictions for a label. Labels on the left represent the predicted labels and the
labels on the bottom represent the actual labels. The model wide accuracy is given above each
confusion matrix, class-wise statistics are provided in table in Table S8.

Figure S12. Bacterial ASV importance plots per random forest classifier. Results for **A)** Drought Treatment, **B)** Soil Inoculum, and **C)** Genotype classifiers. For each random forest classifier, 10-fold cross validation was used to assess the mean decrease in accuracy of the classifier when a particular ASV is withheld. Error bars represent the standard error across the 10-folds. Only ASVs with a mean decrease in accuracy greater than 0.001 are shown.

835
Figure S13. Confusion matrices depicting the classification accuracy of random forest classifiers
trained on fungal data predicting **A)** Soil Inoculum and **B)** Genotype. The models were trained
using an 80:20 data split (80% train, 20% test) with 10-fold cross validation, shading represents
the proportion of predictions for a label. Labels on the left represent the predicted labels and the
labels on the bottom represent the actual labels. The model wide accuracy is given above each
confusion matrix, class-wise statistics are provided in table in Table S9.

Figure S14. Fungal ASV importance plots per random forest classifier. Results for **A)** Drought Treatment, **B)** Soil Inoculum, and **C)** Genotype classifiers. For each random forest classifier, 10-fold cross validation was used to assess the mean decrease in accuracy of the classifier when a particular ASV is withheld. Error bars represent the standard error across the 10-folds. Only ASVs with a mean decrease in accuracy greater than 0.001 are shown.

848
Figure S15. Root growth responses by soil inoculum. Boxplots are colored by soil inoculum for
A) root mass rate and **B)** root/shoot ratio (four R/S data points were removed to reduce the y axis
range and aid in visualization). Separate facets represent drought (D) and well-watered plants (W).
Boxplot hinges represent the 1st and 3rd quartiles; whiskers represent 1.5 times the interquartile
range.

Figure S16. Plant growth responses across longitude for all soil inocula. For each growth
response **A**) shoot mass rate, **B**) root mass rate, **C**) root/shoot ratio we plot the distribution and
trend line for well-watered and droughted plants.

858
859 **Figure S17.** Bacterial families **A)** Corynebacteriaceae and **B)** Streptococcaceae show a significant
860 interaction between CLR transformed abundance and drought treatment for shoot mass rate.

861
Figure S18. Plant growth rates across the collection time points of the study. A) Root mass
rate tended to decrease over the course of the experiment and was modulated by the drought
treatment (ANOVA; *'Timepoint × Treatment' p*-value< 0.01). **B**) Shoot mass rate tended to
increase over the course of the experiment (ANOVA; *'Timepoint' p*-value< 0.01).

Supplemental Tables

Table S1. ANOVA table for maize emergence rate.

	DF	ChiSq	p-value
Intercept	1	1072.80	**<0.01**
Genotype (G)	1	7.28	**0.01**
Soil Inoculum (SI)	5	5.42	0.37
G x SI	5	5.97	0.31

869 Bold values denote significant effects, p-value<0.05

871 **Table S2.** ANOVA table for bacterial alpha diversity metrics of soil samples collected from six

872 locations across the Kansas precipitation gradient.

Factor	DF	log(Chao1)			log(Shannon's Diversity)			log(Inverse Simpson Diversity)		
		SS	F-value	p-value	SS	F-value	p-value	SS	F-value	p-value
Substrate	1	0.57	1.14	0.29	0.08	0.11	0.74	0.01	0.01	0.92
Collection Site	5	15.9	6.37	**<0.01**	14.41	4.24	**<0.01**	10.67	3.29	**0.01**
Sequence Depth	1	81.02	162.34	**<0.01**	58.20	85.58	**<0.01**	36.96	57.06	**<0.01**
Residual	52	25.95			35.36			33.68		

873 Bold values denote significant effects, p-value<0.05

875 **Table S3.** ANOVA table for fungal alpha diversity metrics of soil samples collected from six

876 locations across the Kansas precipitation gradient.

Factor	DF	log(Chao1)			log(Shannon's Diversity)			log(Inverse Simpson Diversity)		
		SS	F-value	p-value	SS	F-value	p-value	SS	F-value	p-value
Substrate	1	10.50	45.07	**<0.01**	0.02	0.08	0.78	2.99	7.16	**0.01**
Collection Site	5	3.07	2.64	**0.03**	2.67	1.75	0.13	5.21	2.5	**0.04**
Sequence Depth	1	14.67	62.99	**<0.01**	0.45	1.47	0.23	1.28	3.07	0.08
Residual	81	18.87			24.69			33.79		

877 Bold values denote significant effects, p-value<0.05

879 **Table S4.** PERMANOVA table for bacterial and fungal community composition of soil and

880 inocula samples collected from six locations across the Kansas precipitation gradient.

Factor	Bacteria					Fungi				
	DF	SS	R^2	F-value	p-value	DF	SS	R^2	F-value	p-value
Substrate	1	44584	0.10	7.93	**<0.01**	1	32134	0.07	8.05	**<0.01**
Collection Site	5	101664	0.23	3.62	**<0.01**	5	101753	0.22	5.1	**<0.01**
Residual	53	297985	0.67			82	327269	0.71		

881 Bold values denote significant effects, p-value<0.05

882 **Table S5.** ANOVA table for bacterial alpha diversity metrics of maize nodal roots.

		Chao1			log(Shannon's Diversity)			log(Inverse Simpson Diversity)		
Factor	DF	SS	F-value	p-value	SS	F-value	p-value	SS	F-value	p-value
Genotype (G)	1	38929	2.23	0.14	0.17	0.39	0.53	0.02	0.06	0.80
Soil Inoculum (SI)	5	128720	1.48	0.2	4.11	1.92	0.09	3.46	1.81	0.11
Drought Treatment (DT)	1	110485	6.34	**0.01**	0.01	0.03	0.87	0.17	0.45	0.50
SI x DT	5	57853	0.66	0.65	2.35	1.10	0.36	2.72	1.43	0.22

883 Bold values denote significant effects, p-value<0.05

885 **Table S6.** ANOVA table for fungal alpha diversity metrics of maize nodal roots.

		Chao1			log(Shannon's Diversity)			log(Inverse Simpson Diversity)		
Factor	DF	SS	F-value	p-value	SS	F-value	p-value	SS	F-value	p-value
Genotype (G)	1	36.25	0.20	0.66	0.17	0.43	0.51	0.17	0.46	0.50
Soil Inoculum (SI)	5	1221.90	1.33	0.26	1.74	0.89	0.49	1.30	0.72	0.61
Drought Treatment (DT)	1	15.85	0.09	0.77	1.04	2.67	0.11	0.77	2.13	0.15
SI x DT	5	1496.80	1.63	0.16	5.95	3.06	**0.01**	5.21	2.89	**0.02**

886 Bold values denote significant effects, p-value<0.05

888 **Table S7.** Optimal hyperparameters for bacterial and fungal machine learning classifiers trained
889 to predict treatment, soil inoculum, and genotype.

	Bacteria		Fungi	
Factor	mtry	Minimum Node Side	mtry	Minimum Node Side
Treatment	559	10	64	10
Soil Inoculum	437	5	46	5
Genotype	559	5	10	1

891 **Table S8.** Output statistics for bacterial machine learning models predicting treatment, genotype,
892 and soil inoculum.

Metric	Treatment	Genotype	Soil Inoculum					
			SVR_P	SVR_{Ag}	HAY_P	TLI_P	TLI_{Ag}	KNZ_P
Precision	0.737	0.714	0.400	0.500	0.400	0.500	0.500	0.714
Recall	0.700	0.714	0.571	0.571	0.250	0.400	0.600	0.625
F1	0.718	0.714	0.471	0.533	0.308	0.444	0.546	0.667
Prevalence	0.500	0.700	0.175	0.175	0.200	0.125	0.125	0.200
Balanced Accuracy	0.725	0.524	0.695	0.725	0.578	0.671	0.757	0.781

894 **Table S9.** Output statistics for fungal machine learning models predicting treatment, genotype,
895 and soil inoculum.

Metric	Treatment	Genotype	Soil Inoculum					
			SVR_P	SVR_{Ag}	HAY_P	TLI_P	TLI_{Ag}	KNZ_P
Precision	0.500	0.625	0.667	0.750	0.400	0.000	0.000	0.444
Recall	0.462	0.8824	0.667	0.600	0.667	0.000	0.000	0.571
F1	0.480	0.7317	0.667	0.667	0.500	NA	NA	0.500
Prevalence	0.500	0.6538	0.115	0.192	0.115	0.269	0.038	0.269
Balanced Accuracy	0.500	0.4412	0.812	0.776	0.768	0.447	0.440	0.654

896